Unexpectedly low recombination rates and presence of hotspots in termite genomes

Turid Everitt[1#], Tilman Rönneburg[1#], Daniel Elsner[2], Anna Olsson[1], Yuanzhen Liu[1], Tuuli

Larva[1], Judith Korb[2,3], Matthew T Webster[1,4*]

1) Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden

2) Evolutionary Biology and Ecology, University of Freiburg, Freiburg, Germany

3) Research Institute for the Environment and Livelihoods, Charles Darwin University,

Casuarina Campus, Darwin, Australia

4) Science for Life Laboratory, Uppsala University, Uppsala, Sweden

# equal contribution

*correspondence to matthew.webster@imbim.uu.se

Abstract

Meiotic recombination is a fundamental evolutionary process that facilitates adaptation and

the removal of deleterious genetic variation. Social Hymenoptera exhibit some of the

highest recombination rates among metazoans, whereas high recombination rates have not

been found among non-social species from this insect order. It is unknown whether

elevated recombination rates are a ubiquitous feature of all social insects. In many

metazoan taxa, recombination is mainly restricted to hotspots a few kilobases in length.

However, little is known about the prevalence of recombination hotspots in insect genomes.

Here we infer recombination rate and its fine-scale variation across the genomes of two

social species from the insect order Blattodea: the termites *Macrotermes bellicosus* and

Cryptotermes secundus. We used linkage-disequilibrium-based methods to infer

recombination rate. We infer that recombination rates are close to 1 cM/Mb in both

species, similar to the average metazoan rate. We also observed a highly punctate

distribution of recombination in both termite genomes, indicative of the presence of

recombination hotspots. We infer the presence of full-length *PRDM9* genes in the genomes

of both species, which suggests recombination hotspots in termites might be determined by

PRDM9, as they are in mammals. We also find that recombination rates in genes are

correlated with inferred levels of germline DNA methylation. The finding of low

recombination rates in termites indicates that eusociality is not universally connected to

elevated recombination rate. We speculate that the elevated recombination rates in social

Hymenoptera are instead promoted by intense selection among haploid males.

Introduction

Meiotic recombination has two fundamental roles: It is essential for correct disjunction of

chromosomes during cell division and it generates new combinations of genetic variants

that form the raw material of evolution (Hartfield and Keightley 2012). However,

recombination can also be deleterious, as it can generate structural mutations and break up

favourable combinations of alleles. Recombination varies among species, with optimal

recombination rate in a genome likely determined by the optimal trade-off of its positive

and negative evolutionary and mechanistic effects (Stapley et al. 2017).

The highest recombination rates in metazoans observed so far are found in social insects

(Wilfert et al. 2007; Stapley et al. 2017). However, until now recombination rate has only

been studied in social insects from the order Hymenoptera, to which the majority of social

insects belong. Among Hymenoptera, the highest rates are found in the genus *Apis*: the

Western honey bee *Apis mellifera* (20.8 cM/Mb) and its relatives *Apis cerana* (17.4 cM/Mb),

Apis florea (20.8 cM/Mb) and *Apis dorsata* (25.1 cM/Mb) (Beye et al. 2006; Shi et al. 2013;

Liu et al. 2015; Wallberg et al. 2015; Rueppell et al. 2016; Kawakami et al. 2019). Other

social Hymenoptera also have high rates including the bumblebee *Bombus terrestris* (8.9

cM/Mb) (Liu et al. 2017; Kawakami et al. 2019), the stingless bee *Frieseomelitta varia* (9.3 –

12.5 cM/Mb) (Waiker et al. 2021), the wasp *Vespula vulgaris* (9.7 cM/Mb) (Sirviö et al.

2011a) and the ants *Pogonomyrmex rugosus* (11.1 cM/Mb) (Sirviö et al. 2011b) and

Acromymex echinatior (6.1 cM/Mb) (Sirviö et al. 2006). In contrast, the solitary bee

Megachile rotundata and the solitary wasp *Nasonia* have relatively low rates (1.0 and 1.5

cM/Mb, respectively) (Niehuis et al. 2010; Jones et al. 2019). For comparison, the average

recombination rate of 15 insect species outside of Hymenoptera is 2.2 cM/Mb (Wilfert et al.

2007).

We can learn about the forces shaping the evolution of recombination rates in general by

understanding why high recombination rates have evolved in social Hymenoptera. Several

hypotheses have been proposed to explain this observation. One set of explanations

focusses on the effects of recombination on intra-colony genetic diversity. For example,

increasing the genetic diversity of a colony could promote task specialization among

workers (Kent et al. 2012; Kent and Zayed 2013). Similarly, elevated genetic variation in a

colony could prevent invasion by pathogens or parasites due to diversification of immune

genes (Fischer and Schmid-Hempel 2005). Recombination could also reduce variance in

relatedness between nestmates, thereby reducing potential kin conflict within colonies

(Sherman 1979; Templeton 1979; Wilfert et al. 2007). Additionally, it has been proposed

that high recombination rates could have facilitated the evolution of eusociality over a

longer evolutionary timescale, because recombination between genes that take on

functions in different castes permits them to evolve more independently (Kent and Zayed

2013). Several studies have addressed these hypotheses (Kent et al. 2012; Liu et al. 2015;

Wallberg et al. 2015; Rueppell et al. 2016; Liu et al. 2017; Jones et al. 2019; Waiker et al.

2021; Kawakami et al. 2019) but so far none is strongly supported. It is also not clear

whether eusociality *per se* selects for higher recombination rates, or whether high rates are

a specific feature of eusocial Hymenoptera.

The main characteristics of eusocial insect species are the presence of castes that forgo

reproduction in order to care for brood or defend other colony members (workers and

soldiers). Eusocial insects are found mainly in Hymenoptera, among bees, wasps and ants,

and also Blattodea, in which all termites (Infraorder: Isoptera) are eusocial but exhibit

varying levels of social complexity. These insect orders likely diverged ~400 million years ago

(Misof et al. 2014) and there are substantial differences between social insects from the two

orders (Korb 2008; Korb and Thorne 2017). Hymenoptera belong to the superorder

Holometabola, which is the most diverse insect superorder and contains 11 orders including

Lepidoptera (butterflies, moths), Coleoptera (beetles), and Diptera (true flies), which all

undergo complete metamorphosis. Blattodea is a hemimetabolous order, which undergoes

partial metamorphosis. Hymenoptera have haplodiploid sex determination in which males

develop from haploid, unfertilized eggs and females derive from diploid, fertilized eggs. In

Blattodea, both males and females are diploid. Members of the worker caste in eusocial

Hymenoptera are all female, whereas eusocial Blattodea workers are of both sexes.

Hymenoptera colonies are headed by one or a small number of queens, whereas Blattodea

colonies contain both a king and a queen.

The hypotheses proposed to explain the high recombination rates observed in eusocial

Hymenoptera predict that recombination should be elevated in all social insects. However,

so far it is unknown whether social insects from other orders also have high rates. As

described above, there are fundamental differences in the sex determination system and

colony compositions of social insects from Hymenoptera and Blattodea. The robustness of

the association between eusociality and high recombination rates can therefore be

addressed by assessing mean genomic recombination rates in social Blattodea (termites).

In addition to variation between species, the rate of recombination also varies along

chromosomes. In mammals, crossover events are mainly restricted to regions that contain

specific motifs that act as binding sites for the PRDM9 (PR/SET domain 9) protein that marks

regions for double-stranded breaks, which are repaired by recombination during meiosis

(Baudat et al. 2010; Myers et al. 2010; Parvanov et al. 2010). Species with an active PRDM9

protein include nearly all mammals and many vertebrate taxa (Baker et al. 2017). In species

without PRDM9, recombination can be directed to hotspots defined by other features, such

as CpG islands and promoters that have open chromatin. Such hotspots have been

characterised in the genomes of diverse taxa including yeast, birds, and dogs (Axelsson et al.

2012; Lam and Keeney 2015; Singhal et al. 2015; Berglund et al. 2014). In insects, the

prevalence of recombination hotspots is not well studied. Fine-scale recombination maps in

the fruit fly *Drosophila melanogaster* and honey bee *A. mellifera* indicate a paucity of

recombination hotspots (Chan et al. 2012; Smukowski Heil et al. 2015; Wallberg et al. 2015),

whereas hotspots appear to be present in the genome of the butterfly *Leptidea sinapis*

(Torres et al. 2023).

Several other factors may also modulate recombination rate along chromosomes, such as

germline DNA methylation. Many vertebrate species have elevated recombination in CpG

islands, which are usually unmethylated in the germline (Axelsson et al. 2012; Singhal et al.

2015; Berglund et al. 2014). In insect genomes, methylation is not always present, but

commonly restricted to gene bodies (Arsala et al. 2022; Wang et al. 2013; Lyko et al. 2010;

Ventós-Alfonso et al. 2020; Bewick et al. 2019; Harrison et al. 2018). In the honey bee *A.*

mellifera and solitary bee *M. rotundata,* recombination is reduced in genes, particularly

those inferred to have the highest levels of germline methylation (Wallberg et al. 2015;

Jones et al. 2019). In addition, some studies have identified correlations between caste-

biases in gene expression and recombination. In honey bees, genes with worker-biased gene

expression have been found to have elevated recombination rates (Kent et al. 2012) but this

is likely an indirect result of differences in germline methylation between sets of genes

(Wallberg et al. 2015).

Here we estimate recombination rate and its fine-scale variation across the genomes of two

distantly related termite species with contrasting social complexities (Korb and Hartfelder

2008; Korb and Thorne 2017). *Macrotermes bellicosus* (Termitidae: Macrotermitinae) is a

fungus-growing termite with a high level of social complexity, with colonies of several

millions of individuals and sterile workers. It belongs to the foraging (multiple-pieces

nesting) termites, where workers leave the nest to forage for food. The drywood termite

Cryptotermes secundus (Kalotermitidae) has a low level of social complexity. Its colonies

consist of a few hundred individuals and totipotent workers from which queens and kings

develop (Hoffmann et al. 2012). It belongs to the wood-dwelling (one-piece nesting)

termites that nest in a single piece of wood that also serves as their food source.

The study has three main aims. Firstly, we aim to test the robustness of the association

between sociality and high recombination rate. If sociality is a universal driver of high

recombination rates, we would expect both termite species to have elevated recombination

rates, and that it is particularly elevated in *M. bellicosus*, which has a higher level of social

complexity. Secondly, we aim to determine whether recombination hotspots exist in the

two termite genomes, and infer whether active full-length *PRDM9* genes are present in their

genomes. Thirdly, we aim to analyse the factors that govern recombination rate variation

across the genome, in particular whether it is associated with patterns of methylation and

gene expression as has been reported in other studies.

Results

Genetic variation in two termite species is typical for social insects

We utilized genome assemblies of two termite species: *M. bellicosus* and *C. secundus*. The

genome assembly of *M. bellicosus* (Qiu et al. 2023) is 1.3 Gbp in total length across 275

scaffolds with scaffold N50 of 22.4 Mbp. The genome assembly of *C. secundus* (Csec_1.0)

(Harrison et al. 2018) is divided into 55,483 scaffolds with a scaffold N50 of 1.2 Mbp and a

total length of 1.0 Gbp.

We sequenced 10 unrelated individuals each of *M. bellicosus* and *C. secundus* to a mean

depth of 34x and 31x per sample, respectively (Supplemental Table S1). We mapped reads

to the appropriate genome assemblies and called variants in each species. In *M. bellicosus*

we called 5.6 million SNPs and in *C. secundus* 15.1 million SNPs (Table 1). One individual of

C. secundus (CS_8) was excluded due to a low proportion of mapped reads, poor mapping

quality and strand bias in mapping. Estimates of variation based on θ_W per bp (Watterson

1975) are 0.14% and 0.44% in *M. bellicosus* and *C. secundus* respectively, which are broadly

comparable to levels of variation in the honey bee *A. mellifera* (0.3 - 0.8%) (Wallberg et al.

2014).

We used these estimates of genetic variation to estimate effective population size (N_E) for

each of the sampled populations (Table 1). These rely on an estimate of mutation rate, μ, in

each species. As there is no estimate of μ for Blattodea, we considered a range of estimates

from insects compiled by (Lynch et al. 2023) based on experimental assays that includes

estimates from Diptera (*Drosophila*) (Keightley et al. 2009, 2014), Hymenoptera (Liu et al.

2017; Yang et al. 2015) and Lepidoptera (Keightley et al. 2015). These estimates range

between a minimum of 2.7 x 10^{-10} bp^{-1}gen^{-1} in *Acyrthosiphon pisum* (Fazalova and Nevado

2020) to a maximum of 8.1 x 10^{-9} bp^{-1}gen^{-1} in *Drosophila pseudoobscura* (Krasovec 2021).

The average of multiple estimates in *Drosophila melanogaster* is 4.5 x 10^{-9} bp^{-1}gen^{-1}, which is

typical of insects (Lynch et al. 2023). A mutation rate of μ = 1.85 x 10^{-9} bp^{-1}gen^{-1} has been

estimated in the orchid mantis, *Hymenopus coronatus* (Huang et al. 2023) based on

inference from levels of interspecific divergence. This species is a member of the order

Mantodea, which contains the closest relatives of Blattodea.

Our estimates of N_E based on θ_W and the various mutation rates are ~43,000 – 1,279,000 in

M. bellicosus, which has high social complexity, and ~137,000 – 4,081,000 in *C. secundus*,

which has a lower level of social complexity (Table 1). Estimates of N_E assuming the typical

mutation rate of 4.5 x 10^{-9} bp^{-1}gen^{-1} are 77,000 and 245,000 respectively. Assuming the

mutation rate of *H. coronatus* gives an N_E of 187,000 for *M. bellicosus* and 596,000 for *C.*

secundus. These estimates are consistent with previous evidence that eusocial species with

large colonies and high social complexity tend to have lower effective population sizes and

the values are broadly comparable to levels of N_E found in social and solitary Hymenoptera

(Leffler et al. 2012; Romiguier et al. 2014).

Low average recombination rate in termite genomes

We next examined the decay of linkage disequilibrium (LD) based on the statistic r^2. Both

termite species show more extensive LD compared to *A. mellifera*, which has an extremely

high recombination rate (Wallberg et al. 2015) (Figure 1). As N_E is broadly similar in all these

species, these differences are expected to largely represent differences in recombination

rates, and indicate that the two termite species do not exhibit the elevated genome average

recombination rate that is observed in honey bees.

We estimated variation in the population recombination rate, ρ, in both datasets using

LDhelmet (Chan et al. 2012) and LDhat (Auton and McVean 2007) (Table 1). Average rates of

ρ/kbp in *M. bellicosus* and *C. secundus* are 1.7 and 4.8 respectively from LDhat and 4.28 and

16.39 respectively from LDhelmet. These estimates can be converted to cM/Mb using

estimates of N_E, which gives average recombination rates of 0.033 - 0.986 cM/Mb in *M.*

bellicosus and 0.030 - 0.885 cM/Mb in *C. secundus* from LDhat and 0.084-2.49 cM/Mb for *M.*

bellicosus and 0.1 - 2.99 cM/Mb for *C. secundus* from LDhelmet. Using a mutation rate close

to the average for insects, the recombination rates in *M. bellicosus* and *C. secundus* are

0.550 and 0.494 cM/Mb respectively from LDhat and 1.39 and 1.67 cM/Mb respectively

from LDhelmet. Using the mutation rate from *H. coronatus*, the recombination rates are

0.23 and 0.20 cM/Mb from LDhat and 0.57 and 0.69 cM/Mb from LDhelmet for *M. bellicosus*

and *C. secundus*, respectively. Even though these estimates vary depending on the

algorithm used for estimating ρ and the assumed mutation rate, they are all substantially

lower than estimates from eusocial Hymenoptera (6-25 cM/Mb; (Sirviö et al. 2006) and

more similar to rates found in other insects (Wilfert et al. 2007). This represents the first

estimation of recombination rate in social insects outside of Hymenoptera and

demonstrates that eusociality is not a universal driver of high recombination rates.

Spearman's correlation between the results from LDhat and LDhelmet, using 10 kbp

windows, was ρ = 0.946 and ρ = 0.947 for *M. bellicosus* and *C. secundus*, respectively, with p

< 2.2 x 10^{-16} for both species. For the remaining analyses, results from the software

LDhelmet were used.

Evidence for active recombination hotspots in termite genomes

Recombination rates are highly variable across the genomes of *M. bellicosus* (Figure 2A) and

C. secundus (Figure 2B). We characterised the distribution of recombination events across

the two termite genomes using a cumulative distribution plot (Figure 3). We found that 50%

of the recombination occurs in 0.4% of the genome in *M. bellicosus* and 0.2% of the genome

in *C. secundus*. This indicates that the majority of recombination events are restricted to a

much smaller portion of the genome than observed in *A. mellifera*, where 50% of

recombination events occur in 32% of the genome (Wallberg et al. 2015).

We further investigated whether recombination hotspots are present in the two termite

genomes by looking at the distribution of ρ in windows of 1 kbp, 10 kbp and 100 kbp across

the genome (Supplemental Figure S1). For all window sizes, a small subset of windows was

observed with ρ greatly exceeding the mean in both termite species. For example, the

percentage of 1 kbp windows with recombination rate at least four standard deviations

above the genome wide mean value is 0.2% for *M. bellicosus* and 0.6% for *C. secundus*. By

contrast, *A. mellifera* has no windows above this limit for any window size, consistent with a

lack of recombination hotspots.

We next defined hotspots as 2 kbp regions with more than 5-fold higher recombination rate

than the surrounding 100 kbp and used STREME (Bailey 2021) to identify 8 to 20-mers that

were enriched in hotspots compared to the rest of the genome. This yielded significantly

enriched motifs in both species (4 motifs for *M. bellicosus* and 1 motif for *C. secundus* after

compensating for multiple hypothesis testing; Supplemental Tables S2 and S3). However,

none of the motifs we identified are present in more than 5% of hotspots or exhibit more

than 2-fold difference in frequency of occurrence relative to the background, suggesting

they are not viable candidate motifs for promoting recombination in hotspots.

Identification and characterization of the PRDM9 gene in termites

We searched the *M. bellicosus* and *C. secundus* genomes for evidence of complete *PRDM9*

genes (Figure 4A), which could potentially govern the presence of recombination hotspots in

these genomes, as it has been shown to do in mammalian genomes (Myers et al. 2010;

Baudat et al. 2010; Parvanov et al. 2010). The *C. secundus* genome contains a *PRDM9*

homolog (XP_023708049.2) with annotated KRAB, SSXRD, SET and zinc finger domains

(Figure 4B). There is no complete homolog of *PRDM9* in the *M. bellicosus* annotation. We

therefore used BLAST to search for sequences homologous to the conserved functional

domains from *C. secundus*. We identified a putative *PRDM9* orthologue on Scaffold 42 of the

M. bellicosus genome assembly, containing all domains in the correct order. We identified

retroviral conserved domains in an intron 5' of the zinc-finger domains in this gene, which

are not predicted to alter the transcript. We confirmed the presence of a full-length

transcript as well as presence and correct order of conserved domains by alignment of the

Zootermopsis nevadensis PRDM9 protein sequence to the *M. bellicosus* genome using

GeneWise (Madeira et al. 2024)(Figure 4C and Supplemental Table S4).

We used a zinc-finger prediction algorithm (Persikov and Singh 2014) to predict DNA binding

motifs from the zinc-finger domain present in PRDM9 in the *M. bellicosus* and *C. secundus*

genome assemblies. The most probable consensus motifs were

TATGGAACGACAGGAACAACGACATCATCAGCCGCCTAAT and TAATAAGTAGAATCGTTTAG for

M. bellicosus (Figure 4D) and *C. secundus* (Figure 4E) respectively. Base probability matrices

for these motifs are shown in Supplemental Table S5. We tested whether the presence of

these motifs corresponded to elevated recombination considering 1 kbp around each motif.

However, recombination rate was not significantly elevated around these predicted motifs

for either species, either when searching based on the most likely motif, or by searching

based on the probability matrices.

Genomic correlates of recombination rate

We analysed genomic correlates of recombination rate in 10 kbp windows. There is a

significant correlation between recombination and $CpG_{O/E}$ in both species with Spearman's

ρ = 0.24, p < 1 x 10^{-4} for *M. bellicosus* and Spearman's ρ = 0.10, p < 1 x 10^{-4} for *C. secundus*

(Supplemental Figure S2). A reduction of recombination rates in regions of low CpG content

is consistent with interference of germline methylation with recombination. However, as

the correlation coefficients are low, the explanatory power is weak. Correlations of similar

magnitudes were observed between recombination and GC content (Spearman's ρ =0.23 for

M. bellicosus, p < 1 x 10^{-4} and Spearman's ρ =0.017 for *C. secundus*, p < 1 x 10^{-4}). This

correlation has been associated with an effect of recombination in driving GC through GC

biased gene conversion but the relatively low correlation coefficients could indicate that this

force is not strong in termites. By contrast, the correlation between GC content and

recombination rate is much stronger in honey bee (R^2 = 0.506), which is likely caused by high

recombination rates and intense GC-biased gene conversion (Wallberg et al. 2015).

There is a weak but significant correlation between overall repeat element density and

recombination rate in *C. secundus* (Spearman's ρ = = 0.02, p < 1 x 10^{-4}) but not for *M.*

bellicosus. There is a weak but significant correlation between simple repeats and

recombination in both genomes (Spearman's ρ = 0.051, p < 2.22 x 10^{-16} for *M. bellicosus* and

Spearman's ρ = 0.055, p < 2.22 x 10^{-16} for *C. secundus*). SINE and LINE elements are also only

weakly associated with recombination rate in both genomes (SINEs: Spearman's ρ = -0.012,

not significant for *M. bellicosus* and Spearman's ρ = 0.02, p < 1 x 10^{-4} for *C. secundus*; LINEs:

Spearman's ρ = -0.02, p < 1 x 10^{-4} for *M. bellicosus* and Spearman's ρ = 0.002 not significant

for *C. secundus*). There is a significant negative correlation between gene density and

recombination for the *M. bellicosus* genome (Spearman's ρ = -0.185, p < 2.22 x 10^{-16}), but no

significant correlation for *C. secundus*. We also tested whether hotspots differ from the

background with respect to GC content. The GC content does not differ substantially

between hotspots and the rest of the genome in either species (means of 40.1% and 41.0%

for *M. bellicosus* and 40.2% and 40.3% for *C. secundus*).

We found a significant correlation between nucleotide diversity (π) and recombination in

both species (Supplemental Figure S2), with Spearman's ρ = 0.48, p < 1 x 10^{-4} for *M.*

bellicosus and Spearman's ρ = 0.31, p < 1 x 10^{-4} for *C. secundus*. This correlation is found

across a wide range of eukaryotic taxa and is generally accepted to be caused by the

interaction between recombination and linked selection (Nachman 2001; Begun and

Aquadro 1992).

Recombination and gene expression patterns correlate with $CpG_{O/E}$

Analysis of the distribution of $CpG_{O/E}$ in both termite species showed that it is bimodally

distributed, with substantially lower values in introns and exons compared to flanking

noncoding regions (Supplemental Figure S3). This indicates that germline DNA methylation

is biased towards gene bodies in the two termite genomes, a pattern that is also found in

honey bees and other insect genomes with functional DNA methylation (Harrison et al.

2018; Elango et al. 2009; Sarda et al. 2012). We found that exons have lower $CpG_{O/E}$

compared to introns, which are both lower than flanking regions (Figure 5, top row; all

comparisons p < 0.001 after Bonferroni correction) consistent with higher levels of germline

methylation in exons (Supplemental Table S6). We next analysed how recombination rate

varies among genomic features. We found that ρ/kbp is also significantly reduced in genes

compared to flanking regions for *C. secundus,* with exons also showing reduced ρ/kbp

compared to introns (Figure 5, bottom row, all p-values <0.005 after Bonferroni correction).

The same trends are observed in *M. bellicosus* although they are not significant. The

correlation between $CpG_{O/E}$ and recombination rate among genes is higher than for the

genome overall (Spearman's ρ = 0.315, p < 2.22 x 10^{-16} for *M. bellicosus*; Spearman's ρ =

0.196, p < 2.22 x 10^{-16} for *C. secundus*).

We next investigated whether patterns of gene expression between castes and sexes

(hereafter, for short 'caste-biased gene expression') were correlated with $CpG_{O/E}$ and

recombination rate using gene expression data from two studies (Lin et al. 2021; Elsner et

al. 2018). It has been proposed that it is evolutionarily advantageous for genes with worker-

biased expression to be located in regions with elevated recombination (Kent et al. 2012).

Comparisons of $CpG_{O/E}$ and recombination rate for the original gene expression categories

reported by (Elsner et al. 2018) for *M. bellicosus* are shown in Supplemental Figure S4. We

also reclassified the genes into the following expression categories: queen-biased, king-

biased, worked-biased, male-biased, female-biased, reproduction-biased, and differentially

expressed (see methods for details). For *C. secundus*, only worker-biased and queen-biased

genes were identified (Lin et al. 2021). We find that mean $CpG_{O/E}$ in gene bodies shows

significant variation between gene expression categories in both *M. bellicosus* and *C.*

secundus (Figure 6). However, this variation is not consistent between species. In *M.*

bellicosus, queen-biased and female-biased genes have low $CpG_{O/E}$ and king-biased and

male-biased genes have high $CpG_{O/E}$. In *C. secundus*, queen-biased genes have higher than

average $CpG_{O/E}$.

The differences in $CpG_{O/E}$ between caste-biased gene expression categories are mirrored by

differences in ρ/kbp (Figure 6). Gene expression categories with elevated $CpG_{O/E}$

consistently show elevated ρ/kbp, and this pattern is found in both species. Studies in the

honey bee have found that worker-biased genes have lower CpG and higher recombination

rates (Kent et al. 2012; Wallberg et al. 2015). In *M. bellicosus,* we find that worker-biased

genes have slightly elevated $CpG_{O/E}$ and ρ/kbp. However, in *C. secundus*, these values are

reduced in worker-biased genes, whereas queen-biased genes have a slightly elevated

$CpG_{O/E}$ and ρ/kbp in this species.

We investigated the relative roles of $CpG_{O/E}$ and gene expression in determining ρ in genes

using linear models, with $CpG_{O/E,}$ ρ in flanking region and expression categories as

explanatory variables. In both *M. bellicosus* and *C. secundus*, we found that $CpG_{O/E}$ was the

strongest predictor of recombination rate variation among genes (Supplemental Tables S7

and S8) in line with the analyses above. We found that caste-biased expression had no

significant impact on the recombination rate. $CpG_{O/E}$ is a better predictor of ρ/kbp in genes

than ρ/kbp in flanking region and variation in ρ/kbp among expression categories mainly

reflects differences in $CpG_{O/E}$.

Discussion

The main findings we present here are 1) that two distantly related termite species with
varying social complexity both have relatively low genomic average rates of meiotic
recombination, 2) both genomes possess recombination hotspots and likely contain full-
length copies of *PRDM9* and, 3) there are reduced levels of recombination in genomic
regions depleted in CpG sites. Our findings contrast with the extremely elevated
recombination rates observed in eusocial Hymenoptera, which are likely to be generated by
a feature that is specific to these taxa. The finding of recombination hotspots is one of the
first in insects. This could reflect a conserved function of PRDM9 in initiating recombination
events in both vertebrates and invertebrates. Our results also support a role for germline
methylation in suppressing recombination.

Low recombination rates in termites

Data from a wide range of taxa indicate that at least one crossover per chromosome tetrad
is essential for correct chromosomal segregation during meiosis (Fernandes et al. 2018). The
chromosome numbers of *M. bellicosus* and *C. secundus* are 2n = 42 and 2n = 40, respectively
(Jankásek et al. 2021). Considering their assembly lengths, this indicates that a minimum
average genomic recombination rate of 0.92 cM/Mb and 1.01 cM/Mb, respectively, would
be necessary for accurate meiosis to occur in the two termite species. Our estimates of
recombination rate per meiosis based on analysis of LD depend on several factors, including
the algorithm to estimate ρ and the estimate of mutation rate. As no estimates of the
mutation rate in termites were available, we considered a range of estimates from other
insects. These estimates vary by more than an order of magnitude between different insect
species (Lynch et al. 2023) and higher estimates of the mutation rate give higher estimates
of the recombination rate. Our estimates based on LDhat are all less than 1 cM/Mb for both
species, even when considering the highest estimate of mutation rate. Using estimates from
LDhelmet and the average of experimentally determined mutation rate estimates in insects
gives estimates of recombination rate of 1-2 cM/Mb. The species most closely related to
termites for which a mutation rate estimate was available was *H. coronatus (Huang et al.*
2023). Using this estimate and ρ from LDhelmet, the recombination rate per meiosis was
estimated to 0.57 cM/Mb for *M. bellicosus* and 0.69 cM/Mb for *C. secundus*, i.e. both
slightly lower than expected based on the requirement of one crossover per chromosome.
Taken together, these results indicate that recombination rates in these two termite species

are substantially lower than the elevated values found in eusocial Hymenoptera (6-25

cM/Mb) (Wilfert et al. 2007).

It should be noted that our estimates of recombination rate are sex-averaged rates. It is

possible that recombination is absent in one of the termite sexes, which would make a

lower sex-averaged recombination rate more feasible. In addition, sex-linked contigs have

not been identified in the termite genome assemblies utilized here. *M. bellicosus* and *C.*

secundus have X1X2/Y1Y2 and X/Y sex-determining systems respectively (Jankásek et al.

2021). The inclusion of sex chromosomes in the analysis leads to errors in determining the

genome-wide recombination rate per meiosis because they have lower N_E than the rest of

the genome. In addition, sex chromosomes are more prone to assembly and read mapping

errors, due to reads aligning with the incorrect copy of the chromosome. However,

considering that sex chromosomes only comprise a minor portion of the genome, this is

unlikely to substantially alter our results. It has also been observed that some chromosomes

form ring or chain structures during meiosis in certain termite species (Bergamaschi et al.

2007), which could plausibly relax the requirement for one crossover per chromosome per

meiosis.

Eusociality is not a universal driver of high rates of meiotic recombination

Elevated genomic rates of recombination are characteristic of all eusocial Hymenoptera so

far investigated. Several hypotheses have been advanced to explain this phenomenon,

which are all based on an evolutionary advantage of recombination in eusocial species

(Wilfert et al. 2007). For instance, it has been suggested that elevated recombination could

increase genetic diversity in a colony. This could promote diversity in the workforce,

enabling more efficient task specialisation among workers (Kent et al. 2012). Alternatively, it

could render the colony less susceptible to invasion by parasites and pathogens (Fischer and

Schmid-Hempel 2005). Elevated recombination might also favour rapid evolution of caste-

specifically expressed genes (Kent and Zayed 2013). In addition, increased recombination

could lead to reduced variance in relatedness, which could prevent kin conflict in colonies

(Sherman 1979; Templeton 1979).

All these hypotheses predict that recombination rate should be elevated in all eusocial

insects. Here, however, in the first report of recombination rate in social insects outside of

Hymenoptera, we find overall low levels of recombination. Both termite species studied

here are eusocial, and in particular *M. bellicosus* has extremely large and complex colonies

characteristic of eusociality (Korb and Thorne 2017). The low recombination rate inferred in

the two termite genomes prompts re-evaluation of hypotheses connecting eusociality with

high recombination rates. Although both termites and social Hymenoptera share many

traits, they differ in others that reflect their different ancestries.

The chromosome numbers of both termite species (*M. bellicosus*, 2n = 42; *C. secundus*, 2n =

40) are typical for termites, although variation exists between taxa (Jankásek et al. 2021).

These values do not differ from those observed in Hymenoptera, which have a similar range

(e.g. *A. mellifera*, 2n = 32) and do not vary between solitary and eusocial taxa (Cardoso et al.

2018; Cunha et al. 2021; Ross et al. 2015). This suggests that differences in chromosome

number are unlikely to be relevant in explaining the differences in recombination rates

among these taxa.

One proposed advantage of sex and recombination is that it is associated with sexual

selection and a higher intensity of selection on males. This could be advantageous because

differential male mating success can reduce mutational load in sexual populations by

causing deleterious mutations to segregate at a lower equilibrium frequency on average

(Siller 2001; Agrawal 2001). Recombination is essential for effective purging of deleterious

alleles as it unlinks deleterious variants from nearby beneficial ones thus preventing

selection interference (Hartfield and Keightley 2012). Selection has been found to be

stronger in males than females across animal species (Janicke et al. 2016; Winkler et al.

2021). Species with a male-biased sex ratio are expected to experience particularly intense

selection on males. It is therefore possible that recombination rate could be particularly

elevated in species in which males face intense competition as this increases the

evolutionary advantage of recombination.

Another proposed advantage of sex and recombination in general is that it is associated

with the intensity of sperm competition. Maynard Smith (1976) presented a model that

explains how sib-competition can produce an immediate evolutionary advantage to sex and

recombination, which occurs when several offspring of a single female compete within a

patch. This model was extended by Manning and Chamberlain (1997) who argued that it is

analogous to intra-ejaculate competition among sperm, in which recombination increases

the variance of fitness. Under these conditions, if inter-ejaculate competition also exists, it is

predicted that there will be selection for increased recombination rate. This model predicts

a significant correlation between recombination rate and gamete redundancy (excess sperm

production), which is observed in mammals (Manning and Chamberlain 1997).

Many social Hymenoptera produce highly male-biased reproductive offspring and

thousands of haploid male drones compete to fertilise a small number of female queens

(Winston 1991; Koeniger et al. 2011; Baer 2005). This situation is similar to sperm

competition as haploid males can be considered analogous to gametes. However, as many

more genes are expressed in adult haploid males compared to gametes there are more

targets of selection. Selection against deleterious alleles is particularly effective in haploid

males because recessive deleterious alleles are exposed (Hedrick and Parker 1997). It is

therefore possible that the male-biased sex ratio observed in many eusocial Hymenoptera

could favour elevated recombination rates in the species so far studied, due to elevated

competition between males.

Male-biased sex ratios are particularly strong in the genus *Apis*, which is also the genus in

which the most extreme recombination rates have been inferred (Beye et al. 2006; Shi et al.

2013; Liu et al. 2015; Wallberg et al. 2015; Rueppell et al. 2016; Kawakami et al. 2019).

Male-biased sex ratios are also observed in other social bees and wasps. Both male and

female biased operational sex ratios have been observed in ants, but are difficult to

measure because workers invest more resources in female offspring and may manipulate

sex ratios. In non-social Hymenoptera, females lay roughly equal numbers of haploid and

diploid eggs (Trivers and Hare 1976). The sex ratio of reproductive males and females is also

relatively even in most termite species (Roisin 2001).

Further research is needed to determine the cause of variation in recombination rates

among social and non-social insects. For example, simulations could be used to determine

the effect of selection on haploid males on recombination rates, considering different sex

ratios. In addition, it is necessary to estimate recombination rate in a wider range of social

and non-social species to determine whether the strength of selection on males is a key

factor in determining these rates.

Full length PRDM9 gene and recombination hotspots in termite genomes

Our results are also indicative of the presence of recombination hotspots in the two termite

genomes. The fine-scale distribution of recombination rate in both termite genomes shows

that 50% of the recombination happens in less than 0.5% of the genome, which is even

more extreme than for humans where this corresponds to ~6% of the genome (Myers et al.

2005). This contrasts to the honey bee and solitary bees where fine-scale maps are available

and recombination is relatively uniformly distributed in the genome (Wallberg et al. 2015;

Jones et al. 2019).

Two main mechanisms have been shown to lead to recombination hotspots. The first, which

is best understood in human and mouse, is governed by the PRDM9 protein. A zinc-finger

domain recognises a specific DNA sequence motif and then performs a histone modification

in the vicinity, which marks the sequence for a DNA double-stranded break that is repaired

by recombination during meiosis (Baudat et al. 2010; Myers et al. 2010; Parvanov et al.

2010). In species with this mechanism, both the PRDM9 zinc-finger domain and the

sequences present in hotspots are fast evolving, which results in a rapid turnover of hotspot

locations (Myers et al. 2010). Full-length PRDM9 orthologues are present among a range of

highly-diverged vertebrates including many fish, reptiles and mammals, but many full or

partial losses of the gene have also been inferred. In a study of 225 vertebrate genomes, a

minimum of six partial and three complete losses were inferred (Baker et al. 2017).

However, it is unclear whether PRMD9-hotspots are common in invertebrates.

A second mechanism is found in vertebrates that lack PRDM9, in which recombination

hotspots are localised to regions of open chromatin such as CpG islands and promoters. This

is found in dogs and in most or all birds, in which PRDM9 has been lost, and results in

hotspots with stable rather than rapidly evolving locations (Axelsson et al. 2012; Singhal et

al. 2015). A similar pattern is also observed in the yeast *Saccharomyces cerevisiae* in which

recombination preferentially occurs in promoter regions (Wu and Lichten 1994). PRDM9-

independent hotspots can be identified in CpG islands in some vertebrate genomes even

when PRDM9 is present (Joseph et al. 2024). In invertebrates such as *Drosophila* and

Caenorhabditis elegans, recombination hotspots appear to be absent (Coop and Przeworski

2007; Chan et al. 2012; Smukowski Heil et al. 2015). In insects, fine-scale maps of social and

solitary bees have not revealed recombination hotspots (Wallberg et al. 2015; Jones et al.
2019), but there is evidence for hotspots in a butterfly genome (Torres et al. 2023).

We find evidence that recombination is directed towards hotspots in both termite species.
We also identify a full-length *PRDM9* gene in both species, containing all of the domains
needed to initiate recombination. However, we are unable to demonstrate that PRMD9 is
responsible for the presence of recombination hotspots in the two genomes. We used a
motif prediction algorithm to predict the zinc-finger binding motif in *M. bellicosus* and *C.*
secundus, but we did not observe elevated recombination rate around instances of the
motif in either of the genomes. Furthermore, we did not identify any common sequence
motifs that were strongly enriched in recombination hotspots in either of the termite
genomes. A plausible interpretation of these observations is that the PRDM9 protein indeed
initiates recombination in hotspots in both species, but that positive selection leads to
frequent shifts in the target motifs, which results in a lack of a strong association between a
specific motif and LD-based hotspots (Myers et al. 2010). It is also possible that the
sequences of the zinc-finger motifs are not correctly represented in the genome assemblies,
which could result from problems with assembly around minisatellites. However, an
unknown mechanism could also be responsible for hotspots in termites.

Evidence that germline DNA methylation suppresses recombination

In insects with a functioning DNA methyltransferase machinery, the $CpG_{O/E}$ statistic shows
substantial variation along the genome, which mainly reflects variation in levels of germline
DNA methylation (Elango et al. 2009). In contrast to vertebrate genomes, methylation is
strongly skewed towards gene bodies in the genomes of insects (Glastad et al. 2011). In the
two termite genomes studied here, we find lowest values of $CpG_{O/E}$ in exons, whereas
noncoding flanking regions are biased towards higher values, indicative of lower levels of
DNA methylation (Figure 5, Supplemental Figure S3). Exons, introns and noncoding flanking
regions all display a predominantly bimodal distribution of $CpG_{O/E}$, likely reflective of the
presence of methylated and unmethylated regions. We find that recombination rate is
reduced in gene bodies compared to flanking regions in one of the two termite genomes,
but the overall genomic correlation between $CpG_{O/E}$ and recombination rate, although
significant, is weak. This could reflect the fact that the majority of the genomes are not
methylated, so that much of the variation in $CpG_{O/E}$ on the genome scale is not strongly

influenced by methylation. Our inference of reduced recombination rates in exons could

also be influenced by lower N_E in these regions due to the effects of linked selection, which

could result in more extensive LD (Charlesworth 2009). However, the observation that

differences in $CpG_{O/E}$ among genes are associated with recombination rate indicates that

linked selection is unlikely to be a major factor driving the observed differences in

recombination rate and that germline DNA methylation is more important.

We identified $CpG_{O/E}$ as the factor with the strongest influence on recombination rate

variation among genes. There were no significant effects of differences in gene expression

patterns across castes, a result that was also observed in the honey bee genome (Wallberg

et al. 2015). This indicates that recombination rate variation is not modulated by differences

in gene expression in these genomes, which might be expected if natural selection favours

increased recombination rates in certain genes. For example, it has been proposed that

natural selection could favour elevated recombination rates in genes with roles in worker

behaviour or in immune function due to their important roles in colony function (Wilfert et

al. 2007; Kent et al. 2012). However, the results presented here suggest that differences in

recombination rate between genes with caste-biased patterns of gene expression are

mainly due to underlying differences in methylation levels among genes.

A direct effect of germline DNA methylation in suppressing recombination could influence

variation in recombination landscapes among vertebrates and invertebrates. Vertebrate

genomes are usually highly methylated with the exception of CpG islands. In vertebrate

genomes that lack a functional *PRDM9* gene, recombination events tend to localise to CpG

islands (for example in dogs and birds) (Singhal et al. 2015; Axelsson et al. 2012). The effect

of DNA methylation on recombination rate variation in invertebrates is less clear, but the

results presented here also support a role of germline methylation in suppressing

recombination.

Materials and Methods

Sample collection

M. bellicosus samples were collected from colonies in Kakpin, next to the Comoé National

Park in Côte d'Ivoire (coordinates 8°39⬚N 3°46⬚W) (Elsner et al. 2018). *C. secundus* colonies

were collected from dead *Ceriops tagal* mangrove trees near Palmerston-Channel Island

(Darwin Harbor, Northern Territory, Australia; 12°30⯑S 131°00⯑E). Colonies were then kept

in *Pinus radiata* wood blocks in climate rooms in Germany, providing 28⯑°C, 70% relative

humidity, and a 12-h day/night cycle (Lin et al. 2021). For each species, each of the 10

collected samples came from a different colony.

Population sequencing

The termites were cut in two, approximately between head and thorax. DNA was extracted

from both parts using the QIAGEN Blood and Tissue Kit following the standard protocol,

including treatment with 4 µL RNase and 25 µL Proteinase K. The DNA extraction product

was cleaned and concentrated with ZymoResearch Genomic DNA Clean & Concentrator kit.

For most samples, the RNase treatment was repeated prior to cleaning and concentrating.

One DNA sample from each individual was chosen, based on the DNA amount and quality,

to be sequenced. Sequencing libraries were produced using the Illumina DNA PCR-free

library preparation kit according to the manufacturers protocol, using an input DNA quantity

of 200 ng per sample. This protocol yields insert sizes of approximately 350 bp. We

performed Illumina short-read sequencing on the libraries using NovaSeq 6000 S4 flow cell

with 2 x 150 bp paired end reads.

Read mapping and variant calling

The sequence data from the two termite species were mapped to the reference genomes

for *M. bellicosus* (Qiu et al. 2023) and *C. secundus* (Csec_1.0) (Harrison et al. 2018)

respectively, using Burrows-Wheeler alignment tool BWA version 0.7.17 (Li & Durbin 2009)

with the BWA-MEM algorithm. Sorting and indexing of the BAM-files was done with the

SAMtools package version 1.17 (Li et al. 2009), followed by adding read groups and marking

duplicate reads with Picard toolkit version 1.118 (http://broadinstitute.github.io/picard/).

After mapping, we excluded all scaffolds < 1 kbp from the *C. secundus* genome, which in

total correspond to 2% of the genome size, as they are potentially of lower quality and not

informative for analysis of linkage disequilibrium or recombination. For *M. bellicosus* there

are no scaffolds shorter than 1 kbp, but we excluded the following five scaffolds, which

together correspond to 3.78% of the genome length, due to unexpectedly high

heterozygosity: 93, 62, 41, 35 and 32.

594 One of the *C. secundus* individuals did not map well to the reference genome (71% mapped

595 reads, unequal distribution between forward and reverse strand and an average mapping

596 quality of 35 while all the other samples had an average mapping quality >45). This

597 individual was excluded from all further analyses and hence the total number of termite

598 individuals analysed in this study is 19, including 10 *M. bellicosus* and 9 *C. secundus*.

599 Variant calling was done using the tools HaplotypeCaller, GenomicsDBImport and

600 GenotypeGVCFs from GATK version 4.3.0.0 as described in their best practices workflow

601 (https://gatk.broadinstitute.org/hc/en-us/articles/360035535932-Germline-short-variant-

602 discovery-SNPs-Indels-, last accessed 2023-07-11). The variants were filtered in multiple

603 steps, first removing extended regions of low mapping quality or deviating read depth as

604 those regions are likely to be unreliable. This was done based on statistics from the

605 SAMtools mpileup and depth tools, respectively, removing 100 kbp windows with mean

606 mapping quality below 70 or mean read depth more than two standard deviations from the

607 genome-wide mean. Indels were removed using VCFtools version 0.1.16 (Danecek et al.

608 2011) before GATK VariantFiltration was run with the following limits, which are all equal to

609 or stricter than the ones recommended (https://gatk.broadinstitute.org/hc/en-

610 us/articles/360035890471-Hard-filtering-germline-short-variants, last accessed 2023-07-11)

611 and adapted to the observed distributions of the corresponding statistics: QD<2; FS>50;

612 MQ<40; ReadPosRankSum< -4; ReadPosRankSum>4; SOR>3; ExcessHet>5 for both species

613 as well as MQRankSum with limits -5 and 5 for *C. secundus* and limits -6 and 4 for *M.*

614 *bellicosus*. This was followed by additional filtering with VCFtools to select only bi-allelic

615 sites with a minimum quality score of 30 and maximum of 40% missing genotypes. The SNPs

616 with mean read depth in the lowest or highest 2.5% of the depth distribution were also

617 filtered out, which translates to a depth below 10 or above 34 for *C. secundus* and below 5

618 or above 40 for *M. bellicosus.* For the recombination rate calculations, the SNPs were also

619 filtered for a minor allele count >= 2, as rare variants appearing on only one chromosome

620 have a relatively high false positive rate and are not informative regarding LD. The average

621 proportion of missing genotypes per individual was 4.5×10^{-5} for *C. secundus* and 1.2×10^{-4}

622 for *M. bellicosus*. After variant calling and filtering, the data was phased and imputed using

623 Beagle version 5.1 (Browning and Browning 2007; Browning et al. 2018).

624 *Estimation of LD decay*

The decay of linkage disequilibrium, r^2, with increasing physical distance was estimated with

the software PopLDDecay version 3.42 (Zhang et al. 2019) over a maximum distance of 10

kbp. For this analysis, sequence data from *A. mellifera scutellata* was included for

comparison (Wallberg et al. 2017). The *A. m. scutellata* sequences were mapped to the

Amel_HAv3.1 reference genome (Wallberg et al. 2019) and processed and filtered in the

same way as described above for the termite data (GATK VariantFiltration with the limits

QD<2, FS>25, MQ<40, MQRankSum<-4, MQRankSum>4, ReadPosRankSum<-4,

ReadPosRankSum>4, SOR>3, ExcessHet>5 and VCFtools to select bi-allelic sites with mean

read depth between 5-13, quality score >= 30, minor allele count >=2 and <=40% missing

genotypes). For *M. bellicosus*, the reference genome contains some gaps of unknown size

and in order to avoid estimating r^2 across those gaps, scaffold breaks were introduced at the

corresponding positions.

Estimation of population recombination rate

We estimated the population-scaled recombination rate, ρ, using two methods: LDhat

(Auton and McVean 2007) and LDhelmet (Chan et al. 2012), which are both based on a

Bayesian reversible-jump Markov Chain Monte Carlo algorithm (rjMCMC).

For LDhat (https://github.com/auton1/Ldhat, downloaded 2023-04-19), the input files were

generated from the filtered vcf-file with VCFtools and the --ldhat option. Then the LDhat

function complete was run in order to create a lookup table, with the estimated value of θ_W

(see below), a maximum ρ value of 100 and 101 grid points, as recommended. This was

followed by the main function, interval, to perform the rjMCMC with 10 million iterations,

sampling every 5000th iteration and a block penalty of 1 as this has been used previously

with LDhat in similar studies (Wallberg et al. 2015; Jones et al. 2019). In order to summarise

the output from interval, the stat function was used, discarding the first 20 samples as burn-

in.

The recombination rate was also estimated using the software LDhelmet version 1.10 (Chan

et al. 2012). In order to prepare the input files for LDhelmet, the filtered vcf-files were

converted into multiple-sequence fasta-files using the vcf2fasta function from vcflib version

2017-04-04 (Garrison et al. 2022) and to the related file formats .snps and .pos with the - -

ldhelmet option from VCFtools (version 0.1.16). The first step of the LDhelmet workflow is

to create haplotype configuration files with the find_confs tool, followed by generation of

likelihood lookup tables with the tool table_gen and computation of Padé coefficients with

the tool pade. These tools were run with the recommended parameters, meaning that

find_confs was run with a window size of 50 SNPs, table_gen was run with a grid of ρ-values

ranging from 0 to 100 with increments of 0.1 up to 10 and increments of 1 for the remaining

grid and pade was run to generate 11 Padé coefficients. Scaffolds shorter than or equal to

the window size of 50 SNPs were removed, corresponding to 1% of the sequence length in

C. secundus and even less for *M. bellicosus*. The tool rjmcmc which contains the main

algorithm was run with a burn-in of 100,000 followed by 1,000,000 iterations with a block

penalty of 50 as recommended (Chan et al. 2012). For each species, the rjmcmc tool was run

three times with different seeds and the convergence between the replicates was evaluated

in terms of Spearman's rank correlation coefficient, which was > 0.95 with p < 2.2 x 10^{-16} for

all pairwise comparisons of both species. The binary output from rjmcmc was converted to

text, extracting the mean ρ-value between each pair of SNPs, using the tool post_to_text.

The outputs from LDhat and LDhelmet give the value of the parameter ρ between each pair

of consecutive SNPs. Those values were converted to equally sized windows across each

scaffold, as well as the coordinates of individual genomic features, by taking a weighted

average of values from overlapping intervals. The *M. bellicosus* scaffolds contain a few gaps

of unknown size and the values between SNPs across such gaps were removed.

The parameter ρ is proportional to the recombination rate r and the effective population

size N_E, which in turn can be estimated based on the mutation rate μ and the number of

segregating sites, K. In order to convert ρ to r, the following equations were used, where n is

the number of haploid sequences. The number of segregating sites (K) was calculated before

the filter on minor allele count >=2 was applied, as this filter likely removes many true SNPs

as well, which could affect the estimates.

θ_W was estimated for each sample using the equation below from (Watterson 1975):

$$\theta_W = K \left/ \sum_{i=1}^{n-1} \frac{1}{i} \right.$$

681   We used the value of θ_W to estimate N_E using:

$$N_E = \frac{\theta_W}{4\mu}$$

682 We estimated N_E using a range of mutation rates estimated from other insect species

683 (Keightley et al. 2009, 2014, 2015; Yang et al. 2015; Liu et al. 2017; Lynch et al. 2023). We

684 then estimated the recombination rate using the following equation, assuming a range of

685 values of N_E.

$$r = \frac{\rho}{4N_E}$$

Cumulative plot of rho ρ/kbp

A cumulative plot of the proportion of recombination events versus the proportion of the

physical distance along the genome was constructed based on estimates of ρ/bp for the two

termite species and previous estimates for *A. mellifera* (Wallberg et al. 2015; PRJNA236426).

The ρ/bp estimates between markers were placed in decreasing order before multiplication

with the respective physical distances between the markers. Then cumulative sums were

calculated for the ρ-values as well as the corresponding physical distances and divided by

the total genome-wide sums.

k-mer enrichment in recombination hotspots

In order to identify *k*-mers associated with elevated recombination rate, we defined

recombination hotspots as 2 kbp segments with at least five-fold elevated recombination

rate compared to the local background, as defined by 50 kbp flanking up and downstream of

the segment, resulting in 39837 regions for *M. bellicosus* and 40894 regions for *C. secundus.*

Then, we used these hotspots to search for motifs associated with elevated recombination

rate by comparing against an equivalent number of randomly chosen 2 kbp regions from the

rest of the genome. This was done using STREME 5.5.4 (Bailey 2021) using a minimum and

maximum size of *k*-mer of 8 and 20, (--minw 8 ; --maxw 20) and otherwise default options.

From all motifs with significant enrichment after adjusting for multiple testing, we then

considered only motifs present in more than 5% of all hotspots and an at least two-fold

enrichment compared to the background as credible candidates.

Identifying PRDM9 orthologues in M. bellicosus and C. secundus

A full-length copy of PRDM9 is present in the *C. secundus* genome annotation, but not in *M.*

bellicosus. Using OrthoFinder (Emms and Kelly 2019), and the genome annotation files for

both species, we identified a single orthologue to the *C. secundus* PRDM9 in *M. bellicosus*.

We analysed this region using the NCBI conserved domain search (Wang et al. 2023). We

also searched the *M. bellicosus* assembly for the PR/SET domains derived from *C. secundus*

and *Zootermopsis nevadensis* using BLAST (Camacho et al. 2009) and subsequently for all

other PRDM9 domains (KRAB, SSXRD, ZF-Casette) to identify hits that contain all other

PRDM9 domains in the vicinity in the correct order. We used GeneWise (Madeira et al.

2024) with default options to perform gapped alignment of a the PRDM9 protein sequence

from *Zoothermopsis nevadensis* against the target region to predict exon-intron borders.

Predicting binding sites from zinc-finger-motifs

Using the ZF-motifs from *C.secundus,* we obtained letter probability matrices for predicted

binding motifs using a webserver (http://zf.princeton.edu/index.php ; (Persikov and Singh

2014) which were used to search for binding sites across the genome using FIMO (Grant et

al. 2011) with default options. Subsequently, the mean rho for the surrounding 1 kbp bin

was used to test for a difference in mean rho between bins associated with a suspected

binding site, and a similar number of randomly chosen bins without association. For

significance testing, we repeated this draw 10.000 times to compare the associated bins

against the 95% confidence interval of the resulting distribution.

Analysis of genomic correlates of recombination

We divided the genome into 10 kbp windows in which average ρ and other genomic

features were estimated. Windows shorter than 5 kbp (half of the specified window size),

which appear in short scaffolds or at the ends at longer scaffolds, were excluded from the

analyses (excluding less than 1% of the sequence for each species). Correlation coefficients

were estimated with Spearman's ρ, with significance estimated by permutation tests with

10,000 iterations.

In order to determine the repeat content across the *M. bellicosus* and *C. secundus* genome

assemblies, we first generated custom repeat libraries for each species with RepeatModeler

(Flynn et al. 2020) using default options. This was then combined with Repbase 29 (Bao et

al. 2015) and Dfam 3.8 (Storer et al. 2021) and used to run RepeatMasker with default

options on each genome. We estimated the proportion of sequence in each repeat class in

10 kbp windows and used Spearman's rank correlation to correlate this with estimates of

recombination rate.

Per gene recombination rate and CpG$_{O/E}$ ratio

The observed/expected frequency of CpG sites (CpG$_{O/E}$) and mean ρ were calculated on a

per-gene basis, as well as for exons and introns, and 50 kbp flanking regions 10 kbp

upstream and downstream of the gene, akin to Wallberg et al (2015). For recombination

rate, estimates of ρ between markers were used, utilising a weighted mean when the

element spanned multiple markers. For visualisation, extreme CpG$_{O/E}$ values due to low GC

content were set to a maximum of 4.

We tested for significant difference in mean between ρ and CpG$_{O/E}$ in flanking regions, exons

and introns using paired *t*-tests corrected for multiple testing using a Bonferroni threshold.

Differential expression data

We classified genes in both species according to their caste-biased patterns of expression

from two studies (Elsner et al. 2018; Lin et al. 2021). Elsner et al. (2018) analysed four castes

of *M. bellicosus*, with two age classes of each: major and minor workers, queens and kings.

They analysed differences in gene expression among these classes using RNA-seq data. We

reclassified the differential expression data by grouping the direct caste-vs-caste

comparisons as "queen-biased", "king-biased", "worker-biased", "reproduction-biased",

"male-biased" or "female-biased" if they were differentially upregulated for this caste (or

group) in comparison to other castes (or groups). Since Elser *et al.* (2018) initially mapped

the *M. bellicosus* reads against the *Macrotermes natalensis* genome, We used OrthoFinder

(Emms and Kelly 2019) to identify the *M. bellicosus* orthologues corresponding to the

differentially expressed genes. Only single-copy genes were considered for this analysis.

Lin et al. (2021), performed RNA-seq on samples of workers and queens. Here, we used the

data from untreated queens and workers. Differentially expressed *C. secundus* genes were

classified as "worker-biased" or "queen-biased" if they were differentially expressed with a

bias to the respective caste, or unbiased if they were not, and subsequently compared

between classes. The lists of genes we identified in the caste-biased categories in both

species are provided as Supplemental Tables S9 and S10.

Correlates of recombination rate variation among genes

In order to assess the relative importance of $CpG_{O/E}$ and differential expression patterns on

recombination rate in genes (genic ρ/kbp), two ordinary least squares models were built

using statsmodels (Seabold and Perktold 2010) for each termite species. Genic $CpG_{O/E}$,

differential expression categories and flanking region ρ/kbp were used as exogenous

variables. In the first model, we included flanking region ρ/kbp and $CpG_{O/E}$. In the second

model, we included all available biased expression categories for each species.

Data access

The Illumina whole-genome sequencing data generated in this study have been submitted

to the NCBI BioProject database (https://www.ncbi.nlm.nih.gov/bioproject/) under

accession number PRJNA1021607. Custom scripts are available on GitHub

(https://github.com/troe27/termite-recombination-rate) and as Supplemental Code.

Competing interest statement

The authors declare no competing interests.

Acknowledgements

We acknowledge grant 2018-03896 from The Swedish Research Council to MTW and from

the Deutsche Forschungsgemeinschaft (DFG) KO-1895/26-1: 417980976 to JK. We

acknowledge support from the National Genomics Infrastructure in Stockholm funded by

Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish

Research Council. The computations were enabled by resources in project NAISS 2023/23-

402 provided by the National Academic Infrastructure for Supercomputing in Sweden

(NAISS) at UPPMAX, funded by the Swedish Research Council through grant agreement no.

2022-06725. We thank Göran Arnqvist for helpful discussions on the evolution of sex and

recombination and Marie Raynaud for assistance running LDhelmet. The Parks and Wildlife

Commission, Northern Territory, and the Department of the Environment, Water, Heritage

and the Arts, Australia, gave permission to collect (Permit number 59044) and export

(PWS2016-001559) the *C. secundus* termites. The Office Ivoirien des Parcs et Réserves

(OIPR) provided sampling and export permits for the *M. bellicosus* samples. The study was

conducted according to the Nagoya protocol. MTW designed and led the research; TE and

TR conducted the analysis; MTW, TE, TR and JK wrote the paper; DE and JK contributed data

and samples; AO, YL, and TL performed lab work.

References

Agrawal AF. 2001. Sexual selection and the maintenance of sexual reproduction. *Nature*
411: 692–695.

Arsala D, Wu X, Yi SV, Lynch JA. 2022. Dnmt1a is essential for gene body methylation and
the regulation of the zygotic genome in a wasp. *PLOS Genetics* **18**: e1010181.

Auton A, McVean G. 2007. Recombination rate estimation in the presence of hotspots.
Genome Research **17**: 1219–27.

Axelsson E, Webster MT, Ratnakumar A, Consortium L, Ponting CP, Lindblad-Toh K. 2012.
Death of PRDM9 coincides with stabilization of the recombination landscape in the
dog genome. *Genome Res* **22**: 51–63.

Baer B. 2005. Sexual selection in Apis bees. *Apidologie* **36**: 187–200.

Bailey TL. 2021. STREME: accurate and versatile sequence motif discovery. *Bioinformatics*
37: 2834–2840.

Baker Z, Schumer M, Haba Y, Bashkirova L, Holland C, Rosenthal GG, Przeworski M. 2017.
Repeated losses of PRDM9-directed recombination despite the conservation of
PRDM9 across vertebrates ed. B. De Massy. *eLife* **6**: e24133.

Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in
eukaryotic genomes. *Mobile DNA* **6**: 11.

Baudat F, Buard J, Grey C, Fledel-Alon A, Ober C, Przeworski M, Coop G, de Massy B. 2010.
PRDM9 is a major determinant of meiotic recombination hotspots in humans and
mice. *Science* **327**: 836–840.

Begun DJ, Aquadro CF. 1992. Levels of naturally occurring DNA polymorphism correlate with
recombination rates in D. melanogaster. *Nature* **356**: 519–520.

Bergamaschi S, Dawes-Gromadzki TZ, Scali V, Marini M, Mantovani B. 2007. Karyology,
mitochondrial DNA and the phylogeny of Australian termites. *Chromosome Res* **15**:
735–753.

Berglund J, Quilez J, Arndt PF, Webster MT. 2014. Germline methylation patterns determine
the distribution of recombination events in the dog genome. *Genome Biol Evol* **7**:
522–530.

Bewick AJ, Sanchez Z, Mckinney EC, Moore AJ, Moore PJ, Schmitz RJ. 2019. Dnmt1 is
essential for egg production and embryo viability in the large milkweed bug,
Oncopeltus fasciatus. *Epigenetics & Chromatin* **12**: 6.

Beye M, Gattermeier I, Hasselmann M, Gempe T, Schioett M, Baines JF, Schlipalius D,
Mougel F, Emore C, Rueppell O, et al. 2006. Exceptionally high levels of
recombination across the honey bee genome. *Genome research* **16**: 1339–44.

Browning BL, Zhou Y, Browning SR. 2018. A One-Penny Imputed Genome from Next-
Generation Reference Panels. *The American Journal of Human Genetics* **103**: 338–
348.

Browning SR, Browning BL. 2007. Rapid and accurate haplotype phasing and missing-data
inference for whole-genome association studies by use of localized haplotype
clustering. *Am J Hum Genet* **81**: 1084–1097.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. *BMC Bioinformatics* **10**: 421.

Cardoso DC, Santos HG, Cristiano MP. 2018. The Ant Chromosome database – ACdb: an
online resource for ant (Hymenoptera: Formicidae) chromosome researchers.
Myrmecological News **27**: 87–91.

Chan AH, Jenkins PA, Song YS. 2012. Genome-Wide Fine-Scale Recombination Rate Variation
in Drosophila melanogaster. *PLOS Genetics* **8**: e1003090.

Charlesworth B. 2009. Fundamental concepts in genetics: effective population size and
patterns of molecular evolution and variation. *Nat Rev Genet* **10**: 195–205.

Coop G, Przeworski M. 2007. An evolutionary view of human recombination. *Nature reviews*
Genetics **8**: 23–34.

Cunha MS, Cardoso DC, Cristiano MP, de Oliveira Campos LA, Lopes DM. 2021. The Bee
Chromosome database (Hymenoptera: Apidae). *Apidologie* **52**: 493–502.

Elango N, Hunt BG, Goodisman MA, Yi SV. 2009. DNA methylation is widespread and
associated with differential gene expression in castes of the honeybee, Apis
mellifera. *Proceedings of the National Academy of Sciences of the United States of*
America **106**: 11206–11.

Elsner D, Meusemann K, Korb J. 2018. Longevity and transposon defense, the case of
termite reproductives. *PNAS* **115**: 5504–5509.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biology* **20**: 238.

Fazalova V, Nevado B. 2020. Low Spontaneous Mutation Rate and Pleistocene Radiation of
Pea Aphids. *Mol Biol Evol* **37**: 2045–2051.

Fernandes JB, Séguéla-Arnaud M, Larchevêque C, Lloyd AH, Mercier R. 2018. Unleashing
meiotic crossovers in hybrid plants. *Proceedings of the National Academy of Sciences*
115: 2431–2436.

Fischer O, Schmid-Hempel P. 2005. Selection by parasites may increase host recombination
frequency. *Biol Lett* **1**: 193–195.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. 2020.
RepeatModeler2 for automated genomic discovery of transposable element families.
Proceedings of the National Academy of Sciences **117**: 9451–9457.

Garrison E, Kronenberg ZN, Dawson ET, Pedersen BS, Prins P. 2022. A spectrum of free
software tools for processing the VCF variant call format: vcflib, bio-vcf, cyvcf2, hts-
nim and slivar. *PLOS Computational Biology* **18**: e1009123.

Glastad KM, Hunt BG, Yi SV, Goodisman M a. D. 2011. DNA methylation in insects: on the
brink of the epigenomic era. *Insect Molecular Biology* **20**: 553–565.

Grant CE, Bailey TL, Noble WS. 2011. FIMO: scanning for occurrences of a given motif.
Bioinformatics **27**: 1017–1018.

Harrison MC, Jongepier E, Robertson HM, Arning N, Bitard-Feildel T, Chao H, Childers CP,
Dinh H, Doddapaneni H, Dugan S, et al. 2018. Hemimetabolous genomes reveal
molecular basis of termite eusociality. *Nature Ecology & Evolution* **2**: 557–566.

Hartfield M, Keightley PD. 2012. Current hypotheses for the evolution of sex and
recombination. *Integrative Zoology* **7**: 192–209.

Hedrick PW, Parker JD. 1997. Evolutionary Genetics and Genetic Variation of Haplodiploids
and X-Linked Genes. *Annual Review of Ecology and Systematics* **28**: 55–83.

Hoffmann K, Foster KR, Korb J. 2012. Nest value mediates reproductive decision making
within termite societies. *Behavioral Ecology* **23**: 1203–1208.

Huang G, Song L, Du X, Huang X, Wei F. 2023. Evolutionary genomics of camouflage
innovation in the orchid mantis. *Nat Commun* **14**: 4821.

Janicke T, Häderer IK, Lajeunesse MJ, Anthes N. 2016. Darwinian sex roles confirmed across
the animal kingdom. *Science Advances* **2**: e1500983.

Jankásek M, Kotyková Varadínová Z, Šťáhlavský F. 2021. Blattodea Karyotype Database. *EJE*
118: 192–199.

Jones JC, Wallberg A, Christmas MJ, Kapheim KM, Webster MT. 2019. Extreme Differences in
Recombination Rate between the Genomes of a Solitary and a Social Bee. *Mol Biol*
Evol **36**: 2277–2291.

Joseph J, Prentout D, Laverré A, Tricou T, Duret L. 2024. High prevalence of PRDM9-
independent recombination hotspots in placental mammals. *Proceedings of the*
National Academy of Sciences **121**: e2401973121.

Kawakami T, Wallberg A, Olsson A, Wintermantel D, de Miranda JR, Allsopp M, Rundlöf M,
Webster MT. 2019. Substantial Heritable Variation in Recombination Rate on
Multiple Scales in Honeybees and Bumblebees. *Genetics* **212**: 1101–1119.

Keightley PD, Ness RW, Halligan DL, Haddrill PR. 2014. Estimation of the Spontaneous
Mutation Rate per Nucleotide Site in a Drosophila melanogaster Full-Sib Family.
Genetics **196**: 313–320.

Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, Davey JW,
Jiggins CD. 2015. Estimation of the Spontaneous Mutation Rate in Heliconius
melpomene. *Mol Biol Evol* **32**: 239–243.

Keightley PD, Trivedi U, Thomson M, Oliver F, Kumar S, Blaxter ML. 2009. Analysis of the
genome sequences of three Drosophila melanogaster spontaneous mutation
accumulation lines. *Genome Res* **19**: 1195–1201.

Kent CF, Minaei S, Harpur BA, Zayed A. 2012. Recombination is associated with the
evolution of genome structure and worker behavior in honey bees. *PNAS* **109**:
18012–18017.

Kent CF, Zayed A. 2013. Evolution of recombination and genome structure in eusocial
insects. *Communicative & Integrative Biology* **6**: e22919.

Koeniger G, Koeniger N, Phiancharoen M. 2011. Comparative Reproductive Biology of
Honeybees. In *Honeybees of Asia* (eds. H.R. Hepburn and S.E. Radloff), pp. 159–206,
Springer, Berlin, Heidelberg https://doi.org/10.1007/978-3-642-16422-4_8 (Accessed
February 13, 2024).

Korb J. 2008. Termites, hemimetabolous diploid white ants? *Frontiers in Zoology* **5**: 15.

Korb J, Hartfelder K. 2008. Life history and development - a framework for understanding
developmental plasticity in lower termites. *Biological Reviews* **83**: 295–313.

Korb J, Thorne B. 2017. Sociality in Termites. In *Comparative Social Evolution* (eds. D.R.
Rubenstein and P. Abbot), pp. 124–153, Cambridge University Press, Cambridge
https://www.cambridge.org/core/books/comparative-social-evolution/sociality-in-
termites/11A80CB9FA947760889676E03899E85C (Accessed August 9, 2023).

Krasovec M. 2021. The spontaneous mutation rate of Drosophila pseudoobscura. *G3*
(Bethesda) **11**: jkab151.

Lam I, Keeney S. 2015. Nonparadoxical evolutionary stability of the recombination initiation
landscape in yeast. *Science* **350**: 932–937.

Leffler EM, Bullaughey K, Matute DR, Meyer WK, Ségurel L, Venkat A, Andolfatto P,
Przeworski M. 2012. Revisiting an Old Riddle: What Determines Genetic Diversity
Levels within Species? *PLoS Biol* **10**: e1001388.

Lin S, Werle J, Korb J. 2021. Transcriptomic analyses of the termite, Cryptotermes secundus,
reveal a gene network underlying a long lifespan and high fecundity. *Commun Biol* **4**:
1–12.

Liu H, Jia Y, Sun X, Tian D, Hurst LD, Yang S. 2017. Direct Determination of the Mutation Rate
in the Bumblebee Reveals Evidence for Weak Recombination-Associated Mutation
and an Approximate Rate Constancy in Insects. *Mol Biol Evol* **34**: 119–130.

Liu H, Zhang X, Huang J, Chen J-Q, Tian D, Hurst LD, Yang S. 2015. Causes and consequences
of crossing-over evidenced via a high-resolution recombinational landscape of the
honey bee. *Genome Biol* **16**: 15.

Lyko F, Foret S, Kucharski R, Wolf S, Falckenhayn C, Maleszka R. 2010. The Honey Bee
Epigenomes: Differential Methylation of Brain DNA in Queens and Workers. *PLoS*
Biol **8**: e1000506.

Lynch M, Ali F, Lin T, Wang Y, Ni J, Long H. 2023. The divergence of mutation rates and
spectra across the Tree of Life. *EMBO reports* **24**: e57561.

Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R, Butcher S.
2024. The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024.
Nucleic Acids Research gkae241.

Manning JT, Chamberlain AT. 1997. Sib competition and sperm competitiveness: an answer
to 'Why so many sperms?' and the recombination/sperm number correlation.
Proceedings of the Royal Society of London Series B: Biological Sciences **256**: 177–
182.

Maynard Smith J. 1976. A short-term advantage for sex and recombination through sib-
competition. *Journal of Theoretical Biology* **63**: 245–258.

Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J, Flouri T,
Beutel RG, et al. 2014. Phylogenomics resolves the timing and pattern of insect
evolution. *Science* **346**: 763–767.

Myers S, Bottolo L, Freeman C, McVean G, Donnelly P. 2005. A fine-scale map of
recombination rates and hotspots across the human genome. *Science* **310**: 321–4.

Myers S, Bowden R, Tumian A, Bontrop RE, Freeman C, Macfie TS, McVean G, Donnelly P.
2010. Drive against hotspot motifs in primates implicates the PRDM9 gene in meiotic
recombination. *Science* **327**: 876–9.

Nachman MW. 2001. Single nucleotide polymorphisms and recombination rate in humans.
Trends in genetics : TIG **17**: 481–5.

Niehuis O, Gibson JD, Rosenberg MS, Pannebakker BA, Koevoets T, Judson AK, Desjardins
CA, Kennedy K, Duggan D, Beukeboom LW, et al. 2010. Recombination and Its Impact
on the Genome of the Haplodiploid Parasitoid Wasp Nasonia. *PLOS ONE* 5: e8597.

Parvanov ED, Petkov PM, Paigen K. 2010. Prdm9 controls activation of mammalian
recombination hotspots. *Science* **327**: 835–835.

Persikov AV, Singh M. 2014. De novo prediction of DNA-binding specificities for Cys2His2
zinc finger proteins. *Nucleic Acids Research* **42**: 97–108.

Qiu B, Elsner D, Korb J. 2023. High-quality long-read genome assemblies reveal evolutionary
patterns of transposable elements and DNA methylation in termites.
2023.10.31.564968.
https://www.biorxiv.org/content/10.1101/2023.10.31.564968v1 (Accessed January
8, 2024).

Roisin Y. 2001. Caste sex ratios, sex linkage, and reproductive strategies in termites. *Insectes*
soc **48**: 224–230.

Romiguier J, Lourenco J, Gayral P, Faivre N, Weinert LA, Ravel S, Ballenghien M, Cahais V,
Bernard A, Loire E, et al. 2014. Population genomics of eusocial insects: the costs of a
vertebrate-like effective population size. *Journal of Evolutionary Biology* **27**: 593–
603.

Ross L, Blackmon H, Lorite P, Gokhman VE, Hardy NB. 2015. Recombination, chromosome
number and eusociality in the Hymenoptera. *Journal of Evolutionary Biology* **28**:
105–116.

Rueppell O, Kuster R, Miller K, Fouks B, Rubio Correa S, Collazo J, Phaincharoen M, Tingek S,
Koeniger N. 2016. A New Metazoan Recombination Rate Record and Consistently
High Recombination Rates in the Honey Bee Genus Apis Accompanied by Frequent
Inversions but Not Translocations. *Genome Biol Evol* **8**: 3653–3660.

Sarda S, Zeng J, Hunt BG, Yi SV. 2012. The Evolution of Invertebrate Gene Body Methylation.
Mol Biol Evol **29**: 1907–1916.

Seabold S, Perktold J. 2010. Statsmodels: Econometric and Statistical Modeling with Python.
In *Proceedings of the 9th Python in Science Conference* (eds. S. Van der Walt and J.
Millman), pp. 92–96.

Sherman PW. 1979. Insect Chromosome Numbers and Eusociality. *The American Naturalist*
113: 925–935.

Shi YY, Sun LX, Huang ZY, Wu XB, Zhu YQ, Zheng HJ, Zeng ZJ. 2013. A SNP Based High-Density
Linkage Map of Apis cerana Reveals a High Recombination Rate Similar to Apis
mellifera. *PLOS ONE* **8**: e76459.

Siller S. 2001. Sexual selection and the maintenance of sex. *Nature* **411**: 689–692.

Singhal S, Leffler EM, Sannareddy K, Turner I, Venn O, Hooper DM, Strand AI, Li Q, Raney B,
Balakrishnan CN, et al. 2015. Stable recombination hotspots in birds. *Science* **350**:
928–932.

Sirviö A, Gadau J, Rueppell O, Lamatsch D, Boomsma JJ, Pamilo P, PAGE Jr. RE. 2006. High
recombination frequency creates genotypic diversity in colonies of the leaf-cutting
ant Acromyrmex echinatior. *Journal of Evolutionary Biology* **19**: 1475–1485.

Sirviö A, Johnston JS, Wenseleers T, Pamilo P. 2011a. A high recombination rate in eusocial
Hymenoptera: evidence from the common wasp Vespula vulgaris. *BMC Genetics* **12**:
95.

Sirviö A, Pamilo P, Johnson RA, Page Jr Robert E, Gadau J. 2011b. ORIGIN AND EVOLUTION
OF THE DEPENDENT LINEAGES IN THE GENETIC CASTE DETERMINATION SYSTEM OF
POGONOMYRMEX ANTS. *Evolution* **65**: 869–884.

Smukowski Heil CS, Ellison C, Dubin M, Noor MAF. 2015. Recombining without Hotspots: A
Comprehensive Evolutionary Portrait of Recombination in Two Closely Related
Species of Drosophila. *Genome Biology and Evolution* **7**: 2829–2842.

Stapley J, Feulner PGD, Johnston SE, Santure AW, Smadja CM. 2017. Variation in
recombination frequency and distribution across eukaryotes: patterns and
processes. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**:
20160455.

Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. 2021. The Dfam community resource of
transposable element families, sequence models, and genome annotations. *Mobile*
DNA **12**: 2.

Templeton AR. 1979. Chromosome Number, Quantitative Genetics and Eusociality. *The*
American Naturalist **113**: 937–941.

Torres AP i, Höök L, Näsvall K, Shipilina D, Wiklund C, Vila R, Pruisscher P, Backström N.
2023. The fine-scale recombination rate variation and associations with genomic
features in a butterfly. *Genome Res* **33**: 810–823.

Trivers RL, Hare H. 1976. Haploidploidy and the Evolution of the Social Insect. *Science* **191**:
249–263.

Ventós-Alfonso A, Ylla G, Montañes J-C, Belles X. 2020. DNMT1 Promotes Genome
Methylation and Early Embryo Development in Cockroaches. *iScience* **23**: 101778.

Waiker P, de Abreu FCP, Luna-Lucena D, Freitas FCP, Simões ZLP, Rueppell O. 2021.
Recombination mapping of the Brazilian stingless bee Frieseomelitta varia confirms
high recombination rates in social hymenoptera. *BMC Genomics* **22**: 673.

Wallberg A, Bunikis I, Pettersson OV, Mosbech M-B, Childers AK, Evans JD, Mikheyev AS,
Robertson HM, Robinson GE, Webster MT. 2019. A hybrid de novo genome assembly

of the honeybee, Apis mellifera, with chromosome-length scaffolds. *BMC Genomics*
20: 275.

Wallberg A, Glémin S, Webster MT. 2015. Extreme recombination frequencies shape
genome variation and evolution in the honeybee, *Apis mellifera. PLoS Genetics* **11**:
e1005189.

Wallberg A, Han F, Wellhagen G, Dahle B, Kawata M, Haddad N, Simões ZLP, Allsopp MH,
Kandemir I, De la Rúa P, et al. 2014. A worldwide survey of genome sequence
variation provides insight into the evolutionary history of the honeybee *Apis*
mellifera. Nat Genet **46**: 1081–1088.

Wallberg A, Schöning C, Webster MT, Hasselmann M. 2017. Two extended haplotype blocks
are associated with adaptation to high altitude habitats in East African honey bees.
PLoS Genetics **13**: e1006792.

Wang J, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu S, Marchler GH, Song JS,
Thanki N, Yamashita RA, et al. 2023. The conserved domain database in 2023.
Nucleic Acids Res **51**: D384–D388.

Wang X, Wheeler D, Avery A, Rago A, Choi J-H, Colbourne JK, Clark AG, Werren JH. 2013.
Function and Evolution of DNA Methylation in Nasonia vitripennis. *PLoS Genet* **9**:
e1003872.

Watterson GA. 1975. On the number of segregating sites in genetical models without
recombination. *Theoretical Population Biology* **7**: 256–276.

Wilfert L, Gadau J, Schmid-Hempel P. 2007. Variation in genomic recombination rates
among animal taxa and the case of social insects. *Heredity (Edinb)* **98**: 189–197.

Winkler L, Moiron M, Morrow EH, Janicke T. 2021. Stronger net selection on males across
animals eds. P.J. Wittkopp and L. Holman. *eLife* **10**: e68316.

Winston ML. 1991. *The Biology of the Honey Bee*. Revised edition. Harvard University Press,
Cambridge, Mass.

Wu T-C, Lichten M. 1994. Meiosis-Induced Double-Strand Break Sites Determined by Yeast
Chromatin Structure. *Science* **263**: 515–518.

Yang S, Wang L, Huang J, Zhang X, Yuan Y, Chen J-Q, Hurst LD, Tian D. 2015. Parent–progeny
sequencing indicates higher mutation rates in heterozygotes. *Nature* **523**: 463–467.

Zhang C, Dong S-S, Xu J-Y, He W-M, Yang T-L. 2019. PopLDdecay: a fast and effective tool for
linkage disequilibrium decay analysis based on variant call format files.
Bioinformatics **35**: 1786–1788.

1076

1077 **Figure Legends:**
1078

Figure 1. Genome-wide decay of LD measured as r^2 versus distance (bp) between SNPs for
the two termite species (*M. bellicosus*, dark red; *C. secundus*, yellow) and honey bees (*A.*
mellifera scutellata, blue; data from Wallberg et al. 2017).

Figure 2. Variation in recombination rate estimated as ρ/kbp across representative genomic
scaffolds in a) *M. bellicosus* and b) *C. secundus*. Estimates of ρ/kbp are calculated in 10 kbp
and 100 kbp windows.

Figure 3. Cumulative plot of proportion of recombination events versus proportion of the
genome for the two termite species *M. bellicosus* and *C. secundus* compared to the honey
bee *A. mellifera* (Wallberg et al. 2015). Dashed lines show the proportion of the genome
where 50 % of the recombination occurs, which for *M. bellicosus* is 0.4 %, *C. secundus* is 0.2
%, and *A. mellifera* is 32 %.

Figure 4. PRMD9 structure and predicted zinc-finger motifs in the *M. bellicosus* and *C.*
secundus genomes. A) Predicted structure of the PRDM9 protein sequence in *C. secundus*. B)
Predicted structure of the *PRDM9* gene in *M. bellicosus* identified by homology searches.
Exons and conserved functional domains are marked. The mapped exons from
Zoothermopsis nevadensis are also shown. C) Structure of *PRDM9* gene in *C. secundus*.
Exons and conserved functional domains are marked. D) Predicted zinc finger binding motif
in *M. bellicosus,* E) Predicted zinc finger binding motif in *C. secundus*.

Figure 5. Boxplots showing variation in levels of $CpG_{O/E}$ and recombination rate (ρ/kbp; rho)
in genic and flanking regions in the *M. bellicosus* and *C. secundus* genomes. The differences
between all categories are significant.

Figure 6. Boxplots showing variation in levels of $CpG_{O/E}$ and recombination rate (ρ/kbp) in
the coding region of genes in the *M. bellicosus* and *C. secundus* genomes classified according
to their patterns of gene expression (Q = queen-biased, K = king-biased, W = worker-biased,
F = female-biased, R = reproductive-biased).

1104

1105

1106 **Table 1.** Estimates of levels genetic variation, effective population size and recombination rate in two termite species.

1107

Species	Assembly length (Mb)	No. SNPs	θ_W/bp	μ bp^{-1} gen^{-1}	N_E	LDhat		LDhelmet	
						ρ/kb	r (cM/Mb)	ρ/kb	r (cM/Mb)
M. bellicosus	1,139	5,581,662	0.14%	2.7 x 10-10	1,278,982	1.69	0.033	4.28	0.084
				4.5 x 10-9	76,876		0.550		1.392
				8.1 x 10-9	42,898		0.986		2.494
C. secundus	992	15,037,468	0.44%	2.7 x 10-10	4,080,725	4.85	0.030	16.39	0.100
				4.5 x 10-9	245,280		0.494		1.671
				8.1 x 10-9	136,869		0.885		2.994

θ_W/bp - Watterson's θ per base pair

ρ/kbp - population recombination parameter, ρ, per kilobase

μ bp^{-1} gen^{-1} - mutation rate per base pair per generation taken from literature. Estimates correspond to *Acyrthosiphon pisum* (2.7 x 10^{-10}), *Drosophila*
melanogaster (4.5 x 10^{-9}) and *Drosophila pseudoobscura* (8.1 x 10^{-9})(Lynch et al. 2023).

N_E - effective population size

r (cM/Mb) - recombination rate in centimorgans per megabase

1114

A *M. bellicosus*

B *C. secundus*