Adaptive evolution of sesquiterpene deoxyphomenone in mycoparasitism by *Hansfordia*

***pulvinata* associated with horizontal gene transfer from *Aspergillus* species**

Kazuya Maeda[1,2], Takuya Sumita[3], Oumi Nishi[3], Hirotoshi Sushida[3,4], Yumiko Higashi[3],

Hiroyuki Nakagawa[5], Tomoko Suzuki[6], Eishin Iwao[1], Much Zaenal Fanani[1], Yoshiaki

Nishiya[2], Yuichiro Iida[1,3,*]

[*] **Correspondence to Y. Iida (yuichiro.iida@setsunan.ac.jp)**

[1] Laboratory of Plant Pathology, Faculty of Agriculture, Setsunan University, Nagaotoge-cho

45-1, Hirakata, Osaka 573-0101, Japan

[2] Laboratory of Environmental Microbiology, Division of Life Science, Graduate School of

Science and Engineering, Setsunan University, 17-8 Ikeda-Nakamachi, Neyagawa, Osaka

572-8508, Japan

[3] National Agriculture and Food Research Organization (NARO), Institute of Vegetable and

Floriculture Science, Kusawa 360, Tsu-city, Mie 514-2392, Japan

[4] Institute of Food Research, National Agriculture and Food Research Organization (NARO),

2-1-12 Kannondai, Tsukuba, Ibaraki 305-8642, Japan

[5] Research center for Advanced Analysis, Core Technology Research Headquarters, National

Agriculture and Food Research Organization (NARO), 2-1-12 Kannondai, Tsukuba, Ibaraki

305-8642, Japan

[6] Department of Chemical Biological Sciences, Faculty of Science, Japan Women's

University, Mejirodai 2-8-1, Bunkyo-ku, Tokyo 112-8681, Japan

Present address:

T. Sumita, Department of biological resources management, School of Environmental

Science, The University of Shiga Prefecture, Hasaka-cho 2500, Hikone, Shiga 522-8533

Oumi Nishi, Laboratory of Insect Pathology and Microbial Control, Institute of Biological

Control, Kyushu University, Motooka 744, Nishi-ku, Fukuoka-city, Fukuoka, 819-0395,

Japan

T. Suzuki, Center for Gene Research, Nagoya University, Furo-cho, Chikusa-ku, Nagoya-

city, Aichi 464-8602, Japan; Mass Spectrometry and Microscopy Unit, Technology Platform

Division, RIKEN Center for Sustainable Resource Science, Yokohama 230-0045, Japan

Abstract

Leaf mold caused by the ascomycete fungus *Cladosporium fulvum* is a devastating disease of tomato plants. The mycoparasitic fungus *Hansfordia pulvinata* is an effective biocontrol agent that parasitizes *C. fulvum* hyphae on leaves and secretes 13-deoxyphomenone, an eremophilane-type sesquiterpene, which was also identified as a sporulation-inducing factor in *Aspergillus oryzae*. Here, we identified deoxyphomenone biosynthesis (*DPH*) gene clusters conserved in both *H. pulvinata* and *Aspergillus* section *Flavi* including *A. oryzae* and *A. flavus*. Functional disruption of *DPH1* orthologous genes encoding sesquiterpene cyclase in *H. pulvinata*, *A. oryzae* and its close relative *A. flavus* revealed that deoxyphomenone in *H. pulvinata* had exogenic antifungal activity against the host fungus *C. fulvum* and controlled endogenic sporulation in *Aspergillus* species. Deoxyphomenone also inhibited mycelial growth of *C. fulvum* and the non-host tomato pathogen *Pseudocercospora fuligena*. Complete *DPH* clusters, highly similar to those in *H. pulvinata*, were exclusive to *Aspergillus* section *Flavi*, while species in other *Aspergillus* sections contained fragmented *DPH* clusters. A comparative genomics analysis revealed that these *DPH* gene clusters share a common origin and are horizontally transferred across large taxonomic distances from an ancestor of *Aspergillus* to *H. pulvinata*. Our results suggest that, after horizontal transfer, *H. pulvinata* maintained the *DPH* cluster as the inhibitory effect of deoxyphomenone on spore germination and mycelial growth contributed to its mycoparasitism on the host fungus *C. fulvum*.

Keywords: mycoparasite, deoxyphomenone, sporogen-AO 1, horizontal gene transfer, secondary metabolite, *Aspergillus flavus*, *Aspergillus oryzae*, *Cladosporium fulvum*, *Fulvia fulva, Hansfordia pulvinata, Dicyma pulvinata*

INTRODUCTION

The biotrophic filamentous fungus *Cladosporium fulvum* (Cooke) (syn. *Passalora fulva*, *Fulvia fulva*), the causal agent of leaf mold of tomato, is an economic problem in most countries that grow tomatoes (Videira et al., 2017; Mesarich et al., 2023). Following spore germination, hyphae enter the leaf through stomata to absorb nutrients from its host in the apoplast. After a few weeks, conidiophores emerge from the stomata and produce numerous spores on the leaves, which appear as brown lesions. Colonization of tomato leaves by *C. fulvum* is facilitated by numerous effector proteins that are secreted into the apoplast by the fungus (de Wit, 2016). To combat *C. fulvum*, breeders have introduced *Cf* resistance genes, which encode receptor-like proteins with extracytoplasmic leucine-rich repeats, into commercial tomato cultivars from wild relatives (de Wit, 2016; Mesarich, 2023). Tomato *Cf* gene encoded receptor-like proteins recognize corresponding effector proteins of *C. fulvum* as virulence factors in the apoplastic region and trigger a hypersensitive response that rapidly kills plant cells near the infection site, inhibiting the growth of the biotrophic pathogen and preventing its spread (Stergiopoulos and de Wit, 2009). However, the occasional loss or mutation of effector genes allows *C. fulvum* to escape the host immune system, resulting in the emergence of new races (de Wit, 2016). Although single resistance genes have been introduced into many tomato cultivars, overuse of resistant cultivars with only one or a few *Cf* genes has led to the development of *C. fulvum* races that have overcome these resistance genes. So far, 13 races have developed in Japan (Iida et al., 2015; Kubota et al., 2015; Yoshida et al., 2021), and *C. fulvum* has evolved to sequentially overcome multiple *Cf* genes (Rosa et al., 2023). The complex race structure of *C. fulvum* in Japan has enabled the fungus to overcome all *Cf* genes present in commercial cultivars (Iida et al., 2015). A further problem is that *C. fulvum* has developed resistance to several fungicides (Yan et al., 2007; Watanabe et al., 2017), and applications of fungicides have to be reduced due to

environmental concerns. Thus, the use of antagonistic microorganisms instead of synthetic

fungicides is desired for sustainable control of leaf mold.

We serendipitously discovered white mycelia growing over lesions of leaf mold-infected

tomato leaves in a greenhouse and found that disease progression was inhibited on these

leaves (**Figure 1A**) (Iida et al., 2018). In further studies, we showed that the fungus

parasitizes hyphae of *C. fulvum* and inhibited all known races of this fungus (Iida et al.,

2018). The mycoparasitic fungus was identified as *Hansfordia pulvinata* (Berk. & M.A.

Curtis) S. Hughes [syn. *Dicyma pulvinata* (Berk. & M.A. Curtis) Arx] (Hughes, 1951;

Deighton, 1972; von Arx, 1981, 1982) based on its morphological characteristics and a

phylogenetic analysis (Iida et al., 2018). The fungus also has biocontrol activity against other

plant parasitic fungi, including *Mycosphaerella berkeleyi*, causing late leaf spot of peanut,

Fusicladium heveae causing South American leaf blight of rubber, and *Epichloë typhina*

causing choke disease of grasses (Peresse and Lepicard, 1980; Mitchell and Taber, 1986;

Mitchell et al., 1986; Mello et al., 2008; Alderman et al., 2010); it thus suppresses foliar

fungal diseases of different host plants. We also demonstrated that isolate 414-3 of *C. fulvum*

suppressed tomato leaf mold most effectively in the greenhouse (Iida et al., 2018). However,

the mechanism of this mycoparasitism of *H. pulvinata* against *C. fulvum* is not yet clear.

H. pulvinata produces 13-deoxyphomenone, an eremophilane-type sesquiterpene, as an

antifungal compound during vegetative growth (Tirilly et al., 1983). Around the same time,

this compound was also found and characterized (but named sporogen-AO 1) from

Aspergillus oryzae, which is used to produce koji that is used for the production of many

Japanese fermented foods such as Japanese sake and soy sauce (Tanaka et al., 1984a, 1984b).

This sesquiterpene is toxic to *C. fulvum*, which is parasitized by *H. pulvinata* but it induces

spore production in *A. oryzae*. In addition to its antifungal and sporogenic activities,

deoxyphomenone is also phytotoxic (Tansakul et al., 2014; Del Valle et al., 2015),

antimalarial (Daengrot et al., 2015), and cytotoxic to cancer cell lines (Tansakul et al., 2017;

Liu et al., 2021). However, its biosynthetic pathway in these fungi, the molecular basis of its

antifungal activity and spectrum, and the specificity its sporogenic activity in different

Aspergillus species have not been elucidated.

Fungal secondary metabolites play important roles in development, pathogenicity,

protection from biotic and abiotic stresses, antagonism, and communication with other

organisms (Keller, 2019). Because deoxyphomenone is produced by most strains of *H.*

pulvinata (Iida et al., 2018) and might be widely produced among *Aspergillus* species, it may

have mycoparasitic and/or ecological functions in these fungi. In this study, we investigated

the roles of deoxyphomenone in *H. pulvinata* and *Aspergillus* species. The antifungal

spectrum of the compound was determined for *C. fulvum* and other tomato fungal pathogens

that are not host fungi of *H. pulvinata*, such as *Pseudocercospora fuligena*, the causal agent

of tomato black leaf mold, which is phylogenetically closely related to *C. fulvum* and the

soilborne tomato pathogen *Fusarium oxysporum* f. sp. *lycopersici*. We identified

deoxyphomenone biosynthetic gene clusters in *H. pulvinata* and *Aspergillus* genome

sequences and functionally disrupted sesquiterpene cyclase genes to reveal the main function

of deoxyphomenone. Furthermore, we used a comparative genomics approach to elucidate

the origin of these gene clusters in *Aspergillus* and *H. pulvinata*.

RESULTS

Antifungal activity of deoxyphomenone against plant pathogens

H. pulvinata 414-3, the most effective biocontrol strain with a published draft genomic

sequence (Sushida et al. 2019) and the highest deoxyphomenone production among several

strains we tested, parasitizes *C. fulvum* hyphae in liquid culture without a carbon source (Iida

et al., 2018). In our present tests, *H. pulvinata* did not parasitize *C. fulvum* when grown on

nylon membranes on nutrient-rich potato dextrose agar (PDA). However, when the

membranes were transferred to nutrient-poor water agar, scanning electron micrographs

showed that hyphae of strain 414-3 penetrated the cell wall of *C. fulvum* (**Figure 1B**).

Although we previously reported that hyphae of *H. pulvinata* coil around the host hyphae, it

does not always do so, and it does not form appressorium-like structures, both common

attributes of the mycoparasitic *Trichoderma* fungi.

Since strain 414-3 produces about 100 µM deoxyphomenone in minimal medium (MM)

broth (Iida et al., 2018), we next evaluated its antifungal activity against *C. fulvum* using 120

µM commercial deoxyphomenone. At 120 µM, the compound inhibited colony formation

and hyphal elongation by more than 90% on MM agar (**Supplementary figure 1A, 1B**).

Deoxyphomenone also strongly inhibited spore germination at 120 µM, which was only 7.2%

after 24 h compared with 76.5% for the control spores treated with 1% (v/v) methanol

(**Figure 1C and Supplementary figure 1C**). When *C. fulvum* spores were treated with the

fungicide captan at 100 µM and after 24 h captan was replaced by distilled water, almost all

failed to germinate, but when deoxyphomenone (120 µM) was replaced by distilled water 24

h after treatment, about half of the treated spores did germinate (**Supplementary figure**

S1C). Spores treated continuously with deoxyphomenone or captan for 48 h failed to

germinate. Thus, the activity of deoxyphomenone is fungistatic; it inhibits the germination of

C. fulvum spores but does not kill them. Deoxyphomenone treatment did not affect the

structure of intracellular organelles or the cell membrane of the spores or induce necrosis in

tomato leaves even at concentrations as high as 80 µM, contrary to a previous report (Tirilly

et al., 1983) (**Supplementary figure S1D, E**). When plated on MM agar, external application

of deoxyphomenone also inhibited colony formation of plant pathogenic fungi that are not

hosts of *H. pulvinata*: the growth of *P. fuligena* phylogenetically closely related to *C. fulvum*,

was inhibited at the same concentrations as used for *C. fulvum*, but growth of *F. oxysporum* f.

sp. *lycopersici* was not affected (**Figure 1D**).

Deoxyphomenone biosynthetic gene clusters are conserved between *H. pulvinata* and

Aspergillus

To identify the genes involved in deoxyphomenone biosynthesis in the genome sequences of

H. pulvinata and *A. oryzae*, we searched for homologous genes using the amino acid

sequence of aristolocene synthase Ari1 from *Penicillium roqueforti* (Proctor and Hohn,

1993), since the structure of deoxyphomenone has an aristolochene-like backbone (**Figure**

1C), and found two candidates (gene IDs, DIP_001954 and DIP_006271) in *H. pulvinata*

414-3 (Sushida et al. 2019) and one (GenBank accession XP_023093357) in *A. oryzae*

RIB40. Phylogenetic comparisons with other fungal sesquiterpene cyclases placed

DIP_001954 and XP_023093357 in the same clade as enzymes that catalyze the 1,10-

cyclization of farnesyl diphosphate, a common sesquiterpene precursor (**Supplementary**

figure S2). The deduced amino acid sequences of DIP_001954 and XP_023093357 had

61.6% similarity (49.1% identity) (**Supplementary figure S3**). We named DIP_001954

"*HpDPH1*" (<u>d</u>eoxy<u>ph</u>omenone biosynthesis gene 1) and the homologous gene

XP_023093357 "*AoDPH1*".

In the nucleotide sequences around *HpDPH1* we found genes encoding short-chain

dehydrogenase (*HpDPH2*), three cytochrome P450s (*HpDPH3, HpDPH4, HpDPH5*), and

major facilitator superfamily transporter (*HpDPH6*), altogether forming the deoxyphomenone

biosynthesis (*DPH*) gene cluster. We also found homologs of these genes in *A. oryzae*

(**Figure 2A**). Also genes encoding a lipase/thioesterase family protein and an integral

membrane protein were conserved in both fungi, but their function in deoxyphomenone

biosynthesis is unknown. The position and orientation of all genes were highly conserved, but

the loci of *AoDPH4* and *AoDPH5* were switched (**Supplementary figure S4**). Comparison

of these genes revealed that the number of exons in the pairs *HpDPH1/AoDPH1* and

HpDPH5/AoDPH5 was identical and that the sequence similarity in each exon was conserved

among all pairs (**Supplementary figure S5**). We presumed that the deoxyphomenone

biosynthetic pathway involved cyclization of farnesyl diphosphate by HpDph1, followed by

oxygenation and epoxidation by cytochrome P450s HpDph3, HpDph4 and HpDph5, and

subsequently hydroxylated by HpDph2 (**Figure 2B**). The putative substrate binding pocket

that is conserved in the major facilitator superfamily transporter (Quistgaard et al., 2016) was

also conserved in HpDph6/AoDph6 (**Supplementary figure S6**).

To obtain a comprehensive view of the distribution of the *DPH* cluster in other fungi, we

extended the search to *HpDPH1* homologues in the MycoCosm database

(mycocosm.jgi.doe.gov; Grigoriev, 2014). Genes partially homologous to *HpDPH1* were

found in 139 fungal genomes of the *Eurotiomycetes*, *Sordariomycetes* and *Dothideomycetes*

(**Supplementary table S1**). Besides the *DPH* gene cluster present in *H. pulvinata*, other *DPH*

genes were detected as a gene cluster in *Aspergillus* species only (*Eurotiomycetes*), but not in

the *Xylariales* (*Sordariomycetes*), to which *H. pulvinata* belongs, which suggests that the

latter fungus has obtained this cluster by horizontal transfer. Notably, the cluster was

completely conserved in *Aspergillus* species section *Flavi* to which *A. oryzae* belongs

(**Supplementary figure S7 and Figure 2A**).

We then analyzed expression of the putative deoxyphomenone biosynthetic genes during

mycoparasitism of *H. pulvinata* 414-3 *in vitro* using quantitative real-time PCR. *H. pulvinata*

never parasitize *C. fulvum* on the nutrient-rich PDA, but it did so on the nutrient-poor water

agar (**Figure 2A**). The expression of the *HpDPH* genes in *H. pulvinata* did not differ when

grown on PDA medium or on water agar (**Figure 3**). In contrast, the expression of *HpDPH*

genes was significantly upregulated when *H. pulvinata* parasitized *C. fulvum* on water agar.

Expression of *HpDPH1*, *HpDPH2* and *HpDPH3* genes, which encode sesquiterpene cyclase,

short-chain dehydrogenase, and cytochrome P450, respectively, were significantly induced

during mycoparasitism (**Figure 3**). Expression of DIP_001952 (lipase/thioesterase family

protein) and DIP_001953 (integral membrane protein) genes also increased, suggesting that

these proteins are associated with deoxyphomenone biosynthesis or mycoparasitism.

Deoxyphomenone is the main antifungal compound of *H. pulvinata* effective against *C.

*fulvum***

Since the *DPH* gene cluster was also found in the genus *Aspergillus*, we assayed for the

presence of deoxyphomenone in the MM culture filtrate from 22 strains of *A. oryzae* and 12

strains of *A. flavus*. In contrast to *H. pulvinata* 414-3, which produced deoxyphomenone as

high as about 80 μM, *A. oryzae* and *A. flavus* strains produced only low levels of

deoxyphomenone: the highest concentrations in *A. oryzae* and *A. flavus* strains are about 0.8

μM and about 3.5 μM, respectively (**Supplementary figure S8**). We then measured

deoxyphomenone by the most productive strains of *A. oryzae* (RIB40) and *A. flavus*

(NBRC114564) in different culturing conditions and found that both strains produced the

compound when grown as still cultures in MM broth (**Supplementary figure S9**).

Interestingly, neither strain produced deoxyphomenone at 35 °C, the optimum temperature

for vegetative growth and sporulation, but showed maximal production at 25 °C, at which

vegetative growth and sporulation are retarded. Similarly, *H. pulvinata* showed the highest

production of deoxyphomenone at 28 °C, a temperature at which vegetative growth is also

retarded (**Supplementary figure S9**).

To determine the function of *HpDPH1* in deoxyphomenone biosynthesis in *H. pulvinata*

414-3, we generated Δ*HpDPH1* knockout mutant strains of 414-3 by homologous

recombination through *Agrobacterium tumefaciens*-mediated transformation

(**Supplementary figure S10**). Quantitative analysis by LC-MS/MS revealed the presence of

deoxyphomenone in the culture filtrate of wild-type 414-3 strain, but none in that of the

ΔHpDPH1 mutant strain KO10 (**Figure 4A**). Strain 414-3 started to produce

deoxyphomenone at 3 d and increased until 15 d, whereas strain KO10 did not produce any

(**Figure 4B**). The culture filtrate of 14-day-old cultures of 414-3 containing deoxyphomenone

strongly inhibited the germination of *C. fulvum* spores, whereas those of the Δ*HpDPH1*

mutant strains KO10 and KO37 showed no inhibitory activity, suggesting that

deoxyphomenone is the main compound contributing to the antifungal activity of *H.*

pulvinata (**Figure 4C**). The antifungal activity was partially restored in the loss-of-function

Δ*HpDPH1* mutant strain (CO10A) after introduction of a functional *AoDPH1* gene (**Figure**

4C). This is most likely explained by restoration of deoxyphomenone production that

accumulated in the culture filtrate of CO10A. The Δ*HpDPH1* mutant KO10 parasitized *C.*

fulvum similar to the wild-type 414-3 strain, with no differences in spore number or

vegetative growth, as we expected previously (Iida et al., 2018) (**Supplementary figure**

S11).

The function of deoxyphomenone differs between *H. pulvinata* and the *Aspergillus*

species

We also generated knockout strains of the homologous gene *AfDPH1* in *A. flavus*

NBRC114564, which produced more deoxyhomenone than *A. oryze* RIB40 (**Supplementary**

figure S8). LC-MS/MS analysis showed that wild-type strain and ectopic transformants

produced deoxyphomenone, while the Δ*AfDPH1* mutant strains produced none (**Figure 5A**).

As mentioned earlier, deoxyphomenone (sporogen-AO 1) was initially identified as a

sporogenic factor in *A. oryzae* (Tanaka et al., 1984b, 1984a). Three Δ*AfDPH1* mutant strains

grown on MM agar produced significantly less spores than the wild-type NBRC114564

(**Figure 5B, C**). Two ectopic transformants produced similar quantities of spores as the wild-

type. Conversely, a Δ*AoDPH1* mutant of *A. oryzae* RIB40 completely lost the ability to

produce deoxyphomenone, but its spore production and colony morphology were similar to

those of the wild-type and ectopic transformants (**Supplementary figure S12**).

We then studied the phenotype of the fungal strains after exposure to deoxyphomenone

on MM. Since *H. pulvinata* 414-3 yielded about 30–100 μM deoxyphomenone during

vegetative growth, we confirmed that deoxyphomenone was fungistatic against *C. fulvum*

over a wide range of concentrations; inhibition by deoxyphomenone was concentration-

dependent; inhibition of spore germination of *C. fulvum* was around 25% at 10 μM, reaching

about 90% at 80 μM (**Figure 6A**). Thus, the concentration of deoxyphomenone secreted by

strain 414-3 seems sufficient to inhibit *C. fulvum* spore germination. As Tanaka et al. (1984b)

treated *A. oryzae* strains with higher concentrations of deoxyphomenone (sporogen-AO 1) to

test sporogenic activity, we also tested concentrations up to 120 μM against *A. oryzae* RIB40

and *A. flavus* NBRC114564. In *Aspergillus*, deoxyphomenone treatment also affected

sporulation in a concentration-dependent manner, but the effect varied among different

species: compared to no treatment, treatment with 120 μM deoxyphomenone reduced spore

production by RIB40, but increased it in NBRC114564 by ~90% (**Figure 6B**). While

deoxyphomenone did not affect colony growth or sporulation of *H. pulvinata* strain 414-3,

colony morphology and formation of aerial hyphae were altered in *Aspergillus* strains, and

colony growth was also slightly inhibited in RIB40, indicating that RIB40 is less tolerant to

this sesquiterpene than 414-3 (**Figure 6B and Supplementary figure S13**). These results

show that deoxyphomenone has quite different activities in different fungi; it has exogenic

antifungal activity against the host fungus and affects endogenic sporogenesis in *Aspergillus*

species.

Deoxyphomenone biosynthetic gene cluster was horizontally transferred from an

Aspergillus* ancestor to *H. pulvinata

Next, we analyzed the commonality and conservation of the *DPH* clusters in genomic

sequences of *Aspergillus* species. Complete *DPH* clusters were detected only in *Aspergillus*

species from the section *Flavi* (Group I, **Figure 2A**), and the protein sequences of HpDph2 to

HpDph6 in these species were highly conserved, with ~70% pairwise similarity to those of *H.*

pulvinata (**Figure 7**). Meanwhile, partial *DPH* clusters were present in *Aspergillus* species of

the section *Circumdati* (Group II: *HpDPH1, HpDPH2, HpDPH3*) and the sections *Nidulantes*

and *Clavati* (Group III: *HpDPH1, HpDPH2, HpDPH4*) (**Figure 7**). Group II species also

contained a lipase/thioesterase family protein- and an integral membrane protein-encoding

genes downstream of *HpDPH1*, the same as in Group I (**Figure 7 and Supplementary**

figure S7). Group III further lacked the commonality and conservation of the *DPH* clusters:

the similarities with *HpDPH1* and *HpDPH2* genes were much lower than in the other two

groups, the position and the order of *DPH2* and *DPH4* genes in the genome sequence were

quite different, and unrelated genes were interspersed in the cluster (**Figure 7** and

Supplementary figure S7). These results suggested that Group II and III species lost parts of

the *DPH* cluster and no longer produced deoxyphomenone.

     To infer the evolutionary origin of the *DPH* clusters in *H. pulvinata* and *Aspergillus*, we

analyzed the evolutionary divergence of the clusters among 60 fungal genomes using

genomic and phylogenetic approaches with the NOTUNG algorithm, which reconciles

differences between species trees and gene trees by inferring gene duplications, transfers, and

losses (Stolzer et al., 2012). Reconciliation of the species tree with the HpDph1 protein tree

detected a transfer event (T1 in **Figure 8**) early in *Aspergillus* section *Circumdati*, indicating

a transfer of *HpDPH1* of *H. pulvinata* from Group II (*Circumdati*) (**Supplementary figure**

S14). Likewise, the HpDph2 and HpDph4 protein trees indicated transfers from Group I

(section *Flavi*) to *H. pulvinata* (T2 in **Figure 8**). Since the *DPH* clusters are highly conserved

in *H. pulvinata* and *Aspergillus*, these results suggest an evolutionary scenario in which the

DPH cluster was transferred from the common ancestor of Groups I and II (sections *Flavi*

and *Circumdati*) to *H. pulvinata* at the same time and that genes were lost in time in

Aspergillus species (**Figure 8**). Since all duplication events detected in *DPH* genes occurred

after the horizontal transfer events (**Supplementary figure S14**), it is unlikely that the

duplication is associated with the transfer of the *DPH* cluster. Altogether these results suggest

that the *DPH* cluster arose before speciation within the *Aspergillus* genus. Finally, because

we inferred the horizontal transfer of the *DPH* cluster from *Aspergillus* to *H. pulvinata*, we

investigated the association of deoxyphomenone with the mycoparasitism. Contrary to our

expectations, *H. pulvinata* 414-3 did not directly penetrate the hyphae of *A. oryzae* RIB40, as

observed with the host fungus *C. fulvum*, but physical contact such as adhesion to or coiling

around the hyphae of the RIB40 was detected (**Supplementary figure S15**).

DISCUSSION

Strict regulation of expression through the physical linkage of genes involved in the same

biosynthetic pathway leads to physiological economization; thus, biosynthetic genes for

secondary metabolites in fungi are often maintained in clusters (Hagee et al., 2020), which

allows for the rapid conversion of potentially toxic or chemically unstable intermediate

products (McGary et al., 2013). In the present study, the *DPH* clusters were highly conserved

between *H. pulvinata* and *Aspergillus* section *Flavi*, suggesting that the final product,

deoxyphomenone, plays critical roles in distinct stages of the life cycle of these fungi. Indeed,

it acts as an antifungal agent of the mycoparasite *H. pulvinata* against the host fungus *C.*

fulvum and controls sporulation at least in *A. flavus*. The correlation of these fungal clusters

can be explained through the most parsimonious evolutionary scenario, which is constructed

by reconciling the species tree with the *DPH* protein tree in consideration of the costs of

evolutionary events (i.e., gene duplications, transfers, and losses) (Stolzer et al., 2012).

Interestingly, we inferred that the *DPH* cluster was horizontally transferred across a large

taxonomic distance from the genus *Aspergillus* (*Eurotiomycetes*) to *H. pulvinata*

(*Sordariomycetes*), suggesting that *H. pulvinata* maintained the cluster as deoxyphomenone

production seemed to be more conductive to mycoparasitism. Horizontal gene transfer is not

rare in fungi (Mehrabi et al. 2011; Rokas et al., 2018); gene clusters for the biosynthesis of

products such as fumonisin (Proctor et al., 2013, Khaldi and Wolfe, 2011), host-specific

toxins in *Alternaria* species (Tsuge et al., 2016), and cercosporin (de Jonge et al., 2018) have

been horizontally transferred across classes, but the final products function in the recipient

fungi basically in a similar way.

We deduced the deoxyphomenone biosynthetic pathway from the component genes of

the *DPH* cluster in *H. pulvinata* and *Aspergillus*. The bicyclic eremophilane-type

sesquiterpene deoxyphomenone is probably formed by the cyclization of farnesyl

pyrophosphate catalyzed by sesquiterpene cyclase (Proctor and Hohn, 1993), followed by

oxygenation and epoxidation of the double bond in *H. pulvinata* and *Aspergillus*. Although

deoxyphomenone has also been detected in *Penicillium* species, which are taxonomically

close to *Aspergillus*, the *DPH* cluster was not detected in the published genome sequences of

Penicillium species; thus, the pathway of deoxyphomenone biosynthesis is likely different

between the two genera. On the basis of a relaxed molecular clock analysis, *Aspergillus* and

Penicillium likely diverged 94.0 million years ago (mya) with *Aspergillus* sections *Flavi* and

Circumdati originating 51.1 mya (Barelli et al., 2015; Steenwyk et al., 2019). Because the

DPH clusters are fully or partially conserved in *Aspergillus* sections *Flavi* and *Circumdati,*

we presume that their horizontal transfer to *H. pulvinata* occurred at least before the common

ancestor of these sections.

It is still unclear how *H. pulvinata* acquired the *DPH* clusters from the ancestral

Aspergillus. Since horizontal transfer of foreign DNA first requires contact between fungal

cells, the involvement of anastomosis between conidia, germ tubes or hyphae, and subsequent

heterokaryon formation, have been proposed (Kubicek et al., 2019; Fleißner et al., 2022;

Bucknell and McDonald, 2023). *H. pulvinata* has a wide host range, including plant

pathogens (Peresse and Lepicard, 1980; Mitchell and Taber, 1986; Mitchell et al., 1986;

Mitchell et al., 1987; Mello et al., 2008; Alderman et al., 2010), and plant pathogens in

Aspergillus sections *Flavi* and *Circumdati* may provide opportunities for contact with *H.*

pulvinata on plants and subsequent gene transfer. Given the conservation of *DPH* clusters

revealed in this study and the nature of the recipient fungus, we propose to add

mycoparasitism as one of the mechanisms that facilitated horizontal gene transfer between

fungi. Mycoparasitic *Trichoderma* species have been suggested to have acquired by

horizontal transfer nearly half its genes encoding plant cell wall-degrading carbohydrate-

active enzymes and auxiliary proteins enabling them to be parasitism on a broad range of

Ascomycota (Druzhinina et al., 2018). *H. pulvinata* 414-3 did not establish a direct parasitic

relationship with *A. oryzae* RIB40 as it did with *C. fulvum*, and no distinct anastomosis was

observed, but it firmly adhered to and coiled around RIB40 hyphae, indicating basic

compatibility facilitating horizontal between the two fungi. Since *H. pulvinata* completely

kills the host *C. fulvum*, *C. fulvum* might not be a stable source of foreign DNA, but rather *H.*

pulvinata might have acquired the gene cluster in direct contact with nonhost fungi. In the

genomic sequence of *H. pulvinata*, a gene cluster derived from *Aspergillus* was found in this

study, but no horizontal gene transfer between host fungi, including *C. fulvum*, has been

found so far. Although transduction by mycoviruses or propagation of retrotransposons could

also have facilitated horizontal gene transfer (Richards et al., 2006), no transposon-like

sequences were detected around the *DPH* cluster.

Several environmental cues such as temperature, light, nutrients, pH, and competing or synergistic organisms influence fungal activity, including transcriptional regulation of secondary metabolism–associated gene clusters (Keller, 2019). Aflatoxin production in *A. flavus* is generally upregulated at a suboptimal 30 °C (Ching'anda et al., 2021). Similarly, we found that deoxyphomenone production was also highest at temperatures unsuitable for growth of *H. pulvinata* and *Aspergillus* species (**Supplementary figure S8**). In *A. flavus*, Δ*AfDPH1* mutants that no longer produced deoxyphomenone formed significantly less spores. These results suggest that in at least *A. flavus*, deoxyphomenone stimulates sporulation in poor nutritional environments. In addition, *A. flavus* strains tended to produce more deoxyphomenone than *A. oryzae* strains, but the level was not sufficiently high for antifungal activity. Gibbons et al. (2012) reported that the expression levels of many secondary metabolite gene clusters of *A. oryzae*, including those for *DPH* and aflatoxin, were repressed compared with those of *A. flavus*. *A. oryzae*, used to ferment rice to produce Japanese sake, is probably derived from *A. flavus*, a pathogen toxic to plants and animals (Gibbons et al., 2012). *A. oryzae* is cultured with the budding yeast *Saccharomyces cerevisiae* during sake brewing and might have lost the ability to produce aflatoxin during its domestication from *A. flavus* (Watarai et al., 2019) because aflatoxin is genotoxic to yeast (Keller-Seitz et al., 2004). Since *A. oryzae* is usually grown under optimal fermentation conditions, the gene encoding deoxyphomenone required for sporulation by *A. flavus* in unfavorable conditions may also have been silenced or lost in *A. oryzae*. In fact, external application of deoxyphomenone inhibited sporulation and colony formation only in *A. oryzae*, indicating that *A. oryzae*, which produces only low levels of the compound, is less tolerant to this sesquiterpene (**Supplementary figure S13**). The mycoparasite *H. pulvinata* also produced deoxyphomenone at temperatures unfavorable for its growth, but in the presence of its host fungus *C. fulvum*, expression of the *HpDPH* genes was strongly induced,

suggesting that deoxyphomenone might be crucial for mycoparasitic activity of *H. pulvinata*.

However, the *ΔHpDPH1* mutant strains of *H. pulvinata* parasitized *C. fulvum*, comparable to

the wild-type strain, supporting that deoxyphomenone is not required directly for its

mycoparasitism (Iida et al., 2018). Spore germination of *C. fulvum* treated with

deoxyhomenone for 24 h was halved, also suggesting an epidemiological effect on the

mycoparasitism of *H. pulvinata*.

Deoxyphomenone did not affect the organelles in *C. fulvum* spores, and the specific

target is still unknown. Although it is not directly involved in the mycoparasitism of *H.*

pulvinata and is not toxic to plants, it had significant fungistatic activity against plant

pathogens—the host *C. fulvum* and the nonhost *P. fuligena* of *H. pulvinata*—which might

indicate a potential effect on biotrophic fungal pathogens of tomato plants. *H. pulvinata*

remains an efficient biocontrol agent that can be used as a biofungicide to suppress races and

fungicide-tolerant strains of *C. fulvum* that cause problems in tomato crops world-wide (Iida

et al., 2015; Mesarich et al., 2023). Among *Aspergillus* species commonly used in

fermentation, the use of deoxyphomenone to control sporulation should promote high

efficiency for industrial fermentation and enzyme production.

MATERIALS AND METHODS

Growth conditions for fungal strains and plants

Strains of *H. pulvinata* 414-3 and *C. fulvum* CF301 were grown on PDA (half-strength, BD

Difco, Franklin Lakes, NJ, USA), MM agar or broth (15 g sucrose, 5 g ammonium tartrate, 1

g NH_4NO_3, 1 g KH_2PO_4, 0.5 g $MgSO_4 \cdot 7H_2O$, 0.1 g NaCl, 0.1 g $CaCl_2 \cdot H_2O$, 25 μL 0.2 mg/mL

biotin, 15 g agar and 1 mL trace elements per liter) at 25 °C in the dark for 1 and 2 weeks,

respectively, then spores were collected in sterile distilled water. *A. oryzae* and *A. flavus*

strains (**Supplementary figure S7**) from NITE-NBRC (www.nite.go.jp/nbrc/catalogue/)

were cultured on PDA, MM agar and broth, and Czapek-Dox agar and broth (3 g NaNO$_3$, 2 g

KCl, 1 g KH$_2$PO$_4$, 0.5 g MgSO$_4$·7H$_2$O 20 g glucose and 15 g agar per liter adjusted to pH 6.5)

at 30 °C in the dark for a week, and spores were collected in sterile distilled water containing

1% (v/v) Tween 20. The concentration of spores was determined using a hemocytometer.

Strains were maintained on PDA at 25 °C for a few months. Spore suspensions containing

10% v/v glycerol of all fungi except *H. pulvinata* were preserved at –80 °C for long-term

storage; *H. pulvinata* was stored in 10% v/v glycerol with 5% w/v trehalose.

For examining mycoparasitic interactions on tomato leaves, tomato cultivar

Moneymaker, which lacks any apparent *Cf* resistance gene, was grown in plastic pots for 3

weeks in a climate chamber at 25 °C with 16 h light/8 h dark. Seedlings were transplanted to

soil in the greenhouse and grown for 3 weeks. The lower sides of tomato leaves were sprayed

with a spore suspension of *C. fulvum* CF301 (1 × 10^5 spores/mL). After 2 weeks, a spore

suspension of *H. pulvinata* 414-3 (1 × 10^5 spores/mL) was sprayed on the brown lesions that

had formed on the abaxial surfaces of the leaves. White mycelial patches of *H. pulvinata*

were observed a week after inoculation.

Electron microscopy

C. fulvum CF301 or *A. oryzae* RIB40 were cultured on a nylon membrane (0.45 μm;

GVS Japan, Tokyo, Japan) placed on PDA in a petri dish and incubated as described above.

The membranes were then transferred to water agar and incubated for a week in the same

conditions. A spore suspension of *H. pulvinata* (1 × 10^5 spores/mL) was sprayed on the test

colony on the membrane, then white colonies were observed 1 week later.

Small pieces (1 × 1 cm^2) were cut from the membrane and fixed in 2.5% (v/v)

glutaraldehyde in 100 mM cacodylate buffer (pH 7.4) overnight at 4 °C, then washed three

times in the buffer, fixed in 1% (w/v) osmium tetroxide in the buffer for 1 h at room

temperature and washed three times in distilled water. Specimens were then dehydrated

through a graded ethanol series, critical-point dried in absolute ethanol using Leica EM

CPD300 critical point dryer (Leica Microsystems, Wetzlar, Germany), then mounted on

stubs, coated with platinum-palladium, and observed with a Hitachi SU-8220 scanning

electron microscope (Tokyo, Japan).

For observing the ultrastructure of *C. fulvum* spores with transmission electron

microscopy (TEM), spores of strain CF301 were treated with deoxyphomenone (120 µM) at

25 °C for 24 h in the dark. Spores were collected by centrifugation, washed three times with

sterile water, then fixed in phosphate-buffered 2% glutaraldehyde at 4 °C overnight. The

samples were again collected by centrifugation, then treated with 2% potassium

permanganate in phosphate buffer for 1 h at room temperature and post-fixed in 2% w/v

osmium tetroxide in phosphate buffer for 2 h at 4 °C. The samples were dehydrated in a

graded ethanol series and embedded in epoxy resin (Fujifilm Wako, Osaka, Japan) at 60 °C

for 48 hours. Ultrathin sections were cut using an ultramicrotome Leica UCT Ultracut (Leica

Microsystems), then stained with uranyl acetate for 15 min and lead solution for 5 min and

observed with a Hitachi TEM H-7600.

Assays of antifungal activity and plant toxicity of deoxyphomenone

C. fulvum spores were treated with deoxyphomenone [(+)-13-deoxyphomenone, sporogen-

AO1; Apollo Scientific, Bredbury, UK] in 1% methanol (v/v) at 25 °C in the dark for 24 h.

Then samples were observed for germination and hyphal elongation using a Nikon E600 light

microscope (Tokyo, Japan).

The antifungal activity of deoxyphomenone was compared with that of an *N*-halo-

alkylthioimide fungicide, captan (orthocide80; Arysta Life Science, Tokyo, Japan). *C. fulvum*

spores (1×10^3 cells/mL) were treated with distilled water, deoxyphomenone (120 µM), or

captan (100 µM) on a hydrophobic TF0808 glass slide (Matsunami Glass, Osaka, Japan) for

24 h at 100% humidity, then each solution was replaced with fresh solution, and samples

incubated another 24 h. Germinated spores were then counted using the Nikon light

microscope. For the controls, dH_2O or 1% methanol (v/v) were used. Each treatment was

done in triplicate, and at least 200 spores and hypha were assessed for each treatment.

Spore suspensions of the strains *C. fulvum* CF301, *P. fuligena* Pf17923, kindly

provided by K. Nakajima and T. Kawakami, and *F. oxysporum* f. sp. *lycopersici* CK3-1 were

adjusted to 1×10^6 spores/mL and 10 µL were dropped on MM agar supplemented with

different concentrations of deoxyphomenone (20 to 120 µM). The strains were cultured at

25 °C in the dark for 2 weeks or 3 days.

To evaluate the toxicity of deoxyphomenone on tomato plants, we injected the abaxial

surface of three leaves on each of two 1-month-old tomato plants with about 20 µL of

deoxyphomenone (20 to 80 µM) or with 1% methanol (v/v) as a control. The plants were then

incubated in a climate chamber for 1 week, then leaves were examined for necrosis.

Phylogenetic analyses of deoxyphomenone biosynthetic gene clusters

Deoxyphomenone biosynthetic genes and *DPH* clusters were initially identified in the

genomic sequences of *H. pulvinata* 414-3 (Sushida et al., 2019) and *A. oryzae* RIB40

(Machida et al., 2005) using the BLAST program (blast.ncbi.nlm.nih.gov). Sequences shown

in **Supplementary table S1 and S2** were aligned using MAFFT online version 7 (Katoh and

Standley, 2013) with default parameters and trimmed using TrimAl version 1.4 (Capella-

Gutiérrez et al., 2009). Maximum likelihood phylogenetic trees were constructed using

RAxML version 8.2.12 with 1000 bootstraps; the PROTCATAUTO and GTRCAT models

were selected for amino acid and nucleotide sequences, respectively (Stamatakis, 2014). Dot

plot alignments were generated from the MAFFT analysis (**Supplementary figure S4**).

Pairwise sequence alignments of *DPH6* homologous genes were performed using EMBOSS

Needle (Madeira et al. 2022).

The phylogenetic species tree in **Figure 8** was constructed using maximum likelihood

based on the 500 monocore genes (a single homolog in each of the species) using CVtree

version 3.0 ($k = 16$) (Qi et al., 2004) by uploading the genomic information for the 60

published fungi shown in **Supplementary table S1**.

Reconciliation analysis between species and gene trees was performed using NOTUNG

v.2.9 (Stolzer et al., 2012) according to the instructions (version 2.8 beta) to infer the

evolutionary trajectories of the *DPH* clusters. Event scores were calculated as the total costs

of duplications, transfers and losses. Costs/weights were set as duplications (D), 1.5; transfers

(T), 8.0; losses (L), 1.0 (ratio D:T:L is 1:5.3:0.67) for amino acid sequences, and D, 1.5; T,

6.0; L, 1.0 (ratio D:T:L is 1:4:0.67) for nucleotide sequences for the gene clusters.

Quantitative detection of deoxyphomenone

H. pulvinata (1×10^6 spores) was cultured in 50 mL of MM broth with shaking at 25 °C in a

light/dark cycle of 16 h/8 h for 2 weeks. *A. oryzae* and *A. flavus* (1×10^6 spores) were

incubated in MM broth without shaking at 25 °C in the dark for 2 weeks. The culture

supernatants were collected by centrifugation and filtered through a 0.45 µm filter (Merk,

Darmstadt, Germany). Deoxyphomenone was quantified using a 4000 QTRAP LC-MS/MS

system (Sciex, MA, US) equipped with a 1290 series HPLC system (Agilent Technologies,

CA, US) as previously reported (Iida et al., 2018). Commercial deoxyphomenone (Apollo

Scientific) was used as a standard.

Construction of transformation vectors

Genomic DNA was extracted from *H. pulvinata* and *A. oryzae* mycelia grown in MM broth

for a week using Nucleo-Mag Plant (TakaraBio, Shiga, Japan) according to the

manufacturer's instructions. The DNA was then amplified by PCR using PrimeSTAR GXL

Premix (TakaraBio) or KOD ONE (TOYOBO, Osaka, Japan) according to the instructions.

For generating the gene knockout vector, the upstream (1.2 kb) and downstream (1.0 kb)

regions of *HpDPH1* were amplified from *H. pulvinata* genomic DNA using primer sets

Dpprx2LF1/Dpprx2LR1 and Dpprx2UF1/Dpprx2UR1, respectively (**Supplementary table**

S3). Geneticin-resistance gene (*gen*) cassettes were amplified from pRM254 (Mehrabi et al.,

2015) using primers pRM254 attL1_F/pRM254 attL1_R. These primers contain a 5′

overhang sequence to overlap the sequences at the ends of the linearized plasmid or the *gen*

cassette (**Supplementary table S3**). Plasmid pPM43GW was amplified using the primers

pPM43GW_RB_F and pPM43GW_RB_R to linearize and remove the *ccdB* gene, which

encodes a product that is lethal to bacterial cells. Four fragments were combined to generate

the gene replacement vector pGW43_dpprx2ko using the In-Fusion EcoDry Cloning Kit

(TakaraBio).

To introduce the *AoDPH1* gene into Δ*HpDPH1* strain KO10, *AoDPH1* (without introns)

and hygromycin resistance gene (*hph*) cassettes were synthesized by VectorBuilder, Inc.

(Chicago, IL, USA). The *AoDPH1* and *hph* genes were driven under the promoters of the *A.*

nidulans trpC gene and *Cochliobolus heterostrophus gpd1* (NCBI accession X02390.1 and

X63516.1), respectively (**Supplementary figure S9A**). Synthesized DNA was inserted into

the blunt end of the PvuII-digested plasmid pPZP-PvuII kindly provided by Prof. Chihiro

Tanaka.

To construct gene replacement vector for *AoDPH1* and *AfDPH1* gene, a plasmid pSH75

(Kimura and Tsuge 1993) was digested with *Bgl* II and *Hin*d III, and the resulting 2.4 kb

fragment was used as a backbone. Marker gene cassettes (4.7 kb) containing a pyrithiamine

resistant gene (*ptrA*) and an *EGFP* gene cassette were amplified from a plasmid

pPTREXeGFP using the primers pPTREXeGFP_F and pPTREXeGFP_R. The upstream (1.5

kb) and downstream (1.5 kb) regions of *AoDPH1* were amplified from genomic DNA of *A.*

oryzae RIB40 using primer sets AoDPH1_LB_F/AoDPH1_LB_R and

AoDPH1_RB_F/AoDPH1_RB_R, respectively (**Supplemental table S3**). These primers

contain a 5′ overhang sequence to overlap the sequences at the ends of the pSH75 fragment

or the marker cassette (**Supplementary table S3**). These four fragments were combined

using the In-Fusion EcoDry Cloning Kit (TakaraBio), then the resulting plasmid was named

pSH75PTRdAoDPH1. The plasmid was linearized by digestion with *Swa* I and used for

fungal transformation.

Plasmid vectors were introduced into *Escherichia coli* DH5α (NIPPON GENE, Tokyo,

Japan) and extracted using MagExtractor -Plasmid- (TOYOBO) according to the instructions.

The correct orientation of the fragments in the final constructs was confirmed by PCR.

Fungal transformation

All transformations were done as previously described (Okmen et al., 2013) with some

modifications for *H. pulvinata*. Briefly, *A. tumefaciens* (*Rhizobium radiobacter*) strain AGL-

1 was transformed with the plasmid vectors used for the *H. pulvinata* transformation using

the Gene Pulser electroporator (Bio-Rad Laboratories, Hercules, CA, USA) according to the

instructions, then grown on LBman agar (10 g tryptone, 10 g NaCl, 5 g yeast extract, 10 g

mannitol, 20 g agar per liter) supplemented with appropriate antibiotic at 28 °C for 3 days.

Colonies were then collected and resuspended in IM broth (1 g glucose, 2.05 g K_2HPO_4, 1.45

g KH_2PO_4, 0.15 g NaCl, 0.50 g $MgSO_4 \cdot 7H_2O$, 0.07 g $CaCl_2 \cdot 2H_2O$, 0.5 g $(NH_4)_2SO_4$, 0.5%

(w/v) glycerol, 8.53 g 2-(*N*-morpholino)ethanesulfonic acid [pH 5.3] per liter) supplemented

with 200 μM acetosyringone at an optical density at 600 nm (OD_{600}) of 0.2. The bacterial

suspension (200 μL) was mixed with 200 μL of a spore suspension of *H. pulvinata* (1.0×10^5

cells/mL), then the suspension was spread on a nylon membrane (GVS Japan) that had been

placed on IM agar (IM; 20 g per liter) in a plate. After 2 days of incubation in the dark at

25 °C, the membranes were transferred to PDA supplemented with either 200 μg/mL

geneticin (G418 sulfate; Fujifilm Wako) or 100 μg/mL hygromycin (Fujifilm Wako), and 50

μg/mL meropenem trihydrate (Fujifilm Wako). Transformed fungal colonies appeared after a

week were transferred to new PDA plates supplemented with geneticin (Fujifilm Wako) or

hygromycin (Fujifilm Wako) and meropenem trihydrate (Fujifilm Wako) at the same

concentration as described above and incubated in the dark at 25 °C for a week. Gene

replacement was confirmed by PCR amplification using the primers listed in **Supplementary**

table S3.

Protoplasts of *A. oryzae* RIB40 and *A. flavus* NBRC114564 were transformed using

polyethylene glycol. First, 2.4×10^8 spores harvested from a 10-d-old culture on Czapek-Dox

agar were resuspended in 200 mL of Czapek-Dox broth and incubated at 30 °C at 160 rpm

for 18 h. The mycelium was then collected using miracloth (Merck, NJ, USA) and washed

once with distilled water, then incubated in 20 mL of a protoplast solution consisted of 10

mM phosphate buffer (pH6.0), 0.8 M NaCl, 10 mg/mL Lysing enzyme (Sigma-Aldrich, MO,

USA), 5 mg/mL Cellulase Onozuka R-10 (Yakult Pharmaceutical, Tokyo, Japan), and 2.5

mg/mL Yatalase (Takara, Shiga, Japan) at 30 °C with shaking at 83 rpm for 3 h. The

protoplasts were filtered through a mesh and centrifuged at $2000 \times g$ and 4 °C for 5 min. The

pellet was washed once in 0.8 M NaCl and centrifuged at $2000 \times g$ and 20 °C for 5 min, then

resuspended in 1.2 mL of solution 1 (Sol1: 9.35 g NaCl, 2 mL 1 M CaCl$_2$, 2 mL 1 M Tris-

HCl per 200 mL). Two handled fourty microliters of solution 2 (Sol2: 40% (w/v) PEG4000,

10 mL 1 M CaCl$_2$, 10 mL 1 M Tris-HCl per 200 mL) was added to the protoplast suspension

and the resulting mixture was dispensed 300 μL each into 14-mL round tubes A linearized

vector DNA was added to the mixture and held on ice for 30 min, then 1 mL of Sol2 was

added and held at 25 °C for 20 min. The protoplasts were washed once in 10 mL of Sol1, and

centrifuged at 2000 × g at 20 °C for 5 min. After the supernatant was discarded, the

protoplasts were resuspended in 300 μL of Sol1, then placed on a regeneration selection

medium consisted of Czapek-Dox agar supplemented with 5% (w/v) NaCl, 0.1 μg/mL

pyrithiamine hydrobromide (Sigma-Aldrich, MO, USA). The resulting plates were overlayed

with 5 mL of soft agar medium consisted of 5% (w/v) NaCl and 0.5% (w/v) agar, and

incubated at 30 °C for 4 d. The transformants grown on the regeneration selection medium

were isolated and transferred to Czapek-Dox agar containing 0.2 μg/mL pyrithiamine

hydrobromide and incubated at 30 °C for another selection. Gene replacement of the

transformants was confirmed by PCR amplification using the primers listed in

Supplementary table S3.

Quantitative real-time PCR

C. fulvum CF301 and *H. pulvinata* 414-3 were cultured on nylon membranes (GVS Japan) on

PDA at 25 °C in the dark for 1 and 2 weeks. The membranes were transferred to water agar

and then incubated in the same conditions for a week. *C. fulvum* colonies were sprayed with a

spore suspension of *H. pulvinata* (1×10^5 spores/mL) and checked for white colony growth

of *H. pulvinata* 1 week later. Total RNA was extracted from the mycelia using the RNeasy

Plant Mini Kit (Qiagen, Hilden, Germany). cDNA libraries were generated using SuperScript

IV VILO Master Mix with ezDNase Enzyme (ThermoFisher Scientific, Waltham, MA, USA)

according to the manufacturer's instructions. Primers for quantitative real-time PCR of 90–

150-bp fragments of cDNA were designed (**Supplementary table S3**), and the PCR was run

on the LightCycler 480 System (Roche, Basel, Switzerland) using the KAPA SYBR Fast

qPCR Kit (Nippon Genetics, Tokyo, Japan) according to the manufacturer's instructions.

Relative expression levels were calculated using the comparative CT ($2^{-\Delta\Delta CT}$) method (Livak

and Schmittgen, 2001). The data were normalized to the transcript level of the *H. pulvinata*

actin gene (**Supplementary table S3**). Transcript levels of target genes in each RNA sample

were measured for three independent experiments, each with two replicates.

In-well assay and observation of mycoparasitism

Green fluorescence protein (GFP)-expressing *C. fulvum* strain CF301gfp (Iida et al., 2018)

and *H. pulvinata* strains were cultured on PDA as mentioned above to prepare spore

suspensions of CF301gfp (5×10^4 spores/mL) and *H. pulvinata* (1×10^6 spores/mL). Then 50

µL of a spore suspension of CF301gfp and of each *H. pulvinata* strain were added to 100 µL

of MM broth without a carbon source (MM without sucrose) in each well of a 48-well plate.

At least 8 wells were used per treatment, and CF301gfp was used as the control. The plates

were incubated at 25 °C in the dark for 3 d. Hyphae were observed using a fluorescence

microscope BZ-X800 (KEYENCE, Osaka, Japan); lack of GFP fluorescence from *C. fulvum*

was correlated with the mycoparasitic activity of *H. pulvinata* using the hybrid cell count

system of the microscope. We named this simple method for detecting mycoparasitism the

IWAO method, derived from "in-well assay and observation" and in honor of the method's

developer, E. Iwao.

Statistical analyses

Means and standard deviations of number of spores, spore germination, and hyphal length

were calculated. Significant differences ($P < 0.05$) in the number of spores produced by the

different fungal strains, percentage spore germination and hyphal lengths of *C. fulvum*, and

colony diameter of *Aspergillus* strains were evaluated using either Tukey's test or Williams'

multiple comparison. Significant differences ($P < 0.05$) in gene expression levels were

determined using Welch's *t*-test followed by Bonferroni–Holm correction for multiple

testing. All data were analyzed in the program R version 4.0.3 (www.r-project.org).

Data availability

The genomic sequence of *H. pulvinata* 414-3 is available in the DDBJ repository under an

accession number BJKQ00000000 (BioProject: PRJDB8178) (Sushida et al., 2019). Other

fungal sequences shown in **Supplementary table S1** are available on the fungal genome

portal MycoCosm (mycocosm.jgi.doe.gov) of the Joint Genome Institute (Lawrence Berkeley

National Institute, Berkeley, CA, USA; Grigoriev, 2014) and in the DDBJ/EMBL/GenBank

repository (www.ncbi.nlm.nih.gov). Auxiliary data are available through Figshare

(doi.org/10.6084/m9.figshare.24559120.v1).

ACKNOWLEDGEMENTS

We are grateful to Kentaro Ikeda and Hiroshi Sakai for providing *H. pulvinata* strains,

Daisuke Hagiwara for *A. oryzae* RIB40 and plasmid vector pPTREXeGFP, and Kaori

Nakajima and Taku Kawakami for the *P. fuligena* strain, Yasuyuki Kubo and Sayo Kodama

for valuable suggestions, Chieko Tanaka for laboratory management, and Pierre J.G.M. de

Wit for critical reading of the manuscript. This work was supported by a Grant-in-Aid for

Scientific Research from JSPS (17H05022 and 20H02993) (Y. Iida).

Conflict of interest

The authors declare that they have no commercial or financial relationships that could be

construed as a potential conflict of interest.

Author contributions

YI designed the study and managed the research funds. KM, HS and YH performed qRT-

PCR and analyzed the data. KM, TaS and ON transformed fungal strains. ToS did the

electron microscopy and EI did the light microscopy. KM, TaS, YH and HN did the mass

spectrometry analyses. MZF and YN predicted the biosynthetic pathways and identified the

genes for enzymes. KM and HS did the phylogenetic analyses. KM and YI wrote the paper.

All authors contributed to the article and approved the submitted version.

Supplementary figures

Figure S1. Effect of deoxyphomenone on *Cladosporium fulvum* and tomato leaves.

Figure S2. Phylogenetic tree of fungal sesquiterpene cyclases.

Figure S3. Alignment and homology of the amino acid sequences of the predicted

sesquiterpene cyclases identified in *Hansfordia pulvinata* and *Aspergillus oryzae* genome

sequences.

Figure S4. Dot plot analysis of the deoxyphomenone biosynthesis (*DPH*) gene clusters from

Hansfordia pulvinata 414-3 and *Aspergillus oryzae* RIB40.

Figure S5. Exon-intron structures of deoxyphomenone biosynthetic (*DPH*) genes.

Figure S6. Alignment of deduced amino acid sequences of *HpDPH6* homologs from

Hansfordia pulvinata and six *Aspergillus* species.

Figure S7. Phylogenetic analysis of genomic sequences containing *HpDPH1* homologous

genes and the upstream region using the maximum likelihood method.

Figure S8. Deoxyphomenone produced by *Hansfordia pulvinata* 414-3 and strains of

Aspergillus oryzae and *A. flavus*.

Figure S9. Deoxyphomenone production by *Hansfordia pulvinata* 414-3, *Aspergillus oryzae*

RIB40 and *A. flavus* NBRC114564 on MM agar or in broth at different temperatures.

Figure S10. PCR detection of the inserted cassettes in transformants and wild-type strains.

Figure S11. *In vitro* assay of *Hansfordia pulvinata* mycoparasitic activity against GFP-

expressing *Cladosporium fulvum*.

Figure S12. Deoxyphomenone production and sporulation of *A. oryzae* RIB40 and

transformants.

Figure S13. Effect of deoxyphomenone on mycelial growth of *Hansfordia pulvinata,*

Aspergillus oryzae and *A. flavus*.

Figure S14. Reconciliation of gene trees with species tree based on *DPH* nucleotide and amino

acid sequences.

Figure S15. Hyphal contact between the mycoparasite *Hansfordia pulvinata* 414-3 and

Aspergillus oryzae RIB40.

Supplemental table S1. Fungal genome sequences used in this study.

Supplemental table S2. Amino acid sequences of *DPH* biosynthesis genes from *Hansfordia*

pulvinata and *Aspergillus* species.

Supplemental table S3. Primer sequences used in this study.

REFERENCES
723
Alderman, S.C., Rao, S., and Martin, R. (2010) First report of *Dicyma pulvinata* on *Epichloë*
typhina and its potential for control of *E. typhina*. *Plant Health Progress*: PHP-2010-
0216-2001-RS.
Barelli, L., Moonjely, S., Behie, S.W., and Bidochka, M.J. (2015) Fungi with multifunctional
lifestyles: endophytic insect pathogenic fungi. *Plant Molecular Biology* 90: 657-664.
Bucknell, A.H., and McDonald, M.C. (2023) That's no moon, it's a Starship: Giant
transposons driving fungal horizontal gene transfer. *Molecular Microbiology* 120: 555-
563.
Capella-Gutiérrez, S., Silla-Martínez, J.M., and Gabaldón, T. (2009) trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:
1972-1973.
Ching'anda, C., Atehnkeng, J., Bandyopadhyay, R., Callicott, K.A., Orbach, M.J., Mehl,
H.L., and Cotty, P.J. (2021) Temperature Influences on Interactions Among
Aflatoxigenic Species of *Aspergillus* Section *Flavi* During Maize Colonization. *Frontiers*
in Fungal Biology 2: 720276.

Christianson, D.W. (2017) Structural and Chemical Biology of Terpenoid Cyclases. *Chemical Reviews* 117: 11570-11648.

Cooke, M.C. (1883) New American fungi. *Grevillea* 12: 22-33.

Daengrot, C., Rukachaisirikul, V., Tansakul, C., Thongpanchang, T., Phongpaichit, S., Bowornwiriyapan, K., and Sakayaroj, J. (2015) Eremophilane Sesquiterpenes and Diphenyl Thioethers from the Soil Fungus *Penicillium copticola* PSU-RSPG138. *Journal of Natural Products* 78: 615-622.

de Jonge, R., Ebert, M. K., Huitt-Roehl, C. R., Pal, P., Suttle, J. C., Spanner, R. E., Neubauer, J. D., Jurick, W. M., 2nd, Stott, K. A., Secor, G. A., Thomma, B. P. H. J., Van de Peer, Y., Townsend, C. A., & Bolton, M. D. (2018) Gene cluster conservation provides insight into cercosporin biosynthesis and extends production to the genus *Colletotrichum*. *Proceedings of the National Academy of Sciences* 115: E5459-E5466.

de Wit, P.J.G.M. (2016) *Cladosporium fulvum* effectors: weapons in the arms race with tomato. *Annual Review of Phytopathology*

Deighton, F.C. (1972) Synonymy of *Hansfordia pulvinata* (Berk. & Curt.) Hughes. *Transactions of the British Mycological Society* 59: 531-536.

Del Valle, P., Figueroa, M., and Mata, R. (2015) Phytotoxic eremophilane sesquiterpenes from the coprophilous fungus *Penicillium* sp. G1-a14. *Journal of Natural Products* 78: 339-342.

Druzhinina, I. S., Chenthamara, K., Zhang, J., Atanasova, L., Yang, D., Miao, Y., Rahimi, M. J., Grujic, M., Cai, F., Pourmehdi, S., Salim, K. A., Pretzer, C., Kopchinskiy, A. G., Henrissat, B., Kuo, A., Hundley, H., Wang, M., Aerts, A., Salamov, A., Lipzen, A., LaButti, K., Barry, K., Grigoriev, I, V., Shen, Q., Kubicek, C. P. (2018) Massive lateral transfer of genes encoding plant cell wall-degrading enzymes to the mycoparasitic fungus *Trichoderma* from its plant-associated hosts. *PLOS Genetics* 14.

Fleißner, A., Oostlander, A.G., and Well, L. (2022) Highly conserved, but highly specific: Somatic cell–cell fusion in filamentous fungi. *Current Opinion in Cell Biology* 79: 102140.

Gibbons, J. G., Salichos, L., Slot, J. C., Rinker, D. C., McGary, K. L., King, J. G., Klich, M. A., Tabb, D. L., McDonald, W. H., & Rokas, A. (2012) The evolutionary imprint of domestication on genome variation and function of the filamentous fungus *Aspergillus oryzae*. *Current Biology* 22: 1403-1409.

Grigoriev, I. V., Nikitin, R., Haridas, S., Kuo, A., Ohm, R., Otillar, R., Riley, R., Salamov, A., Zhao, X., Korzeniewski, F., Smirnova, T., Nordberg, H., Dubchak, I., & Shabalov, I. (2014). MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic acids research*, *42*(Database issue), D699–D704.

Hagee, D., Abu Hardan, A., Botero, J., and Arnone, J.T. (2020) Genomic clustering within functionally related gene families in *Ascomycota* fungi. *Computational and Structural Biotechnology Journal* 18: 3267-3277.

Hughes, S.J. (1951) Studies of micro-fungi XI. Some *Hyphomycetes* which produce phialides. *Mycological Papers* 45: 1-36.

Iida, Y., Ikeda, K., Sakai, H., Nakagawa, H., Nishi, O., and Higashi, Y. (2018) Evaluation of the potential biocontrol activity of *Dicyma pulvinata* against *Cladosporium fulvum*, the causal agent of tomato leaf mold. *Plant Pathology* 67: 1883-1890.

Iida, Y., van 't Hof, P., Beenen, H., Mesarich, C., Kubota, M., Stergiopoulos, I., Mehrabi, R., Notsu, A., Fujiwara, K., Bahkali, A., Abd-Elsalam, K., Collemare, J., & de Wit, P. J. (2015) Novel Mutations Detected in Avirulence Genes Overcoming Tomato Cf Resistance Genes in Isolates of a Japanese Population of *Cladosporium fulvum*. *PLoS One* 10: e0123271.

Katoh, K., and Standley, D.M. (2013) MAFFT multiple sequence alignment software version
7: improvements in performance and usability. *Molecular Biology and Evolution* 30:
772-780.
Keller-Seitz, M.U., Certa, U., Sengstag, C., Würgler, F.E., Sun, M., and Fasullo, M. (2004)
Transcriptional response of yeast to aflatoxin B1: recombinational repair involving
RAD51 and *RAD1*. *Molecular Biology of the Cell* 15: 4321-4336.
Keller, N.P. (2019) Fungal secondary metabolism: regulation, function and drug discovery.
Nature Reviews Microbiology 17: 167-180.
Khaldi, N., & Wolfe, K. H. (2011). Evolutionary Origins of the Fumonisin Secondary
Metabolite Gene Cluster in Fusarium verticillioides and Aspergillus niger. *International
journal of evolutionary biology*, *2011*, 423821.
Kimura, N., and Tsuge, T. (1993) Gene cluster involved in melanin biosynthesis of the
filamentous fungus *Alternaria alternata*. *Journal of bacteriology* 175: 4427-4435.
Kubicek, C. P., Steindorff, A. S., Chenthamara, K., Manganiello, G., Henrissat, B., Zhang, J.,
Cai, F., Kopchinskiy, A. G., Kubicek, E. M., Kuo, A., Baroncelli, R., Sarrocco, S.,
Noronha, E. F., Vannacci, G., Shen, Q., Grigoriev, I. V., & Druzhinina, I. S. (2019)
Evolution and comparative genomics of the most common *Trichoderma* species. *BMC
Genomics* 20: 485.
Kubota, M., Morishima, M., and Iida, Y. (2015) First occurrence of tomato leaf mold caused
by the novel races 2.5.9 and 4.5.9 of *Passalora fulva* in Japan. *Journal of General Plant
Pathology* 81: 320-323.
Liu, X.-J., Li, H.-J., Ma, W.-Z., Zhang, F.-M., Xu, M.-Y., Mahmud, T., and Lan, W.-J.
(2021) Phomaligols F–I, polyoxygenated cyclohexenone derivatives from marine-
derived fungus *Aspergillus flavus* BB1. *Bioorganic Chemistry* 115: 105269.
Livak, K.J., and Schmittgen, T.D. (2001) Analysis of Relative Gene Expression Data Using
Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* 25: 402-408.
Machida, M., Asai, K., Sano, M., Tanaka, T., Kumagai, T., Terai, G. et al. (2005) Genome
sequencing and analysis of *Aspergillus oryzae*. *Nature* 438: 1157-1161.
Madeira, F., Pearce, M., Tivey, A.R.N., Basutkar, P., Lee, J., Edbali, O., Madhusoodanan, N.,
Kolesnikov, A., and Lopez, R. (2022) Search and sequence analysis tools services from
EMBL-EBI in 2022. *Nucleic Acids Research* 50(W1): W276-W279.
McGary, K.L., Slot, J.C., and Rokas, A. (2013) Physical linkage of metabolic genes in fungi
is an adaptation against the accumulation of toxic intermediate compounds. *Proceedings
of the National Academy of Sciences of the United States of America* 110: 11481-11486.
Mehrabi, R., Mirzadi Gohari, A., da Silva, G.F., Steinberg, G., Kema, G.H.J., and de Wit,
P.J.G.M. (2015) Flexible gateway constructs for functional analyses of genes in plant
pathogenic fungi. *Fungal Genetics and Biology* 79: 186-192.
Mehrabi, R., Bahkali, A.H., Abd-Elsalam, K.A., Moslem, M., M'barek, S.B., Gohari, A.M.,
Jashni, M.K., Stergiopoulos, I., Kema, G.H., and de Wit, P.J.G.M. (2011) Horizontal
gene and chromosome transfer in plant pathogenic fungi affecting host range. *FEMS
Microbiology Reviews* 35: 542-554.
Mello, S.C.M., Estevanato, C.E., Brauna, L.M., de Capdeville, G., Queiroz, P.R., and Lima,
L.H.C. (2008) Antagonistic process of *Dicyma pulvinata* against *Fusicladium
macrosporum* on rubber tree. *Tropical Plant Pathology* 33: 5-11.
Mesarich, C. H., Barnes, I., Bradley, E. L., de la Rosa, S., de Wit, P. J. G. M., Guo, Y.,
Griffiths, S. A., Hamelin, R. C., Joosten, M. H. A. J., Lu, M., McCarthy, H. M., Schol, C.
R., Stergiopoulos, I., Tarallo, M., Zaccaron, A. Z., & Bradshaw, R. E. (2023) Beyond the
genomes of *Fulvia fulva* (syn. *Cladosporium fulvum*) and *Dothistroma septosporum*:
New insights into how these fungal pathogens interact with their host plants. *Molecular
plant pathology* 24(5), 474–494.

Mitchell, J.K., and Taber, R.A. (1986) Factors Affecting the Biological Control of
Cercosporidium Leaf Spot of Peanuts by *Dicyma pulvinata*. *Phytopathology* 76: 990-
994.
Mitchell, J.K., Smith, D.H., and Taber, R.A. (1987) Potential for biological control of
Cercosporidium personatum leaf spot of peanuts by *Dicyma pulvinata*. *Canadian*
Journal of Botany-Revue Canadienne De Botanique 65: 2263-2269.
Mitchell, J.K., Taber, R.A., and Pettit, R.E. (1986) Establishment of *Dicyma pulvinata* in
Cercosporidium personatum Leaf Spot of Peanuts: Effect of Spray Formulation,
Inoculation Time, and Hours of Leaf Wetness. *Phytopathology* 76: 1168-1171.
Ökmen, B., Etalo, D. W., Joosten, M. H. A. J., Bouwmeester, H. J., de Vos, R. C. H.,
Collemare, J., & de Wit, P. J. G. M. (2013) Detoxification of alpha-tomatine by
Cladosporium fulvum is required for full virulence on tomato. *The New phytologist* 198:
1203-1214.
Peresse, M., and Lepicard, D. (1980) *Hansfordia pulvinata* mycoparasite destructive to
Cladosporium fulvum. *Mycopathologia* 71: 23-30.
Proctor, R. H., Van Hove, F., Susca, A., Stea, G., Busman, M., van der Lee, T., Waalwijk, C.,
Moretti, A., & Ward, T. J. (2013) Birth, death and horizontal transfer of the fumonisin
biosynthetic gene cluster during the evolutionary diversification of *Fusarium*. *Molecular*
Microbiology 90: 290-306.
Proctor, R.H., and Hohn, T.M. (1993) Aristolochene synthase. Isolation, characterization, and
bacterial expression of a sesquiterpenoid biosynthetic gene (*Ari1*) from *Penicillium*
roqueforti. Journal of Biological Chemistry 268: 4543-4548.
Qi, J., Luo, H., and Hao, B. (2004) CVTree: a phylogenetic tree reconstruction tool based on
whole genomes. *Nucleic Acids Research* 32: W45-W47.
Quistgaard, E. M., Low, C., Guettou, F., and Nordlund, P. (2016) Understanding transport by
the major facilitator superfamily (MFS): structures pave the way. *Nature Reviews*
Molecular Cell Biology 17: 123-132.
Richards, T.A., Dacks, J.B., Jenkinson, J.M., Thornton, C.R., and Talbot, N.J. (2006)
Evolution of Filamentous Plant Pathogens: Gene Exchange across Eukaryotic Kingdoms.
Current Biology 16: 1857-1864.
Rokas, A., Wisecaver, J.H., and Lind, A.L. (2018) The birth, evolution and death of
metabolic gene clusters in fungi. *Nature Reviews Microbiology* 16: 731-744.
Silvia de la Rosa, Christiaan R. Schol, Ángeles Ramos Peregrina, David J. Winter, Anne M.
Hilgers, Kazuya Maeda, Yuichiro Iida, Mariana Tarallo, Ruifang Jia, Henriek G. Beenen,
Mercedes Rocafort, Pierre J.G.M. de Wit, Joanna K. Bowen, Rosie E. Bradshaw,
Matthieu H.A.J. Joosten, Yuling Bai, Carl H. Mesarich (2023) Sequential breakdown of
the complex *Cf-9* leaf mould resistance locus in tomato by *Fulvia fulva*. *bioRxiv*:
2023.2008.2027.554972.
Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis
of large phylogenies. *Bioinformatics* 30: 1312-1313.
Steenwyk, J.L., Shen, X.-X., Lind, A.L., Goldman, G.H., and Rokas, A. (2019) A Robust
Phylogenomic Time Tree for Biotechnologically and Medically Important Fungi in the
Genera *Aspergillus* and *Penicillium*. mBio 10: 10.1128/mbio.00925-00919.
Stergiopoulos, I., and de Wit, P.J. (2009) Fungal effector proteins. *Annual Review of*
Phytopathology 47: 233-263.
Stolzer, M., Lai, H., Xu, M., Sathaye, D., Vernot, B., and Durand, D. (2012) Inferring
duplications, losses, transfers and incomplete lineage sorting with nonbinary species
trees. *Bioinformatics* 28: i409-i415.

Sushida, H., Sumita, T., Higashi, Y., and Iida, Y. (2019) Draft Genome Sequence of *Dicyma*
pulvinata Strain 414-3, a Mycoparasite of *Cladosporium fulvum*, Causal Agent of
Tomato Leaf Mold. *Microbiology Resource Announcements* 8.
Tanaka, S., Wada, K., Katayama, M., and Marumo, S. (1984b) Isolation of Sporogen-AO1, a
Sporogenic Substance, from *Aspergillus oryzae. Agricultural and Biological Chemistry*
48: 3189-3191.
Tanaka, S., Wada, K., Marumo, S., and Hattori, H. (1984a) Structure of sporogen-ao 1, a
sporogenic substance of *Aspergillus oryzae*. T*etrahedron Letters* 25: 5907-5910.
Tansakul, C., Rukachaisirikul, V., Chalothorn, T., Phongpaichit, S., and Sakayaroj, J. (2017)
Synthesis and cytotoxicity against KB and NCI-H187 cell lines of sporogen AO-1
analogues. *Phytochemistry Letters* 22: 128-132.
Tansakul, C., Rukachaisirikul, V., Maha, A., Kongprapan, T., Phongpaichit, S., Hutadilok-
Towatana, N., Borwornwiriyapan, K., & Sakayaroj, J. (2014) A new phenalenone
derivative from the soil fungus *Penicillium herquei* PSU-RSPG93. *Natural Product*
Research 28: 1718-1724.
Tirilly, Y., Kloosterman, J., Sipma, G., and Kettenesvandenbosch, J.J. (1983) A fungitoxic
sesquiterpene from *Hansfordia pulvinata. Phytochemistry* 22: 2082-2083.
Tsuge, T., Harimoto, Y., Hanada, K., Akagi, Y., Kodama, M., Akimitsu, K., and Yamamoto,
M. (2016) Evolution of pathogenicity controlled by small, dispensable chromosomes in
Alternaria alternata pathogens. *Physiological and Molecular Plant Pathology* 95: 27-31.
Videira, S.I.R., Groenewald, J.Z., Nakashima, C., Braun, U., Barreto, R.W., de Wit, P., and
Crous, P.W. (2017) Mycosphaerellaceae - Chaos or clarity? *Studies in Mycology* 87:
257-421.
von Arx, J.A. (1981) The Genera of Fungi Sporulating in Pure Culture. 316-317.
von Arx, J.A. (1982) The genus *Dicyma*, its synonyms and related fungi. *Proc K Nederl Akad*
Wet Ser C 85: 21-28.
Watanabe, H., Horinouchi, H., Muramoto, Y., and Ishii, H. (2017) Occurrence of
azoxystrobin-resistant isolates in *Passalora fulva*, the pathogen of tomato leaf mould
disease. *Plant Pathology* 66: 1472-1479.
Watarai, N., Yamamoto, N., Sawada, K., and Yamada, T. (2019) Evolution of *Aspergillus*
oryzae before and after domestication inferred by large-scale comparative genomic
analysis. *DNA Research* 26: 465-472.
Yan, L., Chen, J., Zhang, C., and Ma, Z. (2007) Molecular characterization of benzimidazole-
resistant isolates of *Cladosporium fulvum. FEMS Microbiology Letters* 278: 242-248.
Yoshida, K., Asano, S., Sushida, H., and Iida, Y. (2021) Occurrence of tomato leaf mold
caused by novel race 2.4.9 of *Cladosporium fulvum* in Japan. *Journal of General Plant*
Pathology 87: 35-38.
923

Figure 1. Deoxyphomenone produced by the mycoparasite *Hansfordia pulvinata* is an

antifungal sesquiterpene active against the tomato leaf mold pathogen *Cladosporium*

***fulvum*.**

(**A**) Leaf mold symptoms (brown lesions) caused by *C. fulvum* and lesions covered by white

mycelia of *H. pulvinata* on a tomato leaf. (**B**) Scanning electron micrographs of hyphae of *H.*

pulvinata parasitizing hyphae of *C. fulvum*. Right: detail of area in dotted box in left image.

Arrow: bulge in hypha of *C. fulvum* caused by hyphae of *H. pulvinata*. (**C**) Germinated

spores of *C. fulvum* 24 h after treatment with 120 µM deoxyphomenone or with 1% methanol

(v/v) as a control (0 µM deoxyphomenone). Germination percentages are in parentheses.

Chemical structure of deoxyphomenone is also shown. (**D**) Antifungal activities of various

concentrations of deoxyphomenone in MM agar on mycelial growth of tomato pathogenic

fungi (*Cladosporium fulvum*, *Pseudocercospora fuligena* and *Fusarium oxysporum* f. sp.

lycopersici). Bars = 10 µm.

Figure 2. Synteny of the deoxyphomenone biosynthetic (*DPH*) gene clusters conserved

in the mycoparasite *Hansfordia pulvinata* and *Aspergillus* section *Flavi*.

(**A**) Gene clusters conserved in *H. pulvinata* and *Aspergillus*. The original gene IDs (DIP) and

names (*HpDPH1* to *HpDHP6*) of *H. pulvinata* are above the genes (detailed in

Supplementary Table S1). Protein annotations are indicated by color. The names of *HpDPH*

homologous genes in *Aspergillus* mentioned in the text are indicated in the arrows. Numbers

indicate the length of the genes and UTRs in base pairs. ORFs (black arrows) were detected

in the upstream region of *HpDPH6* in three *Aspergillus* species, and no homologous gene

sequence was found in the NCBI non-redundant database. (**B**) Predicted biosynthetic pathway

of deoxyphomenone. Intermediates were logically inferred. FPP, farnesyl pyrophosphate.

Figure 3. Relative gene expression levels of the deoxyphomenone biosyntheticsis (*DPH*)

genes in the mycoparasite *Hansfordia pulvinata.*

A spore suspension of *H. pulvinata* 414-3 was sprayed on *Cladosporium fulvum* on water

agar (WA). *H. pulvinata* and *C. fulvum* were grown on PDA or WA as controls. Expression

was quantified by qPCR and normalized using those of *H. pulvinata* actin gene. Values are

the means of three biological replicates ($\pm$ SD). Significant differences ($P < 0.05$) in

expression for a gene among the three treatments are indicated by different letters.

Figure 4. Deoxyphomenone production by wild-type and knock-out mutants of

***Hansfordia pulvinata* and their antifungal activity.**

Wild-type 414-3 of *H. pulvinata,* *ΔHpDPH1* mutants (KO10, KO37) and *AoDHP1*-

complemented KO37 (CO10A) were cultured in MM broth for 2 weeks, and the filtered

culture filtrate collected to (**A**) quantify deoxyphomenone production in 414-3 and mutant

KO10 using LC-MS/MS. The retention time of the highest peak is given for each. (**B**) Time

course of deoxyphomenone production in 414-3 and mutant KO10. (**C**) Germination of

Cladosporium fulvum spores after 24 h of exposure to deoxyphomenone in culture filtrate

(CF) of 414-3 or KO10 (bars = 20 μm). Different letters in graph indicate significant

differences ($P < 0.05$) in germination among treatments. Values are means of three replicates

($\pm$ SD).

Figure 5. Deoxyphomenone production and sporulation by *A. flavus* wild-type

NBRC114564 and Δ*AfDPH1* mutant strains and ectopic mutants.

The wild-type*,* three Δ*AfDPH1* mutants, and two ectopic mutant strains were cultured in MM

broth or agar. Values are means of three replicates ($\pm$ SD). (**A**) LC-MS/MS analysis of

deoxyphomenone in culture filtrates of the various strains. ND: not detected. (**B**) Number of

spores formed on agar. Asterisk indicates significant difference ($P < 0.05$) compared to the

wild-type in Williams' test. (**C**) Colony morphology of the wild-type, Δ*AfDPH1* mutant and

ectopic AfT32 mutant.

Figure 6. Different effects of deoxyphomenone on spore germination of *Cladosporium*

***fulvum* and spore production of two *Aspergillus* species.**

Wild-types *C. fulvum* CF301, *A. oryzae* RIB40 and *A. flavus* NBRC114564 were grown on

MM agar supplemented with deoxyphomenone. (**A**) Inhibition of spore germination of

Cladosporium fulvum. (**B**) Effects on sporulation of RIB40 and NBRC114564. Values are the

mean of three biological replicates. Error bars indicate standard deviations. Asterisks indicate

significant difference ($p < 0.025$) compared to control treatment (0 μM) by Williams' test.

Figure 7. Phylogeny and sequence similarity of deoxyphomenone biosynthetic (*DPH*)

genes in *Hansfordia pulvinata* and *Aspergillus* species.

The phylogenetic species tree based on the fungal genomic sequences (approximately 15 kb)

listed in Supplementary Table S1 was constructed using maximum likelihood. Asterisks

represent species that have been verified to produce deoxyphomenone. Sequence of

Lyophyllum atratum belonging to *Agaricomycetes* was used as an outgroup. *Aspergillus*

species were categorized into three groups based on the presence of *DPH* genes. The heatmap

represents the pairwise similarity (%) of the deduced amino acid sequences of the *DPH*

homologous genes.

Figure 8. Proposed horizontal transfer of the deoxyphomenone biosynthetic (*DPH*) gene

cluster from ancestral *Aspergillus* species to the mycoparasite *Hansfordia pulvinata*.

The phylogenetic species tree was based on 500 monocore genes (a single homolog in each

of the species) from 60 fungal genomes and constructed using maximum likelihood, and

horizontal gene transfer, duplication and loss events inferred in Supplementary figure S14

were described. Horizontal gene transfer events for the *HpDPH1* (T1) and *DPH2/DPH4* (T2)

genes are indicated by yellow stars. Proposed horizontal transfer of the *DPH* cluster from

ancestral *Aspergillus* to *H. pulvinata* is highlighted by a yellow arrow. Gene loss is marked

by an x on the branch. The three *Aspergillus* groups classified by the presence of *DPH* genes

shown are the same as in Figure 7. Fungal species with either a complete or partial *DPH*

cluster are in black, and those without the cluster are in grey. *Aspergillus* sections are given

on the right. *Lyophyllum atratum* was used as an outgroup.

A

B

Cf

Hp

Cf

Hp

Hp

C

0 μM (76.5 ± 3.0)

120 μM (7.2 ± 2.0)

D

C. fulvum

P. fuligena

F. oxysporum

0   20   40   80   120

deoxyphomenone (μM)

Figure 1

A

H. pulvinata 414-3

| DIP_001952 | DIP_001953 | DIP_001954 (*HpDPH1*) | DIP_001955 (*HpDPH2*) | DIP_001956 (*HpDPH3*) | DIP_001957 (*HpDPH4*) | DIP_001958 (*HpDPH5*) | DIP_001959 (*HpDPH6*) |

B

HpDph1 HpDph3 HpDph4 HpDph5 HpDph2

Deoxyphomenone (sporogen-AO 1)

FPP

Figure 2

Legend:
- Lipase/thioesterase family protein
- Integral membrane protein
- Sesquiterpene synthase
- Short-chain dehydrogenase
- Cytochrome P450
- MFS transporter

Figure 3

A

414-3   12.74 min

KO10 (*ΔHpDPH1*)   11.16 min

Retention tme (min)

B

414-3

KO10

Time (day)

C

414-3 CF   KO10 CF

C. fulvum spore germination of (%)

414-3   KO10   KO37   CO10A   dH₂O

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8