1　**Title:** Sediment-associated processes account for most of the spatial variation in ecosystem
2　respiration in the Yakima River basin
3

4　**Authors:** Vanessa A. Garayburu-Caruso[1+*], Matthew Kaufman[1,2+*], Brieanne Forbes[1], Robert O.
5　Hall Jr.[3], Maggi Laan[1], Xingyuan Chen[1], Xinming Lin[1], Stephanie Fulton[1], Lupita Renteria[1], Yilin
6　Fang[1], Kyongho Son[1], James C. Stegen[1,4]
7
[1]Pacific Northwest National Laboratory, Richland, WA 99354
[2]Worcester State University, Worcester, MA 01602
[3]Flathead Lake Biological Station, University of Montana, Polson, MT 59860
[4]Washington State University, Pullman, WA 99163
[+] These authors contributed equally.
*Correspondence: vanessa.garayburu-caruso@pnnl.gov and mkaufman1@worcester.edu

14
15

Abstract: Hyporheic zones (HZ) contribute substantially to total stream ecosystem respiration
(ER_{tot}). We rely on process-based models to understand HZ respiration at watershed scales and
field-based estimates of ER_{tot} at reach scales. These emphasize different but related processes
that have not been jointly explored. Here we evaluate the degree to which spatial variation in HZ
respiration predicted via process-based models can explain spatial variation in field-estimates of
ER_{tot} across 33 sites in the Yakima River basin (YRB). We found that HZ respiration predicted
by process-based models did not explain spatial variation in field estimates of ER_{tot}. Our results
indicate that sediment-associated processes hydrologically connected to the active channel
explained most of the spatial variation of ER_{tot} in the YRB. 88% of sites showed that sediment-
associated respiration contributed to 50% or more of ER_{tot}, while a few sites showed as little as
0% contribution. We propose that ER_{tot} is influenced by processes that integrate contributions
from sediment components beyond the HZ, such as benthic algae and submerged
macrophytes, which are key primary producers in rivers. We encourage conceptual and
process-based models of ER_{tot} to integrate shallow HZ exchange with sediment-associated
primary production as these provide more complete accounting of processes influencing ER_{tot}.

31
Introduction
River corridors contribute substantial amounts of carbon dioxide (CO_2) to the atmosphere[1].
Specifically, areas where groundwater and surface water mix (i.e., hyporheic zones, HZ) can
contribute 3-96% of total ecosystem respiration[2–7]. These contributions are variable and are not
well understood at a basin scale. Field experiments have estimated HZ respiration via
extrapolations of sediment microcosms[8,9], as the difference between ecosystem and benthic
respiration[2], or via end-member mixing analysis using in-situ solute concentration[3]. However,
quantifying HZ respiration in the field across scales is difficult due to high spatial and temporal
heterogeneity in hydrologic conditions and stream geomorphology. Currently, we rely on models
like Networks with Exchange and Subsurface Storage (NEXSS)[10] and the River Corridor Model
(RCM)[11–13] to predict the spatial variation of HZ respiration at watershed to basin scales. These
and other watershed/basin scale models rely on physical stream characteristics (e.g., sediment
texture, geomorphology, discharge) to estimate hydrologic exchange fluxes that, when paired to
reaction network models, provide an estimate of HZ respiration rates[10–13]. While each of these
models provide insights into spatial patterns of HZ respiration, they have not been explored in
conjunction with field estimates of total stream ecosystem respiration.

Ecosystem metabolism and carbon (C) cycling in rivers have been widely studied via two key
processes: C fixation (gross primary production, GPP) and organic C mineralization (ecosystem

respiration, ER)[14]. Field estimates of these processes are often derived from diel dissolved
oxygen (DO) measurements that include the collective aerobic organisms (i.e., autotrophs and
heterotrophs) and habitats (i.e., benthic, planktonic, and hyporheic zones), which together
contribute to C cycling dynamics in streams and rivers[15]. Studying river metabolism allows for
deeper understanding about food webs, energy fluxes, C and nutrient cycling, as well as
predictions of ecosystem responses to environmental change[16–19].

River metabolism is influenced by several biotic and abiotic factors like light, flow, temperature,
nutrients, and hydrogeomorphology[20–22]. Additionally, studies worldwide have evaluated how
environmental factors like climate, elevation, and latitude impact aquatic ecosystems[1,23,24]. A
recent study across 222 rivers in the United States shows that light availability and stream flow
conditions govern spatial variation in both annual GPP and ER[25]. Additionally, a global
assessment of river metabolism shows that GPP and ER increased with latitude and suggests
that spatial variability of site characteristics like land use and vegetation are more influential
than latitudinal temperature effects on river metabolism[26]. These studies identified metabolism
drivers linked to chemical and biological processes across scales, yet they do not explicitly
consider the role of hydrologic exchange, a fundamental process within reactive transport
models that heavily influences HZ biogeochemistry. Thus, we lack conceptual understanding of
the degree to which HZ respiration predicted via process-based models explains spatial
patterns of field estimates of total stream ecosystem respiration (ER_{tot}) at a basin scale.

In this study, we test the hypothesis that spatial variation in HZ respiration predicted via
process-based models explain the spatial variation of field-estimates of ER_{tot} across 33 sites
within the Yakima River Basin (YRB). We examine the relative contributions of water column
respiration (ER_{wc}), and sediment associated respiration (ER_{sed}) to ER_{tot} to test the additional
hypothesis that ER_{tot} is driven primarily by contributions from ER_{sed}, and less so by contributions
from ER_{wc}. Upon finding that ER_{sed} usually accounts for most of ER_{tot}, we further investigate
spatial drivers of ER_{sed} across the basin. Our study highlights the importance of sediment-
associated processes for understanding basin scale spatial variability of ER_{tot} and emphasizes
the need for process-based forward-simulation capabilities that incorporate multiple components
of the river corridor, including water column processes, sediment-associated autotrophic and
heterotrophic metabolism in benthic zones and HZs, and hydrologic exchange.
80
Results and Discussion
Our study provided three primary outcomes: (*i*) Spatial variation in model-predicted HZ
respiration (as viewed through the river corridor model, RCM) did not explain spatial variation in
field-estimates of stream ecosystem respiration (ER_{tot}) at the basin scale. (*ii*) Contributions from
water column respiration (ER_{wc}) and sediment-associated respiration (ER_{sed}) were highly
variable and ER_{sed} was the primary contributor to ER_{tot} across most sites. (*iii*) variation in ER_{sed}
was primarily explained by gross primary production (GPP) and parameters reflecting the
physical environment. These results suggest that there are processes driving spatial variation in
ER_{tot} at a basin scale beyond those present in the RCM. We also infer there are processes in
the RCM that may not be captured in field estimates of ER_{tot}. Our results provide guidance for
which processes could be included in process-based models, such as the RCM, to support
research efforts focused on understanding basin scale patterns and dynamics of ER_{tot} and
ER_{sed}.

Spatial variation in RCM-predicted HZ respiration did not explain spatial variation in field-
estimates of ER_{tot} across the YRB.

Estimates of time averaged ER_{tot} for the deployment period across sites (n = 45) was highly
variable and ranged from -21.50 to 9.26 g O_2 m^{-2} day^{-1} (Fig. 1a). We could not estimate
metabolism (i.e., GPP and ER_{tot}) in 3 sites that failed to meet the assumptions of one station
open channel O_2 budgets due to slower flows and lack of turbulence[27]. In addition, out of the 45
sites where we were able to estimate metabolism, 12 sites displayed positive estimates of ER_{tot}.
These had dissolved O_2 (DO) that rarely went under 100% saturation, which indicates high gas
exchange rates potentially due to too much turbulence and bubbles at higher internal pressure
relative to the atmosphere[28]. Oversaturation hinders the estimation of ER_{tot} using open channel
DO time series because we have no simple way of knowing what the concentration of DO would
be in the river in the absence of any biological activity[29]. We removed sites with positive ER_{tot}
values from our analysis because they have no biological meaning, and their magnitude reflects
misspecification due to bubble-mediated gas exchange and not a biological process. After
removing these values, estimated time averaged ER_{tot} (n = 33) ranged from -21.50 to -0.12 g O_2
m^{-2} day^{-1}.

Field-estimates of ER_{tot} diverged from RCM-predicted HZ respiration, both quantitatively and
qualitatively. The fastest ER_{tot} rates occurred in the lower region of the YRB (Fig. 1a), in
locations with shallow topographic gradients and high order streams (i.e., topographic slopes of
0.0004 - 0.002 and stream orders of 5-7). The fastest HZ respiration rates predicted by the RCM
were, in contrast, in upper parts of the YRB (Fig. 1b). RCM-predicted HZ respiration values (n =
46) ranged from -800 to -0.024 g O_2 m^{-2} day^{-1}. The RCM-predicted rates were much faster than
estimates of ER_{tot} highlighting fundamental differences in the processes represented in the RCM
vs. processes that influence ER_{tot} field estimates. Adding processes to the RCM or making
inferences regarding quantitative differences in the rates is beyond the scope of the current
study. We instead use RCM-predicted spatial patterns as the best current representation for
spatial patterns in HZ respiration rates across the YRB. This approach allows for evaluating a
qualitative relationship between the spatial patterns of RCM-predicted HZ respiration and field-
based ER_{tot}.

134
Spatial variation in RCM-predicted HZ respiration had a weak negative relationship with spatial
variation in ER_{tot}. Before doing associated analyses and given the quantitative difference
between RCM predictions and ER_{tot}, we standardized rates within each dataset as z-scores,
whereby each dataset has a mean of zero and standard deviation of one. This does not change
quantitative relationships within each dataset or the shape of the data distribution, it only puts
the two datasets into the same quantitative scale. Following this normalization, we regressed
RCM-predicted HZ respiration z-scores against z-scores of field-estimates of ER_{tot} (Fig. 2).
These data violated the normality assumption for conducting a parametric regression, yet no

relationship is apparent in the data (Fig. 2a). A weak negative relationship was, however,
observed via Spearman rank-order correlation (Fig. 2b). The negative direction and weakness
of the relationship indicate, however, that HZ respiration—as represented in the RCM—is not a
primary factor explaining spatial variation of ER_{tot} in the YRB. The lack of a strong positive
relationship could be for several reasons, such as ER_{tot} not including respiration occurring
across the full HZ spatial domain that is represented in the RCM, and the RCM not including all
the river corridor components and drivers influencing ER_{tot} estimates.
150

Figure 2. Normalized model-predicted hyporheic zone (HZ) respiration rates do not increase with field-estimates of total
stream ecosystem respiration rates (ER_{tot}; n = 31). (a) Z-score normalized data suggest no relationship. Data distributions are
skewed (i.e., not normal) and thus violate assumptions of parametric regression, so no statistics are given. (b) Spearman rank-order
correlation reveals a weak, negative relationship. The Spearman correlation statistics are provided on the panel.

Field-estimates of ER_{tot} are based on DO dynamics that reflect the reach-scale integrated
outcome of processes occurring across all system components that are hydrologically
connected to the active channel[15]. This estimate includes respiration associated with subsurface
sediments along relatively short HZ flow paths. These HZ flow paths must be fully contained
within the upstream reach that influences observed DO dynamics at a given monitoring location.
If the HZ flow paths are longer, they should not influence observed DO and thus do not
influence estimated ER_{tot}. The length of the reach that a given DO sensor integrates varies
across streams and has been estimated to be roughly three times the turnover length of O_2[15]. It
is plausible that the RCM captures respiration along flow paths that are longer and deeper into
the HZ than what the DO sensors integrate across. If so, respiration in long flow paths—that are
captured in the RCM—could lead to the lack of positive correlation between RCM-predicted HZ
respiration and field-estimates of ER_{tot}. We suspect, however, that this might not be the primary
reason for the lack of a relationship because both RCM-predicted HZ respiration and field-
estimates of ER_{tot} account for short HZ flow paths. These shorter flow paths are likely to be
within the most biogeochemically active sediments, which should in turn be the primary drivers
of total HZ respiration[30]. We postulate that the lack of a relationship (Fig. 2) and divergence in
spatial patterns (Fig. 1a, b) is related to differences in which biogeochemical processes are
captured in the two methods.
174
The RCM represents hydrologic exchange through the HZ and sediment-associated respiration
that is fueled by reactants delivered by that exchange[13]. In the model, reactants delivered to HZ

sediments are not influenced by primary producers (e.g., photoautotrophic algal biofilms) and
the respiration contributed by primary producers is also not accounted for. Respiration occurring
by heterotrophs and photoautotrophic primary producers in the water column is also not part of
the RCM. All these processes can influence observed DO dynamics[31–33] and are thus implicitly
accounted for in field-estimates of ER_{tot}. This highlights that field estimated ER_{tot} is influenced by
several processes that are not represented in the RCM. This is not a critique or rejection of the
RCM, rather, it is a recognition that the divergence in spatial patterns between RCM-predicted
HZ respiration and field-estimates of ER_{tot} are likely due, at least in part, to a difference in the
processes captured by each method.

In addition to the differences in captured processes, RCM-predicted HZ respiration and ER_{tot}
may not be correlated if processes in the RCM do not strongly influence ER_{tot}, if data inputs to
the RCM misrepresent the system, or if there are technical issues associated with estimating
ER_{tot}. Sites with technical issues were removed from the analysis and our estimates of ER_{tot}
largely overlap with literature values[25,34] (Fig. 3a). It is, therefore, likely that the ER_{tot} estimates
represent actual stream respiration suggesting that technical issues with the estimates are not a
cause for the lack of correlation. Thus, the lack of correspondence in spatial patterns suggests
that to use the RCM, or other processed-based models, in future studies to help understand
ER_{tot} it may be useful to expand the breadth of represented processes. Available field estimates
of ER_{tot} allow us to guide which processes could be most important to include in the RCM and,
more generally, to investigate factors associated with spatial patterns in ER_{tot}. To these ends,
we next examine the relative influence of water column- and sediment-associated processes
over ER_{tot}, which provides conceptual understanding of system function and evaluates the need
for each of these in an expanded version of the RCM.

Figure 3. Density plots comparing respiration rates for different river corridor components within the Yakima River basin
(YRB) as well as literature values. (a) Density plot for YRB total ecosystem respiration from this study (ER_{tot}, grey, n = 33) and from
rivers and streams across the Contiguous United States (CONUS) (n = 222 estimated by Appling et al.[34] and Bernhardt et al.[25]
(ER_{tot}(Lit), black), as well as YRB ER_{sed} (brown, n = 33) and the range of YRB water column respiration rates (ER_{wc}, blue n = 48); (b)
Density plot for YRB ER_{wc} (blue). The vertical lines are the medians for each distribution.

The spatial distribution of ER_{tot} was largely explained by variations in ER_{sed} with limited influence
from ER_{wc}.

Our ER_{wc} data, estimated via *in-situ* dark bottle incubations (n = 48) varied greatly across the
basin and ranged from -16.60 to +0.772 g O_2 m^{-2} day^{-1} (Fig. 3b). Contrary to the data cleaning
decisions made for ER_{tot}, we chose to interpret positive ER_{wc} because they reflect instrument
noise paired with extremely slow respiration. In this situation the rate at which O_2 was consumed
was not fast enough to overcome instrument error and resulted in a small positive ER_{wc}

measurement. If these values were excluded it would bias the dataset away from locations with
very slow ER_{wc} rates. Across all 33 field locations with ER_{tot} estimates, ER_{wc} varied greatly and
had fractional contributions of ER_{wc} to ER_{tot} spanning 0-100% (Table 1, Fig. S1a). The range of
measured ER_{wc} values spans a substantial portion of ER_{tot} literature values (Fig. 3a). These
results highlight the diversity of environments in the YRB and suggests that ER_{wc} can be of
similar magnitude to ER_{tot} across Contiguous United States (CONUS) rivers.
223
224 Table 1. Summary statistics (mean, median, and range) for ER_{tot} (g O_2 m^{-2} day^{-1}), ER_{wc} (g O_2 m^{-2} day^{-1}), ER_{sed}, (g O_2
225 m^{-2} day^{-1}), and ER_{wc} and ER_{sed} contributions to ER_{tot} (percent).

	Mean	Median	Range
ER_{tot}	-4.14	-3.15	(-21.50, -0.12)
ER_{wc}	-0.725	-0.0845	(-16.60, 0.772)
ER_{sed}	-3.05	-2.62	(-17.96, 0)
ER_{wc} (%)	8.4%	17.7%	(0%,100%)
ER_{sed} (%)	82.3%	91.7%	(0%, 100%)

235
Spatial patterns in ER_{wc} showed some similarity to spatial patterns in ER_{tot} (Fig. 1a, d, Fig. S1b).
As discussed in Laan et al.[35], ER_{wc} in the YRB spans values that are similar to other streams
and rivers around the world[36–38]. Variable ER_{wc} rates in the YRB could be due to variable
amounts of total suspended solids (TSS), nutrient concentrations, and changes in depth and
surface water residence time, moving from headwaters to larger order streams[35,36,39]. Moreover,
our results indicate that respiration contributions from ER_{wc} can vary widely across the YRB
during base-flow conditions. Within the YRB, 4 locations had ER_{wc} that contributed more than
50% of ER_{tot}. We observed the largest contribution of ER_{wc} to ER_{tot} in 4[th] and 5[th] order streams,
which were also the locations with the slowest ER_{tot}. While ER_{wc} and ER_{tot} rates at these
locations were slow, the water column was responsible for all the observed respiration. These
findings align with previous studies finding large ER_{wc} contributions. For example, working in
mid-sized rivers, Reisinger et al.[38] found that ER_{wc} contributed an average of 35% to ER_{tot}, and
up to 81% in rivers with slow ER_{tot}. Similarly, the rates of other water column processes, like
nutrient uptake and denitrification, may increase with increasing river size, as material
processing shifts from being benthic-dominated to water-column dominated in rivers from 5[th] to
9[th] order[40–43]. These patterns suggest that the relative importance of ER_{wc} is governed by
several stream characteristics, some of which co-vary with stream order.
253
We further examined respiration contributions from river corridor components beyond the water
column (i.e., benthic/streambed sediments and HZs that are hydrologically connected to the
active channel). We define the respiration that comes from these integrated components as
sediment-associated respiration (ER_{sed}). ER_{sed} was calculated as the difference between ER_{tot}
and ER_{wc} at each site with negative ER_{tot} (n = 33) assuming zero ER_{wc} rates if ER_{wc} values were
positive (n = 12). ER_{sed} values ranged from -17.96 to 0 g O_2 m^{-2} day^{-1} (Table 1). The relative
contribution of ER_{sed} to ER_{tot} spanned from 0-100%, with 29 out of 33 locations (88% of
locations) showing ER_{sed} contributions exceeding 50% of ER_{tot} (Table 1, Fig. 4a). The strong
contribution of ER_{sed} to ER_{tot} across the YRB aligns with previous studies that have found large
contributions of HZ and benthic sediments to ecosystem respiration[3,44–46]. It is important to note
that our ER_{sed} measurements integrate multiple ecosystem components, including the HZ,
benthic biofilms, and submerged macrophytes, whereas ER_{wc} specifically represents respiration
from organisms suspended in the water column.
267

ER_{sed} accounted for most of the spatial variability in ER_{tot} across the YRB (Fig. 1a,c). However,
no spatial correlation was observed between ER_{sed} and RCM-predicted HZ respiration (Fig. S2).
While ER_{sed} was fastest in lower parts of the YRB, the RCM predicted fastest HZ respiration in
upper parts of the YRB (Fig.1b, c). The correspondence between ER_{sed} and ER_{tot} spatial
patterns reflected the dominant contribution of ER_{sed} to ER_{tot} (Fig. 4b). This finding is consistent
with previous studies across different stream orders. For example, Naegeli and Uehlinger[2] found
that ER_{sed} contributed 76–96% of ER_{tot} in a gravel bedded river using closed chamber
measurements. Similarly, Fellows et al.[9] reported large HZ contributions to ER_{tot} (40-93%) in two
headwater streams under baseflow conditions. In contrast, Battin et al.[3], found lower HZ
contributions (41% of ER_{tot}) in a third-order piedmont stream, while Roley et al.[47] observed that
water column processes dominated ER_{tot} in the mainstem Columbia River (9th order river).
These varying results, together with our findings, demonstrate that ER_{sed} contributions can span
a wide range across river networks. While these studies provide valuable insights into specific
stream reaches, they highlight a critical knowledge gap in our understanding of what drives
ER_{sed} variability at the basin scale. Our dataset from the YRB provides a useful opportunity to
investigate potential drivers of spatial variation in ER_{sed} across an environmentally diverse river
network.

285
286

287
288
289 **Figure 4. ER_{sed} contributions to and correspondence with ER_{tot}.** (a) Map displaying the fractional contributions of sediment-
290 associated respiration (ER_{sed}) to total ecosystem respiration (ER_{tot}) across the YRB (n = 33). Color is used to indicate the magnitude
291 of the fractional contribution while symbol size is used to indicate magnitude of ER_{sed} respiration rate. (b) Scatterplot relating ER_{tot} to
292 ER_{sed}. The grey dashed line is the one-to-one line.

293
294

Spatial variability in ER_{sed} was best explained by primary production and physical settings.

To investigate factors explaining spatial variation in ER_{sed}, we identified variables known from
previous work to influence ER_{tot} (e.g., gross primary production; GPP) and variables that
influence HZ respiration (e.g., median grain size). We did not include RCM-predicted HZ
respiration as an explanatory variable because it was strongly correlated with median grain size
(i.e., d50), and because we did not find a positive relationship between RCM-predicted HZ

respiration and ER_{tot} or ER_{sed} (Fig. 2, Fig. S2). Our analysis showed that GPP, stream slope,
stream velocity, and total dissolved nitrogen (TDN) explained the most variation in ER_{sed} (Table
2). Other potential drivers (i.e., land cover classes, median grain size, nutrients, depth and water
temperature) had smaller regression coefficients (< 0.1) and were interpreted as being not as
important. The directionality of these relationships further showed that the fastest ER_{sed} rates
(and thus ER_{tot} rates) had the fastest GPP rates (Table 2). This finding is generally consistent
with the observation from Bernhardt et al.[25] who found that ER_{tot} was most strongly linked to
light availability, which should influence GPP. Additionally, faster ER_{sed} rates were observed in
reaches with shallower topographic gradients, lower velocity, and higher TDN concentrations.
Flow velocity controls biofilm scouring, architecture, and functional dynamics[48,49]. Moreover,
stream slope influences flow velocities, sediment transport[50], and nitrogen concentrations, which
can all impact biofilm community structure and C uptake[51]. Together these variables can have a
substantial effect on benthic metabolism and in turn ER_{sed}.
315
Table 2. Regression coefficients for LASSO models explaining variation in ER_{sed}. All input variables from 33 sites
were cube root transformed to reduce the impact of high leverage points in the regression analysis and standardized
as z-scores to enable direct comparison of the regression coefficients. The LASSO regressions were performed over
100 seeds. Reported mean regression coefficients (β) and R^2 were averaged across the 100 seeds. Normalized
regression coefficients were calculated by dividing each β coefficient by the maximum β coefficient in each seed.
Standard deviation (sd) was calculated for each coefficient over the 100 seeds. Values of 0 indicate that the variable
was included as an input to the model but not assigned a β coefficient.

Input Variable	Mean Normalized Regression Coefficient (β)	sd	Mean Regression Coefficient (β)	sd
Slope	1	0	0.293	0.014
Gross Primary Production	-0.697	0.014	-0.204	0.006
Velocity	0.631	0.029	0.185	0.017
Total Dissolved Nitrogen	-0.441	0.027	-0.129	0.003
Percent Agricultural Cover	0.003	0.025	0.001	0.009
Percent Forest Cover	0.003	0.021	0.001	0.007
Percent Shrub Cover	0.002	0.015	0.001	0.005
Drainage Area	0	0	0	0
Aridity	0	0	0	0
Discharge	0	0	0	0
Median Grain Size	0	0	0	0
Total Suspended Solids	0	0	0	0
Dissolved Organic Carbon	0	0	0	0
Mean Depth	0	0	0	0
Water Temperature	0	0	0	0
Mean R^2			0.52 (sd = 0.008)	

323
In addition, ER_{sed} and ER_{tot} covaried linearly with GPP (Fig. 5) providing insight into net
ecosystem production and the metabolic balance of the stream. This information can be
discerned by examining the position of sites relative to the one-to-one line in Fig. 5, where
points below the line indicate net heterotrophy and points above the line indicate net autotrophy.
Our results showed that 22 sites were net autotrophic and 11 sites were net heterotrophic. This
finding contrasts the general paradigm that stream ecosystems are predominantly heterotrophic,
where respiration is sustained by allochthonous C inputs, making rivers net sources of CO_2 to
the atmosphere[15,17,25,52,53]. In autotrophic conditions, rivers fix more carbon than they respire,

potentially storing or exporting this excess carbon downstream[15]. Larger number of sites
exhibiting net autotrophy in the YRB might be due to summer baseflow conditions when light
availability is maximized, combined with stable flow conditions that promote primary
production[54]. Similar patterns of autotrophy have been observed during baseflow in agricultural
systems[55], montane streams with some seasonal and year-to-year variability [56], and managed
pastoral streams[57].
338

Figure 5. ER_{sed} and ER_{tot} as a function of GPP. Scatterplot with linear regression lines depicting the relationship between GPP (x-
axis) and field estimates of sediment respiration (ER_{sed}, brown, y-axis) and field estimates of total respiration (ER_{tot}, dark grey, y-
axis). The grey dashed line is the one-to-one line with slope of –1.

The relatively large LASSO coefficient for GPP (the largest coefficient after slope, Table 2), and
faster ER_{sed} (more negative) with increasing GPP (Fig. 5), suggest an important link between
sediment-associated primary production and ER_{sed}. Primary producers in this case are likely to
include both algal biofilms attached to benthic/streambed sediments and submerged
macrophytes rooted in sediments. We infer that both types of primary producers likely provide
important direct and indirect contributions to ER_{sed}. Direct contributions include phototroph
respiration, and indirect contributions include providing resources (e.g., organic exudates) to
sediment-associated heterotrophs[14,58]. This link between ER_{sed} and GPP in the YRB is
conceptually consistent with Bernhardt et al.[25]'s result that light is a primary explanatory variable
for ER_{tot} across CONUS. Further consistent with Bernhardt et al.[25] is our result that ER_{sed} rates
(and thus ER_{tot} rates) are likely influenced by variables beyond GPP that may be linked to
hydrologic exchange. In the YRB, low-gradient reaches with slow moving water had faster ER_{sed}
(Table 2). This relationship may additionally reflect a shift along the transporter-to-transformer
continuum[59–61] whereby residence time is long enough to enable significant reactions to occur,
thereby enhancing ER_{sed}. Together, these results point to a key role for the ratio of residence
time to reaction time as classically represented in the Damköhler number[62]. That is, we infer that
ER_{sed} hot spots[63] or control points[64] are more likely to occur when the Damköhler number is high
because of low gradient and slow flow causing long residence time and fast GPP causing short
reaction times. These factors are likely predictable from physical features of stream networks,
thereby providing opportunities to develop mechanistic or data-driven models of ER_{sed} hot
spots/control points.
364

365
366 **Conclusions**
367
Spatial patterns of ER_{sed} largely corresponded to ER_{tot} patterns, thus understanding basin-scale
ER_{tot} requires deeper understanding of controls on ER_{sed} spatial variability. Our analysis
revealed that ER_{sed} across the YRB was primarily explained by slope, GPP, velocity and TDN
with the fastest ER_{sed} rates occurring in reaches characterized by high GPP, gentle slopes,
slower velocities, and elevated TDN concentrations. This relationship suggests a complex
coupling between physical, chemical, and biological drivers linked to sediment-associated
primary production and the transporter-to-transformer continuum. Further, the lack of positive
correlation between RCM-predicted HZ respiration and field-estimates of ER_{sed} and ER_{tot}
highlights a need to extend conceptual and process-based models that aim to capture and
predict active channel respiration to represent all processes that have major influences over
those rates. The RCM relies heavily on several spatially variable inputs, including median grain
size (i.e., d50) and dissolved organic carbon (DOC) concentrations. These input data layers,
particularly d50, are generated from sparse field measurements and may not accurately reflect
local conditions in the basin. If the spatial patterns in d50 within the input data deviate from true
spatial patterns in d50, that will cause errors in predicted spatial patterns of HZ respiration.
Additionally, the model does not consider the impact of sediment grain size on reaction rates,
only on exchange rates.
385
To understand spatial patterns in ER_{tot} and ER_{sed} we propose that it is not enough to focus on
HZ respiration as influenced primarily by hydrologic exchange processes. Our results indicate a
need to explicitly represent primary producers and heterotrophs in both the water column and
associated with sediments. Current models can be modified in these ways to generate new
predictions for evaluation with further field deployments designed specifically to test them. We
refer to this approach as hypothesis-based model-experiment (ModEx) iteration, and this study
represents a key step in this approach. A ModEx approach goes beyond data-model integration
whereby models are used to pose new hypotheses that are tested with new data, leading to
modified hypotheses that guide follow-on model development, and so on. We encourage the
application of this approach in other basins and across other flow conditions to evaluate the
transferability of our outcomes. This is key to building predictive models that are useful across
Earth's diverse watershed systems.
398
399
Methods
1. Site selection
We selected the Yakima River basin (YRB) as a test basin for this study. The basin drains an
area of 15,941.4km^2 and spans forested mountainous regions in the west to arid valleys and
plains in the east (Fig 6). The basin has a diversity of land covers and land uses dominated by
dry rangeland, forest, and agriculture[65,66]. Annual precipitation ranges from 203-356 cm in the
west to 25 cm in the east[67].
407

408
409 **Figure 6.** Yakima River basin sampling sites superimposed on a map of catchment cluster analysis results.
410

Field sites spanned a range of stream orders, land cover types, physical settings, and
previously published model-predicted HZ respiration rates[13]. Because model-predictions were
performed at a reach scale for the Columbia River basin, we conducted a clustering analysis
following methods previously described by Laan et al.[35] to locate regions in our test basin with
representative characteristics from the Columbia River basin (Fig. 6). We then combined the
clustering analysis results with logistical considerations regarding site access, safety, and
model-predicted HZ respiration rates to select 48 sites that span a range of clusters, model-
predicted respiration rates and stream orders across the basin (Fig. 6). Table 3 shows a
summary of the cluster analysis results as well as number of sites in the YRB per cluster.
Geospatial variables for each site were extracted primarily from the Environmental Protection
Agency StreamCat Dataset and NHDPlusV2.1 using a custom R script[68]. Land use/land cover
data for all the sites were extracted from StreamCat [69], while catchment area and slope were
extracted via NHDPlusV2.1 with the R package 'nhdplusTools'[70]. The custom R script was also
used to extract aridity data from the Global Aridity Index and Potential Evapotranspiration (ET0)
Database: Version 3[71] with the terra::rast() and terra::extract() functions[72].

Table 3. Cluster analysis results across the Columbia River basin and Yakima River basin with similar biophysical and hydrologic
characteristics and the number and percentage of sites in each.

Cluster	Cluster Name	CRB Drainage Area (%)	YRB Drainage Area (%)	No. YRB Sites Per Cluster	Percent YRB Sites Per Cluster (%)
1	Tree dominated high elevation mesic	22.7	27.4	12	25.0
2	Water dominated	3.4	2.2	0	0
3	Tree dominated high elevation hydric	7.3	2.4	2	4.2
4	Shrub-steppe middle elevation xeric	25.4	28.1	5	10.4
5	Tree dominated middle elevation mesic	17.4	17.3	15	31.3
6	Tree dominated middle elevation xeric	23.9	22.6	14	29.2

"CRB Drainage Area" is the percentage of the total drainage area of the Columbia River basin that was classified in each cluster.
"YRB Drainage Area" is the percentage of the total drainage area of the Yakima River basin that was classified in each cluster.
"YRB Sites Per Cluster" is the total number of field sites in the Yakima River basin (n = 47) located in each cluster. "Percent YRB
Sites Per Cluster" is the percentage of the total number of sampling sites in the Yakima River basin located in each cluster.

2.   Sensor deployment and data collection

During the week of July 25–28, 2022, dissolved oxygen (DO) sensors (miniDOT Logger; PME,
Inc.; Vista, CA, USA) and level loggers (HOBO® Water Level Logger (13') U20L-04; Onset;
Bourne, MA, USA) were deployed with sensor windows facing upstream on the thalweg of the
riverbed at each site (n = 48) for 35 days during baseflow conditions. We were unable to deploy
sensors and sample first order streams due to the lack or low flow during the summer. The
sensors were secured to two bricks with zip ties using several different configurations
depending on streamflow velocity, depth, and channel substrate (Fig. 7b, d). DO sensors
recorded DO and water temperature continuously on a 1-min interval. To minimize sensor
biofouling in warmer, shallower rivers or rivers that received a lot of sunlight, 25 DO sensors
were equipped with wipers (miniWIPER for miniDOT; PME; Vista, CA, USA) and anti-fouling
copper plates (miniDOT Anti-fouling Copper Option; PME; Vista, CA, USA) (Fig 7d). Wipers
were programmed to wipe the sensor face once every 12 hours. The remaining 23 DO sensors
were equipped with full copper kits, which included a copper ring, silicone ring, and a copper
mesh (Fig. 7b). Non-vented level loggers recorded absolute pressure and water temperature
continuously on a 15-min interval. Additional barometric pressure (BP, Rugged BaroTROLL
Data Logger (9 m); In-Situ; Fort Collins, CO, USA) sensors were deployed at a subset of sites (n
= 9) to collect continuous BP and air temperature data throughout the study period (Fig. 7).
Sensors were deployed from a 1-m rope on a tree branch as close as possible to the DO
sensor/level logger deployments and recorded BP and air temperature continuously on a 1-min
interval. BP data was used convert level logger absolute pressure data to depth data
(Supplementary methods).
454
Level loggers and BP sensors were calibrated by the manufacturer and did not need further
calibration. DO sensor calibrations were verified at 100% oxygen saturation in the laboratory
prior to deployment and field-based data were corrected for sensor drift by applying a correction
factor using lab-generated data (Supplementary methods). Approximately mid-way through the
sampling period (August 8–11, 2022), sensors were field-checked for signs of disturbance,
exposure due to decreased flows, and biofouling. DO sensors and level loggers were retrieved
temporarily for maintenance and data downloads and immediately redeployed in the original
location or, if exposed by decreasing flows, redeployed closer to the channel thalweg. Sensors
were retrieved the week of August 29–September 1, 2022, and taken to the laboratory for data
download.
465

466
Figure 7. Example sensor deployment configurations on example study sites. Left panels emphasize the diversity of
environmental settings covered in this study. Right panels show two different sensor configurations used to accommodate different
environmental conditions. (a) Bull Creek (site S63P), Yakama Reservation, Yakima County, Washington, 25 July 2022. (b) DO
sensor and level logger deployed side-by-side on top of two bricks and secured with zip ties, 29 August 2022. DO sensor equipped
with an anti-biofouling copper plate and mesh to inhibit the growth of periphyton on the sensor face. Note the dense accumulation of
periphyton and sediment deposited on the sensor bodies and bricks. (c) Cle Elum River (site S24R) upstream of Cle Elum Lake,
Kittitas County, Washington, 11 August 2022. (d) Level logger (far right) and DO sensor equipped with a wiper kit (white sensor
body on left-hand side) and anti-biofouling copper plate, 01 September 2022. The white sensor body on the left-hand side of the
wiper kit controls the blue flywheel, which is equipped with a brush and programmed to wipe the sensor face 1× every 12 h. Note
the lighter accumulation of periphyton and sediment on the equipment deployed in the Cle Elum River (d) compared to the
equipment deployed in Bull Creek (b).

3. Channel depth, channel width, and discharge data collection

Depth data for sites located on wadeable streams (n = 39) was collected using wading-based
depth transects. For each desired reach length (Supplementary methods), we completed 5-10
depth transects over a period of one hour (a minimum of five transects was desired) using 2-
person teams, starting ~2 m upstream of the depth sensor. Transects were evenly distributed
along the length of the reach. Using a wading rod, depth measurements were collected along
the wetted width of the reach at five roughly equidistance points along a transect perpendicular
to flow. Field personnel manually recorded the start time and the coordinates for the starting
point of each transect; the end time of the last transect was also manually recorded. The
channel width of each transect was measured using a range finder; mean channel width was
calculated by taking the average of the range finder width measurements.
489

Depth data for sites located on non-wadeable rivers (n = 5) was collected using kayak- or
jetboat-mounted sonar (HDS-7 LIVE fish finder with Active Imaging™ 3 in 1 transducer;
Lowrance; Tulsa, Oklahoma, USA) or, in the case of very large or inaccessible rivers (n = 4),
publicly available gauging station data. Kayak-mounted sonar data was collected along each
non-wadeable reach by 3-person kayak teams using a sonar receiver mounted to the bottom of
a fishing-style kayak. Continuous point measurements were collected by turning on the sonar at
the top of each reach and floating downstream using a consistent, zig-zag pattern from bank-to-
bank along the entire length of the reach. Coordinates for the top and bottom of each reach and
sonar start and end times were manually recorded. HOBO level loggers mounted on a meter
stick were used to record depth in reach sections too shallow to collect accurate sonar data.
Mean channel width for non-wadeable rivers where depth data was collected using sonar was
calculated for each reach by averaging a minimum of five width estimates from the sonar map.
For two sites on the Naches River, WA, river depth and channel width were determined from a
rating curve developed by the Washington Department of Ecology
(https://apps.ecology.wa.gov/publications/SummaryPages/1803009.html). For three sites on the
Yakima River, river depth and channel width were determined from a rating curve developed
using data from the United States Geological Survey (USGS) gauging station at Union Gap,
Kiona and Mabton.
508
Stream discharge and velocity for each site were extracted from the National Oceanic and
Atmospheric Administration (NOAA) National Water Model Retrospective Dataset Version 2.1
(NWM V2.1; https://registry.opendata.aws/nwm-archive/). This dataset provides a 42-y
(February 1979 through December 2020) hourly simulation at all sampling locations. We
computed the August monthly average discharge and velocity across these simulation years as
input variables for determining an estimate of gas exchange rate (K600) for streamMetabolizer
model runs (Section 5). Median grain size (d50) data for all the sites were extracted from
NEXSS[10] which uses d50 data from the National Rivers and Streams Assessment and the
Wadeable Stream Assessment (https://www.epa.gov/national-aquatic-resource-surveys/nrsa).

4. Surface water sampling

During the week of August 8–11, 2022, surface water samples were collected in triplicate from
each site for different analysis. A set of samples were collected using a 50-mL syringe and
filtered into 40-mL amber glass vials (ThermoScientific™ Amber Clean Snap Vials) using a
0.22-μm sterivex filter (MilliporeSigmaTM Sterivex™ Sterile Pressure-Driven Devices;
MilliporeSigmaTM; Burlington, Massachusetts, USA). Filtered samples were analyzed for
dissolved organic carbon (DOC) and total dissolved nitrogen (TDN). Additionally, one unfiltered
water sample was collected using a pre-washed 2-L amber bottle (Nalgene™ Rectangular
Amber HDPE Bottles; ThermoFisher Scientific; Waltham, Massachusetts, USA) for total
suspended solids (TSS). All samples were stored on ice in the field and then refrigerated at 4° C
before shipping for analysis at the Pacific Northwest National Laboratory (PNNL) Marine and
Coastal Research Laboratory in Sequim, Washington (DOC, TDN) or transported to PNNL
Biological Sciences Facility Laboratory in Richland, Washington (TSS). TSS samples were
analyzed within one week of collection and DOC and TDN content were measured within two
weeks of collection.
533
Dissolved organic carbon (DOC) and TDN were measured using a Shimadzu TOC-L Total
Organic Carbon Analyzer. DOC was measured as non-purgeable organic carbon and TDN was
measured by converting nitrogen in the samples to nitrogen gas and detected via
chemiluminescence. Visual checks of calibration curves, samples, blanks, and check standard

peaks were performed before exporting data. Concentrations below the limit of detection (LOD)
of the instrument, or below the standard curve, were flagged[73]. The LOD for DOC was 0.17-0.51
mg C L^{-1} and the LOD for TDN was 0.027 mg N L^{-1}. If the concentration of an analyte was below
the lowest standard but above the LOD, the flag for that sample was replaced with half of the
concentration of the lowest standard. If the analyte was below the LOD, the flag for that sample
was replaced with a value corresponding to half of the LOD for that run. The coefficient of
variation (CV) for each replicate set was calculated. Replicates with CV of 30 % or higher were
flagged and the outlier sample was identified by calculating the distance between each of the
replicate samples. Replicates with the highest distance to the other two replicates were flagged
as outliers and removed from the analysis. Mean values for each site were then calculated from
the remaining replicates and inputted as predictor variables in the regression analysis (Section
8).
550
Total suspended solids (TSS) samples were filtered in the laboratory through a pre-weighed and
pre-combusted 4.7 cm, 0.7 µm GF/F glass microfiber filter (Whatman™ glass microfiber filters,
Grade 934-AH®; MilliporeSigma; Burlington, Massachusetts, USA). After water filtration, the
filter and filtration apparatus were rinsed with 30 mL of ultrapure Milli-Q water (Milli-Q® IQ Water
Purification System; MilliporeSigma; Burlington, Massachusetts, USA) to ensure that all residue
was captured by the filter. The filter was placed in foil and oven dried overnight at 45° C. The
filter was allowed to cool in a desiccator prior to weighing. TSS was calculated as the difference
between the weight of the filter before and after filtration of the water sample divided by the
volume of water filtered[74]. Similar to the approach applied to DOC/TN, if the TSS of a sample
was below the LOD, the flag for that sample was replaced with a value corresponding to half of
the LOD (TSS LOD = 0.24 mg L^{-1}) for the TSS method.

5.   Total Ecosystem Respiration (ER$_{tot}$) estimates

We estimated ER$_{tot}$ using the streamMetabolizer R package[27] which estimates ER, GPP and
gas exchange (K600) as parameters fit to a diel model of dissolved oxygen. Model inputs
consisted of time series data for DO, water temperature, barometric pressure (BP), adjusted
time series of depth (Supplementary methods), site ID, site latitude and longitude and a site-
specific prior estimate of gas exchange rate coefficient (K600, day^{-1}). Gas exchange normalized
to a Schmidt number of 600 (K600, day^{-1}) is a free parameter estimated within
streamMetabolizer via Bayesian inference and therefore requires a prior probability estimate.
We first estimated a prior k600 (m day^{-1}) value using the empirical relationship described in
Model Equation 7 from Table 2 in Raymond et al. (2012)[75] as well as site specific slope,
discharge, and velocity (Section 1 and 3). k600 values were then divided by the reach-averaged
depth (Supplementary methods) to obtain a prior probability estimate for K600.
574
Before running the model, the 15-min time series BP and reach-averaged depth were
interpolated (zoo::na.approx function)[76] to match the time interval of the water temperature and
DO. Then the temperature, DO, and BP 1-min time series data were down-sampled to a 15-min
resolution. Down-sampling improved both solution time and K600 pooling. DO time series was
visually inspected and days with odd DO signals or biofouling were removed from the timeseries
prior running the model. Additionally, we flagged and interpolated outliers in the DO time series
using the R function 'auto_data_cleaning' within the 'tsrobprep' package[77] and used interquartile
range for outlier detection. Further, photosynthetically active radiation (PAR) was estimated
using the streamMetabolizer::calc_light() function[27], which is based on solar time, latitude, and
longitude. Saturated DO concentration was calculated for each DO time series using water
temperature and BP times series data and the Garcia and Gordon (1992)[78] equation
(streamMetabolizer::calc_DO_sat function)[27].

For most of the sites, the Bayesian model was configured with normal K600 pooling, which is to
say we partially pooled daily log K600 estimates in a Bayesian multilevel model. We also used
the state-space option in streamMatbolizer to estimate observation and process error for
dissolved O_2. The prior for K600 (K600_daily_meanlog_meanlog) was set as the natural log of
the K600 prior estimate and a K600 scale value of 0.7. Group level variation in log K600 was
K600~normal (0,0.05). We estimated posterior probability distribution for parameters via
Hamiltonian Monte Carlo using the program Stan with 2000 warm-up steps, and 2000 saved
steps across four chains. Some sites failed to converge with the settings mentioned, thus the
K600 daily sigma prior was changed to 0.08, 0.02, 0.01 or 0.05 depending on the site.
StreamMetabolizer outputs comprised ER_{tot} (g O_2 m^{-2} day^{-1}), GPP (g O_2 m^{-2} day^{-1}), and K600
(day^{-1}) as mean daily estimates over the number of days with quality DO data over deployment
time[79]. Finally, we calculated the mean GPP and ER_{tot} for the entire deployment time for each
site.
600
6. Water column respiration (ER_{wc}) and sediment-associated respiration (ER_{sed}) estimates
We estimated water column respiration in triplicate for 75 min at each site during sampling week
using the *in situ* dark bottle incubation approach previously described by Laan et al. (2025)[35].
Briefly, a DO sensor (miniDOT Logger; Precision Measurement Engineering, Inc., Vista, CA,
USA) and a small, battery-powered mixing device (i.e., toy boat motor propellor) were placed
inside a 2-L dark bottle in triplicate (Nalgene™ Rectangular Amber HDPE bottles; ThermoFisher
Scientific, Waltham, Massachusetts, USA). DO sensors and mixing devices were placed in a
cooler with blue ice packs to keep them cool and minimize the time needed at each site for the
sensors to equilibrate with river water temperatures. Bottles with the sensor and mixing device
were rinsed three times with river water, and filled with river water from the thalweg, ensuring no
air bubbles were trapped inside. Once full, the bottles were deployed on the riverbed as close to
the thalweg as possible, depending on river depth and access to the shoreline, and weighted
down with bricks or larger rocks. The sensors recorded DO concentration and temperature at 1-
min intervals during the 75-min incubation period.
615
We extracted the raw DO and temperature sensor data for each site and plotted each variable
against incubation time for each set of triplicate incubations (n = 144). The plots were visually
inspected to a) confirm that temperature sensors were at equilibrium with the river temperature
when the 75-min incubation test period began and b) identify data gaps, outliers, and other data
anomalies. Following the visual inspection of plots, we applied the DO correction factor (as
described in Supplementary Methods), removed the first 5 minutes and trimmed the time-series
to 70 min to account for data anomalies due to emptying and refreshing river water in the bottles
at the beginning and end of the incubation period, and to ensure all sites had the same
incubation time.
625
Water column respiration rates (ER_{wc}) for individual triplicate incubation samples were
calculated as the slope of the linear regression model (R stats::lm function)[80] between DO and
incubation time in units of day^{-1}. The volumetric daily rate was then converted to daily areal
units (g O_2 m^{-2} day^{-1}) by multiplying by the reach-averaged depth (Supplementary methods).
Only sample observations with a normalized root mean square error (NRMSE) ≤ 0.03 were
used for further analysis[81]. Although all triplicate samples met the NRMSE criteria, we removed
a replicate from two sites after visual inspection due to the curve deviating from the other two
replicates. Furthermore, six additional sites only had two replicates due to sensor or data loss.
Median, mean, and range of ER_{wc} values for each site were calculated. 40% of ER_{wc} values
were slightly positive (i.e., ER_{wc} > 0). Although positive respiration rates are biologically

unrealistic, and values greater than 0 g O_2 m^{-2} day^{-1} but less than 0.5 g O_2 m^{-2} day^{-1} are difficult
to distinguish from zero, we retained positive ER_{wc} values to prevent biasing the data.
638
Sediment-associated respiration (ER_{sed}) estimates were performed by combining
streamMetabolizer outputs of mean total ecosystem respiration (ER_{tot}) with in situ dark bottle
incubations to isolate water column respiration (ER_{wc}). Respiration directly and indirectly
associated with sediments (ER_{sed}) was found by the difference between ER_{tot} and ER_{wc},
assuming zero ER_{wc} if the estimated values were positive. These, ER_{sed} estimates include
sediment-associated respiration from all ecosystem components besides the water column.

7. Model-predicted hyporheic zone respiration

Model-predicted HZ respiration values were previously published in Son et al., (2022)[13], and the
data is publicly available on ESS-DIVE[82]. Respiration values by Son et al. (2022)[13] correspond
to the amount of oxygen respiration in the HZ via vertical and lateral hyporheic exchange
between the stream and the HZ, respectively. Therefore, total model-predicted HZ respiration is
a sum of vertical and lateral respiration. The respiration is scaled by the stream surface area.
Respiration values reported in Son et al., (2023)[82] in units of mol O_2 day^{-1}, were converted to g
O_2 m^{-2} day^{-1} using the molar mass of oxygen and RCM stream reach area. We note that model
outputs cover only 46 out of the 48 sites that we studied in the YRB.

8. Data analysis

To explore how physical variables covary with in ER_{sed} at different sites, a LASSO (Least
Absolute Shrinkage and Selection Operator) regression model was built using physical,
chemical and environmental variables (Table 4) as inputs, and ER_{sed} as the target variable. Prior
to conducting the LASSO regression, variables were cube root transformed to reduce the
impact of high leverage points in the regression analysis. Additionally, standardization was
applied to achieve a mean of zero and standard deviation of 1 (i.e. z-score) for all variable to
ensure that each variable is equally and fairly treated without penalizing a variable simply due to
the magnitude or scale of its nature. The LASSO regression was performed over 100 random
seeds. β coefficients were normalized to the maximum β coefficient for each regression, then
averaged over the 100 random seeds for the final, reported value. Both the raw and normalized
mean β coefficient and standard deviation are reported in addition to the R^2.
666
To investigate the correlation between RCM-predicted HZ respiration and field-estimated ER_{tot}
and ER_{sed}, Pearson as well as Spearman's rank correlation were used. In this study, we used
Pearson correlation to investigate the correlation between the z-score normalized RCM-
predicted HZ respiration and ER_{tot}/ER_{sed} (with mean of zero, standard deviation of one), and
Spearman's rank-order correlation to investigate their rank-order correlation. Linear regression
analyses were performed using the 'lm' function in the base R package. All analyses were
performed using R Statistical Software (v4.1.3)[80].

679 Table 4. Variables used as inputs in the LASSO regression model. All NHDPlus, StreamCat, NLCD, and ET0
680 database data were extracted using the custom R script[68].

Variable	Description	Unit	Data Source	Data Source Citation
Water Temperature	HOBO water temperature	C°	Field measured	Delgado et al. (2023)[74]
Mean Depth	average water depth	cm	Field estimated	Delgado et al. (2023)[74]
Slope	stream slope	NA	NHDPlus	McKay et al. (2012)[70]
Velocity	stream velocity	$m\ s^{-1}$	National Water Model	Cosgrove et al. (2024)[83]
Discharge	Discharge	$m^3\ s^{-1}$	National Water Model	Cosgrove et al. (2024)[83]
Dissolved Organic Carbon	non-purgeable organic carbon	$mg\ L^{-1}$	Field measured	Forbes et al. (2023)[73]
Total Dissolved Nitrogen	total dissolved nitrogen	$mg\ L^{-1}$	Field measured	Forbes et al. (2023)[73]
Total Suspended Solids	total suspended solids	$mg\ L^{-1}$	Field measured	Delgado et al. (2023)[74]
Drainage Area	total drainage area	km^2	NHDPlus	McKay et al. (2012)[70]
Percent Forest Cover	percentage of forest as the sum of percentage deciduous, coniferous and mixed forest	%	StreamCat and NLCD	Hill et al. (2016)[69] Jin et al. (2023)[84]
Percent Agricultural Cover	percentage agriculture as the sum of percentage crop and hay	%	StreamCat and NLCD	Hill et al. (2016)[69] Jin et al. (2023)[84]
Percent Shrub Cover	Percentage Shrub	%	StreamCat and NLCD	Hill et al. (2016)[69] Jin et al. (2023)[84]
Aridity	Aridity	NA	ET0 Database	Zomer et al. (2012)[85]
Median grain size (d50)	median grain size	m	National Rivers and Streams Assessment and the Wadeable Stream Assessment	Kaufman et al. (2023)[79]
Gross Primary Production (GPP)	gross primary production	$gO_2\ m^{-2}\cdot day^{-1}$	streamMetabolizer output	Kaufman et al. (2023)[79]

Data Availability:

685 Data collected in the field were published previously in Forbes et al. (2023)[73] and Delgado et al.
686 (2023)[74]. Data generated as part of this manuscript is currently available on GitHub (see below)
687 and will be published on the Environmental System Science Data Infrastructure for a Virtual
688 Ecosystem (ESS-DIVE) repository upon manuscript acceptance. ESS-DIVE repositories and
689 are licensed for reuse under the Creative Commons Attribution 4.0 International License.

Code Availability

691 Scripts necessary to reproduce the primary results of this article are available at
692 https://github.com/river-corridors-sfa/SSS_metabolism.

References:

1. Battin, T. J. *et al.* River ecosystem metabolism and carbon biogeochemistry in a changing world. *Nature* **613**, 449–459 (2023).
2. Naegeli, M. W. & Uehlinger, U. Contribution of the Hyporheic Zone to Ecosystem Metabolism in a Prealpine Gravel-Bed-River. *Journal of the North American Benthological Society* **16**, 794–804 (1997).
3. Battin, T. J., Kaplan, L. A., Newbold, J. D. & Hendricks, S. P. A mixing model analysis of stream solute dynamics and the contribution of a hyporheic zone to ecosystem function*. *Freshwater Biology* **48**, 995–1014 (2003).
4. Gagne-Maynard, W. C. *et al.* Evaluation of Primary Production in the Lower Amazon River Based on a Dissolved Oxygen Stable Isotopic Mass Balance. *Front. Mar. Sci.* **4**, (2017).
5. Fuss, C. & Smock, L. Spatial and temporal variation of microbial respiration rates in a blackwater stream. *Freshwater Biology* **36**, 339–349 (1996).
6. Kaplan, L. A. & Newbold, J. D. 10 - Surface and Subsurface Dissolved Organic Carbon. in *Streams and Ground Waters* (eds. Jones, J. B. & Mulholland, P. J.) 237–258 (Academic Press, San Diego, 2000). doi:10.1016/B978-012389845-6/50011-9.
7. Jones, J. Factors controlling hyporheic respiration in a desert stream. *Freshwater Biology* **34**, 91–99 (1995).
8. Pusch, M. & Schwoerbel, J. Community respiration in hyporheic sediments of a mountain stream (Steina, Black Forest). *Archiv fur Hydrobiologie* **130**, 35–35 (1994).
9. Fellows, C. S., Valett, M. H. & Dahm, C. N. Whole-stream metabolism in two montane streams: Contribution of the hyporheic zone. *Limnology and Oceanography* **46**, 523–531 (2001).
10. Gomez-Velez, J. D., Harvey, J. W., Cardenas, M. B. & Kiel, B. Denitrification in the Mississippi River network controlled by flow through river bedforms. *Nature Geosci* **8**, 941–945 (2015).
11. Fang, Y. *et al.* High-Performance Simulation of Dynamic Hydrologic Exchange and Implications for Surrogate Flow and Reactive Transport Modeling in a Large River Corridor. *Frontiers in Water* **2**, (2020).
12. Fang, Y. *et al.* A multirate mass transfer model to represent the interaction of multicomponent biogeochemical processes between surface water and hyporheic zones (SWAT-MRMT-R 1.0). *Geoscientific Model Development* **13**, 3553–3569 (2020).
13. Son, K., Fang, Y., Gomez-Velez, J. D. & Chen, X. Spatial Microbial Respiration Variations in the Hyporheic Zones Within the Columbia River Basin. *Journal of Geophysical Research: Biogeosciences* **127**, (2022).
14. Odum, H. T. Primary Production in Flowing Waters. *Limnology and Oceanography* **1**, 102–117 (1956).
15. Hall, R. O. & Hotchkiss, E. R. Chapter 34 - Stream Metabolism. in *Methods in Stream Ecology (Third Edition)* (eds. Lamberti, G. A. & Hauer, F. R.) 219–233 (Academic Press, 2017). doi:10.1016/B978-0-12-813047-6.00012-7.
16. Kaylor, M. J., White, S. M., Saunders, W. C. & Warren, D. R. Relating spatial patterns of stream metabolism to distributions of juveniles salmonids at the river network scale. *Ecosphere* **10**, e02781 (2019).
17. Hotchkiss, E. R. *et al.* Sources of and processes controlling CO2 emissions change with the size of streams and rivers. *Nature Geosci* **8**, 696–699 (2015).
18. O'Donnell, B. & Hotchkiss, E. R. Resistance and resilience of stream metabolism to high flow disturbances. *Biogeosciences* **19**, 1111–1134 (2022).
19. Gómez-Gener, L. *et al.* Towards an improved understanding of biogeochemical processes across surface-groundwater interactions in intermittent rivers and ephemeral streams. *Earth-Science Reviews* **220**, 103724 (2021).

20. Bernot, M. J. *et al.* Inter-regional comparison of land-use effects on stream metabolism.
Freshwater Biology **55**, 1874–1890 (2010).
21. Song, C. *et al.* Continental-scale decrease in net primary productivity in streams due to
climate warming. *Nature Geosci* **11**, 415–420 (2018).
22. Arroita, M., Elosegi, A. & Hall Jr., R. O. Twenty years of daily metabolism show riverine
recovery following sewage abatement. *Limnology and Oceanography* **64**, S77–S92 (2019).
23. Tromboni, F. *et al.* High rates of daytime river metabolism are an underestimated
component of carbon cycling. *Commun Earth Environ* **3**, 1–10 (2022).
24. Marzolf, N. S. & Ardón, M. Ecosystem metabolism in tropical streams and rivers: a review
and synthesis. *Limnology and Oceanography* **66**, 1627–1638 (2021).
25. Bernhardt, E. S. *et al.* Light and flow regimes regulate the metabolism of rivers. *Proceedings
of the National Academy of Sciences* **119**, e2121976119 (2022).
26. Gurung, A., Iwata, T., Nakano, D. & Urabe, J. River Metabolism along a Latitudinal Gradient
across Japan and in a global scale. *Sci Rep* **9**, 4932 (2019).
27. Appling, A. P., Hall, R. O., Yackulic, C. B. & Arroita, M. Overcoming Equifinality: Leveraging
Long Time Series for Stream Metabolism Estimation. *Journal of Geophysical Research:
Biogeosciences* **123**, 624–645 (2018).
28. Hall Jr, R. O., Kennedy, T. A. & Rosi-Marshall, E. J. Air–water oxygen exchange in a large
whitewater river. *Limnology and Oceanography: Fluids and Environments* **2**, 1–11 (2012).
29. Hall Jr., R. O. *et al.* Turbidity, light, temperature, and hydropeaking control primary
productivity in the Colorado River, Grand Canyon. *Limnology and Oceanography* **60**, 512–
526 (2015).
30. Roche, K. R. & Dentz, M. Benthic Biolayer Structure Controls Whole-Stream Reactive
Transport. *Geophysical Research Letters* **49**, e2021GL096803 (2022).
31. Schindler, D. E., Jankowski, K., A'mar, Z. T. & Holtgrieve, G. W. Two-stage metabolism
inferred from diel oxygen dynamics in aquatic ecosystems. *Ecosphere* **8**, e01867 (2017).
32. Hotchkiss, E. R. & Hall, Jr., Robert O. High rates of daytime respiration in three streams:
Use of $\delta^{18}OO_2$ and O_2 to model diel ecosystem metabolism. *Limnology and
Oceanography* **59**, 798–810 (2014).
33. del Giorgio, P. A., Cole, J. J. & Cimbleris, A. Respiration rates in bacteria exceed
phytoplankton production in unproductive aquatic systems. *Nature* **385**, 148–151 (1997).
34. Appling, A. P. *et al.* The metabolic regimes of 356 rivers in the United States. *Scientific Data*
5, 180292 (2018).
35. Laan, M. *et al.* Water Column Respiration in the Yakima River Basin is Explained by
Temperature, Nutrients and Suspended Solids. *EGUsphere* **2025**, (2025).
36. Ward, N. D. *et al.* Velocity-amplified microbial respiration rates in the lower Amazon River.
Limnology and Oceanography Letters **3**, 265–274 (2018).
37. Ward, N. D., Sawakuchi, H. O., Richey, J. E., Keil, R. G. & Bianchi, T. S. Enhanced Aquatic
Respiration Associated With Mixing of Clearwater Tributary and Turbid Amazon River
Waters. *Front. Earth Sci.* **7**, (2019).
38. Reisinger, A. J. *et al.* Water column contributions to the metabolism and nutrient dynamics
of mid-sized rivers. *Biogeochemistry* **153**, 67–84 (2021).
39. Ellis, E. E. *et al.* Factors controlling water-column respiration in rivers of the central and
southwestern Amazon Basin. *Limnology and Oceanography* **57**, 527–540 (2012).
40. Reisinger, A. J., Tank, J. L., Rosi-Marshall, E. J., Hall, R. O. & Baker, M. A. The varying role
of water column nutrient uptake along river continua in contrasting landscapes.
Biogeochemistry **125**, 115–131 (2015).
41. Reisinger, A. J., Tank, J. L., Hoellein, T. J. & Hall, R. O. Sediment, water column, and open-
channel denitrification in rivers measured using membrane-inlet mass spectrometry. *Journal
of Geophysical Research: Biogeosciences* **121**, 1258–1274 (2016).

42. del Giorgio, P. A. & Williams, P. J. le B. *Respiration in Aquatic Ecosystems*. (Oxford
University Press, Oxford, United Kingdom, 2005).
43. Gardner, J. R. & Doyle, M. W. Sediment–Water Surface Area Along Rivers: Water Column
Versus Benthic. *Ecosystems* **21**, 1505–1520 (2018).
44. Newcomer, M. E. *et al.* Influence of Hydrological Perturbations and Riverbed Sediment
Characteristics on Hyporheic Zone Respiration of CO_2 and N_2. *Journal of Geophysical
Research: Biogeosciences* **123**, 902–922 (2018).
45. Roche, K. R. *et al.* Benthic biofilm controls on fine particle dynamics in streams. *Water
Resources Research* **53**, 222–236 (2017).
46. Krause, S. *et al.* Ecohydrological interfaces as hot spots of ecosystem processes. *Water
Resources Research* **53**, 6359–6376 (2017).
47. Roley, S. S., Hall Jr., R. O., Perkins, W., Garayburu-Caruso, V. A. & Stegen, J. C. Coupled
primary production and respiration in a large river contrasts with smaller rivers and streams.
Limnology and Oceanography **n/a**, 15 (2023).
48. Hall, R. O., Tank, J. L., Baker, M. A., Rosi-Marshall, E. J. & Hotchkiss, E. R. Metabolism,
Gas Exchange, and Carbon Spiraling in Rivers. *Ecosystems* **19**, 73–86 (2016).
49. Battin, T. J., Kaplan, L. A., Newbold, J. D., Cheng, X. & Hansen, C. Effects of Current
Velocity on the Nascent Architecture of Stream Microbial Biofilms. *Appl Environ Microbiol*
69, 5443–5452 (2003).
50. Cohen, S., Wan, T., Islam, M. T. & Syvitski, J. Global river slope: A new geospatial dataset
and global-scale analysis. *Journal of hydrology* **563**, 1057–1067 (2018).
51. Anderton, C. R. *et al.* Nitrogen Source Governs Community Carbon Metabolism in a Model
Hypersaline Benthic Phototrophic Biofilm. *mSystems* **5**, e00260-20 (2020).
52. Duarte, C. M. & Prairie, Y. T. Prevalence of Heterotrophy and Atmospheric CO_2 Emissions
from Aquatic Ecosystems. *Ecosystems* **8**, 862–870 (2005).
53. Munn, M. D., Konrad, C. P., Miller, M. P. & Jaeger, K. A comparison of spatial and temporal
drivers of stream metabolism. *Freshwater Biology* **68**, 1751–1764 (2023).
54. Carter, A. M. *et al.* Exceptions to the Heterotrophic Rule: Prevalence and Drivers of
Autotrophy in Streams and Rivers. *Ecosystems* **27**, 969–985 (2024).
55. Griffiths, N. A. *et al.* Agricultural land use alters the seasonality and magnitude of stream
metabolism. *Limnology and Oceanography* **58**, 1513–1529 (2013).
56. Summers, B. M. *et al.* Long-term data reveal highly-variable metabolism and transitions in
trophic status in a montane stream. *Freshwater Science* **39**, 241–255 (2020).
57. Young, R. G. & Huryn, A. D. Effects of Land Use on Stream Metabolism and Organic Matter
Turnover. *Ecological Applications* **9**, 1359–1376 (1999).
58. Hoellein, T. J., Bruesewitz, D. A. & Richardson, D. C. Revisiting Odum (1956): A synthesis
of aquatic ecosystem metabolism. *Limnology and Oceanography* **58**, 2089–2100 (2013).
59. Fazekas, H. M., McDowell, W. H., Shanley, J. B. & Wymore, A. S. Climate Variability Drives
Watersheds Along a Transporter-Transformer Continuum. *Geophysical Research Letters*
48, e2021GL094050 (2021).
60. Price, A. N. *et al.* Biogeochemical and community ecology responses to the wetting of non-
perennial streams. *Nat Water* **2**, 815–826 (2024).
61. Ejarque, E. *et al.* Climate-induced hydrological variation controls the transformation of
dissolved organic matter in a subalpine lake. *Limnology and Oceanography* **63**, 1355–1371
(2018).
62. Rehage, H. & Kind, M. The first Damköhler number and its importance for characterizing the
influence of mixing on competitive chemical reactions. *Chemical Engineering Science* **229**,
116007 (2021).
63. McClain, M. E. *et al.* Biogeochemical hot spots and hot moments at the interface of
terrestrial and aquatic ecosystems. *Ecosystems* 301–312 (2003).

64. Bernhardt, E. S. *et al.* Control Points in Ecosystems: Moving Beyond the Hot Spot Hot
Moment Concept. *Ecosystems* **20**, 665–682 (2017).
65. Qiu, J. *et al.* Implications of water management representations for watershed hydrologic
modeling in the Yakima River basin. *Hydrology and Earth System Sciences* **23**, 35–49
(2019).
66. Vaccaro, J. J. & Olsen, T. D. *Estimates of Ground-Water Recharge to the Yakima River
Basin Aquifer System, Washington, for Predevelopment and Current Land-Use and Land-
Cover Conditions.* (2007).
67. Vano, J. A. *et al.* Climate change impacts on water management and irrigated agriculture in
the Yakima River Basin, Washington, USA. *Climatic Change* **102**, 287–317 (2010).
68. Willi, K. & Ross, M. R. V. Geospatial Data Puller for Waters in the Contiguous United States
(Version v1). (2023).
69. Hill, R. A., Weber, M. H., Leibowitz, S. G., Olsen, A. R. & Thornbrugh, D. J. The Stream-
Catchment (StreamCat) Dataset: A Database of Watershed Metrics for the Conterminous
United States. *JAWRA Journal of the American Water Resources Association* **52**, 120–128
(2016).
70. McKay, L. *et al. NHDPlus Version 2: User Guide.* (2012).
71. Zomer, R. & Trabucco, A. Global Aridity Index and Potential Evapotranspiration (ET0)
Database: Version 3. 6086884631 Bytes [object Object]
https://doi.org/10.6084/M9.FIGSHARE.7504448.V6 (2022).
72. Hijmans, R. J., Bivand, R., Pebesma, E. & Sumner, M. D. terra: Spatial Data Analysis.
(2024).
73. Forbes, B. *et al.* WHONDRS River Corridor Dissolved Oxygen, Temperature, Sediment
Aerobic Respiration, Grain Size, and Water Chemistry from Machine-Learning-Informed
Sites across the Contiguous United States (v3). [object Object]
https://doi.org/10.15485/1923689 (2023).
74. Delgado, D. *et al.* Spatial Study 2022: Surface Water Samples, Cotton Strip Degradation,
and Hydrologic Sensor Data across the Yakima River Basin, Washington, USA (v2). (2023).
75. Raymond, P. A. *et al.* Scaling the gas transfer velocity and hydraulic geometry in streams
and small rivers. *Limnology and Oceanography: Fluids and Environments* **2**, 41–53 (2012).
76. Zeileis, A. & Grothendieck, G. **zoo** : *S3* Infrastructure for Regular and Irregular Time Series.
J. Stat. Soft. **14**, (2005).
77. Narajewski, M., Kley-Holsteg, J. & Ziel, F. tsrobprep — an R package for robust
preprocessing of time series data. *SoftwareX* **16**, 100809 (2021).
78. Garcia, H. E. & Gordon, L. I. Oxygen solubility in seawater: Better fitting equations. *Limnol.
Oceanogr.* **37**, 1307–1312 (1992).
79. Kaufman, M. H. *et al.* Spatial Study 2022: Water Column, Sediment, and Total Ecosystem
Respiration Rates across the Yakima River Basin, Washington, USA. (2023).
80. R Core Team. R: The R Project for Statistical Computing. R Foundation for Statistical
Computing (2022).
81. Shcherbakov, M. V. *et al.* A survey of forecast error measures. *World applied sciences
journal* **24**, 171–176 (2013).
82. Son, K., Fang, Y., Gomez-Velez, J. D. & Chen, X. Model Inputs, Outputs, and Scripts
associated with: "Spatial microbial respiration variations in the hyporheic zones within the
Columbia River Basin". [object Object] https://doi.org/10.15485/1962818 (2023).
83. Cosgrove, B. *et al.* NOAA's National Water Model: Advancing operational hydrology through
continental-scale modeling. *J American Water Resour Assoc* 1752–1688.13184 (2024)
doi:10.1111/1752-1688.13184.
84. Jin, S. *et al.* National Land Cover Database 2019: A Comprehensive Strategy for Creating
the 1986–2019 Forest Disturbance Product. *J Remote Sens* **3**, 0021 (2023).

85. Zomer, R. J., Xu, J. & Trabucco, A. Version 3 of the Global Aridity Index and Potential Evapotranspiration Database. *Sci Data* **9**, 409 (2022).

Acknowledgements:
This work was supported by the River Corridor Science Focus Area (SFA) at the Pacific Northwest National Laboratory (PNNL). The SFA is supported by the United States Department of Energy, Office of Biological and Environmental Research (BER), Environmental System Science (ESS) Program. PNNL is operated by Battelle Memorial Institute for the United States Department of Energy under contract no. DE-AC05-76RL01830. We thank the United States Forest Service, Washington Department of Natural Resources, Washington Department of Fish and Wildlife, Confederated Tribes and Bands of the Yakama Nation, and Cowiche Canyon Conservancy for access to field locations where these samples were collected. We also thank the Confederated Tribes and Bands of the Yakama Nation Tribal Council and Yakama Nation Fisheries for working with us to facilitate sample collection and optimization of data usage according to their values and worldview. We thank the field team including: Dillman Delgado, Morgan Barnes, Brandon T. Boehnke, Yunxiang Chen, Kali Cornwell, Brianna I. Gonzalez, Samantha Grieger, Glenn E. Hammond, Peishi Jiang, Bing Li, Zhi Li, Xinming Lin, Sophia A. McKever, Maruti K. Mudunuru, Katherine A. Muller, Opal Otenburg, Aaron Pelly, Kelsey Peta, Alan Roebuck, Joshua M. Torgeson, and Jianqiu Zheng.

Author Contributions:
VAGC, MK, BH and JCS conceptualized the study. JCS, LR, SF, ML conducted the field work. LR and ML analyzed samples. KS, XC and YF configured RCM outputs for this study. XL, ML and BF processed the data and performed analyses with feedback from VAGC, MK, MK, BH and JCS. VAGC and MK carried out metabolism analysis with feedback from BH. BF managed the data. VAGC, MK, SF, BF and JCS drafted the initial manuscript, and all authors contributed to revisions.

Competing interests
All authors declare that there are no competing interests.