Title: Intracellular pH regulates β-catenin with low pHi increasing adhesion and

signaling functions

Condensed title: pH regulates β-catenin stability to alter function

Authors: Brandon J. Czowski[1,2] and Katharine A. White[1,2]

Affiliations:

[1]Department of Chemistry and Biochemistry, University of Notre Dame

[2]Harper Cancer Research Institute, University of Notre Dame

Summary:

Intracellular pH (pHi) regulates the degradation rate of the pH sensor β-catenin, altering

protein abundance, localization, and function in epithelial cells. This work shows pHi

acts as a rheostat to alter both adhesion and signaling functions of β-catenin.

Abstract:

Intracellular pH (pHi) dynamics are linked to cell processes including

proliferation, migration, and differentiation. The adherens junction (AJ) and signaling

protein β-catenin has decreased abundance at high pHi due to increased proteasomal-

mediated degradation. However, the effects of low pHi on β-catenin abundance and

functions have not been characterized. Here, we show that low pHi stabilizes β-catenin

in epithelial cells using population-level and single-cell assays. β-catenin abundance is

increased at low pHi and decreased at high pHi. We also assay single-cell protein

degradation rates to show that β-catenin half-life is longer at low compared to high pHi.

Importantly, we show that AJs are not disrupted by β-catenin loss at high pHi due to

rescue by plakoglobin. Finally, we show that low pHi increases β-catenin transcriptional

activity in single cells and is indistinguishable from a Wnt-on state. This work

characterizes pHi as a rheostat regulating β-catenin abundance, stability, and function

and implicates β-catenin as a molecular mediator of pHi-dependent cell processes.

Introduction

Spatiotemporal pHi dynamics regulate cell processes like differentiation

(Ulmschneider et al., 2016), cell migration (Denker and Barber, 2002), cell cycle

progression (Spear and White, 2023), and cell fate determination (Liu et al., 2023).

While low pHi is required to maintain the adult stem cell niche (Liu et al., 2023;

Ulmschneider et al., 2016) and transient increases in pHi are required for successful

differentiation (Ulmschneider et al., 2016), the molecular drivers of these pHi-dependent

processes are unknown. Recent work has identified β-catenin as a pH-sensitive protein

with decreased abundance at high pHi driven by pH-dependent binding of β-catenin to

the E3 ligase, β-transducin repeat containing protein (β-TrCP) (White, Katharine A.;

Grillo-Hill, Bree K.; Esquivel, Mario; Peralta, Jobelle; Bui, Vivian N.; Chire, Ismahan;

Barber, 2018).

β-catenin is a multifunctional protein that serves both structural and

transcriptional roles inside cells, maintaining adherens junctions (AJs) between

epithelial cells and functioning as the main signal transducer of the Wnt pathway

(Perez-Moreno and Fuchs, 2006; Vleminckx et al., 1999; Wodarz and Nusse, 1998).

Co-localization of β-catenin and E-cadherin at AJs is crucial for AJ integrity (Chen et al.,

1999), protects E-cadherin from degradation (Huber et al., 2001), and recruits α-catenin

to bridge AJs and the cytoskeleton (Drees et al., 2005). In the absence of Wnt ligand,

cytoplasmic β-catenin is bound and degraded by a destruction complex composed of

Axin, Adenomateous polyposis coli (APC), casein kinase 1 α (CK1α), glycogen

synthase kinase 3-β (GSK3β), and β-TrCP (Nusse and Clevers, 2017). Obligate

phosphorylation of β-catenin by CK1α and GSK3β target β-catenin for degradation (Yost

et al., 1996) by the proteasome (Hart et al., 1999). In the Wnt-on case, the destruction

complex is sequestered, reducing kinase activity and dissociating β-TrCP (MacDonald

et al., 2009; Mukherjee et al., 2018). Unphosphorylated β-catenin accumulates in the

cytoplasm and translocates to the nucleus to displace Groucho from the transcriptional

repressive complex (Groucho/TCF/LEF) and forms a transcriptionally active complex (β-

catenin/TCF/LEF) (Daniels and Weis, 2005). This active complex promotes the

expression of several proliferative and morphogenic genes, such as cyclin D1 (Tetsu

and McCormick, 1999), c-myc (He et al., 1998), and SLUG (Moon et al., 2021).

While White et al. previously showed that high pHi was sufficient to reduce whole

cell and junctional β-catenin abundance in the *Drosophila* eye epithelia and in normal

mammalian epithelial cells (White, Katharine A.; Grillo-Hill, Bree K.; Esquivel, Mario;

Peralta, Jobelle; Bui, Vivian N.; Chire, Ismahan; Barber, 2018), there were several

significant limitations to this prior work. First, the effect of low pHi on β-catenin

abundance was not explored due to the lack of methods for reliably lowering pHi in

epithelial cells. Second, the effects of pHi-mediated β-catenin abundance changes on

the integrity and composition of AJs were not completely characterized. Third, the pH-

dependent transcriptional activity of endogenous β-catenin was not investigated. Thus,

our understanding of how pHi regulates β-catenin function is incomplete. Given the

context-dependent function of β-catenin, a full characterization of pH-dependent

abundance of all subcellular pools (nuclear, cytoplasmic, junctional) is needed.

Here, we have developed methods to both raise and lower pHi in Madin Darby

Canine Kidney cells and assay pH-dependent effects on β-catenin abundance, sub-

cellular localization, and degradation dynamics. This enables us to investigate how both

increased and decreased pHi alters the abundance, degradation dynamics, and

functions of β-catenin. Using population level immunoblots we demonstrate pH-

dependent β-catenin abundance, with increased abundance at low pHi compared to

high pHi. We also characterize abundance of distinct subcellular pools of β-catenin

using immunofluorescent microscopy. We show that nuclear, cytoplasmic, and junctional

β-catenin pools are all decreased at high pHi while low pHi stabilizes both nuclear and

cytoplasmic pools of β-catenin. We also used a photoconvertible mMaple3-β-catenin

fusion to determine pH-dependent β-catenin degradation dynamics in single cells. We

showed that low pHi increases β-catenin half-life while high pHi reduces β-catenin half-

life. Collectively, our data demonstrate that pHi regulates β-catenin degradation rates by

functioning as a rheostat to modulate β-catenin abundance.

Importantly, our work also characterizes how pHi regulates β-catenin function by

identifying pH-dependent AJ composition where β-catenin is enriched at low pHi in E-

cadherin containing AJs. High pHi resulted in the loss of β-catenin at E-cadherin

containing AJs but recruitment of plakoglobin maintained AJ integrity and epithelial

morphology. We also demonstrate that modest (~10%) increases in nuclear abundance

of β-catenin at low pHi were sufficient to significantly increase β-catenin transcriptional

activity, phenocopying transcriptional activation observed with activation of the Wnt

pathway. Our results demonstrate that pHi functions as a rheostat to regulate the

abundance, stability, and function of endogenous wild-type β-catenin. This work will

improve our understanding of both normal biological processes (differentiation) and

pathological processes (transformation, cancer stem cells) that are marked by temporal

changes in pHi (Liu et al., 2020; Ulmschneider et al., 2016; White et al., 2017).

Results

Intracellular pH differentially affects subcellular pools of β-catenin

To investigate the pH-dependent regulation of β-catenin in a mammalian cell

model, we used Madin-Darby canine kidney (MDCK) cells. β-catenin has been

extensively studied in MDCK cells, where it was previously shown that increased pHi

was sufficient to reduce overall β-catenin abundance at the population level as well as

nuclear and junctional pools at the single cell level (White et al., 2018). This prior work

established a role for high pHi in regulating β-catenin abundance; however, the lack of

methods to lower pHi in this cell line prevented complete characterization of the role of

pHi in regulating β-catenin abundance, localization, and function.

We first developed a new approach to experimentally lower pHi in MDCK cells,

using a combination treatment of a sodium-proton exchanger 1 (NHE1) inhibitor (5-(N-

ethyl-N-isopropyl)amiloride (EIPA)) and a sodium-bicarbonate cotransporter (NCBn1)

inhibitor (2-chloro-N-[[2′-[(cyanoamino)sulfonyl][1,1′-biphenyl]-4-yl]methyl]-N-[(4-

methylphenyl)methyl]-benzamide (S0859)) (Larsen et al., 2012). To experimentally raise

pHi, we used 15% CO_2 atmospheric culture conditions, which increases the bicarbonate

concentration in the culturing medium and, through the function of NCBn1, results in

increased intracellular bicarbonate and increased pHi (White et al., 2018). Basal pHi in

control MDCK cells was measured at 7.36±0.02, while pHi manipulation treatments

were able to significantly increase pHi to 7.49±0.10 (15% CO_2) or significantly decrease

pHi to 7.17±0.05 (EIPA+S0859) after 24 hours (Figure 1A). Together, these

manipulation methods enable us to study how physiological changes in pHi of 0.13-0.32

pH units regulate β-catenin. These pHi changes are physiologically relevant as pHi

dynamics of similar magnitude occur during cellular differentiation (+0.2 pH units)

(Ulmschneider et al., 2016) and cell division (+0.25 pH units) (Spear and White, 2023).

Importantly, the presence of the proteosome inhibitor MG132 (Lee and Goldberg, 1996)

did not affect the pHi changes achieved with these manipulation protocols (Figure 1A).

We also verified that the treatments used to manipulate pHi for 24 hours did not alter

cell viability (Figure S1). Together, these conditions allow us to study roles for increased

and decreased pHi on β-catenin abundance.

We next used western blot approaches to assay β-catenin abundance at high

and low pHi. We did confirm that high pHi decreased β-catenin abundance compared to

control and we showed that low pHi was sufficient to stabilize β-catenin abundance

compared to high pHi (Figure 1 B&C). Importantly, we demonstrated that pH-dependent

abundance of β-catenin is regulated by proteosome—mediated degradation, as pHi

manipulation in the presence of a proteasome inhibitor (MG132) abrogated pH-

dependent abundance (Figure 1B&C). To ensure that low pHi was not inducing global

protein stabilization, we probed the abundance of the AJ protein, E-cadherin (Figure

1D). We found that high and low pHi conditions had no effect on the overall abundance

of E-cadherin (Figure 1E), demonstrating that the pH-dependent changes in β-catenin

abundance does not result from global protein degradation or stabilization. These data

suggest that pHi functions as a rheostat in regulating β-catenin abundance at the

population level, with increased stability at low pHi compared to high pHi.

150
Figure 1: Decreased pHi stabilizes β-catenin abundance through a proteosome-
dependent mechanism. (A) Box and whisker plots showing population level pHi
measurements in MDCK cells after 24-hour treatments with 1 μM EIPA+ 30 μM S0859
(low) or 15% CO_2 (high) to lower and raise pHi, respectively. Mean and SEM shown.
n=6-8. **(B)** Representative immunoblot of β-catenin (β-cat) and actin under low, control,
and high pHi conditions in the presence and absence of proteosome inhibition (MG132).
(C) Quantification of β-catenin immunoblot data collected as described in **(B)**. Individual
biological replicates were normalized to control MDCK within each experiment. Box and
whisker plots show median (line), 25th-75th percentile (boxes), min and max (whiskers),
and outlier values with points. n=6-8. **(D)** Representative immunoblot of E-cadherin and
actin under low, control, and high pHi conditions. **(E)** Quantification of E-cadherin

immunoblot data collected as described in **(D)** and displayed and normalized as
described in **(C)**. n=6. Statistical analyses in **(A)** were determined using one-way
ANOVA test with Sidak's correction for multiple comparisons. Statistical analyses in **(C)**
and **(E)** were determined with ratio paired t-tests between treatment groups and one-
sample t-tests with a hypothetical mean of 1.0 when comparing to control (which had no
variation due to normalization). *$P<0.05$; **$P<0.01$ ***$P<0.001$; ****$P<0.0001$.
168

Because the function of β-catenin is determined by its subcellular localization,

phosphorylation state, and availability of binding partners (Krieghoff et al., 2006; Morgan

et al., 2019; Valenta et al., 2012), we next investigated how pHi alters the abundance

and distribution of each subcellular pool of β-catenin. We first confirmed that the pHi

manipulation techniques altered pHi in single MDCK cells using live-cell, confocal

microscopy and nigericin standardization (Grillo-Hill et al., 2014; Spear and White,

2023) (Figure S2A). We found that under control conditions the median pHi value of

MDCK cells was 7.39±0.14 (Figure S2B), in line with the previous calculations using

population-level measurements. Importantly, treatment with EIPA+S0859 significantly

lowered single-cell pHi to 7.26±0.17, whereas treatment with 15% CO_2 significantly

raised pHi to 7.42±0.20, relative to control MDCK cells (Figure S2B).

We hypothesized that high pHi would decrease β-catenin levels in each

subcellular pool, whereas low pHi would stabilize each subcellular pool. This hypothesis

is predicated on recent data suggesting subcellular pools of β-catenin can rapidly

exchange and are regulated by the availability of subcellular binding partners (Krieghoff

et al., 2006; Morgan et al., 2019). To test this hypothesis, we developed a 3D image

analysis pipeline to segment normal epithelial cells (MDCK) and separately measure β-

catenin abundance in the nucleus, cytoplasm, and at adherens junctions (Figure S3,

see methods). Briefly, cytoplasmic and nuclear regions of interest were generated in

Nikon Imaging Software-Elements (NIS-Elements), using Hoechst dye (DNA) to identify

nuclear boundaries and an E-cadherin antibody (adherens junctions) to identify cell

boundaries (Figure S3). To specifically quantify junctional pools of β-catenin, we used

IMARIS (Oxford Instruments, see methods) software to generate 3D surfaces

representative of adherens junctions where E-cadherin was used as a marker of

adherens junctions.

Using this segmentation procedure, we quantified β-catenin abundance in each

subcellular pool (cytoplasmic, nuclear, junctional) after 24 hours of pHi manipulation

(Figure 2A). We found that all subcellular pools of β-catenin decreased at high pHi

(Figure 2B), which agrees with the population-level western blot data, and our

hypothesis of increased pHi being sufficient to reduce β-catenin abundance across all

subcellular pools. This data is also in agreement with prior quantification of β-catenin

abundance at junctions and in the nucleus of MDCK cells maintained at high pHi (White,

Katharine A.; Grillo-Hill, Bree K.; Esquivel, Mario; Peralta, Jobelle; Bui, Vivian N.; Chire,

Ismahan; Barber, 2018). At low pHi, we found that the cytoplasmic and nuclear pools of

β-catenin were significantly increased compared to control and high pHi conditions

(Figure 2A&B). In our initial analysis, low pHi was not sufficient to increase the total

abundance of β-catenin at adherens junctions compared to control MDCK cells (Figure

2B). However, contrary to our population level measurements (Fig 1C), we observed a

significant decrease in E-cadherin at MDCK junctions under low pHi compared to

control MDCK (Fig 2A&C). Importantly, this loss of E-cadherin at low pHi does not alter

cell morphology. We further quantified the ratio of β-catenin to E-cadherin at MDCK

junctions and observed strong pH-dependence with increased β-catenin localizing in E-

cadherin-containing AJs at low pHi compared to control and high pHi (Fig 2D). This

suggests that the level of β-catenin saturation at AJs changes with pHi, with less β-

catenin in E-cadherin-containing AJs at high pHi and more β-catenin in E-cadherin-

containing AJs at low pHi, even as total junctional E-cadherin is reduced.

215
Figure 2: β-catenin abundance across subcellular pools is differentially regulated by pHi. (A) Representative confocal images of MDCK cells fixed and stained for β-catenin (β-cat) and E-cadherin (E-cad). β-catenin is pseudocolored and E-cadherin is shown in inverse mono according to each scale. Scale bar: 20 µM. **(B)** Quantification of cytoplasmic, nuclear, and junctional pools of β-catenin. Individual cells and surfaces

were normalized to the median of control conditions within each biological replicate.
Individual cytoplasms, nuclei, and junctions are shown as gray dots with various
shading for each biological replicate. Red dots represent the median for each biological
replicate with interquartile ranges shown with bars. **(C)** Quantification of junctional E-
cadherin displayed and normalized as in **(B)**. **(D)** Calculation of β-cat/E-cad ratio from
quantifications in **(B)** and **(C)** that are normalized and displayed as in **(B)**. Individual
cytoplasms and nuclei, n=171-221; junctions, n=934-1194, from 3 biological replicates.
Statistical analyses in **(B-D)** were performed using the Kruskal-Wallis test with Dunn's
multiple comparison correction. ***P<0.001, ****P<0.0001.

Taken together, our data suggest that subcellular pools of β-catenin are

exchangeable, where high pHi causes cytoplasmic pools to be reduced, resulting in

concomitant loss of nuclear and junctional β-catenin. Similarly, our data suggest low pHi

results in cytoplasmic and nuclear accumulation and a relative increase in β-catenin at

E-cadherin containing AJs. Collectively, these data demonstrate that manipulating pHi

for 24 hours in MDCK cells significantly alters β-catenin abundance in single cells.

Quantification of each subcellular pool of β-catenin under each condition provides the

first glimpse as to how β-catenin distribution and abundance change with respect to pHi

dynamics.

Endogenous plakoglobin compensates for pHi-dependent loss of β-catenin to maintain

AJs at high pHi

In the immunofluorescence assays of β-catenin, we observed no change in

MDCK colony morphology with altered pHi, even when junctional pools of β-catenin

were significantly decreased at high pHi (Figure 2A&B). This suggests that other AJ

proteins may be able to rescue the loss of junctional β-catenin to maintain AJ integrity.

While β-catenin is well-known for maintaining AJs, plakoglobin serves adhesive

functions in both AJs and desmosomes, with preferential incorporation into the latter

(Cowin et al., 1986). Work by Fukunaga and colleagues demonstrated the functional

redundancy of β-catenin and plakoglobin in maintaining AJs, where cell-cell contacts

can be rescued with plakoglobin expression in a β-catenin null background (Fukunaga

et al., 2005). Furthermore, knockout of β-catenin in mouse models resulted in significant

recruitment of plakoglobin to AJs, compensating for the loss of β-catenin (Wickline et al.,

2011). A similar effect was observed in mouse models using conditional β-catenin

knockdown in cardiomyocytes, where β-catenin was reduced by 90% and plakoglobin

was subsequently enriched at junctions (Zhou et al., 2007). We therefore hypothesized

that in MDCK cells, the homologous protein plakoglobin can rescue β-catenin loss at

junctions at high pHi to preserve epithelial morphology.

To determine whether plakoglobin compensates for the pHi-dependent loss of β-

catenin from AJs at high pHi, we performed immunofluorescent confocal microscopy of

plakoglobin using the same segmentation pipeline outlined in Figure S3 after 24 hours

of pHi manipulation. We found that junctional plakoglobin was significantly increased at

high pHi and significantly decreased at low pHi compared to control MDCK (Figure

S4A&B). Importantly, overall abundance of plakoglobin is unchanged by pHi (Figure

S4C). In summary, while β-catenin co-localization with E-cadherin is reduced at high pHi

compared to control (Figure 2D), plakoglobin co-localization with E-cadherin is

increased at high pHi (Figure S4B). This suggests that when β-catenin is lost from

junctions at high pHi, plakoglobin is recruited to stabilize AJs and attenuate morphology

changes due to β-catenin loss.

Characterization of pH-dependent β-catenin degradation

Prior work has already extensively examined roles for high pHi in reducing β-

catenin stability (White, Katharine A.; Grillo-Hill, Bree K.; Esquivel, Mario; Peralta,

Jobelle; Bui, Vivian N.; Chire, Ismahan; Barber, 2018), but the effects of low pHi on β-

catenin stability has not been characterized. We next measured the temporal

localization and stability dynamics of endogenous β-catenin, with the prediction that low

pHi increases β-catenin stability resulting in the observed increased β-catenin

abundance at low pHi. We quantified pH-dependent stability of β-catenin by performing

a cycloheximide chase assay (Kao et al., 2015) after 24 hours of pHi manipulation in

MDCK cells, collecting samples every 30 minutes for 2.5 hours (Figure S5A). The half-

life of endogenous β-catenin was determined to be 1.58 hours in control MDCK cells, in

agreement with prior work in MDCK cells (White et al, 2018). Importantly, we found that

β-catenin half-life was increased at low pHi (2.79 hours) and decreased at high pHi

(1.22 hours) compared to control (Figure S5B). Of note, White et al. used different

treatment conditions to raise pHi (ammonium chloride compared to our method of 15%

CO_2) (Whtie et al, 2018). Thus, our work indicates that high pHi accelerates β-catenin

degradation, regardless of the method of pHi manipulation. Finally, this work is the first

to report an extension of β-catenin half-life in cells under low pHi conditions.

An experimental approach to monitor β-catenin degradation rates in single cells

The population-level analyses of β-catenin stability using cycloheximide

demonstrated pH-dependence, where low pHi increased and high pHi decreased β-

catenin half-life relative to control. Although these data revealed altered rates of β-

catenin degradation with pHi manipulation, the population data require discontinuous

measurements across cell populations and lack both single-cell and temporal resolution.

Single-cell analyses are important for answering biological questions, particularly for

elucidating roles for pH-dependent β-catenin regulation in pH-dependent processes that

occur at the single-cell level, such as differentiation (Ulmschneider et al., 2016). One

approach for quantifying live- and single-cell protein dynamics is through monitoring a

photoconvertible fluorescent protein (PCFP) fused to a protein of interest. PCFPs are a

subfamily of fluorescent proteins that undergo an excitation and emission spectra shift

upon stimulation with a specific wavelength of light (Shcherbakova and Verkhusha,

2014). This approach enables the photoconversion of a small subpopulation of the

PCFP-tagged protein, allowing for specific tracking of the photoconverted protein

fluorescence intensity as a direct readout of protein localization changes and

degradation (after accounting for photobleaching effects). This approach also eliminates

the need to average bulk cell responses as with cycloheximide chase assays, while

investigating how pHi regulates protein degradation in single cells.

We selected mMaple3, a PCFP that transitions from green-to-red fluorescence

upon exposure to 405 nm light, for these assays to monitor live-cell β-catenin protein

degradation rates. Derived from the parent protein mMaple, mMaple3 has reduced

dimerization tendency, superior brightness in the converted channel, and improved

photoconversion efficiency (defined as the proportion of protein that yields detectable

fluorescence signal) (McEvoy et al., 2012; Wang et al., 2014; Zhang et al., 2007). To

validate that we could measure differences in protein degradation rates of

photoconverted mMaple3 using confocal microscopy, we transiently expressed

mMaple3 alone or N-terminally fused to β-catenin (mMaple3-β-catenin) and determined

fitted half-life values of median cell traces to compare relative protein stability. This

assay does not permit determination of the absolute half-life of β-catenin, as loss of

fluorescent signal includes protein degradation, diffusion, and photobleaching of the

PCFP. However, this approach does allow us to compare relative degradation rates

across conditions. In MDCK cells expressing mMaple3 alone, the half-life was >1201

seconds, while the half-life of mMaple3-β-catenin was significantly reduced to just 290.9

seconds (Figure S6). This suggests that the shorter mMaple3-β-catenin half-life is being

driven by ubiquitination and degradation of β-catenin in the fusion protein. Confirming

this, we showed that treatment with proteasome inhibitor (MG132) significantly

increased the half-life of mMaple3-β-catenin to 686.9 seconds (Figure S6). Taken

together, these data demonstrate mMaple3-β-catenin is i) degraded much faster than

mMaple3 alone and ii) can be stabilized by inhibiting proteasome activity. Thus, using

this single-cell system we can monitor mMaple3-β-catenin degradation dynamics in

single cells and compare relative half-life values between conditions. Next, we

measured mMaple3-β-catenin degradation rates under pHi manipulation conditions in

single MDCK cells.

Intracellular pH titrates the degradation rates of β-catenin in single epithelial cells

We altered pHi in mMaple3-β-catenin expressing MDCK cells and

photoconverted cytoplasmic mMaple3-β-catenin using a spatially restricted region of

interest (Supplemental Video 1). We then monitored relative degradation rates of

cytoplasmic β-catenin in single cells (Figure 3A) and observed a reduced median half-

life at high pHi (200.2 seconds) and increased median half-life at low pHi (294.4

seconds) compared to control MDCK cells (263.4 seconds). Importantly, this approach

allows us to determine mMaple3-β-catenin half-life in individual cells. We similarly found

that cells with low pHi had significantly stabilized mMaple3-β-catenin compared to high

pHi MDCK cells (Figure 3B). These experiments demonstrate that pHi titrates the

degradation rate of cytoplasmic β-catenin in single cells, with faster degradation at high

pHi and slower degradation at low pHi compared to control.

349
Figure 3: mMaple3-β-catenin can be used to compare relative degradation rates of
β-catenin that are regulated by pHi. (A) Raw median cell traces for each condition
were fit using nonlinear regression to obtain half-life values. LOW, n=14; CRL, n=13;
HIGH, n=11; N=3. **(B)** Single-cell half-life values were obtained by fitting individual cell
traces as in **(A)**. Median shown as line with interquartile range shown as bars. Outliers
were removed using the ROUT method (Q=1%). LOW, n=14; CRL, n=13; HIGH, n=8
from 3 biological replicates. Statistical significance in **(B)** was determined using Kruskal-
Wallis test with Dunn's correction for multiple comparisons. *P<0.05, **P<0.01.
358

In the photoconversion experiments described above, we performed cytoplasmic

photoconversion and specific tracking of the cytoplasmic pool. This approach also

allows us to monitor subcellular redistribution of photoconverted cytoplasmic mMaple3-

β-catenin into the nucleus. We measured nuclear accumulation of photoconverted

mMaple3-β-catenin following cytoplasmic photoconversion and observed significantly

increased nuclear accumulation of mMaple3-β-catenin at low pHi compared to high pHi

cells (Figure S7). These data suggest that pHi modulates nuclear accumulation of

photoconverted β-catenin and supports the observation that nuclear β-catenin

abundance is increased at low pHi compared to control and high pHi (Figure 2B). While

the function of β-catenin is regulated by its localization, recent work revealed that

nuclear localization alone is not sufficient to determine β-catenin function (Van Der Wal

and Van Amerongen, 2020). Our work suggests that low pHi significantly increases β-

catenin abundance in the cytoplasm and nucleus from attenuated degradation in the

cytoplasm. A remaining unanswered question is whether the transcriptional activity of

wild-type, endogenous β-catenin is altered as a consequence of pH-dependent

degradation of cytoplasmic protein pools, as this was not addressed in prior work

investigating pH-dependent β-catenin stability (White, Katharine A.; Grillo-Hill, Bree K.;

Esquivel, Mario; Peralta, Jobelle; Bui, Vivian N.; Chire, Ismahan; Barber, 2018). To

address this, we tested the hypothesis that low pHi increases the transcriptional activity

of β-catenin whereas high pHi reduces transcriptional output.

Decreased pHi is sufficient to increase transcriptional activity of β-catenin

We first used population-level reverse transcriptase quantitative polymerase

chain reaction (RT-qPCR) of endogenous β-catenin transcriptional targets after 24

hours of pHi manipulation in MDCK cells. We identified MYC and SLUG as direct

transcriptional targets of β-catenin and verified primer specificity of each target (see

Methods) (Martínez-Estrada et al., 2006). At low pHi we found an increase in transcript

abundance for MYC and SLUG, while high pHi resulted in a decrease for each of these

targets (Figure S8). Similarly to the population-level β-catenin abundance analysis, we

only observed a modest pH-dependent trend for each target (Figure S8). Therefore, we

next compared the level of β-catenin transcriptional activity achieved at low pHi to an

active Wnt-on cell state. Stimulation of MDCK cells with WNT3a ligand for 24 hours

resulted in a similar increase of MYC and SLUG transcript abundance that observed

under low pHi conditions (Figure S8). Importantly, the observed level of pathway

activation agrees with the magnitude of response previously observed in WNT3a

stimulated MDCK cells (Howard et al., 2011). These data suggest that low pHi is

sufficient to induce β-catenin transcriptional activation that is indistinguishable from

WNT3a stimulation in MDCK cells.

A recent study has shown that Wnt activity is heterogenous within a tumor, and

single-cell analysis of Wnt pathway activation correlated with higher expression of target

genes (Yi et al., 2020). Thus, we reasoned that Wnt activation may be heterogenous

even in a clonal population of MDCK cells. To investigate this, we next performed

single-cell analysis of pH-dependent β-catenin transcriptional activity using a highly

specific LEF-1/TCF transcriptional reporter plasmid that drives expression of

destabilized GFP (TOPdGFP) (Reya et al., 2003). Cells transiently expressing the

plasmid were treated with pHi manipulation media for 24 hours prior to live-cell imaging

using confocal microscopy (Figure 4A). The single-cell transcriptional response was

heterogeneous in clonal normal epithelial cells, but we observed a statistically

significant increase in reporter fluorescence at low pHi compared to high pHi (Figure

4B). We again compared the extent of pH-dependent signaling upregulation to MDCK

cells treated with WTN3a. We found that WNT3a treatment resulted in an increase in

reporter intensity compared to control MDCK and that the extent of reporter transcription

with low pHi was statistically indistinguishable from Wnt3a (Figure 4A&B). These single-

cell analyses show that low pHi alone can recapitulate the Wnt-active state, highlighting

low pHi as a driver of β-catenin transcriptional activity in normal epithelial cells.

414
Figure 4: Low pHi phenocopies WNT3a-mediated activation of β-catenin
transcription. (A) Representative images of MDCK cells expressing a TOPdGFP
reporter plasmid shown according to scale with cell membranes visualized with Cell
Mask (magenta). Scale bar = 20μm. **(B)** Quantification of single-cell reporter intensities
after 24 hours of pHi manipulation or WNT3a treatment (40 ng/mL). All cells in each field
of view were quantified and were plotted if the average intensity was two-fold higher
than the average of mock-transfected cells. Individual cells are shown as block dots with
median and interquartile ranges shown in red lines and bars, respectively. LOW, n=103,
N=5; CRL, n=181, N=7; HIGH, n=147, N=5; WNT3a, n=103, N=3. Statistical
significance in **(B)** was determined using Kruskal-Wallis test with Dunn's correction for
multiple comparisons. *P<0.05.
426

Discussion

Our data highlight an important role for pHi in serving as a rheostat to regulate β-

catenin abundance, where low pHi increases β-catenin stability to promote high

adhesion and transcriptional activity whereas high pHi reduces β-catenin stability to

promote reduced β-catenin at AJs and lower transcriptional activity compared to control

cells. This significantly expands on prior work that only explored effects of increased pHi

on β-catenin abundance and did not fully characterize pH-dependent regulation of β-

catenin function. Our work demonstrates how changes in pHi within the physiological

range can regulate the degradation, and thus abundance and function, of endogenous

β-catenin.

We also found that subcellular pools of β-catenin were differentially regulated by

pHi. We show that pHi-mediated loss of β-catenin from AJs can be rescued by

plakoglobin to maintain AJ integrity. While genetic knockdown of β-catenin was

previously shown to recruit plakoglobin to adherens junctions (Fukunaga et al., 2005),

here we demonstrate the pHi-mediated loss of β-catenin is sufficient to produce the

same compensatory phenotype at AJs.

Our data suggests that pHi can alter not only β-catenin membrane localization,

but also redistribute the cadherin and catenin composition of adherens junctions. We

have demonstrated the plakoglobin is able to rescue the loss of β-catenin at adherens

junctions under high pHi. Additionally, we have found that the composition of adherens

junctions changes with pHi, with a preference for β-catenin at low pHi and a preference

for plakoglobin at high pHi. This presents a potential novel mechanism of pH-dependent

AJ remodeling that may be a previously unrecognized reinforcer of pH-dependent cell

polarity (Frantz et al., 2007) or EMT (Amith et al., 2016). Moreover, the increased ratio

of plakoglobin to E-cadherin at high pHi may be indicative of changes in desmosome

formation, as plakoglobin-containing AJs are a prerequisite for their establishment

(Lewis et al., 1997). Future work investigating differences in AJ strength of

plakoglobin/E-cadherin and β-catenin/E-cadherin complexes will further explore how the

physical properties of AJ change with pHi.

Collectively, our population and single-cell level analyses of β-catenin

transcriptional activity demonstrate that modest stabilization of nuclear β-catenin is

sufficient to drive significant changes in transcriptional activity. We found that low pHi

increased β-catenin transcriptional activity and phenocopied transcriptional outcomes of

Wnt3a treatment. Importantly, while plakoglobin can rescue the AJ function of β-catenin

at high pHi, our data suggest plakoglobin cannot rescue the reduced signaling function

of β-catenin at high pHi. This fits with prior work showing β-catenin/TCF4 complexes are

transcriptionally active but that plakoglobin/TCF4 complexes are unable to bind DNA

(Roura et al., 1999; Simcha et al., 1998). Our data show that pHi directly regulates β-

catenin signaling and transcriptional activity in the absence of Wnt pathway activation

and indicates that pH-dependent regulation of β-catenin signaling may reinforce stem

cell phenotypes previously associated with low pHi (Liu et al., 2023) and high Wnt

signaling (Nusse, 2008).

Decades of research have been dedicated to understanding the dynamics and

regulation of β-catenin in normal physiology and disease (Nusse and Clevers, 2017;

Wodarz and Nusse, 1998). Our work provides new mechanistic insight in the context of

pH-dependent differentiation, where stem cells have low pHi and high β-catenin

abundance and pHi increases are necessary for differentiation, accompanied by the

loss of β-catenin (Fevr et al., 2007; Liu et al., 2023). Future work will explore roles for

pHi dynamics in regulating β-catenin abundance through pH-dependent stability to

support both maintenance of pluripotency (stabilized β-catenin at low pHi) and

facilitation of differentiation (destabilized β-catenin at high pHi).

Finally, our work may shed light on how pHi regulates β-catenin abundance and

transcriptional activity in cancer. Constitutively increased pHi is an emerging hallmark of

cancer that is associated with several cancer cell behaviors (White et al., 2017). While

pHi has been shown to increase with metastatic potential in tissue-matched cell lines

(Spear and White, 2023), spatial pHi heterogeneity exists in 3D cancer cell cultures

(Dovmark et al., 2017; Tafech et al., 2023). Concurrent work in our lab has revealed that

microenvironment can alter pHi and concomitantly modulate β-catenin abundance in

cancer cell models. In combination with these data, this work suggests that pHi changes

may serve as a physiological cue to regulate β-catenin abundance and activity to drive

transcriptional and adhesion changes associated with cancer adaptation. Future work

will focus on simultaneously quantifying pHi and β-catenin activity in more complex

models to determine if pH-dependent regulation of the Wnt pathway (through β-catenin

regulation) alters tumorigenic phenotypes. This will create opportunities to investigate

context-specific avenues for altering β-catenin abundance and regulatory pathways in

order to develop improved cancer therapeutics.

Abbreviations:

AJ (adherens junctions)

β-TrCP (Beta-Transducin repeat containing protein)

β-cat (Beta-catenin)

E-cad (E-cadherin)

EIPA (5-(N-ethyl-N-isopropyl)amiloride)

MDCK (Madin Darby canine kidney)

PCFP (photoconvertible fluorescent protein)

pHi (intracellular pH)

Plak (plakoglobin)

S0859 (2-chloro-N-[[2′-[(cyanoamino)sulfonyl][1,1′-biphenyl]-4-yl]methyl]-N-[(4-

methylphenyl)methyl]-benzamide)

Materials and Methods:

Cell Culture and Conditions

Madin-Darby Canine Kidney (MDCK) cells were cultured in EMEM (Quality Biological,

112-018-101) supplemented with 10% Fetal Bovine Serum (FBS; Peak Serum, cat: PS-

FB3). Cells were maintained at 37º C in a humidified incubator maintained at 5% CO_2

(unless indicated otherwise). Cells were authenticated and tested for mycoplasma in

November 2022.

Intracellular pH manipulation

Cells were plated 24 hours prior to being treated for pHi manipulation. 24 hours after

plating, the conditioned medium was aspirated and replaced with medium containing 1

µM EIPA + 30 µM S0859 (Cf) (low pHi) or fresh medium and placed in an incubator

maintained at 5% CO_2 (low pHi and control) or 15% CO_2 (high pHi) for 24 hours. Where

indicated, proteasome activity was inhibited 18 hours after pHi manipulation treatment

using 10 µM MG132 (Selleck, S2619) for a total of 6 hours prior to sample analysis or

collection.

Acquisition and analysis of microscopy images

Plates were imaged using a Nikon Ti-2 spinning disk confocal microscope with a stage-

top incubator (Tokai Hit), a spinning disk confocal head (Yokogawa; CSU-X1), a Ti2-S-

SE stage, multi-point perfect focus system (PFS), an Orca Flash 4.0 CMOS camera,

four laser lines (405, 488, 561, 647 nm) with appropriate filter sets (DAPI: 405 nm laser

excitation, 455/50 nm emission; GFP: 488 nm laser excitation, 525/36 nm emission;

TxRed: 561 nm laser excitation, 605/52 nm emission; SNARF: 561 nm laser excitation,

705/72 nm emission; Cy5: 647 nm laser excitation, 705/72 nm emission) were used.

Immunofluorescent and live-cell transcriptional reporter images were collected on a 40X

oil immersion objective (Nikon, CFI PLAN FLUOR NA 1.30). Photoconversion images

were collected on a 60X oil immersion objective (Nikon, CFI PLAN APO OIL 1.40). Live-

cell imaging was performed after 24 hours of pHi manipulation. The stage top chamber

was equilibrated to 37º C and 5% CO_2 (or 15% CO_2 when capturing experimental

images for high pHi) prior to loading plates on microscope.

Trypan blue exclusion assay

MDCK cells were plated at 1×10^3 cells per well in a 24-well plate. Cells were cultured for

24 hours after plating before being treated for pHi manipulation as described above.

After the 24-hour treatment period, the medium was aspirated, the cells were washed

with 1 mL pre-warmed 1X Dulbecco's Phosphate-Buffered Saline (DPBS) (Quality

Biological, cat: 114-057-101) prior to trypsinization. Cells were trypsinized in 100 µL of

trypsin (Corning, cat: 25-053-CI) for 8-10 minutes to ensure detachment from the plates.

Each well then received 100 µL pre-warmed culture medium to inactivate the trypsin

and resuspend the trypsinized cells to a single-cell suspension. Immediately before

counting cell suspensions, 50 µL of the suspension was mixed with 50 µL trypan blue

0.4% solution (Gibco, cat: 15250-061) in a separate Eppendorf tube. Cells were

counted using 10 µL of the 1:1 cell suspension:trypan blue mixture in a hemocytometer

for healthy (no dye uptake) and unhealthy/dying (blue) cells. Each biological replicate is

the average of three wells (technical replicates) plated, treated, and counted on

independent days.

Population-level pHi measurement using BCECF-AM

MDCK cells were plated at 1×10^3 cells/mL in 24-well plate 48 hours prior to performing

the assay on the plate reader. Cells were treated for pHi manipulation as outlined

above. After 24 hours of treatment, 2'-7'-Bis-(2-carboxyethyl)-5-(and-)-

carboxyfluorescein, Acetoxymethyl Ester (BCECF-AM) (VWR, 891390244) was

prepared as a 100 µM stock solution in 10% DMSO in DPBS. Cells were dye loaded

with 1 µM BCECF-AM for 30 minutes in respective culturing conditions. Pre-warmed

HEPES-based wash buffer (30 mM HEPES, 145 mM NaCl, 5 mM KCl, 10 mM glucose,

1 mM $MgSO_4$, 1mM $KHPO_4$, 2mM $CaCl_2$, pH 7.4) and Nigericin buffer (25 mM HEPES,

105 mM KCl, 1 mM $MgCl_2$ supplemented with 10 µM Nigericin (Fisher, N1495)) are

prepared the day of the assay. Nigericin buffer is used to prepare three standard buffers

with pH values ~7.8, ~7.2 and ~6.5, where pH was accurately recorded to hundredths

place for each point and each biological replicate for pHi back-calculation. After dye-

loading, cells are washed 3x5minutes with HEPES buffer containing the appropriate

treatment at 37º C. Experimental fluorescence intensities were measured on a Biotek

Cytation5 plate reader (440ex/535em) and (490ex/535em) with kinetic reads taken

every 39 seconds for 7 minutes (11 reads). HEPES wash buffer was aspirated and

replaced with Nigericin buffer, pH ~7.8 and incubated for 7 minutes at 37º C before

starting the kinetic read. This procedure was repeated for the other two Nigericin

buffers. A data reduction step calculates the mean 490/440 intensity for experimental

and standardized pH reads where slope and y-intercept are calculated using Nigericin

standard buffer pH values to the hundredth place. Population level pHi values are back

calculated using experimental ratios and Nigericin standard curve values.

Whole-cell lysate preparation & protein quantification

After pHi manipulation (see above section) plates were removed from the incubator and

placed on ice. The media was aspirated, and the plates were washed two times with

ice-cold DPBS. Plates were wrapped with parafilm and stored at -80º C if not prepared

immediately. To washed or thawed plates on ice, 100 µL of lysis buffer (500 mM NaCl,

10 mM EDTA, 500mM HEPES-free acid, 10 mM EGTA, 10% Triton X-100, and protease

inhibitor cocktail (Pierce™ Protease Inhibitor Tablets, A32965; 1 tablet/50 mL lysis

buffer), pH 7.4) was added to each plate. Cells were removed using cell scrapers and

transferred to pre-chilled microfuge tubes, briefly vortexed, then incubated on ice for 15

minutes. Lysates were centrifuged at 14,000 x g for 15 minutes at 4º C. Clarified lysates

were isolated to separate microfuge tubes on ice. The Bradford assay was used to

quantify the total protein content of lysates (Bradford, 1976) using Coomassie Protein

reagent (Thermo Scientific, cat: 1856209). Lysates were stored at -80º C if not

immediately being used for SDS-PAGE sample preparation.

Immunoblotting

Protein samples were prepared in 6X Laemilli (reducing) buffer (Thermo Scientific, cat:

J61337.AC) and boiled for 10 minutes at 100º C. A total of 5 µg of MDCK lysate was

added to each lane for immunoblot analysis. Protein samples were loaded onto 10%

SDS-PAGE gels, which were run at 125V and 0.08 Amps for 15 minutes to stack

samples. Protein samples were separated by running the gel at 175 V and 0.08 Amps

for an hour and a half. Polyvinylidene difluoride (PVDF) membranes were pre-soaked in

100% ethanol for 5 minutes and rinsed 3X in water. Wet transfers were performed on

ice at a constant amperage of 0.04 A (100V) for an hour and half in transfer buffer (Tris-

Glycine buffer, 10% methanol, 0.01% SDS). Membranes were blocked for 1 hour in 5%

milk in 1X Tris-buffered Saline + 0.1% Tween® 20 (TBS-T) at room temperature with

shaking. Membranes were washed 3x5 min in 1X TBS-T before being cut and incubated

with primary antibodies. The following primary antibodies were used: β-catenin (BD

Biosciences, 610154; 1:5000 dilution in 3% BSA in 1X TBS-T), plakoglobin (Cell

Signaling, 2309S; 1:2000 dilution in 3% BSA in 1X TBS-T) and actin (Santa Cruz, sc-

58673; 1:500 diluted in 3% BSA in 1X TBS-T). Primary antibodies were incubated

overnight at 4º C with shaking. Primary antibody solutions were removed, and

membranes were washed 3x5 minutes in 1X TBS-T before incubating secondary

antibodies (Goat anti-mouse/rabbit HRP-conjugated antibodies; 1:10,000 diluted in 5%

milk in 1X TBS-T) for 1 hour at room temperature with shaking. Secondary antibody

solutions were removed, and membranes were washed 3x5 minutes in 1X TBS-T before

being developed using Pierce SuperSignal West Pico PLUS (Thermo Scientific, cat:

34580) (β-catenin, actin) or Pierce SuperSignal West Femto PLUS (Thermo Scientific,

cat: 34095) (plakoglobin). Chemiluminescence and colorimetric images were acquired

using a BioRad ChemiDoc imager.

Western Blot Analysis

Raw files were exported from ChemiDoc imager and opened in ImageJ. Band

intensities were acquired by densitometry analysis (area under the curve of individual

bands in respective lanes). For each replicate blot, target protein intensities were

normalized to the intensity of loading controls before normalizing each condition ratio to

that of the control within biological replicates.

Single-cell pHi calculations using SNARF-AM and confocal microscopy

Cells were plated at 5×10^3 cells per well (4-well glass bottom dish) or 2×10^4 cells per

dish (35 mm dish with 14 mm glass cover slip) in complete medium 48 hours prior to

imaging. Cells were treated for pHi manipulation as described above. Fields of view

(FOVs) were selected by viewing cells in differential interference contract (DIC). Cells

were dye loaded in conditioned media with 10 µM of 5-(and-6)-Carboxy SNARF ™-1

Acetoxymethyl Ester, Acetate (Fisher Scientific, cat: C1272) prepared as a 100 µM

stock solution dissolved in a 10% DMSO in DPBS for 10 minutes on the microscope

stage. Dye-containing media was removed, and plates were washed three times with

pre-warmed complete growth medium containing appropriate treatment. A 10-minute

equilibration period was used prior to capturing images in SNARF (20% laser power,

400 ms), TxRed (20% laser power, 400 ms), and DIC (200 ms) channels for each FOV

with five z-stacks (+/- 1 µm; 0.5 µm steps). SNARF dye was calibrated using three pre-

warmed Nigericin buffers (as described above). Complete medium was removed after

the initial capture and the pH ~7.8 Nigericin buffer was used to wash the plate and

incubated for 10 minutes prior to capture. This procedure was repeated for standard

points at pH ~7.2 and ~6.5. After data acquisition the NIS-Elements Analysis software

was used to define single-cell regions of interest (ROIs) after background subtraction for

each channel. The mean fluorescence intensity for each ROI in each channel was

exported to Microsoft Excel. Ratios were calculated by dividing the SNARF mean

intensity by the TxRed mean intensity for each capture. The three ratios for each

nigericin point and their corresponding pH values were used to calculate slope and y-

intercept. These values and the experimental ratio of SNARF/TxRed were used to back

calculate single-cell pHi values.

Immunofluorescent Staining

Cells were cultured in 35 mm imaging dishes containing a 14 mm glass coverslip and

plated for experiments 48 hours prior to fixation. Cells were treated for pHi manipulation

as described above. Intracellular pH was manipulated for 24 hours before removing

culture dishes from the incubator and washing them with ice-cold DPBS. After being

washed, cells were fixed in 3.7% paraformaldehyde in DPBS for 10 minutes at room

temperature with shaking. The fixative solution was removed, and cells were washed

3x2 minutes in DPBS. Cells were permeabilized for 10 minutes at room temperature in

a 0.1% Triton X-100 in 1X DBPS. Permeabilization buffer was removed and washed 3x2

minutes before blocking for 1 hour at room temperature in a 1% Bovine Serum Albumin

(BSA) (Millipore Sigma, cat: 2910-OP) in DPBS solution. The blocking solution was

removed and washed 3x2 minutes in DPBS. Primary antibodies (see Immunoblotting

methods 1.3.7 for vendor information) were diluted in antibody solution (1% BSA, 0.1%

Triton X-100 in DPBS) with the following dilutions: mouse anti-β-catenin (1:200), rabbit

anti-plakoglobin (1:400), and rat anti-E-cadherin (1:200). Primary antibody solutions

were incubated in dishes at 4º C overnight with rocking. After primary antibody

incubation, the plates were washed 3x2 minutes prior to incubating Alexa-Fluor

conjugated secondary antibodies diluted 1:1000 in antibody solution for 1 hour at room

temperature while being protected from light. Secondary antibody solutions were

removed and washed from the plates 3x2 minutes using DPBS. Cells were

counterstained with Hoechst dye (DAPI; Thermo Scientific, cat: 62249) diluted 1:10,000

in antibody solution for 10 minutes at room temperature protected from light.

Counterstain was removed and washed 3x2 minutes in DPBS. Imaging dishes were

stored and imaged in DPBS. Image acquisition details are outlined above. Z-stacks

were collected 2 μm above and below a relative z-plane that captured the center of the

cell determined using the DAPI channel. Nuclear pools of proteins were identified using

Nikon Elements Analysis software by auto-detecting individual regions of interest (ROI)

in the DAPI channel that represent individual nuclei. Whole cell regions of interest were

hand drawn using E-cadherin as a membrane marker to differentiate single-cells,

drawing the ROI interior to E-cadherin staining to prevent inclusion of membrane signal.

Nuclear ROIs were subtracted from hand-drawn whole cell ROIs to produce cytoplasmic

ROIs, excluding both cell membranes and nuclei. The mean intensity for each channel

was exported for each z-plane that had the largest mean intensity in the DAPI which

indicated the center of the individual cells. Junctional protein levels were quantified

using IMARIS software (Oxford Instruments, version 9.5.1) by generating surfaces

based on E-cadherin intensity. Mean intensities for all channels within each surface

were exported and analyzed for statistical significance using GraphPad Prism software.

Cycloheximide chase assay

Cells were plated at 1.5×10^4 cells per well in 6 well plate 48 hours prior to collecting

lysates. 24 hours after plating, cells were treated for pHi manipulation as described

above. Stock solutions of cycloheximide (VWR, 9427) were prepared fresh for each

experiment. After 20 hours of pHi manipulation, all plates except time-0 samples were

treated with 50 µg/mL cycloheximide and returned to the appropriate incubator.

Immediately after treatment, lysates from time-0 plates were prepared as described

previously and the remaining samples were lysed every 30 minutes for 2.5 hours.

Immunoblotting and analysis of β-catenin and actin were performed as described

above. The ratio of β-catenin to actin for each time point was normalized to time=0 h

within each condition and for each replicate. Normalized data for each biological

replicate was imported to Prism where one-phase decay was used to determine half-life

values.

Quantifying protein degradation and re-localization using mMaple3 constructs via

photoconversion and confocal microscopy

MDCK cells were seeded at 3×10^6 cells in 10 cm dishes and allowed to adhere to the

dish for 8 hours. 8 hours after seeding, cells were transfected with 15 µg mMaple3

containing plasmid (mMaple3 alone, mMaple3-β-catenin WT or H36R) using

Lipofectamine 3000 according to the manufacturer's instructions. Transfections

proceeded for 18 hours prior to the removal of transfection media, rinsing of the plate

with 10 mL of warm DPBS, and addition of 1 mL trypsin. Cells were trypsinized at 37° C

for 8-10 minutes. After all the cells were detached, complete medium was added to the

dish to create a single cell suspension. Using the transfected single cell suspension, 35

mm imaging dishes were seeded at 2.5×10^4 cell per dish and adhered for 8 hours. After

the cells adhered for 8 hours, pHi manipulation was performed as described above.

Images were collected as described above. Cell nuclei were visualized via SPY650-

DNA (Cytoskeleton, cat. #: CY-SC501) probe that was incubated for 2 hours at 37º C at

a 1:1,000 dilution in complete media. Photoconversion was performed using a digital

micromirror device (DMD) patterned illumination system (Polygon 4000) at 405 nm. For

each experiment, an extra plate was prepared to calibrate the DMD device for a fixed

stimulation ROI of 6x16 pixels. Pre-stimulation images were acquired with using GFP

(pre-conversion mMaple3; 30% laser power, 500 ms) and Cy5 (DNA; 25% laser power,

200 ms), which were used to place the stimulation ROI. Within the Nikon Elements

software, an ND Stimulation protocol was used to stimulate at 405 nm within a user-

defined stimulation ROI (6x16 pixels) and then acquire images across the entire FOV

using TxRed (converted mMaple3; 30% laser, 500 ms). For these experiments, a pre-

conversion image was acquired followed by photoconversion of mMaple3 or mMaple3-

β-catenin constructs using 405 nm light at 100% LED power for 90 seconds. Then the

converted (TxRed) signal was monitored every 30 seconds for 10 minutes post-

conversion. ROIs for nuclei and cytoplasm were generated using GFP (unconverted

mMaple3) and Cy5 (DNA probe) images and applied to the photoconverted movies to

track nuclear and cytoplasmic signal over time. Cytoplasmic tracking data were

exported to excel and normalized for mMaple3 and mMaple3-β-catenin -/+ MG132 at

t=185 seconds. Normalized traces from individual cell cytoplasmic ROIs were imported

to Prism and fit using nonlinear regression comparing one-phase and two-phase decay

models with a constrained Y_o equal to 1.0. Two-phase decay models fit mMaple3-β-

catenin -/+ MG132 curves significantly better and were used to compare mMaple3 and

mMaple3-β-catenin -/+ MG132. Raw cytoplasmic traces for mMaple3-β-catenin under

low, control, high pHi conditions were imported to Prism for visualization and nonlinear

regression. Comparing goodness of fit between one-phase and two-phase decay

models with a constrained Y_o equal to 1.0 showed that two-phase models better

represent curves under low and high pHi conditions. Even though the one-phase model

was preferred for control conditions, the half-life parameter was nonsensically higher

(204-fold higher) whereas the two-phase half-life parameter was only 10% higher

compared to previous experiments. For that reason, one-phase decay models were

used to compare mMaple3-β-catenin degradation under low, control, and high pHi

conditions. Nuclear re-localization was quantified by generating nuclear ROIs based on

SPY650-DNA intensity (see above) and quantifying nuclear intensity over the same time

course used to monitor cytoplasmic degradation, in matched cells. Nuclear intensity was

normalized to t=185 seconds and median traces were imported to Prism and fit using

nonlinear regression and a one-phase association model with a constrained Y_o equal to

1.0 to account for normalization. Overall nuclear accumulation was readout by

comparing plateau values between pHi conditions, where statistical comparison

demonstrated that the same plateau value cannot adequately fit each dataset.

RNA extraction from MDCK cells

MDCK cells were trypsinized and pellets of 1×10^6 cells were frozen at -80º C. Pellets

were thawed on ice and the QIAGEN RNeasy Plus kit was used to extract total RNA.

Yields were quantified using a nanodrop. Extracted RNA was stored at -80º C until used

for reverse transcriptase reaction. cDNA was synthesized using 1000 ng extracted RNA

and the SuperScript IV reverse transcriptase (Thermo Fisher, cat. # 18090010) following

the manufacturer's instructions.

Real-time quantitative polymerase chain reaction (RT-qPCR) of endogenous β-catenin

transcriptional targets

Primers for qPCR (Table 1) were confirmed using standard PCR using DreamTaq

master mix (Thermo Fisher, cat: K1082), 500 ng cDNA, and 500 nM forward and

reverse primers. qPCR was performed on a BioRad CFX Connect using applied

biosystems PowerUp SYBR Green Master Mix (Thermo Fisher Scientific, cat:

100029284). Analysis of MYC and SLUG were performed with 500 nM of each primer

and 10 ng of cDNA and normalized to RPLP2 using the same concentration of primer

and cDNA. Each 20 µL reaction contained 10 µL of SYBR Green, the listed primer and

cDNA concentrations, and diluted with water. The thermocycling conditions were as

follows: pre-denaturation at 95º C for 2 minutes followed by 40 cycles of denaturation at

95º C for 15 sec, annealing at 55º C for 15 sec, and extension at 72º C for 60 sec.

Analysis was performed using the $2^{-\Delta\Delta Cq}$ method (Livak and Schmittgen, 2001). Each

target was analyzed using technical triplicate (n=3) wells and averaged to obtain

biological replicate Cq value relative to the normalizing gene, RPLP2, and each

condition was normalized to control. Five biological replicate (N=5) Cq values were

averaged to calculate the change in expression of each target relative to control.

Table 1: Primer sequences used in RT-qPCR

Target	Sequence 5' → 3'
RPLP2-F	GCTACGTCGCCTCCTACCTG
RPLP2-R	GCTCGCTGATGACCTTGTTG
MYC-F	GGTCTTCGGACACGGAGGAG
MYC-R	TGGACGGACAGGATGTACGC
SLUG-F	AGTGATTATTTCCCCATATCTCTATGA
SLUG-R	GTAGTCTTTCCTCTTCATCACTAATGG

Single-cell β-catenin transcriptional activity assays using live-cell microscopy

Lentiviral-TOP/FOP-dGFP-reporters (wild-type consensus plasmid TOP: Addgene

plasmid #14715; http://n2t.net/addgene:14715 ; RRID:Addgene_14715; inverted

consensus plasmid FOP: Addgene plasmid # 14885 ; http://n2t.net/addgene:14885 ;

RRID:Addgene_14885) were a gift from Tannishtha Reya. MDCK cells were seeded at

$3x10^6$ cells in 10 cm dishes and allowed to adhere to the dish for 8 hours. 8 hours after

seeding, cells were transfected with 0 µg (mock transfection) or 15 µg TOP-dGFP-

reporter/FOP-dGFP-reporter plasmid using Lipofectamine 3000 according to the

manufacturer's instructions. Transfections proceeded for 18 hours prior to the removal

of transfection media, rinsing of the plate with 10 mL of warm DPBS, and addition of 1

mL trypsin. Cells were trypsinized at 37° C for 8-10 minutes. After all the cells were

detached, complete medium was added to the dish to create a single cell suspension.

Using the transfected single cell suspension, 35 mm imaging dishes were seeded at

1.5×10^4 cell per dish and adhered for 8 hours. After the cells adhered for 8 hours, the

plating medium was removed and replaced with complete media used for pHi

manipulations (see section above) or Wnt stimulation (40 ng/mL (Cf) murine WNT3a

(Peprotech; 315-20)). Images were acquired as outlined in the above sections. Cell

nuclei and cell membranes were visualized via Hoechst dye (DAPI; Thermo Scientific,

cat: 62249; 1:10,000) and CellMask Deep Red (Thermo Fisher, C10046; 1:20,000),

respectively, incubated for 15 minutes at 37° C in complete media. Fields of view were

selected by visualizing nuclei (DAPI) and images were collected in the DAPI (20% laser

power, 400 ms), GFP (30% laser power, 500 ms), Cy5 (20% laser power, 400 ms), and

DIC (32.6 DIA, 200 ms) channels. Whole-cell regions of interest (ROIs) were drawn

within individual cells using cell mask as a membrane marker and the average GFP

intensity for individual cells were exported to Excel. Single-cell intensities that were

greater than 2X the average intensity of mock transfected cells was imported to

GraphPad Prism for statistical analysis and visualization. None of the cells expressing

FOP-dGFP that were analyzed met the 2-fold mock cutoff to be included in the final

analysis.

Statistics

GraphPad Prism (10.0) was used to prepare graphs and perform statistical comparisons

as indicated in each figure legend. Normality tests were performed on all datasets after

performing outlier analysis on pooled data (ROUT method; Q=1%), where normally

distributed data is shown with means and non-normally distributed data is shown with

medians. "N" indicates the number of biological replicates performed and "n" represents

the number of technical replicates or individual measurements collected.

Online Supplementary Materials: Online supplementary materials include Figure S1-

S8 and Supplemental Video S1.

Acknowledgements: We would like to thank Tannishtha Reya for TOP/FOP-dGFP-

reporter plasmids (TOP: Addgene plasmid #14715; http://n2t.net/addgene:14715 ;

RRID:Addgene_14715; FOP: Addgene plasmid # 14885 ; http://n2t.net/addgene:14885 ;

RRID:Addgene_14885). We would also like to thank members of the White lab for their

helpful conversations and feedback during figure and manuscript preparation. The

spinning disk confocal microscope used in this work is a part of the Notre Dame

Integrated Imaging Facility (NDIIF).

Author contributions:

Conceptualization: BJC, KAW; Methodology: BJC, KAW; Validation: BJC, KAW; Formal

analysis: BJC, KAW; Investigation: BJC; Resources: K.A.W.; Data collection: BJC; Data

curation: BJC, KAW; Writing – original draft: BJC; Writing - review & editing: BJC, KAW;

Visualization: BJC, KAW; Supervision: BJC, KAW; Project administration: KAW;

Funding acquisition: KAW

Competing Interests: Authors declare no competing interests.

Funding:

This work was supported by an National Science Foundation NSF CAREER Award (

MCB-2238694) and the Henry Luce Foundation. KAW was additionally supported by the

NIH Director's New Innovator Award (1DP2CA260416-01).

References

Amith, S.R., J.M. Wilkinson, and L. Fliegel. 2016. Na+/H+ exchanger NHE1 regulation modulates metastatic potential and epithelial-mesenchymal transition of triple-negative breast cancer cells. *Oncotarget*. 7:21091–21113. doi:10.18632/oncotarget.8520.

Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*. 72:248– 254. doi:10.1016/0003-2697(76)90527-3.

Chen, Y.-T., D.B. Stewart, and W.J. Nelson. 1999. Coupling Assembly of the E-Cadherin/β- Catenin Complex to Efficient Endoplasmic Reticulum Exit and Basal-lateral Membrane Targeting of E-Cadherin in Polarized MDCK Cells. *Journal of Cell Biology*. 144:687– 699. doi:10.1083/jcb.144.4.687.

Cowin, P., H.-P. Kapprell, W.W. Franke, J. Tamkun, and R.O. Hynes. 1986. Plakoglobin: A protein common to different kinds of intercellular adhering junctions. *Cell*. 46:1063– 1073. doi:10.1016/0092-8674(86)90706-3.

Daniels, D.L., and W.I. Weis. 2005. β-catenin directly displaces Groucho/TLE repressors from Tcf/Lef in Wnt-mediated transcription activation | Nature Structural & Molecular Biology. *Nature Structural & Molecular Biology*. 12:364–371.

Denker, S.P., and D.L. Barber. 2002. Cell migration requires both ion translocation and cytoskeletal anchoring by the Na-H exchanger NHE1. *Journal of Cell Biology*. 159:1087–1096. doi:10.1083/jcb.200208050.

Dovmark, T.H., M. Saccomano, A. Hulikova, F. Alves, and P. Swietach. 2017. Connexin-43 channels are a pathway for discharging lactate from glycolytic pancreatic ductal adenocarcinoma cells. *Oncogene*. 36:4538–4550. doi:10.1038/onc.2017.71.

Drees, F., S. Pokutta, S. Yamada, W.J. Nelson, and W.I. Weis. 2005. α-Catenin Is a Molecular Switch that Binds E-Cadherin-β-Catenin and Regulates Actin-Filament Assembly. *Cell*. 123:903–915. doi:10.1016/j.cell.2005.09.021.

Fevr, T., S. Robine, D. Louvard, and J. Huelsken. 2007. Wnt/β-Catenin Is Essential for Intestinal Homeostasis and Maintenance of Intestinal Stem Cells. *Mol Cell Biol*. 27:7551–7559. doi:10.1128/MCB.01034-07.

875   Frantz, C., A. Karydis, P. Nalbant, K.M. Hahn, and D.L. Barber. 2007. Positive feedback
876   between Cdc42 activity and H+ efflux by the Na-H exchanger NHE1 for polarity of
877   migrating cells. *Journal of Cell Biology*. 179:403–410. doi:10.1083/jcb.200704169.

878   Fukunaga, Y., H. Liu, M. Shimizu, S. Komiya, M. Kawasuji, and A. Nagafuchi. 2005. Defining
879   the roles of β-catenin and plakoglobin in cell-cell adhesion: Isolation of β-
880   catenin/plakoglobin-deficient F9 cells. *Cell Structure and Function*. 30:25–34.
881   doi:10.1247/csf.30.25.

882   Grillo-Hill, B.K., B.A. Webb, and D.L. Barber. 2014. Chapter 23 - Ratiometric Imaging of pH
883   Probes. *In* Methods in Cell Biology. J.C. Waters and T. Wittman, editors. Academic
884   Press. 429–448.

885   Hart, M., J.-P. Concordet, I. Lassot, I. Albert, R. del los Santos, H. Durand, C. Perret, B.
886   Rubinfeld, F. Margottin, R. Benarous, and P. Polakis. 1999. The F-box protein β-TrCP
887   associates with phosphorylated β-catenin and regulates its activity in the cell. *Current
888   Biology*. 9:207–211. doi:10.1016/S0960-9822(99)80091-8.

889   He, T.-C., A.B. Sparks, C. Rago, H. Hermeking, L. Zawel, L.T. da Costa, P.J. Morin, B.
890   Vogelstein, and K.W. Kinzler. 1998. Identification of c-MYC as a Target of the APC
891   Pathway. *Science*. 281:1509–1512. doi:10.1126/science.281.5382.1509.

892   Howard, S., T. Deroo, Y. Fujita, and N. Itasaki. 2011. A Positive Role of Cadherin in Wnt/β-
893   Catenin Signalling during Epithelial-Mesenchymal Transition. *PLOS ONE*. 6:e23899.
894   doi:10.1371/journal.pone.0023899.

895   Huber, A.H., D.B. Stewart, D.V. Laurents, W.J. Nelson, and W.I. Weis. 2001. The Cadherin
896   Cytoplasmic Domain Is Unstructured in the Absence of β-Catenin: A POSSIBLE
897   MECHANISM FOR REGULATING CADHERIN TURNOVER *. *Journal of Biological
898   Chemistry*. 276:12301–12309. doi:10.1074/jbc.M010377200.

899   Kao, S.-H., W.-L. Wang, C.-Y. Chen, Y.-L. Chang, Y.-Y. Wu, Y.-T. Wang, S.-P. Wang, A.I.
900   Nesvizhskii, Y.-J. Chen, T.-M. Hong, and P.-C. Yang. 2015. Analysis of Protein Stability
901   by the Cycloheximide Chase Assay. *Bio Protoc*. 5:e1374. doi:10.21769/BioProtoc.1374.

902   Krieghoff, E., J. Behrens, and B. Mayr. 2006. Nucleo-cytoplasmic distribution of β-catenin is
903   regulated by retention. *Journal of Cell Science*. 119:1453–1463. doi:10.1242/jcs.02864.

904   Larsen, A.M., N. Krogsgaard-Larsen, G. Lauritzen, C.W. Olesen, S. Honoré Hansen, E.
905   Boedtkjer, S.F. Pedersen, and L. Bunch. 2012. Gram-Scale Solution-Phase Synthesis of
906   Selective Sodium Bicarbonate Co-transport Inhibitor S0859: in vitro Efficacy Studies in
907   Breast Cancer Cells. *ChemMedChem*. 7:1808–1814. doi:10.1002/cmdc.201200335.

908   Lee, D.H., and A.L. Goldberg. 1996. Selective Inhibitors of the Proteasome-dependent and
909   Vacuolar Pathways of Protein Degradation in Saccharomyces cerevisiae. *Journal of
910   Biological Chemistry*. 271:27280–27284. doi:10.1074/jbc.271.44.27280.

Lewis, J.E., J.K. Wahl, K.M. Sass, P.J. Jensen, K.R. Johnson, and M.J. Wheelock. 1997. Cross-
Talk between Adherens Junctions and Desmosomes Depends on Plakoglobin. *J Cell Biol*.
136:919–934.

Liu, Y., E. Reyes, D. Castillo-Azofeifa, O.D. Klein, T. Nystul, and D.L. Barber. 2023.
Intracellular pH dynamics regulates intestinal stem cell lineage specification. *Nat*
Commun. 14:3745. doi:10.1038/s41467-023-39312-9.

Liu, Y., K.A. White, and D.L. Barber. 2020. Intracellular pH Regulates Cancer and Stem Cell
Behaviors: A Protein Dynamics Perspective. *Frontiers in Oncology*. 10.

Livak, K.J., and T.D. Schmittgen. 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*. 25:402–408.
doi:10.1006/meth.2001.1262.

MacDonald, B.T., K. Tamai, and X. He. 2009. Wnt/β-Catenin Signaling: Components,
Mechanisms, and Diseases. *Developmental Cell*. 17:9–26.
doi:10.1016/j.devcel.2009.06.016.

Martínez-Estrada, O.M., A. Cullerés, F.X. Soriano, H. Peinado, V. Bolós, F.O. Martínez, M.
Reina, A. Cano, M. Fabre, and S. Vilaró. 2006. The transcription factors Slug and Snail
act as repressors of Claudin-1 expression in epithelial cells1. *Biochemical Journal*.
394:449–457. doi:10.1042/BJ20050591.

McEvoy, A.L., H. Hoi, M. Bates, E. Platonova, P.J. Cranfill, M.A. Baird, M.W. Davidson, H.
Ewers, J. Liphardt, and R.E. Campbell. 2012. mMaple: A Photoconvertible Fluorescent
Protein for Use in Multiple Imaging Modalities. *PLOS ONE*. 7:e51314.
doi:10.1371/journal.pone.0051314.

Moon, J.H., S.H. Lee, and Y.C. Lim. 2021. Wnt/β-catenin/Slug pathway contributes to tumor
invasion and lymph node metastasis in head and neck squamous cell carcinoma. *Clin Exp*
Metastasis. 38:163–174. doi:10.1007/s10585-021-10081-3.

Morgan, R.G., J. Ridsdale, M. Payne, K.J. Heesom, M.C. Wilson, A. Davidson, A. Greenhough,
S. Davies, A.C. Williams, A. Blair, M.L. Waterman, A. Tonks, and R.L. Darley. 2019.
LEF-1 drives aberrant β-catenin nuclear localization in myeloid leukemia cells.
Haematologica. 104:1365–1377. doi:10.3324/haematol.2018.202846.

Mukherjee, A., N. Dhar, M. Stathos, D.V. Schaffer, and R.S. Kane. 2018. Understanding How
Wnt Influences Destruction Complex Activity and β-Catenin Dynamics. *iScience*. 6:13–
21. doi:10.1016/j.isci.2018.07.007.

Nusse, R. 2008. Wnt signaling and stem cell control. *Cell Res*. 18:523–527.
doi:10.1038/cr.2008.47.

Nusse, R., and H. Clevers. 2017. Wnt/β-Catenin Signaling, Disease, and Emerging Therapeutic
Modalities. *Cell*. 169:985–999. doi:10.1016/j.cell.2017.05.016.

Perez-Moreno, M., and E. Fuchs. 2006. Catenins: Keeping Cells from Getting Their Signals
Crossed. *Developmental Cell*. 11:601–612. doi:10.1016/j.devcel.2006.10.010.

Reya, T., A.W. Duncan, L. Ailles, J. Domen, D.C. Scherer, K. Willert, L. Hintz, R. Nusse, and
I.L. Weissman. 2003. A role for Wnt signalling in self-renewal of haematopoietic stem
cells. *Nature*. 423:409–414. doi:10.1038/nature01593.

Roura, S., S. Miravet, J. Piedra, A.G. de Herreros, and M. Duñach. 1999. Regulation of E-
cadherin/Catenin Association by Tyrosine Phosphorylation *. *Journal of Biological*
Chemistry. 274:36734–36740. doi:10.1074/jbc.274.51.36734.

Shcherbakova, D.M., and V.V. Verkhusha. 2014. Chromophore chemistry of fluorescent proteins
controlled by light. *Current Opinion in Chemical Biology*. 20:60–68.
doi:10.1016/j.cbpa.2014.04.010.

Simcha, I., M. Shtutman, D. Salomon, J. Zhurinsky, E. Sadot, B. Geiger, and A. Ben-Ze'ev.
1998. Differential nuclear translocation and transactivation potential of β- Catenin and
plakoglobin. *Journal of Cell Biology*. 141:1433–1448. doi:10.1083/jcb.141.6.1433.

Spear, J.S., and K.A. White. 2023. Single-cell intracellular pH dynamics regulate the cell cycle
by timing the G1 exit and G2 transition. *Journal of Cell Science*. 136:jcs260458.
doi:10.1242/jcs.260458.

Tafech, A., P. Jacquet, C. Beaujean, A. Fertin, Y. Usson, and A. Stéphanou. 2023.
Characterization of the Intracellular Acidity Regulation of Brain Tumor Cells and
Consequences for Therapeutic Optimization of Temozolomide. *Biology*. 12:1221.
doi:10.3390/biology12091221.

Tetsu, O., and F. McCormick. 1999. β-Catenin regulates expression of cyclin D1 in colon
carcinoma cells. *Nature*. 398:422–426. doi:10.1038/18884.

Ulmschneider, B., B.K. Grillo-Hill, M. Benitez, D.R. Azimova, D.L. Barber, and T.G. Nystul.
2016. Increased intracellular pH is necessary for adult epithelial and embryonic stem cell
differentiation. *Journal of Cell Biology*. 215:345–355. doi:10.1083/jcb.201606042.

Valenta, T., G. Hausmann, and K. Basler. 2012. The many faces and functions of β-catenin.
EMBO J. 31:2714–2736. doi:10.1038/emboj.2012.150.

Van Der Wal, T., and R. Van Amerongen. 2020. Walking the tight wire between cell adhesion
and WNT signalling: A balancing act for β-catenin: A balancing act for CTNNB1. *Open*
Biology. 10. doi:10.1098/rsob.200267.

Vleminckx, K., R. Kemler, and A. Hecht. 1999. The C-terminal transactivation domain of β-
catenin is necessary and sufficient for signaling by the LEF-1/β-catenin complex in
Xenopus laevis. *Mechanisms of Development*. 81:65–74. doi:10.1016/S0925-
4773(98)00225-1.

Wang, S., J.R. Moffitt, G.T. Dempsey, X.S. Xie, and X. Zhuang. 2014. Characterization and
development of photoactivatable fluorescent proteins for single-molecule-based
superresolution imaging. *Proceedings of the National Academy of Sciences of the United*
States of America. 111:8452–8457. doi:10.1073/pnas.1406593111.

White, K.A., B.K. Grillo-Hill, and D.L. Barber. 2017. Cancer cell behaviors mediated by
dysregulated pH dynamics at a glance. *Journal of Cell Science*. 130:663–669.
doi:10.1242/jcs.195297.

White, Katharine A.; Grillo-Hill, Bree K.; Esquivel, Mario; Peralta, Jobelle; Bui, Vivian N.;
Chire, Ismahan; Barber, D.L. 2018. β-Catenin is a pH sensor with decreased stability at
higher intracellular pH. *Journal of Cell Biology*. 217:3965–3976.
doi:10.1083/jcb.201712041.

Wickline, E.D., P.K. Awuah, J. Behari, M. Ross, D.B. Stolz, and S.P.S. Monga. 2011.
Hepatocyte γ-catenin compensates for conditionally deleted β-catenin at adherens
junctions. *Journal of Hepatology*. 55:1256–1262. doi:10.1016/j.jhep.2011.03.014.

Wodarz, A., and R. Nusse. 1998. Mechanisms of Wnt Signaling in Development. *Annual Review*
of Cell and Developmental Biology. 14:59–88. doi:10.1146/annurev.cellbio.14.1.59.

Yi, H., G. Li, Y. Long, W. Liang, H. Cui, B. Zhang, Y. Tan, Y. Li, L. Shen, D. Deng, Y. Tang,
C. Mao, S. Tian, Y. Cai, Q. Zhu, Y. Hu, W. Chen, and L. Fang. 2020. Integrative multi-
omics analysis of a colon cancer cell line with heterogeneous Wnt activity revealed
RUNX2 as an epigenetic regulator of EMT. *Oncogene*. 39:5152–5164.
doi:10.1038/s41388-020-1351-z.

Yost, C., M. Torres, J.R. Miller, E. Huang, D. Kimelman, and R.T. Moon. 1996. The axis-
inducing activity, stability, and subcellular distribution of beta-catenin is regulated in
Xenopus embryos by glycogen synthase kinase 3. *Genes Dev.* 10:1443–1454.
doi:10.1101/gad.10.12.1443.

Zhang, L., N.G. Gurskaya, E.M. Merzlyak, D.B. Staroverov, N.N. Mudrik, O.N. Samarkina,
L.M. Vinokurov, S. Lukyanov, and K.A. Lukyanov. 2007. Method for real-time
monitoring of protein degradation at the single cell level. *BioTechniques*. 42:446–450.
doi:10.2144/000112453.

Zhou, J., J. Qu, P.Y. Xian, K. Graber, L. Huber, X. Wang, A.M. Gerdes, and F. Li. 2007.
Upregulation of γ-catenin compensates for the loss of β-catenin in adult cardiomyocytes.
American Journal of Physiology - Heart and Circulatory Physiology. 292:270–276.
doi:10.1152/ajpheart.00576.2006.

1015