Autophagy at Crossroads: Modulating Responses to Combined Stresses, Unveiling

Metabolic Shifts and Bacterial Dependencies

Short title: Autophagy: Autophagy at the Nexus of Combined Stresses

Heike Seybold[#]*, Ella Katz, Yoram Soroka and Tamar Avin-Wittenberg*

The Hebrew University of Jerusalem, Department of Plant and Environmental Sciences,

Alexander Silberman Institute of Life Sciences, Edmond Safra Campus, Jerusalem, Israel.

Current address: Federal Research Centre for Cultivated Plants, Julius Kühn-Institut,

Institute for Strategies and Technology Assessment, Kleinmachnow, Germany.

* Corresponding authors

tamar.avin-wittenberg@mail.huji.ac.il

heike.seybold@mail.huji.ac.il

Abstract

Plants face diverse stresses in natural environments, necessitating complex responses for survival. Abiotic and biotic stress responses are typically counteractive, posing challenges for breeding crops resilient to multiple stresses. Autophagy, a cellular transport process, plays a vital role in plant stress response, facilitating the degradation of cellular components and enabling nutrient recycling. Here, we asked what the role of autophagy is in combined abiotic (heat) and biotic (bacterial infection) stress. We introduce a conceptual framework based on assays monitoring autophagy activation, bacterial infection, and metabolic profiling.

Our findings reveal the engagement of members of the autophagy-related 8 (ATG8) protein family in stress-specific activation. Notably, we observed that heat stress facilitates bacterial growth in an autophagy-dependent manner. Metabolic profiling highlights effector-dependent shifts in nutrient availability during stress, influencing bacterial performance. Our study challenges the assumption that combined stresses are simply the sum of individual responses as exemplified by activation of the autophagic pathway. Instead, it establishes autophagy as a pivotal link connecting environmental factors and plant-microbe interactions. Insights for our study can present a novel perspective for designing strategies to enhance crop resilience in the face of multifaceted challenges.

Introduction

Field-grown plants face many stresses, both abiotic (e.g., drought, temperature) and biotic (e.g., pathogens), requiring intricate physiological and molecular strategies for survival. Understanding these strategies has become more urgent due to climate change and the spread of new pests and diseases linked with these changes (Luck et al., 2011; Cavicchioli et al., 2019). Moreover, the nature and severity of stresses encountered by crops are predicted to worsen (Bebber, 2015; Hey et al., 2010). Abiotic and biotic stress responses are typically perceived to be counteractive, thus posing a significant challenge for breeding crops resilient to multiple stresses (Zhang and Sonnewald, 2017). Despite their distinct nature, there are intriguing similarities in the signaling patterns of biotic and abiotic stress responses, such as changes in phytohormone levels, calcium influx, reactive oxygen species (ROS) production, and the involvement of specific transcription factors for stress-induced gene expression (Atkinson and Urwin, 2012; Fraire-Velazquez et al., 2011; Jones and Dangl, 2006; Glazebrook, 2005; Dodds and Rathjen, 2010; Islam et al., 2018; Ohama et al., 2017; Zhu,

2016). Another common feature of these stresses is the redirection of energy resources from growth to stress responses (Signorelli et al., 2019).

Autophagy, a conserved eukaryotic mechanism, is vital for cellular component recycling and energy sourcing (Yang and Klionsky, 2009; Feng et al., 2014; Masclaux-Daubresse et al., 2020; Liu and Bassham, 2012). In plants, three types of autophagy have been described: Micro-, Macro-, and Megautophagy (Liu and Bassham, 2012; van Doorn and Papini, 2013; Marshall and Vierstra, 2018). Macroautophagy (hereafter called autophagy) is extensively studied and involves the formation of cytosolic double-membrane vesicles called autophagosomes. Autophagosomes deliver cytosolic cargo, which can be randomly (bulk autophagy) or selectively acquired (selective autophagy), to the vacuole for degradation (Liu and Bassham, 2012). Though autophagy functions in both plant biotic and abiotic stress response (Signorelli et al., 2019), little is known regarding its role in combined stresses. Here, we focused on the role of autophagy in the combination of heat stress (abiotic stress) and bacterial infection (biotic stress).

Autophagy progression is divided into distinct steps and, in plants, involves more than 40 autophagy-related (ATG) proteins (Liu and Bassham, 2012; Yang and Klionsky, 2010; Chen and Klionsky, 2011; Chung, 2019; Wang et al., 2021). *ATG* genes are regulated on a transcriptional level, and *ATG* transcript availability and timing are critical for an appropriate stress response (Magen et al., 2022). Plant transcription factors were shown to confer abiotic stress resistance by inducing *ATG* gene expression and, consequently, autophagy (Wang et al., 2015; Zhu et al., 2018). The WRKY transcription factor family is particularly interesting regarding combined stress. WRKY33 induces autophagy in response to both abiotic and biotic stresses by activating the transcription of *ATG5* and *ATG7*. It is involved in heat tolerance in different plant species and, in addition to *ATG* genes, induces the expression of defense-related genes in *Arabidopsis thaliana* (Arabidopsis) (Lai et al., 2011; Schmidt et al., 2020; Mao et al., 2011; Lippok et al., 2007; Zhou et al., 2014; Zheng et al., 2006). Arabidopsis knockout mutants of *ATG* genes (*atg* mutants) are hypersensitive to different stresses, while overexpression of *ATG* genes improves plant fitness and performance (Liu and Bassham, 2012; Slavikova et al., 2008; Minina et al., 2018; Magen et al., 2022; Zhang et al., 2021a). ATG proteins critical for stress response include ATG5, ATG7, and the selective autophagy receptor NBR1.

Heat can have cytotoxic effects resulting in cellular metabolism and homeostasis perturbations due to protein denaturation and aggregation. Plant autophagy can remove heat-induced protein aggregates (Baxter et al., 2014; Zhou et al., 2013; Zhou et al., 2014). In

addition, NBR1-mediated selective autophagy was shown to play a crucial role in the recovery from heat stress by regulating heat stress memory (Thirumalaikumar et al., 2021). NBR1 recruits cargo to the autophagosome by interaction with ATG8, a small ubiquitin-like protein, which undergoes lipidation by Phosphatidylethanolamine upon autophagy induction (Svenning et al., 2011; Zhou et al., 2013). The inner autophagosomal membrane, associated proteins (ATG8, NBR1), and cargo undergo degradation within the vacuole. The number of ATG8 isoforms varies among plant species, expanding from one isoform in algae to multiple isoforms in vascular plants (Zess et al., 2019). Notably, ATG8 exhibits a non-canonical function in response to heat stress, aiding the Golgi apparatus reassembly in Arabidopsis after acute heat stress (Zhou et al., 2023).

The role of autophagy in biotic stress response is more complicated and depends on the pathogen's lifestyle (Sertsuvalkul et al., 2022). Bacterial plant pathogens, residing extracellularly, employ a type 3 secretion system (T3SS) to deliver effectors into the plant cell, manipulating cellular processes, including autophagy, for their benefit (Üstün et al., 2016; Lal et al., 2020; Leong et al., 2022b). NBR1-mediated selective autophagy can antagonize the effector-driven virulence (Leong et al., 2022b; Üstün et al., 2018). Besides bacterial pathogens, the oomycete *Phytophthora infestans* inhibits autophagy through effectors, countered by NBR1-mediated selective autophagy (Pandey et al., 2021; Dagdas et al., 2016; Dagdas et al., 2018). In addition, overexpression of *ATG5* or *ATG7* increased resistance against the necrotrophic fungal pathogen *Alternaria brassicicola*, and other examples support a positive role for autophagy in defense responses against necrotrophic plant pathogens (Minina et al., 2018; Lai et al., 2011; Lenz et al., 2011; Patel and Dinesh-Kumar, 2008). In line with these findings, autophagy plays a negative role in defense responses against obligate biotrophic pathogens (Wang et al., 2011). In contrast to oomycetes and bacterial pathogens, plant viruses, present intracellularly, are removed directly through NBR1-dependent selective autophagy (Hafrén et al., 2017; Hafrén et al., 2018).

Despite the significance of combined stresses in real-world natural and agricultural settings, research on their impact, particularly involving autophagy, remains limited. Understanding how plants respond to combined stresses is crucial for ensuring crop fitness and productivity in the face of climate change. The interaction between plant stress responses can be synergistic. Still, in most cases, responses to combined stresses differ significantly from those to single stresses (Rasmussen et al., 2013; Pandey et al., 2015; Ramegowda and Senthil-Kumar, 2015; Zhang and Sonnewald, 2017). Plant autophagy as a pathway involved in various stress responses has an exciting potential to increase plant tolerance towards

combined stress scenarios with broad agricultural implications. Here, we investigate the pivotal role of autophagy in orchestrating the response to combined stresses. We chose a dual stress paradigm involving elevated temperature followed by infection with the bacterial pathogen *Xanthomonas campestris* pv. *vesicatoria* (*Xcv*), the causal agent of bacterial spot disease on pepper and tomato. Autophagy was shown to contribute to plant defense against *Xanthomonas* pathogens, while *Xcv* targets autophagy to inhibit appropriate defense responses (Yan et al., 2017; Leong et al., 2022b).

Results

Stress-specific activation of ATG8 isoforms

Plants face diverse stresses requiring specific responses, with autophagy playing a crucial role (Avin-Wittenberg, 2019; Sertsuvalkul et al., 2022). Understanding stress-specific activation of autophagy remains largely unclear. We focused on the ATG8 protein family, which functions in target recognition and cargo binding. We speculated that ATG8 isoforms participate in stress-specific autophagy, which is yet to be determined (Kellner et al., 2017). Degradation of GFP-ATG8 fusion proteins in the vacuole and release of free GFP was investigated as a proxy of autophagic flux (Klionsky et al., 2021). We investigated the seven *St*ATG8 isoforms in potato (*Solanum tuberosum*), exploring their roles in stress-induced autophagy.

GFP-tagged versions of all seven *St*ATG8 isoforms were transiently expressed in *Nicotiana benthamiana*. The plants were then subjected to different autophagy-inducing stress conditions (darkness, heat, *Xcv* infection) (Avin-Wittenberg et al., 2015; Leong et al., 2022b; Zhou et al., 2014). Prolonged darkness leads to carbon starvation, widely known to induce autophagy, presumably in a non-selective manner (Barros et al., 2017). In contrast, pathogen infection or extreme temperatures require more specific plant responses. Carbon starvation was induced by three days of darkness, leading to increased degradation of all GFP-*St*ATG8 isoforms and high levels of free GFP (Fig1A). Heat stress reduced protein levels of all GFP-*St*ATG8 isoforms after 6 h and 8 h of 40°C (Fig1B). However, ATG8 degradation differed in timing and intensity. While some isoforms responded quickly (after 6 h), others showed a slower reduction upon stress (after 8 h). GFP-*St*ATG8-3.1 and GFP-*St*ATG8-4 showed a quick, strong, and lasting decrease in protein levels.

We next tested the effect of infection with *Xcv*. Although *N. benthamiana* is not a natural host of *Xcv*, the *Nicotiana benthamiana* mutant line Δ*roq1* (Grützner et al., 2021) can be infected by the bacteria. Only a subset of GFP-*St*ATG8 isoforms was degraded upon *Xcv* challenge

(Fig1C). The GFP-*St*ATG8-2 isoforms, GFP-*St*ATG8-3.1, and in some but not all experimental replicates, GFP-*St*ATG8-4 displayed increased degradation upon *Xcv* infection. GFP-*St*ATG8-1 isoforms showed the opposite effect. Interestingly, the *Xcv*-specific activation of GFP-*St*ATG8 isoforms was independent of bacterial effectors. Infection with the T3SS mutant *Xcv* Δ*hrcV* (hereafter called *hrcV*), lacking the effector secretion system, resulted in a GFP-ATG8 degradation pattern similar to that of *Xcv*. Surprisingly, the detected degradation band was not free GFP, unlike observed for carbon starvation. These results indicate a specific autophagic response to stress, particularly to *Xcv*/bacterial infection, unlike the non-specific reaction to general stressors like carbon starvation.

Differential autophagy activation under combined stress compared to single stresses

As *St*ATG8-3.1 was degraded upon heat stress as well as after inoculation with *Xcv*, we were interested in the degradation of this isoform in response to a combined heat-*Xcv* challenge. As performed in the single-stress experiments, GFP-*St*ATG8-3.1 was transiently expressed in *N. benthamiana* Δ*roq1* plants. The plants were then subjected to 6 h heat stress or 6 h *Xcv* infection. In addition, 6 h heat stress was followed by subsequent 6 h *Xcv* or a 6 h recovery period without *Xcv* inoculation (Fig2A). Although *Xcv* inoculation was the second and, therefore, more recent treatment, GFP-*St*ATG8-3.1 degradation after combined stress treatment resembled heat stress instead of *Xcv* infection (Fig2B).

ATG genes are transcriptionally regulated upon stress. We tested native transcript levels in tobacco plants subjected to single and combined stresses. *NbATG7* mRNA levels were reduced after 6 h heat stress and 6 h *Xcv*, respectively. They were further reduced after combined heat-*Xcv* stress. Unfortunately, this trend, though visibly striking, was not statistically significant (FigS1). In contrast, *NbJOKA2* (the *NBR1* homolog in solanaceous species) showed no transcriptional effect on the single or combined stress. These findings affirm that combined stresses surpass the sum of individual stress effects, highlighting autophagy's role in responding to such combined stress.

Heat stress facilitates bacterial growth in an autophagy-dependent manner

Since the degradation pattern of GFP-*St*ATG8-3.1 after combined heat-*Xcv* stress showed no signs of response to the bacterial infection, we next asked how a preceding heat stress affects bacterial infection. We syringe-infiltrated *Xcv* into *N. benthamiana* Δ*roq1* as well as its natural host tomato (*Solanum lycopersicum*) M82 plants. Bacterial titers were checked 3 days post-

infection (dpi) (Fig3A). *Xcv* grew better in heat-stressed plants in both species (Fig3B and C).

This beneficial effect of preceding heat stress on bacterial growth was dependent on

autophagy. In the presence of the autophagy inhibitor 3-Methyadenine (3-MA), preceding

heat stress did not affect bacterial growth. In addition, the T3SS mutant *hrcV* did not benefit

from previous heat stress in plants (Fig3D), making the increase in bacterial growth after heat

stress dependent on bacterial effector delivery. Since *NbATG7* showed a transcriptional

reduction after combined heat-*Xcv* stress (FigS1), we tested bacterial performance with and

without preceding heat stress in plants transiently expressing *AtATG7*. In plants transiently

overexpressing *AtATG7*, *Xcv* grew similarly in heat-stressed and unstressed plants (Fig3E).

These results confirm our previously observed reduced immune response to bacterial infection

in heat-stressed plants. The results also confirm that autophagy is involved in the response to

combined stress.

Differential *At*ATG protein degradation upon combined stress is autophagy-dependent

To test how changes in autophagic activity affect the response to combined stresses, we

changed the plant system and used available *ATG* overexpression and knockout Arabidopsis

lines (Minina et al., 2018). Since Arabidopsis is not a host for *Xcv*, we used the bacterial

elicitor flg22, a 22 amino acid peptide of the bacterial protein Flagellin, known to induce

autophagy (Rodriguez et al., 2020). Transiently expressed *St*ATG8-3.1 showed a reduced

response to the biotic stress (*Xcv*) when preceded by heat stress (Fig2B). To test the

conservation of this response, we checked GFP-*At*ATG8a degradation in stable

overexpression lines treated with heat stress, biotic stress (flg22), and a stress combination

(Fig4A). As seen for GFP-*St*ATG8-3.1 in *N. benthamiana*, *At*ATG8a was degraded in

Arabidopsis to a similar level in the combined stress scenario and the single heat stress

(Fig4B).

We next examined the response of other ATG proteins following combined stress.

Overexpressed *At*NBR1-TAP was degraded differentially after heat stress and flg22 (Fig4C).

While heat stress led to an accumulation of *At*NBR1, flg22 treatment reduced *At*NBR1 protein

levels. After combined stress, *At*NBR1-TAP band intensities were mid-range between heat

and flg22 stress (Fig4C). In wildtype plants, protein levels of native *At*ATG7 were lowest

under combined stress (Fig4D). This effect was more pronounced in *AtATG5* and *AtNBR1*-

TAP overexpression lines and absent in *atg5-1* plants. The results suggest that *At*ATG7 is

degraded in an autophagy-dependent manner upon combined stress. Taken together, our

results suggest a dynamic involvement of different components of the autophagy pathway in

response to combined stress. Depending on the component, combined stress either mimics or

intensifies the reaction to one of the individual stresses or shows an intermediate response

between the single stresses.

Autophagy affects *FRK1* gene expression under single and combined stresses

Exploring the impact of autophagic activity on stress response transcription, we assessed the

expression of stress marker genes in stable Arabidopsis *ATG* overexpression and knockout

lines under single and combined stress conditions. Despite highly consistent results across two

independent experiments, the lack of significance may be attributed to substantial biological

replicate variation. In wildtype plants, the flg22 marker gene *AtFRK1* (At2g19190) increased

1 h after flg22 treatment, as expected (Winter et al., 2007)(http://bar.utoronto.ca/efp/cgi-

bin/efpWeb.cgi) and showed a pronounced increase in samples that received the combined

stress (FigS2B). *AtFRK1* expression increased strongly upon flg22 treatment in lines

overexpressing GFP-*At*ATG8a, *At*ATG5, and *At*NBR1-TAP. Expression did not increase

further or was even reduced in samples treated with the heat-flg22 combination (FigS2B).

Overexpression of *At*ATG7, however, did not lead to changes in *AtFRK1* gene expression

compared to the wildtype. All knockout lines in this experiment displayed a wildtype-like

AtFRK1 gene expression pattern (FigS2C). Hence, the biotic stress marker gene expression

proved to be autophagy-independent but responsive to increased autophagic activity. The heat

stress marker gene *AtHSP70* (At3g12580) was equally expressed in wildtype samples that

received heat stress and samples treated with the heat-flg22 combination (FigS2D). In

contrast, all overexpression lines responded to heat stress with highly increased *HSP70*

expression, while combined heat-flg22 stress did not lead to an increase in *HSP70* expression

(FigS2D). This effect was, however, also observed in *atg5-1* and *nbr1* lines (FigS2E),

highlighting the importance of autophagy in maintaining *HSP70* expression under combined

stress.

Heat stress recovery fails to halt bacterial growth

247 The Arabidopsis experiments underscored autophagy's pivotal role in combating combined

248 stress, suggesting benefits from heightened autophagic activity in ATG overexpression lines.

249 Given the limitations of Arabidopsis for *Xcv* experiments, we reverted to the *N. benthamiana*

250 Δ*roq1* system. We hypothesized that increased bacterial growth in previously heat-stressed

251 plants (see Fig3B) resulted from the exhaustion of autophagic resources for defense against

Xcv. The autophagic capacity, diverted to handle heat-induced damage, could not defend against bacterial effector proteins. To test our hypothesis, we repeated bacterial growth experiments with heat-stressed plants undergoing a 24-h recovery phase (Fig5A). We expected restored autophagic capacity after a 24-h recovery following a mild 6-h heat stress. However, comparable to results without a recovery period, previous heat treatment resulted in increased bacterial growth even after the recovery period (unpaired t-test p value 0.0067), although not significantly different in the ANOVA (Fig5B). As seen without the recovery period, this effect was again autophagy-dependent. These results suggest that the cause for increased bacterial growth was not pre-occupation of the autophagy machinery, or dependent on effects lasting over 24 h.

Metabolic shift caused by *Xcv* is treatment-resilient

Recognizing autophagy's role in plant nutrient availability (Signorelli et al., 2019), we theorized that its activation during heat stress might reshape metabolic content to favor bacterial growth. We performed metabolic profiling via GC-MS on previously *N. benthamiana* heat-stressed leaf tissue infected with bacteria. We concentrated on primary metabolites, representing nutrients potentially utilized by the bacteria. Principal component analysis (PCA) showed a major shift between non-treated plants (red) and samples from heat-treated plants (yellow) (Fig5C). *Xcv*-inoculation led to another shift. All samples that received *Xcv*-treatment clustered together independent of additional treatments. The complete analysis also included samples from *hrcV*-infected plants. Interestingly, some *hrcV*-treated samples clustered away from the *Xcv*-treated samples in the PCA, pointing to an effector-dependent shift of the metabolome (FigS3).

Generally, heat-treated plants as well as *Xcv*-infected plants, showed strong differences from control plants for most identified compounds (Fig5D). In particular, sugars and amino acids were reduced under heat stress and after infection, respectively. In contrast, the sugar alcohol Myo-inositol increased, specifically after heat challenge (see below). The phytohormone Abscisic acid (ABA), which is involved in signaling after many abiotic stresses, including heat (Larkindale and Knight, 2002; Zhu, 2016), increased especially under heat-*Xcv* combinations. Collectively, our findings indicate significant stress-induced metabolic shifts, with stress-specific patterns yet common alterations in energy sources. Notably, the response to *Xcv* infection exhibited remarkable resilience to additional factors or stresses.

Changes in organic acids are independent of autophagy

We next focused on specific metabolic changes induced by stress combinations and their dependence on autophagy. Two compounds of the TCA cycle, succinic acid and malic acid, were reduced only upon heat treatment but not infection (FigS4E,F). However, while levels of malic acid stayed low in all plants that previously received heat treatment, *Xcv* but not *hrcV* infection restored control levels of succinic acid in heat-stressed plants in an autophagy-independent manner. The levels of a third detected member of the TCA cycle, citric acid, did not change in response to any stress treatment (FigS4G). No changes in shikimic acid levels were detected in response to heat stress, but levels were reduced significantly in response to infection or a combined heat-pathogen stress (FigS4A). This response to infection was independent of autophagy and similar in *Xcv* and *hrcV*. Nicotinic acid showed a mild reduction under combined stress compared to the single stresses, which was significantly lower than in control plants (FigS4B). Again, the response to combined stress was independent of autophagy. Together, our results suggest that changes in organic acids in response to stress combinations are not dependent on autophagy.

Shift in amino acid levels is independent of bacterial effectors

We also investigated metabolites functioning as energy sources. Amino acids are a major energy source that can be mobilized under stress (Signorelli et al., 2019). We detected three amino acids, which were generally reduced upon bacterial infection as well as heat stress, especially in the case of glycine (Fig6A,B). *Xcv* and *hrcV* showed a similar effect. Asparagine was reduced upon bacterial infection and further reduced in the presence of 3-MA under control conditions (Fig6C). This effect was also observed in heat-stressed plants but to a lesser extent. Thus, the shifts in amino acid levels are not a result of bacterial manipulation but rather a general response of the plant to stress conditions. Urea levels, a metabolite directly linked to amino acid catabolism (Witte, 2011), stayed stable upon heat challenge but were reduced upon *Xcv* infection, independent of temperature (Fig6D). Interestingly, *hrcV* treatment significantly increased urea levels in an autophagy-dependent and heat-sensitive manner.

Effector-dependent reduction of sugars is heat-sensitive and modulated by autophagy

Sugars are a main energy source under stress conditions (Signorelli et al., 2019). Most of the mono- and disaccharides were globally decreasing upon heat- and *Xcv*-treatment compared to

control samples, with sucrose showing only a decrease upon infection, not heat (Fig7A-C, E-G). This effect was not influenced by 3-MA treatment. Samples treated with *hrcV* displayed a more minor sugar reduction than *Xcv* infection (significant only for Glucose). However, when autophagy was inhibited by 3-MA treatment, *hrcV* infection reduced sugar levels to a similar extent as after *Xcv* infection. The effect of *Xcv* treatment was not further intensified by 3-MA treatment, as seen for *hrcV*. We, therefore, suggest that the decrease in sugar levels upon infection is partially dependent on effector(s) targeting autophagy. Interestingly, the effect observed for *hrcV* was only present in control conditions and absent in heat-stressed plants where *Xcv* and *hrcV* infection had similar, autophagy-independent effects.

The only detected Monosaccharide that did not fit into the described pattern is Xylose/Lyxose (Fig7D). Levels of this sugar were not reduced upon *Xcv* infection or heat treatment but only upon inhibition of autophagy, pointing to a role of autophagy in maintaining Xylose/Lyxose levels under stress conditions.

Myo-inositol did not change upon *Xcv* infection (Fig7H). However, levels increased significantly upon *hrcV*-treatment in an autophagy-dependent manner. Heat stress led to a general increase in Myo-inositol levels, a well-documented response to heat and other abiotic stresses in different plant species (Khurana et al., 2017; Sharma et al., 2020; Boominathan et al., 2004). Combined heat-*Xcv* but not heat-*hrcV* stress reduced these levels significantly in an autophagy-dependent manner (ANOVA performed in the absence of control *hrcV* samples). The massive shifting of sugars is, at least partially, the result of an effector-mediated, heat-sensitive inhibition of autophagy by *Xcv*. Taken together, our detailed metabolic profiling showed a strong effect of the applied stress (combinations) on metabolites mainly involved in primary (energy) metabolism. For most sugars and urea, the impact of *Xcv* infection was effector-dependent and absent in *hrcV*. This effector-dependent effect was mimicked in *hrcV* by inhibition of autophagy.

Discussion

The simultaneous occurrence of environmental stresses, such as heat, with pathogen infection poses a widespread threat to plant health. It is crucial to comprehend how plants navigate these dual challenges to enhance crop breeding initiatives aimed at creating robust varieties. In this study, we highlight autophagy as a fundamental mechanism that bestows stress-specific capabilities upon plants. Bacterial pathogens target autophagy to promote disease, and this influence is evident in metabolic patterns during combined heat-pathogen stress.

ATG8 confers specificity to combined plant stress responses

Autophagy can selectively remove damaged or unwanted cellular components. The selectivity is conferred by target binding to autophagy receptors, which, in turn, bind ATG8 (Stephani and Dagdas, 2020). Thus, although many conditions induce autophagy, the autophagic cargo and the resulting degradome are expected to be unique to each condition. Since plants have multiple ATG8 isoforms, it has been proposed that these isoforms play a crucial role in shaping the stress-specific response and selectivity of autophagy (Kellner et al., 2017; Lin et al., 2023; Zess et al., 2019). There are two main methods to study autophagy activation using GFP-ATG8. The GFP tag enables autophagosome visualization by confocal microscopy. In addition, the ratio between GFP-ATG8 and free GFP by Western blot allows us to estimate ongoing autophagic flux (Klionsky et al., 2021). We could not detect free GFP in all our *N. benthamiana* Western blots. In fact, we often and repeatedly only detected an unknown but GFP-specific degradation band. Similar bands were reported in other plant species (Zhou et al., 2023), yet their existence was not explained. This novel band is clearly connected to the degradation of GFP-*St*ATG8, and is absent without GFP-*St*ATG8 protein. Therefore, we decided to use this degradation band instead of free GFP to calculate the ratio between its intensity and GFP-*St*ATG8.

In our study, the seven StATG8 isoforms exhibited stress-specific responses based on the type of stress (Fig1). Carbon starvation triggered a uniform activation of all isoforms, whereas *Xcv* infection or heat differed in ATG8 activation pattern. Bacterial infection is hampered by removing effector proteins via autophagy (Leong et al., 2022b). The process requires selective autophagy receptors e.g., NBR1, which exhibits a preference for specific ATG8 isoforms (Svenning et al., 2011). Accordingly, only selected *St*ATG8 isoforms were activated by *Xcv* infection. By contrast, under heat stress, *St*ATG8 isoforms differed in activation timing. Heat stress generally causes protein aggregation, but proteins vary in their ability to withstand elevated temperatures. We suggest *St*ATG8 isoforms are activated at different times during heat stress to account for the variability in protein heat sensitivity. One of the unique structural features of ATG8 proteins is an N-terminal helix domain (Bu et al., 2020). This domain determines the binding specificity to ATG8-interacting proteins and its sequence is not conserved in plants (Zess et al., 2019). Thus, preferential binding of ATG8 isoforms to different targets and selective autophagy receptors seems only logical (Kellner et al., 2017). Besides differential activation of ATG8 proteins, expression of *AtATG8* genes differed when

exposed to starvation or (a)biotic stress (Wang et al., 2020; Chiu et al., 2023; Leong et al., 2022b), further confirming our finding of stress-specific involvement of *St*ATG8 isoforms.

Response to combined abiotic-biotic stress is very complex, involving the interaction of two living organisms (the plant and the pathogen) as well as the additional level of abiotic stress (Kissoudis et al., 2014). Subjecting the plants to heat while concurrently exposing them to *Xcv* implies that both the plants and the bacteria are experiencing above-optimum temperatures. We tried to reduce the level of complexity by applying the combined heat-*Xcv* stress as a consecutive stress where the bacteria are not experiencing direct heat stress. Studies investigating combined stresses usually focus on a single time point; however, we believe that our experimental system of a consecutive stress setup reflects a natural situation more accurately where stress may also occur sequentially rather than simultaneously.

GFP-*St*ATG8-3.1 was involved in the response to heat stress and *Xcv* infection (Fig1). Yet, when exposed to a combined heat-*Xcv* challenge, the activation did not mimic a combination of both stresses (Fig2). Hence, the assumption that combined stresses equal the sum of the involved single stresses falls short of the complexity of natural systems. There is a possibility that combined stress activates another suite of *St*ATG8 isoforms than the single stresses. However, degradation of GFP-*At*ATG8a after a combined heat-flg22 treatment showed similar behavior with degradation mimicking heat treatment, not flg22 treatment (Fig4). Infection assays with *Xcv* further confirmed the observation made for GFP-*St*ATG8-3.1 after combined heat-*Xcv* stress. The bacteria grew better in previously heat-stressed plants (Fig3). The effect was absent in *hrcV* infections and when autophagy was inhibited, confirming the results of GFP-*St*ATG8-3.1 activation during the heat-*Xcv* stress combination. Enhanced growth of another bacterial phytopathogen, *Pseudomonas Pst* DC300, at moderately elevated temperature during infection has been described before (Huot et al., 2017). As in our case, the increased growth depended on the delivery of bacterial effector proteins. We can now link this enhanced growth to bacterial manipulation of autophagy.

Autophagy components other than ATG8 fine-tune the response to combined stress

While ATG8 contributes to stress-specificity, monitoring other ATGs can give a more detailed insight into the activation of autophagy during combined stress. The degradation of *At*NBR1 and *At*ATG7 proteins in response to single and combined stresses gives a more complex picture than the degradation of ATG8 proteins. In contrast to GFP-*St*ATG8s in tobacco, *At*NBR1-TAP degradation in Arabidopsis in response to combined heat-flg22 stress

showed an intermediate response compared to the two single stresses (Fig4). Although a role for NBR1 was documented in immunity against various pathogens, including bacteria (Dagdas et al., 2016; Üstün et al., 2018; Hafrén et al., 2017), it is possibly not involved in the autophagic response activated by flg22 treatment. flg22 is recognized by the immune receptor kinase FLAGELLIN-SENSING 2 (FLS2) (Chinchilla et al., 2006). FLS2 is known to be recycled, and in Arabidopsis, orosomucoid (ORM) proteins were described as selective autophagy receptors specifically targeting FLS2, but not other immune receptors (Beck et al., 2012; Yang et al., 2019). Nevertheless, *At*NBR1-TAP was reduced upon flg22 challenge compared to the control.

In contrast, overexpressed *At*NBR1-TAP accumulated upon heat treatment. Although to a lesser extent than in *atg* mutants, accumulation of overexpressed *At*NBR1-TAP proteins has been documented after 6 to 9 h of severe heat stress in Arabidopsis (Zhou et al., 2013), similar to what we observed. This accumulation vanished when the heat stress was followed by flg22 challenge, contradicting the results from GFP-*St*ATG8-3.1 where combined stress mimicked the heat stress (Fig2). Interestingly, in the same *At*NBR1-TAP overexpression samples, both flg22 and combined heat-flg22 treatment led to an autophagy-dependent reduction of *At*ATG7 protein levels, which was not observed after heat treatment.

A similar pattern of *At*ATG7 protein was found in wildtype plants and an *At*ATG5 overexpression line (Fig4). The reduction of *At*ATG7 protein levels after flg22 and combined stress matches the results obtained for *NbATG7* mRNA levels (FigS1). However, we did not see an accumulation of *NbATG7* mRNA after heat stress as we did for *At*ATG7 protein, making it unlikely that the effects observed on the protein level result from transcriptional changes. We postulate that the changes in *At*ATG7 protein were caused by modulated protein degradation. While this degradation depends on autophagy, it is possible that *At*ATG7 was degraded via other pathways (e.g. the proteasome) as shown for SH3P2 upon *Xcv* infection. Here, the pathogen blocks autophagy by increasing SH3P2 degradation to enhance its pathogenicity (Leong et al., 2022b). Though our results aline with enhanced degradation of ATG proteins by bacterial manipulation, it is important to note that the response we observed was for heat and a bacterial elicitor. We cannot exclude that combining heat stress and infection with an actual pathogen might yield a different response. Still, our observations underline the importance of monitoring not only one but several components of the autophagic pathway to get a clearer picture of the highly dynamic involvement of autophagy in combined stress response.

Besides the degradation of ATG proteins, transcriptional induction of *ATG* genes and selective

autophagy receptors plays a key role in the regulation of autophagy (Magen et al., 2022;

Hachez et al., 2014; Stephani and Dagdas, 2020). *NbATG7* mRNA levels were downregulated

in response to both moderate heat stress of 40 °C and *Xcv* infection alone and the combination

(FigS1). Regulation of *ATG*s can differ drastically depending on the plant species and severity

of the heat stress treatment (Zhou et al., 2014; Zhou et al., 2013). In contrast, *NbJOKA2*

mRNA levels remained relatively stable (FigS1), although some instances of responsiveness

to heat stress have been reported for *NBR1* (Zhou et al., 2013; Zhou et al., 2014). Studies on

autophagy and pathogen defense also show contrasting effects. While *AtATGs* are induced by

necrotrophic fungus *B. cinerea* and contribute to resistance, autophagy has a negative role in

defense against biotrophic pathogens (Lai et al., 2011; Minina et al., 2018; Wang et al., 2011;

Üstün et al., 2018; Leong et al., 2022a). Inhibition of *ATG* gene expression could be an

essential strategy for bacterial pathogens to avoid autophagy-dependent immune responses.

However, it remains unclear why heat also leads to the downregulation of *NbATG7*

transcription, emphasizing caution when comparing heat stress of varying intensities across

plant species with differing tolerance to high temperatures. Besides the plant species, timing is

important when studying stress-related transcriptional regulation. Studies, including ours,

often focus on one or a few time points, missing the dynamic regulation of transcription.

We also examined the expression of stress marker genes like *HSP70* for heat stress and *FRK1*

for biotic stress upon combined stress (FigS2). *HSP70* was upregulated in wildtype upon

single and combined heat-pathogen stress. In lines with altered autophagic activity, this

response shifted towards a stronger response to the single stress and a sharply reduced

response to combined heat-pathogen stress, underlining autophagy's role in *HSP70*

transcriptional regulation during combined stress. In contrast, *FRK1* mRNA levels were

induced by autophagy. Notably, *FRK1* gene expression does not depend on autophagy, as

evidenced by wildtype-like induction upon flg22 perception in *atg* mutant plants. Similar

results were observed for the Arabidopsis defensin gene, *AtPDF1.2* (Lai et al., 2011). This

robust response is essential as pathogens evolved to undermine components of the immune

response network (Tsuda et al., 2013). It is also of special interest when it comes to combined

stress scenarios.

Despite reproducibility, the lack of significant differences was disappointing in the large-scale

RT-qPCR experiment involving up to eight lines, five treatments, and a minimum of five

biological replicates per treatment. The procedure was carefully controlled, including chamber

transfers for temperature treatment and manual syringe infiltration for flg22 treatment.

However, potential changes in light quality and humidity between the experimental chamber and the cultivation room might have influenced marker gene expression. Additionally, the time-consuming tasks of infiltrating numerous plants and harvesting two leaves per plant made precise 1-hour (60 min) sample collection impractical, resulting in a more realistic time range of 55 to 75 mins. Considering the dynamic nature of *FRK1* gene expression (Winter et al., 2007)(http://bar.utoronto.ca/efp/cgi-bin/efpWeb.cgi), with a 44-fold increase at 1 hour and more than a 90-fold increase at 4 hours after flg22 treatment, variations in marker gene expression may be attributed, at least partially, to these factors.

Changes in metabolites as an underestimated effect of autophagy influencing bacterial performance

Nutrient availability plays a major role in pathogen growth. We thus asked whether previous heat stress affected nutrient availability for the invading pathogen. Nutrients in the apoplast available for bacterial nutrition are mainly sugars and amino acids (Rico and Preston, 2008; Fatima and Senthil-Kumar, 2015). Sugars and amino acids were generally reduced in the tested leaf tissue as a result of heat stress and also after infection. This observation is not surprising, given that stress responses are energetically costly (Signorelli et al., 2019). While we see this reduction in amino acids upon *Xcv* infection (Fig6), we cannot confirm it results from bacterial manipulation of autophagy or bacterial consumption of these metabolites. Urea, linked to amino acid catabolism (Witte, 2011), displayed an opposite response to stress. While amino acid levels decreased upon infection, heat stress alone had no major effect on urea levels, and *Xcv* infection or combined stress even slightly reduced urea levels (Fig6). Interestingly, this reduction resulted from bacterial manipulation as levels increased significantly following *hrcV* infection. More importantly, this increase was absent when *hrcV* infection was combined with autophagy inhibition. Amino acids are recycled from proteins degraded via autophagy.

In addition, components targeted by bacterial effectors to manipulate metabolite recycling via autophagy might be heat-sensitive. When *hrcV* infection followed heat stress, urea levels were reduced, resembling *Xcv* infection. The absence of increased urea in combined *hrcV*-heat stress can also be explained on a more general immune response level. Salicylic acid (SA) is the dominant phytohormone involved in the plant's defense against (hemi)biotrophic pathogens. This SA response is heat sensitive in Arabidopsis and tobacco (Wang et al., 2009). Loss of pathogen-induced SA was described as a central mechanism underlying enhanced

bacterial growth of *Pseudomonas Pst* DC300 at elevated temperatures (Huot et al., 2017). Without an energy-intensive immune response, amino acids would not be channeled into energy production, failing to increase urea levels.

Xcv infection at least partially manipulates sugar levels in an autophagy-dependent manner, which seems to be mimicked by heat stress (Fig7). During infection, phytopathogens can adjust their metabolic machinery towards the most abundant nutrients in the host. *Xcv* was shown to modulate its manner of active nutrient acquisition (Fatima and Senthil-Kumar, 2015). So-called TBDTs (TonB-dependent transporters) are involved in uptake of carbohydrates, with genomes of several *Xanthomonas* species having a high copy number of TBDT genes (Blanvillain et al., 2007; Dugé de Bernonville et al., 2014; Pieretti et al., 2012). Moreover, pathogens like *Xcv*, colonizing the apoplast, release effector proteins to manipulate plant cells and optimize nutrient availability, e.g. by facilitating sugar efflux through induction of gene expression of sugar transporters (Chen et al., 2010; Eom et al., 2015; Zhou et al., 2015).

In addition to effectors influencing nutrient efflux, pathogens target autophagy directly (Üstün et al., 2016; Lal et al., 2020; Leong et al., 2022a). The reduction of sugar levels is partially dependent on effector proteins, as the *hrcV* mutant was unable to reduce sugar levels comparable to *Xcv*. Inhibition of autophagy by 3-MA led to *Xcv*-like sugar levels after *hrcV* infection, suggesting that the reduced sugar levels after Xcv infection resulted from autophagy inhibition by bacterial effectors. The autophagy-dependent manipulation of sugar levels is no longer observable when *hrcV* infection follows heat stress. Since heat stress affects autophagic activity (Zhou et al., 2014) and thereby also metabolite recycling, manipulation of autophagy might be obsolete under these conditions. Another interpretation is that, as mentioned above, the effector targets are heat sensitive, and this manipulation is no longer needed when the infection follows a heat stress. However, the effect of *Xcv* on sugar levels does not solely depend on autophagy, as autophagy inhibition by 3-MA did not alter the *Xcv* sugar phenotype. We thus postulate that manipulation of autophagy is not the only mechanism underlying pathogen success. This observation also explains why *Xcv* growth benefits from preceding heat stress whereas *hrcV* growth does not (Fig3). While the *hrcV* sugar phenotype can be compensated by preceding heat stress, the *hrcV* growth phenotype is unaffected, presumably because there are other effectors contributing to pathogen success than those targeting autophagy. As *hrcV* can not secrete any effectors to the plant, future studies using more specific effector mutants might be able to discern the particular effects of bacterial effectors in autophagy-dependent nutrient availability.

Our findings underscore the specificity inherent in the autophagic response, particularly
regarding ATG8 isoforms. We demonstrate that combined stresses do not equal the sum of
their counterparts, specifically focusing on autophagy at the nexus of simultaneous
challenges. It will be interesting to see if the specificity of the autophagic response, especially
in the case of stress combinations, also translates to the autophagic cargo level. The influence
of heat stress on autophagy, seen in its facilitation of bacterial growth, establishes it as a key
link between environmental factors and plant-microbial interactions. This connection
underscores the broader ecological implications of autophagy in plant defense and unveils
potential targets for enhancing stress resilience. As autophagy functions in many aspects of
plant life, it is not possible to distinguish its roles in nutrient supply, abiotic stress, and
pathogen defense. As our results show, it is possible that other, still unknown factors
contribute to autophagy-dependent bacterial growth following heat stress.

Materials and Methods

Biological material and growth conditions

All *Arabidopsis thaliana* plants used in this study were of the Columbia ecotype (Col-0). *A.*
thaliana lines overexpressing GFP-*AtATG8a* in Col-0, *atg5-1*, *atg7-2*, *AtATG5* OE and
AtATG7 OE backgrounds were kindly provided by Elena Minina (Swedish University of
Agricultural Sciences, Sweden; (Minina et al., 2018)). *At*NBR1-TAP OE and *nbr1* CRISPR
lines were kindly provided by Agnieszka Sirko (Polish Academy of Sciences, Poland;
(Tarnowski et al., 2020)). *Nicotiana benthamiana* mutant line Δ*roq1* was kindly provided by
Johannes Stuttmann (CEA, France; (Grützner et al., 2021)). Transient expression constructs
were kindly provided by Yasin Dagdas (Gregor Mendel Institute, Austria; GFP-*StATG8*
isoforms) and Elena Minina (Swedish University of Agricultural Sciences, Sweden; *AtATG7*).
Xanthomonas campestris pv. *vesicatoria* (*Xcv*) strains 85-10 and 85-10 Δ*hrcV* were kindly
provided by Guido Sessa (Tel Aviv University, Israel).
Solanum lycopersicum M82 and *N. benthamiana* were grown under a 16h light/8h darkness
cycle (long day, LD), 22 °C. Tomato plants were used for stress experiments at an age of five
weeks, tobacco plants were transformed and subsequently stressed at an age of approximately
four weeks. *A. thaliana* plants were grown under 8h light/16h darkness cycle (short day, SD),
22°C. Six-week-old plants were used stress experiments. All plant were grown in a mixture of
commercial plant soil, perlit, and vermiculite (2:1:1).

Transient expression in tobacco

To study the stress specificity of ATG8 isoforms, *Agrobacterium tumefaciens* strain GV3101::pMP90 was transformed by electroporation with constructs of N-terminally GFP-tagged versions of all seven *St*ATG8 isoforms (Zess et al., 2019), respectively. For transient expression in *N. benthamiana Δroq1* plants, bacteria were grown in LB medium at 28 °C overnight, resuspended in infiltration buffer (10 mM MES, pH 5.7; 10 mM $MgCl_2$; 150 μM Acetosyringone), adjusted to OD_{600} 0.5 and incubated for 2 hours at room temperature. The bacteria were subsequently pressure-infiltrated into *N. benthamiana* leaf tissue using a needle-less syringe. After an initial phase of 18 h under LD conditions for expression of constructs, the plants were subjected to the respective stress conditions.

Stress treatments

To induce carbon starvation, plant trays were covered with aluminum foil for three days but kept in the same climate chamber as control plants without carbon starvation. For biotic stress treatments, *Xcv* suspensions of OD_{600} 0.2 in 10 mM $MgCl_2$ were pressure-infiltrated into leaf tissue of *N. benthamiana* using a needle-less syringe. Alternatively, 200 nM flg22 peptide was pressure-infiltrated into leaf tissue of *A. thaliana*. Plants were placed into a Percival chamber at 22 °C (control temperature) for 6 h (tobacco) or 1 h (Arabidopsis), respectively. For heat stress treatments, tobacco and tomato plants were placed into a second Percival chamber and subjected to 40 °C for 6 h while Arabidopsis were incubated at 38 °C for 3 h.

Protein extraction, SDS-PAGE, and Western blot

Equal amounts of plant leaf tissue were ground in liquid nitrogen using metal beads. Total proteins were extracted using a protein extraction buffer (50 mM Tris, pH 8; 20 mM NaCl; 10 % glycerol; 0.5 % Triton X-100; 1x Protease Inhibitor). Samples were mixed thoroughly, incubated on ice for 10 min, and centrifuged (14 000 rpm, 10 min, 4 °C). The supernatant was collected, mixed with SDS-PAGE sample buffer, and incubated at 80 °C for 5 min. Total proteins were separated by SDS-PAGE and transferred to a polyvinylidene difluoride membrane. Equal loading was validated using Coomassie Brilliant Blue staining of the membrane before the remaining unspecific protein binding sites were blocked with blocking solution (3 % milk + 2 % BSA in 1x PBST). For immunodetection, primary antibodies (anti-GFP: Abcam ab290; anti-NBR1: Agrisera AS142805; anti-ATG7: Agrisera AS194277) from rabbit were used in combination with a secondary HRP-conjugated anti-rabbit antibody from goat. For repeated detection of proteins from one membrane, membranes were first dried and then re-wetted with 1x PBST before Coomassie Brilliant Blue staining. The membranes were

stripped by incubation in stripping buffer (200 mM glycine; 3.5 mM SDS; 1 % Tween 20;

pH 2.2) for 20 min and subsequent washing with 1x PBST (3x for 5 min). Membranes were

re-blocked with blocking solution prior to re-probing with the new antibody.

Bacterial growth assay

Xcv was grown in NYGB medium at 28 °C overnight. For measuring bacterial growth, leaves

of five-week old tomato plants or four-week old tobacco plants were pressure-infiltrated with

Xcv OD_{600} 0.0004 in 10 mM $MgCl_2$ with or without 5 mM 3-Methyladenine (3-MA) using a

needle-less syringe. Leaf disk samples were taken immediately after infiltration to ensure

equal infiltration densities. To check bacterial performance, leaf disk samples were taken three

days after inoculation. Bacterial growth was monitored by serial dilution plating of ground

leaf discs.

Gene expression analysis by quantitative reverse transcription polymerase chain

reaction (RT-qPCR)

To analyze transcript abundance, total RNA was extracted from leaf tissue using the TRIzol

method. Total RNA was treated with TURBO DNA-free kit (Invitrogen). The integrity of the

RNA was checked on 1% (w/v) agarose gels and the concentration was measured using a

Nanodrop spectrophotometer. Intron-specific primers (see Table S1) were used to exclude

contamination with genomic DNA. Total RNA was reverse transcribed with SensiFAST

cDNA Synthesis Kit (Bioline) according to the manufacturer's recommendations. Real-time

quantitative PCR analysis was performed in a final volume of 10 μl in 96-well plates using

the CFXDuet system (Bio-Rad). Amplification specificity was evaluated by post-

amplification dissociation curves. *ACTIN* was used as the internal control for quantification of

gene expression. Primer sequences are listed in Table S1.

GC-MS analysis

Plant metabolites from *N. benthamiana* were extracted following an established protocol for

GC-MS-based metabolite profiling (Lisec et al., 2006), with minor modifications. In short, the

samples (25 mg leaf tissue per sample with six biological replicates per treatment) were

extracted in methanol containing an internal standard (Pentaerythritol, stock concentration

0.6 mg/ml in DDW, final concentration 0.016 mg/ml), shaken for 15 min at 70 °C, and

centrifuged at 14 000 rpm for 10 min. The supernatant was transferred to new tubes, and

chloroform and double-distilled water were added. The mixture was centrifuged at

14 000 rpm for 15 min, and aliquots of 200 μL of the upper phase of each sample were

transferred to new tubes and dried by Speedvac, overnight. Derivatization was carried out as

previously described (Lisec et al., 2006). Polar metabolites were measured by the Agilent

7200B GC/Q-TOF. The injection and separation procedures were performed according to

Dahtt using the DB-35MS column (Dhatt et al., 2019). Metabolite detection and annotation

were performed by Quant software (Agilent, Santa Clara, CA, USA), according to an accurate

mass library of known markers generated by our group and run in the same system. Following

blank subtraction, the peak area of each metabolite was normalized to the internal standard

(i.e., Pentaerythritol) in each sample, as well as to the fresh weight of the respective sample.

For more information see Table S2.

Statistical analysis and data visualization

Data were statistically examined using analysis of variance (Ordinary one-way ANOVA) and

tested for significant differences using unpaired *t* tests using GraphPad Prism 8.0.1 software.

Graphs and heatmaps were generated using GraphPad Prism 8.0.1 while Principal component

analysis (PCA) of the metabolome was performed using Metaboanalyst, an open-source, web-

based tool for metabolomics data analysis (https://www.metaboanalyst.ca/). Western blots

were processed to maximize readers accessibility to the data. Original blots can be found in

the supplemental data (FigS5).

Acknowledgments

We thank the Rachel Green and Omri Finkel groups at HUJI for assistance with heatable

phytochambers and Elena Minina, Agnieszka Sirko, Johannes Stuttmann, Yasin Dagdas and

Guido Sessa for sharing transgenic seeds, bacterial strains and plasmids. Fig. 2A, 3A, 4A and

5A were created with Biorender (https://www.biorender.com). The project was funded by a

Walter-Benjamin Fellowship to HS by Deutsche Forschungsgemeinschaft (Projektnummer

446674370).

Author Contributions

678 HS and TAW designed the research; HS, EK, YS and TAW performed research experiments;

679 HS, YS and TAW analyzed data; HS and TAW wrote the manuscript with input from all

680 authors.

References

Atkinson, N.J., and Urwin, P.E. (2012). The interaction of plant biotic and abiotic stresses: From genes to the field. Journal of experimental botany **63** (10): 3523–3543.

Avin-Wittenberg, T. (2019). Autophagy and its role in plant abiotic stress management. Plant, cell & environment **42** (3): 1045–1053.

Avin-Wittenberg, T., Bajdzienko, K., Wittenberg, G., Alseekh, S., Tohge, T., Bock, R., Giavalisco, P., and Fernie, A.R. (2015). Global analysis of the role of autophagy in cellular metabolism and energy homeostasis in Arabidopsis seedlings under carbon starvation. The Plant cell **27** (2): 306–322.

Barros, J.A.S., Cavalcanti, J.H.F., Medeiros, D.B., Nunes-Nesi, A., Avin-Wittenberg, T., Fernie, A.R., and Araújo, W.L. (2017). Autophagy Deficiency Compromises Alternative Pathways of Respiration following Energy Deprivation in Arabidopsis thaliana. Plant physiology **175** (1): 62–76.

Baxter, A., Mittler, R., and Suzuki, N. (2014). ROS as key players in plant stress signalling. Journal of experimental botany **65** (5): 1229–1240.

Bebber, D.P. (2015). Range-expanding pests and pathogens in a warming world. Annual review of phytopathology **53**: 335–356.

Beck, M., Zhou, J., Faulkner, C., MacLean, D., and Robatzek, S. (2012). Spatio-temporal cellular dynamics of the Arabidopsis flagellin receptor reveal activation status-dependent endosomal sorting. The Plant cell **24** (10): 4205–4219.

Blanvillain, S., Meyer, D., Boulanger, A., Lautier, M., Guynet, C., Denancé, N., Vasse, J., Lauber, E., and Arlat, M. (2007). Plant carbohydrate scavenging through tonB-dependent receptors: A feature shared by phytopathogenic and aquatic bacteria. PloS one **2** (2): e224.

Boominathan, P., Shukla, R., Kumar, A., Manna, D., Negi, D., Verma, P.K., and Chattopadhyay, D. (2004). Long term transcript accumulation during the development of dehydration adaptation in Cicer arietinum. Plant physiology **135** (3): 1608–1620.

Bu, F., Yang, M., Guo, X., Huang, W., and Chen, L. (2020). Multiple Functions of ATG8 Family Proteins in Plant Autophagy. Frontiers in cell and developmental biology **8**: 466.

Cavicchioli, R., Ripple, W.J., Timmis, K.N., Azam, F., and Bakken, L.R., et al. (2019). Scientists' warning to humanity: Microorganisms and climate change. Nature reviews. Microbiology **17** (9): 569–586.

Chen, L.-Q., Hou, B.-H., Lalonde, S., Takanaga, H., Hartung, M.L., Qu, X.-Q., Guo, W.-J., Kim, J.-G., Underwood, W., Chaudhuri, B., Chermak, D., Antony, G., White, F.F.,

Somerville, S.C., Mudgett, M.B., and Frommer, W.B. (2010). Sugar transporters for
intercellular exchange and nutrition of pathogens. Nature **468** (7323): 527–532.

Chen, Y., and Klionsky, D.J. (2011). The regulation of autophagy - unanswered questions.
Journal of cell science **124** (Pt 2): 161–170.

Chinchilla, D., Bauer, Z., Regenass, M., Boller, T., and Felix, G. (2006). The Arabidopsis
receptor kinase FLS2 binds flg22 and determines the specificity of flagellin perception.
The Plant cell **18** (2): 465–476.

Chiu, C.-Y., Lung, H.-F., Chou, W.-C., Lin, L.-Y., Chow, H.-X., Kuo, Y.-H., Chien, P.-S.,
Chiou, T.-J., and Liu, T.-Y. (2023). Autophagy-Mediated Phosphate Homeostasis in
Arabidopsis Involves Modulation of Phosphate Transporters. Plant & cell physiology **64**
(5): 519–535.

Chung, T. (2019). How phosphoinositides shape autophagy in plant cells. Plant science an
international journal of experimental plant biology **281**: 146–158.

Dagdas, Y.F., Belhaj, K., Maqbool, A., Chaparro-Garcia, A., Pandey, P., Petre, B.,
Tabassum, N., Cruz-Mireles, N., Hughes, R.K., Sklenar, J., Win, J., Menke, F.,
Findlay, K., Banfield, M.J., Kamoun, S., and Bozkurt, T.O. (2016). An effector of the
Irish potato famine pathogen antagonizes a host autophagy cargo receptor. eLife **5**.

Dagdas, Y.F., Pandey, P., Tumtas, Y., Sanguankiattichai, N., Belhaj, K., Duggan, C.,
Leary, A.Y., Segretin, M.E., Contreras, M.P., Savage, Z., Khandare, V.S., Kamoun, S.,
and Bozkurt, T.O. (2018). Host autophagy machinery is diverted to the pathogen
interface to mediate focal defense responses against the Irish potato famine pathogen.
eLife **7**.

Dhatt, B.K., Abshire, N., Paul, P., Hasanthika, K., Sandhu, J., Zhang, Q., Obata, T., and
Walia, H. (2019). Metabolic Dynamics of Developing Rice Seeds Under High Night-
Time Temperature Stress. Frontiers in plant science **10**: 1443.

Dodds, P.N., and Rathjen, J.P. (2010). Plant immunity: Towards an integrated view of plant-
pathogen interactions. Nature reviews. Genetics **11** (8): 539–548.

Dubiella, U., Seybold, H., Durian, G., Komander, E., Lassig, R., Witte, C.-P., Schulze,
W.X., and Romeis, T. (2013). Calcium-dependent protein kinase/NADPH oxidase
activation circuit is required for rapid defense signal propagation. Proceedings of the
National Academy of Sciences of the United States of America **110** (21): 8744–8749.

Dugé de Bernonville, T., Noël, L.D., SanCristobal, M., Danoun, S., Becker, A., Soreau, P.,
Arlat, M., and Lauber, E. (2014). Transcriptional reprogramming and phenotypical

changes associated with growth of Xanthomonas campestris pv. campestris in cabbage

xylem sap. FEMS microbiology ecology **89** (3): 527–541.

Eom, J.-S., Chen, L.-Q., Sosso, D., Julius, B.T., Lin, I.W., Qu, X.-Q., Braun, D.M., and

Frommer, W.B. (2015). SWEETs, transporters for intracellular and intercellular sugar

translocation. Current opinion in plant biology **25**: 53–62.

Fatima, U., and Senthil-Kumar, M. (2015). Plant and pathogen nutrient acquisition

strategies. Frontiers in plant science **6**: 750.

Feng, Y., He, D., Yao, Z., and Klionsky, D.J. (2014). The machinery of macroautophagy.

Cell research **24** (1): 24–41.

Fraire-Velazquez, S., Rodriguez-Guerra, R., and Sanchez-Caldero, L. (2011). Abiotic and

Biotic Stress Response Crosstalk in Plants. In Abiotic Stress Response in Plants -

Physiological, Biochemical and Genetic Perspectives, A. Shanker, ed (InTech).

Glazebrook, J. (2005). Contrasting mechanisms of defense against biotrophic and

necrotrophic pathogens. Annual review of phytopathology **43**: 205–227.

Grützner, R., Martin, P., Horn, C., Mortensen, S., Cram, E.J., Lee-Parsons, C.W.T.,

Stuttmann, J., and Marillonnet, S. (2021). High-efficiency genome editing in plants

mediated by a Cas9 gene containing multiple introns. Plant communications **2** (2):

100135.

Guerra, T., Schilling, S., Hake, K., Gorzolka, K., Sylvester, F.-P., Conrads, B.,

Westermann, B., and Romeis, T. (2020). Calcium-dependent protein kinase 5 links

calcium signaling with N-hydroxy-l-pipecolic acid- and SARD1-dependent immune

memory in systemic acquired resistance. The New phytologist **225** (1): 310–325.

Hachez, C., Veljanovski, V., Reinhardt, H., Guillaumot, D., Vanhee, C., Chaumont, F.,

and Batoko, H. (2014). The Arabidopsis abiotic stress-induced TSPO-related protein

reduces cell-surface expression of the aquaporin PIP2;7 through protein-protein

interactions and autophagic degradation. The Plant cell **26** (12): 4974–4990.

Hafrén, A., Macia, J.-L., Love, A.J., Milner, J.J., Drucker, M., and Hofius, D. (2017).

Selective autophagy limits cauliflower mosaic virus infection by NBR1-mediated

targeting of viral capsid protein and particles. Proceedings of the National Academy of

Sciences of the United States of America **114** (10): E2026-E2035.

Hafrén, A., Üstün, S., Hochmuth, A., Svenning, S., Johansen, T., and Hofius, D. (2018).

Turnip Mosaic Virus Counteracts Selective Autophagy of the Viral Silencing Suppressor

HCpro. Plant physiology **176** (1): 649–662.

Hey, S.J., Byrne, E., and Halford, N.G. (2010). The interface between metabolic and stress
signalling. Annals of botany **105** (2): 197–203.

Huot, B., Castroverde, C.D.M., Velásquez, A.C., Hubbard, E., Pulman, J.A., Yao, J.,
Childs, K.L., Tsuda, K., Montgomery, B.L., and He, S.Y. (2017). Dual impact of
elevated temperature on plant defence and bacterial virulence in Arabidopsis. Nature
communications **8** (1): 1808.

Islam, M.R., Feng, B., Chen, T., Tao, L., and Fu, G. (2018). Role of Abscisic Acid in
Thermal Acclimation of Plants. J. Plant Biol. **61** (5): 255–264.

Jones, J.D.G., and Dangl, J.L. (2006). The plant immune system. Nature **444** (7117): 323–
329.

Kellner, R., La Concepcion, J.C. de, **Maqbool, A., Kamoun, S., and Dagdas, Y.F.** (2017).
ATG8 Expansion: A Driver of Selective Autophagy Diversification? Trends in plant
science **22** (3): 204–214.

Khurana, N., Sharma, N., and Khurana, P. (2017). Overexpression of a heat stress
inducible, wheat myo-inositol-1-phosphate synthase 2 (TaMIPS2) confers tolerance to
various abiotic stresses in Arabidopsis thaliana. Agri Gene **6**: 24–30.

Kissoudis, C., van de Wiel, C., Visser, R.G.F., and van der Linden, G. (2014). Enhancing
crop resilience to combined abiotic and biotic stress through the dissection of
physiological and molecular crosstalk. Frontiers in plant science **5**: 207.

Klionsky, D.J., Abdel-Aziz, A.K., Abdelfatah, S., Abdellatif, M., and Abdoli, A., et al.
(2021). Guidelines for the use and interpretation of assays for monitoring autophagy (4[th]
edition)1. Autophagy **17** (1): 1–382.

Lai, Z., Wang, F., Zheng, Z., Fan, B., and Chen, Z. (2011). A critical role of autophagy in
plant resistance to necrotrophic fungal pathogens. The Plant journal for cell and molecular
biology **66** (6): 953–968.

Lal, N.K., Thanasuwat, B., Huang, P.-J., Cavanaugh, K.A., Carter, A., Michelmore,
R.W., and Dinesh-Kumar, S.P. (2020). Phytopathogen Effectors Use Multiple
Mechanisms to Manipulate Plant Autophagy. Cell host & microbe **28** (4): 558-571.e6.

Larkindale, J., and Knight, M.R. (2002). Protection against heat stress-induced oxidative
damage in Arabidopsis involves calcium, abscisic acid, ethylene, and salicylic acid. Plant
physiology **128** (2): 682–695.

Lenz, H.D., Haller, E., Melzer, E., Kober, K., Wurster, K., Stahl, M., Bassham, D.C.,
Vierstra, R.D., Parker, J.E., Bautor, J., Molina, A., Escudero, V., Shindo, T., van der
Hoorn, R.A.L., Gust, A.A., and Nürnberger, T. (2011). Autophagy differentially

controls plant basal immunity to biotrophic and necrotrophic pathogens. The Plant journal for cell and molecular biology **66** (5): 818–830.

Leong, J.X., Langin, G., and Üstün, S. (2022a). Selective autophagy: Adding precision in plant immunity. Essays in biochemistry **66** (2): 189–206.

Leong, J.X., Raffeiner, M., Spinti, D., Langin, G., Franz-Wachtel, M., Guzman, A.R., Kim, J.-G., Pandey, P., Minina, A.E., Macek, B., Hafrén, A., Bozkurt, T.O., Mudgett, M.B., Börnke, F., Hofius, D., and Üstün, S. (2022b). A bacterial effector counteracts host autophagy by promoting degradation of an autophagy component. The EMBO journal **41** (13): e110352.

Lin, L.-Y., Chow, H.-X., Chen, C.-H., Mitsuda, N., Chou, W.-C., and Liu, T.-Y. (2023). Role of autophagy-related proteins ATG8f and ATG8h in the maintenance of autophagic activity in Arabidopsis roots under phosphate starvation. Frontiers in plant science **14**: 1018984.

Lippok, B., Birkenbihl, R.P., Rivory, G., Brümmer, J., Schmelzer, E., Logemann, E., and Somssich, I.E. (2007). Expression of AtWRKY33 encoding a pathogen- or PAMP-responsive WRKY transcription factor is regulated by a composite DNA motif containing W box elements. Molecular plant-microbe interactions MPMI **20** (4): 420–429.

Lisec, J., Schauer, N., Kopka, J., Willmitzer, L., and Fernie, A.R. (2006). Gas chromatography mass spectrometry-based metabolite profiling in plants. Nature protocols **1** (1): 387–396.

Liu, Y., and Bassham, D.C. (2012). Autophagy: Pathways for self-eating in plant cells. Annual review of plant biology **63**: 215–237.

Luck, J., Spackman, M., Freeman, A., Tre¿bicki, P., Griffiths, W., Finlay, K., and Chakraborty, S. (2011). Climate change and diseases of food crops. Plant Pathology **60** (1): 113–121.

Magen, S., Seybold, H., Laloum, D., and Avin-Wittenberg, T. (2022). Metabolism and autophagy in plants-a perfect match. FEBS letters **596** (17): 2133–2151.

Mao, G., Meng, X., Liu, Y., Zheng, Z., Chen, Z., and Zhang, S. (2011). Phosphorylation of a WRKY transcription factor by two pathogen-responsive MAPKs drives phytoalexin biosynthesis in Arabidopsis. The Plant cell **23** (4): 1639–1653.

Marshall, R.S., and Vierstra, R.D. (2018). Autophagy: The Master of Bulk and Selective Recycling. Annual review of plant biology **69**: 173–208.

Masclaux-Daubresse, C., d'Andrea, S., Bouchez, I., and Cacas, J.-L. (2020). Reserve lipids and plant autophagy. Journal of experimental botany **71** (10): 2854–2861.

Minina, E.A., Moschou, P.N., Vetukuri, R.R., Sanchez-Vera, V., Cardoso, C., Liu, Q.,
Elander, P.H., Dalman, K., Beganovic, M., Lindberg Yilmaz, J., Marmon, S.,
Shabala, L., Suarez, M.F., Ljung, K., Novák, O., Shabala, S., Stymne, S., Hofius, D.,
and Bozhkov, P.V. (2018). Transcriptional stimulation of rate-limiting components of the
autophagic pathway improves plant fitness. Journal of experimental botany **69** (6): 1415–
1432.

Ohama, N., Sato, H., Shinozaki, K., and Yamaguchi-Shinozaki, K. (2017). Transcriptional
Regulatory Network of Plant Heat Stress Response. Trends in plant science **22** (1): 53–65.

Pandey, P., Leary, A.Y., Tumtas, Y., Savage, Z., Dagvadorj, B., Duggan, C., Yuen, E.L.,
Sanguankiattichai, N., Tan, E., Khandare, V., Connerton, A.J., Yunusov, T.,
Madalinski, M., Mirkin, F.G., Schornack, S., Dagdas, Y., Kamoun, S., and Bozkurt,
T.O. (2021). An oomycete effector subverts host vesicle trafficking to channel starvation-
induced autophagy to the pathogen interface. eLife **10**.

Pandey, P., Ramegowda, V., and Senthil-Kumar, M. (2015). Shared and unique responses
of plants to multiple individual stresses and stress combinations: Physiological and
molecular mechanisms. Frontiers in plant science **6**: 723.

Patel, S., and Dinesh-Kumar, S.P. (2008). Arabidopsis ATG6 is required to limit the
pathogen-associated cell death response. Autophagy **4** (1): 20–27.

Pieretti, I., Royer, M., Barbe, V., Carrere, S., Koebnik, R., Couloux, A., Darrasse, A.,
Gouzy, J., Jacques, M.-A., Lauber, E., Manceau, C., Mangenot, S., Poussier, S.,
Segurens, B., Szurek, B., Verdier, V., Arlat, M., Gabriel, D.W., Rott, P., and
Cociancich, S. (2012). Genomic insights into strategies used by Xanthomonas albilineans
with its reduced artillery to spread within sugarcane xylem vessels. BMC genomics **13**:
658.

Ramegowda, V., and Senthil-Kumar, M. (2015). The interactive effects of simultaneous
biotic and abiotic stresses on plants: Mechanistic understanding from drought and
pathogen combination. Journal of plant physiology **176**: 47–54.

Rasmussen, S., Barah, P., Suarez-Rodriguez, M.C., Bressendorff, S., Friis, P.,
Costantino, P., Bones, A.M., Nielsen, H.B., and Mundy, J. (2013). Transcriptome
responses to combinations of stresses in Arabidopsis. Plant physiology **161** (4): 1783–
1794.

Rico, A., and Preston, G.M. (2008). Pseudomonas syringae pv. tomato DC3000 uses
constitutive and apoplast-induced nutrient assimilation pathways to catabolize nutrients

that are abundant in the tomato apoplast. Molecular plant-microbe interactions MPMI **21**

(2): 269–282.

Rodriguez, E., Chevalier, J., Olsen, J., Ansbøl, J., Kapousidou, V., Zuo, Z., Svenning, S.,

Loefke, C., Koemeda, S., Drozdowskyj, P.S., Jez, J., Durnberger, G., Kuenzl, F.,

Schutzbier, M., Mechtler, K., Ebstrup, E.N., Lolle, S., Dagdas, Y., and Petersen, M.

(2020). Autophagy mediates temporary reprogramming and dedifferentiation in plant

somatic cells. The EMBO journal **39** (4): e103315.

Schmidt, A., Mächtel, R., Ammon, A., Engelsdorf, T., Schmitz, J., Maurino, V.G., and

Voll, L.M. (2020). Reactive oxygen species dosage in Arabidopsis chloroplasts can

improve resistance towards Colletotrichum higginsianum by the induction of WRKY33.

The New phytologist **226** (1): 189–204.

Sertsuvalkul, N., DeMell, A., and Dinesh-Kumar, S.P. (2022). The complex roles of

autophagy in plant immunity. FEBS letters **596** (17): 2163–2171.

Sharma, N., Chaudhary, C., and Khurana, P. (2020). Wheat Myo-inositol phosphate

synthase influences plant growth and stress responses via ethylene mediated signaling.

Scientific reports **10** (1): 10766.

Signorelli, S., Tarkowski, Ł.P., van den Ende, W., and Bassham, D.C. (2019). Linking

Autophagy to Abiotic and Biotic Stress Responses. Trends in plant science **24** (5): 413–

430.

Slavikova, S., Ufaz, S., Avin-Wittenberg, T., Levanony, H., and Galili, G. (2008). An

autophagy-associated Atg8 protein is involved in the responses of Arabidopsis seedlings to

hormonal controls and abiotic stresses. Journal of experimental botany **59** (14): 4029–

4043.

Stephani, M., and Dagdas, Y. (2020). Plant Selective Autophagy-Still an Uncharted Territory

With a Lot of Hidden Gems. Journal of molecular biology **432** (1): 63–79.

Sung, D.Y., Vierling, E., and Guy, C.L. (2001). Comprehensive expression profile analysis

of the Arabidopsis Hsp70 gene family. Plant physiology **126** (2): 789–800.

Svenning, S., Lamark, T., Krause, K., and Johansen, T. (2011). Plant NBR1 is a selective

autophagy substrate and a functional hybrid of the mammalian autophagic adapters NBR1

and p62/SQSTM1. Autophagy **7** (9): 993–1010.

Tarnowski, L., Rodriguez, M.C., Brzywczy, J., Piecho-Kabacik, M., Krčkova, Z.,

Martinec, J., Wawrzynska, A., and Sirko, A. (2020). A selective autophagy cargo

receptor NBR1 modulates abscisic acid signalling in Arabidopsis thaliana. Scientific

reports **10** (1): 7778.

Thirumalaikumar, V.P., Gorka, M., Schulz, K., Masclaux-Daubresse, C., Sampathkumar, A., Skirycz, A., Vierstra, R.D., and Balazadeh, S. (2021). Selective autophagy regulates heat stress memory in Arabidopsis by NBR1-mediated targeting of HSP90.1 and ROF1. Autophagy **17** (9): 2184–2199.

Tsuda, K., Mine, A., Bethke, G., Igarashi, D., Botanga, C.J., Tsuda, Y., Glazebrook, J., Sato, M., and Katagiri, F. (2013). Dual regulation of gene expression mediated by extended MAPK activation and salicylic acid contributes to robust innate immunity in Arabidopsis thaliana. PLoS genetics **9** (12): e1004015.

Üstün, S., Hafrén, A., Liu, Q., Marshall, R.S., Minina, E.A., Bozhkov, P.V., Vierstra, R.D., and Hofius, D. (2018). Bacteria Exploit Autophagy for Proteasome Degradation and Enhanced Virulence in Plants. The Plant cell **30** (3): 668–685.

Üstün, S., Sheikh, A., Gimenez-Ibanez, S., Jones, A., Ntoukakis, V., and Börnke, F. (2016). The Proteasome Acts as a Hub for Plant Immunity and Is Targeted by Pseudomonas Type III Effectors. Plant physiology **172** (3): 1941–1958.

van Doorn, W.G., and Papini, A. (2013). Ultrastructure of autophagy in plant cells: A review. Autophagy **9** (12): 1922–1936.

Wang, P., Nolan, T.M., Yin, Y., and Bassham, D.C. (2020). Identification of transcription factors that regulate ATG8 expression and autophagy in Arabidopsis. Autophagy **16** (1): 123–139.

Wang, P., Wang, T., Han, J., Li, M., Zhao, Y., Su, T., and Ma, C. (2021). Plant Autophagy: An Intricate Process Controlled by Various Signaling Pathways. Frontiers in plant science **12**: 754982.

Wang, Y., Bao, Z., Zhu, Y., and Hua, J. (2009). Analysis of temperature modulation of plant defense against biotrophic microbes. Molecular plant-microbe interactions MPMI **22** (5): 498–506.

Wang, Y., Cai, S., Yin, L., Shi, K., Xia, X., Zhou, Y., Yu, J., and Zhou, J. (2015). Tomato HsfA1a plays a critical role in plant drought tolerance by activating ATG genes and inducing autophagy. Autophagy **11** (11): 2033–2047.

Wang, Y., Nishimura, M.T., Zhao, T., and Tang, D. (2011). ATG2, an autophagy-related protein, negatively affects powdery mildew resistance and mildew-induced cell death in Arabidopsis. The Plant journal for cell and molecular biology **68** (1): 74–87.

Winter, D., Vinegar, B., Nahal, H., Ammar, R., Wilson, G.V., and Provart, N.J. (2007). An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets. PloS one **2** (8): e718.

Witte, C.-P. (2011). Urea metabolism in plants. Plant science an international journal of
experimental plant biology **180** (3): 431–438.

Yan, Y., Wang, P., He, C., and Shi, H. (2017). MeWRKY20 and its interacting and
activating autophagy-related protein 8 (MeATG8) regulate plant disease resistance in
cassava. Biochemical and biophysical research communications **494** (1-2): 20–26.

**Yang, F., Kimberlin, A.N., Elowsky, C.G., Liu, Y., Gonzalez-Solis, A., Cahoon, E.B., and
Alfano, J.R.** (2019). A Plant Immune Receptor Degraded by Selective Autophagy.
Molecular plant **12** (1): 113–123.

Yang, Z., and Klionsky, D.J. (2009). An overview of the molecular mechanism of autophagy.
Current topics in microbiology and immunology **335**: 1–32.

Yang, Z., and Klionsky, D.J. (2010). Eaten alive: A history of macroautophagy. Nature cell
biology **12** (9): 814–822.

Zess, E.K., Jensen, C., Cruz-Mireles, N., La Concepcion, J.C. de, **Sklenar, J., Stephani,
M., Imre, R., Roitinger, E., Hughes, R., Belhaj, K., Mechtler, K., Menke, F.L.H.,
Bozkurt, T., Banfield, M.J., Kamoun, S., Maqbool, A., and Dagdas, Y.F.** (2019). N-
terminal β-strand underpins biochemical specialization of an ATG8 isoform. PLoS biology
17 (7): e3000373.

Zhang, B., Shao, L., Wang, J., Zhang, Y., Guo, X., Peng, Y., Cao, Y., and Lai, Z. (2021a).
Phosphorylation of ATG18a by BAK1 suppresses autophagy and attenuates plant
resistance against necrotrophic pathogens. Autophagy **17** (9): 2093–2110.

Zhang, H., and Sonnewald, U. (2017). Differences and commonalities of plant responses to
single and combined stresses. The Plant journal for cell and molecular biology **90** (5):
839–855.

Zhang, Y., Min, H., Shi, C., Xia, G., and Lai, Z. (2021b). Transcriptome analysis of the role
of autophagy in plant response to heat stress. PloS one **16** (2): e0247783.

Zheng, Z., Qamar, S.A., Chen, Z., and Mengiste, T. (2006). Arabidopsis WRKY33
transcription factor is required for resistance to necrotrophic fungal pathogens. The Plant
journal for cell and molecular biology **48** (4): 592–605.

**Zhou, J., Ma, J., Yang, C., Zhu, X., Li, J., Zheng, X., Li, X., Chen, S., Feng, L., Wang, P.,
Ho, M.I., Ma, W., Liao, J., Li, F., Wang, C., Zhuang, X., Jiang, L., Kang, B.-H., and
Gao, C.** (2023). A non-canonical role of ATG8 in Golgi recovery from heat stress in
plants. Nature plants **9** (5): 749–765.

**Zhou, J., Peng, Z., Long, J., Sosso, D., Liu, B., Eom, J.-S., Huang, S., Liu, S., Vera Cruz,
C., Frommer, W.B., White, F.F., and Yang, B.** (2015). Gene targeting by the TAL

effector PthXo2 reveals cryptic resistance gene for bacterial blight of rice. The Plant

journal for cell and molecular biology **82** (4): 632–643.

Zhou, J., Wang, J., Cheng, Y., Chi, Y.-J., Fan, B., Yu, J.-Q., and Chen, Z. (2013). NBR1-

mediated selective autophagy targets insoluble ubiquitinated protein aggregates in plant

stress responses. PLoS genetics **9** (1): e1003196.

Zhou, J., Wang, J., Yu, J.-Q., and Chen, Z. (2014). Role and regulation of autophagy in

heat stress responses of tomato plants. Frontiers in plant science **5**: 174.

Zhu, J.-K. (2016). Abiotic Stress Signaling and Responses in Plants. Cell **167** (2): 313–324.

Zhu, T., Zou, L., Li, Y., Yao, X., Xu, F., Deng, X., Zhang, D., and Lin, H. (2018).

Mitochondrial alternative oxidase-dependent autophagy involved in ethylene-mediated

drought tolerance in Solanum lycopersicum. Plant biotechnology journal **16** (12): 2063–

2076.

Figure 1: GFP-*St*ATG8 isoforms show a stress-specific degradation pattern. All seven GFP-*St*ATG8 isoforms were transiently expressed in *N. benthamiana*. The plants were subjected to different stresses 1 day after syringe-infiltration of GFP-*St*ATG8 constructs. Western blots show the degradation pattern after 3 days of darkness (**A**), 6/8 h of 40 °C (**B**) and 6 h after *Xcv/hrcV* infection (**C**). Degradation was quantified via measurement of band intensities of GFP-*St*ATG8 (**B**) or GFP-*St*ATG8 and free GFP (**A**) or a specific degradation band (**C**), respectively. Band intensities were normalized to the protein loading control and ratios of GFP-*St*ATG8 and a degradation band (if applicable) were calculated. Original blots can be found in the supplemental data. The experiments were carried out independently three times (B, C)/twice (A) with similar results.

1010

Figure 2: Autophagy is differentially activated under combined stress. (**A**) Schematic representation of the experimental setup to test the effect of combined heat-*Xcv* stress of GFP-*St*ATG8-3.1 degradation and gene expression. (**B**) Degradation pattern of transiently expressed GFP-*St*ATG8-3.1 upon heat, *Xcv,* and a combined heat-*Xcv* challenge. The original blot can be found in the supplemental data. The experiment was carried out independently three times with similar results.

1019

Figure 3: ***Xcv*** **growth is benefited by preceding heat stress in an autophagy-dependent**
manner. (**A**) Schematic representation of the experimental setup to test bacterial growth in
previously heat-stressed plants. (**B-D**) Bacterial growth (*Xcv* (**B**, **C**) or *hrcV* (**D**)) was tested in
tobacco (**B**, **D**) and tomato (**C**) plants that were previously subjected to heat or control
temperature and co-infiltrated with the autophagy inhibitor 3-MA. (**E**) Bacterial growth (*Xcv*)
in heat- and control-treated tobacco plants transiently expressing *AtATG7*. Experiments were
carried out independently at least three times (B, D)/twice (C, E) with similar results. Number
of biologically independent replicates: *Xcv* in tobacco ($n = 9$), *Xcv* in tomato ($n = 12$), *hrcV*
($n = 9$), *At*ATG7 ($n = 6$). Statistical analysis was performed by One-way ANOVA and Turkey's
multiple comparison test.

1030

Figure 4: *At*ATG protein degradation upon combined stress is dependent on autophagy.
(**A**) Schematic representation of the experimental setup to test protein levels and degradation of overexpressed and native *At*ATGs in response to single and combined stress. (**B-C**) Band intensities and degradation pattern of GFP-*At*ATG8a (**B**) and *At*NBR1-TAP (**C**) in stable overexpression lines. (**D**) Comparison of *At*ATG7 band intensities in different Arabidopsis overexpression and knockout lines. All band intensities were normalized to the protein loading controls. Original blots can be found in the supplemental data. Experiments were carried out independently three times with similar results.

Figure 5: A recovery phase does not affect bacterial growth or the plant metabolome. (A) Schematic representation of the experimental setup to test the effect of a recovery phase on bacterial growth and the metabolome. **(B)** Bacterial growth of *Xcv* was tested in tobacco plants previously subjected to heat or control temperature followed by a 24 h recovery period at control temperature and co-infiltrated with the autophagy inhibitor 3-MA (*n* = 9). Statistical analysis was performed by One-way ANOVA and Turkey's multiple comparison test. **(C)** Principal component analysis of the metabolome in response to heat treatment or bacterial infection with and without a recovery phase and 3-MA treatment. **(D)** Heat maps displaying median values in log2 scale of the response of 4 different groups of metabolites to stress treatment as in (A). Red color refers to upregulation compared to the control; blue color refers to downregulation compared to the control.

1054

Figure 6: Detailed overview on effects of *Xcv* infection on the levels of three amino acids and urea. Boxplots showing levels of Glycine (**A**), Aspartic acid (**B**), Asparagine (**C**), and urea (**D**) in response to *Xcv*, *hrcV*, and heat treatment, respectively (*n* = 6). Statistical analysis was performed by One-way ANOVA and Turkey's multiple comparison test.

1059

1060

1061 **Figure 7: Detailed overview on effector-dependent effects of *Xcv* infection on sugar**

1062 **levels.** Boxplots showing levels of (**A-D**) Mono-, (**E-G**) Disaccharides and (**H**) sugar alcohol

1063 in response to *Xcv*, *hrcV*, and heat treatment, respectively (*n* = 6). Statistical analysis was

1064 performed by One-way ANOVA and Turkey's multiple comparison test.

1065

1066

1067

1068

1069 **Supplemental Figure 1:** Transcript levels of native *NbATG7* and *NbJOKA2* upon single and

1070 combined stress. Error bars, SEM (*n* = 4-6); Statistical analysis was performed by One-way

1071 ANOVA, Turkey's multiple comparisons test.

1072

1073

Supplemental Figure 2: Transcriptional response to biotic stress is independent of autophagy but can be induces by increased autophagic activity. Schematic representation of the experimental setup to test the effect of autophagy on stress marker gene transcript levels (**A**). Transcript levels of the biotic stress marker gene *AtFRK1* (**B**, **C**) and the heat stress marker gene *AtHSP70* (**D**, **E**) in wildtype and transgenic overexpression and knockout lines of Arabidopsis. Error bars, SEM (*n* = 3-4); Statistical analysis was performed by One-way ANOVA, Turkey's multiple comparisons test. The experiment was carried out independently twice with similar results.

1082

1083

1084   **Supplemental Figure 3:** Principal component analysis of the metabolome as in Figure 5C

1085   including *hrcV* samples

1086

Supplemental Figure 4: Detailed overview on effects of *Xcv* infection and heat on the levels of organic compounds. Boxplots showing levels of Shikimic acid (**A**), Nicotinic acid (**B**), Phosphoric acid (**C**), Glycerol (**D**), Succinic acid (**E**), Malic acid (**F**) Citric acid (**G**), and Abscisic acid (**H**) in response to *Xcv*, *hrcV*, and heat treatment, respectively (*n* = 6). Statistical analysis was performed by One-way ANOVA and Turkey's multiple comparison test.

1094

1095 **Supplemental Figure 5:** Original blots

1096

1097   **Table S1: Primers used for RT-qPCR analysis**

Primer	Sequence	Organism	Gene ID	Reference
qAtActin2_fwd	TCCCTCAGCACATTCCAGCAGAT	A. thaliana	At3g18780	(Dubiella et al., 2013)
qAtActin2_rev	AACGATTCCTGGACCTGCCTCATC			
qIntron_fwd	TTTTTTGCCCCCTTCGAATC	A. thaliana	At5g65080	(Guerra et al., 2020)
qIntron_rev	ATCTTCCGCCACCACATTGTAC			
qAtHsp70_fwd	TCAAGCGGATAAGAGTCACT	A. thaliana	At3g12580	(Sung et al., 2001; Zhang et al., 2021b)
qAtHsp70_rev	CTCGTCCGGGTTAATGCT			
qAtFRK1_fwd	CGGTCAGATTTCAACAGTTGTC	A. thaliana	At2g19190	(Dubiella et al., 2013)
qAtFRK1_rev	AATAGCAGGTTGGCCTGTAATC			
qNbActin_fwd	AAAGACCAGCTCATCCGTGGAGAA	N. benthamiana		(Leong et al., 2022b)
qNbActin_rev	TGTGGTTTCATGAATGCCAGCAGC			
qNbJoka2_fwd	CGTTGTGATGGTTGTGGTGT	N. benthamiana		(Leong et al., 2022b)
qNbJoka2_rev	AGGGACGCCGGTAAGTTAAA			
qNbATG7_fwd	ATGGCGGATAGTGGAAGAGG	N. benthamiana		(Leong et al., 2022b)
qNbATG7_rev	CCTGACTAGCTAGCCGAGAC			

1098

Parsed Citations

Atkinson, N.J., and Urwin, P.E. (2012). The interaction of plant biotic and abiotic stresses: From genes to the field. Journal of experimental botany 63 (10): 3523–3543.
  Google Scholar: Author Only Title Only Author and Title

Avin-Wittenberg, T. (2019). Autophagy and its role in plant abiotic stress management. Plant, cell & environment 42 (3): 1045–1053.
  Google Scholar: Author Only Title Only Author and Title

Avin-Wittenberg, T., Bajdzienko, K., Wittenberg, G., Alseekh, S., Tohge, T., Bock, R., Giavalisco, P., and Fernie, A.R. (2015). Global analysis of the role of autophagy in cellular metabolism and energy homeostasis in Arabidopsis seedlings under carbon starvation. The Plant cell 27 (2): 306–322.
  Google Scholar: Author Only Title Only Author and Title

Barros, J.A.S., Cavalcanti, J.H.F., Medeiros, D.B., Nunes-Nesi, A., Avin-Wittenberg, T., Fernie, A.R., and Araújo, W.L. (2017). Autophagy Deficiency Compromises Alternative Pathways of Respiration following Energy Deprivation in Arabidopsis thaliana. Plant physiology 175 (1): 62–76.
  Google Scholar: Author Only Title Only Author and Title

Baxter, A., Mittler, R., and Suzuki, N. (2014). ROS as key players in plant stress signalling. Journal of experimental botany 65 (5): 1229–1240.
  Google Scholar: Author Only Title Only Author and Title

Bebber, D.P. (2015). Range-expanding pests and pathogens in a warming world. Annual review of phytopathology 53: 335–356.
  Google Scholar: Author Only Title Only Author and Title

Beck, M., Zhou, J., Faulkner, C., MacLean, D., and Robatzek, S. (2012). Spatio-temporal cellular dynamics of the Arabidopsis flagellin receptor reveal activation status-dependent endosomal sorting. The Plant cell 24 (10): 4205–4219.
  Google Scholar: Author Only Title Only Author and Title

Blanvillain, S., Meyer, D., Boulanger, A., Lautier, M., Guynet, C., Denancé, N., Vasse, J., Lauber, E., and Arlat, M. (2007). Plant carbohydrate scavenging through tonB-dependent receptors: A feature shared by phytopathogenic and aquatic bacteria. PloS one 2 (2): e224.
  Google Scholar: Author Only Title Only Author and Title

Boominathan, P., Shukla, R., Kumar, A., Manna, D., Negi, D., Verma, P.K., and Chattopadhyay, D. (2004). Long term transcript accumulation during the development of dehydration adaptation in Cicer arietinum. Plant physiology 135 (3): 1608–1620.
  Google Scholar: Author Only Title Only Author and Title

Bu, F., Yang, M., Guo, X., Huang, W., and Chen, L. (2020). Multiple Functions of ATG8 Family Proteins in Plant Autophagy. Frontiers in cell and developmental biology 8: 466.
  Google Scholar: Author Only Title Only Author and Title

Cavicchioli, R., Ripple, W.J., Timmis, K.N., Azam, F., and Bakken, L.R., et al. (2019). Scientists' warning to humanity: Microorganisms and climate change. Nature reviews. Microbiology 17 (9): 569–586.
  Google Scholar: Author Only Title Only Author and Title

Chen, L.-Q., Hou, B.-H., Lalonde, S., Takanaga, H., Hartung, M.L., Qu, X.-Q., Guo, W.-J., Kim, J.-G., Underwood, W., Chaudhuri, B., Chermak, D., Antony, G., White, F.F., Somerville, S.C., Mudgett, M.B., and Frommer, W.B. (2010). Sugar transporters for intercellular exchange and nutrition of pathogens. Nature 468 (7323): 527–532.
  Google Scholar: Author Only Title Only Author and Title

Chen, Y., and Klionsky, D.J. (2011). The regulation of autophagy - unanswered questions. Journal of cell science 124 (Pt 2): 161–170.
  Google Scholar: Author Only Title Only Author and Title

Chinchilla, D., Bauer, Z., Regenass, M., Boller, T., and Felix, G. (2006). The Arabidopsis receptor kinase FLS2 binds flg22 and determines the specificity of flagellin perception. The Plant cell 18 (2): 465–476.
  Google Scholar: Author Only Title Only Author and Title

Chiu, C.-Y., Lung, H.-F., Chou, W.-C., Lin, L.-Y., Chow, H.-X., Kuo, Y.-H., Chien, P.-S., Chiou, T.-J., and Liu, T.-Y. (2023). Autophagy-Mediated Phosphate Homeostasis in Arabidopsis Involves Modulation of Phosphate Transporters. Plant & cell physiology 64 (5): 519–535.
  Google Scholar: Author Only Title Only Author and Title

Chung, T. (2019). How phosphoinositides shape autophagy in plant cells. Plant science an international journal of experimental plant biology 281: 146–158.
  Google Scholar: Author Only Title Only Author and Title

Dagdas, Y.F., Belhaj, K., Maqbool, A., Chaparro-Garcia, A., Pandey, P., Petre, B., Tabassum, N., Cruz-Mireles, N., Hughes, R.K.,

Sklenar, J., Win, J., Menke, F., Findlay, K., Banfield, M.J., Kamoun, S., and Bozkurt, T.O. (2016). An effector of the Irish potato famine pathogen antagonizes a host autophagy cargo receptor. eLife 5.
Google Scholar: Author Only Title Only Author and Title

Dagdas, Y.F., Pandey, P., Tumtas, Y., Sanguankiattichai, N., Belhaj, K., Duggan, C., Leary, A.Y., Segretin, M.E., Contreras, M.P., Savage, Z, Khandare, V.S., Kamoun, S., and Bozkurt, T.O. (2018). Host autophagy machinery is diverted to the pathogen interface to mediate focal defense responses against the Irish potato famine pathogen. eLife 7.
Google Scholar: Author Only Title Only Author and Title

Dhatt, B.K., Abshire, N., Paul, P., Hasanthika, K., Sandhu, J., Zhang, Q., Obata, T., and Walia, H. (2019). Metabolic Dynamics of Developing Rice Seeds Under High Night-Time Temperature Stress. Frontiers in plant science 10: 1443.
Google Scholar: Author Only Title Only Author and Title

Dodds, P.N., and Rathjen, J.P. (2010). Plant immunity: Towards an integrated view of plant-pathogen interactions. Nature reviews. Genetics 11 (8): 539–548.
Google Scholar: Author Only Title Only Author and Title

Dubiella, U., Seybold, H., Durian, G., Komander, E., Lassig, R., Witte, C.-P., Schulze, W.X., and Romeis, T. (2013). Calcium-dependent protein kinase/NADPH oxidase activation circuit is required for rapid defense signal propagation. Proceedings of the National Academy of Sciences of the United States of America 110 (21): 8744–8749.
Google Scholar: Author Only Title Only Author and Title

Dugé de Bernonville, T., Noël, L.D., SanCristobal, M., Danoun, S., Becker, A, Soreau, P., Arlat, M., and Lauber, E. (2014). Transcriptional reprogramming and phenotypical changes associated with growth of Xanthomonas campestris pv. campestris in cabbage xylem sap. FEMS microbiology ecology 89 (3): 527–541.
Google Scholar: Author Only Title Only Author and Title

Eom, J.-S., Chen, L.-Q., Sosso, D., Julius, B.T., Lin, I.W., Qu, X.-Q., Braun, D.M., and Frommer, W.B. (2015). SWEETs, transporters for intracellular and intercellular sugar translocation. Current opinion in plant biology 25: 53–62.
Google Scholar: Author Only Title Only Author and Title

Fatima, U., and Senthil-Kumar, M. (2015). Plant and pathogen nutrient acquisition strategies. Frontiers in plant science 6: 750.
Google Scholar: Author Only Title Only Author and Title

Feng, Y., He, D., Yao, Z, and Klionsky, D.J. (2014). The machinery of macroautophagy. Cell research 24 (1): 24–41.
Google Scholar: Author Only Title Only Author and Title

Fraire-Velazquez, S., Rodriguez-Guerra, R., and Sanchez-Caldero, L. (2011). Abiotic and Biotic Stress Response Crosstalk in Plants. In Abiotic Stress Response in Plants - Physiological, Biochemical and Genetic Perspectives, A. Shanker, ed (InTech).
Google Scholar: Author Only Title Only Author and Title

Glazebrook, J. (2005). Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. Annual review of phytopathology 43: 205–227.
Google Scholar: Author Only Title Only Author and Title

Grützner, R., Martin, P., Horn, C., Mortensen, S., Cram, E.J., Lee-Parsons, C.W.T., Stuttmann, J., and Marillonnet, S. (2021). High-efficiency genome editing in plants mediated by a Cas9 gene containing multiple introns. Plant communications 2 (2): 100135.
Google Scholar: Author Only Title Only Author and Title

Guerra, T., Schilling, S., Hake, K., Gorzolka, K., Sylvester, F.-P., Conrads, B., Westermann, B., and Romeis, T. (2020). Calcium-dependent protein kinase 5 links calcium signaling with N-hydroxy-l-pipecolic acid- and SARD1-dependent immune memory in systemic acquired resistance. The New phytologist 225 (1): 310–325.
Google Scholar: Author Only Title Only Author and Title

Hachez, C., Veljanovski, V., Reinhardt, H., Guillaumot, D., Vanhee, C., Chaumont, F., and Batoko, H. (2014). The Arabidopsis abiotic stress-induced TSPO-related protein reduces cell-surface expression of the aquaporin PIP2;7 through protein-protein interactions and autophagic degradation. The Plant cell 26 (12): 4974–4990.
Google Scholar: Author Only Title Only Author and Title

Hafrén, A, Macia, J.-L., Love, A.J., Milner, J.J., Drucker, M., and Hofius, D. (2017). Selective autophagy limits cauliflower mosaic virus infection by NBR1-mediated targeting of viral capsid protein and particles. Proceedings of the National Academy of Sciences of the United States of America 114 (10): E2026-E2035.
Google Scholar: Author Only Title Only Author and Title

Hafrén, A, Üstün, S., Hochmuth, A, Svenning, S., Johansen, T., and Hofius, D. (2018). Turnip Mosaic Virus Counteracts Selective Autophagy of the Viral Silencing Suppressor HCpro. Plant physiology 176 (1): 649–662.
Google Scholar: Author Only Title Only Author and Title

Hey, S.J., Byrne, E., and Halford, N.G. (2010). The interface between metabolic and stress signalling. Annals of botany 105 (2): 197–203.

Google Scholar: Author Only Title Only Author and Title

Huot, B., Castroverde, C.D.M., Velásquez, A.C., Hubbard, E., Pulman, J.A., Yao, J., Childs, K.L., Tsuda, K., Montgomery, B.L., and He, S.Y. (2017). Dual impact of elevated temperature on plant defence and bacterial virulence in Arabidopsis. Nature communications 8 (1): 1808.
Google Scholar: Author Only Title Only Author and Title

Islam, M.R., Feng, B., Chen, T., Tao, L., and Fu, G. (2018). Role of Abscisic Acid in Thermal Acclimation of Plants. J. Plant Biol. 61 (5): 255–264.
Google Scholar: Author Only Title Only Author and Title

Jones, J.D.G., and Dangl, J.L. (2006). The plant immune system. Nature 444 (7117): 323–329.
Google Scholar: Author Only Title Only Author and Title

Kellner, R., La Concepcion, J.C. de, Maqbool, A., Kamoun, S., and Dagdas, Y.F. (2017). ATG8 Expansion: A Driver of Selective Autophagy Diversification? Trends in plant science 22 (3): 204–214.
Google Scholar: Author Only Title Only Author and Title

Khurana, N., Sharma, N., and Khurana, P. (2017). Overexpression of a heat stress inducible, wheat myo-inositol-1-phosphate synthase 2 (TaMIPS2) confers tolerance to various abiotic stresses in Arabidopsis thaliana. Agri Gene 6: 24–30.
Google Scholar: Author Only Title Only Author and Title

Kissoudis, C., van de Wiel, C., Visser, R.G.F., and van der Linden, G. (2014). Enhancing crop resilience to combined abiotic and biotic stress through the dissection of physiological and molecular crosstalk. Frontiers in plant science 5: 207.
Google Scholar: Author Only Title Only Author and Title

Klionsky, D.J., Abdel-Aziz, A.K., Abdelfatah, S., Abdellatif, M., and Abdoli, A, et al. (2021). Guidelines for the use and interpretation of assays for monitoring autophagy (4th edition)1. Autophagy 17 (1): 1–382.
Google Scholar: Author Only Title Only Author and Title

Lai, Z, Wang, F., Zheng, Z, Fan, B., and Chen, Z (2011). A critical role of autophagy in plant resistance to necrotrophic fungal pathogens. The Plant journal for cell and molecular biology 66 (6): 953–968.
Google Scholar: Author Only Title Only Author and Title

Lal, N.K., Thanasuwat, B., Huang, P.-J., Cavanaugh, K.A, Carter, A, Michelmore, R.W., and Dinesh-Kumar, S.P. (2020). Phytopathogen Effectors Use Multiple Mechanisms to Manipulate Plant Autophagy. Cell host & microbe 28 (4): 558-571.e6.
Google Scholar: Author Only Title Only Author and Title

Larkindale, J., and Knight, M.R. (2002). Protection against heat stress-induced oxidative damage in Arabidopsis involves calcium, abscisic acid, ethylene, and salicylic acid. Plant physiology 128 (2): 682–695.
Google Scholar: Author Only Title Only Author and Title

Lenz, H.D., Haller, E., Melzer, E., Kober, K., Wurster, K., Stahl, M., Bassham, D.C., Vierstra, R.D., Parker, J.E., Bautor, J., Molina, A, Escudero, V., Shindo, T., van der Hoorn, R.A.L., Gust, A.A., and Nürnberger, T. (2011). Autophagy differentially controls plant basal immunity to biotrophic and necrotrophic pathogens. The Plant journal for cell and molecular biology 66 (5): 818–830.
Google Scholar: Author Only Title Only Author and Title

Leong, J.X., Langin, G., and Üstün, S. (2022a). Selective autophagy: Adding precision in plant immunity. Essays in biochemistry 66 (2): 189–206.
Google Scholar: Author Only Title Only Author and Title

Leong, J.X., Raffeiner, M., Spinti, D., Langin, G., Franz-Wachtel, M., Guzman, A.R., Kim, J.-G., Pandey, P., Minina, A.E., Macek, B., Hafrén, A, Bozkurt, T.O., Mudgett, M.B., Börnke, F., Hofius, D., and Üstün, S. (2022b). A bacterial effector counteracts host autophagy by promoting degradation of an autophagy component. The EMBO journal 41 (13): e110352.
Google Scholar: Author Only Title Only Author and Title

Lin, L.-Y., Chow, H.-X., Chen, C.-H., Mitsuda, N., Chou, W.-C., and Liu, T.-Y. (2023). Role of autophagy-related proteins ATG8f and ATG8h in the maintenance of autophagic activity in Arabidopsis roots under phosphate starvation. Frontiers in plant science 14: 1018984.
Google Scholar: Author Only Title Only Author and Title

Lippok, B., Birkenbihl, R.P., Rivory, G., Brümmer, J., Schmelzer, E., Logemann, E., and Somssich, I.E. (2007). Expression of AtWRKY33 encoding a pathogen- or PAMP-responsive WRKY transcription factor is regulated by a composite DNA motif containing W box elements. Molecular plant-microbe interactions MPMI 20 (4): 420–429.
Google Scholar: Author Only Title Only Author and Title

Lisec, J., Schauer, N., Kopka, J., Willmitzer, L., and Fernie, A.R. (2006). Gas chromatography mass spectrometry-based metabolite profiling in plants. Nature protocols 1 (1): 387–396.
Google Scholar: Author Only Title Only Author and Title

Liu, Y., and Bassham, D.C. (2012). Autophagy: Pathways for self-eating in plant cells. Annual review of plant biology 63: 215–237.

Google Scholar: Author Only Title Only Author and Title

Luck, J., Spackman, M., Freeman, A, Tre̦bicki, P., Griffiths, W., Finlay, K., and Chakraborty, S. (2011). Climate change and diseases of food crops. Plant Pathology 60 (1): 113–121.
 Google Scholar: Author Only Title Only Author and Title

Magen, S., Seybold, H., Laloum, D., and Avin-Wittenberg, T. (2022). Metabolism and autophagy in plants-a perfect match. FEBS letters 596 (17): 2133–2151.
 Google Scholar: Author Only Title Only Author and Title

Mao, G., Meng, X., Liu, Y., Zheng, Z, Chen, Z, and Zhang, S. (2011). Phosphorylation of a WRKY transcription factor by two pathogen-responsive MAPKs drives phytoalexin biosynthesis in Arabidopsis. The Plant cell 23 (4): 1639–1653.
 Google Scholar: Author Only Title Only Author and Title

Marshall, R.S., and Vierstra, R.D. (2018). Autophagy: The Master of Bulk and Selective Recycling. Annual review of plant biology 69: 173–208.
 Google Scholar: Author Only Title Only Author and Title

Masclaux-Daubresse, C., d'Andrea, S., Bouchez, I., and Cacas, J.-L. (2020). Reserve lipids and plant autophagy. Journal of experimental botany 71 (10): 2854–2861.
 Google Scholar: Author Only Title Only Author and Title

Minina, E.A, Moschou, P.N., Vetukuri, R.R., Sanchez-Vera, V., Cardoso, C., Liu, Q., Elander, P.H., Dalman, K., Beganovic, M., Lindberg Yilmaz, J., Marmon, S., Shabala, L., Suarez, M.F., Ljung, K., Novák, O., Shabala, S., Stymne, S., Hofius, D., and Bozhkov, P.V. (2018). Transcriptional stimulation of rate-limiting components of the autophagic pathway improves plant fitness. Journal of experimental botany 69 (6): 1415–1432.
 Google Scholar: Author Only Title Only Author and Title

Ohama, N., Sato, H., Shinozaki, K., and Yamaguchi-Shinozaki, K. (2017). Transcriptional Regulatory Network of Plant Heat Stress Response. Trends in plant science 22 (1): 53–65.
 Google Scholar: Author Only Title Only Author and Title

Pandey, P., Leary, A.Y., Tumtas, Y., Savage, Z, Dagvadorj, B., Duggan, C., Yuen, E.L., Sanguankiattichai, N., Tan, E., Khandare, V., Connerton, A.J., Yunusov, T., Madalinski, M., Mirkin, F.G., Schornack, S., Dagdas, Y., Kamoun, S., and Bozkurt, T.O. (2021). An oomycete effector subverts host vesicle trafficking to channel starvation-induced autophagy to the pathogen interface. eLife 10.
 Google Scholar: Author Only Title Only Author and Title

Pandey, P., Ramegowda, V., and Senthil-Kumar, M. (2015). Shared and unique responses of plants to multiple individual stresses and stress combinations: Physiological and molecular mechanisms. Frontiers in plant science 6: 723.
 Google Scholar: Author Only Title Only Author and Title

Patel, S., and Dinesh-Kumar, S.P. (2008). Arabidopsis ATG6 is required to limit the pathogen-associated cell death response. Autophagy 4 (1): 20–27.
 Google Scholar: Author Only Title Only Author and Title

Pieretti, I., Royer, M., Barbe, V., Carrere, S., Koebnik, R., Couloux, A, Darrasse, A, Gouzy, J., Jacques, M.-A, Lauber, E., Manceau, C., Mangenot, S., Poussier, S., Segurens, B., Szurek, B., Verdier, V., Arlat, M., Gabriel, D.W., Rott, P., and Cociancich, S. (2012). Genomic insights into strategies used by Xanthomonas albilineans with its reduced artillery to spread within sugarcane xylem vessels. BMC genomics 13: 658.
 Google Scholar: Author Only Title Only Author and Title

Ramegowda, V., and Senthil-Kumar, M. (2015). The interactive effects of simultaneous biotic and abiotic stresses on plants: Mechanistic understanding from drought and pathogen combination. Journal of plant physiology 176: 47–54.
 Google Scholar: Author Only Title Only Author and Title

Rasmussen, S., Barah, P., Suarez-Rodriguez, M.C., Bressendorff, S., Friis, P., Costantino, P., Bones, A.M., Nielsen, H.B., and Mundy, J. (2013). Transcriptome responses to combinations of stresses in Arabidopsis. Plant physiology 161 (4): 1783–1794.
 Google Scholar: Author Only Title Only Author and Title

Rico, A, and Preston, G.M. (2008). Pseudomonas syringae pv. tomato DC3000 uses constitutive and apoplast-induced nutrient assimilation pathways to catabolize nutrients that are abundant in the tomato apoplast. Molecular plant-microbe interactions MPMI 21 (2): 269–282.
 Google Scholar: Author Only Title Only Author and Title

Rodriguez, E., Chevalier, J., Olsen, J., Ansbøl, J., Kapousidou, V., Zuo, Z, Svenning, S., Loefke, C., Koemeda, S., Drozdowskyj, P.S., Jez, J., Durnberger, G., Kuenzl, F., Schutzbier, M., Mechtler, K., Ebstrup, E.N., Lolle, S., Dagdas, Y., and Petersen, M. (2020). Autophagy mediates temporary reprogramming and dedifferentiation in plant somatic cells. The EMBO journal 39 (4): e103315.
 Google Scholar: Author Only Title Only Author and Title

Schmidt, A, Mächtel, R., Ammon, A, Engelsdorf, T., Schmitz, J., Maurino, V.G., and Voll, L.M. (2020). Reactive oxygen species

dosage in Arabidopsis chloroplasts can improve resistance towards Colletotrichum higginsianum by the induction of WRKY33. The New phytologist 226 (1): 189–204.
Google Scholar: Author Only Title Only Author and Title

Sertsuvalkul, N., DeMell, A., and Dinesh-Kumar, S.P. (2022). The complex roles of autophagy in plant immunity. FEBS letters 596 (17): 2163–2171.
Google Scholar: Author Only Title Only Author and Title

Sharma, N., Chaudhary, C., and Khurana, P. (2020). Wheat Myo-inositol phosphate synthase influences plant growth and stress responses via ethylene mediated signaling. Scientific reports 10 (1): 10766.
Google Scholar: Author Only Title Only Author and Title

Signorelli, S., Tarkowski, Ł.P., van den Ende, W., and Bassham, D.C. (2019). Linking Autophagy to Abiotic and Biotic Stress Responses. Trends in plant science 24 (5): 413–430.
Google Scholar: Author Only Title Only Author and Title

Slavikova, S., Ufaz, S., Avin-Wittenberg, T., Levanony, H., and Galili, G. (2008). An autophagy-associated Atg8 protein is involved in the responses of Arabidopsis seedlings to hormonal controls and abiotic stresses. Journal of experimental botany 59 (14): 4029–4043.
Google Scholar: Author Only Title Only Author and Title

Stephani, M., and Dagdas, Y. (2020). Plant Selective Autophagy-Still an Uncharted Territory With a Lot of Hidden Gems. Journal of molecular biology 432 (1): 63–79.
Google Scholar: Author Only Title Only Author and Title

Sung, D.Y., Vierling, E., and Guy, C.L. (2001). Comprehensive expression profile analysis of the Arabidopsis Hsp70 gene family. Plant physiology 126 (2): 789–800.
Google Scholar: Author Only Title Only Author and Title

Svenning, S., Lamark, T., Krause, K., and Johansen, T. (2011). Plant NBR1 is a selective autophagy substrate and a functional hybrid of the mammalian autophagic adapters NBR1 and p62/SQSTM1. Autophagy 7 (9): 993–1010.
Google Scholar: Author Only Title Only Author and Title

Tarnowski, L., Rodriguez, M.C., Brzywczy, J., Piecho-Kabacik, M., Krčkova, Z, Martinec, J., Wawrzynska, A., and Sirko, A. (2020). A selective autophagy cargo receptor NBR1 modulates abscisic acid signalling in Arabidopsis thaliana. Scientific reports 10 (1): 7778.
Google Scholar: Author Only Title Only Author and Title

Thirumalaikumar, V.P., Gorka, M., Schulz, K., Masclaux-Daubresse, C., Sampathkumar, A, Skirycz, A, Vierstra, R.D., and Balazadeh, S. (2021). Selective autophagy regulates heat stress memory in Arabidopsis by NBR1-mediated targeting of HSP90.1 and ROF1. Autophagy 17 (9): 2184–2199.
Google Scholar: Author Only Title Only Author and Title

Tsuda, K., Mine, A, Bethke, G., Igarashi, D., Botanga, C.J., Tsuda, Y., Glazebrook, J., Sato, M., and Katagiri, F. (2013). Dual regulation of gene expression mediated by extended MAPK activation and salicylic acid contributes to robust innate immunity in Arabidopsis thaliana. PLoS genetics 9 (12): e1004015.
Google Scholar: Author Only Title Only Author and Title

Üstün, S., Hafrén, A, Liu, Q., Marshall, R.S., Minina, E.A, Bozhkov, P.V., Vierstra, R.D., and Hofius, D. (2018). Bacteria Exploit Autophagy for Proteasome Degradation and Enhanced Virulence in Plants. The Plant cell 30 (3): 668–685.
Google Scholar: Author Only Title Only Author and Title

Üstün, S., Sheikh, A, Gimenez-Ibanez, S., Jones, A, Ntoukakis, V., and Börnke, F. (2016). The Proteasome Acts as a Hub for Plant Immunity and Is Targeted by Pseudomonas Type III Effectors. Plant physiology 172 (3): 1941–1958.
Google Scholar: Author Only Title Only Author and Title

van Doorn, W.G., and Papini, A (2013). Ultrastructure of autophagy in plant cells: A review. Autophagy 9 (12): 1922–1936.
Google Scholar: Author Only Title Only Author and Title

Wang, P., Nolan, T.M., Yin, Y., and Bassham, D.C. (2020). Identification of transcription factors that regulate ATG8 expression and autophagy in Arabidopsis. Autophagy 16 (1): 123–139.
Google Scholar: Author Only Title Only Author and Title

Wang, P., Wang, T., Han, J., Li, M., Zhao, Y., Su, T., and Ma, C. (2021). Plant Autophagy: An Intricate Process Controlled by Various Signaling Pathways. Frontiers in plant science 12: 754982.
Google Scholar: Author Only Title Only Author and Title

Wang, Y., Bao, Z, Zhu, Y., and Hua, J. (2009). Analysis of temperature modulation of plant defense against biotrophic microbes. Molecular plant-microbe interactions MPMI 22 (5): 498–506.
Google Scholar: Author Only Title Only Author and Title

Wang, Y., Cai, S., Yin, L., Shi, K., Xia, X., Zhou, Y., Yu, J., and Zhou, J. (2015). Tomato HsfA1a plays a critical role in plant drought tolerance by activating ATG genes and inducing autophagy. Autophagy 11 (11): 2033–2047.
Google Scholar: Author Only Title Only Author and Title

Wang, Y., Nishimura, M.T., Zhao, T., and Tang, D. (2011). ATG2, an autophagy-related protein, negatively affects powdery mildew resistance and mildew-induced cell death in Arabidopsis. The Plant journal for cell and molecular biology 68 (1): 74–87.
Google Scholar: Author Only Title Only Author and Title

Winter, D., Vinegar, B., Nahal, H., Ammar, R., Wilson, G.V., and Provart, N.J. (2007). An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets. PloS one 2 (8): e718.
Google Scholar: Author Only Title Only Author and Title

Witte, C.-P. (2011). Urea metabolism in plants. Plant science an international journal of experimental plant biology 180 (3): 431–438.
Google Scholar: Author Only Title Only Author and Title

Yan, Y., Wang, P., He, C., and Shi, H. (2017). MeWRKY20 and its interacting and activating autophagy-related protein 8 (MeATG8) regulate plant disease resistance in cassava. Biochemical and biophysical research communications 494 (1-2): 20–26.
Google Scholar: Author Only Title Only Author and Title

Yang, F., Kimberlin, A.N., Elowsky, C.G., Liu, Y., Gonzalez-Solis, A, Cahoon, E.B., and Alfano, J.R. (2019). A Plant Immune Receptor Degraded by Selective Autophagy. Molecular plant 12 (1): 113–123.
Google Scholar: Author Only Title Only Author and Title

Yang, Z., and Klionsky, D.J. (2009). An overview of the molecular mechanism of autophagy. Current topics in microbiology and immunology 335: 1–32.
Google Scholar: Author Only Title Only Author and Title

Yang, Z., and Klionsky, D.J. (2010). Eaten alive: A history of macroautophagy. Nature cell biology 12 (9): 814–822.
Google Scholar: Author Only Title Only Author and Title

Zess, E.K., Jensen, C., Cruz-Mireles, N., La Concepcion, J.C. de, Sklenar, J., Stephani, M., Imre, R., Roitinger, E., Hughes, R., Belhaj, K., Mechtler, K., Menke, F.L.H., Bozkurt, T., Banfield, M.J., Kamoun, S., Maqbool, A, and Dagdas, Y.F. (2019). N-terminal β-strand underpins biochemical specialization of an ATG8 isoform. PLoS biology 17 (7): e3000373.
Google Scholar: Author Only Title Only Author and Title

Zhang, B., Shao, L., Wang, J., Zhang, Y., Guo, X., Peng, Y., Cao, Y., and Lai, Z. (2021a). Phosphorylation of ATG18a by BAK1 suppresses autophagy and attenuates plant resistance against necrotrophic pathogens. Autophagy 17 (9): 2093–2110.
Google Scholar: Author Only Title Only Author and Title

Zhang, H., and Sonnewald, U. (2017). Differences and commonalities of plant responses to single and combined stresses. The Plant journal for cell and molecular biology 90 (5): 839–855.
Google Scholar: Author Only Title Only Author and Title

Zhang, Y., Min, H., Shi, C., Xia, G., and Lai, Z. (2021b). Transcriptome analysis of the role of autophagy in plant response to heat stress. PloS one 16 (2): e0247783.
Google Scholar: Author Only Title Only Author and Title

Zheng, Z, Qamar, S.A, Chen, Z., and Mengiste, T. (2006). Arabidopsis WRKY33 transcription factor is required for resistance to necrotrophic fungal pathogens. The Plant journal for cell and molecular biology 48 (4): 592–605.
Google Scholar: Author Only Title Only Author and Title

Zhou, J., Ma, J., Yang, C., Zhu, X., Li, J., Zheng, X., Li, X., Chen, S., Feng, L., Wang, P., Ho, M.I., Ma, W., Liao, J., Li, F., Wang, C., Zhuang, X., Jiang, L., Kang, B.-H., and Gao, C. (2023). A non-canonical role of ATG8 in Golgi recovery from heat stress in plants. Nature plants 9 (5): 749–765.
Google Scholar: Author Only Title Only Author and Title

Zhou, J., Peng, Z, Long, J., Sosso, D., Liu, B., Eom, J.-S., Huang, S., Liu, S., Vera Cruz, C., Frommer, W.B., White, F.F., and Yang, B. (2015). Gene targeting by the TAL effector PthXo2 reveals cryptic resistance gene for bacterial blight of rice. The Plant journal for cell and molecular biology 82 (4): 632–643.
Google Scholar: Author Only Title Only Author and Title

Zhou, J., Wang, J., Cheng, Y., Chi, Y.-J., Fan, B., Yu, J.-Q., and Chen, Z. (2013). NBR1-mediated selective autophagy targets insoluble ubiquitinated protein aggregates in plant stress responses. PLoS genetics 9 (1): e1003196.
Google Scholar: Author Only Title Only Author and Title

Zhou, J., Wang, J., Yu, J.-Q., and Chen, Z. (2014). Role and regulation of autophagy in heat stress responses of tomato plants. Frontiers in plant science 5: 174.
Google Scholar: Author Only Title Only Author and Title

Zhu, J.-K. (2016). Abiotic Stress Signaling and Responses in Plants. Cell 167 (2): 313–324.
Google Scholar: Author Only Title Only Author and Title

Zhu, T., Zou, L., Li, Y., Yao, X., Xu, F., Deng, X., Zhang, D., and Lin, H. (2018). Mitochondrial alternative oxidase-dependent autophagy involved in ethylene-mediated drought tolerance in Solanum lycopersicum. Plant biotechnology journal 16 (12): 2063–2076.
Google Scholar: Author Only Title Only Author and Title