Surface Hydrophilicity Promotes Bacterial Twitching Motility

Megan T. O'Hara[a], Tori M. Shimozono[a], Keane J. Dye[ab], David Harris[ac], and Zhaomin Yang[a]

[a]Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia, USA

[b]Current address: Uniformed Services University of the Health Sciences, Bethesda, Maryland,

USA

[c]Current address: Department of Biochemistry, Virginia Tech, Blacksburg, Virginia, USA

#Address correspondence to Zhaomin Yang at zmyang@vt.edu

Keywords: Twitching motility, *Pseudomonas aeruginosa*, *Acinetobacter*, bile salts, detergents,

surface property, hydrophilicity

Abstract word count: 241

Importance word count: 147

Main text word count: 4,570

Abstract

Twitching motility is a form of bacterial surface translocation powered by the type IV pilus (T4P). It is frequently analyzed by interstitial colony expansion between agar and the polystyrene surfaces of Petri dishes. In such assays, the twitching motility of *Acinetobacter nosocomialis* was observed with MacConkey but not Luria-Bertani (LB) agar media. One difference between these two media is the presence of bile salts as a selective agent in MacConkey but not in LB. Here, we demonstrate that the addition of bile salts to LB allowed *A. nosocomialis* to display twitching. Similarly, bile salts enhanced the twitching of *Acinetobacter baumannii* and *Pseudomonas aeruginosa* in LB. These observations suggest that there is a common mechanism whereby bile salts enhance bacterial twitching and promote interstitial colony expansion. Bile salts disrupt lipid membranes and apply envelope stress as detergents. Surprisingly, their stimulatory effect on twitching appears not to be related to a bacterial physiological response to stressors. Rather it is due to their ability to alter the physicochemical properties of a twitching surface. We observed that while other detergents promoted twitching like bile salts, stresses applied by antibiotics, including the outer membrane-targeting polymyxin B, did not enhanced twitching motility. More importantly, bacteria displayed increased twitching on hydrophilic surfaces such as those of glass and tissue culture-treated polystyrene plastics, and bile salts no longer stimulated twitching on these surfaces. Together, our results show that altering the hydrophilicity of a twitching surface significantly impacts T4P functionality.

Importance

The bacterial type IV pilus (T4P) is a critical virulence factor for many medically important

pathogens, some of which are prioritized by the World Health Organization for their high levels

of antibiotic resistance. The T4P is known to propel bacterial twitching motility, providing a

convenient assay for T4P functionality. Here, we show that bile salts and other detergents

augment the twitching of multiple bacterial pathogens. We identified the underlying mechanism

as the alteration of surface hydrophilicity by detergents. Consequently, hydrophilic surfaces such

as those of glass or plasma-treated polystyrene promote bacterial twitching, bypassing the

requirement for detergents. The implication is that surface properties, such as those of tissues and

medical implants, significantly impact the functionality of bacterial T4P as a virulence

determinant. This offers valuable insights for developing countermeasures against the

colonization and infection by bacterial pathogens of critical importance to human health on a

global scale.

Introduction

Twitching motility is a form of non-flagellated bacterial locomotion that allows bacteria

to move on or between solid surfaces (1-4). It is powered by the bacterial type IV pilus (T4P)

which can be assembled and disassembled by the supramolecular T4P machinery (T4PM) (1, 5-

7). The current model proposes that it is the recurrent cycles of T4P assembly and disassembly,

or extension and retraction, that powers this form of bacterial surface motility (8, 9). The T4PM

assembles the long T4P filament that protrudes from a cell into its surroundings. When the tip of

an extended T4P attaches to a solid substratum, the retraction of the T4P by the T4PM moves a

bacterium toward the point of attachment. This translocation of bacterial cells on or between

solid surfaces, results in bacterial twitching motility.

Of relevance to human health, the T4P plays a crucial role in the pathogenesis of many

important bacterial pathogens (10-15). These include *Pseudomonas aeruginosa* and

Acinetobacter baumannii, both on the list of priority pathogens per the World Health

Organization (WHO) (16). One of the primary functions of the T4P as a virulence factor is for

adherence to human cells or tissues to initiate colonization and invasion (4, 17, 18).

Acinetobacter nosocomialis, a close relative of *A. baumannii,* is an opportunistic pathogen

primarily causing nosocomial or hospital-acquired infections (19). The *A. nosocomialis* M2

strain has been used as a model for studies of *Acinetobacter* pathogenesis and T4P functionality

(18, 20-22). Despite the lack of flagella and the acineto- or non-motile designation for this genus,

many *Acinetobacter* species are, in fact, motile by T4P-dependent twitching motility (23-26). As

such, twitching motility provides a convenient assay for investigating the functionality of the

bacterial T4P in these medically important pathogens.

Twitching motility is routinely analyzed by observing interstitial colony expansion

between the lower surface of solidified nutrient agar and that of plastic Petri dishes made of

polystyrene (25, 27). Such stab assays involve the inoculation of the interstitial space by stabbing

through the agar, and this method has been used for the identification of T4P or *pil* genes by the

isolation of *P. aeruginosa* mutants that were defective in twitching motility (3). T4P genes

encode the core components of the T4PM and their functions in twitching motility are conserved

among *P. aeruginosa* and many Gram-negative and Gram-positive bacteria (2, 10, 28-31). These

include PilA, the major pilin, as well as PilB, the T4P extension ATPase, and PilT, the T4P

retraction ATPase. Along with other T4P proteins, the PilB and the PilT ATPases polymerize

and depolymerize pilins into or from the T4P filament, respectively. Bacterial translocation by

twitching motility over distances longer than the length of an extended pilus depends on the

dynamic nature of T4P assembly and disassembly coordinated by the T4PM (8, 9).

The regulation of bacterial motility by environmental cues has been studied most

extensively in flagellated bacteria (32-35). Besides chemotactic responses (36), the biogenesis of

bacterial flagella is modulated through gene expression and flagellar assembly by signals such as

nutrient and surface availability (32-35). In addition, an alternative sigma factor, which is

responsive to envelope and other environmental stressors, transcriptionally regulates the

expression of flagellar genes in many bacteria (37-39). Although the T4P has been investigated

to a lesser extent, there is clear evidence that its biogenesis and function are influenced by

regulatory mechanisms and environmental factors. In many T4P or *pil* gene clusters, there are

conserved two-component systems including PilS and PilR (2, 31). In selected organisms, these

regulators have been demonstrated to affect the expression of T4P genes (40-43). Signals of both

chemical and physical nature are known to influence T4P-mediated motility (44-46). For

example, lactate can induce PilT-dependent T4P retraction in *Neisseria meningitidis* whereas

both temperature and blue light were shown to influence bacterial twitching motility (47, 48).

It was observed previously that *A. nosocomialis* exhibited twitching motility only with

MacConkey but not with Luria-Bertani (LB) agar media on polystyrene Petri dishes (22, 49). In

our current study, we investigated the underlying reasons for the observed differences in

bacterial twitching between these two media. We determined that bile salts are the key

component that allows *A. nosocomialis* to twitch in MacConkey media. This is because the

addition of bile salts to LB allowed *A. nosocomialis* to twitch to a similar extent as with

MacConkey. We also observed similar stimulatory effects of bile salts on the twitching of *P.*

aeruginosa and *A. baumannii*. Bile salts are anionic detergents that can apply membrane stress to

bacteria (50-53). Our results further demonstrate that other detergents likewise can enhance

bacterial twitching on polystyrene surfaces. Antibiotics, including the outer membrane-targeting

polymyxin B, do not increase *P. aeruginosa* twitching motility. This suggests that the

mechanism for the stimulatory effect of bile salts is unlikely related to a bacterial response to the

presence of a general or envelope stressor. Instead, we suspected that bile salts and other

detergents increased the hydrophilicity of polystyrene surfaces, and it was this increase in

hydrophilicity that promoted bacterial twitching. Indeed, we observed that glass surfaces, which

are more hydrophilic than polystyrene, significantly promoted twitching motility. In contrast,

increasing the hydrophobicity of glass surfaces by coating attenuated twitching. Like glass,

plasma-treatment of polystyrene surfaces is known to increase their hydrophilicity for culturing

tissues or cells. We observed that tissue culture (TC)-treated polystyrene surfaces significantly

increased bacterial twitching. Moreover, the addition of bile salts no longer stimulated twitching

on glass or TC-treated polystyrene surfaces. Our results here suggest that bacterial pathogens

may have evolved mechanisms to differentially interact with surfaces that have varying

physicochemical properties to optimize host recognition, colonization, and infections.

Results

Bile salts enable *Acinetobacter* twitching motility in stab assays. It has been reported in the

literature that *A. nosocomialis* displays twitching motility with MacConkey, but not LB agar, in stab

assays (22, 49). In these assays, bacterial cells are stab-inoculated through the agar to form an interstitial

colony between the Petri dish and the agar media (25, 27). After a period of incubation, the size of an

interstitial colony can be measured to quantify twitching motility. As shown in Figure 1A, the *A.*

nosocomialis M2 strain shows clear twitching with MacConkey, but not with LB agar. These results

confirmed that MacConkey but not LB media allows *A. nosocomialis* to display twitching motility as

observed previously (22, 49).

We compared the composition of these two commonly used bacterial growth media (Table

S1). Notwithstanding their commonalities, LB lacks peptone, lactose, and bile salts that are

present in MacConkey. Peptone is a proteinous nutrient source and lactose is a carbon and

energy source. Bile salts are cholesterol derivatives with aliphatic side chains (51, 54) that

regulate various biological processes in vertebrates and their microbiomes (50, 54-57). The

amphipathic nature of bile salts allows them to interact with and disrupt membranes, resulting in

envelope stress in bacteria as detergents (51, 52). Both Gram-positive and Gram-negative

bacteria can respond to the presence of bile salts, leading to changes in gene expression and

cellular physiology (52, 58, 59). Generally, bile salts are more inhibitory of Gram-positive

bacteria because of the lack of an outer membrane. Thus, they are included in MacConkey as a

selective agent against Gram-positive bacteria in favor of enteric bacteria.

We supplemented LB agar with peptone, lactose, or bile salts at the same concentration

present in MacConkey agar to determine if one of these could enable *A. nosocomialis* to twitch

in LB. As shown in Figure 1A, the addition of neither lactose nor peptone led to any discernible

twitching motility in *A. nosocomialis*. In contrast, the supplementation of bile salts resulted in *A.*

nosocomialis twitching motility in LB comparable to what was observed with MacConkey agar.

These results indicated that bile salts are the component that specifically stimulates twitching

motility of *A. nosocomialis* as analyzed by stab assays with polystyrene Petri dishes.

Twitching motility has been observed in *A. baumannii* (24, 60), a closely related

Acinetobacter species and a WHO priority pathogen (16). The twitching motility of this

bacterium is similarly noted in MacConkey agar as analyzed by a similar assay (22, 49).

However, variable motility phenotypes were observed with LB media for different clinical

isolates (60), suggesting an effect of media composition on *A. baumannii* twitching. We tested

two *A. baumannii* strains, AYE and AB0057 (61, 62), in LB supplemented with bile salts in

comparison with *A. nosocomialis* M2. As shown in Figure 1B, while none of these strains

displayed twitching motility with the LB agar, supplementation of bile salts elicited twitching

motility of both *A. baumannii* strains similarly to *A. nosocomialis*. These results suggest that the

stimulatory effects of bile salts on twitching motility is a more general phenomenon in the

Acinetobacter genus.

Bile salts stimulated *P. aeruginosa* twitching motility. The above observations prompted us to

investigate if bile salts enhanced bacterial twitching in other bacteria. *P. aeruginosa*, another WHO

priority pathogen (16), has been used as a model for studies of bacterial twitching (8, 63, 64). Its

twitching motility has been routinely analyzed using stab assays with LB instead of MacConkey agar

plates (65-68). *P. aeruginosa* PAO1, a frequently used laboratory strain, exhibits twitching motility in

LB agar (69, 70). However, the addition of bile salts to LB significantly increased its twitching motility

(Figure 2A). Further, we examined the dose response of PAO1 twitching to bile salts. As shown in

Figure 2B, the stimulation of twitching motility shows concentration dependency, with a plateau

between 0.1% and 0.4% of bile salts. At higher concentrations, bile salts start to inhibit *P. aeruginosa*

growth and reduce its twitching motility in this assay (data not shown). We additionally examined the

twitching motility of PA14, another commonly used *P. aeruginosa* strain in the literature (71-73). It was

observed that the twitching motility of PA14 was stimulated by bile salts in LB media like that of PAO1

(Figure S1). These results demonstrate that the stimulatory effect of bile salts on twitching is applicable

to both *Acinetobacter* species and *P. aeruginosa* isolates. For the remainder of this study, we mostly

used *P. aeruginosa* PAO1 as the model organism to investigate the mechanisms by which bile salts

stimulate bacterial twitching motility.

Detergents stimulate bacterial twitching. Bile salts, produced from cholesterol metabolism, are

anionic detergents (50-52, 55). They are known to apply membrane or envelope stress in bacteria (51,

52, 74). It is possible that bile salts function as a detergent to apply envelope or general stress to cells

and it was the cellular stress response that underlies the stimulatory effects of bile salts on bacterial

twitching motility. To examine this possibility, we investigated the effect of other detergents on the

twitching motility of *P. aeruginosa.* To avoid the complications between growth inhibition and

twitching motility, we determined the maximum non-inhibitory concentrations of detergents

experimentally (Table S2) to guide their use in our twitching motility assays. For this experiment, we

supplemented the LB media with the anionic detergent sodium dodecyl sulfate (SDS), or the non-ionic

detergents Triton X-100 and Triton X-114 (Table S2). As shown in Figure 3A, all the detergents

examined, whether anionic or non-ionic, significantly stimulated the twitching motility of *P. aeruginosa*

much like bile salts. These results support the notion that the promotional effects of bile salts on

twitching are related to their amphipathic properties as detergents.

The stimulatory effects of bile salts and other detergents on twitching motility could be explained by

a physiological response of a bacterium to envelope stress applied by these amphipathic molecules (51,

52, 74, 75) or a general stress response to various environmental stressors (76). To investigate this, we

tested antibiotics with different modes of action at their maximum non-inhibitory concentrations as

stressors. These included ampicillin, gentamicin, and ciprofloxacin, which target cell wall biosynthesis,

ribosome function and DNA topology, respectively. We first determined the maximum non-inhibitory

concentrations of these antibiotics by testing their effect on *P. aeruginosa* growth at different

concentrations (Table S3). We then tested these antibiotics at their respective maximum non-inhibitory

concentrations for their effect on *P. aeruginosa* twitching. As shown in Figure 3B, none of these above

antibiotics affected *P. aeruginosa* twitching motility significantly. This suggested that the stimulation of

twitching by detergents was unlikely the result of a physiological response to general stressors.

Moreover, we tested the effect of polymyxin B, which applies envelope stress, as do bile salts, by

targeting the outer membrane of Gram-negative bacteria (53). Somewhat unexpectedly, this antibiotic

showed no stimulatory effect on *P. aeruginosa* twitching (Figure 3B). These results suggested that the

observed stimulation of twitching motility by bile salts and other detergents (Figures 2 and 3A) might

not be related to a response to general or envelope stress.

Glass enhances *P. aeruginosa* twitching in comparison to polystyrene. Detergents such as bile

salts are amphipathic molecules with both polar and non-polar moieties (50, 54, 56, 57, 77). As such,

they can change the physicochemical properties of a surface (50, 78-80). In stab assays for twitching

motility, bacteria cells translocate in the interstitial space between the solidified agar media and the

hydrophobic surface of a polystyrene Petri dish. We considered the possibility that bile salts in a growth

media may interact with the hydrophobic surface of the polystyrene Petri dishes to alter its

physicochemical properties. Such interactions may allow bile salts to make the polystyrene surface more

hydrophilic to possibly facilitate twitching motility. In comparison with polystyrene, glass Petri dishes

present a more hydrophilic surface. We therefore examined *P. aeruginosa* twitching with LB media

using glass in comparison with polystyrene Petri plates. As shown in Figure S2, *P. aeruginosa* was

observed to twitch significantly more on glass Petri dishes than polystyrene ones in LB without the

addition of bile salts. These results are consistent with the proposition that surface hydrophobicity or

hydrophilicity plays crucial roles in bacterial twitching.

Bile salts do not enhance *P. aeruginosa* twitching on glass surfaces. We reproduced the above

observation on glass (Figure S2) with a modified twitching assay where a glass or a polystyrene

microscope slide was used as the twitching surface (see Materials and Methods). In this assay, the slides

were cleaned and sterilized before they were placed in a polystyrene Petri dish. Molten LB agar media

was then poured into the Petri dish. Twitching motility was analyzed as before, except that the

incubation time was shortened to limit the twitching zone to be within the boundaries of the width of the

microscope slide. As shown in Figure 4A, PAO1 twitched significantly more on glass slides than on

polystyrene ones as was observed with Petri dishes (Figure S2). As expected, the addition of bile salts

significantly stimulated twitching on the polystyrene slide (Figure 4A). In contrast, the supplementation

of bile salts showed no promotional effect on twitching with the glass slide (Figure 4A). A *pilA* mutant,

which is non-piliated, was used as the non-twitching control, and it showed no twitching motility on all

surfaces with or without bile salts (Figure 4). It is also noteworthy that the twitching motility on the

polystyrene slide in the presence of bile salts showed no statistical difference from that on the glass

slides with or without bile salts. These results indicate that surface hydrophilicity likely enhances

twitching motility, and the effects of bile salts on twitching could be attributed to their ability to change

a hydrophobic surface to a more hydrophilic one. This is consistent with the observation that the

stimulatory effect of bile salts is no longer observed on the more hydrophilic glass surface in contrast

with polystyrene ones.

Increase in hydrophobicity of glass surfaces reduces *P. aeruginosa* twitching. Next, we modified

the surface of the glass slides to be more hydrophobic using a chemical treatment. For this, we

pretreated the glass slides with a polydimethylsiloxane (PDMS) solution before the analysis of twitching

motility. PDMS is known to coat glass surfaces to make them more hydrophobic (81). As shown in

Figure 4B, the treatment of the glass surface with PDMS significantly reduced *P. aeruginosa* twitching

to a level that is not significantly different from that on a polystyrene slide. In comparison, PDMS

treatment did not impact twitching motility of *P. aeruginosa* on polystyrene slides (Figure 4B), ruling

out any inhibitory effects by PDMS. The *P. aeruginosa pilA* mutant showed no twitching under all

experimental conditions as expected (Figure 4). These results are consistent with the idea that

hydrophilicity of surfaces enhance *P. aeruginosa* twitching, and that bile salts and other detergents

stimulate twitching motility on hydrophobic polystyrene surfaces by making them more hydrophilic.

Increase in hydrophilicity of polystyrene surfaces drastically enhances bacterial twitching

motility. While natural polystyrene surfaces are hydrophobic, they can be treated with plasma gas to

increase their hydrophilicity for tissue culture (TC) purposes (82). The surfaces of plasma- or TC-treated

plates are therefore more hydrophilic than non-treated ones. We compared *P. aeruginosa* twitching

motility with 6-well polystyrene plates either TC-treated or non-treated (Figure 5A). The *P. aeruginosa*

PAO1 strain exhibited significantly increased twitching motility on plasma-treated surfaces over the

non-treated ones in LB media (Figure 5A). The magnitude of increase in this case is about 2- to 3-fold.

This increase is more pronounced than on glass surfaces which led to an increase of 1-fold or less

(Figures 4A). While the addition of bile salts significantly enhanced *P. aeruginosa* twitching on

untreated plates, it had no stimulatory effect on twitching with the TC-treated surfaces.

We examined whether the drastic increase in twitching motility with TC-treated polystyrene surfaces

with *P. aeruginosa* could be extended to *A. nosocomialis*. As shown in Figure 5B, *A. nosocomialis* M2

displayed no twitching motility in LB media with the non-treated plates. This is expected because these

plates are made of polystyrene like the Petri dishes routinely used for twitching motility assays. TC-

treated plates drastically increased M2 twitching with LB media by almost 50-fold without the addition

of bile salts or detergents. An isogenic *pilA* mutant was used as a control, and it displayed no twitching

under all experimental treatments (Figure 5B). The results here indicate the enhancement of twitching

motility by hydrophilic surfaces is not confined to *P. aeruginosa.* Similar enhancement in *A.*

nosocomialis suggests a more general phenomenon where hydrophilic surfaces promote interactions that

are more favorable for bacterial twitching as mediated by the T4P as a motility apparatus.

Discussion

The interaction with surfaces is essential for the survival and proliferation of bacteria in

their natural environment as well as in health and disease. In their natural habitats, most bacteria

exist in multicellular ensembles known as biofilms, the establishment of which depends on

bacterial attachment to surfaces (83-86). During bacterial infection of a host, one of the earliest

steps in the process is the adhesion of a pathogen to the surfaces of host cells, tissues, and

medical implants. From a bacterial perspective, such interactions rely on the timely biogenesis

and proper functioning of adhesins on their surfaces. One of the structures critical for bacterial

adhesion to both biotic and abiotic surfaces is the bacterial T4P which is prevalent in both Gram-

positive and Gram-negative bacteria (2, 13, 28, 29, 31). It is an important virulence factor in

many pathogens including *P. aeruginosa* and *A. baumannii* which are both on the WHO priority

pathogens list (16). In these bacteria, as well as *A. nosocomialis* and others, the T4P is known to

power bacterial twitching motility, which provides a convenient assay for the investigation of

T4P biogenesis and function.

Here we described an unexpected mechanism by which bile salts and other detergents can

stimulate bacterial twitching motility. It was previously observed that *A. nosocomialis* exhibits

twitching motility with MacConkey but not Luria-Bertani (LB) agar media (16). This

phenomenon was observed using stab assays to visualize interstitial colony expansion with

polystyrene Petri dishes. After confirming this observation, we identified bile salts as the

component in MacConkey responsible for eliciting *A. nosocomialis* twitching motility (Figure

1A). The stimulatory effects of bile salts on twitching are not limited to *A. nosocomialis* as we

made similar observations with multiple strains of *A. baumannii* (Figure 1B) and *P. aeruginosa*

(Figures 2A and S1). Additionally, our results indicated that other detergents, whether anionic or

non-ionic, likewise promoted *P. aeruginosa* twitching motility (Figure 3A). Surprisingly, this

stimulation of twitching is likely not due to a physiological change in response to the presence of

bile salts and other detergents in the growth medium. Instead, it is the ability of detergents to

alter the physicochemical properties of a surface that enhances twitching motility in multiple

bacterial species.

The above conclusion is based on a few lines of experimental evidence from this study.

First, antibiotics with various modes of actions failed to enhance *P. aeruginosa* twitching

motility (Figure 3B). These included polymyxin B, which mimics the effects of bile salts and

other detergents in applying envelope stress. These results suggested that the enhancement of

twitching motility by detergents is likely not a bacterial response to a general or envelope

stressor in a growth media. Second, we observed that surfaces of glass, which are more

hydrophilic than that of polystyrene, significantly enhanced twitching motility (Figures S2 and

4A). We further demonstrated that the use of glass surfaces abrogated the stimulatory effect of

bile salts such that the addition of bile salts no longer promoted *P. aeruginosa* twitching motility

on these surfaces (Figure 4A). When the surface properties of glass were changed by the

application of a hydrophobic coating, the enhancement of bacterial twitching by glass was

reversed (Figure 4B). These results show that hydrophilic surfaces promote twitching, whereas

hydrophobic ones suppress it. Because bile salts in the growth media only enhance twitching

motility on polystyrene but not on glass surfaces, we conclude that bile salts likely function to

modify natural polystyrene surfaces to be more hydrophilic, which promote bacterial motility.

Lastly, we performed experiments with TC-treated and nontreated polystyrene surfaces (Figure

5). The TC-treated surfaces, which are more hydrophilic, significantly enhanced bacterial

twitching motility. As similarly observed on glass surfaces, the addition of bile salts no longer

displayed a stimulatory effect on *P. aeruginosa* and *A. nosocomialis* twitching on TC-treated

plates. These results support our conclusion that the physicochemical properties of a surface

significantly impact the effectiveness of T4P-powered twitching motility in bacteria. On

hydrophilic surfaces, bacteria twitch more, and on hydrophobic surfaces, they twitch less. We

further conclude that the promotional effects of bile salts and other detergents on bacterial

twitching are largely due to their ability to change the properties of a surface over which bacteria

translocate by twitching motility.

Categorically, there are two possible mechanistic explanations for the observed effects of

the physicochemical properties of a surface on bacterial twitching. There have been reports that

bacteria attach better to hydrophobic surfaces in the context of biofilm formation or otherwise

(87-90). It follows that reducing hydrophobicity may lead to alteration of the interactions of a

bacterial cell or its pilus with a subsurface over which a bacterium translocates by twitching

motility. Sustained twitching movement relies on the recurrence of a multi-step process. These

steps include the unobstructed extension of a T4P, the subsequent attachment of the pilus

through its distal end for anchoring, followed by a successful T4P retraction event. It is

conceivable that tampering with any of these steps through physicochemical interactions with a

surface can lead to changes in bacterial twitching behaviors. Alternatively, surface sensing has

been demonstrated to mediate changes in cell physiology and behavior (6, 91-96). It is possible

that the physicochemical properties of a surface may be detected by a bacterium through surface

sensing to modulate T4P biogenesis and its functions, leading to alterations in twitching motility.

Further investigation is necessary to determine if the above scenarios or others, either alone or in

combination, are the underlying reasons for the observed enhancement of T4P-powered bacterial

twitching by surface hydrophilicity.

Materials and Methods

Strains and culture conditions. The bacterial strains used in this study are listed in Table 1. These

include *A. nosocomialis* M2, *A. baumannii* strains Ab0057 and AYE, and *P. aeruginosa* strains PAO1

and PA14. When appropriate, isogenic *pilA* mutants were used as controls. *P. aeruginosa* strains were

grown at 37°C on 1.5% Luria-Bertani agar (LBA) while *A. nosocomialis* and *A. baumannii* strains were

grown at 37°C on 1.5% MacConkey agar (Oxoid). Oxoid bile acids were used in this study when

indicated.

Twitching motility assays. Twitching motility was analyzed by three different protocols using the agar

stab methods (27) (97) with 1.2% agar media. For all stab assays, the inoculum was prepared by

spreading a loopful of bacteria from the outside edge of an overnight plate culture onto a fresh 1.5%

LBA plate to create a thin layer of cells. Using a toothpick, bacteria were picked up to inoculate a plate

by stabbing through the agar to touch either the polystyrene or glass surface below.

The first protocol uses a standard 100 mm × 15 mm polystyrene or glass Petri dish (Fisher

Scientific) as previously described (27, 97). In brief, plates with 25 ml agar media were prepared a day

before the assays. These plates were then dried in a biosafety cabinet for 20 minutes before stab

inoculation. After 48 hours of incubation at 37°C in a humidity chamber, the agar media was removed,

and the twitching zone was visualized by staining with 1% crystal violet. Twitching areas were

determined using the NIH ImageJ software (98).

The second protocol uses either a glass (Opto-Edu) or a polystyrene (VWR International)

microscope slide (1″ × 3″) inside a standard polystyrene Petri dish for analyzing twitching motility. The

microscope slides were first submerged in a filter-sterilized polydimethylsiloxane (PDMS) solution

(RainX) (99) or in 70% ethanol as a control. These slides were air dried on a rack at 40°C before being

placed at the bottom of a polystyrene Petri dish. 25 ml of molten agar media was poured over the

microscope slides into the Petri dish a day before. A twitching assay is initiated by stab-inoculation as

described above except that the incubation is shortened to 18 hours to limit the twitching zone within the

boundaries of the microscope slide. One additional modification is that the twitching zone in this case

was visualized by incident light and traced with a permanent marker without removing the agar from the

Petri dish. Twitching area was determined using ImageJ as above.

The third protocol uses 6-well polystyrene plates with or without TC-treatment (Falcon). Each

well of a plate contains 2 ml agar media and twitching motility assays were initiated by stab inoculation

as described above. The twitching area was analyzed as described for the microscope slide-based assay

except that the incubation time was extended to 24 hours.

Acknowledgements

The research in the Yang laboratory is partially supported by the National Science Foundation grants MCB-1919455 and EFMA-2318093, as well as the CHRB award 208-07-23 to Z.Y. The Yang laboratory additionally received a pilot grant from the Center for Emerging, Zoonotic and Arthropod-borne Pathogens. M.T.O. was the recipient of the Presidential Scholarship Initiative from Virginia Tech and the David Lyerly Undergraduate Microbiology Research Fellowship from the Department of Biological Sciences at Virginia Tech. K.J.D was the recipient of a GSDA and the Lewis Edward Goyette Fellowships as well as the Liberati Scholarship from Virginia Tech.

M.T.O., K.J.D., and Z.Y. designed research and analyzed data. M.T.O. and Z.Y wrote the manuscript. M.T.O., T.M.S., and D.H. performed experiments. T.M.S. aided in figure preparation and protocol development.

We thank Dr. Kurt Piepenbrink, Dr. Robert Bonomo, Dr. Helen Zgurskara, and Dr. George O'Toole for providing bacterial strains. We additionally thank Ghazal Soleymani for proof-reading the manuscript.

There is no conflict of interest to declare.

References

391   1.   Craig L, Forest KT, Maier B. 2019. Type IV pili: dynamics, biophysics and functional
392   consequences. Nat Rev Microbiol 17:429-440.
393   2.   Wall D, Kaiser D. 1999. Type IV pili and cell motility. Mol Microbiol 32:1-10.
394   3.   Mattick JS. 2002. Type IV pili and twitching motility. Annu Rev Microbiol 56:289-314.
395   4.   Shi W, Sun H. 2002. Type IV pilus-dependent motility and its possible role in bacterial
396   pathogenesis. Infect Immun 70:1-4.
397   5.   Chang YW, Rettberg LA, Treuner-Lange A, Iwasa J, Sogaard-Andersen L, Jensen GJ.
398   2016. Architecture of the type IVa pilus machine. Science 351:aad2001.
399   6.   Chang CY. 2017. Surface sensing for biofilm formation in *Pseudomonas aeruginosa*.
400   Front Microbiol 8:2671.
401   7.   Ligthart K, Belzer C, de Vos WM, Tytgat HLP. 2020. Bridging bacteria and the gut:
402   functional aspects of type IV pili. Trends Microbiol 28:340-348.
403   8.   Tala L, Fineberg A, Kukura P, Persat A. 2019. *Pseudomonas aeruginosa* orchestrates
404   twitching motility by sequential control of type IV pili movements. Nat Microbiol 4:774-
405   780.
406   9.   Wadhwa N, Berg HC. 2022. Bacterial motility: machinery and mechanisms. Nat Rev
407   Microbiol 20:161-173.
408   10.   Craig L, Pique ME, Tainer JA. 2004. Type IV pilus structure and bacterial pathogenicity.
409   Nat Rev Microbiol 2:363-78.
410   11.   McKee RW, Aleksanyan N, Garrett EM, Tamayo R. 2018. Type IV pili promote
411   *Clostridium difficile* adherence and persistence in a mouse model of infection. Infect
412   Immun 86.
413   12.   Dos Santos Souza I, Maissa N, Ziveri J, Morand PC, Coureuil M, Nassif X, Bourdoulous
414   S. 2020. Meningococcal disease: A paradigm of type-IV pilus dependent pathogenesis.
415   Cell Microbiol 22:e13185.
416   13.   Martini AM, Moricz BS, Woods LJ, Jones BD. 2021. Type IV pili of *Streptococcus*
417   *sanguinis* contribute to pathogenesis in experimental infective endocarditis. Microbiol
418   Spectr 9:e0175221.
419   14.   Persat A, Inclan YF, Engel JN, Stone HA, Gitai Z. 2015. Type IV pili
420   mechanochemically regulate virulence factors in *Pseudomonas aeruginosa*. Proc Natl
421   Acad Sci U S A 112:7563-8.
422   15.   Inclan YF, Persat A, Greninger A, Von Dollen J, Johnson J, Krogan N, Gitai Z, Engel JN.
423   2016. A scaffold protein connects type IV pili with the Chp chemosensory system to
424   mediate activation of virulence signaling in *Pseudomonas aeruginosa*. Mol Microbiol
425   101:590-605.
426   16.   WHO. 2017. WHO publishes list of bacteria for which new antibiotics are urgently
427   needed, *on* WHO. Accessed
428   17.   Colquhoun JM, Rather PN. 2020. Insights into mechanisms of biofilm formation in
429   *Acinetobacter baumannii* and implications for uropathogenesis. Front Cell Infect
430   Microbiol 10:253.
431   18.   Piepenbrink KH, Lillehoj E, Harding CM, Labonte JW, Zuo X, Rapp CA, Munson RS,
432   Jr., Goldblum SE, Feldman MF, Gray JJ, Sundberg EJ. 2016. Structural diversity in the
433   type IV pili of multidrug-resistant *Acinetobacter*. J Biol Chem 291:22924-22935.

19. Chusri S, Chongsuvivatwong V, Rivera JI, Silpapojakul K, Singkhamanan K, McNeil E,
Doi Y. 2014. Clinical outcomes of hospital-acquired infection with *Acinetobacter*
nosocomialis and *Acinetobacter pittii*. Antimicrob Agents Chemother 58:4172-9.
20. Harding CM, Kinsella RL, Palmer LD, Skaar EP, Feldman MF. 2016. Medically relevant
Acinetobacter species require a type II secretion system and specific membrane-
associated chaperones for the export of multiple substrates and full virulence. PLoS
Pathog 12:e1005391.
21. Harding CM, Nasr MA, Kinsella RL, Scott NE, Foster LJ, Weber BS, Fiester SE, Actis
LA, Tracy EN, Munson RS, Jr., Feldman MF. 2015. *Acinetobacter* strains carry two
functional oligosaccharyltransferases, one devoted exclusively to type IV pilin, and the
other one dedicated to O-glycosylation of multiple proteins. Mol Microbiol 96:1023-41.
22. Vo N, Sidner BS, Yu Y, Piepenbrink KH. 2023. Type IV pilus-mediated inhibition of
Acinetobacter baumannii biofilm formation by phenothiazine compounds. Microbiol
Spectr 11:e0102323.
23. Harding CM, Tracy EN, Carruthers MD, Rather PN, Actis LA, Munson RS, Jr. 2013.
Acinetobacter baumannii strain M2 produces type IV pili which play a role in natural
transformation and twitching motility but not surface-associated motility. mBio 4.
24. Corral J, Perez-Varela M, Sanchez-Osuna M, Cortes P, Barbe J, Aranda J. 2021.
Importance of twitching and surface-associated motility in the virulence of *Acinetobacter*
baumannii. Virulence 12:2201-2213.
25. Biswas I, Machen A, Mettlach J. 2019. *In vitro* motility assays for *Acinetobacter* species.
Methods Mol Biol 1946:177-187.
26. Leong CG, Bloomfield RA, Boyd CA, Dornbusch AJ, Lieber L, Liu F, Owen A, Slay E,
Lang KM, Lostroh CP. 2017. The role of core and accessory type IV pilus genes in
natural transformation and twitching motility in the bacterium *Acinetobacter baylyi*.
PLoS One 12:e0182139.
27. Turnbull L, Whitchurch CB. 2014. Motility assay: twitching motility. Methods Mol Biol
1149:73-86.
28. Melville S, Craig L. 2013. Type IV pili in Gram-positive bacteria. Microbiol Mol Biol
Rev 77:323-41.
29. Piepenbrink KH, Sundberg EJ. 2016. Motility and adhesion through type IV pili in Gram-
positive bacteria. Biochem Soc Trans 44:1659-1666.
30. Henrichsen J. 1975. The occurrence of twitching motility among Gram-negative bacteria.
Acta Pathol Microbiol Scand B 83:171-8.
31. Pelicic V. 2008. Type IV pili: e pluribus unum? Mol Microbiol 68:827-37.
32. Chilcott GS, Hughes KT. 2000. Coupling of flagellar gene expression to flagellar
assembly in *Salmonella enterica* serovar *typhimurium* and *Escherichia coli*. Microbiol
Mol Biol Rev 64:694-708.
33. Kalir S, McClure J, Pabbaraju K, Southward C, Ronen M, Leibler S, Surette MG, Alon
U. 2001. Ordering genes in a flagella pathway by analysis of expression kinetics from
living bacteria. Science 292:2080-3.
34. Utsey K, Keener JP. 2020. A mathematical model of flagellar gene regulation and
construction in *Salmonella enterica*. PLoS Comput Biol 16:e1007689.
35. Zhu S, Kojima S, Homma M. 2013. Structure, gene regulation and environmental
response of flagella in *Vibrio*. Front Microbiol 4:410.

36. Wadhams GH, Armitage JP. 2004. Making sense of it all: bacterial chemotaxis. Nat Rev
Mol Cell Biol 5:1024-37.
37. Ancona V, Li W, Zhao Y. 2014. Alternative sigma factor RpoN and its modulation
protein YhbH are indispensable for *Erwinia amylovora* virulence. Mol Plant Pathol
15:58-66.
38. Sporing I, Felgner S, Preusse M, Eckweiler D, Rohde M, Haussler S, Weiss S, Erhardt
M. 2018. Regulation of flagellum biosynthesis in response to cell envelope stress in
Salmonella enterica serovar *Typhimurium*. mBio 9.
39. Tague JG, Hong J, Kalburge SS, Boyd EF. 2022. Regulatory small RNA Qrr2 is
expressed independently of sigma factor-54 and can function as the sole Qrr small RNA
to control quorum sensing in *Vibrio parahaemolyticus*. J Bacteriol 204:e0035021.
40. Zhou Y, Lepp D, Carere J, Yu H, Yang C, Gong J. 2021. A novel PilR/PilS two-
component system regulates necrotic enteritis pilus production in *Clostridium*
perfringens. J Bacteriol 203:e0009621.
41. Xu K, Shen D, Yang N, Chou SH, Gomelsky M, Qian G. 2021. Coordinated control of
the type IV pili and c-di-GMP-dependent antifungal antibiotic production in *Lysobacter*
by the response regulator PilR. Mol Plant Pathol 22:602-617.
42. Hobbs M, Collie ES, Free PD, Livingston SP, Mattick JS. 1993. PilS and PilR, a two-
component transcriptional regulatory system controlling expression of type 4 fimbriae in
Pseudomonas aeruginosa. Mol Microbiol 7:669-82.
43. Wu SS, Kaiser D. 1995. Genetic and functional evidence that type IV pili are required for
social gliding motility in *Myxococcus xanthus*. Mol Microbiol 18:547-58.
44. Vesel N, Blokesch M. 2021. Pilus production in *Acinetobacter baumannii* is growth
phase dependent and essential for natural transformation. J Bacteriol 203.
45. Mattingly AE, Weaver AA, Dimkovikj A, Shrout JD. 2018. Assessing travel conditions:
environmental and host influences on bacterial surface motility. J Bacteriol 200:e00014-
18.
46. Huang B, Whitchurch CB, Mattick JS. 2003. FimX, a multidomain protein connecting
environmental signals to twitching motility in *Pseudomonas aeruginosa*. J Bacteriol
185:7068-76.
47. Sigurlasdottir S, Lidberg K, Zuo F, Newcombe J, McFadden J, Jonsson AB. 2021.
Lactate-induced dispersal of *Neisseria meningitidis* microcolonies is mediated by
changes in cell density and pilus retraction and is influenced by temperature change.
Infect Immun 89:e0029621.
48. Bitrian M, Gonzalez RH, Paris G, Hellingwerf KJ, Nudel CB. 2013. Blue-light-
dependent inhibition of twitching motility in *Acinetobacter baylyi* ADP1: additive
involvement of three BLUF-domain-containing proteins. Microbiology (Reading)
159:1828-1841.
49. Ronish LA, Lillehoj E, Fields JK, Sundberg EJ, Piepenbrink KH. 2019. The structure of
PilA from *Acinetobacter baumannii* AB5075 suggests a mechanism for functional
specialization in *Acinetobacter* type IV pili. J Biol Chem 294:218-230.
50. Monte MJ, Marin JJ, Antelo A, Vazquez-Tato J. 2009. Bile acids: chemistry, physiology,
and pathophysiology. World J Gastroenterol 15:804-16.
51. Urdaneta V, Casadesus J. 2017. Interactions between bacteria and bile salts in the
gastrointestinal and hepatobiliary tracts. Front Med (Lausanne) 4:163.

52.   Begley M, Gahan CG, Hill C. 2005. The interaction between bacteria and bile. FEMS
Microbiol Rev 29:625-51.

53.   Velkov T, Roberts KD, Nation RL, Thompson PE, Li J. 2013. Pharmacology of
polymyxins: new insights into an 'old' class of antibiotics. Future Microbiol 8:711-24.

54.   Hofmann AF, Hagey LR, Krasowski MD. 2010. Bile salts of vertebrates: structural
variation and possible evolutionary significance. J Lipid Res 51:226-46.

55.   Winston JA, Theriot CM. 2020. Diversification of host bile acids by members of the gut
microbiota. Gut Microbes 11:158-171.

56.   Fleishman JS, Kumar S. 2024. Bile acid metabolism and signaling in health and disease:
molecular mechanisms and therapeutic targets. Signal Transduct Target Ther 9:97.

57.   de Buy Wenniger LM, Tusl T, Beuers U. 2013. Bile salts, p 159-163. *In* Lennarz WJ,
Lane MD (ed), Encyclopedia of Biological Chemistry II, 2nd ed. Academic Press.

58.   Ridlon JM, Gaskins HR. 2024. Another renaissance for bile acid gastrointestinal
microbiology. Nat Rev Gastroenterol Hepatol 21:348-364.

59.   Larabi AB, Masson HLP, Baumler AJ. 2023. Bile acids as modulators of gut microbiota
composition and function. Gut Microbes 15:2172671.

60.   Vijayakumar S, Rajenderan S, Laishram S, Anandan S, Balaji V, Biswas I. 2016. Biofilm
formation and motility depend on the nature of the *Acinetobacter baumannii* clinical
isolates. Front Public Health 4:105.

61.   Hamidian M, Venepally P, Hall RM, Adams MD. 2017. Corrected genome sequence of
Acinetobacter baumannii strain AB0057, an antibiotic-resistant isolate from lineage 1 of
global clone 1. Genome Announc 5.

62.   Adams MD, Goglin K, Molyneaux N, Hujer KM, Lavender H, Jamison JJ, MacDonald
IJ, Martin KM, Russo T, Campagnari AA, Hujer AM, Bonomo RA, Gill SR. 2008.
Comparative genome sequence analysis of multidrug-resistant *Acinetobacter baumannii*.
J Bacteriol 190:8053-64.

63.   Kuhn MJ, Tala L, Inclan YF, Patino R, Pierrat X, Vos I, Al-Mayyah Z, Macmillan H,
Negrete J, Jr., Engel JN, Persat A. 2021. Mechanotaxis directs *Pseudomonas aeruginosa*
twitching motility. Proc Natl Acad Sci U S A 118.

64.   Burrows LL. 2012. *Pseudomonas aeruginosa* twitching motility: type IV pili in action.
Annu Rev Microbiol 66:493-520.

65.   Wen F, Wu Y, Yuan Y, Yang X, Ran Q, Gan X, Guo Y, Wang X, Chu Y, Zhao K. 2024.
Discovery of psoralen as a quorum sensing inhibitor suppresses *Pseudomonas aeruginosa*
virulence. Appl Microbiol Biotechnol 108:222.

66.   Pei TT, Luo H, Wang Y, Li H, Wang XY, Zhang YQ, An Y, Wu LL, Ma J, Liang X, Yan
A, Yang L, Chen C, Dong T. 2024. Filamentous prophage Pf4 promotes genetic
exchange in *Pseudomonas aeruginosa*. ISME J 18.

67.   Schroven K, Putzeys L, Kerremans A, Ceyssens PJ, Vallino M, Paeshuyse J, Haque F,
Yusuf A, Koch MD, Lavigne R. 2023. The phage-encoded PIT4 protein affects multiple
two-component systems of *Pseudomonas aeruginosa*. Microbiol Spectr 11:e0237223.

68.   Gur M, Erdmann J, Will A, Liang Z, Andersen JB, Tolker-Nielsen T, Haussler S. 2023.
Challenges in using transcriptome data to study the c-di-GMP signaling network in
Pseudomonas aeruginosa clinical isolates. FEMS Microbes 4:xtad012.

69.   Kearns DB, Robinson J, Shimkets LJ. 2001. *Pseudomonas aeruginosa* exhibits directed
twitching motility up phosphatidylethanolamine gradients. J Bacteriol 183:763-7.

70. Li J, Wan SJ, Metruccio MME, Ma S, Nazmi K, Bikker FJ, Evans DJ, Fleiszig SMJ.
2019. DMBT1 inhibition of *Pseudomonas aeruginosa* twitching motility involves its N-
glycosylation and cannot be conferred by the scavenger receptor cysteine-rich bacteria-
binding peptide domain. Sci Rep 9:13146.
71. He J, Baldini RL, Deziel E, Saucier M, Zhang Q, Liberati NT, Lee D, Urbach J,
Goodman HM, Rahme LG. 2004. The broad host range pathogen *Pseudomonas*
aeruginosa strain PA14 carries two pathogenicity islands harboring plant and animal
virulence genes. Proc Natl Acad Sci U S A 101:2530-5.
72. O'Toole GA, Kolter R. 1998. Flagellar and twitching motility are necessary for
Pseudomonas aeruginosa biofilm development. Mol Microbiol 30:295-304.
73. Nolan LM, McCaughey LC, Merjane J, Turnbull L, Whitchurch CB. 2020. ChpC
controls twitching motility-mediated expansion of *Pseudomonas aeruginosa* biofilms in
response to serum albumin, mucin and oligopeptides. Microbiology (Reading) 166:669-
678.
74. Heuman DM. 1995. Bile salt-membrane interactions and the physico-chemical
mechanisms of bile salt toxicity. Ital J Gastroenterol 27:372-5.
75. Henriksen JR, Andresen TL, Feldborg LN, Duelund L, Ipsen JH. 2010. Understanding
detergent effects on lipid membranes: a model study of lysolipids. Biophys J 98:2199-
205.
76. Gipson KS, Nickerson KP, Drenkard E, Llanos-Chea A, Dogiparthi SK, Lanter BB,
Hibbler RM, Yonker LM, Hurley BP, Faherty CS. 2020. The great ESKAPE: exploring
the crossroads of bile and antibiotic resistance in bacterial pathogens. Infect Immun 88.
77. Camilleri M. 2022. Bile acid detergency: permeability, inflammation, and effects of
sulfation. Am J Physiol Gastrointest Liver Physiol 322:G480-G488.
78. Forooqi Motlaq V, Ortega-Holmberg M, Edwards K, Gedda L, Lyngso J, Pedersen JS,
Bergstrom LM. 2021. Investigation of the enhanced ability of bile salt surfactants to
solubilize phospholipid bilayers and form mixed micelles. Soft Matter 17:7769-7780.
79. di Gregorio MC, Travaglini L, Del Giudice A, Cautela J, Pavel NV, Galantini L. 2019.
Bile salts: natural surfactants and precursors of a broad family of complex amphiphiles.
Langmuir 35:6803-6821.
80. Pigliacelli C, Belton P, Wilde P, Qi S. 2019. Probing the molecular interactions between
pharmaceutical polymeric carriers and bile salts in simulated gastrointestinal fluids using
NMR spectroscopy. J Colloid Interface Sci 551:147-154.
81. Kaneda S, Ono K, Fukuba T, Nojima T, Yamamoto T, Fujii T. 2012. Modification of the
glass surface property in PDMS-glass hybrid microfluidic devices. Anal Sci 28:39-44.
82. Lerman MJ, Lembong J, Muramoto S, Gillen G, Fisher JP. 2018. The evolution of
polystyrene as a cell culture material. Tissue Eng Part B Rev 24:359-372.
83. Kreve S, Reis ACD. 2021. Bacterial adhesion to biomaterials: What regulates this
attachment? A review. Jpn Dent Sci Rev 57:85-96.
84. Muhammad MH, Idris AL, Fan X, Guo Y, Yu Y, Jin X, Qiu J, Guan X, Huang T. 2020.
Beyond risk: bacterial biofilms and their regulating approaches. Front Microbiol 11:928.
85. Ciolacu L, Zand E, Negrau C, Jaeger H. 2022. Bacterial attachment and biofilm
formation on antimicrobial sealants and stainless steel surfaces. Foods 11.
86. Palmer J, Flint S, Brooks J. 2007. Bacterial cell attachment, the beginning of a biofilm. J
Ind Microbiol Biotechnol 34:577-88.

87.   De-la-Pinta I, Cobos M, Ibarretxe J, Montoya E, Eraso E, Guraya T, Quindos G. 2019.
615   Effect of biomaterials hydrophobicity and roughness on biofilm development. J Mater Sci
616   Mater Med 30:77.
617   88.   Krasowska A, Sigler K. 2014. How microorganisms use hydrophobicity and what does
618   this mean for human needs? Front Cell Infect Microbiol 4:112.
619   89.   Mirani ZA, Fatima A, Urooj S, Aziz M, Khan MN, Abbas T. 2018. Relationship of cell
620   surface hydrophobicity with biofilm formation and growth rate: A study on *Pseudomonas*
621   *aeruginosa, Staphylococcus aureus,* and *Escherichia coli*. Iran J Basic Med Sci 21:760-
622   769.
623   90.   Khatoon Z, McTiernan CD, Suuronen EJ, Mah TF, Alarcon EI. 2018. Bacterial biofilm
624   formation on implantable devices and approaches to its treatment and prevention.
625   Heliyon 4:e01067.
626   91.   Chuang SK, Vrla GD, Frohlich KS, Gitai Z. 2019. Surface association sensitizes
627   *Pseudomonas aeruginosa* to quorum sensing. Nat Commun 10:4118.
628   92.   Ellison CK, Kan J, Dillard RS, Kysela DT, Ducret A, Berne C, Hampton CM, Ke Z,
629   Wright ER, Biais N, Dalia AB, Brun YV. 2017. Obstruction of pilus retraction stimulates
630   bacterial surface sensing. Science 358:535-538.
631   93.   Snyder RA, Ellison CK, Severin GB, Whitfield GB, Waters CM, Brun YV. 2020. Surface
632   sensing stimulates cellular differentiation in *Caulobacter crescentus*. Proc Natl Acad Sci
633   U S A 117:17984-17991.
634   94.   Zheng S, Bawazir M, Dhall A, Kim HE, He L, Heo J, Hwang G. 2021. Implication of
635   surface properties, bacterial motility, and hydrodynamic conditions on bacterial surface
636   sensing and their initial adhesion. Front Bioeng Biotechnol 9:643722.
637   95.   Laventie BJ, Jenal U. 2020. Surface sensing and adaptation in bacteria. Annu Rev
638   Microbiol 74:735-760.
639   96.   Hofer U. 2018. Bacterial physiology: Touching base on bacterial surface sensing. Nat
640   Rev Microbiol 16:3.
641   97.   Dye KJ, Vogelaar NJ, O'Hara M, Sobrado P, Santos W, Carlier PR, Yang Z. 2022.
642   Discovery of two inhibitors of the type IV pilus assembly ATPase PilB as potential
643   antivirulence compounds. Microbiol Spectr 10:e0387722.
644   98.   Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of
645   image analysis. Nat Methods 9:671-5.
646   99.   Shea TM. 2008. Durable hydrophobic surface coatings using silicone resins. USA.
647   100.   Niu C, Clemmer KM, Bonomo RA, Rather PN. 2008. Isolation and characterization of an
648   autoinducer synthase from *Acinetobacter baumannii*. J Bacteriol 190:3386-92.
649   101.   Stover CK, Pham XQ, Erwin AL, Mizoguchi SD, Warrener P, Hickey MJ, Brinkman FS,
650   Hufnagle WO, Kowalik DJ, Lagrou M, Garber RL, Goltry L, Tolentino E, Westbrock-
651   Wadman S, Yuan Y, Brody LL, Coulter SN, Folger KR, Kas A, Larbig K, Lim R, Smith
652   K, Spencer D, Wong GK, Wu Z, Paulsen IT, Reizer J, Saier MH, Hancock RE, Lory S,
653   Olson MV. 2000. Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an
654   opportunistic pathogen. Nature 406:959-64.
655   102.   Held K, Ramage E, Jacobs M, Gallagher L, Manoil C. 2012. Sequence-verified two-allele
656   transposon mutant library for *Pseudomonas aeruginosa* PAO1. J Bacteriol 194:6387-9.
657

658   **Table 1. Bacterial strains used in this study.**

Strain		Description	Reference
Acinetobacter nosocomialis	M2	Clinical isolate	(100)
	M2 Δ*pilA*	*pilA* deletion and insertion of kanamycin resistant cassette (Δ*pilA*::kan)	(23)
Acinetobacter baumannii	Ab0057	Clinical isolate	(61)
	AYE	Clinical isolate	(62)
Pseudomonas aeruginosa	PAO1	Wildtype	(101)
	PW8622	PAO1 *pilA*-H02::ISphoA/hah	(102)

Figure Legends

Figure 1. Bile salts enable *Acinetobacter* to twitch. A. Bile salts allow *A. nosocomialis* M2 to

twitch in LB media. The twitching motility of *A. nosocomialis* M2 was analyzed with

MacConkey (MC) or Luria-Bertani (LB) media without or with 0.5% bile salts (+BSs), 1%

lactose (+Lac), or 2% peptone (+Pep) with standard polystyrene Petri dishes as described in

Materials and Methods. Data shown are the averages from three biological experiments each

performed in triplicate. **B.** Bile salts provoke *A. baumannii* twitching in LB media. *A. baumannii*

strains AB0057 and AYE were analyzed for twitching motility in LB agar without (-) or with (+)

0.5% bile salts on standard polystyrene Petri dishes with *A. nosocomialis* (M2) as a control. Data

shown are from three biological experiments, represented by different symbols, each performed

in triplicate. Single (*), double (**), and quadruple (****) asterisks indicate two values are

statistically different with P<0.05, P<0.01, and P<0.0001 by the Student's T-test, respectively.

Figure 2. Bile salts enhance *P. aeruginosa* twitching motility. A. Bile salts increase *P.*

aeruginosa twitching. Twitching motility of PAO1 was analyzed without (-) or with (+) 0.5%

bile salts as in Figure 1B with data similarly presented. Quadruple asterisks (****) indicates

values that are statistically different with P<0.0001 by the Student's T-test. **B.** Dose effect of bile

salts on PAO1 twitching. PAO1 twitching was analyzed with the standard polystyrene Petri dish

protocol as in (A) with varying concentrations (w/v %) of bile salts as indicated. Data presented

in both panels are from three biological experiments each performed in triplicate. Data points

from the same experiment are represented by the same symbols in color and shape.

Figure 3. Detergents, but not antibiotics, promote *P. aeruginos*a twitching. A. Effects of

detergents. PAO1 twitching was analyzed with the standard Petri dish protocol as in Figure 2A

with LB agar without modification (-) or with bile salts (BSs) (5 mg/ml), Triton X-100 (TX100)

(75 µg/ml), Triton X-114 (TX114) (75 µg/ml), or sodium dodecyl sulfate (SDS) (850 µg/ml). **B.**

Effects of antibiotics. PAO1 twitching was analyzed as in (A) with ampicillin (Amp) (313

ng/ml), ciprofloxacin (Cipro) (31 ng/ml), gentamicin (Gent) (31 ng/ml), or polymyxin B (PB)

(313 ng/ml). Data presented in both panels are from three biological experiments each

performed in triplicate. Data points from the same experiment are represented by the same

symbols in color and shape. Triple (***) and quadruple (****) asterisks indicate two values are

statistically different with $P<0.001$ and $P<0.0001$ by the Student's T-test, respectively.

Antibiotics resulted in values that are not significantly (ns) different with $P>0.05$.

694
Figure 4. Glass surfaces increase *P. aeruginosa* twitching motility. A. Glass surfaces

stimulate PAO1 twitching motility and eliminate the stimulatory effect of bile salts. The

twitching motility of PAO1 and its isogenic *pilA* mutant were analyzed with polystyrene (PS) or

glass microscope slides in LB without or with 0.5% bile salts (BSs) after 18 hours of incubation

(see Materials and Methods). **B.** Hydrophobic coating of glass reduces twitching motility.

Experiments were performed as in A, except that the microscope slides were coated without or

with polydimethylsiloxane (PDMS) (see Materials and Methods). Data presented in both panels

are from three biological experiments each performed in triplicate. Double asterisks (**) indicate

two values are statistically different with $P<0.01$ by the Student's T-test.

704
Figure 5. TC-treated plates ameliorate twitching and abolish the effects of bile salts. A. *P.*

aeruginosa twitching on TC-treated polystyrene. The twitching motility of *P. aeruginosa* PAO1

and its isogenic *pilA* mutant strain was analyzed with 6-well polystyrene plates (see Materials

and Methods) either TC-treated or non-treated in LB agar without or with 0.5% bile salts (BSs).

B. *A. nosocomialis* twitching on TC-treated polystyrene. The twitching motility of *A.*

nosocomialis M2 and its isogenic *pilA* deletion mutant was analyzed as in A. Data presented in

both panels are from three biological experiments each performed in triplicate. Double (**) and

quadruple (****) asterisks indicate that two values are statistically different with P<0.01 and

P<0.0001 by the Student's T-test, respectively.

Figure 1. Bile salts enable *Acinetobacter* to twitch. A. Bile salts allow *A. nosocomialis* M2 to twitch in LB media. The twitching motility of *A. nosocomialis* M2 was analyzed with MacConkey (MC) or Luria-Bertani (LB) media without or with 0.5% bile salts (+BSs), 1% lactose (+Lac), or 2% peptone (+Pep) with standard polystyrene Petri dishes as described in Materials and Methods. Data shown are the averages from three biological experiments each performed in triplicate. **B.** Bile salts provoke *A. baumannii* twitching in LB media. *A. baumannii* strains AB0057 and AYE were analyzed for twitching motility in LB agar without (-) or with (+) 0.5% bile salts on standard polystyrene Petri dishes with *A. nosocomialis* (M2) as a control. Data shown are from three biological experiments, represented by different symbols, each performed in triplicate. Single (*), double (**), and quadruple (****) asterisks indicate two values are statistically different with P<0.05, P<0.01, and P<0.0001 by the Student's T-test, respectively.

Figure 2. Bile salts enhance *P. aeruginosa* twitching motility. A. Bile salts increase *P. aeruginosa* twitching. Twitching motility of PAO1 was analyzed without (-) or with (+) 0.5% bile salts as in Figure 1B with data similarly presented. Quadruple asterisks (****) indicates values that are statistically different with P<0.0001 by the Student's T-test. **B.** Dose effect of bile salts on PAO1 twitching. PAO1 twitching was analyzed with the standard polystyrene Petri dish protocol as in (A) with varying concentrations (w/v %) of bile salts as indicated. Data presented in both panels are from three biological experiments each performed in triplicate. Data points from the same experiment are represented by the same symbols in color and shape.

Figure 3. Detergents, but not antibiotics, promote *P. aeruginosa* twitching. A. Effects of detergents. PAO1 twitching was analyzed with the standard Petri dish protocol as in Figure 2A with LB agar without modification (-) or with bile salts (BSs) (5 mg/ml), Triton X-100 (TX100) (75 μg/ml), Triton X-114 (TX114) (75 μg/ml), or sodium dodecyl sulfate (SDS) (850 μg/ml). **B.** Effects of antibiotics. PAO1 twitching was analyzed as in (A) with ampicillin (Amp) (313 ng/ml), ciprofloxacin (Cipro) (31 ng/ml), gentamicin (Gent) (31 ng/ml), or polymyxin B (PB) (313 ng/ml). Data presented in both panels are from three biological experiments each performed in triplicate. Data points from the same experiment are represented by the same symbols in color and shape. Triple (***) and quadruple (****) asterisks indicate two values are statistically different with P<0.001 and P<0.0001 by the Student's T-test, respectively. Antibiotics resulted in values that are not significantly (ns) different with P>0.05.

Figure 4. Glass surfaces increase *P. aeruginosa* twitching motility. A. Glass surfaces stimulate PAO1 twitching motility and eliminate the stimulatory effect of bile salts. The twitching motility of PAO1 and its isogenic *pilA* mutant were analyzed with polystyrene (PS) or glass microscope slides in LB without or with 0.5% bile salts (BSs) after 18 hours of incubation (see Materials and Methods). **B.** Hydrophobic coating of glass reduces twitching motility. Experiments were performed as in A, except that the microscope slides were coated without or with polydimethylsiloxane (PDMS) (see Materials and Methods). Data presented in both panels are from three biological experiments each performed in triplicate. Double asterisks (**) indicate two values are statistically different with P<0.01 by the Student's T-test.

Figure 5. TC-treated plates ameliorate twitching and abolish the effects of bile salts.
A. *P. aeruginosa* twitching on TC-treated polystyrene. The twitching motility of *P. aeruginosa* PAO1 and its isogenic *pilA* mutant strain was analyzed with 6-well polystyrene plates (see Materials and Methods) either TC-treated or non-treated in LB agar without or with 0.5% bile salts (BSs). **B.** *A. nosocomialis* twitching on TC-treated polystyrene. The twitching motility of *A. nosocomialis* M2 and its isogenic *pilA* deletion mutant was analyzed as in A. Data presented in both panels are from three biological experiments each performed in triplicate. Double (**) and quadruple (****) asterisks indicate that two values are statistically different with P<0.01 and P<0.0001 by the Student's T-test, respectively.