MANUSCRIPT TITLE

Genome-wide analysis of mRNA decay in Arabidopsis shoot and root reveals the importance of co-translational mRNA decay in the general mRNA turnover.

AUTHORS

Carpentier Marie-Christine[1,2], Receveur Anne-Elodie[1,2], Boubegtitene Alexandre[1,2], Cadoudal Adrien[1,2], Bousquet-Antonelli Cécile[1,2] and Merret Rémy[1,2]*

[1] CNRS-LGDP UMR 5096, 58 avenue Paul Alduy 66860 Perpignan, France

[2] Université de Perpignan Via Domitia, LGDP-UMR5096, 58 avenue Paul Alduy, 66860 Perpignan, France

- To whom correspondence should be addressed. Tel: +33(0)468662247; Fax: +33(0)468668499; Email: remy.merret@univ-perp.fr

ABSTRACT

Until recently, the general 5'-3' mRNA decay was placed in the cytosol after the mRNA was released from ribosomes. However, the discovery of an additional 5' to 3' pathway, the Co-Translational mRNA Decay (CTRD), changed this paradigm. Up to date, defining the real contribution of CTRD in the general mRNA turnover has been hardly possible as the enzyme involved in this pathway is also involved in cytosolic decay. Here we overcame this obstacle and created an Arabidopsis line specifically impaired for CTRD called XRN4ΔCTRD line. Through a genome-wide analysis of mRNA decay rate in shoot and root, we tested the importance of CTRD in mRNA turnover. First, we observed that mRNAs tend to be more stable in root than in shoot. Next, using XRN4ΔCTRD line, we demonstrated that CTRD is a major determinant in mRNA turnover. In shoot, the absence of CTRD leads to the stabilization of thousands of transcripts while in root its absence is highly compensated resulting in faster decay rates. We demonstrated that this faster decay rate is partially due to XRN4 cytosolic decay. Finally, we correlated this organ-specific effect with XRN4ΔCTRD line phenotypes revealing a crucial role of CTRD in mRNA homeostasis and proper organ development.

INTRODUCTION

The tight control and dynamic fine-tuning of gene expression are sine qua non conditions for life necessary to maintain proper cellular homeostasis and identity in response to developmental and environmental cues. The reprograming of gene expression is exerted at many steps: transcriptionally, post-transcriptionally at the level of messenger RNA (mRNA) and post-translationally at the protein level. All these processes are studied intensively in a global effort to understand basic mechanisms regulating gene expression in eukaryotes. Regulations at the mRNA level emerged as potent means to modify gene expression pattern to swiftly adapt the cellular activity and allow organisms to acclimate to and survive stress (1).

In the cytoplasm, over its entire lifetime, any mRNA is in balance between translation, storage and decay. The spatiotemporal regulation of this equilibrium is critical for transcriptome adjustment in response to developmental and environmental cues in plants (2). In *Arabidopsis thaliana*, dysfunction in mRNA decay due to the loss of function of key factors often results in post-embryonic lethality, severe growth defects or impairment of the stress response (3–5). As an example, the loss of function of XRN4, the major cytoplasmic 5′-3′ exoribonuclease, induces defects in leaf morphology, root growth and stress response such as dark response, nitrogen deprivation or heat stress (6–9). These pleiotropic phenotypes factors strongly support the importance of mRNA decay across development and response to stress in plants.

Degradation of mRNAs is an important component of gene expression that controls the steady state concentration of functional transcripts in the cell. This process is well conserved in eukaryotes and can initiate from the 5′ or 3′ extremities of mRNA (10). The so-called general 5′ to 3′ mRNA turnover that takes place in the cytoplasm occurs along three steps. Removal of the poly(A) tail of mRNA by deadenylases is the first and rate-limiting step in mRNA degradation. Following poly(A) tail shortening (deadenylation), the decapping complex (composed of VCS, DCP1 and DCP2 in *Arabidopsis thaliana*) hydrolyses the mRNA cap structure. Then, the 5′-phosphate end of the decapped mRNA is attacked by the cytoplasmic exoribonuclease (XRN4 in *Arabidopsis thaliana*, XRN1 in *Saccharomyces cerevisiae*) which digests the body of the transcript. The 3′ to 5′ mRNA can act through the activity of the RNA exosome complex or the suppressor of varicose, SOV in *Arabidopsis thaliana* (11, 12). The biological relevance of these pathways was addressed in *Arabidopsis thaliana*, revealing a large

contribution of 5′-3′ mRNA decay in the general mRNA turnover (13). A defect in one of these pathways can be compensated by a feedback mechanism to maintain proper mRNA homeostasis. However, the molecular mechanism governing this feedback is still unclear (10, 13). The redundancy of the 5′-3′ and 3′-5′ decay pathways was also demonstrated. Indeed, dysfunction of bidirectional RNA decay pathways results in an accumulation of spurious siRNAs and causes several developmental defects (14), demonstrating the importance of mRNA decay in plant development.

While initially thought to be mutually exclusive, data from several organisms including *Saccharomyces cerevisiae* and *Arabidopsis thaliana* now proved that mRNA decay and translation are actually tightly intertwined (15–17). In particular, the existence of an evolutionarily conserved turnover process, the 5′-3′ co-translational mRNA decay (CTRD) was proposed, where mRNAs are degraded from 5′ to 3′ while still engaged in translation. The existence of a CTRD was first proposed in yeast (18). The authors demonstrated using a set of reporter genes that uncapped mRNAs can be detected associated with ribosomes. Next the existence of this pathway in *Arabidopsis thaliana* was also proposed (7, 19). Heat stress induces a slowing down of ribosome elongation triggering the CTRD of transcripts coding for proteins with hydrophobic N-termini (19). This process was subsequently found to globally shape the whole transcriptome of many eukaryotes such as yeast, mammalian cells and plants (15–17, 20, 21). Recently, this pathway was proposed to be conserved in Angiosperms (21).

This transcriptome-wide effect of CTRD was revealed by sequencing of RNA decay intermediates using high-throughput degradome (or 5′P-Seq) approaches (20, 22–24). These approaches revealed that mRNA decay intermediates follow an XRN1/4-dependent, three-nucleotide periodicity. This periodicity can be explained by the fact that XRN1/4 chases the last translating ribosome in a codon-by-codon manner, and since it is a processive enzyme, only degradation intermediates protected by ribosomes can be captured. Consequently, each of these degradome approaches gives a snapshot of CTRD and also reveals ribosome dynamics and how degradation impacts this dynamic (15). Since the development of 5′P-Seq approaches an increasing number of articles reported the existence of CTRD, identified actors of this process and provided evidence that CTRD activity is variable across conditions (16, 20–22, 25–27). However, the actual respective contributions of the 5′-3′ cytosolic and 5′-3′ CTRD

decays to the general 5'-3' mRNA turnover have never been properly appreciated in any organisms.

Until now, it has been difficult to distinguish the relative contributions of 5'-3' cytosolic and co-translational decays in mRNA turnover since both processes involve the same enzyme : XRN1/XRN4. Here, we overcame this obstacle and created an Arabidopsis line specifically impaired for CTRD called XRN4ΔCTRD line. In this line, XRN4 can still decay mRNAs in the cytosol but can no longer perform CTRD. Using this line in comparison to WT and *xrn4* mutant, we performed a genome-wide mRNA decay analysis in both root and shoot. First, we demonstrated that mRNAs expressed in both organs tend to be more stable in root than in shoot. Next, we demonstrated that CTRD is the main 5'-3' mRNA decay pathway in shoot. Indeed, 70% of transcripts targeted by 5'-3' mRNA decay are targeted by CTRD with most of them exclusively targeted by this pathway. Finally, in root, we observed that an RNA decay feedback mechanism takes place specifically in the absence of CTRD, with a faster decay rate. We demonstrated that part of this feedback is mediated by XRN4 cytosolic decay. This mechanism induces specific root phenotypes in the XRN4ΔCTRD line opposite to those observed in the *xrn4* mutant.

MATERIAL AND METHODS

Sequence analysis

Sequence alignment was performed with Uniprot protein sequence of ScXRN1 (P22147), CeXRN1 (Q9BHK7), HsXRN1 (Q8IZH2), ScXRN2 (Q02792), AtXRN4 (Q9FQ04), HsXRN2 (Q9H0D6), CeXRN2 (Q9U299) and DmXRN1 (Q9VWI1).

Generation of AtXRN4ΔCTRD and AtXRN4ΔCTRD($R_{118}A/R_{119}A$) transgenic lines

The mEGFP coding sequence was inserted in frame after the triplet encoding for E263 within the XRN4 CDS sequence (Supplemental Figure 1). This sequence was then subcloned in a gateway vector containing XRN4 promoter sequence (1100 bp before XRN4 start codon). XRN4ΔCTRD was transformed in *xrn4.5* (SAIL_ 681_E01) background using the *Agrobacterium tumefaciens* based floral dip technique (28). Independent lines expressing XRN4ΔCTRD were

screened by western-blot using XRN4 specific antibody (7). Two independent lines expressing XRN4 at levels similar to WT were selected for further analysis. To generate AtXRN4ΔCTRD($R_{118}A/R_{119}A$) transgenic lines, the same procedure was followed, except that Arg118 and Arg119 were mutated in Alanine prior to cloning steps.

Growth conditions

Analyses were carried out with Columbia-0 line as Wild-type (WT), XRN4ΔCTRD line and *xrn4-5* mutant (SAIL_681_E01). Seeds were sown on a 245x245mm square plate in a single row. Plantlets were grown vertically on synthetic Murashige and Skoog medium (Duchefa) containing 1% Sucrose and 0.8% plant agar at 22°C under a 16-h-light/8-h-dark regime.

Transcriptional arrest time course.

Transcriptional inhibition was performed on 15-d-old seedlings as described previously (13). The time-course experiment was performed 2 hours after daybreak. Plantlets were transferred horizontally on incubation buffer (15 mM sucrose, 1 mM Pipes pH 6.25, 1 mM KCl, 1 mM sodium citrate, 1 mM cordycepin) in a single row to easily separate shoots and roots. Roots and shoots were collected separately at 0, 7.5, 15, 30, 60, 120, 240 and 480 min after transcription arrest. Roots and shoots were separated using a scissor at the basis of the hypocotyl and rapidly transferred in liquid nitrogen prior to RNA extraction. For RNAseq, four biological replicates per genotype were collected.

Total RNA extraction, RNA sequencing

Total RNA was isolated using Monarch Total RNA Miniprep Kit according to manufacturer's instructions. Total RNA was then subjected to ribosomal RNA depletion using QIAseq FastSelect rRNA Plant Kit according to manufacturer's instructions. RNA library preparation was performed using QIAseq Stranded RNA Library Kit. The samples were multiplexed and sequenced in PE 2x150. At least 20 million of clusters were obtained per sample. After filtering out reads corresponding to chloroplastic, mitochondrial, ribosomal, and small RNA sequences, reads were mapped against the TAIR10 genome using Hisat2 v2.03 and the gtf Araport11 annotation file with standard parameters. Read counts by gene were performed by htseq-count in RPM. Only transcripts with at least 1 RPM at T_0 in all genotypes and all replicates were kept. Differential expression analyses were performed using DESeq2.

mRNA decay rate analysis

mRNA decay rate analysis was performed according to (13). Data normalization, as well as the modelling of mRNA decay and genotype effect, were performed using the Bioconductor RNAdecay package (13, 29). Data normalization was performed using the mean fold increase of 50 stable genes in all genotypes (Supplemental Table 1). mRNA half-life ($t_{1/2}$) for each transcript in each genotype is presented on Supplemental Table 2.

Polysome profile analysis

Polysome profiles and western-blot analysis were performed as described previously (16). To quantify XRN4 association to ribosomes, fractions corresponding to monosome and polysomes were pooled. To quantify free XRN4, fractions corresponding to free mRNPs were pooled. Proteins were then precipitated from both fractions by adding 2 volumes of absolute ethanol. After 6 hours at 4°C and centrifugation, protein pellets were washed 5 times with ethanol 70%. Finally, pellets were resuspending in Laemmli 4X buffer. For western-blot analysis, the same percentage of each fraction (Input, free mRNPs and ribosome-bound fractions) were loaded. Quantification was performed using Vilber software on 4 biological replicates. All blots were prepared and immunoblotted in parallel and simultaneously exposed for chemiluminescence quantification.

Immunoblotting

XRN4 antibody (7) was used at 1/1000th dilution. UGPase antibody was purchased (Agrisera) and used at 1/5000th dilution. Antibodies against DCP5, PAT1, PAT1H1, PAT1H2 were produced in rabbits using the Eurogentec double X immunization program. The DCP5 protein was detected using the serum affinity-purified against peptide PNGHSFPNHNGYRGRG at 1/1000th dilution; the PAT1 protein was detected using the serum affinity-purified against peptide VEQRIPDRTKLYPEP at 1/1000th dilution; the PAT1H1 protein was detected using the serum affinity-purified against peptide PGNRSPQASPGNLHR at 1/1000th dilution; the PAT1H2 protein was detected using the serum affinity-purified against peptide VPPRVSNHGNPNDGL at 1/1000th dilution. Primary antibody was incubated overnight at 4°C under constant agitation. A horseradish peroxidase-coupled secondary antibody was used and revealed with the Immobilon-P kit from Millipore.

RT-ddPCR

A spike-in Luciferase RNA was first transcribed *in vitro* using HiScribe™ T7 Quick High Yield RNA Synthesis Kit (New England Biolabs) according to Manufacturer's instructions. The FLuc plasmid provided in the kit was used as template. Luciferase RNA was then purified using Monarch RNA cleanup kit (New England Biolabs) according to Manufacturer's instructions. For each sample, 500 ng of total RNA was spiked with 10 pg of Luciferase RNA and reverse-transcribed using SuperScript IV kit using random primers (Thermo Scientific). cDNAs were then diluted 50-fold prior to ddPCR analysis. ddPCR was performed as described previously (30). Luciferase quantification in each sample was used for normalization. Primers used for ddPCR are listed on Supplemental Table 3.

5'P sequencing

5'Pseq library was prepared as described previously (31). Raw reads were trimmed to 50 pb before mapping. Metagene analysis was performed using FIVEPSEQ software v1.0.0 (22). "meta_counts_START.txt" and "meta_counts_STOP.txt" files were used to analyse 5'P reads accumulation around start and stop positions. The translational termination stalling index (TSI) was defined as the ratio of the number of 5'P read ends at the ribosome boundary (16–17 nt upstream from stop codon) to the mean number of 5'P read ends within the flanking 100 nt. Transcripts with a TSI value higher than 3 in WT were used to assess CTRD activity in *xrn4* and XRN4ΔCTRD line.

GO terms analysis

GO terms analysis was performed using clusterProfiler R package.

Phenotypic analysis

For leaf phenotypes, plants were cultivated in soil in a growth chamber under long day conditions (16h light at 21°C/8h dark at 16°C) at 65% relative humidity. Number of leaves and leaf fresh weight were calculated on 34 days-old plants. For root phenotypes, plants were cultivated *in vitro* on synthetic MS containing 1% Sucrose and 0.8% plant agar at 22°C under a 16-h-light/8-h-dark regime. For NaCl treatment, 4-d-old seedlings were transferred to MS

media supplemented with 125 mM NaCl. Root length was then determined on 15-d-old seedlings.

RESULTS

Generation of AtXRN4ΔCTRD transgenic line.

As XRN4 catalyses both cytosolic and co-translational mRNA decay, it is challenging to uncouple these pathways and determine their respective contributions. To tackle this question, we took advantage of a recent structural analysis of the yeast ScXRN1 physical interactions with the 80S ribosome (32). ScXRN1 carries several ribosome contact areas, including three loops (L1, L2 and L3), with the flexible loop L3 located in between the CR1 and CR2 conserved structured regions that form the catalytic domain, that is essential for ribosome binding (Figure 1A) (32). Tessina et al., showed that disruption of loop L3, through insertion of a monomeric GFP (mEGFP) abolishes ScXRN1 association with polysomes without affecting its exoribonucleolytic activity. We reasoned that such mutant is likely to not be able to degrade mRNA co-translationally anymore while retaining a cytosolic mRNA decay property. As we are interested in understanding the molecular mechanisms of 5'-3' mRNA degradation in *Arabidopsis thaliana,* we set to construct a line solely expressing an *XRN4* allele unable to bind polysomes. As we and others previously showed (7), the first half of XRNs, encompassing the CR1-CR2 catalytic core is highly conserved at the primary sequence level, and an AlphaFold-based 3D structure prediction supports the existence of loop L3, at the end of CR1 in AtXRN4 as in ScXRN1 (Figure 1B). In yeast, the mEGFP was inserted into loop L3, after serine 235, nine amino acids after the conserved CR1 region (Figure 1A), we hence chose to disrupt L3 in AtXRN4, also by inserting an mEGFP nine amino acids into loop 3, hence after glutamic acid 263 (AtXRN4 numbering) (Figure 1A). We next generated a stable transgenic line expressing this modified XRN4 under the control of its own promoter in an *xrn4* loss-of-function (lof) mutant (*xrn4-5*, SAIL_681_E01). We called this line XRN4ΔCTRD and identified two independent transgenic plants expressing XRN4ΔCTRD at levels similar to that of the endogenous protein in both shoot and root (Figure 1C). Several approaches were then deployed to demonstrate that this line is impaired in co-translational mRNA decay (Figure 2 and Supplemental Figures 2-3). First, we quantified XRN4 association to ribosomes by polysome fractionation followed by western-blotting using XRN4 specific antibodies. Fractions

corresponding to monosomes and polysomes were pooled and referred to as "Ribosome-bound fractions". Fractions corresponding to free mRNPs were also pooled and used to quantify free XRN4. XRN4 signal was compared in both fractions relative to an input fraction prior to polysome fractionation (Figure 2A-B). Interestingly, in both shoot and root, XRN4 signal is significantly higher in the ribosome-bound fractions than in the free mRNPs fractions suggesting that CTRD activity is important in both organs. We performed a similar analysis with the XRN4ΔCTRD line and found that XRN4 signal in ribosome-bound fractions drastically decreased (Figure 2A-B). In contrast, a GFP insertion at the C-terminus of XRN4 does not affect XRN4 distribution (Figure 2A-B).

Then, to test CTRD activity in XRN4ΔCTRD lines, we ran 5'Pseq assays and compared results to that of wild-type (WT) and *xrn4* lof mutant through monitoring of read accumulation around stop and start codons (Figure 2C and Supplemental Figure 2). As expected, a meta-transcriptome analysis shows a strong overaccumulation of reads at position 16/17nt before the stop codon in wild-type, a hallmark of active CTRD (16). In the XRN4ΔCTRD line, the minus 16/17nt peak shows a dramatic decrease that is almost identical to that of the *xrn4-5* lof mutant, supporting a drastic inactivation of CTRD (Figure 2C). Conversely, before the start codon, a clear over-accumulation (as compared to WT) of 5'P reads is observed in both *xrn4-5* and XRN4ΔCTRD lines (Supplemental Figure 2), another hallmark of 5'-3' mRNA decay impairment. We suppose that the higher accumulation observed in *xrn4-5* compared to XRN4ΔCTRD line is the consequence of the loss of both the cytosolic and co-translational mRNA decays. Recently, it was proposed that the terminational stalling index (TSI) can be used to identify CTRD activity at individual transcript levels (21, 33). Using this TSI metric, we demonstrated that CTRD activity is reduced in the XRN4ΔCTRD line to levels similar to that of a complete loss of XRN4 (Figure 2D-E). Finally, we observed through confocal microscopy that the XRN4ΔCTRD shows a subcellular distribution similar to that of a C-terminally tagged XRN4-GFP fusion and is still present in the cytosol (Supplemental Figure 3). All together, these data support that the insertion of a monomeric GFP into AtXRN4 loop 3, does not affect the total levels of the protein nor does it impair its cytoplasmic accumulation. But that it blocks its ability to associate to polysomes, and inactivates the CTRD pathway as efficiently as total loss of the XRN4 protein. We hence conclude that the XRN4ΔCTRD lines only retains the ability to

degrade mRNAs outside of polysomes, providing a useful tool to evaluate the contribution of CTRD to the 5'-3' turnover of transcripts.

Strategy for mRNA half-life determination and CTRD contribution

To determine the contribution of CTRD in mRNA turnover, we took advantage of the transgenic line XRN4ΔCTRD. A genome-wide mRNA decay analysis was carried on WT (Col0, active cytosolic and CTRD degradations), *xrn4* lof mutant (*xrn4-5*, SAIL_681_E01, inactive cytosolic and CTRD degradations) and two independent XRN4ΔCTRD lines (active cytosolic and inactive CTRD degradations) 15-d-old seedlings as we have shown that CTRD is highly active at this stage (16). As little is known about the importance of CTRD at the organ level, the analysis was performed on root and shoot separately (Supplemental Figure 4A). After transcriptional inhibition by addition of cordycepin, roots and shoots were collected separately at 0, 7.5, 15, 30, 60, 120, 240 and 480 min. In order to capture both deadenylated and adenylated mRNA decay intermediates, rRNA depletion was performed prior to RNA library preparation. RNA libraries were then multiplexed and sequenced in Paired-Ends 2X150bp. RNAseq was performed on four biological samples, revealing reproducible differences between genotypes and time-points (Supplemental Figure 5). Only transcripts with at least 1 RPM at T_0 in all genotypes and all replicates were kept. Prior to mRNA decay analysis, a DESeq analysis was performed between the two XRN4ΔCTRD lines (Supplemental Figure 6). Among the 16 comparisons, only 782 unique transcripts were detected as significantly up- or down-regulated demonstrating the reproducibility of both lines. These transcripts were removed from the analysis. Finally, to simplify subsequent analyses, the mean of RPM values of both XRN4ΔCTRD lines was performed resulting in one XRN4ΔCTRD line. The decay analysis was thus initiated on 12 306 transcripts (Supplemental Figure 4B). To determine the mRNA half-life for each transcript, we took advantage of the development of an RNA decay pipeline based on a mathematical modeling approach (13, 29). After data normalization and mathematical modelling, an mRNA half-life ($t_{1/2}$) was obtained for the 12 306 transcripts in WT, *xrn4* and XRN4ΔCTRD lines in both root and shoot (Supplemental Table 2).

mRNA decay landscape between shoot and root in WT

The study was first initiated in WT between root and shoot. The analysis of mRNA decay rates in both organs reveals a wide range of decay from very fast (few minutes) to very slow (several

hours). Interestingly, mRNAs expressed in both organs tend to be more stable in root than in shoot (171 minutes versus 103 minutes) (Figure 3A-B, Supplemental Table 2). The mRNA half-life distribution shows that root presents less unstable ($t_{1/2}$ < 100 minutes) and more stable mRNAs ($t_{1/2}$ > 100 minutes) compared to shoot. We next tested if mRNA stability in each organ could be correlated with corresponding biological functions. For each transcript, the fold change between its half-life in shoot versus root (FC $t_{1/2 \text{ shoot}}$ / $t_{1/2 \text{ root}}$) was determined followed by a GO slim analysis (Figure 2C). Interestingly, some biological functions present a higher stability in shoot than in root and vice-versa. mRNAs stability in *Arabidopsis thaliana* was correlated with some *cis*-elements such as intron numbers and 5'UTRs A/G content (13). We analyzed the distribution of these two features and also observed a strong correlation between intron number, 5'UTR content and mRNA stability at the organ level (Supplemental Figure 7). Intron number, 5'UTR increased A content and 5'UTR decreased G content were positively correlated with mRNA stability both in root and in shoot. As median mRNA half-life is much higher in root than in shoot, we then tested if this difference could be due to different levels of key components of the 5'-3' and 3'-5' mRNA decay machineries. Interestingly, the *SKI2* mRNA (an RNA helicase, subunit of the SKI complex involved in the 3'-5' decay process) is much more stable in shoot than in root (787 minutes in shoot and 218 minutes in root), what was confirmed by an independent approach (Supplemental Figure 8), suggesting that the 3'-5' mRNA decay could be less active in root than in shoot. Regarding the 5'-3' mRNA decay components, we analyzed published quantitively proteomic data on shoot and root and found that PAT1H1 and PAT1H2 (cofactors of decapping) are less accumulated in root compared to shoot (34). Using specific antibodies against PAT1, PAT1H1 and PATH2, we tested protein accumulation in our conditions (Supplemental Figure 9). While XRN4 and DCP5 seemingly accumulate at similar levels in root and shoot, we found that PAT1, PAT1H1 and PAT1H2 proteins under accumulate in root compared to shoot. These data could suggest that co-factors of decapping are less abundant in root resulting in a lower 5'-3' decay activity.

Contribution of CTRD in the general 5'-3' mRNA turnover

The contribution of XRN4 in CTRD and in the general 5'-3' mRNA turnover is presented in Figure 4. In shoot, the medians half-life in *xrn4* and XRN4ΔCTRD lines are similar (146 and 147 minutes respectively) and significantly higher than WT (Figure 4A, Supplemental Table 2) indicating that CTRD contributes to decay of many mRNAs and supporting the idea of an

important role in 5'-3' mRNA decay in this tissue. In root, the median half-life is also significantly higher in *xrn4* (218 minutes compared to 171 minutes in WT) but surprisingly, the median half-life in XRN4ΔCTRD line is significantly lower (112 minutes versus 171 minutes in WT). Majority of decay rates are faster in XRN4ΔCTRD (Figure 4B, Supplemental Table 2). To exclude a transgenic line effect, we repeated mRNA half-life determination on the two independent XRN4ΔCTRD lines. The same phenomenon is observed with the two lines, supporting a biological effect due to the absence of CTRD in root (Supplemental Figure 10).

To determine the contribution of the two decay pathways for each mRNA in root and shoot, we used the likelihood function presents in the RNAdecay package to assign a statistical genotypic effect to each transcript (13, 29). Modelling considered 5 combinations (called α groups) of genotypic effects (Supplemental Figure 11). An mRNA might show a different half-life in each genotype (α group 1) or show a similar half-life in two genotypes (α groups 2, 3 and 4), or show the same half-life in all genotypes (α group 5). In α groups 1 to 4, different subgroups exist based on the effects of *xrn4* and XRN4ΔCTRD lines on mRNA half-life (Supplemental Figure 11). Interestingly, the distribution of mRNAs in α groups are different between root and shoot suggesting distinct decay mechanisms in each organ. The α group 5.1 is the most abundant in shoot and root with 3 332 and 2 679 transcripts respectively, representing 27% (3 332/12 306) and 22% (2 679/12 306) of the population. These transcripts are not affected in *xrn4* and XRN4ΔCTRD lines (Supplemental Figure 11).

According to *xrn4* and XRN4ΔCTRD lines genotypic effects, we categorized subgroups in two main groups, 'XRN4 targets' and 'RNA decay feedback targets'. We considered that the XRN4 targets group is composed by transcripts with slower decay (*e.g.* longer $t_{1/2}$) in the *xrn4-5* and XRN4ΔCTRD lines than in WT (subgroups 1.4, 1.6, and 2.2) or with slower decay in *xrn4-5* and not affected in the XRN4ΔCTRD line (subgroup 4.2) (Supplemental Figure 11, Supplemental Table 2). The RNA decay feedback group encompasses transcripts with faster decay in *xrn4-5* or XRN4ΔCTRD lines compared to WT (subgroups 1.1, 1.2, 1.3, 1.5, 2.1, 3.1 and 4.1). Finally, since the biological relevance of the subgroup 3.2 ($t_{1/2}$ XRN4ΔCTRD > $t_{1/2}$ *xrn4*-5 = $t_{1/2}$ WT) was difficult to interpret, its transcripts were removed as well as those from subgroup 5.1 (not affected in the *xrn4-5* nor in XRN4ΔCTRD lines). We hereby retained 7 882 and 9 291 transcripts from shoot and root respectively, for further analyses. The distribution of the 'XRN4' and 'RNA decay feedback' target groups is presented in Figure 5A-B. Strikingly, the proportions of the

two groups are totally different between organs, with 71% (5 634) of XRN4 targets and 29% (2 248) of RNA decay feedback in shoot compared to 39% (3 648) and 61% (5 643) in root, here again suggesting distinct decay regulations between shoot and root.

To identify transcripts exclusively targeted by cytosolic or CTRD pathways, we reasoned as follows: 1- A transcript only targeted by CTRD pathway will have a similar increased half-life in both XRN4ΔCTRD and *xrn4* lines (subgroup 2.2). As increased half-life in XRN4ΔCTRD line is slightly higher than *xrn4* in subgroup 1.4, these transcripts were also considered as CTRD specific targets, 2- A transcript only targeted by cytosolic pathway will have an increased half-life in *xrn4* and an half-life not affected in XRN4ΔCTRD line (subgroup 4.2), 3- A transcript targeted by both pathways will have an increased half-life in both XRN4ΔCTRD and *xrn4* lines but at lower amplitude in XRN4ΔCTRD line (subgroup 1.6).

In shoot, 32% of XRN4 targets (1 794 out of 5 634 XRN4 targets, subgroup 4.2) are only affected in *xrn4* and not in XRN4ΔCTRD line, suggesting that these transcripts are targeted only by XRN4 cytosolic decay. *In contrario*, 50% are exclusively targeted by the CTRD pathway (2 815, subgroups 2.2 and 1.4) while 18% (1 025) are targeted by both pathways (subgroup 1.6). These data are consistent with $t_{1/2}$ distribution observed in *xrn4* and XRN4ΔCTRD lines in shoot (Figure 4A) and suggest that CTRD is the major 5'-3' mRNA decay pathway in shoot.

In root the situation is more complex. XRN4 targets represent only 39% of the population (3 648/9 291) with most of them targeted by cytosolic decay (2 282/3 648, subgroup 4.2). On the other hand, the majority of transcripts (61%, 5 643/9 291) presents a faster decay in *xrn4* and/or XRN4ΔCTRD line (Figure 5B). This phenomenon was already observed in *Arabidopsis thaliana sov* mutant (13) and suggests that an RNA decay feedback mechanism takes place in the absence of cytosolic and/or CTRD decays. The most important effect is in subgroup 3.1 (2 381). Indeed, transcripts of this group are decayed faster only in XRN4ΔCTRD line. This subgroup was only represented by 469 transcripts in shoot. Interestingly, subgroups in which transcripts decay is faster in XRN4ΔCTRD line (1.1, 1.2, 1.5, 2.1, 3.1) are the only subgroups that increase in number between shoot and root suggesting that a specific feedback mechanism takes place in absence of CTRD in root.

Next, we compared CTRD- and cytosolic-specific targets between organs and observed only a very limited overlap, suggesting that identical mRNAs are turned over by distinct means in different organs (Figure 5C-D).

In order to determine biological processes targeted by 5'-3' mRNA decay in root and shoot, a Gene Ontology (GO) analysis was performed on XRN4 CTRD specific targets (subgroups 1.4 and 2.2), cytosolic specific targets (subgroup 4.2) and RNA decay feedback targets (subgroups 1.1, 1.2, 1.3, 1.5, 2.1, 3.1 and 4.1) (Figure 6). In both organs, XRN4 CTRD and cytosolic specific targets are part of very distinct biological processes. In shoot, mRNAs coding for various RNA processes such as ribosome biogenesis or ribonucleoprotein complex biogenesis are targeted by CTRD while mRNAs coding for vesicle localization, phosphate starvation or alpha-amino acid metabolic process are targeted by cytosolic decay (Figure 6).

Part of the feedback mechanism is mediated by XRN4 cytosolic decay pathway

Many studies report the existence of a buffering system in which defects in turnover are compensated by a transcriptional readjustment, to maintain even steady-state mRNA levels (13, 35–37). To assess whether the faster decay rates observed in XRN4ΔCTRD line in root induce transcriptional feedback, we compared RNA abundances at T_0 in XRN4ΔCTRD line and WT (Supplemental Figure 12). Using the whole dataset, no clear differences are observed between XRN4ΔCTRD line and WT. By contrast, CTRD specific targets present higher abundance in XRN4ΔCTRD line while RNA decay feedback targets present lower abundance suggesting that the loss of CTRD does not lead to transcriptional feedback (Supplemental Figure 12). Faster decay rates observed in the root of the XRN4ΔCTRD line suggest that an alternative decay mechanism takes place in the absence of CTRD. To test the possibility of a feedback mechanism mediated by the XRN4 cytosolic pathway, we created an additional XRN4ΔCTRD line but impaired in its 5'-3' exonucleolytic activity. To do so, we performed a protein sequence alignment between AtXRN4 and diverse XRN orthologues (Supplemental Figure 13A). In *Drosophila melanogaster*, it was reported that the substitution of arginines R100 and R101 into alanines abolishes DmXRN1 activity. These two arginines are conserved and present in AtXRN4 at positions 118 and 119. Thus, we created the *AtXRN4ΔCTRD(R$_{118}$A/R$_{119}$A)* allele, that we expressed under the control of its own promoter in the *xrn4-5* background. We selected a transformant line expressing transgene levels similar to that of endogenous XRN4

and transgenic XRN4ΔCTRD (Supplemental Figure 13B). We next measured half-lives of candidate transcripts, through ddPCR-monitoring of RNA levels following cordycepin treatment (30), in wild-type, *xrn4* lof, XRN4ΔCTRD and XRN4ΔCTD($R_{118}A/R_{119}A$) backgrounds (Figure 7 and Supplemental Figure 14). First, to confirm that the XRN4ΔCTRD allele retains normal and similar catalytic activities both in root and shoot, we selected three transcripts amongst the list of those identified as only targeted by the XRN4 cytosolic decay in both organs (subgroup 4.2, Figure 5A-B). For all three of them, mRNA half-life measurement revealed similar half-lives in WT and XRN4ΔCTRD lines in both shoot and root while, their half-lives significantly increased at the same amplitude in *xrn4-5* and XRN4ΔCTRD($R_{118}A/R_{119}A$) lines (Figure 7A, Supplemental Figure 14A). This first analysis confirms that insertion of an mEGFP into AtXRN4 loop L3, does not affect its catalytic activity, as is the case for ScXRN1. It also comes as a validation of the transcriptome-wide assays that identified the mRNAs from subgroup 4.2 are cytosolic XRN4 targets. Next, we selected transcripts from subgroup 1.5 to validate and analyse the feedback mechanism observed in root (Figure 7B, Supplemental Figure 14B). Interestingly for transcripts of subgroup 1.5, the feedback mechanism observed in XRN4ΔCTRD line is totally abolished in XRN4ΔCTRD($R_{118}A/R_{119}A$) line suggesting that XRN4 cytosolic decay can compensate partially the absence of CTRD in root (Figure 5B).

XRN4ΔCTRD line presents contrasting phenotypes in shoot and root

As mRNAs half-life distribution in *xrn4* and XRN4ΔCTRD lines reveals similar profiles in shoot and opposite profiles in root (Figure 4), we tested if this discrepancy could be associated with different phenotypes at the organ level. Several phenotypes were already observed in *xrn4*, such as defects in root growth under various conditions (6, 8) or leaf development (9). We analyzed these different phenotypes in XRN4ΔCTRD line in comparison to WT, *xrn4* and a complemented XRN4-GFP line (Figure 8). At leaf level, XRN4ΔCTRD line presents similar phenotypes than *xrn4* such as serrated leaves, higher leaf fresh weight and greater number of leaves (Figure 8A-C). However, at the root level, XRN4ΔCTRD line presents opposite phenotypes than *xrn4* (Figure 8D-E). Primary root is longer in XRN4ΔCTRD line compared to WT while *xrn4* presents a shorter primary root (Figure 8D). Interestingly, this phenomenon is still observed under stress conditions such as salt stress (Figure 8E). To correlate these phenotypes to transcripts stability, we focused on transcripts associated with GO terms "root morphology" and "root development". Interestingly, among them, 111 transcripts present

distinct mRNA half-life between *xrn4* and XRN4ΔCTRD line : 41 transcripts have opposite half-life between *xrn4* and XRN4ΔCTRD line (subgroup 1.5 or subgroup 1.3) and 70 present faster decay in XRN4ΔCTRD line and are not affected in *xrn4* (Supplemental Table 4). These data could suggest that the misregulation of mRNA stability for transcripts involved in root morphogenesis/development could affect root growth. All together, these phenotypic analyses reveal a strong correlation between decay rates alteration and organ phenotype.

DISCUSSION

mRNA decay plays an important role in the post-transcriptional regulation of gene expression. In eukaryotes, two major pathways can together turn over the transcriptome, the 5'-3' mRNA decay and the 3'-5' mRNA decay (10). In plants, defects in 5'-3' mRNA decay can induce drastic developmental and stress response phenotypes revealing the importance of this pathway (3, 4, 6, 7, 9, 38). The general 5'-3' mRNA decay was placed in the cytosol after ribosomes release. However, for many years, it has also been proposed that mRNA decay can occur on translated mRNAs through a 5'-3' co-translational mRNA decay mechanism (15, 16, 20). While players, enzymes and cofactors of the 5'-3' turnover are well recognized, the importance either cellular or physiological and the relative contribution of each pathway has never been assessed in any organism. As a matter of fact, uncoupling the 5'-3' decay pathways proved difficult because they are catalysed by the same enzymes, the main one being XRN4 in *Arabidopsis thaliana*. Here, we developed a transgenic line impaired in CTRD that allows the discrimination between 5'-3' cytosolic and 5'-3' CTRD pathways. Using a genome-wide mRNA decay analysis, we determined the importance of CTRD in both shoot and root and provide evidence that it plays a crucial role in root development.

mRNA decay analysis in WT in shoot and root reveals diverse range of decay rates (from minutes to hours) with a respective median value of 103 and 171 minutes. This range is quite similar to previous mRNA decay analysis made in *Arabidopsis thaliana* (13, 39) but reveals an organ-specific pattern. This organ-specific mRNA stability was already observed in different organisms. As an example, in drosophila, stabilization of transcripts involved in cell-fate decisions and axonogenesis is an important process for neural development (40). In human, changes in mRNA stability contribute to fibroblast quiescence maintenance (41). By analyzing

cis- and *trans-* elements, we found that SKI2 transcript is less stable in root compared to shoot. We also found that PAT mRNA decapping factors (PAT1, PAT1H1, PAT1H2) are less accumulated in root compared to shoot. The SKI complex (composed of SKI2, SKI3 and SKI8) is involved in the unfolding of mRNAs to assist exosome-mediated decay (42). The SKI complex is important at the leaf level as together with uridylation, they regulate RNA degradation pathway important for photosynthesis (43). This is consistent with the extreme stability of SKI2 mRNA in our shoot half-life dataset. In *Arabidopsis thaliana*, PAT1 has been implicated in pathogen response where it activates decay of specific transcripts (44). More recently, it was proposed that PAT1H1, PAT1H2 harbor specific and overlapping functions with PAT1 during plant growth (45). However, their importance at the organ level was never tested. More investigations will be needed to better assess the importance of decay factors in mRNA stability at the organ level.

The importance of the 5′-3′ mRNA decay was revealed by the distribution of mRNA half-lives in *xrn4*. In *xrn4*, the median half-life was 43 and 77 minutes higher than WT in shoot and root respectively indicating that XRN4 contributes to decay of many mRNAs. These data are in agreement with the contribution of the decapping (through VCS) in the decay of the transcriptome where 68% of transcripts present a VCS-dependent decay (13). The importance of CTRD was revealed by the distribution of mRNA half-lives in the XRN4ΔCTRD line. In shoot, the contribution is clear with similar half-life distributions in *xrn4* and XRN4ΔCTRD lines (Figure 4A). The α group pattern in shoot reveals that most of the 5′-3′ degradation is mediated by CTRD (Figure 5). Interestingly, few transcripts are targeted by CTRD and cytosolic decay (α group 1.6) meaning that transcripts are mostly targeted by one of the two pathways.

The specific inactivation of the XRN4-mediated CTRD in root results for a large fraction of the root and to a lesser extent of the shoot transcriptomes, in a counter intuitive phenomenon where thousands of transcripts decay faster (Figure 4 and Figure 5A-B). Such phenomenon was previously reported in yeast and *Arabidopsis thaliana* and recognized as a buffering process, on RNA decay defects (36, 37, 46). In yeast, a defect in decay rate induces the reduction of the transcription rates to maintain mRNA steady-state at proper levels (37, 46). This mechanism was called RNA buffering and is mediated by the nucleocytoplasmic shuttling of ScXRN1 (37). In fact, two NLS sequences were recently identified in ScXRN1 that are important for its import to the nucleus and for transcription rate modulation (35). In plants, no

direct ortholog of XRN1 exists. *Arabidopsis thaliana* indeed carries three XRNs, AtXRN2, AtXRN3 and AtXRN4 that are structurally similar to ScXRN2/RAT1 (47). The deletion of a bipartite NLS during duplication may have given rise to an AtXRN4 cytoplasmic exoribonuclease that have cytoplasmic functions similar to that of ScXRN1 (47). Thus, AtXRN4 is probably not able to perform nucleocytoplasmic shuttling explaining the absence of transcriptional feedback in the XRN4ΔCTRD line (Supplemental Figure 12).

Faster decay rates in root of XRN4ΔCTRD line suggest that another decay mechanisms take place in absence of CTRD. Using XRN4ΔCTRD(R_{118}A/R_{119}A) line, we demonstrated that part of this feedback is mediated by XRN4 cytosolic decay (subgroup 1.5, Figure 7C). However, transcripts from other subgroups present also RNA decay feedback. We and others proposed that XRN4 is not the only exoribonuclease involved in CTRD (16, 48). In fact, in *xrn4* mutant, the 3-nt periodicity and 5'P reads accumulation -16/-17 nt before stop codons are not totally abolished (16). Interestingly, in *fry1* mutant, 5'-3' mRNA decay seems more repressed than in *xrn4* (48). In *fry1* mutant, a constitutive accumulation of 3'-phosphoadenpsione 5'-phosphate (PAP) is observed resulting in inhibition of XRN activity (49). These data suggest that in addition to XRN4, another exoribonuclease sensitive to PAP, could be involved in CTRD. Whether the feedback mechanism is mediated by another co-translational decay pathway is still an open question.

Why and how at the molecular level, the compensation phenomenon to the loss of CTRD is significantly more important in root than in shoot is another interesting question that emerges from our work that will need further studies to answer. However, consistent with the differences in compensation and the distinct contributions of the CTRD to overall 5'-3' mRNA turnover, we report that the CTRD takes on distinct physiological importance in root and shoot (Figure 8). Indeed, the phenotypes observed in the XRN4ΔCTRD line correlate with alteration of decay rates. In shoot, *xrn4* and XRN4ΔCTRD lines present similar alterations of decay rate resulting in similar phenotypes while in root, opposite alterations of decay rate are observed resulting in opposite phenotypes. Taken together, our data report that CTRD plays important roles in the fine-tuning of mRNA turnover and is essential for proper plant development. The development of this XRN4ΔCTRD line will give to the community the opportunity to analyze deeper the importance of CTRD in plant development and stress response.

DATA AVAILABILITY

The RNAseq and 5'Pseq datasets generated in this study have been deposited in the Short Read Archive with the accession code PRJNA1018408 and PRJNA1023397 respectively.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR online.

ACKNOWLEDGEMENTS

This study is set within the framework of the "Laboratoires d'Excellences (LABEX)" TULIP (ANR-10-LABX-41) and of the "École Universitaire de Recherche (EUR)" TULIP-GS (ANR-18-EURE-0019). The authors gratefully acknowledge Jean-Marc Deragon for critical reading of the manuscript.

FUNDING

Agence Nationale de la Recherche [ANR-21-CE20-0003 to R.M.]

Agence Nationale de la Recherche [ANR-17-CE20-0007 to C.B.A.]

CONFLICT OF INTEREST

None declared

REFERENCES

1. Matsui,A., Nakaminami,K. and Seki,M. (2019) Biological Function of Changes in RNA Metabolism in Plant Adaptation to Abiotic Stress. *Plant Cell Physiol*, **60**, 1897–1905.

2. Chantarachot,T. and Bailey-Serres,J. (2018) Polysomes, stress granules, and processing bodies: A dynamic triumvirate controlling cytoplasmic mRNA fate and function. *Plant Physiol*, **176**, 254–269.

3. Xu,J. and Chua,N.-H. (2009) Arabidopsis decapping 5 is required for mRNA decapping, P-body formation, and translational repression during postembryonic development. *Plant Cell*, **21**, 3270–9.

4. Xu,J., Yang,J.-Y., Niu,Q.-W. and Chua,N.-H. (2006) Arabidopsis DCP2, DCP1, and VARICOSE form a decapping complex required for postembryonic development. *Plant Cell*, **18**, 3386–98.

5. Zhang,W., Murphy,C. and Sieburth,L.E. (2010) Conserved RNaseII domain protein functions in cytoplasmic mRNA decay and suppresses Arabidopsis decapping mutant phenotypes. *Proc Natl Acad Sci U S A*, **107**, 15981–15985.

6. Kawa,D., Meyer,A.J., Dekker,H.L., Abd-El-Haliem,A.M., Gevaert,K., Van De Slijke,E., Maszkowska,J., Bucholc,M., Dobrowolska,G., De Jaeger,G., *et al.* (2020) SnRK2 Protein Kinases and mRNA Decapping Machinery Control Root Development and Response to Salt. *Plant Physiol*, **182**, 361–377.

7. Merret,R., Descombin,J., Juan,Y. ting, Favory,J.J., Carpentier,M.C., Chaparro,C., Charng,Y. yung, Deragon,J.M. and Bousquet-Antonelli,C. (2013) XRN4 and LARP1 are required for a heat-triggered mRNA decay pathway involved in plant acclimation and survival during thermal stress. *Cell Rep*, **5**, 1279–1293.

8. Nagarajan,V.K., Kukulich,P.M., von Hagel,B. and Green,P.J. (2019) RNA degradomes reveal substrates and importance for dark and nitrogen stress responses of Arabidopsis XRN4. *Nucleic Acids Res*, **47**, 9216–9230.

9. Windels,D. and Bucher,E. (2018) The 5'-3' exoribonuclease XRN4 regulates auxin response via the degradation of auxin receptor transcripts. *Genes (Basel)*, **9**, 1–12.

10. Sieburth,L.E. and Vincent,J.N. (2018) Beyond transcription factors: Roles of mrna decay in regulating gene expression in plants [version 1; referees: 3 approved]. *F1000Res*, **7**, 1–10.

11. Januszyk,K. and Lima,C.D. (2014) The eukaryotic RNA exosome. *Curr Opin Struct Biol*, **24**, 132–140.

12. Goeres,D.C., Van Norman,J.M., Zhang,W., Fauver,N.A., Spencer,M. Lou and Sieburth,L.E. (2007) Components of the Arabidopsis mRNA decapping complex are required for early seedling development. *Plant Cell*, **19**, 1549–1564.

13. Sorenson,R.S., Deshotel,M.J., Johnson,K., Adler,F.R. and Sieburth,L.E. (2018) Arabidopsis mRNA decay landscape arises from specialized RNA decay substrates, decapping-mediated feedback, and redundancy . *Proceedings of the National Academy of Sciences*, **115**, E1485–E1494.

14. Zhang,X., Zhu,Y., Liu,X., Hong,X., Xu,Y., Zhu,P., Shen,Y., Wu,H., Ji,Y., Wen,X., *et al.* (2015) Suppression of endogenous gene silencing by bidirectional cytoplasmic RNA decay in Arabidopsis. *Science (1979)*, **348**, 120–123.

15. Pelechano,V., Wei,W. and Steinmetz,L.M. (2015) Widespread co-translational RNA decay reveals ribosome dynamics. *Cell*, **161**, 1400–1412.

16. Carpentier,M.C., Deragon,J.M., Jean,V., Seng Hour Vichet,B., Bousquet-Antonelli,C. and Merret,R. (2020) Monitoring of XRN4 targets reveals the importance of cotranslational decay during arabidopsis development. *Plant Physiol*, **184**, 1251–1262.

17. Tuck,A.C., Rankova,A., Arpat,A.B., Liechti,L.A., Hess,D., Iesmantavicius,V., Castelo-Szekely,V., Gatfield,D. and Bühler,M. (2020) Mammalian RNA Decay Pathways Are Highly Specialized and Widely Linked to Translation. *Mol Cell*, **77**, 1222-1236.e13.

18. Hu,W., Sweet,T.J., Chamnongpol,S., Baker,K.E. and Coller,J. (2009) Co-translational mRNA decay in Saccharomyces cerevisiae. *Nature*, **461**, 225–229.

19. Merret,R., Nagarajan,V.K., Carpentier,M.C., Park,S., Favory,J.J., Descombin,J., Picart,C., Charng,Y.Y., Green,P.J., Deragon,J.M., *et al.* (2015) Heat-induced ribosome pausing triggers mRNA co-translational decay in Arabidopsis thaliana. *Nucleic Acids Res*, **43**, 4121–4132.

20. Yu,X., Willmann,M.R., Anderson,S.J. and Gregory,B.D. (2016) Genome-Wide Mapping of Uncapped and Cleaved Transcripts Reveals a Role for the Nuclear mRNA Cap-Binding Complex in Cotranslational RNA Decay in Arabidopsis. *Plant Cell*, **28**, 2385–2397.

21. Guo,R., Yu,X. and Gregory,B.D. (2023) The identification of conserved sequence features of co-translationally decayed mRNAs and upstream open reading frames in angiosperm transcriptomes. *Plant Direct*, **7**, e479.

22. Nersisyan,L., Ropat,M. and Pelechano,V. (2020) Improved computational analysis of ribosome dynamics from 5'P degradome data using fivepseq. *NAR Genom Bioinform*, **2**, 1–12.

23. German,M.A., Pillay,M., Jeong,D.H., Hetawal,A., Luo,S., Janardhanan,P., Kannan,V., Rymarquis,L.A., Nobuta,K., German,R., *et al.* (2008) Global identification of microRNA-target RNA pairs by parallel analysis of RNA ends. *Nat Biotechnol*, **26**, 941–946.

24. Willmann,M.R., Berkowitz,N.D. and Gregory,B.D. (2014) Improved genome-wide mapping of uncapped and cleaved transcripts in eukaryotes-GMUCT 2.0. *Methods*, **67**, 64–73.

25. Garre,E., Pelechano,V., Sánchez del Pino,M., Alepuz,P. and Sunnerhagen,P. (2018) The Lsm1-7/Pat1 complex binds to stress-activated mRNAs and modulates the response to hyperosmotic shock. *PLoS Genet*, **14**, 1–30.

26. Rodríguez-Galán,O., García-Gómez,J.J., Rosado,I. V, Wei,W., Méndez-Godoy,A., Pillet,B., Alekseenko,A., Steinmetz,L.M., Pelechano,V., Kressler,D., *et al.* (2021) A functional connection between translation elongation and protein folding at the ribosome exit tunnel in Saccharomyces cerevisiae . *Nucleic Acids Res*, **49**, 206–220.

27. Zhang,Y., Xu,P., Xue,W., Zhu,W. and Yu,X. (2023) Diurnal gene oscillations modulated by RNA metabolism in tomato. *The Plant Journal*, **116**, 728–743.

28. Clough,S.J. and Bent,A.F. (1998) Floral dip: A simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant Journal*, **16**, 735–743.

29. Sorenson R, Johnson K, Adler F and Sieburth L (2022) RNAdecay: Maximum Likelihood Decay Modeling of RNA Degradation Data. *R Package*.

30. Boubegtitene,A. and Merret,R. (2022) Monitoring mRNA Half-Life in Arabidopsis Using Droplet Digital PCR. *Plants*, **11**, 1–9.

31. Carpentier,M.-C., Bousquet-Antonelli,C. and Merret,R. (2021) Fast and Efficient 5 ' P Degradome Library Preparation for Analysis of Co-Translational Decay in Arabidopsis. *Plants*, **10**, 1–9.

32. Tesina,P., Heckel,E., Cheng,J., Fromont-Racine,M., Buschauer,R., Kater,L., Beatrix,B., Berninghausen,O., Jacquier,A., Becker,T., *et al.* (2019) Structure of the 80S ribosome–Xrn1 nuclease complex. *Nat Struct Mol Biol*, **26**, 275–280.

33. Zhang,Y., Xu,P., Xue,W., Zhu,W. and Yu,X. (2023) Diurnal gene oscillations modulated by RNA metabolism in tomato. *Plant Journal*, 10.1111/tpj.16400.

34. Song,G., Hsu,P.Y. and Walley,J.W. (2018) Assessment and Refinement of Sample Preparation Methods for Deep and Quantitative Plant Proteome Profiling. *Proteomics*, **18**, 1800220.

35. Chattopadhyay,S., Garcia-Martinez,J., Haimovich,G., Fischer,J., Khwaja,A., Barkai,O., Chuartzman,S.G., Schuldiner,M., Elran,R., Rosenberg,M.I., *et al.* (2022) RNA-controlled nucleocytoplasmic shuttling of mRNA decay factors regulates mRNA synthesis and a novel mRNA decay pathway. *Nat Commun*, **13**, 1–17.

36. Sun,M., Schwalb,B., Pirkl,N., Maier,K.C., Schenk,A., Failmezger,H., Tresch,A. and Cramer,P. (2013) Global analysis of Eukaryotic mRNA degradation reveals Xrn1-dependent buffering of transcript levels. *Mol Cell*, **52**, 52–62.

37. Haimovich,G., Medina,D.A., Causse,S.Z., Garber,M., Millán-Zambrano,G., Barkai,O., Chávez,S., Pérez-Ortín,J.E., Darzacq,X. and Choder,M. (2013) XGene expression is circular: Factors for mRNA degradation also foster mRNA synthesis. *Cell*, **153**, 1000–1011.

38. Wawer,I., Golisz,A., Sulkowska,A., Kawa,D., Kulik,A. and Kufel,J. (2018) mRNA decapping and 5'-3' decay contribute to the regulation of ABA signaling in Arabidopsis thaliana. *Front Plant Sci*, **9**, 1–12.

39. Narsai,R., Howell,K.A., Millar,A.H., O'Toole,N., Small,I. and Whelan,J. (2007) Genome-wide analysis of mRNA decay rates and their determinants in Arabidopsis thaliana. *Plant Cell*, **19**, 3418–3436.

40. Burow,D.A., Umeh-Garcia,M.C., True,M.B., Bakhaj,C.D., Ardell,D.H. and Cleary,M.D. (2015) Dynamic regulation of mRNA decay during neural development. *Neural Dev*, **10**, 1–16.

41. Johnson,E.L., Robinson,D.G. and Coller,H.A. (2017) Widespread changes in mRNA stability contribute to quiescence-specific gene expression patterns in a fibroblast model of quiescence. *BMC Genomics*, **18**, 1–9.

42. Lange,H. and Gagliardi,D. (2022) Catalytic activities, molecular connections, and biological functions of plant RNA exosome complexes. *Plant Cell*, **34**, 967–988.

43. Wang,X., Kong,W., Wang,Y., Wang,J., Zhong,L., Lao,K., Dong,X., Zhang,D., Huang,H., Mo,B., *et al.* (2022) Uridylation and the SKI complex orchestrate the Calvin cycle of photosynthesis through RNA surveillance of TKL1 in Arabidopsis. *Proceedings of the National Academy of Sciences*, **119**, e2205842119.

44. Roux,M.E., Rasmussen,M.W., Palma,K., Lolle,S., Regué,À.M., Bethke,G., Glazebrook,J., Zhang,W., Sieburth,L., Larsen,M.R., *et al.* (2015) The mRNA decay factor PAT 1 functions in a pathway including MAP kinase 4 and immune receptor SUMM 2 . *EMBO J*, **34**, 593–608.

45. Zuo,Z., Edna Roux,M., Dagdas,Y.F., Rodriguez,E. and Petersen,M. (2022) PAT mRNA decapping factors function specifically and redundantly during development in Arabidopsis. *bioRxiv*, 10.1101/2022.07.06.498930.

46. Sun,M., Schwalb,B., Schulz,D., Pirkl,N., Etzold,S., Larivière,L., Maier,K.C., Seizl,M., Tresch,A. and Cramer,P. (2012) Comparative dynamic transcriptome analysis (cDTA) reveals mutual feedback between mRNA synthesis and degradation. *Genome Res*, **22**, 1350–1359.

47. Kastenmayer,J.P. and Green,P.J. (2000) Novel features of the XRN-family in Arabidopsis: Evidence that AtXRN4, one of several orthologs of nuclear Xrn2p/Rat1p, functions in the cytoplasm. *Proc Natl Acad Sci U S A*, **97**, 13985–13990.

48. Han,W.-Y., Hou,B.-H., Lee,W.-C., Chan,T.-C., Lin,T.-H. and Chen,H.-M. (2023) Arabidopsis mRNA decay landscape shaped by XRN 5'-3' exoribonucleases. *The Plant Journal*, **114**, 895–913.

49. Gy,I., Gasciolli,V., Lauressergues,D., Morel,J.B., Gombert,J., Proux,F., Proux,C., Vaucheret,H. and Mallory,A.C. (2007) Arabidopsis FIERY1, XRN2, and XRN3 are endogenous RNA silencing suppressors. *Plant Cell*, **19**, 3451–3461.

TABLE AND FIGURES LEGENDS

Figure 1. Generation of AtXRN4ΔCTRD transgenic line. A. Alignment of the *Arabidopsis thaliana* XRN4 protein with orthologs from Ce, *Caenorhabditis elegans*; Hs, *Homo sapiens*; Sc, *Saccharomyces cerevisiae*. Only the first 360 amino acids are only represented. The position of the L3 loop in *Saccharomyces cerevisiae* is presented (positions 227-244). To generate the AtXRN4ΔCTRD line, a mEGFP was inserted after the position E263. Red arrows indicate positions of S235 in ScXRN1 and E263 in AtXRN4. B. Superposition of ScXRN1 and AtXRN4 AlphaFold prediction. Regions corresponding to L1, L2 and L3 loops in ScXRN1 are only represented. Red asterisks indicate the location of mEGFP insertion. C. XRN4 accumulation in XRN4ΔCTRD lines compared to Wild-type (Col0) and *xrn4.5* using XRN4 specific antibody. The fold change relative to Wild-type is indicated. N=4 biological replicates. A t-test was performed to test significance. n.s., not significant.

Figure 2. AtXRN4CTRD transgenic line is impaired in co-translational mRNA decay. A. Distribution of XRN4 in free mRNPs and ribosome-bound fractions in WT, XRN4ΔCTRD line and XRN4-GFP complemented line using XRN4 specific antibody. The same percentage of each fraction was loaded. UGPase antibody was used as a negative control. B. Proportion of XRN4 in free mRNPs (light colored bar) and ribosome-bound (dark colored bar) fractions in

each line. N=4 biological replicates. A t-test was performed to test significance. All blots were prepared and immunoblotted in parallel and simultaneously exposed for chemiluminescence quantification. C. Metagene analysis of 5'P reads accumulation around stop codons in both shoot and root. D, E. Distribution of Terminational Stalled index (TSI) in both shoot and root. N = 3 biological replicates. A Wilcoxon test was performed to test significance.

Figure 3. Analysis of mRNAs stability in WT root and shoot. A. Boxplots of mRNA half-lives in shoot and root (N = 12 306). B. Distribution of mRNA half-lives in shoot (green bars) and root (red bars). C. Distribution of the fold change between $t_{1/2}$ shoot and $t_{1/2}$ root separated into GO slim groups. The median value of the whole dataset is represented by a red line. Only groups with at least 10 members were used for the analysis.

Figure 4. RNA half-life distributions in *xrn4* and XRN4ΔCTRD lines. Data are presented in shoot (A) or in root (B). A Wilcoxon test was performed to test significance. N=12 306.

Figure 5. Identification of XRN4 co-translational and cytosolic targets and RNA decay feedback in shoot and root. A. Heat map representing $t_{1/2}$ in XRN4ΔCTRD and *xrn4* relative to $t_{1/2}$ in WT. Transcripts are organized in α subgroups (indicated on each cluster) and ordered in 'XRN4 targets' and 'RNA decay feedback targets' groups. N= 7 882 in shoot and 9 291 in root. B. Comparison of XRN4 CTRD specific targets (subgroups 1.4 and 2.2) in shoot and root and their behaviour in the respective organ. C. Comparison of XRN4 cytosolic specific targets (subgroup 4.2) in shoot and root and their behaviour in the respective organ. Both decays : subgroup 1.6; CTRD specific decay : subgroups 1.4 and 2.2; Cytosolic specific decay : subgroup 4.2; RNA decay feedback targets : subgroups 1.1, 1.2, 1.3, 1.5, 2.1, 3.1, and 4.1; Non targets : 5.1.

Figure 6. GO term analysis reveals specific biological functions targeted by cytosolic and co-translational decays. GO term analysis was performed on XRN4 CTRD specific targets (subgroups 1.4 and 2.2), XRN4 cytosolic specific targets (subgroup 4.2) and RNA decay feedback targets (subgroups 1.1; 1.2; 1.3; 1.5; 2.1; 3.1; 4.1) on shoot (A) and root (B).

Figure 7. Part of the feedback mechanism is mediated by cytosolic XRN4. mRNA half-life was determined according to data presented on Supplemental Figure 14. mRNA stability was determined *in vivo* after cordycepin treatment followed by RT-ddPCR on selected transcripts

from subgroups 4.2 (A) and 1.5 (B). N = 3 biological replicates. Normalization was performed using Luciferase RNA spike-in. Mean ± SD. The significance was tested against WT and is indicated below each mRNA half-life. The color chart indicates slower or faster decay as on Figure 5.

Figure 8. XRN4ΔCTRD line presents similar phenotypes as *xrn4* at leaf level and opposite phenotypes at root level. WT, *xrn4*, XRN4ΔCTRD line, and XRN4-GFP complemented line rosette morphology (A), rosette fresh weight (B), number of leaves (C), primary root length under normal (D) and salt stress conditions (E). Red arrows indicate serrated leaves. A t.test was performed to test significance. N > 18.

Supplemental Figure 1. The complete DNA sequence encoding for XRN4ΔCTRD. Red : Sequence encoding XRN4 CDS. Green : Sequence encoding for mEGFP. Blue : Sequence corresponding to promoter region and 5'UTR.

Supplemental Figure 2. Metagene analysis of 5'P reads accumulation around start codon. A. Shoot. B. Root. Blue line : WT, Orange line : XRN4ΔCTRD n°1, grey line : *xrn4*.

Supplemental Figure 3. Subcellular localisation of XRN4 in AtXRN4ΔCTRD (A) and XRN4-GFP lines (B). White lines correspond to 10 µm.

Supplemental Figure 4. Experimental design and workflow used to determine mRNA half-life. A. Experimental Design. 15-d-old seedlings from WT, XRN4ΔCTRD n°1, XRN4ΔCTRD n°2, and *xrn4* genotypes were treated *in vivo* with cordycepin. Roots and shoots were collected separately at T_0, $T_{7.5}$, T_{15}, T_{30}, T_{60}, T_{120}, T_{240} and T_{480} minutes. B. Workflow used to determine mRNA half-life ($t_{1/2}$) and alpha groups.

Supplemental Figure 5. RNAseq data reproducibility. Principal component analysis for WT (A), *xrn4* (B), XRN4ΔCTRD n°1 (C) and XRN4ΔCTRD n°2 (D) RNAseq samples. N = 4 biological replicates per genotype.

Supplemental Figure 6. Volcano plot representing transcripts significantly up- or down-regulated between XRN4ΔCTRD lines. Comparisons were performed between each time-point using DESeq pipeline. FC >2 or <0.5 and p-value <0.05. The number of differentially expressed genes is indicated on each graph.

Supplemental Figure 7. Short mRNA half-lives present a small intron numbers and are correlated with increased A and decreased G content in the 5' UTR in both shoot and root. A. Distribution of mRNA half-life in WT according to numbers of introns in shoot and root respectively. B. Correlation between 5'UTR composition and mRNA half-life in both shoot and root. Black line indicates a Loess smoothing function.

Supplemental Figure 8. Analysis of SKI2 mRNA stability in shoot and root. After transcriptional inhibition by addition of cordycepin, roots and shoots were collected separately at 0, 30, 60, 120, 240 and 480 min. After RNA extraction and reverse-transcription using random primers, SKI2 mRNA stability was analysed by ddPCR using specific primers. Data were normalized according to Luciferase RNA spike-in signal. Mean ± SD. N = 3 biological replicates.

Supplemental Figure 9. Protein accumulation of some 5'-3' decay components in shoot and root. Total proteins extracted were analysed by immunoblotting. An equivalent of 5 mg of tissue powder was analysed in both root and shoot. Specific antibodies against XRN4, DCP5, PAT1, PAT1H1, PAT1H2 and UGPase were used.

Supplemental Figure 10. The two independent XRN4ΔCTRD lines present an RNA decay feedback mechanism in root. RNA half-life distributions in XRN4ΔCTRD line n°1 (A) and line n°2 (B) in comparison to WT and *xrn4*.

Supplemental Figure 11. Genotypic effects on decay rate in shoot and root. According to the impact of the genotype on mRNA half-life, transcripts are categorized in different α groups. α group 1 corresponds to transcripts with a different half-life in each genotype. α group 2 corresponds to transcripts with similar half-life in *xrn4* and XRN4ΔCTRD. α group 3 corresponds to transcripts with similar half-life in *xrn4* an WT. α group 4 corresponds to transcripts with similar half-life in XRN4ΔCTRD and WT. α group 5 corresponds to transcripts with identical half-life in each genotype. The number of transcripts per group is indicated. Red boxes : α groups considered as 'XRN4 targets'. Blue boxes : α groups considered as "RNA decay feedback targets". White box : α group considered as "non XRN4-targets". N = 12 306.

Supplemental Figure 12. Loss of CTRD does not lead to transcriptional feedback. Distribution of fold change between XRN4ΔCTRD and WT from T_0 root samples. whole : whole

dataset, CTRD specific targets : subgroups 1.4 and 2.2, Cytosolic specific targets : subgroups 4.2, RNA decay feedback targets : subgroups 1.1, 1.2, 1.3, 1.5, 2.1, 3.1 and 4.1.

Supplemental Figure 13. Generation of AtXRN4ΔCTRD (R118A/R119A) line. A. Alignment of the *Arabidopsis thaliana* XRN4 protein with orthologs from Sc, *Saccharomyces cerevisiae*; Dm, *Drosophila melanogaster*; Hs, *Homo sapiens* and Ce, Caenorhabditis elegans. The first 160 amino acids are only represented. The position of the L3 loop in *Saccharomyces cerevisiae* is presented (positions 227-244). To generate the AtXRN4ΔCTRD ($R_{118}A/R_{119}A$) line, Alanine substitutions were performed on R118 and R119 in XRN4. The red stars indicate the two Arginine residues conserved in the different XRNs. B. XRN4 accumulation in AtXRN4ΔCTRD (R118A/R119A) line compared to Wild-type (Col0) and AtXRN4ΔCTRD line using XRN4 specific antibody.

Supplemental Figure 14. mRNA half-life determination on selected transcripts. mRNA stability was determined *in vivo* after cordycepin treatment followed by RT-ddPCR on selected transcripts from subgroup 4.2 (A) (from left to right : At1g55280, At5g15820, At4g16780) and from subgroup 1.5 (B) (from left to right : At2g04690, At4g11600, At5g49500). N = 3 biological replicates. Normalization was performed using Luciferase RNA spike-in. Mean ± SD.

Supplemental Table 1. List of the 50 genes used for data normalization.

Supplemental Table 2. mRNA decay analysis results.

Supplemental Table 3. Primers used in this study.

Supplemental Table 4. Transcripts associated with GO slim terms "Root morphology" and "Root development" in subgroups 1.3, 1.5 and 3.1.

Figure 1. Generation of AtXRN4ΔCTRD transgenic line. A. Alignment of the *Arabidopsis thaliana* XRN4 protein with orthologs from Ce, *Caenorhabditis elegans*; Hs, *Homo sapiens*; Sc, *Saccharomyces cerevisiae*. Only the first 360 amino acids are only represented. The position of the L3 loop in *Saccharomyces cerevisiae* is presented (positions 227-244). To generate the AtXRN4ΔCTRD line, a mEGFP was inserted after the position E263. Red arrows indicate positions of S235 in ScXRN1 and E263 in AtXRN4. B. Superposition of ScXRN1 and AtXRN4 AlphaFold prediction. Regions corresponding to L1, L2 and L3 loops in ScXRN1 are only represented. Red asterisks indicate the location of mEGFP insertion. C. XRN4 accumulation in XRN4ΔCTRD lines compared to Wild-type (Col0) and *xrn4.5* using XRN4 specific antibody. The fold change relative to Wild-type is indicated. N=4 biological replicates. A t-test was performed to test significance. n.s., not significant.

Figure 2. AtXRN4CTRD transgenic line is impaired in co-translational mRNA decay. A. Distribution of XRN4 in free mRNPs and ribosome-bound fractions in WT, XRN4ΔCTRD line and XRN4-GFP complemented line using XRN4 specific antibody. The same percentage of each fraction was loaded. UGPase antibody was used as a negative control. B. Proportion of XRN4 in free mRNPs (light colored bar) and ribosome-bound (dark colored bar) fractions in each line. N=4 biological replicates. A t-test was performed to test significance. All blots were prepared and immunoblotted in parallel and simultaneously exposed for chemiluminescence quantification. C. Metagene analysis of 5'P reads accumulation around stop codons in both shoot and root. D, E. Distribution of Terminational Stalled index (TSI) in both shoot and root. N = 2 biological replicates. A Wilcoxon test was performed to test significance.

Figure 3. Analysis of mRNAs stability in WT root and shoot. A. Boxplots of mRNA half-lives in shoot and root (N = 12 306). B. Distribution of mRNA half-lives in shoot (green bars) and root (red bars). C. Distribution of the fold change between $t_{1/2}$ shoot and $t_{1/2}$ root separated into GO slim groups. The median value of the whole dataset is represented by a red line. Only groups with at least 10 members were used for the analysis.

Figure 4. RNA half-life distributions in *xrn4* and XRN4ΔCTRD lines. Data are presented in shoot (A) or in root (B). A Wilcoxon test was performed to test significance. N=12 306.

Figure 5. Identification of XRN4 co-translational and cytosolic targets and RNA decay feedback in shoot and root. Heat map representing $t_{1/2}$ in XRN4ΔCTRD and *xrn4* relative to $t_{1/2}$ in WT in shoot (A) or in root (B). Transcripts are organized in α subgroups (indicated on each cluster) and ordered in 'XRN4 targets' and 'RNA decay feedback targets' groups. N= 7 882 in shoot and 9 291 in root. C. Comparison of XRN4 CTRD specific targets (subgroups 1.4 and 2.2) in shoot and root. D. Comparison of XRN4 cytosolic specific targets (subgroup 4.2) in shoot and root.

Figure 6. GO term analysis reveals specific biological functions targeted by cytosolic and co-translational decays. GO term analysis was performed on XRN4 CTRD specific targets (subgroups 1.4 and 2.2), XRN4 cytosolic specific targets (subgroup 4.2) and RNA decay feedback targets (subgroups 1.1, 1.2, 1.3, 1.5, 2.1, 3.1 and 4.1) on shoot (A) and root (B).

A

	Shoot mRNA half-life (minutes ± SD)				Root mRNA half-life (minutes ± SD)			
	Subgroup 4.2							
	WT	XRN4ΔCTRD	*xrn4*	XRN4ΔCTRD (R118A/R119A)	WT	XRN4ΔCTRD	*xrn4*	XRN4ΔCTRD (R118A/R119A)
At1g55280	24.3 ± 1.2	23.7 ± 3.3 n.s.	109.6 ± 29.6 *	156.3 ± 66.4 *	24.3 ± 1.2	23.7 ± 3.3 n.s.	109.6 ± 29.6 *	156.3 ± 66.4 *
At5g15820	23.8 ± 2.3	28.2 ± 1.1 n.s.	80 ± 5.9 ***	90.2 ± 9.5 ***	31.2 ± 3.0	33 ± 1.9 n.s.	53.9 ± 6.7 **	62.3 ± 14.1 *
At4g16780	44.3 ± 1.8	46.7 ± 3.5 n.s.	170 ± 10.1 **	285.6 ± 71.4 *	35.3 ± 1	36.5 ± 1.4 n.s.	74.1 ± 5.3 **	69.1 ± 7.9 **

B

	Root mRNA half-life (minutes ± SD)			
	Subgroup 1.5			
	WT	XRN4ΔCTRD	*xrn4*	XRN4ΔCTRD (R118A/R119A)
At2g04690	61.0 ± 4.0	38.7 ± 5.1 **	123.6 ± 19.8 *	149.7 ± 48.0 *
At4g11600	52.4 ± 0.4	35.8 ± 3.4 **	158.7 ± 39.4 *	161.1 ± 23.3 **
At5G49500	60.4 ± 1.8	33.9 ± 4.0 **	110.7 ± 8.2 **	133.3 ± 32.7 *

Figure 7. Part of the feedback mechanism is mediated by cytosolic XRN4. mRNA half-life determination on selected transcripts. mRNA half-life was determined according to data presented on Supplemental Figure 14. mRNA stability was determined *in vivo* after cordycepin treatment followed by RT-ddPCR on selected transcripts from subgroups 4.2 (A) and 1.5 (B). N = 3 biological replicates. Normalization was performed using Luciferase RNA spike-in. Mean ± SD. The significance was tested against WT and is indicated below each mRNA half-life. The color chart indicates slower or faster decay as on Figure 5.

Figure 8. XRN4ΔCTRD line presents similar phenotypes as *xrn4* at leaf level and opposite phenotypes at root level. WT, *xrn4*, XRN4ΔCTRD line, and XRN4-GFP complemented line rosette morphology (A), rosette fresh weight (B), number of leaves (C), primary root length under normal (D) and salt stress conditions (E). Red arrows indicate serrated leaves. A t.test was performed to test significance. N > 18.