Spatial Transcriptomic Analysis Identifies Epithelium-Macrophage Crosstalk in Endometriotic Lesions

Gregory W. Burns[1], Zhen Fu[2], Erin L. Vegter[1], Zachary B. Madaj[2], Erin Greaves[3], Idhaliz Flores[4], and Asgerally T. Fazleabas[1]*

[1]Department of Obstetrics, Gynecology and Reproductive Biology, Michigan State University, Grand Rapids, MI, USA

[2]Bioinformatics and Biostatistics, Van Andel Institute, Grand Rapids, MI, USA

[3]Division of Biomedical Sciences, University of Warwick, Coventry, UK

[4]Department of Basic Sciences, Ponce Research Institute, Ponce Health Sciences University, Ponce, PR, USA

*Corresponding author
Email: fazleaba@msu.edu
Address: 400 Monroe Ave NW, GRRC 3020, Grand Rapids, MI 49503 USA
Telephone: +1 616-234-0981

Abstract

The mechanisms underlying the pathophysiology of endometriosis, characterized by the presence of endometrium-like tissue outside the uterus, remain poorly understood. This study aimed to identify cell type-specific gene expression changes in superficial peritoneal endometriotic lesions and elucidate the crosstalk among the stroma, epithelium, and macrophages compared to patient-matched eutopic endometrium. Surprisingly, comparison between lesions and eutopic endometrium revealed transcriptional similarities, indicating minimal alterations in the sub-epithelial stroma and epithelium of lesions. Spatial transcriptomics highlighted increased signaling between the lesion epithelium and macrophages, emphasizing the role of the epithelium in driving lesion inflammation. We propose that the superficial endometriotic lesion epithelium orchestrates inflammatory signaling and promotes a pro-repair phenotype in macrophages, providing a new role for Complement 3 in lesion pathobiology. This study underscores the significance of considering spatial context and cellular interactions in uncovering mechanisms governing disease in endometriotic lesions.

Introduction

Endometriosis affects around 10% of reproductive age women and is defined by the presence of endometrium-like tissue growing outside of the uterus [1-3]. The disease is clinically associated with life-altering pain and infertility and up to 50% of infertile women are diagnosed with endometriosis [4, 5]. Endometriotic lesions are estrogen-dependent and the only effective non-surgical treatment is suppression of ovarian function to limit estrogen production. Endometriotic lesions in the peritoneal cavity are classified by location with lesions present on the visceral or parietal peritoneum termed peritoneal lesions and those on the ovary termed endometriomas. Peritoneal lesions are further separated into superficial or deep infiltrating according to the invasion depth of underlying tissue with deep infiltrating lesions invading more than 5 mm [6, 7]. Superficial peritoneal endometriotic lesions likely develop from endometrial tissue fragments that are refluxed into the peritoneal cavity during menstruation [8, 9]. This hypothesis is supported by the baboon model of endometriosis in which menstrual tissue inoculated into the peritoneal cavity results in endometriotic lesions and disease that persists for at least 15 months [10-12]. Endometriosis can also be induced in mice with tissue collected from an artificial menstruation cycle [13, 14].

Superficial peritoneal lesions generally consist of endometrium-like epithelium and stroma, fibroblasts that express smooth muscle actin (ACTA2), and immune cells [15]. The presence of ACTA2+ fibroblasts is positively correlated with fibrotic lesions where the presence of epithelium is less common, in contrast to lesions of endometrium-like stroma and epithelium [16, 17]. During lesion development, activation of pro-inflammatory immune populations, specifically macrophages, provide a hospitable inflammatory environment to protect the ectopic endometrial tissue [18]. Previous bulk and single-cell RNA-sequencing analyses have uncovered thousands of differentially expressed genes (DEG) in endometriotic lesions, notably including inflammation- and immune-related genes [19-23]. While bulk RNA-sequencing offers

comprehensive data, it lacks the ability to discern information about individual cell types.

Conversely, single-cell RNA-sequencing provides insights into cell type-specific expression but

relies on post-hoc cell type identification using known markers and lacks spatial context. The

emergence of spatial transcriptomics has addressed these limitations by providing a powerful tool

to understand the spatial context of gene expression within tissues, particularly when coupled

with canonical cell type immunostaining. Two of the most prominent platforms for spatial

transcriptomics, NanoString GeoMx and 10X Visium, offer distinct advantages and limitations

depending on experimental design. In this study, we chose the GeoMx platform based on the

ability to provide cell type specific data from formalin fixed paraffin embedded (FFPE) tissues with

high specificity based on antibody-based fluorescent segmentation [24-26]. Our aim was to

determine the genes and pathways altered in the glandular epithelium, stroma, and macrophages

within endometriotic lesions compared to matched endometrium with spatial transcriptomics. We

hypothesized that gene expression changes in superficial peritoneal endometriotic lesions would

manifest in a cell type-specific manner, with disruption of the crosstalk among the stroma,

epithelium, and macrophages being evident compared to patient-matched eutopic endometrium.

Results

Spatial Transcriptomics of Human Superficial Peritoneal Lesions

We sought to determine the contributions of epithelium, stroma, and myeloid cell

compartments of endometriotic lesions to altered genes and pathways in endometriotic lesions

by spatial transcriptomic analyses. Superficial peritoneal lesions and eutopic uterine tissue were

collected from women in the secretory phase of the menstrual cycle. Matched lesion and eutopic

tissues from five women (n = 10) were selected based on the histological presence of endometrial

gland-like structures in lesions confirmed by trichrome staining (Figure 1). The average age of the

patients was 42 (Table 1, SD = 6.1). Tissues were stained for pan-cytokeratin, smooth muscle

actin, and CD68 to identify epithelium (pan-cytokeratin+), stroma (ACTA2) and myeloid-lineage

cells (CD68+, monocytes and macrophages). Myeloid cells were present in both eutopic

endometrium and endometriotic lesions within stromal and epithelial compartments (Figure 1).

ACTA2+ cells were present in the vasculature and stroma, although not in the immediate sub-

epithelial stroma. Duplicate regions of interest were selected for each tissue and segmented

based on fluorescence intensity for macrophages, epithelium, and stroma, resulting in 60

segments.

Based on the limit of quantification, 10 segments were removed with a gene detection rate

of lower than 10% (Figure 2A). Read counts were normalized for the remaining 50 segments

using the quartile 3 (Q3) method. Separation between the geometric mean of the negative control

probes, used to calculate the limit of detection, and Q3 of all gene read counts was confirmed at

the segment and distribution levels to ensure compatibility with Q3 normalization (Figure 2B).

Next, genes detected in less than 10% of segments were removed leaving 7,945 genes (Figure

2C). Overall gene expression across segments was visualized with a Uniform Manifold

Approximation and Projection (UMAP) plot, which resulted in two major clusters (Figure 2D).

Epithelial segments were contained in one cluster and the second cluster included both stroma

and macrophage segments. Notably, segments from the same patient were found in close

proximity within each cluster.

Epithelium and Stroma Marker Genes Are Similar in Lesions and Endometrium

We compared the epithelium and stroma segments by tissue type to confirm cell type

separation and identify differences in the cell identity in the lesion tissue. The eutopic endometrial

epithelium was compared to matching stroma and 1,372 DEG corresponding to cell type were

found (Figure 3A). Similarly, there were 1,313 DEG in lesion epithelium versus stroma (Figure 3B)

and 14 of the top 20 genes were in common. Gene expression patterns were similar in the

endometrium and lesion tissues with a correlation coefficient of 0.81 ($p < 2.2 \times 10^{-16}$, Figure 3C).

Stromal and epithelial biological process terms were distinct and consistent with cell type (Figure

4D). Vascularization, extracellular matrix, and cell adhesion terms were enriched in the stroma

while epithelial cell differentiation and cell-cell junction terms were enriched in the epithelium.

Terms were very similar in the stroma with 9 of 10 terms common between the eutopic

endometrium and lesions. However, five terms were unique to the lesion epithelium, related to

cell-cell adhesion and cell projections or cilia, and two terms to the eutopic endometrial epithelium,

cellular transition metal ion homeostasis and circulatory system process.

Endometriotic Stroma is Minimally Altered

The sub-epithelial stroma from lesions was compared to matched eutopic endometrium to

determine transcriptome alterations in endometriotic lesions. Separation of eutopic stroma from

the lesions was strongest on PC3, accounting for 12.37% of variation in the dataset (Figure 4A).

Two downregulated genes were identified in lesions (Figure 4B), *DHRS1*,

dehydrogenase/reductase 1, and *SCGB2A1*, secretoglobin family 2A member 1, both related to

steroid metabolism. Six hallmark gene sets [27] were significant in a GSEA, four increased and

two decreased (Figure 4C). The increased gene sets were inflammation-related, TNFα signaling

via NF-κB, interferon gamma response, interferon alpha response, and inflammatory response,

and the decreased gene sets were MYC targets V1 and epithelial mesenchymal transition.

Immune Response is Increased in the Endometriotic Lesion Epithelium

The endometriotic epithelium was compared to the matched eutopic glandular epithelium

and was most separated from the eutopic endometrial epithelium on PC3, representing 9.45% of

variation in gene expression (Figure 5A). Differential expression analysis found 76 DEG with 33

increased in the lesion epithelium and 43 decreased (Figure 5B). Notably, multiple immune-

related genes were increased including, *C3*, or complement 3, *CD74*, *CLU*, or clusterin, and major

histocompatibility complex class II members *HLA-DRB1* and *HLA-DPB1*. There were 10

significant hallmark gene sets from a GSEA with four terms increased and six decreased (Figure

5C). Among the increased gene sets were interferon gamma response and allograft rejection, which matched the increase in immune-related genes in the volcano plot. Genes decreased in response to damage by UV radiation, a source of tissue damage and inflammation, were decreased in lesions. Compared to the eutopic endometrium, glycolysis, angiogenesis, androgen response, and epithelial mesenchymal transition gene sets were decreased in the lesion epithelium. Genes increased in the lesion epithelium were enriched for 33 gene ontology (GO) biological process terms and nine of the top 10 were related to immune response (Figure 5D). In fact, nine of the 33 upregulated genes were associated with positive regulation of immune response: *CD74*, *HLA-DRB1*, *CLU*, *C3*, *HLA-DPB1*, *RPS19*, *HLA-DQB1*, *CFB*, and *VTCN1*.

Neither Stroma nor Epithelium Clustered by Tissue

Hierarchical clustering dendrograms were plotted for the stromal and epithelial segments (Figures 6A-B) as an unsupervised measure of segment similarity. Neither cell type clustered by eutopic or lesion origin. Segments from two patients were clustered in the stroma and epithelium, that is lesion segments were more like eutopic tissue from the same patient than another lesion. Principal component analyses of the stroma and epithelium (Figures 6C-D) segments revealed that PC1 was driven by patient ID, representing 23.28% and 16.68% of variation in gene expression, respectively.

Macrophages are Altered in Peritoneal Lesions

Lesion macrophages were most separated from eutopic endometrium macrophages on PC3, representing 8.87% of variation in gene expression (Figure 7A). Differential expression analysis found 148 DEG with 35 increased and 113 decreased in the lesion macrophages (Figure 7B). Nine hallmark gene sets were significantly enriched from a GSEA with seven terms increased and two decreased (Figure 7C). The seven increased terms were associated with inflammation and included TNFα signaling via NF-κB, inflammatory response, allograft rejection, and IL2 STAT5 signaling. Gene sets that were decreased were myogenesis and epithelial mesenchymal

transition. Neither macrophage number nor proximity to the epithelium was altered in lesions

compared to eutopic endometrium (Supplemental Figure 1).

To provide context for our findings, we sought to compare our results with bulk sequencing

data from human peritoneal lesions [21]. Human peritoneal lesions were separated from eutopic

endometrium on principal component (PC) one (39.43% of variation) and 3,656 differentially

expressed genes (DEG) were identified in an analysis of bulk RNA-sequencing data

(Supplemental Figures 2A-C). Gene set enrichment analysis (GSEA) [28] found TNFα signaling

via NF-κB, myogenesis, immune response, and adipogenesis hallmark genes [27] increased,

while estrogen response, glycolysis, and multiple cell cycle gene sets were decreased in

endometriotic lesions (Supplemental Figure 1D).

Ligand-Receptor Analysis Identified Epithelium-Macrophage Signaling Axis in

Lesions

CellChat [29] ligand-receptor analysis was performed to determine the cell-cell

communication landscape in endometriotic lesions and eutopic endometrium. Based on 976

significant ligand-receptor interactions, 29 signaling pathways were identified in total. Twenty-six

pathways were present in lesions, 24 in the endometrium, and 21 pathways were common to

both. Three pathways were unique to the endometrium, CXCL, SPP1, and TGF-β, and five unique

to lesions, CD45, CD46, ITGB2, PTPRM, and SEMA3. A heatmap of signaling pathways

consisting of all possible connections, inter- and intra-tissue, ranked by signaling strength (Figure

8A) found collagen signaling to be the strongest followed by MHC-II, laminin, macrophage

migration inhibitory factor (MIF) and complement. The three top interacting cell types were lesion

stroma, endometrium stroma, and lesion macrophages. Visualization of the cell-cell

communication networks in the endometrium and lesions by weight in circle plots showed strong

connections from the stroma compartment to the epithelium in both tissue types (Figure 8B).

Based on combined network weights, confined to tissue type, the top senders were endometrial

(0.172) and lesion stroma (0.170), and the top receivers were lesion macrophages (0.184) and

epithelium (0.104). Particularly striking was a 3.70-fold increase in epithelium-to-macrophage

signaling in lesions compared to eutopic endometrium. Interaction strength heatmaps (Figure 8C),

confirmed the strong connection from the endometrial stroma received by the epithelium. In

lesions, the strongest interaction was macrophage-macrophage signaling followed by epithelium-

macrophage, surpassing stroma to epithelium signaling. These results point to a strengthened

epithelium-macrophage signaling axis in endometriotic lesions.

Given that multiple inflammatory pathways were increased in lesions, including MHC-II

and complement, and since increased signaling was noted between the lesion epithelium and

macrophages, we sought to determine the source of inflammation-related ligands in lesions. A

ligand-receptor bubble plot (Figure 9A) demonstrated that complement and MHC-II ligands were

originating in the lesion epithelium and likely binding to receptors on macrophages. Within the

stroma, MIF was the only visualized inflammatory pathway with likely communication probability

indicating that inflammatory signals originated largely in the lesion epithelium. Indeed, a chord

diagram of complement signaling (Figure 9B) highlighted the increased complement signaling

present in lesions compared to the endometrium and the lesion epithelium as the source.

Discussion

Endometriosis is a complex reproductive disorder marked by the presence of

endometrium-like tissue outside the uterus. Focusing on superficial peritoneal lesions, and

excluding deep infiltrating or ovarian endometrioma disease, we hypothesized that endometrium-

like tissue in these lesions would display an altered molecular phenotype influenced by the

peritoneal microenvironment. To explore this hypothesis, we utilized spatial transcriptomic,

receptor-ligand, and bulk RNA-sequencing analyses of endometriotic lesions compared to eutopic

endometrium.

Bulk RNA-sequencing unveiled thousands of differentially expressed genes (DEGs) and

numerous altered inflammatory pathways in lesions compared to the eutopic endometrium,

highlighting the inflammatory nature of these lesions and the presence of fibrotic tissue. This

observation supported the notion that endometrium-like tissue within the peritoneum would differ

significantly from its uterine counterpart. Inflammatory cytokines are increased in the peritoneal

fluid of women with endometriosis [30], likely influencing gene expression. Indeed, miRNAs,

Notch signaling, aromatase enzymes, prostaglandins, and progestin responsiveness are altered

in lesions compared to matched endometrial tissues [31-42].

Contrary to our initial expectations, spatial analysis revealed transcriptomic similarities

between the three major cell types found in endometrium-like regions of lesions and patient-

matched eutopic endometrium, with macrophages displaying the most substantial transcriptional

difference. This lack of transcriptional divergence in the epithelium and stroma supports the idea

that superficial peritoneal endometriotic lesions are derived from the endometrium following

retrograde menstruation [8, 9]. Further, patient rather than tissue type, that is lesion or

endometrium, was the primary determinant of gene expression based on principal component

and UMAP plots. The differences observed in macrophage gene expression are likely due to the

presence of peritoneal and monocyte-derived macrophages in lesions, compared to tissue

resident macrophages in the eutopic endometrium [43-45].

The discrepancy between bulk RNA-seq and spatial analyses points to the inherent

differences in methodologies. While bulk RNA-seq provides a comprehensive view of gene

expression changes across the entire homogenized tissue, spatial transcriptomics offers a

targeted approach focusing on localized gene expression patterns within distinct regions of

interest, such as the endometrium-like structures of lesions. This targeted approach resulted in

comparatively few DEG, indicating that the high difference in transcriptomic profiles between

lesions and eutopic endometrium found in bulk RNA-seq may not be attributable to endometrium-

like tissue in lesions but from the peripheral tissue. This highlights the importance of spatially

resolved data in heterogenous tissues like endometriotic lesions. For example, we expect many

of the DEG identified in bulk RNA-seq originate from the fibrotic regions of lesions and reflect

sustained inflammation and the presence of activated fibroblasts [9, 16]. Interestingly, our findings

suggest that the sub-epithelial stroma in superficial peritoneal lesions remains largely unaltered

and could serve as an internal control for future spatial experiments, bypassing the requirement

for matched eutopic stromal tissue. The paired design of this study increased power with a limited

sample size. However, the lack of healthy control endometrium and unknown time since symptom

onset are notable caveats and may limit the generalizability of the findings, particularly for the

initiation of disease. As such, further research is necessary to elucidate the molecular

mechanisms underlying the early pathogenesis of superficial peritoneal endometriosis.

The spatial transcriptomic technique, sequencing hybridized probes from specific areas of

FFPE tissue, resulted in an overall reduced transcriptome size after removing genes with low

counts across segments. The spatial transcriptome included 7,945 genes while the bulk RNA-

sequencing analysis included 17,405 genes. Consequently, many of the lowest expression genes

from the bulk RNA-sequencing data were filtered from the spatial transcriptome before DEG

analysis. The advantage of spatial transcriptomics, however, is gaining information from genes

expressed only by a subset of cells or a single cell type that would be lost in the bulk tissue. This

is evidenced by the inclusion of 551 genes in the spatial transcriptome that were considered as

not expressed in the bulk RNA-sequencing analysis. Further, spatial transcriptomics indicated

increased proliferation of the lesion epithelium while bulk RNA-sequencing found an overall

reduction in cell cycle gene sets. The epithelium of lesions is likely to be a minor contributor to

gene expression from homogenized tissue since lesions often have little or even no remaining

epithelial component. Interestingly, these results suggest the lesion epithelium may proliferate in

response to inflammatory peritoneal environment while stroma reduces proliferation and may

even become senescent [46] . A growing body of evidence has found that senescence plays a

role in multiple pathologies and may contribute to endometriotic lesions [47-49].

Receptor-ligand analysis identified an almost four-fold increase in signaling between

epithelium and macrophages in lesions compared to the endometrium, including multiple

inflammatory pathways. MHC-II ligands expressed in the lesion epithelium were predicted to

communicate to macrophages through the CD4 receptor. Notably, human monocytes and

macrophages express CD4, unlike the mouse, serving as an important target for HIV-1 infection

[50-52]. The canonical macrophage marker from this study, CD68, is expressed by monocytes

and macrophages [53]. Although monocytes rapidly differentiate into macrophages within a tissue,

some monocytes were likely included in the spatial transcriptome analysis. Canonically, MHC-II

binding of the CD4 receptor on monocytes promotes differentiation into macrophages [51], thus

increased MHC-II in the lesion epithelium would drive macrophage differentiation from circulating

monocytes homing to areas of hypoxia and inflammation. Increased expression of MHC class II

expression in the epithelium aligns with recent single cell RNA-sequencing analysis of ovarian

endometriosis, which found increased MHC class II expression in the epithelium [54].

Macrophage migration inhibitory factor (MIF) is constitutively expressed by many cell

types and has pleiotropic autocrine and paracrine proinflammatory effects on tissues [55]. MIF

signaling in macrophages sustains proinflammatory function by inhibiting apoptosis,

counteracting the immunosuppressive effects of glucocorticoids, and promoting secretion of

prostaglandin E2 [55, 56]. Additionally, MIF is released by monocytes, macrophages, and other

immune cells upon exposure to proinflammatory mediators, such as C3, to reinforce a local

proinflammatory environment [57-59].

Complement 3 expression was increased in bulk RNA-sequencing results (3.7-fold, FDR

$p = 0.0003$) and found in the lesion epithelium from spatial transcriptomic analysis (4.6-fold, FDR

$p = 0.003$). Endometriotic lesions were originally reported to express Complement 3 (C3) by

Weed and Arquembourg in 1980 [60] and later found to be secreted by the lesion epithelium [61,

62]. In fact, serum levels of C3 are elevated in patients with endometriosis and C3 was proposed

as a potential diagnostic biomarker [63]. The complement system is a signaling cascade that plays

a pleiotropic role in immune response, including production of proinflammatory mediators and

recognition and clearance of pathogens by lysis and enhanced phagocytic uptake by immune

cells [64]. The cascade of over 50 proteins is characterized by a series of protein cleavage events

that produce inflammatory cytokines and the components for subsequent steps. Multiple stimuli

can activate the complement system, but all pathways converge on activation of C3 by a

convertase. Although the presence of C3 in endometriotic lesions has been known for 40 years,

its potential contribution to the disease is unclear [65]. Following cleavage, the resulting fragments

are referred to by appending the letter "a", for the smaller fragment, or "b", for the larger fragment,

to the protein name, for example, C3a.

C3a and C3b are produced following activation of C3 by a C3 convertase, itself the product

of prior cleavage events. C3b induces phagocytic activity of macrophages, but lesion tissue is

refractory to complement targeting through multiple mechanisms. CD47, otherwise known as the

"don't eat me signal", is expressed by tissue and binds to the SIRPα receptor on macrophages,

inhibiting phagocytosis [66, 67]. Complement regulatory proteins, such as CD35, CD46, and

CD55, inhibit complement attack by accelerating breakdown or direct cleavage of complement

components critical for continuation of the cascade [64]. In the event of complement activation,

CD59, another complement regulatory protein, inhibits formation of the membrane attack complex

and prevents cell lysis [68]. Thus, multiple mechanisms exist within the lesion to prevent direct

complement attack but complement signaling promotes a localized proinflammatory environment.

The proinflammatory cytokine C3a, released following C3 cleavage, can activate a pro-

repair or tumor-associated macrophage (TAM) phenotype by binding to C3AR1, as found in the

ligand-receptor analysis [69]. This provides a mechanism by which C3 activation from the

superficial peritoneal endometriotic lesion epithelium promotes the pro-lesion macrophage

phenotype described as pro-repair, tumor-associated, or M2. These macrophages then contribute

to lesion tissue remodeling including neovascularization and fibrosis [18, 70, 71]. Additionally,

semaphorin 3B was found in the lesion epithelium signaling to macrophages and could further

promote the pro-disease TAM phenotype since SEMA3B increased expression of pro-repair

macrophage markers in a model of rheumatoid arthritis [72].

Activated complement signaling in tissue induces a diffuse inflammatory state that

includes increased prostaglandins, also upregulated by MIF [65]. Increased prostaglandins can

activate steroidogenic enzymes and, potentially the production of estradiol and estrone by the

lesion [31, 33, 35, 73]. Finally, increased estrogens can activate complement in lesion tissue [74]

resulting a positive feedback loop, common in pathogenic signaling. In this manner, the

complement signaling is sustained by local estrogenic activity and could sustain local

inflammation and the presence of pro-repair macrophages until such time as the lesion becomes

completely fibrotic with a tissue stiffness that no longer supports cell growth and the lesion

becomes scar tissue and contributes to peritoneal adhesions. In support of complement signaling

promoting lesion pathogenesis, C3 null uterine tissue formed fewer lesions after 21 days in a

mouse model of endometriosis [75].

The comparison of lesions to eutopic endometrium yielded intriguing insights into the

molecular and cellular dynamics underlying lesion pathophysiology. Firstly, despite previous

findings from bulk tissue analysis suggesting significant differences between endometrium-like

tissue in lesions and eutopic endometrium, our spatial transcriptomic data reveal a surprising

similarity between these two tissues. The lack of drastic differences may be attributed to the

composition of the excised lesions used in bulk RNA-sequencing, with endometrium-like tissue

comprising only a small proportion of the lesion, such that differential expression was largely

attributable to the peripheral stroma and fibrotic tissue. Notably, this difference highlights the

importance of employing advanced techniques, like spatial transcriptomics, to gain an improved

understanding of lesion biology.

Secondly, spatial transcriptomics has provided novel insights into the intricate cellular

crosstalk within lesions. Our data indicate signaling between the stroma and epithelium, as well

as between the lesion epithelium and macrophages. Notably, the epithelium emerged as a central

player in driving inflammation within the lesion, promoting inflammatory signaling independent of

the stromal compartment. Moreover, the presence of CD68+ cells amid epithelial cells highlights

the physiological relevance of cell-cell interactions across different cell types.

Drawing from these observations, we present a model wherein the superficial lesion

epithelium orchestrates inflammatory signaling within the lesion while promoting a pro-repair

phenotype in infiltrating macrophages. Spatial transcriptomics facilitated the identification of

inflammatory signaling emanating from the epithelium, including C3, MIF, and MHC class II. This

novel insight sheds light on the pathological consequence of C3 expressed by the epithelium in

modulating macrophage phenotype to support lesion development. In summary, our study

underscores the critical role of considering spatial context and cellular interactions within lesions

to unravel the complexities of superficial peritoneal endometriotic lesions. Further investigation is

imperative to elucidate the specific mechanisms governing epithelium-driven inflammation and its

implications for disease progression and therapeutic intervention.

Methods
Human Tissues

All protocols involving tissue collection were approved by the institutional review board

(IRB) Committee of Ponce Health Sciences University, School of Medicine (PHSU). Endometriotic

lesions and matched eutopic uterine tissues from 10 patients were obtained as archived

deidentified formalin-fixed paraffin-embedded (FFPE) tissue blocks [76-78]. All tissues were

previously evaluated by a pathologist to confirm the diagnosis of endometriosis and menstrual

cycle stage of eutopic endometrium using Noyes criteria [79].

Histology

Tissue blocks were sectioned and trichrome stained (StatLab #KTTRBPT) to confirm the

presence of defined epithelial and stromal compartments in lesions. Blocks were serial sectioned

at 6 μm thickness and floated onto positively charged microscope slides. Tissues were attached

to the slides prior to processing by heating on a slide warmer for 15 minutes. Tissues were then

deparaffinized and rehydrated through three changes each of xylenes for five minutes and 100%

ethanol for one minute before a rinsing in running tap water for one minute. Slides were incubated

in Bouin's fluid at room temperature overnight. Slides were rinsed in running tap water for three

minutes and immersed in modified Mayer's hematoxylin stain for four minutes at room

temperature. After a three-minute rinse in running tap water, slides were immersed in one step

trichrome stain for five minutes and rinsed in running tap water for five seconds. Slides were then

dehydrated through three changes of 100% ethanol for one minute, cleared in three changes of

xylenes for one minute, and a coverslip was applied with Permount mounting medium (Fisher

Scientific #SP15).

Spatial Transcriptomics

Human tissue slide preparation, library generation, and sequencing for digital spatial

profiling (DSP) on the NanoString GeoMx DSP instrument were performed by the Van Andel

Institute Histology and Genomics Cores. Sections were cut at 5 μm thickness and mounted on

plus-charged slides (Epredia Colormark Plus #CM-4951WPLUS-001). Slides were baked at 60°C

for one hour and stored at 4°C in a vacuum-sealed container containing desiccant for up to two

weeks. All subsequent steps were performed under RNase-free conditions with DEPC-treated

water. Slides were deparaffinized with three sequential five-minute washes in xylenes, followed

by two washes in 100% ethanol for five minutes, one wash in 95% ethanol, and one wash in 1X

PBS. Antigen retrieval was performed in target retrieval reagent (EDTA, pH 9.0; Invitrogen #00-

4956-58) diluted to 1X in the BioGenex EZ-Retriever System for 10 minutes at 95°C. Slides were

then washed with 1X PBS for five minutes. Slides were then incubated in 1 µg/mL proteinase K

10 minutes at 37°C and washed in 1X PBS for five minutes at room temperature. Slides were

fixed for five minutes in 10% neutral buffered formalin followed by two washes in NBF stop buffer,

for five minutes each, and one wash in 1X PBS for five minutes. Slides were then incubated with

UV-photocleavable Human Whole Transcriptome Atlas hybridization probes diluted in buffer R

(GeoMx RNA Slide Prep FFPE-PCLN kit, #121300313) in a hybridization oven at 37°C for 16-20

hours. Following probe incubation, slides were washed with stringent buffer (1:1, formamide:4X

SSC buffer; Thermo Fisher #AM9342; Sigma-Aldrich #S6639) at 37°C twice for 25 minutes each.

Then slides were washed twice in 2X SSC buffer. Slides were blocked in 200 µL buffer W (GeoMx

RNA Slide Prep FFPE-PCLN kit) for 30 minutes and incubated at 4ºC overnight with SYTO 13

nucleic acid stain (500 nM; Thermo Fisher Scientific #S7575) and antibodies for pan-cytokeratin

(panCK; 1 µg/mL; Novus Biologicals #NBP2-33200AF488), smooth muscle actin (ACTA2; 1.25

µg/mL; Abcam #ab202368), and CD68 (0.5 µg/mL; Santa Cruz Biotechnology #sc-20060AF647)

diluted in buffer W. Slides were washed four times in 2X SSC buffer for 3 minutes each and placed

in the NanoString GeoMx DSP instrument.

Duplicate 500 µm^2 regions of interest (ROI) were selected for each based on fluorescent

imaging on the GeoMx DSP instrument (NanoString Technologies). ROI were drawn in the central

portion of the endometrium stratum functionalis, to include glandular and not luminal epithelium,

for eutopic uterine tissue and in lesions areas with defined epithelium. ROIs were then segmented

in order of macrophages (CD68+), epithelium (panCK+), and stroma (ACTA2-) areas of

illumination (AOI) for individual collection. AOI were processed sequentially on the instrument

using UV light focused through each AOI that released the photocleavable oligos into wells of a

96 well plate. After AOI collection, indexing primers were hybridized during NanoString library

preparation. Quality and quantity of the finished library pools were assessed using a combination

of Agilent DNA High Sensitivity chips (Agilent Technologies, Inc.) and the QuantiFluor dsDNA

System (Promega Corp.). Paired-end, 50 bp sequencing was performed on an Illumina Novaseq

6000 sequencer using an S2, 100 cycle sequencing kit v1.5 (Illumina Inc.). Base calling was done

by Illumina RTA3 and output was demultiplexed and converted to fastq format with bcl2fastq

v1.9.0.

Spatial Data Analysis

Fastq files were converted to digital count conversion (DCC) files using GeoMx NGS

Pipeline v2.3.3.10. Sample annotation, PKC configuration, and DCC files were imported with

GeomxTools (v3.2.0) in R to generate a NanoStringGeoMxSet object. Read counts with a value

of zero were shifted to one for downstream transformations and quality control (QC) analyses.

Each AOI, or segment, was assessed for sequencing and tissue quality using the parameters

listed in **Box 1**. Outlier probes were then excluded before counts were collapsed to gene-level

data (18,677 genes) by geometric mean. The limit of quantification (LOQ) was determined per

segment based on the distribution of negative control probes and set at two geometric standard

deviations above the geometric mean with a minimum value of two. Segments with a gene

detection rate less than 10% were filtered before removing genes detected in less than 10% of

segments. Separation between the upper quartile 3 (Q3) of gene counts and the geometric mean

of the negative probes was confirmed before count normalization using the quartile 3 (Q3)

method.

Normalized read counts were $\log_2$-transformed and dimension reduction was performed

by uniform manifold approximation and projection (UMAP) [80] with the umap package (v0.2.10)

with the random state set to 42. Differential expression analyses were completed using $\log_2$-

transformed normalized read counts with a linear model approach in the limma package (v3.54.2)

[81]. A linear model was fit for each gene with lmFit before an empirical Bayes test was completed

using patient as a covariate. P-values were corrected for multiple testing with the Benjamini

Hochberg method included in the decideTests function of limma. Genes were considered

differentially expressed when the adjusted p-value was below 0.05. Principal component analysis,

volcano plots, gene set enrichment analysis, and over-representation tests were completed as

described for bulk RNA-sequencing data.

Macrophage Quantification

The macrophage count distance between from epithelium was manually measured for

each segment using the line tool in ImageJ software [82]. The global scale was set by measuring

the scale bar on the image (1 pixel = 2.51 μm). For each image, the green epithelium and red

macrophage channels were merged for analysis. A line was drawn from the center of each

macrophage to the center of the closest epithelial cell and the measurements. Based on these

measurements the macrophages count and the average distance for each segment was

calculated. Data were analyzed in Prism (v10.2.2, GraphPad Software LLC) following

confirmation of normality (Shapiro-Wilk, $p > 0.1$). To determine if there was a relative increase in

lesions compared to the matched endometrium, macrophage count was tested using a ratio

paired t-test. Macrophage distance to the nearest epithelium was tested using a paired t-test.

Bulk RNA-sequencing Analysis

Read counts per gene were downloaded from eutopic endometrium (n = 7) and peritoneal

endometriotic lesion samples (n = 6) for series GSE179640 [21] from the NCBI Gene Expression

Omnibus repository. Files were parsed with R (version 4.2.0) and all subsequent analyses, unless

otherwise noted, were completed in R using RStudio [83]. Differential expression analysis was

conducted using the edgeR-robust method (v4.0.3) [84, 85]. Genes with low counts per million

(CPM) were removed using the filterByExpr function from edgeR. Multidimensional scaling plots,

generated with the plotMDS function of edgeR, were used to investigate group separation prior

to statistical analysis. Principal component analysis of normalized $\log_2$-transformed CPM was

conducted with the PCAtools package (v2.10). The biplot and pairsplot functions were used to visualize sample separation across principal components one through five. Differentially expressed genes (DEG) were identified based on Benjamini-Hochberg false discovery rate (FDR) p-values less than 0.05. The hclust function was used for hierarchical clustering (ward.D2) of Euclidean distances and plotted with the dendextend package (v1.17.1). A volcano plot was generated with EnhancedVolcano (v1.16) and the top 20 DEG, based on FDR p-value, were labeled. Gene set enrichment and over-representation analyses were completed with clusterProfiler package (version 4.6.0) [86] using mSigDB [27] reference databases.

Receptor-Ligand Analysis

Quantile-normalized read counts were used for receptor-ligand analysis with the CellChat package (1.6.10) in R [29]. Significant genes and interactions were identified using the functions `identifyOverExpressedGenes` and `identifyOverExpressedInteractions`. Communication probabilities were calculated between cell types with raw input counts option enabled, as recommended for bulk-sequencing data. The resulting interactions were visualized using the included plotting functions. Weighted communication probabilities were subset for eutopic endometrium or lesions before calculating tissue-specific communication interaction strengths.

Data Availability

Sequence data from the spatial transcriptomics experiment were deposited in the NCBI Gene Expression Omnibus and available under series GSE263897.

Acknowledgements

The authors would like to sincerely thanks all members of the Fazleabas laboratory and the Van Andel Genomics core, particularly Becca Siwicki, for their contributions to this project. This work was supported by NanoString Technologies, Incorporated (GWB) and National Institutes of Health grants F32HD104478 (GWB), R01HD099090 (ATF), R01HD050559 (IF),

476 U56CA126379 (IF), and T32HD087166 (GWB, ATF). The content is solely the responsibility of

477 the authors and does not necessarily represent the official views of the National Institutes of

478 Health.

Author contributions

Conceptualization, G.W.B., E.G., and A.T.F.; Data curation, G.W.B, Z.F., Z.B.M, and I.F.;

Formal analysis, G.W.B, Z.F., E.L.V, and Z.B.M.; Funding acquisition, G.W.B, I.F, and A.T.F.;

Investigation, G.W.B and E.L.V.; Methodology, G.W.B, Z.F., E.L.V, and Z.B.M; Resources, I.F. and

A.T.F; Software, G.W.B, Z.F., and Z.B.M; Supervision, E.G. and A.T.F; Visualization, G.W.B;

Writing – original draft, G.W.B and A.T.F.; Writing – review & editing, G.W.B, Z.F., E.G., I.F., and

A.T.F.

Declaration of interests

487   The authors declare no competing interests.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used ChatGPT (v3.5) for proofreading and

to improve readability. After using this service, the authors reviewed and edited the content as

needed and take full responsibility for the content of the publication.

Figures

Figure 1

Segmentation strategy for spatial transcriptomics. Superficial peritoneal endometriotic lesions and

matched endometrium (n = 10) were visualized with trichrome staining to confirm the presence of

epithelial and stromal compartments. Tissues were immunostained for pan-cytokeratin, CD68,

and ACTA2 for segmentation and spatial transcriptome collection on the GeoMx platform.

Representative regions of interest (ROI) are shown for each tissue type. Duplicate ROI were

collected for each tissue and stroma, epithelium, and macrophages were segmented from each

for a total of 20 ROI and 60 segments. Scale bar = 100 μm.

Figure 2

Spatial transcriptomics results from eutopic endometrium superficial peritoneal endometriotic

lesions. (A) Bar plot of all 60 segments separated by gene detection rate. Segments with a

detection rate of less than 10% were removed. (B) Confirmation of gene and negative probe count

separation by segment in a scatter plot and by distribution for each cell type in histograms.

Separation after gene and segment filtering is required for effective Q3 normalization. (C) Bar plot

of detectable genes per segment used for filtering genes found in less than 10% of segments

leaving 7,945 genes for further analysis. (D) Dimensionality reduction UMAP plot of all segments

coded by shape and colored by patient. Epithelium clustered separately from stroma and

macrophages.

Figure 3

Epithelium versus stroma in eutopic endometrium and lesions. Volcano plots of differentially expressed genes (DEG) in the (A) endometrium and (B) lesions. The top DEG were similar in volcano plots with canonical epithelial genes like *MUC1*, *PAX8*, and *WFDC2* increased in both epithelia. (C) Genes enriched for each compartment were highly correlated in the eutopic endometrium and lesions. (D) Enriched gene ontology biological process terms were consistent with cell type, including extracellular matrix genes in the stroma and cell-cell junction genes in the epithelium, in the eutopic endometrium and lesions. Several terms related to cell-projections and cilia were unique, however, to the lesion epithelium.

Figure 4

Endometriotic sub-epithelial stroma is minimally altered compared to eutopic endometrium. (A)

Principal component (PC) plot of lesion stroma compared to eutopic endometrium showed group

separation most clearly on PC3. (B) Volcano plot of two down-regulated genes in the

endometriotic stroma. (C) Gene set enrichment analysis of hallmark pathways found four

increased inflammatory gene sets and two decreased.

Figure 5

Immune response genes are increased in the endometriotic lesion epithelium. (A) Principal

component (PC) plot of lesion epithelium compared to eutopic endometrium showed group

separation most clearly on PC3. (B) Volcano plot of 76 differentially expressed genes (DEG) in

the endometriotic epithelium. Of note, complement 3, *C3*, was one of the top increased DEG

and several others were major histocompatibility complex class II related genes, like *CD74* and

HLA-DRB1. (C) Gene set enrichment analysis of hallmark pathways found four increased gene

sets and six decreased, indicating increased inflammation and proliferation. (D) Nine of the top

10 enriched gene ontology biological process terms were immune related.

Figure 6

Neither stroma nor epithelium cluster by lesion versus eutopic endometrium. Hierarchical clustering dendrograms of (A) stroma and (B) epithelium segments found no clustering of segments by tissue of origin (endometrium = black bar). Principal component (PC) plots demonstrated that PC1, the largest component of variation, was correlated with patient identity.

Figure 7

Macrophages are altered in endometriotic lesions. A) Principal component (PC) plot of

lesion macrophages compared to eutopic endometrium showed group separation most clearly on

PC3. (B) Volcano plot of 148 differentially expressed genes (DEG) in the endometriotic

macrophages. (C) Gene set enrichment analysis of hallmark pathways found seven increased

gene sets and two decreased, indicating increased inflammation likely reflective of the presence

of monocyte-derived macrophages in the lesion tissue.

Figure 8

CellChat ligand-receptor analysis identified increased communication from the lesion epithelium to macrophages. (A) Signaling pathway heatmap of all 29 significant pathways identified by CellChat sorted by signaling strength, indicated by the gray bar. Interactions, combined incoming and outgoing, totaled for each cell type in the bar plot at the top of the heatmap and scaled per row. (B) Circle plots of communication networks including 24 pathways in the endometrium and 26 in lesions where line thickness corresponds to strength. Stroma to epithelium communication was strong in the eutopic endometrium and similarly present in the lesions. However, epithelium to macrophage communication was increased 3.7-fold in lesions. (C) Interaction strength heatmaps highlight the gain of epithelium to macrophage signals in lesions and strong macrophage-macrophage interactions.

Figure 9

Inflammatory ligands originate in the endometriotic lesion epithelium. (A) Ligand – receptor dot plot of inflammatory pathways in endometriotic lesions. Pathway names are indicated on the left side of the plot followed by specific ligand and receptor interactions. All signals originating from the lesion epithelium are in the left three columns, macrophages in the center, and stroma on the right. Complement and major histocompatibility complex class II (MHC-II) pathways originated in the lesion epithelium and not the stroma. Macrophage migration inhibitory factor (MIF) signaling was strongest in the epithelium, but also sent from the stroma and macrophage cell types. (B) Complement signaling chord diagram of the weighted complement signaling pathway indicating increased signaling from the lesion epithelium compared to the eutopic endometrium epithelium.

```
minSegmentReads = 1000,
percentTrimmed = 80,
percentStitched = 80,
percentAligned = 75,
percentSaturation = 40,
minNegativeCount = 1,
maxNTCCount = 9000,
minNuclei = 20,
minArea = 1000
```

573   Box 1

574   Spatial transcriptomics quality control parameters for segment filtering.

Supp Figure 1

Supplemental Figure 1

Quantification of macrophages in spatial transcriptomic segments from eutopic endometrium and superficial peritoneal lesions. Data are shown by group and individual patient. (A) Macrophage number was not increased in lesions (p = 0.98) and (B) did not appear to be located closer to the lesion epithelium (p = 0.59).

Supplemental Figure 2

Transcriptome alterations in superficial endometriotic lesions compared to eutopic endometrium

as measured by bulk RNA-sequencing. (A) Principal component (PC) plot of superficial peritoneal

endometriotic lesions and eutopic endometrium (n = 13) from GSE179640. Samples were

separated across PC1 by tissue type, representing 39% of variation in gene expression. (B)

Hierarchical clustering dendrogram confirmed separation of lesions from eutopic endometrium.

(C) Volcano plot of 3,656 differentially expressed genes in lesions versus endometrium. (D) Gene

set enrichment analysis of hallmark pathways found inflammation, represented by TNFα signaling

via NF-κB and inflammatory response, increased in lesions while three cell cycle-related gene

sets, MYC targets, G2M checkpoint, and E2F targets were decreased. This indicates increased

inflammation and decreased proliferation in lesions compared to eutopic endometrium.

Tables

Table 1

Patient demographics, indications for surgery, and post-operative diagnosis.

Patient	Age	Indication(s)	Diagnosis
2756	47	Uterine Leiomyoma Adenomyosis	Endometriosis L ovary and oviduct
4909	36	Uterine Leiomyoma Pelvic pain	Endometriosis, Leiomyoma
5322	49	Uterine Leiomyoma	Endometriosis L&R ovary, Leiomyoma
9997	36	Pelvic pain	Endometriosis R oviduct, Endometrioma L ovary, Pelvic adhesions
12529	44	Uterine Leiomyoma	Endometriosis L&R oviduct, Leiomyoma

References

1. Eskenazi B, Warner ML. Epidemiology of endometriosis. Obstet Gynecol Clin North Am. 1997;24(2):235-58. Epub 1997/06/01. PubMed PMID: 9163765.

2. Bulun SE. Endometriosis. N Engl J Med. 2009;360(3):268-79. Epub 2009/01/16. doi: 10.1056/NEJMra0804690. PubMed PMID: 19144942.

3. Giudice LC. Clinical practice. Endometriosis. N Engl J Med. 2010;362(25):2389-98. Epub 2010/06/25. doi: 10.1056/NEJMcp1000274. PubMed PMID: 20573927; PubMed Central PMCID: PMCPMC3108065.

4. Practice Committee of the American Society for Reproductive Medicine. Treatment of pelvic pain associated with endometriosis. Fertil Steril. 2008;90(5 Suppl):S260-9. Epub 2008/11/26. doi: 10.1016/j.fertnstert.2008.08.057. PubMed PMID: 19007642.

5. Endometriosis and infertility: a committee opinion. Fertil Steril. 2012;98(3):591-8. Epub 2012/06/19. doi: 10.1016/j.fertnstert.2012.05.031. PubMed PMID: 22704630.

6. Koninckx PR, Martin DC. Deep endometriosis: a consequence of infiltration or retraction or possibly adenomyosis externa? Fertil Steril. 1992;58(5):924-8. Epub 1992/11/01. doi: 10.1016/s0015-0282(16)55436-3. PubMed PMID: 1426377.

7. Koninckx PR, Ussia A, Adamyan L, Wattiez A, Donnez J. Deep endometriosis: definition, diagnosis, and treatment. Fertil Steril. 2012;98(3):564-71. Epub 2012/09/04. doi: 10.1016/j.fertnstert.2012.07.1061. PubMed PMID: 22938769.

8. Sampson JA. Peritoneal endometriosis due to the menstrual dissemination of endometrial tissue into the peritoneal cavity. American Journal of Obstetrics & Gynecology. 1927;14(4):422-69. doi: 10.1016/S0002-9378(15)30003-X.

9. Zondervan KT, Becker CM, Koga K, Missmer SA, Taylor RN, Vigano P. Endometriosis. Nat Rev Dis Primers. 2018;4(1):9. Epub 2018/07/22. doi: 10.1038/s41572-018-0008-5. PubMed PMID: 30026507.

10. Hastings JM, Fazleabas AT. A baboon model for endometriosis: implications for fertility. Reprod Biol Endocrinol. 2006;4 Suppl 1:S7. Epub 2006/11/23. doi: 10.1186/1477-7827-4-S1-S7. PubMed PMID: 17118171; PubMed Central PMCID: PMCPMC1775067.

11. Jones CJ, Denton J, Fazleabas AT. Morphological and glycosylation changes associated with the endometrium and ectopic lesions in a baboon model of endometriosis. Hum Reprod. 2006;21(12):3068-80. Epub 2006/10/05. doi: 10.1093/humrep/del310. PubMed PMID: 17018533.

12. Harirchian P, Gashaw I, Lipskind ST, Braundmeier AG, Hastings JM, Olson MR, et al. Lesion
kinetics in a non-human primate model of endometriosis. Hum Reprod. 2012;27(8):2341-51.
Epub 2012/06/08. doi: 10.1093/humrep/des196. PubMed PMID: 22674203; PubMed Central
PMCID: PMCPMC3398680.

13. Greaves E, Cousins FL, Murray A, Esnal-Zufiaurre A, Fassbender A, Horne AW, et al. A novel
mouse model of endometriosis mimics human phenotype and reveals insights into the
inflammatory contribution of shed endometrium. Am J Pathol. 2014;184(7):1930-9. Epub
2014/06/10. doi: 10.1016/j.ajpath.2014.03.011. PubMed PMID: 24910298; PubMed Central
PMCID: PMCPMC4076466.

14. Dorning A, Dhami P, Panir K, Hogg C, Park E, Ferguson GD, et al. Bioluminescent imaging in
induced mouse models of endometriosis reveals differences in four model variations. Dis
Model Mech. 2021;14(8). Epub 2021/08/13. doi: 10.1242/dmm.049070. PubMed PMID:
34382636; PubMed Central PMCID: PMCPMC8419713.

15. Vigano P, Ottolina J, Bartiromo L, Bonavina G, Schimberni M, Villanacci R, et al. Cellular
Components Contributing to Fibrosis in Endometriosis: A Literature Review. J Minim Invasive
Gynecol. 2020;27(2):287-95. Epub 2019/12/01. doi: 10.1016/j.jmig.2019.11.011. PubMed
PMID: 31785417.

16. Vigano P, Candiani M, Monno A, Giacomini E, Vercellini P, Somigliana E. Time to redefine
endometriosis including its pro-fibrotic nature. Hum Reprod. 2018;33(3):347-52. Epub
2017/12/06. doi: 10.1093/humrep/dex354. PubMed PMID: 29206943.

17. Brosens I, Puttemans P, Benagiano G. Which cell defines endometriosis? Hum Reprod.
2018;33(3):357-60. Epub 2018/02/09. doi: 10.1093/humrep/dey016. PubMed PMID:
29420768.

18. Capobianco A, Rovere-Querini P. Endometriosis, a disease of the macrophage. Front
Immunol. 2013;4:9. Epub 2013/02/02. doi: 10.3389/fimmu.2013.00009. PubMed PMID:
23372570; PubMed Central PMCID: PMCPMC3556586.

19. Song Y, Burns GW, Joshi NR, Arora R, Kim JJ, Fazleabas AT. Spheroids as a model for
endometriotic lesions. JCI Insight. 2023;8(11). Epub 2023/04/27. doi:
10.1172/jci.insight.160815. PubMed PMID: 37104033.

20. Marla S, Mortlock S, Heinosalo T, Poutanen M, Montgomery GW, McKinnon BD. Gene
expression profiles separate endometriosis lesion subtypes and indicate a sensitivity of
endometrioma to estrogen suppressive treatments through elevated ESR2 expression. BMC
Med. 2023;21(1):460. Epub 2023/11/24. doi: 10.1186/s12916-023-03166-1. PubMed PMID:
37996888; PubMed Central PMCID: PMCPMC10666321.

21. Tan Y, Flynn WF, Sivajothi S, Luo D, Bozal SB, Davé M, et al. Single-cell analysis of
endometriosis reveals a coordinated transcriptional programme driving immunotolerance and
angiogenesis across eutopic and ectopic tissues. Nat Cell Biol. 2022. Epub 2022/07/22. doi:
10.1038/s41556-022-00961-5. PubMed PMID: 35864314.

22. Fonseca MAS, Haro M, Wright KN, Lin X, Abbasi F, Sun J, et al. Single-cell transcriptomic
analysis of endometriosis. Nat Genet. 2023;55(2):255-67. Epub 2023/01/10. doi:
10.1038/s41588-022-01254-1. PubMed PMID: 36624343.

23. Li R, Tran DN, Lessey BA, Young SL, Kim TH, Jeong JW. Transcriptomic changes in eutopic
endometrium and ectopic lesions during endometriosis progression in a mouse model. F S
Sci. 2024. Epub 2024/02/12. doi: 10.1016/j.xfss.2024.02.001. PubMed PMID: 38342342.

24. Zollinger DR, Lingle SE, Sorg K, Beechem JM, Merritt CR. GeoMx™ RNA Assay: High
Multiplex, Digital, Spatial Analysis of RNA in FFPE Tissue. Methods in molecular biology.
2020;2148:331-45. Epub 2020/05/13. doi: 10.1007/978-1-0716-0623-0_21. PubMed PMID:
32394392.

25. Blumhagen RZ, Kurche JS, Cool CD, Walts AD, Heinz D, Fingerlin TE, et al. Spatially distinct
molecular patterns of gene expression in idiopathic pulmonary fibrosis. Respir Res.
2023;24(1):287. Epub 2023/11/18. doi: 10.1186/s12931-023-02572-6. PubMed PMID:
37978501; PubMed Central PMCID: PMCPMC10655274.

26. Pennel KAF, Hatthakarnkul P, Wood CS, Lian GY, Al-Badran SSF, Quinn JA, et al. JAK/STAT3
represents a therapeutic target for colorectal cancer patients with stromal-rich tumors. J Exp
Clin Cancer Res. 2024;43(1):64. Epub 2024/03/01. doi: 10.1186/s13046-024-02958-4.
PubMed PMID: 38424636; PubMed Central PMCID: PMCPMC10905886.

27. Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular
Signatures Database (MSigDB) hallmark gene set collection. Cell Syst. 2015;1(6):417-25.
Epub 2016/01/16. doi: 10.1016/j.cels.2015.12.004. PubMed PMID: 26771021; PubMed
Central PMCID: PMCPMC4707969.

28. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. Proc Natl Acad Sci U S A. 2005;102(43):15545-50. Epub 2005/10/04. doi:
10.1073/pnas.0506580102. PubMed PMID: 16199517; PubMed Central PMCID:
PMCPMC1239896.

29. Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, et al. Inference and
analysis of cell-cell communication using CellChat. Nat Commun. 2021;12(1):1088. Epub

2021/02/19. doi: 10.1038/s41467-021-21246-9. PubMed PMID: 33597522; PubMed Central
PMCID: PMCPMC7889871.

30. Harada T, Iwabe T, Terakawa N. Role of cytokines in endometriosis. Fertility and Sterility.
2001;76(1):1-10. doi: https://doi.org/10.1016/S0015-0282(01)01816-7.

31. Noble LS, Simpson ER, Johns A, Bulun SE. Aromatase expression in endometriosis. J Clin
Endocrinol Metab. 1996;81(1):174-9. Epub 1996/01/01. doi: 10.1210/jcem.81.1.8550748.
PubMed PMID: 8550748.

32. Attia GR, Zeitoun K, Edwards D, Johns A, Carr BR, Bulun SE. Progesterone receptor isoform
A but not B is expressed in endometriosis. J Clin Endocrinol Metab. 2000;85(8):2897-902.
Epub 2000/08/18. doi: 10.1210/jcem.85.8.6739. PubMed PMID: 10946900.

33. Bulun SE, Yang S, Fang Z, Gurates B, Tamura M, Sebastian S. Estrogen production and
metabolism in endometriosis. Annals of the New York Academy of Sciences. 2002;955:75-85;
discussion 6-8, 396-406. Epub 2002/04/13. doi: 10.1111/j.1749-6632.2002.tb02767.x.
PubMed PMID: 11949967.

34. Fazleabas AT, Brudney A, Chai D, Langoi D, Bulun SE. Steroid receptor and aromatase
expression in baboon endometriotic lesions. Fertil Steril. 2003;80 Suppl 2:820-7. Epub
2003/09/25. doi: 10.1016/s0015-0282(03)00982-8. PubMed PMID: 14505759.

35. Attar E, Tokunaga H, Imir G, Yilmaz MB, Redwine D, Putman M, et al. Prostaglandin E2 via
steroidogenic factor-1 coordinately regulates transcription of steroidogenic genes necessary
for estrogen synthesis in endometriosis. J Clin Endocrinol Metab. 2009;94(2):623-31. Epub
2008/11/13. doi: 10.1210/jc.2008-1180. PubMed PMID: 19001523; PubMed Central PMCID:
PMCPMC2646521.

36. Yilmaz BD, Bulun SE. Endometriosis and nuclear receptors. Hum Reprod Update.
2019;25(4):473-85. doi: 10.1093/humupd/dmz005. PubMed Central PMCID:
PMCPMC6601390.

37. Ohlsson Teague EM, Van der Hoek KH, Van der Hoek MB, Perry N, Wagaarachchi P,
Robertson SA, et al. MicroRNA-regulated pathways associated with endometriosis. Mol
Endocrinol. 2009;23(2):265-75. Epub 2008/12/17. doi: 10.1210/me.2008-0387. PubMed
PMID: 19074548; PubMed Central PMCID: PMCPMC5419313.

38. Nothnick WB, Falcone T, Joshi N, Fazleabas AT, Graham A. Serum miR-451a Levels Are
Significantly Elevated in Women With Endometriosis and Recapitulated in Baboons (Papio
anubis) With Experimentally-Induced Disease. Reproductive sciences (Thousand Oaks,
Calif). 2017;24(8):1195-202. Epub 2016/12/07. doi: 10.1177/1933719116681519. PubMed
PMID: 27920341.

39. Rekker K, Tasa T, Saare M, Samuel K, Kadastik U, Karro H, et al. Differentially-Expressed
miRNAs in Ectopic Stromal Cells Contribute to Endometriosis Development: The Plausible
Role of miR-139-5p and miR-375. Int J Mol Sci. 2018;19(12). Epub 2018/11/30. doi:
10.3390/ijms19123789. PubMed PMID: 30487429; PubMed Central PMCID:
PMCPMC6321240.

40. Pavone ME, Grover AR, Confino R, Pearson EK, Malpani S, Cheng YH, et al. Retinoic acid
action is altered within endometrium of baboons affected with endometriosis. Journal of
Endometriosis and Pelvic Pain Disorders. 2022:22840265211062008. doi:
10.1177/22840265211062008. PubMed PMID: WOS:000740968400001.

41. Song Y, Su RW, Joshi NR, Kim TH, Lessey BA, Jeong JW, et al. Interleukin-6 (IL-6) Activates
the NOTCH1 Signaling Pathway Through E-Proteins in Endometriotic Lesions. J Clin
Endocrinol Metab. 2020;105(5). Epub 2020/03/03. doi: 10.1210/clinem/dgaa096. PubMed
PMID: 32119078; PubMed Central PMCID: PMCPMC7096313.

42. Kai K, Joshi NR, Burns GW, Hrbek SM, Vegter EL, Ochoa-Bernal MA, et al. MicroRNA-210-
3p Regulates Endometriotic Lesion Development by Targeting IGFBP3 in Baboons and
Women with Endometriosis. Reproductive sciences (Thousand Oaks, Calif).
2023;30(10):2932-44. Epub 2023/05/16. doi: 10.1007/s43032-023-01253-5. PubMed PMID:
37188982; PubMed Central PMCID: PMCPMC10556147.

43. Hogg C, Panir K, Dhami P, Rosser M, Mack M, Soong D, et al. Macrophages inhibit and
enhance endometriosis depending on their origin. Proc Natl Acad Sci U S A. 2021;118(6).
Epub 2021/02/05. doi: 10.1073/pnas.2013776118. PubMed PMID: 33536334.

44. Hogg C, Horne AW, Greaves E. Endometriosis-Associated Macrophages: Origin, Phenotype,
and Function. Front Endocrinol (Lausanne). 2020;11:7. Epub 2020/02/11. doi:
10.3389/fendo.2020.00007. PubMed PMID: 32038499; PubMed Central PMCID:
PMCPMC6989423.

45. Henlon Y, Panir K, McIntyre I, Hogg C, Dhami P, Cuff AO, et al. Single-cell analysis identifies
distinct macrophage phenotypes associated with pro-disease and pro-resolving functions in
the endometriotic niche. bioRxiv. 2024:2024.03.07.583861. doi: 10.1101/2024.03.07.583861.

46. Croasdell Lucchini A, Gachanja NN, Rossi AG, Dorward DA, Lucas CD. Epithelial Cells and
Inflammation in Pulmonary Wound Repair. Cells. 2021;10(2). Epub 2021/02/11. doi:
10.3390/cells10020339. PubMed PMID: 33562816; PubMed Central PMCID:
PMCPMC7914803.

47. Ren JL, Pan JS, Lu YP, Sun P, Han J. Inflammatory signaling and cellular senescence. Cell
Signal. 2009;21(3):378-83. Epub 2008/11/11. doi: 10.1016/j.cellsig.2008.10.011. PubMed
PMID: 18992324; PubMed Central PMCID: PMCPMC2870712.

48. Zou L, Meng L, Xu Y, Wang K, Zhang J. Revealing the diagnostic value and immune infiltration
of senescence-related genes in endometriosis: a combined single-cell and machine learning
analysis. Front Pharmacol. 2023;14:1259467. Epub 2023/10/20. doi:
10.3389/fphar.2023.1259467. PubMed PMID: 37860112; PubMed Central PMCID:
PMCPMC10583561.

49. Muter J, Kong CS, Brosens JJ. The Role of Decidual Subpopulations in Implantation,
Menstruation and Miscarriage. Front Reprod Health. 2021;3:804921. Epub 2022/10/29. doi:
10.3389/frph.2021.804921. PubMed PMID: 36303960; PubMed Central PMCID:
PMCPMC9580781.

50. Crocker PR, Jefferies WA, Clark SJ, Chung LP, Gordon S. Species heterogeneity in
macrophage expression of the CD4 antigen. J Exp Med. 1987;166(2):613-8. Epub
1987/08/01. doi: 10.1084/jem.166.2.613. PubMed PMID: 3110358; PubMed Central PMCID:
PMCPMC2189597.

51. Zhen A, Krutzik SR, Levin BR, Kasparian S, Zack JA, Kitchen SG. CD4 ligation on human
blood monocytes triggers macrophage differentiation and enhances HIV infection. J Virol.
2014;88(17):9934-46. Epub 2014/06/20. doi: 10.1128/jvi.00616-14. PubMed PMID:
24942581; PubMed Central PMCID: PMCPMC4136363.

52. Stevenson M. HIV-1 pathogenesis. Nat Med. 2003;9(7):853-60. Epub 2003/07/02. doi:
10.1038/nm0703-853. PubMed PMID: 12835705.

53. Betjes MG, Haks MC, Tuk CW, Beelen RH. Monoclonal antibody EBM11 (anti-CD68)
discriminates between dendritic cells and macrophages after short-term culture.
Immunobiology. 1991;183(1-2):79-87. Epub 1991/09/01. doi: 10.1016/s0171-2985(11)80187-
7. PubMed PMID: 1834546.

54. Yan J, Zhou L, Liu M, Zhu H, Zhang X, Cai E, et al. Single-cell analysis reveals insights into
epithelial abnormalities in ovarian endometriosis. Cell Rep. 2024;43(3):113716. Epub
2024/02/27. doi: 10.1016/j.celrep.2024.113716. PubMed PMID: 38412094.

55. Calandra T, Roger T. Macrophage migration inhibitory factor: a regulator of innate immunity.
Nat Rev Immunol. 2003;3(10):791-800. Epub 2003/09/23. doi: 10.1038/nri1200. PubMed
PMID: 14502271; PubMed Central PMCID: PMCPMC7097468.

56. Mitchell RA, Metz CN, Peng T, Bucala R. Sustained mitogen-activated protein kinase (MAPK)
and cytoplasmic phospholipase A2 activation by macrophage migration inhibitory factor (MIF).

Regulatory role in cell proliferation and glucocorticoid action. J Biol Chem.
1999;274(25):18100-6. Epub 1999/06/11. doi: 10.1074/jbc.274.25.18100. PubMed PMID:
10364264.

57. Schenkein HA, Rutherford B. C3-mediated release of prostaglandin from human monocytes.
Behaviour in short-term culture. Immunology. 1984;51(1):83-91. Epub 1984/01/01. PubMed
PMID: 6690398; PubMed Central PMCID: PMCPMC1454415.

58. Calandra T, Bernhagen J, Mitchell RA, Bucala R. The macrophage is an important and
previously unrecognized source of macrophage migration inhibitory factor. J Exp Med.
1994;179(6):1895-902. Epub 1994/06/01. doi: 10.1084/jem.179.6.1895. PubMed PMID:
8195715; PubMed Central PMCID: PMCPMC2191507.

59. Bacher M, Metz CN, Calandra T, Mayer K, Chesney J, Lohoff M, et al. An essential regulatory
role for macrophage migration inhibitory factor in T-cell activation. Proc Natl Acad Sci U S A.
1996;93(15):7849-54. Epub 1996/07/23. doi: 10.1073/pnas.93.15.7849. PubMed PMID:
8755565; PubMed Central PMCID: PMCPMC38837.

60. Weed JC, Arquembourg PC. Endometriosis: can it produce an autoimmune response
resulting in infertility? Clin Obstet Gynecol. 1980;23(3):885-93. Epub 1980/09/01. PubMed
PMID: 7418287.

61. Isaacson KB, Coutifaris C, Garcia CR, Lyttle CR. Production and secretion of complement
component 3 by endometriotic tissue. J Clin Endocrinol Metab. 1989;69(5):1003-9. Epub
1989/11/01. doi: 10.1210/jcem-69-5-1003. PubMed PMID: 2793987.

62. Tao XJ, Sayegh RA, Isaacson KB. Increased expression of complement component 3 in
human ectopic endometrium compared with the matched eutopic endometrium. Fertil Steril.
1997;68(3):460-7. Epub 1997/10/07. doi: 10.1016/s0015-0282(97)00254-9. PubMed PMID:
9314915.

63. Hasan A, Rahim A, Afzal M, Naveed AK, Ayub S, Jahan S. Serum Albumin and C3
Complement Levels in Endometriosis. J Coll Physicians Surg Pak. 2019;29(8):702-5. Epub
2019/07/31. doi: 10.29271/jcpsp.2019.08.702. PubMed PMID: 31358086.

64. Young SL. The complement system and its role in the endometrium. In: Aplin JD, Fazleabas
AT, Glasser SR, Giudice LC, editors. The Endometrium: Molecular, Cellular and Clinical
Perspectives, Second Edition: Taylor & Francis; 2008. p. 568-78.

65. Agostinis C, Balduit A, Mangogna A, Zito G, Romano F, Ricci G, et al. Immunological Basis of
the Endometriosis: The Complement System as a Potential Therapeutic Target. Front
Immunol. 2020;11:599117. Epub 2021/01/29. doi: 10.3389/fimmu.2020.599117. PubMed
PMID: 33505394; PubMed Central PMCID: PMCPMC7829336.

66. Brown EJ, Frazier WA. Integrin-associated protein (CD47) and its ligands. Trends Cell Biol.
2001;11(3):130-5. Epub 2001/04/18. doi: 10.1016/s0962-8924(00)01906-1. PubMed PMID:
11306274.

67. Gresham HD, Dale BM, Potter JW, Chang PW, Vines CM, Lowell CA, et al. Negative
regulation of phagocytosis in murine macrophages by the Src kinase family member, Fgr. J
Exp Med. 2000;191(3):515-28. Epub 2000/02/09. doi: 10.1084/jem.191.3.515. PubMed
PMID: 10662797; PubMed Central PMCID: PMCPMC2195814.

68. Meri S, Morgan BP, Davies A, Daniels RH, Olavesen MG, Waldmann H, et al. Human protectin
(CD59), an 18,000-20,000 MW complement lysis restricting factor, inhibits C5b-8 catalysed
insertion of C9 into lipid bilayers. Immunology. 1990;71(1):1-9. Epub 1990/09/01. PubMed
PMID: 1698710; PubMed Central PMCID: PMCPMC1384213.

69. Yang H, Li L, Liu X, Zhao Y. High Expression of the Component 3a Receptor 1 (C3AR1) Gene
in Stomach Adenocarcinomas Infers a Poor Prognosis and High Immune-Infiltration Levels.
Med Sci Monit. 2021;27:e927977. Epub 2021/02/05. doi: 10.12659/msm.927977. PubMed
PMID: 33539329; PubMed Central PMCID: PMCPMC7871482.

70. Zhu S, Wang A, Xu W, Hu L, Sun J, Wang X. The heterogeneity of fibrosis and angiogenesis
in endometriosis revealed by single-cell RNA-sequencing. Biochim Biophys Acta Mol Basis
Dis. 2023;1869(2):166602. Epub 2022/11/19. doi: 10.1016/j.bbadis.2022.166602. PubMed
PMID: 36400338.

71. Sunderkötter C, Steinbrink K, Goebeler M, Bhardwaj R, Sorg C. Macrophages and
angiogenesis. J Leukoc Biol. 1994;55(3):410-22. Epub 1994/03/01. doi: 10.1002/jlb.55.3.410.
PubMed PMID: 7509844.

72. Martínez-Ramos S, Rafael-Vidal C, Malvar-Fernández B, Pérez N, Mouriño C, Pérez SG, et
al. Semaphorin3B promotes an anti-inflammatory and pro-resolving phenotype in
macrophages from rheumatoid arthritis patients in a MerTK-dependent manner. Front
Immunol. 2023;14:1268144. Epub 2024/01/29. doi: 10.3389/fimmu.2023.1268144. PubMed
PMID: 38283352; PubMed Central PMCID: PMCPMC10811190.

73. Bulun SE, Fang Z, Imir G, Gurates B, Tamura M, Yilmaz B, et al. Aromatase and
endometriosis. Semin Reprod Med. 2004;22(1):45-50. Epub 2004/04/15. doi: 10.1055/s-
2004-823026. PubMed PMID: 15083380.

74. Ullah K, Rahman TU, Pan HT, Guo MX, Dong XY, Liu J, et al. Serum estradiol levels in
controlled ovarian stimulation directly affect the endometrium. J Mol Endocrinol.
2017;59(2):105-19. Epub 2017/05/26. doi: 10.1530/jme-17-0036. PubMed PMID: 28539318;
PubMed Central PMCID: PMCPMC5510595.

75. Agostinis C, Zorzet S, Balduit A, Zito G, Mangogna A, Macor P, et al. The Inflammatory Feed-
Forward Loop Triggered by the Complement Component C3 as a Potential Target in
Endometriosis. Front Immunol. 2021;12:693118. Epub 2021/09/08. doi:
10.3389/fimmu.2021.693118. PubMed PMID: 34489939; PubMed Central PMCID:
PMCPMC8418148.

76. Ruiz LA, Baez-Vega PM, Ruiz A, Peterse DP, Monteiro JB, Bracero N, et al. Dysregulation of
Lysyl Oxidase Expression in Lesions and Endometrium of Women With Endometriosis.
Reproductive sciences (Thousand Oaks, Calif). 2015;22(12):1496-508. Epub 2015/05/13. doi:
10.1177/1933719115585144. PubMed PMID: 25963914; PubMed Central PMCID:
PMCPMC5933196.

77. Colon-Diaz M, Baez-Vega P, Garcia M, Ruiz A, Monteiro JB, Fourquet J, et al. HDAC1 and
HDAC2 are differentially expressed in endometriosis. Reproductive sciences (Thousand
Oaks, Calif). 2012;19(5):483-92. Epub 2012/02/22. doi: 10.1177/1933719111432870.
PubMed PMID: 22344732; PubMed Central PMCID: PMCPMC3343094 research, authorship,
and/or publication of this article.

78. Colón-Caraballo M, García M, Mendoza A, Flores I. Human Endometriosis Tissue Microarray
Reveals Site-specific Expression of Estrogen Receptors, Progesterone Receptor, and Ki67.
Appl Immunohistochem Mol Morphol. 2019;27(7):491-500. Epub 2018/04/10. doi:
10.1097/pai.0000000000000663. PubMed PMID: 29629944; PubMed Central PMCID:
PMCPMC6174114.

79. Noyes RW, Hertig AT, Rock J. Dating the endometrial biopsy. American journal of obstetrics
and gynecology. 1975;122(2):262-3. Epub 1975/05/01. doi: 10.1016/s0002-9378(16)33500-
1. PubMed PMID: 1155504.

80. McInnes L, Healy J, Melville J. Umap: Uniform manifold approximation and projection for
dimension reduction. arXiv preprint arXiv:180203426. 2018.

81. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential
expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.
2015;43(7):e47. Epub 2015/01/22. doi: 10.1093/nar/gkv007. PubMed PMID: 25605792;
PubMed Central PMCID: PMCPMC4402510.

82. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.
Nat Methods. 2012;9(7):671-5. Epub 2012/08/30. doi: 10.1038/nmeth.2089. PubMed PMID:
22930834; PubMed Central PMCID: PMCPMC5554542.

83. Team R. RStudio: Integrated Development for R. Boston, MA: RStudio, PBC.; 2020. Available
from: http://www.rstudio.com/.

84. McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor RNA-Seq

experiments with respect to biological variation. Nucleic Acids Res. 2012;40(10):4288-97.

Epub 2012/01/31. doi: 10.1093/nar/gks042. PubMed PMID: 22287627; PubMed Central

PMCID: PMCPMC3378882.

85. Zhou X, Lindsay H, Robinson MD. Robustly detecting differential expression in RNA

sequencing data using observation weights. Nucleic Acids Res. 2014;42(11):e91. Epub

2014/04/23. doi: 10.1093/nar/gku310. PubMed PMID: 24753412; PubMed Central PMCID:

PMCPMC4066750.

86. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment tool

for interpreting omics data. Innovation (Camb). 2021;2(3):100141. Epub 2021/09/25. doi:

10.1016/j.xinn.2021.100141. PubMed PMID: 34557778; PubMed Central PMCID:

PMCPMC8454663.