Bacterial defenses and their trade-off with growth are not

ubiquitous but depend on ecological contexts

Zhi-Ling Liu[1,2], Jia Liu[1], Deng-Ke Niu[1,2*]

[1]MOE Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life Sciences,

Beijing Normal University, Beijing, 100875, China

[2]Beijing Key Laboratory of Gene Resource and Molecular Development, College of Life Sciences,

Beijing Normal University, Beijing, 100875, China

*Correspondence author:

Deng-Ke Niu, College of Life Sciences, Beijing Normal University, Beijing, 100875, China.

dkniu@bnu.edu.cn, dengkeniu@hotmail.com

ORCID:

Zhi-Ling Liu, https://orcid.org/0000-0001-6579-6642

Deng-Ke Niu, https://orcid.org/0000-0003-4503-2658

Competing interests

23 The authors declare no competing interests.

Abstract

Bacteriophages, also known as bacterial viruses, exert significant influence on microbial ecosystems due to their widespread presence. In response to the threat of phage predation, bacteria have developed an extensive array of antiviral defense mechanisms. This study investigates the intricate relationship between bacterial defense mechanisms and growth rates across various ecological contexts. First, we found that bacteria lacking defense systems show prolonged doubling times. By integrating phylogenetic eigenvectors into the ecological feature matrix, we employed a phylogenetic random forest model to determine the most influential ecological features on the presence and abundance of bacterial defenses. Further phylogenetic regressions unveiled nuanced dependencies of bacterial defenses on specific environmental factors. Contrary to conventional assumptions, our findings challenge the idea of a universal distribution of bacterial defense systems, instead highlighting their reliance on subtle ecological characteristics. Particularly noteworthy are observations that symbiotic and endosymbiotic bacteria often exhibit reduced defense systems and negative correlations between minimal doubling time and defense system abundance. Conversely, as per conventional assumptions, free-living and motile bacteria displayed significant positive correlations between minimal doubling time and defense system abundance. Overall, our results suggest that the trade-off between growth and defense strategies is not universally applicable but rather contingent on context. Moreover, our study illuminates the critical role of ecological variables such as habitat specificity, motility, and nutrient availability in shaping bacterial growth rates and defense mechanisms. These findings underscore the complexity of microbial interactions and stress the importance of considering the ecological context in understanding bacterial defense strategies.

Keywords

Defense system, minimal doubling time, trade-off, trade-up, symbiosis, endosymbiosis, gut microbiome.

Introduction

Bacterial viruses, also known as phages or bacteriophages, represent the most abundant biological entities on Earth. They significantly outnumber bacterial cells across various samples and microbiomes, with ratios reaching up to ~15-25 fold, as observed in freshwater plankton (López-García et al. 2023; Roux and Brum 2023). With few exceptions, such as intracellular parasitic and symbiotic bacteria, the majority of bacterial species fall prey to phages (Koonin et al. 2017). In environments like seawater, this constant predation results in a substantial daily death rate of bacterioplankton, estimated between 25-40% (Wommack and Colwell 2000; Suttle 2007). To combat phage attacks, bacteria have evolved an arsenal of defense tactics targeting every stage of the phage's life cycle. These tactics include diverse mechanistic approaches such as the restriction-modification (RM) system, the Clustered Regularly Interspaced Short Palindromic Repeat and associated protein (CRISPR-Cas) system, and the Abortive Infection (Abi) system (Stern and Sorek 2011; Koonin et al. 2017; Bernheim and Sorek 2020), collectively referred to as antiviral defense systems.

Despite the potential for horizontal gene transfer, bacteria typically only possess a subset of the available anti-phage mechanisms (Rostøl and Marraffini 2019). For instance, the CRISPR-Cas systems, which provide adaptive immunity by targeting previously encountered pathogens, are found in approximately 40% of bacteria (Makarova et al. 2020). Similarly, Abi systems are sporadically distributed across bacterial phylogeny (Stern and Sorek 2011), while RM systems are widespread but absent in about 4% of bacterial genomes (Roberts et al. 2015). Additionally, the uneven distribution of antiviral defense systems extends to the archaeal lineage (Roberts et al. 2015; Makarova et al. 2020), reflecting trade-offs between the costs of maintaining such systems and the benefits gained in phage resistance (Winter et al. 2010; Ferenci 2016; Pančić and Kiørboe 2018; Rostøl and Marraffini 2019; Thingstad 2022; Liu et al. 2023).

In biological systems, traits often exhibit trade-offs due to constraints on resources such as energy, habitat, or time (Roff and Fairbairn 2007; Garland Jr 2014; Garland et al. 2022). For example, energy allocated for growth cannot simultaneously be used for defense or other energy-intensive processes, leading to necessary compromises among traits. While the concept of growth-defense

trade-offs is well-established in plants and animals, its relevance in virus-infected prokaryotes is less

recognized (Herms and Mattson 1992; Ferenci 2016).

Laboratory research on phage resistance has extensively explored trade-offs using model

bacteria and phages such as *Escherichia coli* and *Pseudomonas aeruginosa*. Phage resistance

evolution significantly impacts bacterial fitness components, including slowed cell division rates,

reduced nutrient uptake, decreased virulence, and impaired cell motility (Lenski 1988; Bohannan and

Lenski 2000; Bohannan et al. 2002; Buckling et al. 2005; Lennon et al. 2007; Jessup and Bohannan

2008; Quance and Travisano 2009; Hall et al. 2012; Ferenci 2016; Burmeister and Turner 2020;

Castledine et al. 2022). Notably, some phage-resistant strains exhibit trade-ups, where selection

improves multiple fitness components simultaneously, including antibiotic resistance (Moulton-

Brown and Friman 2018; Burmeister et al. 2020).

However, resistance to phages does not consistently correlate negatively with bacterial

growth rates in natural environments. Studies examining marine *Synechococcus* strains and phage

isolates, as well as bacteria and phages from horse chestnut leaves, have shown conflicting results

regarding the growth-defense trade-off (Lennon et al. 2007; Koskella et al. 2011). Recent research on

marine bacterial communities has observed a negative correlation between growth rates and resistance

to protists and viruses, supporting the growth-defense trade-off hypothesis (Yang et al. 2023).

To gain a comprehensive understanding of the trade-off between prokaryotic growth and

defense, investigations encompassing diverse environments and species are essential. Recent research

analyzing growth rates and adaptive immunity across bacterial domains revealed a significant positive

correlation, indicating evolutionary trade-offs between growth and adaptive defense against virulent

phages (Liu et al., 2023). Additionally, besides CRISPR-Cas systems, prokaryotes possess numerous

other antiviral mechanisms, such as the RM and Abi systems, warranting further exploration (Koonin

et al., 2017; Bernheim and Sorek, 2020; Georjon and Bernheim, 2023; Tesson and Bernheim, 2023).

Understanding the relationship between growth and antiviral defense across different environments

and prokaryotic lineages holds promise for advancing our knowledge of evolution and informing the

development of antimicrobial strategies.

To comprehend the trade-off between prokaryotic growth and defense, it's imperative to adopt

a holistic perspective. Despite occasional inconsistencies, our aim is to attain a global understanding

of whether phage-resistant lineages generally exhibit slower growth rates compared to phage-sensitive

ones across bacterial domains. Such insights could profoundly influence our comprehension of

evolutionary processes and facilitate the development of novel antimicrobial strategies.

Recently, we conducted an investigation into the relationship between growth rate and

adaptive immunity in 4142 bacteria, utilizing predicted minimal doubling times retrieved from the

EGGO database and CRISPR-Cas systems annotated with CRISPRCasFinder (Couvin et al. 2018;

Weissman et al. 2021; Liu et al. 2023). Our findings unveiled a significant positive correlation,

indicating evolutionary trade-offs between growth and adaptive defense against virulent phages

within the bacterial domain.

In addition to CRISPR-Cas systems, prokaryotes have evolved various other antiviral

mechanisms, including the RM system, the Abi system, and numerous others that warrant further

exploration (Koonin et al. 2017; Bernheim and Sorek 2020; Georjon and Bernheim 2023; Tesson and

Bernheim 2023). Notably, certain prokaryotes, such as the delta proteobacteria *Desulfonema limicola*,

can harbor over 50 defense systems in their genomes (Tesson et al. 2022). Recently, Tesson et al.

(2022) developed DefenseFinder, a novel antivirus-system detection program based on the HMM

mapping algorithm, which enables the identification of known antiviral systems across 21,000 fully

sequenced prokaryotic genomes. Leveraging this tool, we explored the relationship between growth

and antiviral defense across bacterial domains, as well as within specific environments.

Materials and Methods

The minimal doubling times of bacteria and archaea were extracted from the EGGO database

(accessed on 2021-11-26), encompassing over 200,000 predictions of estimated growth rates

(Weissman et al. 2021). These values underwent a log transformation using the natural logarithm as

the base. Due to its smaller sample size, we opted not to utilize the dataset of prokaryotic empirical

minimal doubling times provided by Madin et al. (2020) in this study. While empirical data typically

offer higher accuracy compared to computationally generated predictions, predictive methodologies

have their own advantages (Liu et al. 2023). Empirical minimal doubling periods may offer superior

advantages over computationally generated predictions due to the inherent limitations of predictive

methodologies achieving 100% precision. However, the wide spectrum of genetic diversity observed

among strains of individual bacterial species necessitates empirical data encompassing various factors,

all procured from equivalent strains, a requirement that is not consistently met. Predicted phenotype

values, including defense systems and growth rate, for a strain originate from the genome sequences

of that very strain, potentially circumventing faults stemming from polymorphism within each species.

The antiviral defense systems of over 21,000 complete microbial genomes were sourced from

Tesson et al. (2022). Phylogenetic relationships among the analyzed prokaryotes were obtained from

the Genome Taxonomy Database (GTDB) (Parks et al., 2022) (accessed 2022-04-08). GTDB offers

superior taxonomic resolution and more precise classification compared to traditional phylogenetic

trees by leveraging genomic data.

The intersecting samples among the three aforementioned datasets consisted of 3532 bacterial

genomes and 237 archaeal genomes, some of which belonged to the same species. However, upon

intersecting with subsequent datasets, only a minimal number of archaeal species were acquired,

rendering most analyses impractical. Consequently, this paper focuses solely on the study of bacteria.

Ecological characteristics of prokaryotes were obtained from the ProTraits database (Brbić et

al. 2016), accessed on October 2, 2022, via "ProTraits_precisionScores.txt" available at

http://protraits.irb.hr/data.html. This comprehensive database integrates data from various repositories

such as GOLD, NCBI, and KEGG, utilizing automated text mining and comparative genomic

techniques to associate microorganisms with phenotypes. Our study focuses on ecological features

potentially influencing defense system distribution, excluding physiological traits like glycerol and

glutamate. Additionally, features with ambiguous biological significance, automated physiological

features from machine learning, and those with missing P values exceeding 10% of the overall sample

were discarded, retaining 125 prominent ecological features for analysis. Each ecological feature in

the database is represented by two columns, providing precision scores for both the presence and

absence of the feature, respectively. To facilitate analysis, these scores were converted into an overall

maximal confidence score ranging from 0 to 1, following the methodology of Weissman et al. (2019).

Missing values in the ProTraits database were imputed using phylogenetic information (Debastiani et

al. 2021). The PVRdecomp function of the R package PVR (Version 0.3) (Santos 2018) was

employed to extract eigenvectors from the phylogenetic distance matrix. Subsequently, the missForest

function of the R package missForest (Version 1.5) (Stekhoven and Bühlmann 2011) was utilized to

estimate missing values.

Random forest, an integrated algorithm of decision trees, was utilized for regression tasks in

this study. This method optimizes regression tasks, effectively handles high dimensionality, mitigates

overfitting risks, accommodates both categorical and numerical variables, and consistently delivers

reliable outcomes, making it superior to alternative techniques. The eigenvectors extracted from the

phylogenetic distance matrix were incorporated into the ecological feature matrix to control

phylogenetic relationships within the random forest analysis. The randomForest package (version 4.7-

1.1) (Liaw and Wiener 2002) was implemented in R (version 4.2.0).

Phylogenetic generalized least-squares (PGLS) analysis of correlations was conducted using

the R package phylolm (version 2.6.2) (Ho and Ane 2014). The Pagel λ model was employed due to

its generally lower AIC values in our dataset. Differences in minimal doubling times between the two

groups were compared using the phylANOVA function (Garland et al. 1993), while phylogenetic

logistic regression analysis with dichotomous variables as dependent variables was performed using

the phyloglm function (Ives and Garland 2010). When multiple correlation analyses were performed

on the same data, *P* values were corrected using the Benjamini & Hochberg method via the p.adjust

function in R (Version 4.0.2).

Results

Phylogenetic Signals in Analyzed Features and Their Implications for Comparative

Analysis

Due to shared ancestry among bacterial samples, feature values may exhibit non-independence,

violating fundamental assumptions of conventional statistical methods. As such, using phylogenetic

comparative methods becomes imperative to mitigate these effects (Felsenstein 1985; Symonds and

Blomberg 2014). We evaluated the phylogenetic signals of all analyzed features and observed

significant signals in nearly all cases, with minimal doubling time demonstrating the strongest signal

(Supplementary Table 1). Consequently, we will employ phylogenetic comparative methods

whenever applicable.

A recent study by Chen et al. (2023) highlighted the potential for conflicting outcomes in

PGLS analysis when interchanging dependent and independent variables. While PGLS cannot

establish causality but rather identifies correlations, Chen et al. utilized simulated data and a gold

standard to establish criteria for selecting the dependent variable. Their findings suggested that

choosing the feature with a stronger phylogenetic signal as the dependent variable led to more

accurate correlation detection than randomly selected variables. In line with this approach, we

adopted the feature exhibiting a stronger phylogenetic signal as the dependent variable in all PGLS

analyses conducted in this study.

Ecological Factors Influencing the Presence of Antiviral Defense Systems in Bacteria

Our analysis stems from a comprehensive program capable of detecting all known antiviral systems in

prokaryotic genomes (Tesson et al. 2022). However, among the 3532 bacterial genomes examined, 7%

(243) were found to lack any defense systems. Phylogenetic ANOVA analysis unveiled that these

bacteria exhibited significantly longer minimal doubling times compared to those encoding one or

more defense systems (Mean $\pm$ SD: 5.6 $\pm$ 3.1 h vs. 3.2 $\pm$ 4.4 h. P = 0.0011). Surprisingly, bacteria

devoid of defense systems did not demonstrate faster proliferation rates, contradicting the anticipated

trade-off between growth and defense. It is plausible that these bacteria inhabit specific ecological

niches where neither defense mechanisms nor rapid growth are necessary.

To delve into the factors influencing the presence or absence of bacterial defense systems, we

leveraged ecological data from the ProTraits database. Since the database primarily comprises

bacterial species rather than individual strains or genomes, we aggregated some of the 3532 genomes

into species. To capture species-level characteristics, we computed median values of defense system

abundance and minimal doubling time for genomes within the same species, identifying 1044 distinct

bacterial species.

Utilizing a random forest model, we ranked the importance of 125 ecological features.

Remarkably, factors such as "known habitats = insect endosymbiont," "known habitats =

endosymbiont," "biotic relationship = symbiont," and "host = insects general" emerged as top

influencers, underscoring the significance of symbiotic and endosymbiotic relationships (Fig. 1).

Given that the random forest model cannot discern the directional effect of features on the

presence or absence of antiviral defense systems, we conducted phylogenetic logistic regression

(Table 1). Results indicated a negative correlation between antiviral defense presence and "known

habitats = insect endosymbiont," "biotic relationship = symbiont," and "host = insects general" ($P <$

10^{-8} for all), while "known habitats = endosymbiont" showed no significant correlation (P = 0.452).

Ecological Determinants of Defense System Abundance: Insights from Symbiosis and

Environmental Dynamics

The preceding study explored factors influencing the presence or absence of defense systems. Here,

we delve deeper into the variables shaping defense system abundance. Bacteria lacking defense

mechanisms will be recorded as having zero systems, ensuring inclusivity in our analysis.

Among 125 ecological features, "known habitats = insect endosymbiont," "known habitats =

endosymbiont," and "biotic relationship = symbiont" emerged as top influencers, underscoring the

importance of symbiotic associations (Fig. 2). Furthermore, symbiotic relationships with hosts were

supported by additional features within the top rankings, including "ecosystem type = circulatory

system," "ecosystem subtype = blood," "habitat = host associated," "host = insects general,"

"ecosystem category = mammals," "known habitats = host," and "ecosystem = host associated." If

symbiotic bacteria tend to have fewer defense systems for certain reasons, free-living bacteria should

exhibit a comparatively higher abundance of defense systems. Indeed, "habitat = free living" ranks

within the top 19.

Additionally, aquatic environments and energy sources emerged as key factors. "Habitat =

aquatic" and "energy source = photoautotroph" ranked prominently, indicating their influence on

defense system abundance (Fig. 2). Additionally, polyhabitat, filamentous shape, mobility, and

bioremediation also appear to influence defense system abundance significantly.

Further PGLS analysis revealed negative correlations between symbiotic characteristics and

defense system abundance, while free-living bacteria exhibited a positive association (Table 2). While

some correlations may not achieve statistical significance, all other ecological characteristics

positively correlate with the abundance of defense systems (Table 2). Notably, bacteria inhabiting

multiple habitats, exhibiting motility, or being involved in environmental bioremediation tend to

possess a higher abundance of defense systems. Given their exposure to complex or fluctuating

environments, these bacteria are more susceptible to encountering invasive elements such as phages.

Another intriguing finding is that filamentous bacteria harbor significantly more defense systems,

likely due to their filamentous morphology providing increased sites for phage attachment.

Ecological Factors Impacting Bacterial Growth Rates: Insights from Endosymbiosis and

Habitat Dynamics

We investigate whether ecological factors influencing defense system abundance affect bacterial

growth rates. Using PGLS regression, we correlated minimal doubling time with overall maximal

confidence scores for significant ecological features.

Endosymbiosis and symbiosis with insects differ in their effects on bacterial growth rates

compared to symbiosis with mammals (Table 3). "Known habitats = insect endosymbiont," "Known

habitats = endosymbiont," and "Host = insects general" exhibit significantly positive correlations with

minimal doubling time. Conversely, features related to symbiosis with mammals show negative

correlations, albeit not statistically significant. Ambiguous features like "Biotic relationship =

symbiotic" show a significant positive correlation, while others like "Ecosystem = host associated,"

"Known habitats = host," and "Habitat = host associated" exhibit negative correlations.

Aquatic habitat and energy source features positively correlate with minimal doubling time

(Table 3), suggesting slower growth rates in bacteria residing in aquatic environments and utilizing

natural energy sources. Additionally, filamentous morphology correlates with slower growth rates,

while bacteria inhabiting multiple habitats tend to grow faster (Table 3).

Exploring Growth-Defense Relationships Across Ecological Niches

In our analysis utilizing the ProTraits database, each species received an overall maximal confidence score indicating the likelihood of possessing specific ecological features. We categorized species into presence or absence groups based on a threshold of 0.5, compiling species counts for the top 30 influential ecological features affecting defense system abundance. Features with fewer than 20 species were filtered out.

To investigate potential growth-defense trade-offs within these ecological niches, we conducted PGLS regression analysis, correlating bacterial minimal doubling time with defense system abundance (Table 4). Highly significant correlations ($P < 0.01$) were primarily observed in two niches: "Habitat = free living" and "Mobility," with extensive sample sizes comprising hundreds of bacterial species.

Among the features related to symbiosis, significant correlations between growth and defense were identified in three features, each representing samples of over 400 species: "Ecosystem = host associated," "Known habitats = host," and "Habitat = host associated." Accounting for phylogenetic data, closely related lineages with similar characteristics were treated as nearly one effective sample. Thus, the effective sample size in phylogenetic comparative analysis was substantially lower than the total number of analyzed lineages. Consequently, the conventional statistical rule of thumb regarding small and large samples ($n = 30$) should not be applied in phylogenetic comparative studies (Hu et al. 2022).

Nevertheless, the findings presented in Table 4 suggest the presence of not only a trade-off but also a trade-up between bacterial growth and defense.

Contextual Dependency in Growth-Defense Correlations Across Bacterial Environments

In Table 4, diverse correlations emerge between bacterial growth and defense across different environments. Positive correlations are observed in some bacteria within specific habitats, while

negative correlations are apparent in others; meanwhile, no correlation is discernible in some

instances. Consequently, within a large sample, the prevalence of each bacterial type dictates the

observed correlation pattern: a predominance of bacteria exhibiting positive correlations yields a

positive overall correlation, a balance of the three types results in no discernible correlation, and a

prevalence of bacteria with negative correlations leads to an overall negative correlation. Thus, the

significance of analyzing growth-defense correlations on a global scale diminishes.

Nevertheless, we conducted PGLS regression analysis to explore the growth-defense

correlation in our dataset of 3532 samples, yielding statistically insignificant results ($P = 0.398$).

Despite the lack of statistical significance, our analysis underscores the complex and context-

dependent nature of the relationship between bacterial growth rates and defense mechanisms across

diverse ecological contexts.

Discussion

The analysis of bacterial defense mechanisms, whether assessed by the presence or absence of

defense systems, or their abundance, yielded notable findings: ecological factors associated with

symbiosis and endosymbiosis emerged as the primary influencers of bacterial defenses (Figs. 1, 2).

Furthermore, phylogenetic regression analyses demonstrated a trend wherein symbiotic or

endosymbiotic bacteria tend to exhibit fewer defense systems or lack them altogether (Table 1, 2).

This observation aligns with prior findings noting the scarcity of defense systems among insect

endosymbionts (Makarova et al. 2013; Koonin et al. 2017). Our study unveils statistically significant

associations between symbiosis/endosymbiosis and defense mechanisms, suggesting that such

bacteria have reduced the necessity for robust defensive strategies. Endosymbiotic bacteria, residing

within host cells, enjoy physical insulation from aggressive phages. Similarly, evidence suggests that

many symbiotic bacteria are spatially separated from such phages. Notably, the gut microbiome

represents a well-studied example of symbiotic bacteria, comprising the majority of human symbionts

(Sender et al. 2016a; Sender et al. 2016b). Within the mammalian small intestine and colon,

specialized epithelial cells secrete a mucosal layer (Donaldson et al. 2016), which serves as a

formidable barrier against the spread and access of phage populations, providing a refuge for bacteria

(Guerin and Hill 2020; Lourenço et al. 2020; Shkoporov et al. 2022). Consequently, the phage-

bacterial ratios in the human gut are significantly lower compared to other ecosystems, often ranging

from 1:1 to 0.1:1 as opposed to 10:1(Shkoporov and Hill 2019).

However, the impact of endosymbiosis on bacterial growth rates in insects differs

significantly from symbiosis in mammals (Table 3). The proliferation of endosymbionts is not solely

determined by their inherent ability to replicate but is also influenced by regulatory constraints

imposed by their hosts (Whittle et al. 2021; Whittle et al. 2023). In contrast, the gut presents a highly

complex environment for bacteria. Nutrient levels fluctuate considerably on a daily basis, and the

presence of antimicrobial molecules and oxygen secreted by the epithelium strongly restricts potential

microbial inhabitants (Donaldson et al. 2016), creating a gradient of suitability for bacterial growth

from the lumen to the mucosa. Bacteria residing in isolated microhabitats such as crypts may be

tightly controlled by the host epithelium, while those dispersed into the gut lumen could proliferate in

response to nutrient availability. We hypothesize that the overall impact of gut symbiosis on bacterial

growth rate depends on the distribution of bacteria across different microhabitats.

Due to insufficient sample size, no statistical results have been yielded regarding the

relationship between growth and defense in endosymbiotic bacteria. However, a weak negative

correlation between minimal doubling time and the abundance of defense systems was observed in

symbiotic bacteria (Table 4). This suggests that symbiotic bacteria with robust defense mechanisms

tend to exhibit rapid growth. Rather than a trade-off, there appears to be a potential trade-up between

growth and defense in symbiotic bacteria. We speculate that, despite potential fluctuations in nutrient

levels, hosts typically provide a eutrophic environment for symbiotic bacteria, where energy and

resources are not limiting factors, thus rendering the trade-off between growth and defense

unnecessary. The gut lumen may favor r-selected bacteria, which prioritize rapid growth and

reproduction in unstable environments.

Several ecological features, such as free-living habitat, phototrophic lifestyle, occupation of

multiple habitats, motility, filamentous morphology, and involvement in bioremediation, have been

found to be positively associated with the abundance of defense systems (Table 2). The trade-off

between growth and defense has been statistically demonstrated in free-living and motile bacteria.

These findings suggest that certain ecological niches may favor the allocation of resources towards

defense mechanisms, potentially at the expense of growth rates, in order to cope with environmental

challenges such as predation by bacteriophages. The positive association between defense system

abundance and ecological features like free-living habitat and motility implies that bacteria thriving in

diverse and dynamic environments may require heightened defense strategies to survive and

proliferate. However, further research is needed to explore the specific mechanisms underlying these

associations and to determine how ecological factors influence the evolution of bacterial defense

strategies.

363 Supplementary Information

364 The supplementary material is submitted alongwith the main manuscript online.

366 Acknowledgements

367 This study was funded by the National Natural Science Foundation of China (grant number

368 31671321). We are grateful to Tiao-Hao Zhao for his help in machine learning techniques.

370 Author Contributions

371 DKN conceived the research hypothesis; ZLL and JL analyzed the data. ZLL and DKN drafted the

372 manuscript; all authors contributed equally to the final version and reviewed the manuscript.

374 Data Availability

375 The datasets analyzed during the current study are available in the Supplementary Files.

377 References

Bernheim A, Sorek R (2020) The pan-immune system of bacteria: antiviral defence as a community resource.
Nat Rev Microbiol 18:113-119
Bohannan BJM, Kerr B, Jessup CM, Hughes JB, Sandvik G (2002) Trade-offs and coexistence in microbial
microcosms. Antonie van Leeuwenhoek 81:107-115

Bohannan BJM, Lenski RE (2000) Linking genetic change to community evolution: insights from studies of
bacteria and bacteriophage. Ecol Lett 3:362-377
Brbić M, Piškorec M, Vidulin V, Kriško A, Šmuc T, Supek F (2016) The landscape of microbial phenotypic
traits and associated genes. Nucleic Acids Res 44:10074-10090
Buckling A, Wei Y, Massey RC, Brockhurst MA, Hochberg ME (2005) Antagonistic coevolution with parasites
increases the cost of host deleterious mutations. Proc R Soc B 273:45-49
Burmeister AR, Fortier A, Roush C, Lessing AJ, Bender RG, Barahman R, Grant R, Chan BK, Turner PE (2020)
Pleiotropy complicates a trade-off between phage resistance and antibiotic resistance. Proc Natl Acad
Sci USA 117:11207-11216
Burmeister AR, Turner PE (2020) Trading-off and trading-up in the world of bacteria-phage evolution. Curr
Biol 30:R1120-R1124
Castledine M, Padfield D, Sierocinski P, Soria Pascual J, Hughes A, Mäkinen L, Friman V-P, Pirnay J-P,
Merabishvili M, de Vos D, Buckling A (2022) Parallel evolution of Pseudomonas aeruginosa phage
resistance and virulence loss in response to phage treatment in vivo and in vitro. eLife 11:e73679
Chen Z-L, Guo H-J, Niu D-K (2023) Dependent variable selection in phylogenetic generalized least squares
regression analysis under Pagel's lambda model. bioRxiv:2023.2005.2021.541623
Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, Rocha EPC, Vergnaud G,
Gautheret D, Pourcel C (2018) CRISPRCasFinder, an update of CRISRFinder, includes a portable
version, enhanced performance and integrates search for Cas proteins. Nucleic Acids Res 46:W246-
W251
Debastiani VJ, Bastazini VAG, Pillar VD (2021) Using phylogenetic information to impute missing functional
trait values in ecological databases. Ecol Inform 63:101315
Donaldson GP, Lee SM, Mazmanian SK (2016) Gut biogeography of the bacterial microbiota. Nat Rev
Microbiol 14:20-32
Felsenstein J (1985) Phylogenies and the comparative method. Am Nat 125:1-15
Ferenci T (2016) Trade-off mechanisms shaping the diversity of bacteria. Trends Microbiol 24:209-223
Garland Jr T (2014) Trade-offs. Curr Biol 24:R60-R61
Garland T, Downs CJ, Ives AR (2022) Trade-offs (and constraints) in organismal biology. Physiol Biochem
Zool 95:82-112
Garland T, Jr., Dickerman AW, Janis CM, Jones JA (1993) Phylogenetic analysis of covariance by computer
simulation. Syst Biol 42:265-292
Georjon H, Bernheim A (2023) The highly diverse antiphage defence systems of bacteria. Nat Rev Microbiol
21:686-700
Guerin E, Hill C (2020) Shining Light on Human Gut Bacteriophages. Front Cell Infect Microbiol 10
Hall AR, De Vos D, Friman V-P, Pirnay J-P, Buckling A (2012) Effects of sequential and simultaneous
applications of bacteriophages on populations of *Pseudomonas aeruginosa* in vitro and in wax moth
larvae. Appl Environ Microbiol 78:5646-5652
Herms DA, Mattson WJ (1992) The dilemma of plants: to grow or defend. Q Rev Biol 67:283-335
Ho LST, Ane C (2014) A linear-time algorithm for Gaussian and non-Gaussian trait evolution models. Syst Biol
63:397-408
Hu E-Z, Lan X-R, Liu Z-L, Gao J, Niu D-K (2022) A positive correlation between GC content and growth
temperature in prokaryotes. BMC Genomics 23:110
Ives AR, Garland T, Jr. (2010) Phylogenetic logistic regression for binary dependent variables. Syst Biol 59:9-
26
Jessup CM, Bohannan BJM (2008) The shape of an ecological trade-off varies with environment. Ecol Lett
11:947-959
Koonin EV, Makarova KS, Wolf YI (2017) Evolutionary genomics of defense systems in archaea and bacteria.
Annu Rev Microbiol 71:233-261
Koskella B, Taylor TB, Bates J, Buckling A (2011) Using experimental evolution to explore natural patterns
between bacterial motility and resistance to bacteriophages. ISME J 5:1809-1817
Lennon JT, Khatana SAM, Marston MF, Martiny JBH (2007) Is there a cost of virus resistance in marine
cyanobacteria? ISME J 1:300-312
Lenski RE (1988) Experimental studies of pleiotropy and epistasis in *Escherichia coli*. I. Variation in
competitive fitness among mutants resistant to virus T4. Evolution 42:425-432
Liaw A, Wiener M (2002) Classification and regression by randomForest. R News 2:18-22
Liu Z-L, Hu E-Z, Niu D-K (2023) Investigating the relationship between CRISPR-Cas content and growth rate
in bacteria. Microbiol Spectr 11:e03409-03422
López-García P, Gutiérrez-Preciado A, Krupovic M, Ciobanu M, Deschamps P, Jardillier L, López-Pérez M,
Rodríguez-Valera F, Moreira D (2023) Metagenome-derived virus-microbe ratios across ecosystems.
ISME J 17:1552-1563

Lourenço M, Chaffringeon L, Lamy-Besnier Q, Pédron T, Campagne P, Eberl C, Bérard M, Stecher B,
443    Debarbieux L, De Sordi L (2020) The spatial heterogeneity of the gut limits predation and fosters
444    coexistence of bacteria and bacteriophages. Cell Host Microbe 28:390-401.e395
445 Madin JS, Nielsen DA, Brbic M, Corkrey R, Danko D, Edwards K, Engqvist MKM, Fierer N, Geoghegan JL,
446    Gillings M, Kyrpides NC, Litchman E, Mason CE, Moore L, Nielsen SL, Paulsen IT, Price ND, Reddy
447    TBK, Richards MA, Rocha EPC, Schmidt TM, Shaaban H, Shukla M, Supek F, Tetu SG, Vieira-Silva
448    S, Wattam AR, Westfall DA, Westoby M (2020) A synthesis of bacterial and archaeal phenotypic trait
449    data. Scientific Data 7:170
450 Makarova KS, Wolf YI, Iranzo J, Shmakov SA, Alkhnbashi OS, Brouns SJJ, Charpentier E, Cheng D, Haft DH,
451    Horvath P, Moineau S, Mojica FJM, Scott D, Shah SA, Siksnys V, Terns MP, Venclovas C, White MF,
452    Yakunin AF, Yan W, Zhang F, Garrett RA, Backofen R, van der Oost J, Barrangou R, Koonin EV
453    (2020) Evolutionary classification of CRISPR-Cas systems: a burst of class 2 and derived variants. Nat
454    Rev Microbiol 18:67-83
455 Makarova KS, Wolf YI, Koonin EV (2013) Comparative genomics of defense systems in archaea and bacteria.
456    Nucleic Acids Res 41:4360-4377
457 Moulton-Brown CE, Friman V-P (2018) Rapid evolution of generalized resistance mechanisms can constrain
458    the efficacy of phage–antibiotic treatments. Evol Appl 11:1630-1641
459 Pančić M, Kiørboe T (2018) Phytoplankton defence mechanisms: traits and trade-offs. Biol Rev 93:1269-1303
460 Quance MA, Travisano M (2009) Effects of temperature on the fitness cost of resistance to bacteriophage T4 in
461    *Escherichia coli*. Evolution 63:1406-1416
462 Roberts RJ, Vincze T, Posfai J, Macelis D (2015) REBASE—a database for DNA restriction and modification:
463    enzymes, genes and genomes. Nucleic Acids Res 43:D298-D299
464 Roff DA, Fairbairn DJ (2007) The evolution of trade-offs: where are we? J Evol Biol 20:433-447
465 Rostøl JT, Marraffini L (2019) (Ph)ighting phages: how bacteria resist their parasites. Cell Host Microbe
466    25:184-194
467 Roux S, Brum JR (2023) Counting dots or counting reads? Complementary approaches to estimate virus-to-
468    microbe ratios. ISME J 17:1521-1522
469 Santos T (2018) Phylogenetic eigenvectors regression and phylogentic signal-representation curve R Cran
470    Project
471 Sender R, Fuchs S, Milo R (2016a) Are We Really Vastly Outnumbered? Revisiting the Ratio of Bacterial to
472    Host Cells in Humans. Cell 164:337-340
473 Sender R, Fuchs S, Milo R (2016b) Revised Estimates for the Number of Human and Bacteria Cells in the Body.
474    PLoS Biol 14:e1002533
475 Shkoporov AN, Hill C (2019) Bacteriophages of the human gut: the "known unknown" of the mcrobiome. Cell
476    Host Microbe 25:195-209
477 Shkoporov AN, Turkington CJ, Hill C (2022) Mutualistic interplay between bacteriophages and bacteria in the
478    human gut. Nat Rev Microbiol 20:737-749
479 Stekhoven DJ, Bühlmann P (2011) MissForest—non-parametric missing value imputation for mixed-type data.
480    Bioinformatics 28:112-118
481 Stern A, Sorek R (2011) The phage-host arms race: Shaping the evolution of microbes. BioEssays 33:43-51
482 Suttle CA (2007) Marine viruses - major players in the global ecosystem. Nat Rev Microbiol 5:801-812
483 Symonds MRE, Blomberg SP (2014) A primer on phylogenetic generalised least squares. In: Garamszegi LZ
484    (ed) Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology:
485    Concepts and Practice. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 105-130
486 Tesson F, Bernheim A (2023) Synergy and regulation of antiphage systems: toward the existence of a bacterial
487    immune system? Curr Opin Microbiol 71:102238
488 Tesson F, Hervé A, Mordret E, Touchon M, d'Humières C, Cury J, Bernheim A (2022) Systematic and
489    quantitative view of the antiviral arsenal of prokaryotes. Nat Commun 13:2561
490 Thingstad TF (2022) Competition-defense trade-offs in the microbial world. Proc Natl Acad Sci USA
491    119:e2213092119
492 Weissman JL, Hou S, Fuhrman JA (2021) Estimating maximal microbial growth rates from cultures,
493    metagenomes, and single cells via codon usage patterns. Proc Natl Acad Sci USA 118:e2016810118
494 Weissman JL, Laljani RMR, Fagan WF, Johnson PLF (2019) Visualization and prediction of CRISPR incidence
495    in microbial trait-space to identify drivers of antiviral immune strategy. ISME J 13:2589-2602
496 Whittle M, Barreaux AMG, Bonsall MB, Ponton F, English S (2021) Insect-host control of obligate,
497    intracellular symbiont density. Proc R Soc B 288:20211993
498 Whittle M, Bonsall MB, Barreaux AMG, Ponton F, English S (2023) A theoretical model for host-controlled
499    regulation of symbiont density. J Evol Biol 36:1731-1744
500 Winter C, Bouvier T, Weinbauer MG, Thingstad TF (2010) Trade-offs between competition and defense
501    specialists among unicellular planktonic organisms: the "killing the winner" hypothesis revisited.

Microbiol Mol Biol Rev 74:42-57

Wommack KE, Colwell RR (2000) Virioplankton: Viruses in aquatic ecosystems. Microbiol Mol Biol Rev

64:69-114

Yang JW, Chang FH, Yeh YC, Tsai AY, Chiang KP, Shiah FK, Gong GC, Hsieh CH (2023) Trade-offs

between competitive ability and resistance to top-down control in marine microbes. Msystems

8:e01017-01022

Figure 1. Importance ranking of ecological features for defense system presence/absence in bacteria.

The vertical axis represents the names of ecological features, with higher ranks indicating a greater

impact on the presence or absence of the defense system. MeanDecreaseGini, displayed on the

horizontal axis, quantifies feature importance, with higher values indicating greater significance. The

figure displays the top thirty features in the model. Features starting with the letter "C" represent

eigenvectors representing phylogenetic relationships.

Figure 2. Importance ranking of ecological features for defense system abundance in bacteria. The

vertical axis represents the names of ecological features, with higher ranks indicating a greater impact

on the abundance of the defense system. IncNodePurity, displayed on the horizontal axis, quantifies

feature importance, with higher values indicating greater significance. The figure displays the top

thirty features in the model. Features starting with the letter "C" represent eigenvectors representing

phylogenetic relationships.

Table 1. Phylogenetic logistic regression of defense system presence/absence on bacterial habitats

Dependent variable ~ independent variable	Slope	P	$P_{adjusted}$
Presence/absence of defense ~ Known habitats = insect endosymbiont	−15.785	4.634×10^{-14}	1.854×10^{-13}
Presence/absence of defense ~ Known habitats = endosymbiont	0.286	0.452	0.452
Presence/absence of defense ~ Biotic relationship = symbiotic	−7.344	1.54×10^{-10}	3.080×10^{-10}
Presence/absence of defense ~ Host = insects general	−8.215	3.0×10^{-9}	4.0×10^{-9}

The default parameter logistic MPLE was used in the phylogenetic logistic regression of the

relationships. In the regression analysis, the presence of defense systems was assigned as 1 and the

absence was assigned as 0. The habitats are measured by the overall maximal confidence score

described in Materials and Methods section. The ratio of samples with defense systems to those

without related systems was 962 vs 82. The $P_{adjusted}$ value represents the corrected P-value computed

using the Benjamini & Hochberg method.

Table 2. PGLS regressions of ecological features on the defense system abundance in bacteria

	Slope	P	P_{adjusted}
Known habitats = insect endosymbiont	−0.001	0.001	**0.004**
Known habitats = endosymbiont	−0.001	4.4×10^{-4}	**0.002**
Biotic relationship = symbiotic	−0.002	0.022	**0.031**
Ecosystem type = circulatory system	−0.001	0.019	**0.029**
Ecosystem subtype = blood	−0.001	0.068	0.084
Habitat = host associated	−0.002	0.002	**0.005**
Host = insects general	−0.002	0.002	**0.005**
Ecosystem category = mammals	−0.002	0.015	**0.024**
Known habitats = host	−0.002	0.008	**0.017**
Ecosystem = host associated	−0.002	0.011	**0.019**
Habitat = free living	0.002	0.007	**0.016**
Habitat = aquatic	0.001	0.054	0.071
Known habitats = freshwater	0.001	0.190	0.210
Ecosystem category = aquatic	0.001	0.284	0.284
Energy source = photoautotroph	0.001	0.251	0.264
Energy source = phototroph	0.003	8.5×10^{-6}	**9.0×10^{-5}**
Energy source = chemolithotroph	0.001	0.105	0.123
Habitat = multiple	0.002	0.001	**0.004**
Shape = filamentous	0.002	3.8×10^{-6}	**8.0×10^{-5}**

Mobility	0.002	0.010	**0.019**
Ecosystem category = bioremediation	0.002	3.9×10^{-5}	**2.7×10^{-4}**

Due to the weaker phylogenetic signal of defense system abundance in bacteria compared to

ecological features (Table S1), it was utilized as the independent variable in the PGLS regression

analysis. Ecological features were measured using the overall maximal confidence score described in

the Materials and Methods section. A total of 1044 bacterial species were included in these

regressions. The $P_{adjusted}$ value represents the corrected P-value computed using the Benjamini &

Hochberg method. For readability, we have highlighted $P_{adjusted}$ values less than 0.05 in bold font.

PGLS, phylogenetic generalized least-squares. AIC, Akaike information criterion.

Table 3. PGLS regressions of bacterial minimal doubling time on ecological features

	Slope	P	$P_{adjusted}$
Known habitats = insect endosymbiont	1.809	1.0×10^{-5}	**3.0×10^{-5}**
Known habitats = endosymbiont	0.799	0.004	**0.006**
Host = insects general	1.07	3.9×10^{-9}	**4.1×10^{-8}**
Biotic relationship = symbiotic	0.609	3.3×10^{-5}	**8.7×10^{-5}**
Ecosystem = host associated	−0.859	8.4×10^{-8}	**5.9×10^{-7}**
Known habitats = host	−0.625	9.2×10^{-5}	**1.9×10^{-4}**
Habitat = host associated	−0.478	0.004	**0.006**
Ecosystem type = circulatory system	−0.197	0.325	0.341
Ecosystem subtype = blood	−0.27	0.097	0.113
Ecosystem category = mammals	−0.111	0.503	0.503
Habitat = free living	0.253	0.089	0.110
Habitat = aquatic	0.907	9.7×10^{-7}	**3.4×10^{-6}**
Known habitats = freshwater	0.693	6.7×10^{-5}	**1.6×10^{-4}**
Ecosystem category = aquatic	0.92	7.7×10^{-7}	**3.2×10^{-6}**
Energy source = photoautotroph	1.235	7.5×10^{-7}	**3.2×10^{-6}**
Energy source = phototroph	0.422	0.035	**0.049**
Energy source = chemolithotroph	1.518	2.5×10^{-12}	**5.3×10^{-11}**
Habitat = multiple	−0.656	2.2×10^{-4}	**4.2×10^{-4}**
Shape = filamentous	0.845	0.002	**0.004**

Mobility	−0.305	0.081	0.106
Ecosystem category = bioremediation	0.293	0.311	0.341

Since the minimal doubling time of bacteria exhibits a stronger phylogenetic signal than the

ecological features (Table S1), we employed it as the dependent variable in the PGLS regression

analysis. Ecological features were measured using the overall maximal confidence score described in

the Materials and Methods section. A total of 1044 bacterial species were included in these

regressions. The $P_{adjusted}$ value represents the corrected P-value computed using the Benjamini &

Hochberg method. For readability, we have highlighted $P_{adjusted}$ values less than 0.05 in bold font.

PGLS, phylogenetic generalized least-squares. AIC, Akaike information criterion.

Table 4. PGLS regressions of bacterial minimal doubling time on the abundance of defense systems

	n	Slope	*P*
Host = insects general	30	0.092	0.478
Biotic relationship = symbiotic	98	−0.009	0.210
Ecosystem = host associated	499	−0.015	**0.028**
Known habitats = host	518	−0.016	**0.021**
Habitat = host associated	434	−0.017	**0.020**
Ecosystem type = circulatory system	52	−0.012	0.194
Ecosystem subtype = blood	92	−0.018	0.104
Ecosystem category = mammals	81	−0.019	0.129
Habitat = free living	605	0.013	**0.008**
Habitat = aquatic	194	0.007	0.265
Known habitats = freshwater	147	0.001	0.901
Ecosystem category = aquatic	302	0.007	0.243
Energy source = photoautotroph	20	0.028	**0.023**
Energy source = chemolithotroph	22	0.01	0.524
Mobility	410	0.02	**0.002**
Habitat = multiple	206	0.01	0.264
Shape = filamentous	22	−0.001	0.837

Since the minimal doubling time of bacteria exhibits a stronger phylogenetic signal than the

abundance of defense systems (Table S1), we used it as the dependent variable in the PGLS

regression analysis. We discarded the ecological features where less than 20 bacterial samples were

obtained. We did adjust the *P*-values in this table using the Benjamini & Hochberg method because

the multiple regressions were not performed on the same dataset. For readability, we have highlighted

P values less than 0.05 in bold font. PGLS, phylogenetic generalized least-squares. AIC, Akaike

information criterion.