Possibilities of multiple disease progression pathways in female Alzheimer's
disease patients inferred from transcriptome and epigenome data of the
dorsolateral prefrontal cortex

Short title: Progression pathways of AD

Kousei Honda[1], Akinori Awazu[1,2,*]

[1] Graduate School of Integrated Sciences for Life, Hiroshima University,
Higashihiroshima, Hiroshima 739-8526, Japan
[2] Research Center for the Mathematics on Chromatin Live Dynamics, Hiroshima
University, Higashihiroshima, Hiroshima 739-8526, Japan

* Send correspondence to: Akinori Awazu; Email: awa@hiroshima-u.ac.jp

Keywords: Alzheimer's disease, transcriptome**,** epigenome

ABSTRACT

Late-onset Alzheimer's disease (AD) is a typical type of dementia for which therapeutic strategies have not yet been established. Based on an analysis of publicly available data of women patients, this study found that the pathology of AD was suggested to be nonuniform; however, several different typical forms of cognitive deterioration were observed. First, cluster analysis was performed on individuals diagnosed with "No Cognitive Impairment (NCI)", "Mild Cognitive Impairment (MCI)", and "Alzheimer's Disease (AD)" stages in clinical trials using gene expression and epigenome data. NCI was found to contain at least two substages, whereas MCI and AD were found to contain four and six substages, respectively. An inference of adjacency networks among substages, evaluated via partition-based graph abstraction using the gene expression profiles of individuals, suggested the possibilities of multiple typical disease progression pathways from NCI through different MCI substages and then to different AD substages. These results will aid future research on the pathological features and mechanisms of AD and induce changes in the strategic bases of conventional AD prevention and treatment.

INTRODUCTION

Alzheimer's disease (AD) is the most common cause of dementia and accounts for 60–70 % of dementia cases in older adults (Gatz et al., 2006; World Health Organization, 2017). AD can be classified into the familial type, which is strongly influenced by genetic factors, and the late-onset (sporadic) type, which is strongly influenced by aging and environmental factors, with the latter accounting for the majority of AD cases (Suresh et al., 2023). No treatment or preventive methods for AD have yet been established because the fundamental pathogenic mechanism of sporadic AD remains unclear.

Plaques of abnormally aggregated amyloid β (Aβ) are often observed in the brains of patients with AD and are considered to be related to the onset and progression of the disease (Hardy and Higgins, 1992; Hardy and Selkoe, 2002; Jin et al., 2011; Selkoe et al., 2016). However, cases have been reported in which AD develops without the accumulation of Aβ (Sekiya et al., 2018) and inhibiting Aβ production even worsened symptoms (Francesca et al., 2023). Factors other than Aβ plaque formation such as the abnormal aggregation of tau protein (Arriagada et al., 1992), the expression of apolipoprotein E variant APOE4 (Strittmatter et al., 1993), herpes virus and

coronaviruses (including COVID-19) infections (Itzhaki et al., 1997; Osaka et al., 2022; Readhead et al., 2018; Wang et al., 2022), diabetes (Kandimalla et al., 2017; Ott et al., 1999), brain inflammation (McGeer et al., 1995), oxidative damage in cells (Hirai et al., 2001; Nunomura et al., 2001; Wang et al., 2014), endoplasmic reticulum and ciliary dysplasia (Chakravarthy et al., 2012; Morelli et al., 2017; Plácido et al., 2014; Vorobyeva et al., 2018), myelin dysplasia (Bartzokis, 2004, 2011), and retrotransposons activities (Cheng et al., 2021; Macciardi et al., 2022; Ravel et al., 2021) have also been reported to influence the onset and progression of AD. Owing to the involvement of the aforementioned factors, the causes and symptoms of AD are thought to vary widely from patient to patient. Further, the variations in symptoms and disease progression rates among patients have been previously reported (Duara et al., 2021; Komarova et al., 2011; Younas et al., 2020, 2024).

In recent years, transcriptome and genome-wide analysis data of nucleotide polymorphisms have been collected from patients with AD, and various differentially expressed genes and patient-specific SNPs have been identified in healthy individuals and patients with AD (Jansen et al., 2019; Wightman et al., 2021; Yuen et al., 2020). It is notable that these characteristics are not always seen equally in all studies. One reason for this is the variations in technical factors among studies. However, the diversity of molecular conditions and gene expression patterns in the brain among patients with AD could also be a major contributor (Bellenguez et al., 2022; Johnson et al., 2022; Neff et al., 2021; Tijms et al., 2020, 2021).

Thus, AD progression is expected to exhibit various disease states. Classifying and characterizing these disease states and progression pathways is necessary to enable appropriate state-dependent treatment. In addition to a comparison of the features of the entire transcriptome, a comparison of the entire epigenome status among various patients should be useful for such classification and characterization of disease diversity. The database of the Rush Alzheimer's Disease study by ENCODE consortium (https://www.encodeproject.org/) (ENCODE Project Consortium, 2012; Luo et al., 2020) contains transcriptome data and various epigenome data such as histone modifications (H3K27ac, H3K27me3, and H3K4me3) and CCCTC-binding factor (CTCF) binding distribution of individuals in "No Cognitive Impairment (NCI)," "Mild Cognitive Impairment (MCI)," and "Alzheimer's Disease (AD)" groups that are available for unlimited public use (in the following, call Rush AD database). Therefore, the analyses of these publicly available data of patients with different cognitive

89 impairment stages are expected to elucidate the diversity in the onset and progression of
90 late-onset AD in more detail.
91
However, the Rush AD database seemed to contain a satisfactory amount of data for
women individuals but the amount of data for men seemed insufficient, particularly for
MCI and AD individuals, where the number of women individuals in NCI, MCI, AD
groups was 25, 29, and 28, respectively, whereas that of men was 14, 4, and 7,
respectively. Therefore, in this study, the classification of disease status and progression
pathway of women AD patients were performed using transcriptome and epigenome
data of women individuals in the Rush AD database as a first step of the analysis of the
diversity of AD patients using multi-omics data. Of course, this study would be
expected to be useful not only for female patients but also for understanding symptoms
in male patients, even though some features such as disease progression rate were
known to be different between male and female patients (Davis et al., 2020; Lopez-Lee
et al., 2024).
104
In this study, patient diversity was considered by assessing the deviation from healthy
individuals and the diversity of disease progression. For this purpose, we first searched
for gene sets that characterize the differences between healthy individuals and patients.
The gene set searched here was that which contained the largest number of genes from
gene sets that could actually classify healthy subjects from patients using cluster
analysis. The candidate genes included therein were selected through statistical tests of
differences in expression levels. Second, we focused on the variety of the expression
patterns of those gene sets between patients to show the existence of diverse conditions
and subgroups within patients. The existence of these subgroups was also supported by
comparisons of epigenomic data between individuals. Third, the relationship between
subgroups and the structure of the disease progression pathway was predicted based on
the similarity of gene expression between subgroups.
117
118

RESULTS
Fundamental features of the analyzed transcriptome and epigenome data
RNA sequencing (RNA-seq) data (read count and TPM) of the dorsolateral prefrontal
cortex of female individuals belonging to the NCI, MCI, and AD groups were acquired
from the Rush AD database. In this dataset, the number of RNA-seq data of individuals

in the NCI, MCI, and AD groups were 25, 29, and 28, respectively (Table 1, Table S1a-
S1b).

To examine the quality and bias of analyzed data, we estimated one of the size factors
of the RNA-seq data, that is the median of the ratios (MOR) defined in the recent
literature for DESeq (Anders et al., 2010), using read count data (Table S1a). The ratio
[standard deviation of MOR over individuals]/[average MOR over individuals] was
0.1075. Similarly, we evaluated the first and third quartiles of the ratio (1OR and 3OR)
between the average and standard deviation, which were 0.1147 and 0.1006,
respectively. These facts suggested that the size factor would not considerably input
bias among samples or affect the comparison of either the read count data or TPM
values.

When examining the significance of the difference in gene expression levels between
individuals aged <85 years and those >90 years, we did not detect any genes exhibiting
a false discovery rate (FDR) < 0.1 (likelihood ratio test and two-tailed t-test) (Table
S1a-S1b). This suggested that the variations in gene expression levels among the NCI,
MCI, and AD groups was not due to the influence of age, but rather due to differences
in clinical stages.

From the Rush AD database, in addition to RNA-seq data, the narrow peak data on
genome-wide histone modifications (H3K27ac, H3K27me3, and H3K4me3) and CTCF
binding distribution obtained by chromatin immunoprecipitation sequencing (ChIP-seq)
could be acquired for 39 individuals: 11, 11, and 17 individuals of NCI, MCI, and AD
patients, respectively (Table S1c). Here, for ChIP-seq data of H3K27ac, H3K27me3,
H3K4me3, and CTCF, the correlation coefficients between the total read number
mapped to the human genome and the number of narrow peaks were obtained as 0.193
(P-value = 0.239), 0.146 (P-value = 0.374), 0.027 (P-value = 0.872), and 0.333 (P-value
= 0.038), respectively. These results suggested the number of narrow peaks of
H3K27ac, H3K27me3, and H3K4me3 seemed to be independent of the total number of
mapped reads. For the data of CTCF, the correlation between the total number of
mapped reads and the number of narrow peaks was not considered 0 but expected
sufficiently small with which the total number of mapped reads did not influence the
analytic results of narrow peaks seriously.

Cluster analysis of individuals' provided gene set that characterized the differences in major MCI individuals from others

To extract a set of genes that might characterize the differences among the NCI, MCI, and AD groups, we performed the likelihood ratio test, which is one of the most familiar tests for estimating differentially expressed genes (DEGs) from read count data, for each gene using the DESeq2 software (Love et al., 2014) (Table S1a). Consequently, we performed hierarchical clustering of individuals using genes with P-values below the threshold value P^D_{NMA} = 0.01, 0.02, 0.03, 0.04, and 0.05 (Fig. 1, Fig. S1a-S1e). When $P^D_{NMA} \leq 0.05$, a cluster with significantly more MCI individuals (MCI cluster) and a cluster with NCI, AD, and a small number of MCI individuals was formed by the first divergence of the dendrogram ((P < 0.05, Fisher's exact test; Fig. S1a-S1e). However, even though the gene set for clustering contained only genes with smaller P values, individuals with NCI, MCI, and AD were not from their respective cluster; no clusters comprising solely of individuals with NCI, MCI, or AD were formed by the first divergence of the dendrogram.

We then performed the same analysis using analysis of variance (ANOVA), which is the most familiar test for evaluating differences of characteristic values among three or more groups, to determine the P-values using TPM values (Table S1b), where P^A_{NMA} was defined instead of P^D_{NMA}. Here, ANOVA was generally better for detecting differences and was expected to identify more candidate genes in a gene set than the likelihood ratio test. As a result, an MCI cluster and a cluster with NCI, AD, and a small number of MCI individuals were formed, as in the previous case (Fig. S1f-S1j). Additionally, when the P^A_{NMA} was 0.04, a cluster comprising solely of individuals with MCI was formed by the first divergence of the dendrogram (Fig. 1a). In contrast, we did not observe any clear separation between participants with NCI and those with AD, regardless of the P^A_{NMA} value.

It should be noted that we did not obtain any gene set by the P^D_{NMA} and P^A_{NMA} values that enabled the complete discrimination of NCI, MCI, and AD individuals. Additionally, depending on the P^D_{NMA} and P^A_{NMA} values, the individuals with MCI who deviated from the MCI cluster varied; however, 24 MCI individuals were always included in the MCI cluster (Fig. S1). This suggested that the MCI cluster consisted of 24 MCI individuals, named MCI_{Major} individuals, whereas the five other MCI individuals (M3, M12, M14, M16, and M28) who tended to deviate from the MCI_{Major} individuals, were named MCI_{NCI_AD} individuals. We assumed that the MCI_{Major} and

MCI_{NCI_AD} individuals were at the substages named MCI_{Major} and MCI_{NCI_AD},

respectively.

Moreover, the set of genes with P-values of the above ANOVA $\leq P^A_{NMA} = 0.04$ was

named the "M-NA set" as it characterized the differences between majority of MCI

individuals from NCI and AD individuals (Fig. 1b, Table S2a). In a 3-dimensional U-

map using M-NA set, we clearly observed the deviation of the MCI_{Major} individuals

from individuals at other substages, where MCI_{NCI_AD} were found in a mixed group of

NCI and AD individuals (Fig. 1c).

Note that the classification result of MCI individuals into MCI_{Major} and MCI_{NCI_AD}

individuals seemed to be independent of the ages of individuals. For example, the

numbers of individuals over and under 90 years old in MCI_{Major} were 19 and 5, and

those in MCI_{NCI_AD} were 3 and 2, respectively (Table S1a-S1b), by which the P-value of

Fisher's exact test was obtained 0.5688.

Functional features of genes in the M-NA set

We also considered the functional features of genes in the M-NA set. In this set, we

obtained 594 genes (290 upregulated and 304 downregulated genes in MCI_{Major}), of

which 349 genes (164 upregulated and 185 downregulated in MCI_{Major}) were annotated

(Fig. 1b) as protein coding or noncoding RNA genes using Entrez (Schuler et al., 1996).

Herein, both sets of upregulated and downregulated genes in MCI_{Major} contained some

previously reported AD risk-associated genes; the upregulated gene set contained

CYP19A1, GSTM1, HSD11B1, IL10, MT-RNR1, PCK1, TRAK2, UBE2D1, and *XBP1*,

whereas the downregulated gene set contained *MT-ATP8* and *NME8* (Hu et al. 2017).

Additionally, we performed enrichment analysis of all genes including genes with

higher and those with lower expression levels in the MCI_{Major} individuals (Table S2b–

S2d). We identified that genes with higher expression levels in MCI_{Major} individuals

were enriched in GO terms for the functions "response to lipopolysaccharide,"

"response to molecule of bacterial origin," "Toll-like receptor signaling pathway,"

"regulation of MHC class II biosynthetic process," "vitamin metabolic process," "water-

soluble vitamin metabolic process," "chemotaxis," and "taxis" (P-value $< 10^{-3}$ and FDR

< 0.5) (Table S2c). Whereas, we identified that genes with lower expression levels in

MCI_{Major} individuals were enriched in GO terms for the functions "mRNA 5'-splice site

recognition" and "mRNA cis splicing, via spliceosome" (P-value $< 10^{-3}$ and FDR < 0.5)

(Table S2d). Recent studies have suggested that some of these functional features

characterize the pathology of AD; for example, the lipopolysaccharide found in the wall

of all Gram-negative bacteria was reported to play a role in causing AD (Singh et al.,

2022; Zhan et al., 2018), Toll-like receptors participate in neuronal cell death by

apoptosis (Adhikarla et al., 2021; Wu et al., 2022), MHC class II-expressing cells may

be involved in the degradation of neurons (Parachikova, et al., 2007), vitamins affect the

generation and clearance of Aβ (Grimm et al., 2016), and microglia exhibit chemotaxis

for the clearance of Aβ (Lau et al., 2023; Rogers et al., 2001).

240 Gene sets that differentiate groups of NCI and AD individuals

To extract a gene set that may characterize the difference between the NCI and AD

individuals, we performed hierarchical clustering of individuals using the genes

satisfying the following criteria: mean expression level (TPM) was higher in AD

individuals than in NCI individuals, with smaller P-values in the likelihood ratio test

than the P^D_{NA} threshold, or mean expression level (TPM) was lower in AD individuals

than in NCI individuals, with smaller P-values in the likelihood ratio test than the P^D_{AN}

threshold. Subsequently, we searched the sets of P^D_{NA} and P^D_{NA} values that divided the

NCI and AD individuals into different clusters in the first branch of the dendrogram.

Within the range $P^D_{NA} \leq 0.05$ and $P^D_{AN} \leq 0.05$, we identified the finite but narrow area

of P^D_{NA} and P^D_{AN}, $P^D_{NA} = 0.01$ and $P^D_{AN} = 0.02$, $P^D_{NA} = 0.01$ and $P^D_{AN} = 0.03$, $P^D_{NA} =$

0.02 and $P^D_{AN} = 0.03$, $P^D_{NA} = 0.03$ and $P^D_{AN} = 0.03$, and $P^D_{NA} = 0.05$ and $P^D_{AN} = 0.02$,

where the NCI individuals and AD individuals completely separated (Fig. 2a). Thus, we

obtained the largest set of genes separating the NCI and AD groups at $P^D_{NA} = 0.05$ and

$P^D_{AN} = 0.02$ (Fig. S2a, Table S3a); this gene set was named the N-A$_{D1}$ set. It should also

be noted that the number of genes with a lower mean expression level (TPM) in the AD

individuals than in the NCI individuals was larger when $P^D_{NA} = 0.03$ and $P^D_{AN} = 0.03$

than when $P^D_{NA} = 0.05$ and $P^D_{AN} = 0.02$ (Fig. S2b, Table S3b). Of course, this gene set

contained all genes in other gene sets with smaller P^D_{NA} and P^D_{AN} than 0.03 that could

separate the NCI and AD groups. Thus, we focused on the gene set obtained at $P^D_{NA} =$

0.03 and $P^D_{AN} = 0.03$ and named it the "N-A$_{D2}$ set."

We performed the same analysis using two-tailed *t*-tests to determine the P-values,

defining the P^t_{NA} and P^t_{AN} instead of the P^D_{NA} and P^D_{AN}, respectively. Here, the t-test

was generally better for detecting differences and was expected to identify more

candidate genes in a gene set than the likelihood ratio test. Searching in the ranges P^t_{NA}

≤ 0.05 and $P^t_{AN} \leq 0.05$, the NCI individuals and the AD individuals were completely

separated in 12 of 15 cases of P^t_{NA} and P^t_{AN} set for $P^t_{NA} \leq P^t_{AN}$ (Fig. 2a). When we

searched the wider range $P^t_{NA} \leq 0.1$ and $P^t_{AN} \leq 0.1$, the NCI and AD individuals were

completely separated in 37 of 55 cases of P^t_{NA} and P^t_{AN} set for $P^t_{NA} \leq P^t_{AN}$ (Fig. 2a). We

thus obtained the largest set of genes separating the NCI and AD groups at $P^t_{NA} = 0.08$,

and $P^t_{AN} = 0.1$ (Fig. S2c, Table S3c). This gene set was named the "N-A_t set."

Hierarchical clustering using a combination of the expression levels of all genes in the

N-A_t and N-A_{D1} sets also separated the NCI and AD individuals (Fig. 2b, Table S3d).

Conversely, the expression of all genes in the N-A_t and N-A_{D2} sets did not separate the

NCI and AD individuals. However, clustering using a combination of the expression

levels of genes in the N-A_t set and a part of the N-A_{D2} set with low P-values completely

separated the NCI and AD individuals (Fig. 2c, Table S3e). Such gene sets are expected

to involve various typical genes that characterize the differences between NCI and AD

individuals. In this study, the gene set consisting of the N-A_t and N-A_{D1} sets, which

could completely separate the NCI and AD individuals, was named the "N-A1 set"

(Table S3d), whereas the gene set consisting of the N-A_t set and the maximum number

of genes in the N-A_{D2} set with low P-values was named the "N-A2 set" (Table S3e).

AD individuals can be classified into three groups

From the results of the hierarchical clustering for NCI and AD individuals performed

using the N-A1 set, we noticed that three subgroups of AD individuals emerged (Fig.

2b, Fig. 3a). Thus, assuming that AD individuals could be classified into these three

subgroups, we performed ANOVA on the gene expression levels of genes whose

expression levels were higher in AD individuals than in NCI individuals among the

three assumed subgroups (1020 genes). We identified 367 genes whose expression

levels differed significantly (FDR < 0.1) among the subgroups. Performing k-means

cluster analysis of these genes using their expression levels in AD individuals, we

identified two characteristic clusters: a cluster consisting of 96 genes whose expression

levels were low in only one subgroup (Cluster CA1) and another consisting of 271

genes whose expression levels were high in only one subgroup (Cluster CB1) (Fig. 3a).

Thereafter, we performed a similar analysis for the NCI individuals. Based on the shape

of the dendrogram, the NCI individuals were divided into three or four subgroups (Fig.

2b, Fig. 3a). However, when the NCI individuals were assumed to be divided into three

subgroups, ANOVA revealed only nine genes with FDR < 0.1 among the three

subgroups (Table S3d). Additionally, when NCI individuals were divided into four

subgroups, ANOVA revealed only two genes with FDR < 0.1 among the three

subgroups (Table S3d). Based on these values, the number of genes in clusters CA1 and

CB1 was sufficiently large.

Collectively, these results suggested that the gene expression patterns of AD

individuals, which can be classified into three types, were highly nonuniform. AD

individuals exhibiting high gene expression levels in both clusters CA1 and CB1 were

regarded as individuals in the AD_{HH} subgroup (named AD_{HH} individuals), those

exhibiting low gene expression levels in both CA1 and CB1 were regarded as

individuals in the AD_{LL} subgroup (named AD_{LL} individuals), and those exhibiting high

gene expression levels in cluster CA1 and low gene expression levels in cluster CB1

were regarded as individuals in the AD_{HL} subgroup (named AD_{HL} individuals).

Hierarchical clustering of the NCI and AD individuals using the N-A2 set also exhibited

three possible subgroups (Fig. 2c, Fig. 3b). By performing the same analysis described

above, we identified gene clusters similar to cluster CA1 containing 86 genes (cluster

CA2) and cluster CB1 containing 234 genes (cluster CB2). Similar to a previous

method, AD individuals were divided into AD_{HH}, AD_{LL}, and AD_{HL} groups based on the

expression profiles of genes in clusters CA2 and CB2. The members of the AD_{HH},

AD_{LL}, and AD_{HL} groups were the same in both the N-A1 and N-A2 sets (Fig. S3a).

Importantly, the AD_{HH}, AD_{HL}, and AD_{LL} individuals generally varied when different

P^t_{NA} and P^t_{AN} values were employed to determine the genes in the N-At set. However,

for small changes in the P^t_{NA} and P^t_{AN} values, less exchange of individuals was observed

among subgroups. For example, considering the case where $P^t_{NA} = 0.06$ and $P^t_{AN} = 0.1$

were employed instead of $P^t_{NA} = 0.08$ and $P^t_{AN} = 0.1$ to determine the N-At. Herein, the

gene set named N-A3 was defined as that consisting of genes in the $N-A_t$ set and the

maximum number of genes in $N-A_{D1}$ set with low P-values that could completely

separate the NCI and AD individuals, whereas the gene set named N-A4 was defined as

that consisting of genes in the $N-A_t$ set and the maximum number of genes in $N-A_{D4}$ set

with low P-values that could completely separate the NCI and AD individuals. The

classification of AD individuals into the AD_{HH}, AD_{LL}, and AD_{HL} groups was similar to

the results obtained for N-A1, N-A2, N-A3, and N-A4, where 24 of the 28 individuals

belonged to the same subgroups (Fig. S3). This suggested the robustness of our method

in classifying the AD_{HH}, AD_{HL}, and AD_{LL} individuals.

Additionally, note that the classification result of AD individuals into AD_{HH}, AD_{HL}, and

AD_{LL} individuals seemed to be independent of the ages of individuals. For example, the

numbers of individuals over and under 90 years old in AD_{HH} were 8 and 2, those in

AD_{LL} were 6 and 4, and those in AD_{HL} were 4 and 4, respectively, by which the P-value

of Fisher's exact test was obtained 0.4850.

Common functional features of the N-A1 and N-A2 sets

We also considered the common functional features of the N-A1 and N-A2 sets. The N-

A1 set contained 1477 genes (1020 upregulated and 457 downregulated, AD), and 933

genes (683 upregulated and 250 downregulated) were annotated as protein coding or

noncoding RNA genes using Entrez (Schuler et al., 1996) (Fig. 3a). The N-A2 set

contained 1424 genes (947 upregulated and 477 downregulated, AD), and 889 genes

(619 upregulated and 270 downregulated) were annotated as protein coding or

noncoding RNA genes using Entrez (Schuler et al., 1996) (Fig. 3b). The upregulated

genes in both the N-A1 and N-A2 set contained *AR*, *CBS*, *CLU*, *CYP19A1*, *DBH*,

GNA11, *HLA-A*, *IQCK* , *MPO*, *MS4A4E*, *MX1*, *NLRP3*, *SDC2*, *TAP2*, and *ZFHX3*

(Table S3d-e), all of which were reported to be AD risk-associated genes in recent

studies (Hu et al. 2017; Xiang et al. 2018; Rahman et al. 2019; Yang et al. 2022).

Similarly, the downregulated genes in both the N-A1 and N-A2 set contained *COL25A1*

and *TARDBP* (Table S3d-e), which were also reported to be AD risk-associated genes

(Hu et al. 2017).

Additionally, the clusters CA1, CB1, CA2, and CB2 contained 71, 215, 64, and 181

annotated genes, respectively (Fig. 3). We found *CLU*, *SDC2*, and *ZFHX3* in both the

CA1 and CA2 clusters, whereas *DBH*, *GNA11*, *IQCK*, *MPO*, and *NLRP3* were found in

both the CB1 and CB2 clusters (Table S3d-e), thereby suggesting that these genes are

responsible for the differences in the features among the three substages of AD (Hu et

al. 2017; Yang et al. 2022).

Next, we performed enrichment analysis of all genes, including those with higher and

those with lower expression levels in the AD individuals for both the N-A1 and N-A2

set (Table S3f–S3o). We identified that genes with higher expression levels in AD

individuals in both the N-A1 and N-A2 set were commonly enriched in GO terms for

"forebrain dorsal/ventral pattern formation," "mRNA 5'-splice site recognition," and

"negative regulation of NF-kappaB transcription factor activity" (P-value < 10^{-3} and

FDR < 0.5) (Table S3g, S3j). Whereas, genes with lower expression levels in AD

individuals in both the N-A1 and N-A2 set were enriched in GO terms for the functions
"negative regulation of intrinsic apoptotic signaling pathway in response to DNA
damage by p53 class mediator," "negative regulation of DNA damage response, signal
transduction by p53 class mediator," and "coronary vasculature morphogenesis" (P-
value $< 10^{-3}$ and FDR < 0.5) (Table S3h, S3k). In the nervous system, NF-κB is known
as one of the crucial components in the molecular switch that converts short- to long-
term memory (Kaltschmidt et al., 2015). Thus, the upregulation of genes that suppress
NF-κB expression is expected to exacerbate the pathology of AD. Additionally, the
downregulation of genes that suppress the apoptosis in response to DNA damage would
also promote AD progression through the enhancement of cell death in the central
nervous system (Lin et al., 2020; Ranganathan et al., 2022).
386
We found that genes in both the CA1 and CA2 clusters were enriched in GO terms for
the functions "response to xenobiotic stimulus", "cellular response to lipid", "negative
regulation of small molecule metabolic process", "lung development", "cellular
response to external stimulus", "negative regulation of carbohydrate metabolic process",
"neuron migration", and "positive regulation of epithelial cell differentiation" (P-value <
10^{-3} and FDR < 0.5) (Table S3l, S3n). Whereas, genes in both the CB1 and CB2 clusters
were enriched in GO terms for the functions "mitotic spindle assembly checkpoint
signaling" and "L-alpha-amino acid transmembrane transport" (P-value $< 10^{-3}$ and FDR
< 0.5) (Table S3m, S3o). Notably, recent studies suggested that the perturbed cerebral
carbohydrate metabolism contributes to the pathogenesis of AD (Chen et al., 2013;
Cunnane et al., 2010), whereas the neuron migration ability contributes to neurogenesis
in brain (Esteve et al., 2022; Young-Pearse et al., 2007). These facts suggested that the
three substages of AD may be characterized by differences in these functional features.
400
401 Additionally, the chromosomal distribution of genes that belong to CA1 or CB1 cluster
402 in N-A1 set, and that to CA2 or CB2 clusters in N-A2 set, were evaluated (Fig. 4).
403 Compared to the chromosomal distribution of all human coding and noncoding genes,
404 both genes set belonging to CA1 or CB1 clusters and that to CA2 or CB2 clusters
405 seemed to accumulate at Chr. 1, 7, 9, 16, 19, and 22 (Fig. 4).
406
Epigenome data analysis supported classification of AD individuals into three
groups
The data from the dorsolateral prefrontal cortex of Rush AD analyzed in the present
study also included data on genome-wide distributions of epigenomic markers such as

histone modification (H3K4me3, H3K27me3, and H3K27ac) and CTCF binding sites
distributions for approximately half of all individuals (Table S1). Subsequently, we
evaluated the similarity of such epigenomic features among individuals using Jaccard
coefficients of the genome-wide distributions of epigenome status to classify them
based on other transcriptome features. At the first divergence of the dendrogram
according to the hierarchical clustering of individuals using Jaccard coefficients,
individuals were divided into two groups: a group with a typical distribution of
epigenome status and that with an untypical one, for respective epigenomic markers
(Fig. 5-6, Fig S4). The Jaccard coefficients of H3K27ac, H3K27me3, and CTCF among
individuals suggested that more AD_{HL} individuals exhibited untypical patterns
compared with those in other AD individuals (Fig. 5, Table 2a–2c, Fig. S4).
Additionally, a larger number of AD_{HH} exhibited untypical genome-wide distribution
patterns of H3K4me3 compared with those in other AD individuals (Fig. 6, Table 2d).
These facts suggested that the AD_{HH}, AD_{HL}, and AD_{LL} individuals exhibited different
epigenomic features.
426
Furthermore, we evaluated the epigenetic differences between individuals using the
density of each epigenomic marker on each chromosome. As a result, we observed a
large difference in the density of histone modification on chromosomes 17, 19, and X
between Typical and Untypical groups, and the density of CTCF binding mainly on
chromosomes 17 and 19. This fact suggested that the density of H3K27ac and
H3K27me3 on chromosomes 17, 19, and X, and CTCF on chromosomes 17, 19 differed
between AD_{HL} individuals and those at other substages. Additionally, H3K4me3 on
chromosomes 17, 19, and X also differed between AD_{HH} individuals and those at other
substages.
436
437 Notably, chromosomes 17, 19, and X contain the Tau, APOE, and KDM6A genes,
438 respectively, which are deeply involved in the progression of Alzheimer's disease.
439 However, they were found neither M-NA set, N-A1 set, nor N-A2 set.
440
Adjacency network among substages
The data on the expression levels of genes in the M-NA set revealed the differences
between typical MCI, NCI, and AD individuals, whereas those in the N-A1 or N-A2
sets revealed the differences between NCI and AD individuals. To clarify the
relationships among disease substages and the pathways of disease progression through
these substages, we described the adjacency networks among substages using these

gene sets by performing the partition-based graph abstraction (PAGA) method (Wolf et
al., 2019). Using PAGA, we generated a graph of substages in which the edge weights
represented confidence in the presence of connections. Notably, the confidence of
connections between NCI and substages in AD often involved finite values (Table S4);
however, the direct connection between NCI and substages in AD seemed unnatural in
disease progression. Therefore, we assumed that only connections with confidence
values sufficiently larger than the maximum confidence values of the connection
between NCI and substages in AD (MD_{N_A}) were effective for the adjacency network
among substages (Fig. S5). Subsequently, we used connections with confidence values
larger than twice that of MD_{N_A} to construct the adjacency network (Table S4).

Changes in gene expression patterns that characterize disease states are considered to
occur gradually as the disease progresses. Therefore, the disease appeared to progress
from NCI to AD along the previously obtained adjacency network. Note that the
adjacency network obtained using the M-NA and N-A1 sets (Fig S5a) and that
constructed using the M-NA and N-A2 sets (Fig. S5b) exhibited a little different
structure because the contained genes in these sets were a little different. Thus, we
focused on the network constructed using common connections obtained from both
networks constructed using the M-NA and N-A1 sets (Fig S5a) and the M-NA and N-
A2 sets (Fig. S5b) (Fig. 7). We did not obtain any connections from NCI to MCI_{Major},
which is the major group of MCI individuals, in any of the networks (Fig. 7, Fig S5a-b),
which seemed unnatural for the progression of AD. Thus, we performed more detailed
subgroup classifications of NCI, MCI, and AD individuals to reveal the relationship
between NCI and MCI_{Major} as described below.

Detailed classification of MCI by comparing between NCI and MCI and between
MCI and AD
To classify MCI individuals more comprehensively, we conducted comparisons
between NCI and MCI individuals and between MCI and AD individuals. First, we
performed hierarchical clustering of individuals using the genes satisfying the following
criteria: mean expression level (TPM) was higher in the MCI group than in the NCI
group with smaller P-values in the likelihood ratio test than the P^D_{MN} threshold or the
mean expression level (TPM) was lower in the MCI group than in the NCI group, with
smaller P-values in the likelihood ratio test than the P^D_{NM} threshold. We performed
cluster analysis using each gene set within the range $P^D_{NM} \leq 0.05$ and $P^D_{MN} \leq 0.05$.
However, we did not identify any cases in which the NCI and MCI individuals were

completely separated (Fig. 8a, Fig. S6a). However, clusters with significantly more NCI

individuals (NCI clusters) and MCI clusters were formed within all cases of P^D_{NM} and

P^D_{MN} except a case of $P^D_{NM} = 0.05$ and $P^D_{MN} = 0.01$ (Fig. 8a).

When we used two-tailed t-tests instead of the likelihood ratio test to determine the P-

values, the results were qualitatively the same. P^t_{MN} and P^t_{NM} were defined instead of

P^D_{MN} and P^D_{NM}, respectively. Analysis using each gene set within the ranges $P^t_{MN} \leq 0.05$

and $P^t_{NM} \leq 0.05$ revealed no cases in which the NCI and MCI individuals were

completely separated (Fig. 8a, Fig. S6a). However, the NCI and MCI clusters were

formed within all cases of P^t_{NM} and P^t_{MN} (Fig. 8a).

Next, we performed the same analyses for the MCI and AD individuals. In this case,

P^D_{MA}, P^D_{AM}, P^t_{MA}, and P^t_{AM} were given as values corresponding to P^D_{NM}, P^D_{MN}, P^t_{NM},

and P^t_{MN}, respectively. Similar to the above results, we did not identify any cases in

which the MCI and AD individuals were completely separated (Fig. 8b, Fig. S6b).

However, the MCI and AD clusters were formed within all cases of P^D_{MA}, P^D_{AM}, P^t_{MA},

and P^t_{AM} except a case of $P^D_{MA} = 0.05$ and $P^D_{AM} = 0.01$ and that of $P^t_{MA} = 0.05$ and P^t_{AM}

$= 0.01$ (Fig. 8b).

In this study, the MCI individuals who deviated from the MCI cluster varied depending

on P^D_{NM}, P^D_{MN}, P^t_{NM}, P^t_{MN}, P^D_{MA}, P^D_{AM}, P^t_{MA}, and P^t_{AM} values. For each individual with

MCI, we counted the number of times they belonged to the NCI cluster upon changing

the P^D_{NM}, P^D_{MN}, P^t_{NM}, and P^t_{MN} values, and those belonging to the AD cluster upon

changing the P^D_{MA}, P^D_{AM}, P^t_{MA}, and P^t_{AM} values (Fig. 8c). Consequently, individuals

M3, M4, M5, M8, M12, M13, M14, M16, M28, and M29 belonged to the NCI cluster

more than once, whereas individuals M3, M5, M9, M12, M14, M16, M23, M27, and

M28 belonged to the AD cluster more than once.

Notably, individuals M3, M12, M14, M16, and M28, (same as those in MCI_{NCI_AD}),

clearly exhibited a larger number of deviations from NCI and AD than those in MCI_{Major}

(Fig. 8c). To classify the MCI_{Major} individuals in more detail, we performed k-means

clustering using the number of times each individual belonged to the NCI cluster and

that belonging to the AD cluster (Fig. 8c, Fig. S6c). As a result, the MCI_{Major} individuals

were divided into three subtypes: $MCI_{Typical}$ individuals (17 individuals), who rarely or

never belonged to other clusters; MCI_{NCI} individuals (four individuals), who may

belong to NCI clusters only; and MCI_{AD} individuals (three individuals), who may

belong to AD clusters only (Table S1a-S1b). Thus, the MCI stage was divided into four

substages: $MCI_{Typical}$, MCI_{NCI}, MCI_{AD}, and MCI_{NCI_AD}.

Detailed classification of NCI by comparing between NCI and MCI individuals

The analysis performed in the previous section also showed that individuals N11, N12,

N18, N19, N21, and N22 belonged to the MCI cluster more than once when the P^D_{NM},

P^D_{MN}, P^t_{NM}, and P^t_{MN} values were changed (Fig. 9a). After k-means clustering of the

NCI individuals using the number of times each individual belonged to the MCI cluster,

the NCI individuals were divided into two clusters: one comprising N11, N21, and N22,

and the other comprising NCI individuals who never belonged to the MCI cluster (Fig.

9a, Fig. S7a). Therefore, these individuals were classified as typical NCI individuals

belonging to the MCI cluster and named $NCI_{Typical}$ individuals, and were assumed to be

in the $NCI_{Typical}$ substage. Individuals N12, N18, and N19 were classified as NCI

individuals close to MCI and named NCI_{MCI} individuals, and were assumed to be in the

NCI_{MCI} substage (Table S1a-S1b).

Detailed classification of AD by comparing between AD and MCI individuals

Similarly, 17 AD individuals belonged to the MCI cluster more than once, when the

P^D_{MA}, P^D_{AM}, P^t_{MA}, and P^t_{AM} values were changed (Fig. 9b). Using k-means clustering of

AD individuals based on the number of times each individual belonged to the MCI

cluster in the previous section, AD individuals were divided into three clusters, with

individuals A10, A14, A15, A17, A18, A19, A22, and other AD individuals who never

belonged to the MCI cluster being included in the same cluster (Fig. 9b, Fig S7b).

Therefore, these participants were expected to be in one of the typical AD substages:

AD_{HH}, AD_{HL}, or AD_{LL} (Table S1a-S1b). In contrast, individuals A2, A3, A4, A6, A7,

A8, A9, A11, A20, and A21 were assumed to be at one of the AD substages close to

MCI and named AD_{HH_MCI}, AD_{HL_MCI}, or AD_{LL_MCI} (Table S1a-S1b).

Adjacency network among substages obtained after detailed classifications of

individuals suggests multiple disease progression pathways from NCI to AD

Detailed classifications of individuals suggested that NCI was classified into $NCI_{Typical}$

and NCI_{MCI}; MCI was classified into $MCI_{Typical}$, MCI_{NCI}, MCI_{AD}, and MCI_{NCI_AD}; and

AD was classified into AD_{HH}, AD_{HH_MCI}, AD_{HL}, AD_{HL_MCI}, AD_{LL}, and AD_{LL_MCI}. We

constructed adjacency networks among these 12 substages using gene expression data

from the M-NA and N-A1 sets (Fig. S8a, Table S5a) and the M-NA and N-A2 sets (Fig.

S8b, Table S5b) as previously described. Subsequently, we focused on the adjacency

network that consisted of common connections between the network from the data in

the M-NA and N-A1 sets and that from the data in the M-NA and N-A2 sets (Fig. 10).

We identified four or more progression paths from NCI to AD that passed through

different substages, as follows: I) from NCI ($NCI_{Typical}$ or NCI_{MCI}) to AD_{LL} through

MCI_{NCI_AD}, II) from NCI to MCI_{NCI} and from MCI_{NCI} to AD_{LL} through AD_{HH_MCI} and

AD_{LL_MCI}, III) from NCI to MCI_{NCI} and from MCI_{NCI} to AD_{HH} through $MCI_{Typical}$ or

AD_{HH_MCI}, and IV) from NCI to AD_{HL} or AD_{LL} via MCI_{NCI}, MCI_{AD}, and AD_{HL_MCI}. This

suggested that at least four disease progression pathways are responsible for the

progression of the three classified AD substages.

Results for comparisons with recent studies

Many recent studies performed transcriptome analyses and identified various significant

differentially expressed genes (significant DEGs) between healthy (control) individuals

and patients with AD as "AD risk genes" (Hu et al. 2017; Xiang et al. 2018; Rahman et

al. 2019; Yang et al. 2022). The cluster analysis of NCI, MCI, and AD individuals and

that of NCI and AD individuals was performed using these AD risk genes (Fig. S9,

Table S6). The results differed from the individual clustering results in Fig. 1, 2, and 3.

It is noteworthy that most of the data of AD risk genes used here were obtained from

research on the hippocampus, while the data analyzed in the previous sections of this

study were obtained from the prefrontal cortex. One of the recent studies has shown that

gene expression patterns differ between the hippocampus and the prefrontal cortex (Nef

et al., 2021). These facts were expected to be the reason for such inconsistency.

From the data set analyzed in the previous sections, no significant DEGs between NCI

and AD individuals and only 9 significant DEGs among NCI, MCI, and AD individuals

were obtained (Table S1a-S1b, Table S7). Here, significant DEGs were evaluated by the

likelihood test using DEseq2 with FDR < 0.01. Notably, the abovementioned study

reported that not many significant DEGs were yielded in areas belonging to the

prefrontal cortex (Nef et al., 2021). These facts suggested to support the consistency

between the results of recent and present studies.

DISCUSSION

In this study, by the analysis of the data from women individuals in Rush AD database,

the pathological progression pathway of AD was suggested to be nonuniform, with

different typical forms of deterioration. First, cluster analysis of individuals using gene

expression data from those diagnosed with NCI, MCI, and AD through clinical trials

suggested that NCI contained at least two substages, whereas MCI and AD contained

four and six substages, respectively. The differences among AD substages were

characterized based on epigenomic features such as the genome-wide distributions of

histone modifications and CTCF binding. Additionally, some of the genes and gene sets

with respective GO terms that were suggested as AD-associated in recent studies

seemed to contribute to the characterization of each AD substage. Second, an inference

of adjacency networks among the different substages suggested the existence of four

typical NCI → AD disease progressions from different MCI to different AD substages.

These typical pathological progression pathways were unknown, so these findings on

disease progression features will contribute to the development of AD research.

Additionally, they are expected to induce changes in the strategic bases of AD

prevention and treatment.

Recent studies using hippocampal and frontal lobe transcriptomes have also suggested

the existence of multiple AD subtypes. However, the relationship between multiple

states, such as their connection in series on the same disease progression path or on

different paths, has not been clarified. Furthermore, these studies did not focus on the

details of the diversity of NCI and MCI. In contrast, in this study, both NCI and MCI

individuals were divided into multiple subgroups at different substages based on their

respective gene expression characteristics. Additionally, estimating the connectivity

among substages clearly revealed multiple pathological pathways for disease

progression.

This study analyzed data from the dorsolateral prefrontal cortex of Rush AD database,

rather than the hippocampus, which has been the focus of many previous studies. The

reason for the analysis of this dataset in this study was that, for some individuals, this

dataset included not only transcriptome data but also data of genome-wide distributions

of histone modification (H3K4me3, H3K27me3, H3K27ac) and CTCF binding sites

(Table S1). Such datasets can potentially reveal the relationship between a wide range

of transcriptional regulatory states and disease conditions. Recently, a relationship

between changes in CTCF binding and transcription activities associated with changes

in pathological conditions was found by analyzing such datasets (Patel et al., 2023). The

analysis of these epigenome data in the present study suggested that individuals in three

different AD substages exhibited different respective epigenomic features, particularly

in chromosomes 17, 19, and X.

It is also notable that these three subgroups of AD were classified according to the

difference in the expression level of genes that tended to be enriched on Chr. 1, 7, 9, 16,

19, and 22 (Fig. 4). Here, the differences in expression levels of genes on chromosome

19 among three AD groups may be directly related to the different epigenetic status of

chromosome 19 among three AD groups. However, the expression of genes on other

chromosomes may be indirectly influenced by the differences in the epigenetic status of

chromosomes 17, 19, and X. For these facts, we proposed the following hypothesis

regarding the possible relationships between the differences in epigenome status and

those in expression levels of genes among three AD subgroups.

First, changes in gene expression are strongly influenced by histone modifications in the

promoters and enhancers of each gene but are not significantly influenced by

modifications in other regions (even gene regions) in general. Therefore, it is not

unnatural that differences in the distribution of histone modifications across the genome

of chromosomes 17, 19, and X. Moreover, the distribution of CTCF binding that

controls them did not directly lead to significant differences in the expression of genes

on those chromosomes.

In contrast, changes in the distribution of histone modifications throughout the genome

change the distribution of heterochromatin along the sequence of each genome, which

changes the physical structure and properties of the genome (Komoto et al., 2022;

Fujishiro et al., 2022). Such changes in the structure and properties of each chromosome

are also known to dramatically change the intranuclear location of each genomic region

(Finan et al., 2011; Komoto et al., 2022; Fujishiro et al., 2022). Furthermore, it is known

that such differences in the intranuclear location of chromosomes and local genomic

regions change the gene expression of each chromosome through the formation and

deformation of transcription factories (Iborra et al., 1996; Osborne et al., 2004). Based

on these findings, the differences in the genome-wide distribution of histone

modifications in chromosomes 17, 19, and X among the three AD subgroups induced

different tendencies in the arrangement of each chromosome within the nucleus among

the three AD subgroups. These differences may result in differences in the expression of

genes contained in chromosomes 1, 7, 9, 16, 19, and 22.

To verify this hypothesis, it is necessary to carry out fluorescent imaging of intranuclear

chromosomes in brain cells of various AD patients in the future.

Additionally, the following hypothesis can also be proposed. The histone modifications
and CTCF bindings influence transcript splicing and transcriptional start sites, thus,
contributing to various isoform formations (Alharbi et al., 2021; Awazu et al., 2023;
Podlaha et al., 2014; Zhou et al., 2013). Each isoform may have different functions or
may have opposite effects on the regulation of the expression of other genes
(Yamamoto et al., 2016; Stevens et al., 2019; Awazu et al., 2023). Therefore,
differences in the epigenetic status of chromosomes 17, 19, and X may alter the
proportions of each isoform produced by the genes on these chromosomes, thereby
influencing gene expression on other chromosomes.

However, the transcriptome data analyzed in the present study were obtained from short
read sequencing, so that the estimation of the transcription levels of each isoform had
various difficulties. In the future, long read sequencing should be performed for the
precise estimation of the transcription levels of each isoform.

We also expected that, it may be possible to clarify the relationship between the
differences in gene expression and the differences in the epigenomic state in
chromosomes, such as 17, 19 and X, by examining the differences in each gene
expression and the differences in the epigenetic state of each promoter and enhancer
through a more detailed analysis of the epigenomic profile and gene expression profile.
We consider that such considerations are also an important future challenge.

It is noted that many recent studies performed transcriptome analyses to identify
significant DEGs between healthy and AD individuals. However, by focusing on the
expression patterns of more widespread candidate genes rather than the significant
DEGs, we found that different characteristics appeared in the dorsolateral prefrontal
cortex between healthy and AD individuals and that disease substages occur in response
to changes in the disease state.

Additionally, many studies focused on gene expression patterns in the hippocampus.
One such previous research suggested that multiple states exist in the hippocampus of
patients with AD based on differences in the gene expression profiles (Neff et al.,
2021). Between the hippocampus and prefrontal cortex, as mentioned previously, gene
expression patterns and their variations between healthy individuals and AD patients
differ, and then, different results would be obtained even from the same analysis.

Importantly, the hippocampal data analyzed in previous studies can be analyzed in the
same way as in this study; connections between substages can be extracted and
pathological routes can be estimated, which are crucial issues that need to be addressed
in the future.
703
However, the present study also had some limitations. We classified female individuals
through the analyses of transcriptome and epigenomic data of women in the Rush AD
database. The insights are expected to be useful not only for female patients but also for
understanding the rapid progression of symptoms in male patients. To validate this
expectation, the dataset from male individuals should be analyzed. In this dataset,
however, the number of women patients in the NCI, MCI, and AD groups was 25, 29,
and 28, respectively, whereas that of men was 14, 4, and 7, respectively, which was 3
times smaller than that of female individuals in total (7 times and 4 times smaller than
those on female individuals for MCI and AD). Additionally, the number of men
individuals with MCI and AD in the database (4 and 7) was fewer than that of
individuals in each of the three AD groups obtained for women individuals. With such
little data on male individuals, it was impossible to perform similar analyses on male
individuals or sex-independent analyses. To perform such analyses, obtaining more data
on male patients is one of the essential future issues that must be resolved for
eliminating the potential sex-related bias.
719
To evaluate the differences in RNA-seq data between different groups when searching
for gene sets for characterizing and classifying individuals, the likelihood ratio test
using DESeq2, which is one of the typical methods, and t-test were used. The t-test is
generally better for detecting differences and was expected to identify more candidate
genes in a gene set than the likelihood ratio test or other nonparametric testing methods.
However, the t-test produces more false positives than other testing methods when the
fluctuations in the data deviate from a Gaussian distribution. The gene sets used in this
study were expected to contain few false-positive genes because they were chosen under
imposing the additional condition that the healthy individuals and patients could be
clearly distinguished through cluster analysis. However, the possibility that
inappropriate genes were retained is a valid case. For more precise consideration,
verification is necessary using various nonparametric tests.
732
In addition to the transcriptome of each individual, the database targeted for the present
analysis also contained various epigenome data; however, other information on each

individual, such as postmortem interval, brain PH, and medical history was not
included. Therefore, although this study has yielded unprecedented knowledge, the
validation and characterization of each substage through comparisons with other
physiological data could not be presented. It is possible that such analysis could be
processed using meta-analysis combined with databases focusing on other feature
groups.
741
EXPERIMENTAL PROCEDURES
Data processing
Gene expression data (read count and TPM) from the dorsolateral prefrontal cortex of
NCI, MCI, and AD individuals were acquired from the Rush AD database. In the
present analysis, the data of genes with average expression levels (TPM) in NCI, MCI,
and AD individuals were smaller than 0.1, while those of spike-in-RNAs were
eliminated.
749
Note that the data provided by the Rush AD study in ENCODE does not publish
information on RNA quality such as the postmortem interval of samples and RIN
values. However, any specific alerts regarding data quality according to ENCODE
standards were not given to the RNA-seq data used in this study. Thus, the quality of
the data was assumed to be guaranteed. In fact, the set of genes whose expression
changes between NCI and AD and among NCI, MCI, and AD, involved many genes
obtained from previous studies as AD risk genes. This also supports the lack of data
bias that would contradict previous reports.
758
Statistical analyses
Two-tailed t-tests and ANOVA were performed using Python (Ver.3.9.16). All t-tests in
this study were Welch's t-tests. The false discovery rate (FDR) was estimated through a
set of P-values using the Benjamini–Hochberg procedure (Benjamini et al., 1995)
implemented in MultiPy (https://puolival.github.io/multipy/) (Puoliväli et al., 2020).
The search for DEGs via the likelihood ratio test using DESeq2 and the calculation of
P-values for the Fisher's exact test were performed using R (Ver. 4.1.3).
766
Cluster analyses
Cluster analyses were performed using the values of $X = \log_2 (TPM + 1) - C$, where C
was chosen to satisfy the average of X over individuals = 0 for each gene. iDEP.96
(http://bioinformatics.sdstate.edu/idep96/) was used for hierarchical clustering (GE et

al., 2018). For hierarchical clustering, the group-average method, using the correlation
coefficient as the distance, was used to create a dendrogram. If the cluster formed by the
first branch of hierarchical clustering satisfied the following criteria for a certain group
(NCI, MCI, and AD), then that cluster was regarded as a cluster in which there were
significantly more individuals belonging to that group. These criteria were as follows: i)
containing more than 2/3 of the individuals belonging to that group , and ii) Fisher exact
test resulting in a P-value < 0.05. K-means cluster analysis was performed using the
Python package scikit-learn (Ver.1.0.2) (Pedregosa et al., 2011). The number of clusters
in each k-means cluster analysis was determined using the elbow method.
The construction of UMAP was performed using sc.tl.pca() (Pedregosa et al., 2011)
with default parameters, scanpy.pp.neighbors() with "n_neighbors" = 30 (McInnes et
al., 2018), and the scanpy.tl.umap() functions with n_components = 3 (Wolf et al.,
2019) of scanpy (Ver.1.9.3) according to the literature on basic workflows for clustering
analysis using Scanpy (https://scanpy-tutorials.readthedocs.io/en/latest/pbmc3k.html).
All principal components were used for clustering using the scanpy.pp.neighbors()
function. The obtained results were qualitatively independent of the parameters of these
applications.
788
Epigenomic data analysis
The Jaccard coefficient of the genome-wide distributions of epigenomic markers among
individuals was estimated as follows. The bed formatted files of the narrow peaks of
these epigenome markers and their bam formatted files of the read mapping results were
acquired for individuals whose ChIP-seq data of H3K27ac, H3K27me3, H3K4me3, and
CTCF exist in Rush AD database. The total number of mapped reads was counted from
the bam formatted file using samtools (Ver. 1.20). The correlation between the number
of narrow peaks and the total number of mapped reads for each ChIP-seq data was
evaluated by Pearson correlation coefficient. The Jaccard coefficients between each pair
of individuals were estimated using Bedtools (Ver. 2.31.0). The density of each
epigenomic marker on each chromosome was estimated according to the following
equation: [Total length of genome regions with narrow peak of ChIP-seq data in
chromosome obtained by bed formatted file (bp)]/[Total length of chromosome
obtained by GRCh38 (bp)].
803
804   **Enrichment analysis**
805   Enrichment analysis was performed for the M-NA, N-A1, and N-A2 sets using
806   Metascape (v3.5.20240101) (https://metascape.org/gp/index.html#/main/step1) (Zhou et

807 al., 2019). The biological process terms in Gene Ontology were the main focus of this
808 study.
809

Construction of an adjacency network among substages
The adjacency network among the substages was estimated via PAGA (Wolf et al.,
2019) using the gene expression data of each individual. Each group at each substage
was regarded as a partition in PAGA. PAGA can generate a graph of substages in which
the edge weights represent confidence in the presence of connections. PAGA was
performed using sc.tl.pca() (Pedregosa et al., 2011) with default parameters,
scanpy.pp.neighbors() with "n_neighbors" = 2 (McInnes et al., 2018), and the
scanpy.tl.paga() functions (Wolf et al., 2019) of scanpy (Ver.1.9.3) according to the
literature for trajectory inference using transcriptome data (https://scanpy-
tutorials.readthedocs.io/en/latest/paga-paul15.html). All principal components were
used for clustering using the scanpy.pp.neighbors() function. Additionally, to extract the
essential network among substages, the parameter "n_neighbors" of the function
scanpy.pp.neighbors() was chosen as 2, which is the minimum value in the range of this
parameter. Thus, a network with nonredundant connections among the substages was
expected to be obtained.
825

AUTHOR CONTRIBUTIONS
Kousei Honda and Akinori Awazu conceived and designed the study. Kousei Honda
performed the analyses. Kousei Honda and Akinori Awazu drafted the manuscript.

ACKNOWLEDGEMENTS
This research was supported by JST Moonshot R&D Grant Number JPMJMS2021 and
a JSPS KAKENHI grant (award number 21K06124, 24K09248). Computations were
partially performed on the NIG supercomputer at the ROIS National Institute of
Genetics.

REFERENCES
Adhikarla, S. V., Jha, N. K., Goswami, V. K., Sharma, A., Bhardwaj, A., Dey, A., ... &
Jha, S. K. (2021). TLR-mediated signal transduction and neurodegenerative
disorders. *Brain sciences*, *11*(11), 1373. https://doi.org/10.3390/brainsci11111373

Alharbi, A. B., Schmitz, U., Bailey, C. G., & Rasko, J. E. (2021). CTCF as a regulator
of alternative splicing: new tricks for an old player. *Nucleic acids research*, *49*(14),
7825-7838. https://doi.org/10.1093/nar/gkab520

Alves, F., Kalinowski, P., & Ayton, S. (2023). Accelerated brain volume loss caused by
anti-β-amyloid drugs: a systematic review and meta-analysis. *Neurology*, *100*, e2114–
e2124. https://doi.org/10.1212/wnl.0000000000207156

Anders, S., & Huber, W. (2010). Differential expression analysis for sequence count
data. *Genome Biology*, *11*, R106. https://doi.org/10.1186/gb-2010-11-10-r106

Arriagada, P. V., Growdon, J. H., Hedley-Whyte, E. T., & Hyman, B. T. (1992).
Neurofibrillary tangles but not senile plaques parallel duration and severity of
Alzheimer's disease. *Neurology*, *42*, 631–631. https://doi.org/10.1212/wnl.42.3.631

Awazu, A., Takemoto, D., Watanabe, K., & Sakamoto, N. (2023). Possibilities of skin
coat color-dependent risks and risk factors of squamous cell carcinoma and deafness of
domestic cats inferred via RNA-seq data. *Genes to Cells*, *28*(12), 893-905.
https://doi.org/10.1111/gtc.13076

Bartzokis, G. (2004). Age-related myelin breakdown: a developmental model of
cognitive decline and Alzheimer's disease. *Neurobiology of Aging*, *25*, 5–18.
https://doi.org/10.1016/j.neurobiolaging.2003.03.001

Bartzokis, G. (2011). Alzheimer's disease as homeostatic responses to age-related
myelin breakdown. *Neurobiology of Aging*, *32*, 1341–1371.
https://doi.org/10.1016/j.neurobiolaging.2009.08.007

Bellenguez, C., Küçükali, F., Jansen, I. E., Kleineidam, L., Amin, N., Naj, A. C., ...
Lambert, J. (2022). New insights into the genetic etiology of Alzheimer's disease and
related dementias. *Nature Genetics*, *54*, 412–436. https://doi.org/10.1038/s41588-022-
01024-z

Benjamini, Y., Hochberg, Y. (1995). Controlling the false discovery rate: a practical
and powerful approach to multiple testing. *Journal of the Royal Statistics Society Series*
B, *57*, 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Chakravarthy, B., Gaudet, C., Ménard, M., Brown, L., Atkinson, T., LaFerla, F. M.,... &
Whitfield, J. (2012). Reduction of the immunostainable length of the hippocampal
dentate granule cells' primary cilia in 3xAD-transgenic mice producing human Aβ1-42
and tau. *Biochemical and Biophysical Research Communications*, *427*(1), 218–222.
https://doi.org/10.1016/j.bbrc.2012.09.056

Cheng, Y., Saville, L., Gollen, B., Veronesi, A. A., Mohajerani, M., Joseph, J. T., &
Zovoilis, A. (2021). Increased Alu RNA processing in Alzheimer brains is linked to
gene expression changes. *EMBO Reports*, *22*, e52255.
https://doi.org/10.15252/embr.202052255

Chen, Z., & Zhong, C. (2013). Decoding Alzheimer's disease from perturbed cerebral
glucose metabolism: implications for diagnostic and therapeutic strategies. *Progress in*
neurobiology, *108*, 21-43. https://doi.org/10.1016/j.pneurobio.2013.06.004

Cunnane, S., Nugent, S., Roy, M., Courchesne-Loyer, A., Croteau, E., Tremblay, S., ...
& Rapoport, S. I. (2011). Brain fuel metabolism, aging, and Alzheimer's
disease. *Nutrition*, *27*(1), 3-20. https://doi.org/10.1016/j.nut.2010.07.021

Davis, E. J., Broestl, L., Abdulai-Saiku, S., Worden, K., Bonham, L. W., Miñones-
Moyano, E., ... & Dubal, D. B. (2020). A second X chromosome contributes to
resilience in a mouse model of Alzheimer's disease. *Science translational*
medicine, *12*(558), eaaz5677. https://doi.org/aaz5677

Duara, R., & Barker, W. (2021). Heterogeneity in Alzheimer's disease diagnosis and
progression rates: implications for therapeutic trials. *Neurotherapeutics*, *19*, 8–25.
https://doi.org/10.1007/s13311-022-01185-z

Esteve, D., Molina-Navarro, M. M., Giraldo, E., Martínez-Varea, N., Blanco-Gandia,
M. C., Rodríguez-Arias, M., ... & Lloret, A. (2022). Adult neural stem cell migration is
impaired in a mouse model of Alzheimer's disease. *Molecular Neurobiology*, 1-15.
https://doi.org/10.1007/s12035-021-02620-6

Finan, K., Cook, P. R., & Marenduzzo, D. (2011). Non-specific (entropic) forces as
major determinants of the structure of mammalian chromosomes. *Chromosome*
research, *19*, 53-61. https://doi.org/10.1007/s10577-010-9150-y

Fujishiro, S., & Sasai, M. (2022). Generation of dynamic three-dimensional genome
structure through phase separation of chromatin. *Proceedings of the National Academy*
of Sciences, *119*(22), e2109838119. https://doi.org/10.1073/pnas.2109838119

Gatz, M., Reynolds, C. A., Fratiglioni, L., Johansson, B., Mortimer, J. A., Berg, S.,... &
Pedersen, N. L. (2006). Role of genes and environments for explaining Alzheimer
disease. *Archives of General Psychiatry*, *63*, 168–174.
https://doi.org/10.1001/archpsyc.63.2.168

Ge, S.X., Son, E.W., & Yao, R. (2018). iDEP: an integrated web application for
differential expression and pathway analysis of RNA-Seq data. *BMC Bioinformatics*,
19, 534. https://doi.org/10.1186/s12859-018-2486-6

Grimm, M. O., Mett, J., & Hartmann, T. (2016). The impact of vitamin E and other fat-
soluble vitamins on Alzheimer s disease. *International journal of molecular*
sciences, *17*(11), 1785. https://doi.org/10.3390/ijms17111785

Hardy, J. A., & Higgins, G. A. (1992). Alzheimer's disease: the amyloid cascade
hypothesis. *Science*, *256*, 184–185. https://doi.org/10.1126/science.1566067

Hardy, J., & Selkoe, D. J. (2002). The amyloid hypothesis of Alzheimer's disease:
progress and problems on the road to therapeutics. *Science*, *297*, 353–356.
https://doi.org/10.1126/science.1072994

Higginbotham, L., Ping, L., Dammer, E. B., Duong, D. M., Zhou, M., Gearing, M., …
Seyfried, N. T. (2020). Integrated proteomics reveals brain-based cerebrospinal fluid
biomarkers in asymptomatic and symptomatic Alzheimer's disease. *Science Advances*,
6, eaaz9360. https://doi.org/10.1126/sciadv.aaz9360

Hirai, K., Aliev, G., Nunomura, A., Fujioka, H., Russell, R. L., Atwood, C. S.,... Smith,
M. A. (2001). Mitochondrial abnormalities in Alzheimer's disease. *Journal of*
Neuroscience, *21*, 3017–3023. https://doi.org/10.1523/jneurosci.21-09-03017.2001

Iborra, F. J., Pombo, A., Jackson, D. A., & Cook, P. R. (1996). Active RNA

polymerases are localized within discrete transcription 'factories' in human

nuclei. *Journal of cell science*, *109*(6), 1427-1436.

https://doi.org/10.1242/jcs.109.6.1427

Itzhaki, R. F., Lin, W. R., Shang, D., Wilcock, G. K., Faragher, B., & Jamieson, G. A.

(1997). Herpes simplex virus type 1 in brain and risk of Alzheimer's disease. *The*

Lancet, *349*, 241–244. https://doi.org/10.1016/s0140-6736(96)10149-5

Jansen, I. E., Savage, J. E., Watanabe, K., Bryois, J., Williams, D. M., Steinberg, S.,...

Posthuma, D. (2019). Genome-wide meta-analysis identifies new loci and functional

pathways influencing Alzheimer's disease risk. *Nature Genetics*, *51*, 404–413.

https://doi.org/10.1038/s41588-018-0311-9

Jin, M., Shepardson, N., Yang, T., Chen, G., Walsh, D., & Selkoe, D. J. (2011). Soluble

amyloid β-protein dimers isolated from Alzheimer cortex directly induce Tau

hyperphosphorylation and neuritic degeneration. *Proceedings of the National Academy*

of Sciences, *108*, 5819–5824. https://doi.org/10.1073/pnas.1017033108

Johnson, E. C., Carter, E. K., Dammer, E. B., Duong, D. M., Gerasimov, E. S., Liu,

Y.,... Seyfried, N. T. (2022). Large-scale deep multi-layer analysis of Alzheimer's

disease brain reveals strong proteomic disease-related changes not observed at the RNA

level. *Nature Neuroscience*, *25*, 213–225. https://doi.org/10.1038/s41593-021-00999-y

Kaltschmidt, B., & Kaltschmidt, C. (2015). NF-KappaB in long-term memory and

structural plasticity in the adult mammalian brain. *Frontiers in molecular*

neuroscience, *8*, 69. https://doi.org/10.3389/fnmol.2015.00069

Kandimalla, R., Thirumala, V., & Reddy, P. H. (2017). Is Alzheimer's disease a type 3

diabetes? A critical appraisal. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of*

Disease, *1863*, 1078–1089. https://doi.org/10.1016/j.bbadis.2016.08.018

Komarova, N. L., & Thalhauser, C. J. (2011). High degree of heterogeneity in

Alzheimer's disease progression patterns. *PLoS Computational Biology*, *7*, e1002251.

https://doi.org/10.1371/journal.pcbi.1002251

Komoto, T., Fujii, M., & Awazu, A. (2022). Epigenetic-structural changes in X
chromosomes promote Xic pairing during early differentiation of mouse embryonic
stem cells. *Biophysics and Physicobiology*, *19*, e190018.
https://doi.org/10.2142/biophysico.bppb-v19.0018

Lau, S. F., Wu, W., Wong, H. Y., Ouyang, L., Qiao, Y., Xu, J., ... & Ip, N. Y. (2023).
The VCAM1–ApoE pathway directs microglial chemotaxis and alleviates Alzheimer's
disease pathology. *Nature Aging*, *3*(10), 1219-1236.
https://doi.org/10.1038/s43587-023-00491-1

Lin, X., Kapoor, A., Gu, Y., Chow, M. J., Peng, J., Zhao, K., & Tang, D. (2020).
Contributions of DNA damage to Alzheimer's disease. *International Journal of*
Molecular Sciences, *21*(5), 1666. https://doi.org/10.3390/ijms21051666

Lopez-Lee, C., Torres, E. R. S., Carling, G., & Gan, L. (2024). Mechanisms of sex
differences in Alzheimer's disease. *Neuron*.
https://doi.org/10.1016/j.neuron.2024.01.024

1002

1003 Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
1004 dispersion for RNA-seq data with DESeq2. *Genome biology*, *15*, 1-21.
1005 https://doi.org/10.1186/s13059-014-0550-8

1006

1007 Luo, Y., Hitz, B. C., Gabdank, I., Hilton, J. A., Kagda, M. S., Lam, B.,... Cherry, J. M.
1008 (2020). New developments on the Encyclopedia of DNA Elements (ENCODE) data
1009 portal. *Nucleic Acids Research*, 48, D882–D889. https://doi.org/10.1093/nar/gkz1062

1010

1011 Macciardi, F., Giulia Bacalini, M., Miramontes, R., Boattini, A., Taccioli, C., Modenini,
1012 G.,... Mapstone, M. (2022). A retrotransposon storm marks clinical phenoconversion to
1013 late-onset Alzheimer's disease. *GeroScience*, *44*, 1525–1550.
1014 https://doi.org/10.1007/s11357-022-00580-w

1015

1016 McGeer, P. L., & McGeer, E. G. (1995). The inflammatory response system of brain:
1017 implications for therapy of Alzheimer and other neurodegenerative diseases. *Brain*
1018 *Research Reviews*, *21*, 195–218. https://doi.org/10.1016/0165-0173(95)00011-9

1019

McInnes, L., Healy, J., & Melville, J. UMAP: Uniform manifold approximation and
projection for dimension reduction. arXiv:1802.03426.
https://doi.org/10.48550/arXiv.1802.03426

Morelli, A., Sarchielli, E., Guarnieri, G., Coppi, E., Pantano, D., Comeglio, P.,...
Vannelli, G. B. (2017). Young human cholinergic neurons respond to physiological
regulators and improve cognitive symptoms in an animal model of Alzheimer's disease.
Frontiers in Cellular Neuroscience, *11*, 339. https://doi.org/10.3389/fncel.2017.00339

Neff, R. A., Wang, M., Vatansever, S., Guo, L., Ming, C., Wang, Q.,... Zhang, B.
(2021). Molecular subtyping of Alzheimer's disease using RNA sequencing data reveals
novel mechanisms and targets. *Science Advances*, *7*, eabb5398.
https://doi.org/10.1126/sciadv.abb5398

Nunomura, A., Perry, G., Aliev, G., Hirai, K., Takeda, A., Balraj, E. K.,... Smith, M. A.
(2001). Oxidative damage is the earliest event in Alzheimer disease. *Journal of*
Neuropathology & Experimental Neurology, *60*, 759–767.
https://doi.org/10.1093/jnen/60.8.759

Osaka, R., Kobayashi, N., Shimada, K., Ishii, A., Oka, N., & Kondo, K. (2022). VP26, a
herpes simplex virus type 1 capsid protein, increases DNA methylation in COASY
promoter region. *Brain, Behavior, & Immunity - Health*, *26*, 100545.
https://doi.org/10.1016/j.bbih.2022.100545

Osborne, C. S., Chakalova, L., Brown, K. E., Carter, D., Horton, A., Debrand, E., ... &
Fraser, P. (2004). Active genes dynamically colocalize to shared sites of ongoing
transcription. *Nature genetics*, *36*(10), 1065-1071. https://doi.org/10.1038/ng1423

Ott, A., Stolk, R. P., van Harskamp, F., Pols, H. A. P., Hofman, A., & Breteler, M. M.
B. (1999). Diabetes mellitus and the risk of dementia: The Rotterdam
Study. *Neurology*, *53*, 1937–1937. https://doi.org/10.1212/wnl.53.9.1937

Parachikova, A., Agadjanyan, M. G., Cribbs, D. H., Blurton-Jones, M., Perreau, V.,
Rogers, J., ... & Cotman, C. W. (2007). Inflammatory changes parallel the early stages
of Alzheimer disease. *Neurobiology of aging*, *28*(12), 1821-1833.
https://doi.org/10.1016/j.neurobiolaging.2006.08.014

Patel, P. J., Ren, Y., & Yan, Z. (2023). Epigenomic analysis of Alzheimer's disease brains reveals diminished CTCF binding on genes involved in synaptic organization. *Neurobiology of Disease*, 184, 106192. https://doi.org/10.1016/j.nbd.2023.106192

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., … Duchesnay, É. (2011). Scikit-learn: machine learning in Python. *JMLR*, *12*, 2825–2830.

Plácido, A. I., Pereira, C. M. F., Duarte, A. I., Candeias, E., Correia, S. C., Santos, R. X.,... Moreira, P. I. (2014). The role of endoplasmic reticulum in amyloid precursor protein processing and trafficking: implications for Alzheimer's disease. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, *1842*, 1444–1453. https://doi.org/10.1016/j.bbadis.2014.05.003

Podlaha, O., De, S., Gonen, M., & Michor, F. (2014). Histone modifications are associated with transcript isoform diversity in normal and cancer cells. *PLoS computational biology*, *10*(6), e1003611. https://doi.org/10.1371/journal.pcbi.1003611

Puoliväli, T., Palva, S., & Palva, J.M. (2020). Influence of multiple hypothesis testing on reproducibility in neuroimaging research: A simulation study and Python-based software. *Journal of Neuroscience Methods*, *337*, 108654. https://doi.org/10.1016/j.jneumeth.2020.108654

Ranganathan, R., Sapozhnikov, G., Ni, W., Li, S., & Song, Y. (2023). Recent developments in the role of DNA damage response and understanding its implications for new therapeutic approaches in Alzheimer's disease. *Translational Medicine of Aging*. https://doi.org/10.1016/j.tma.2023.06.004

Ravel-Godreuil, C., Znaidi, R., Bonnifet, T., Joshi, R. L., & Fuchs, J. (2021). Transposable elements as new players in neurodegenerative diseases. *FEBS Letters*, *595*, 2733–2755. https://doi.org/10.1002/1873-3468.14205

Readhead, B., Haure-Mirande, J. V., Funk, C. C., Richards, M. A., Shannon, P., Haroutunian, V.,... Dudley, J. T. (2018). Multiscale analysis of independent Alzheimer's cohorts finds disruption of molecular, genetic, and clinical networks by human herpesvirus. *Neuron*, *99*, 64–82. https://doi.org/10.1016/j.neuron.2018.05.023

1092

1093 Rogers, J., & Lue, L. F. (2001). Microglial chemotaxis, activation, and phagocytosis of

1094 amyloid β-peptide as linked phenomena in Alzheimer's disease. *Neurochemistry*

1095 *international*, *39*(5-6), 333-340. https://doi.org/10.1016/S0197-0186(01)00040-7

1096

1097 Schuler, G. D., Epstein, J. A., Ohkawa, H., & Kans, J. A. (1996). Entrez: Molecular

1098 biology database and retrieval system. In Methods in enzymology (Vol. 266, pp. 141-

1099 162). Academic Press. https://doi.org/10.1016/S0076-6879(96)66012-1

1100

1101 Sekiya, M., Wang, M., Fujisaki, N., Sakakibara, Y., Quan, X., Ehrlich, M. E.,... Iijima,

1102 K. M. (2018). Integrated biology approach reveals molecular and pathological

1103 interactions among Alzheimer's Aβ42, Tau, TREM2, and TYROBP in Drosophila

1104 models. *Genome Medicine*, *10*(1), 1–20. https://doi.org/10.1186/s13073-018-0530-9

1105

1106 Selkoe, D. J., & Hardy, J. (2016). The amyloid hypothesis of Alzheimer's disease at 25

1107 years. *EMBO Molecular Medicine*, *8*(6), 595–608.

1108 https://doi.org/10.15252/emmm.201606210

1109

1110 Singh, S., Sahu, K., Singh, C., & Singh, A. (2022). Lipopolysaccharide induced altered

1111 signaling pathways in various neurological disorders. *Naunyn-Schmiedeberg's Archives*

1112 *of Pharmacology*, *395*(3), 285-294. https://doi.org/10.1007/s00210-021-02198-9

1113

1114 Stevens, M., & Oltean, S. (2019). Modulation of the apoptosis gene Bcl-x function

1115 through alternative splicing. *Frontiers in Genetics*, *10*, 804.

1116 https://doi.org/10.3389/fgene.2019.00804

1117

1118 Strittmatter, W. J., Saunders, A. M., Schmechel, D., Pericak-Vance, M., Enghild, J.,

1119 Salvesen, G. S., & Roses, A. D. (1993). Apolipoprotein E: high-avidity binding to beta-

1120 amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer

1121 disease. *Proceedings of the National Academy of Sciences*, *90*, 1977–1981.

1122 https://doi.org/10.1073/pnas.90.5.1977

1123

1124 Suresh, S., Singh, A., Rushendran, R., Vellapandian, C., & Prajapati, B. (2023).

1125 Alzheimer's disease: the role of extrinsic factors in its development, an investigation of

1126 the environmental enigma. *Frontiers in Neurology*, *14*, 1303111.

1127 https://doi.org/10.3389/fneur.2023.1303111

1128

1129 Tijms, B. M., Gobom, J., Reus, L., Jansen, I., Hong, S., Dobricic, V.,... Visser, P. J.

1130 (2020). Pathophysiological subtypes of Alzheimer's disease based on cerebrospinal

1131 fluid proteomics. *Brain*, *143*, 3776–3792. https://doi.org/10.1093/brain/awaa325

1132

1133 Tijms, B. M., Gobom, J., Teunissen, C., Dobricic, V., Tsolaki, M., Verhey, F.,... Visser,

1134 P. J. (2021). CSF proteomic Alzheimer's disease-predictive subtypes in cognitively

1135 intact amyloid negative individuals. *Proteomes*, *9*, 36.

1136 https://doi.org/10.3390/proteomes9030036

1137

1138 Vorobyeva, A. G., & Saunders, A. J. (2018). Amyloid-β interrupts canonical Sonic

1139 hedgehog signaling by distorting primary cilia structure. *Cilia*, *7*(1), 1–11.

1140 https://doi.org/10.1186/s13630-018-0059-y

1141

1142 Wang, L., Davis, P. B., Volkow, N. D., Berger, N. A., Kaelber, D. C., & Xu, R. (2022).

1143 Association of COVID-19 with new-onset Alzheimer's disease. *Journal of Alzheimer's*

1144 *Disease*, *89*(2), 1–4. https://doi.org/10.3233/jad-220717

1145

1146 Wang, X., Wang, W., Li, L., Perry, G., Lee, H. G., & Zhu, X. (2014). Oxidative stress

1147 and mitochondrial dysfunction in Alzheimer's disease. *Biochimica et Biophysica Acta*

1148 *(BBA)-Molecular Basis of Disease*, *1842*, 1240–1247.

1149 https://doi.org/10.1016%2Fj.bbadis.2013.10.015

1150

1151 Wightman, D. P., Jansen, I. E., Savage, J. E., Shadrin, A. A., Bahrami, S., Holland,

1152 D.,... Posthuma, D. (2021). A genome-wide association study with 1,126,563

1153 individuals identifies new risk loci for Alzheimer's disease. *Nature Genetics*, *53*, 1276–

1154 1282. https://doi.org/10.1038%2Fs41588-021-00921-z

1155

1156 Wu, L., Xian, X., Xu, G., Tan, Z., Dong, F., Zhang, M., & Zhang, F. (2022). Toll-like

1157 receptor 4: a promising therapeutic target for alzheimer's disease. *Mediators of*

1158 *inflammation*, *2022*. https://doi.org/10.1155/2022/7924199

1159

1160 Wolf, F. A., Hamey, F. K., Plass, M., Solana, J., Dahlin, J. S., Göttgens, B.,... Theis, F.

1161 J. (2019). PAGA: graph abstraction reconciles clustering with trajectory inference

1162 through a topology preserving map of single cells. *Genome Biology*, *20*, 1–9.

1163 https://doi.org/10.1186%2Fs13059-019-1663-x

1164

1165 World Health Organization. (2017). Global action plan on the public health response to
1166 dementia 2017–2025.

1167

1168 Yamamoto, H., Rundqvist, H., Branco, C., & Johnson, R. S. (2016). Autocrine VEGF
1169 isoforms differentially regulate endothelial cell behavior. *Frontiers in cell and*
1170 *developmental biology*, *4*, 99. https://doi.org/10.3389/fcell.2016.00099

1171

1172 Younas, A., Younas, N., Iqbal, M. J., Ferrer, I., & Zerr, I. (2024). Comparative
1173 interactome mapping of Tau-protein in classical and rapidly progressive Alzheimer's
1174 disease identifies subtype-specific pathways. *Neuropathology and Applied*
1175 *Neurobiology*, *50*, e12964. https://doi.org/10.1111/nan.12964

1176

1177 Younas, N., Zafar, S., Shafiq, M., Noor, A., Siegert, A., Arora, A. S., … Zerr, I. (2020).
1178 SFPQ and Tau: Critical factors contributing to rapid progression of Alzheimer's
1179 disease. *Acta Neuropathologica*, *140*, 317–339.
1180 https://doi.org/10.1007/s00401-020-02178-y

1181

1182 Young-Pearse, T. L., Bai, J., Chang, R., Zheng, J. B., LoTurco, J. J., & Selkoe, D. J.
1183 (2007). A critical function for β-amyloid precursor protein in neuronal migration
1184 revealed by in utero RNA interference. *Journal of Neuroscience*, *27*(52), 14459-14469.
1185 https://doi.org/10.1523/JNEUROSCI.4701-07.2007

1186

1187 Yuen, S. C., Zhu, H., & Leung, S. W. (2020). A systematic bioinformatics workflow
1188 with meta-analytics identified potential pathogenic factors of Alzheimer's disease.
1189 *Frontiers in Neuroscience*, *14*, 209. https://doi.org/10.3389/fnins.2020.00209

1190

1191 Zhan, X., Stamova, B., & Sharp, F. R. (2018). Lipopolysaccharide associates with
1192 amyloid plaques, neurons and oligodendrocytes in Alzheimer's disease brain: a
1193 review. *Frontiers in aging neuroscience*, *10*, 42.
1194 https://doi.org/10.3389/fnagi.2018.00042

1195

1196 Zhou, H. L., Luo, G., Wise, J. A., & Lou, H. (2013). Regulation of alternative splicing
1197 by local histone modifications: potential roles for RNA-guided mechanisms. *Nucleic*
1198 *acids research*, *42*(2), 701-713. https://doi.org/10.1093/nar/gkt875

1199
1200   Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., …

1201   Chanda, S. K. (2019). Metascape provides a biologist-oriented resource for the analysis

1202   of systems-level datasets. *Nature Communications*, *10*(1),1523.

1203   https://doi.org/10.1038/s41467-019-09234-6

1204

1205

1206 **Table 1**. Summary of the individuals included in this study (see Table S1a and S1b for
1207 more details).

Clinical stage	No Cognitive Impairment (NCI)	Mild Cognitive Impairment (MCI)	Alzheimer's Disease (AD)
# of individuals	25	29	28
Individual's index	N1 – N25	M1 – M29	A1 – A28

1208

Table 2. Number of individuals exhibiting typical genome-wide distributions of each
epigenetic state, E_T, and that exhibiting the untypical one, E_U, in the three AD substages
(AD_{HH}, AD_{LL}, and AD_{HL}); H3K27ac (a), H3K27me3 (b), CTCF (c), and H3K4me3 (d).
The E_T and E_U in each substage were obtained from Figures 5, 6, and S4. P-values
obtained using the Fisher's exact test for each table were 0.04328 (a), 0.04299 (b),
0.04328 (c), and 0.1829 (d).

1215

1216 (a)

H3K27ac	AD_{HH}	AD_{LL}	AD_{HL}
# of individuals with typical (E_T)	6	4	3
# of individuals with untypical (E_U)	0	0	4

1217

1218 (b)

H3K27me3	AD_{HH}	AD_{LL}	AD_{HL}
# of individuals with typical (E_T)	5	4	2
# of individuals with untypical (E_U)	1	0	5

1219

1220 (c)

CTCF	AD_{HH}	AD_{LL}	AD_{HL}
# of individuals with typical (E_T)	6	4	3
# of individuals with untypical (E_U)	0	0	4

1221

1222 (d)

H3K4me3	AD_{HH}	AD_{LL}	AD_{HL}
# of individuals with typical (E_T)	2	2	6
# of individuals with untypical (E_U)	4	2	1

1223

1224

(a)

1225

(b)

(c)

1226

Fig. 1. Re

obtained

case of the

NA set of map

of the expression levels of each gene was described using $X = \log_2(\text{TPM} + 1) - C$,

where C was chosen to satisfy the average of X over individuals = 0 for each gene. The

$\text{MCI}_{\text{Major}}$ individuals occupy the right region in the dendrogram. N_G and N_{kG} were the

total number of genes and number of genes annotated as protein coding or noncoding

RNA genes with higher (UP) and lower (DOWN) expression levels in the $\text{MCI}_{\text{Major}}$

individuals, respectively. Scatter plot of individuals in 3-dimensional space constructed

using U-MAP 1, 2, and 3 axes shown from various directions (c).

1238

(a)

1239

(b)

N-A1 set (P^D_{NA} = 0.05, P^D_{AN} = 0.02, P^t_{NA} = 0.08, P^t_{AN} = 0.10)

■ NCI ■ AD

1240

(c)

N-A2 set (P^D_{NA} = 0.03, P^D_{AN} = 0.03, P^t_{NA} = 0.08, P^t_{AN} = 0.10)

■ NCI ■ AD

1241

Fig. 2 Phase diagrams of the hierarchical clustering of NCI and AD individuals,

and the results of the hierarchical clustering of individuals of the N-A1 and N-A2

sets. Phase diagram of the results of hierarchical clustering using genes where $p \leq P^D_{NA}$

and $p \leq P^D_{AN}$ (a, left), and that using genes where $p \leq P^t_{NA}$ and $p \leq P^t_{AN}$ (a, right).

Circles indicate where NCI and AD individuals were completely divided into different

clusters at the first branch of the dendrogram. The dendrograms obtained using the N-

A1 (b) and N-A2 sets (c) are shown. Broken lines indicate the boundary between the

AD_{HL}, AD_{LL}, and AD_{HH}.

1250

 Fig. 3 Results of the hierarchical clustering of individuals of the N-A1 and N-A2

 sets and expression level of each gene. The dendrograms and expression level of each

 gene in the N-A1 set of each individual (a) (see Table S3d for more details) and that in

 the N-A2 set of each individual was plotted (b) (See Table S3e for more details). The

 heatmap of the expression levels of each gene was described using $X = \log_2(TPM + 1) -$

 C, where C was chosen to satisfy the average of X over individuals = 0 for each gene.

 Broken lines indicate the boundary between the AD_{HL}, AD_{LL}, and AD_{HH}.

Chromosomal distribution of genes belonging to cluster CA1 or CB1 (%)

Chromosomal distribution of genes belonging to cluster CA2 or CB2 (%)

Chromosomal distribution of coding and non-coding genes (%)

1260

1261 **Fig. 4** Chromosomal distribution of genes belonging to the cluster CA1 or CB1 (Fig.

1262 3a), and that to the cluster CA2 or CB2 (Fig. 3b). Chromosomal distribution of coding

1263 and non-coding genes was estimated by human genome data GRCh38.

1264

1265

Fig. 5 Jaccard coefficient of the genome-wide distributions of H3K27ac among

individuals. Individuals were divided into two groups at the first divergence of the

1269   dendrogram. The epigenetic state exhibited by individuals belonging to the group

1270   containing a larger number of individuals was regarded as the "typical" epigenetic state,

1271   whereas that containing fewer individuals was regarded as the "untypical" epigenetic

1272   state. Dashed lines indicate the boundaries between groups.

1273

H3K4me3

1276 **Fig. 6** Jaccard coefficient of the genome-wide distributions of H3K4me3 among

1277 individuals. Each object in this figure have the same meaning as that in Fig. 5.

1279

1280

1281 **Fig. 7 Adjacency network among substages.** The network constructed using common
1282 connections obtained from both the network constructed using the M-NA and N-A1 sets
1283 (Fig S5a) and that constructed using the M-NA and N-A2 sets (Fig. S5b). The width of
1284 each edge indicates the confidence in the presence of a connection between each pair of
1285 connected nodes.
1286

1287

(a)

		# of genes					
		77	145	211	286	360	
		0.01	0.02	0.03	0.04	0.05	P^D_{MN}
70	0.01	0, 5	0, 5	0, 5	0, 5	0, 5	
147	0.02	1, 5	0, 5	0, 5	0, 5	2, 5	
224	0.03	0, 5	0, 3	0, 5	0, 5	0, 5	
317	0.04	4, 5	1, 5	0, 5	0, 5	0, 5	
414	0.05	×	0, 5	0, 5	0, 5	0, 5	

# of genes (vertical axis), P^D_{NM}

		# of genes					
		49	95	158	219	286	
		0.01	0.02	0.03	0.04	0.05	P^t_{MN}
78	0.01	1, 3	0, 3	0, 3	1, 4	0, 5	
164	0.02	1, 4	2, 2	0, 2	1, 5	0, 5	
271	0.03	1, 4	0, 6	1, 2	0, 6	2, 3	
392	0.04	1, 4	0, 5	0, 4	0, 8	0, 6	
504	0.05	0, 8	0, 7	0, 6	0, 5	0, 5	

# of genes (vertical axis), P^t_{NM}

1288

1289

(b)

		# of genes					
		131	256	388	512	638	
		0.01	0.02	0.03	0.04	0.05	P^D_{AM}
138	0.01	3, 5	4, 3	5, 0	2, 9	3, 7	
247	0.02	5, 0	5, 0	4, 1	6, 0	2, 5	
372	0.03	5, 0	5, 0	6, 0	7, 0	3, 6	
478	0.04	5, 9	4, 0	5, 0	5, 0	5, 2	
593	0.05	×	4, 0	2, 7	4, 0	5, 0	

# of genes (vertical axis), P^D_{MA}

		# of genes					
		65	140	216	307	389	
		0.01	0.02	0.03	0.04	0.05	P^t_{AM}
92	0.01	4, 0	3, 3	4, 2	4, 2	5, 2	
183	0.02	5, 1	4, 1	5, 0	5, 0	4, 1	
302	0.03	3, 2	3, 2	4, 0	3, 0	4, 0	
419	0.04	2, 2	3, 2	4, 0	3, 0	3, 0	
526	0.05	×	3, 1	5, 0	5, 0	4, 0	

# of genes (vertical axis), P^t_{MA}

1290

(c)

1291

Fig. 8 Detailed classification of the MCI individuals. Number of NCI individuals who belong to the MCI cluster (left value) and that of MCI individuals who belong to the NCI cluster (right value) as a function of P^D_{NM} and P^D_{MN} (a, left), and as a function of P^t_{NM} and P^t_{MN} (a, right). Number of MCI individuals who belong to the AD cluster (left value) and that of AD individuals who belong to the MCI cluster (right value) as a

function of P^D_{MA} and P^D_{AM} (b, left), and as a function of P^t_{MA} and P^t_{AM} (b, right).

Number of deviations of each MCI individual from the MCI to the NCI (c, upper left)

and to AD (c, lower left) cluster. Broken lines indicate boundaries between the

MCI_{NCI_AD} and MCI_{Major} individuals. Scatter plot of the MCI_{Major} individuals as a

function of the number of times that each individual belonged to the NCI and AD

cluster (c, right). Using k-means clustering (Fig. S6c), the MCI_{Major} individuals were

divided into $MCI_{Typical}$, MCI_{NCI}, and MCI_{AD}.

1304

1305

1306

1307 **Fig. 9 Detailed classification of the NCI individuals.** Number of deviations of each

1308 NCI individual from the NCI to the MCI cluster (a), and that of each AD individual

1309 from the AD to the MCI cluster (b). Broken lines indicate boundaries among clusters of

1310 individuals obtained via k-means clustering (Fig. S7a and S7b).

1311

1312

1313

1314 **Fig. 10 Adjacency networks among substages.** A network containing common

1315 connections from both networks obtained using the data in the M-NA and N-A1 sets

1316 (Fig. S8a) and that in the M-NA and N-A2 sets (Fig. S8b). The width of each edge

1317 indicates the confidence in the presence of a connection between each pair of connected

1318 nodes. Arrows indicated the inferred direction of disease progression.