Potential multiple disease progression pathways in female patients with

Alzheimer's disease inferred from transcriptome and epigenome data of the

dorsolateral prefrontal cortex

Short title: Progression pathways of AD

Kousei Honda[1], Akinori Awazu[1,2,*]

[1] Graduate School of Integrated Sciences for Life, Hiroshima University,

Higashihiroshima, Hiroshima 739-8526, Japan

[2] Research Center for the Mathematics on Chromatin Live Dynamics, Hiroshima

University, Higashihiroshima, Hiroshima 739-8526, Japan

* Send correspondence to: Akinori Awazu; Email: awa@hiroshima-u.ac.jp

Keywords: Alzheimer's disease, transcriptome, epigenome

ABSTRACT

Late-onset Alzheimer's disease (AD) is a typical type of dementia for which therapeutic strategies have not yet been established. The database of the Rush Alzheimer's Disease study by the ENCODE consortium contains transcriptome and various epigenome data. Although the Rush AD database may contain a satisfactory amount of data for women, the amount of data for men remains insufficient. Here, based on an analysis of publicly available data from female patients, this study found that AD pathology appears to be nonuniform; several different typical forms of cognitive deterioration were observed. First, cluster analysis was performed on individuals diagnosed with "No Cognitive Impairment (NCI)," "Mild Cognitive Impairment (MCI)," and "Alzheimer's Disease (AD)" stages in clinical trials using gene expression. NCI was found to contain at least two substages, whereas MCI and AD contained four and six substages, respectively. The epigenome data, in particular genome-wide H3k4me3 distribution data, also supported the existence of multiple AD substages. However, *APOE* gene polymorphisms of individuals seemed to not correlate with disease stage. An inference of adjacency networks among substages, evaluated via partition-based graph abstraction using the gene expression profiles of individuals, suggested the possibility of multiple typical disease progression pathways from NCI to different AD substages through various MCI substages. These results will aid future research on the pathological features and mechanisms of AD and may induce changes in the strategic bases of conventional AD prevention and treatment.

INTRODUCTION

Alzheimer's disease (AD) is the most common cause of dementia, accounting for 60–70 % of dementia cases in older adults [1, 2]. AD can be classified into the familial type, which is strongly influenced by genetic factors, and the late-onset (sporadic) type, which is strongly influenced by aging and environmental factors, with the latter accounting for the majority of AD cases [3]. No treatment or preventive methods for AD have yet been established because the fundamental pathogenic mechanism of sporadic AD remains unclear.

Plaques of abnormally aggregated amyloid β (Aβ) are often observed in the brains of patients with AD and are considered to be related to the onset and progression of the disease [4-7]. However, cases have been reported in which AD developed without the accumulation of Aβ [8] and inhibiting Aβ production even worsened symptoms

(Francesca et al., 2023). Factors other than Aβ plaque formation, such as the abnormal

aggregation of tau protein [9], the expression of apolipoprotein E variant APOE4 [10],

herpes virus and coronaviruses (including SARS-CoV2) infections [11-14], diabetes

[15, 16], brain inflammation [17], oxidative damage in cells [18-20], endoplasmic

reticulum and ciliary dysplasia [21-24], myelin dysplasia [25, 26], and retrotransposons

activities [27-29] have also been reported to influence the onset and progression of AD.

Owing to the involvement of the aforementioned factors, the causes and symptoms of

AD are thought to vary widely from patient to patient. Further, variations have been

previously reported in symptoms and disease progression rates among patients [30-33].

In recent years, transcriptome and genome-wide analysis data of nucleotide

polymorphisms have been collected from patients with AD, and various differentially

expressed genes and patient-specific SNPs have been identified in healthy individuals

and patients with AD [34-36]. Notably, these characteristics are not always seen equally

across all studies. One reason for this is the variations in technical factors among

studies. However, the diversity in molecular conditions and gene expression patterns in

the brain among patients with AD could also be a major contributor [37-41].

Thus, AD progression is expected to exhibit various disease states. Classifying and

characterizing these disease states and progression pathways is necessary to enable

appropriate state-dependent treatment. In addition to a comparison of the features of the

entire transcriptome, a comparison of the entire epigenome status among patients should

be useful for such classification and characterization of disease diversity. The database

of the Rush Alzheimer's Disease study by the ENCODE consortium

(https://www.encodeproject.org/) [42, 43] contains transcriptome and various

epigenome data such as histone modifications (H3K27ac, H3K27me3, and H3K4me3)

and CCCTC-binding factor (CTCF) binding site distribution from individuals in the

"No Cognitive Impairment (NCI)," "Mild Cognitive Impairment (MCI)," and

"Alzheimer's Disease (AD)" groups that are available for unlimited public use (termed

Rush AD database thereafter). Therefore, analyzing these publicly available data of

patients with different cognitive impairment stages is expected to elucidate the diversity

in the onset and progression of late-onset AD in more detail.

However, although the Rush AD database seemed to contain a satisfactory amount of

data for women, the amount of data for men seemed insufficient, particularly for

patients with MCI and AD, where the number of female patients in NCI, MCI, and AD

groups was 25, 29, and 28, respectively, whereas that of male patients was 14, 4, and 7, respectively. Therefore, in this study, the classification of disease status and AD progression pathways were performed using transcriptome and epigenome data of female patients in the Rush AD database as a first step of the analysis of the diversity of patients with AD using multiomics data. Our findings are expected to be useful not only for female patients but also for understanding symptoms in male patients, even though some features such as disease progression rate are different between male and female patients [44, 45].

In this study, patient diversity was considered by assessing the deviation from healthy individuals and diversity of disease progression. For this purpose, we first searched for differentially expressed gene sets between healthy individuals and patients. The gene sets searched here were those which contained the largest number of genes from gene sets that could actually distinguish healthy individuals from patients using cluster analysis. The candidate genes included therein were selected by identifying statistically significant differences in their expression levels. Second, we focused on the variety in the expression patterns of those gene sets between patients to show the existence of diverse conditions and subgroups within patients. The existence of these subgroups was also supported by comparisons of epigenomic data between individuals. Third, the relationship between subgroups and structure of the disease progression pathway was predicted based on the similarity in gene expression between subgroups.

RESULTS

Fundamental features of the analyzed transcriptome and epigenome data

RNA sequencing (RNA-seq) data (read count and TPM) of the dorsolateral prefrontal cortex of female individuals belonging to the NCI, MCI, and AD groups were acquired from the Rush AD database. In this dataset, the number of RNA-seq data of individuals in the NCI, MCI, and AD groups were 25, 29, and 28, respectively (Table 1, Table S1a–S1b).

To examine the quality and bias of analyzed data, we estimated one of the size factors of the RNA-seq data, that is, the median of the ratios (MOR) defined in the recent literature for DESeq [46], using read count data (Table S1a). The ratio [standard deviation of MOR over individuals]/[average MOR over individuals] was 0.1075. Similarly, we evaluated the first and third quartiles of the ratio (1OR and 3OR) between

126 the average and standard deviation, which were 0.1147 and 0.1006, respectively. These
127 results suggested that the size factor would not considerably input bias among samples
128 or affect the comparison of either the read count data or TPM values.
129
When examining the significance of the difference in gene expression levels between
individuals aged <85 years and those >90 years, we did not detect any genes exhibiting
a false discovery rate (FDR) < 0.1 (likelihood ratio test and two-tailed t-test) (Table
S1a-S1b). This suggested that the variations in gene expression levels among the NCI,
MCI, and AD groups was not due to the influence of age but rather due to differences in
clinical stages.
136
In addition to RNA-seq data, the bigwig format data of fold change over control (FC)
on genome-wide histone modifications (H3K4me3, H3K27ac, and H3K27me3) and
CTCF binding site distribution obtained through chromatin immunoprecipitation
sequencing (ChIP-seq) were acquired from the Rush AD database for 39 individuals:
11, 11, and 17 individuals from the NCI, MCI, and AD groups, respectively (Table
S1c). In this study, the regions with FC > 5 were defined as the regions of peak
intensities of these epigenome marks [47].
144
Cluster analysis of differentially expressed gene sets in the majority of patients
with MCI
To extract a differentially expressed set of genes among patients with NCI, MCI, or AD,
we performed the likelihood ratio test, which is one of the most familiar tests for
estimating differentially expressed genes (DEGs) from read count data, for each gene
using the DESeq2 software [48] (Table S1a). Consequently, we performed hierarchical
clustering of individuals using genes with P-values below the threshold value P^D_{NMA} =
0.01, 0.02, 0.03, 0.04, and 0.05 (Figs. 1 and S1a–S1e). When $P^D_{NMA} \leq 0.05$, a cluster
with significantly more patients with MCI (MCI cluster) and a cluster with patients with
NCI, AD, and a small number of patients with MCI was formed via the first divergence
of the dendrogram (P < 0.05, Fisher's exact test; Fig. S1a–S1e). However, even though
the gene set for clustering contained only genes with smaller P values, patients with
NCI, MCI, and AD were not from their respective cluster; no clusters comprising solely
of patients with NCI, MCI, or AD were formed through the first divergence of the
dendrogram.
160

We then performed the same analysis using analysis of variance (ANOVA), which is

the most familiar test for evaluating differences of characteristic values among three or

more groups, to determine the P-values using TPM values (Table S1b), where P^A_{NMA}

was defined instead of P^D_{NMA}. Herein, ANOVA was generally better for detecting

differences and was expected to identify more candidate genes in a gene set than the

likelihood ratio test. Consequently, an MCI cluster and a cluster with patients with NCI,

AD, and a small number of patients with MCI were formed, as in the previous case (Fig.

S1f-S1j). Additionally, when the P^A_{NMA} was 0.04, a cluster comprising solely patients

with MCI was formed through the first divergence of the dendrogram (Fig. 1a). In

contrast, we did not observe any clear separation between patients with NCI and those

with AD, regardless of the P^A_{NMA} value.

Notably, we did not obtain any gene set when applying the P^D_{NMA} and P^A_{NMA} values

that enabled the complete discrimination of patients with NCI, MCI, and AD. In

addition, depending on the P^D_{NMA} and P^A_{NMA} values, the patients with MCI who

deviated from the MCI cluster varied; however, 24 patients with MCI were always

included in the MCI cluster (Fig. S1). This suggests that the MCI cluster consisted of 24

patients with MCI, named MCI_{Major} individuals, whereas the 5 other patients with MCI

(M3, M12, M14, M16, and M28) who tended to deviate from the MCI_{Major} individuals,

were named MCI_{NCI_AD} individuals. We assumed that the MCI_{Major} and MCI_{NCI_AD}

individuals were classified to the substages named MCI_{Major} and MCI_{NCI_AD},

respectively.

Moreover, the set of genes with P-values of the above ANOVA $\leq P^A_{NMA} = 0.04$ was

named the "M-NA set" as it significantly differed between the majority of patients with

MCI from those with NCI and AD (Fig. 1b, Table S2a). In a 3-dimensional U-map

using the M-NA set, we clearly observed the deviation of the MCI_{Major} individuals from

those at other substages, with MCI_{NCI_AD} detected in a mixed group of patients with NCI

and AD (Fig. 1c).

The classification result of patients with MCI into MCI_{Major} and MCI_{NCI_AD} individuals

seemed to be independent of the age of individuals. For example, the numbers of

patients over and under 90 years old in MCI_{Major} were 19 and 5, whereas those in

MCI_{NCI_AD} were 3 and 2, respectively (P-value of the Fisher's exact test = 0.5688;

Table S1a–S1b).

Functional features of genes in the M-NA set

We also considered the functional features of genes in the M-NA set. In this set, we obtained 594 genes (290 upregulated and 304 downregulated genes in MCI_{Major}); of them, 349 genes (164 upregulated and 185 downregulated in MCI_{Major}) were annotated (Fig. 1b) as protein coding or noncoding RNA genes using Entrez [49]. Herein, both sets of upregulated and downregulated genes in MCI_{Major} contained some previously reported AD risk-associated genes; in particular, the upregulated gene set contained *CYP19A1, GSTM1, HSD11B1, IL10, MT-RNR1, PCK1, TRAK2, UBE2D1*, and *XBP1*, whereas the downregulated gene set contained *MT-ATP8* and *NME8* [50].

In addition, we performed enrichment analysis of all genes including genes with higher and lower expression levels in MCI_{Major} individuals (Table S2b–S2d). We identified that genes with higher expression levels in MCI_{Major} individuals were enriched in GO terms for the functions "response to lipopolysaccharide," "response to molecule of bacterial origin," "Toll-like receptor signaling pathway," "regulation of MHC class II biosynthetic process," "vitamin metabolic process," "water-soluble vitamin metabolic process," "chemotaxis," and "taxis" (P-value $< 10^{-3}$ and FDR < 0.5) (Table S2c). Whereas, we identified that genes with lower expression levels in MCI_{Major} individuals were enriched in GO terms for the functions "mRNA 5′-splice site recognition" and "mRNA cis splicing, via spliceosome" (P-value $< 10^{-3}$ and FDR < 0.5) (Table S2d). Recent studies have suggested that some of these functional features are characteristic of AD pathology; for example, the lipopolysaccharide found in the wall of all Gram-negative bacteria was reported to play a role in causing AD [51, 52], Toll-like receptors were found to participate in neuronal cell death by apoptosis [53, 54], MHC class II-expressing cells were suggested to be involved in the degradation of neurons [55], vitamins were reported to affect the generation and clearance of Aβ [56], and microglia were shown to exhibit chemotaxis for Aβ clearance [57, 58].

Differentially expressed gene sets between patients with NCI and AD

To extract a differentially expressed gene set between patients with NCI and those with AD, we performed hierarchical clustering of individuals using the genes satisfying the following criteria: mean expression level (TPM) was higher in patients with AD than in those with NCI, with smaller P-values in the likelihood ratio test than the P^D_{NA} threshold, or mean expression level (TPM) was lower in patients with AD than in those with NCI, with smaller P-values in the likelihood ratio test than the P^D_{AN} threshold. Subsequently, we searched the sets of P^D_{NA} and P^D_{AN} values that divided the patients

with NCI and those with AD into different clusters in the first branch of the

dendrogram. Within the range $P^D_{NA} \leq 0.05$ and $P^D_{AN} \leq 0.05$, we identified the finite but

narrow area of P^D_{NA} and P^D_{AN}, $P^D_{NA} = 0.01$ and $P^D_{AN} = 0.02$, $P^D_{NA} = 0.01$ and $P^D_{AN} =$

0.03, $P^D_{NA} = 0.02$ and $P^D_{AN} = 0.03$, $P^D_{NA} = 0.03$ and $P^D_{AN} = 0.03$, and $P^D_{NA} = 0.05$ and

$P^D_{AN} = 0.02$, where patients with NCI were completely separated from those with AD

(Fig. 2a). Thus, we obtained the largest set of genes separating the NCI and AD groups

at $P^D_{NA} = 0.05$ and $P^D_{AN} = 0.02$ (Fig. S2a, Table S3a); this gene set was named the N-

A_{D1} set. Notably, the number of genes with a lower mean expression level (TPM) in

patients with AD than in those with NCI was larger when $P^D_{NA} = 0.03$ and $P^D_{AN} = 0.03$

than when $P^D_{NA} = 0.05$ and $P^D_{AN} = 0.02$ (Fig. S2b, Table S3b). Notably, this gene set

contained all genes in other gene sets with P^D_{NA} and P^D_{AN} smaller than 0.03 that could

separate the NCI and AD groups. Thus, we focused on the gene set obtained at $P^D_{NA} =$

0.03 and $P^D_{AN} = 0.03$ and named it the "N-A_{D2} set."

We performed the same analysis using two-tailed t-tests to determine the P-values,

defining the P^t_{NA} and P^t_{AN} instead of the P^D_{NA} and P^D_{AN}, respectively. Herein, the t-test

was generally better for detecting differences and was expected to identify more

candidate genes in a gene set than the likelihood ratio test. Searching in the ranges P^t_{NA}

≤ 0.05 and $P^t_{AN} \leq 0.05$, patients with NCI were completely separated from those with

AD in 12 of 15 cases of P^t_{NA} and P^t_{AN} set for $P^t_{NA} \leq P^t_{AN}$ (Fig. 2a). When we searched

the wider range $P^t_{NA} \leq 0.1$ and $P^t_{AN} \leq 0.1$, patients with NCI were completely separated

from those with AD in 37 of 55 cases of P^t_{NA} and P^t_{AN} set for $P^t_{NA} \leq P^t_{AN}$ (Fig. 2a). We

thus obtained the largest set of genes separating the NCI and AD groups at $P^t_{NA} = 0.08$,

and $P^t_{AN} = 0.1$ (Fig. S2c, Table S3c). This gene set was named the "N-A_t set."

Hierarchical clustering using a combination of the expression levels of all genes in the

N-A_t and N-A_{D1} sets also separated patients with NCI from those with AD (Fig. 2b,

Table S3d). Conversely, the expression of all genes in the N-A_t and N-A_{D2} sets did not

separate patients with NCI from those with AD. However, clustering using a

combination of the expression levels of genes in the N-A_t set and a part of the N-A_{D2} set

with low P-values completely separated patients with NCI from those with AD (Fig. 2c,

Table S3e). Such gene sets are expected to involve various typical differentially

expressed genes between patients with NCI and those with AD. In this study, the gene

set consisting of the N-A_t and N-A_{D1} sets, which could completely separate patients with

NCI from those with AD, was named the "N-A1 set" (Table S3d), whereas the gene set

consisting of the N-A$_t$ set and maximum number of genes in the N-A$_{D2}$ set with low P-
values was named the "N-A2 set" (Table S3e).

Patients with AD can be classified into three groups

From the results of the hierarchical clustering for patients with NCI and AD performed

using the N-A1 set, we noticed the emergence of three subgroups of patients with AD

(Figs. 2b and 3a). Thus, we performed ANOVA on the expression levels of genes that

were higher in patients with AD than in those with NCI among the three assumed

subgroups (1020 genes). We identified 367 genes whose expression levels differed

significantly (FDR < 0.1) among the 3 subgroups. Performing k-means cluster analysis

of the expression levels of these genes in patients with AD, we identified two

characteristic clusters: a cluster consisting of 96 genes the expression levels of which

were low in only one subgroup (Cluster CA1) and another consisting of 271 genes the

expression levels of which were high in only one subgroup (Cluster CB1) (Fig. 3a).

Thereafter, we performed a similar analysis for patients with NCI. Based on the shape

of the dendrogram, patients with NCI were divided into three or four subgroups (Fig.

2b, Fig. 3a). However, when patients with NCI were divided into three subgroups,

ANOVA revealed only nine genes with FDR < 0.1 among the three subgroups (Table

S3d). Moreover, when patients with NCI were divided into four subgroups, ANOVA

revealed only two genes with FDR < 0.1 among the four subgroups (Table S3d). Based

on these values, the number of genes in clusters CA1 and CB1 was sufficiently large.

Collectively, these results suggested that the gene expression patterns in patients with

AD, which can be classified into three types, were highly nonuniform. Patients with AD

exhibiting high gene expression levels in both clusters, CA1 and CB1, were classified to

the AD$_{HH}$ subgroup (named AD$_{HH}$ individuals), those exhibiting low gene expression

levels in both CA1 and CB1 were classified in the AD$_{LL}$ subgroup (named AD$_{LL}$

individuals), whereas those exhibiting high gene expression levels in cluster CA1 and

low gene expression levels in cluster CB1 were classified in the AD$_{HL}$ subgroup (named

AD$_{HL}$ individuals).

Hierarchical clustering of patients with NCI and AD using the N-A2 set also revealed

three possible subgroups (Figs. 2c and 3b). By performing the same analysis described

above, we identified gene clusters similar to cluster CA1 containing 86 genes (cluster

CA2) and cluster CB1 containing 234 genes (cluster CB2). Accordingly, patients with

AD were divided into AD_{HH}, AD_{LL}, and AD_{HL} groups based on the expression profiles

of genes in clusters CA2 and CB2. The members of the AD_{HH}, AD_{LL}, and AD_{HL} groups

were the same in both the N-A1 and N-A2 sets (Fig. S3a).

Importantly, variations were observed among the AD_{HH}, AD_{HL}, and AD_{LL} individuals

when different P^t_{NA} and P^t_{AN} values were employed to determine the genes in the N-At

set. However, for small changes in the P^t_{NA} and P^t_{AN} values, less variation in individuals

was observed among subgroups; for example, when using $P^t_{NA} = 0.06$ and $P^t_{AN} = 0.1$

instead of $P^t_{NA} = 0.08$ and $P^t_{AN} = 0.1$ to determine the N-At. Herein, the gene set named

N-A3 was defined as that consisting of genes in the N-A$_t$ set and maximum number of

genes in N-A$_{D1}$ set with low P-values that could completely separate patients with NCI

from those with AD, whereas the gene set named N-A4 was defined as that consisting

of genes in the N-A$_t$ set and maximum number of genes in N-A$_{D4}$ set with low P-values

that could completely separate patients with NCI from those with AD. The classification

of patients with AD into the AD_{HH}, AD_{LL}, and AD_{HL} groups was similar to that obtained

for N-A1, N-A2, N-A3, and N-A4, where 24 of the 28 individuals belonged to the same

subgroups (Fig. S3). This suggested the robustness of our method in classifying the

AD_{HH}, AD_{HL}, and AD_{LL} individuals.

In addition, the classification result of patients with AD into AD_{HH}, AD_{HL}, and AD_{LL}

individuals seemed to be independent of the ages of individuals. For example, the

numbers of individuals aged over and under 90 years in AD_{HH} were eight and two, those

in AD_{LL} were six and four, and those in AD_{HL} were four and four, respectively (P-value

= 0.4850).

Common functional features of the N-A1 and N-A2 sets

We also considered the common functional features of the N-A1 and N-A2 sets. The N-

A1 set contained 1477 genes (1020 upregulated and 457 downregulated, AD), with 933

genes (683 upregulated and 250 downregulated) annotated as protein coding or

noncoding RNA genes using Entrez [49] (Fig. 3a). The N-A2 set contained 1424 genes

(947 upregulated and 477 downregulated, AD), with 889 genes (619 upregulated and

270 downregulated) annotated as protein coding or noncoding RNA genes using Entrez

[49] (Fig. 3b). The upregulated genes in both the N-A1 and N-A2 set contained *AR*,

CBS, *CLU*, *CYP19A1*, *DBH*, *GNA11*, *HLA-A*, *IQCK* , *MPO*, *MS4A4E*, *MX1*, *NLRP3*,

SDC2, *TAP2*, and *ZFHX3* (Table S3d–e), all of which have been reported to be AD

risk-associated genes in recent studies [50, 59-61]. Similarly, the downregulated genes

in both the N-A1 and N-A2 set contained *COL25A1* and *TARDBP* (Table S3d–e), which

have also been reported to be AD risk-associated genes [50].

Additionally, the clusters CA1, CB1, CA2, and CB2 contained 71, 215, 64, and 181

annotated genes, respectively (Fig. 3). We identified *CLU*, *SDC2*, and *ZFHX3* in both

the CA1 and CA2 clusters, whereas *DBH*, *GNA11*, *IQCK*, *MPO*, and *NLRP3* were

identified in both the CB1 and CB2 clusters (Table S3d–e), thereby suggesting that

these genes are responsible for the differences in the features among the three AD

subgroups [50, 61].

Next, we performed enrichment analysis of all genes, including those with higher and

lower expression levels in patients with AD for both the N-A1 and N-A2 set (Table

S3f–S3o). We identified that genes with higher expression levels in patients with AD in

both the N-A1 and N-A2 set were commonly enriched in GO terms for "forebrain

dorsal/ventral pattern formation," "mRNA 5′-splice site recognition," and "negative

regulation of NF-kappaB transcription factor activity" (P-value < 10^{-3} and FDR < 0.5)

(Tables S3g and S3j). Whereas, genes with lower expression levels in patients with AD

in both the N-A1 and N-A2 set were enriched in GO terms for "negative regulation of

intrinsic apoptotic signaling pathway in response to DNA damage by p53 class

mediator," "negative regulation of DNA damage response, signal transduction by p53

class mediator," and "coronary vasculature morphogenesis" (P-value < 10^{-3} and FDR <

0.5) (Table S3h, S3k). In the nervous system, NF-κB is known as one of the crucial

components in the molecular switch that converts short- to long-term memory [62].

Thus, the upregulation of genes that suppress NF-κB expression is expected to

exacerbate AD pathology. In addition, the downregulation of genes that suppress

apoptosis in response to DNA damage would also promote AD progression through the

enhancement of cell death in the central nervous system [63, 64].

We found that genes in both the CA1 and CA2 clusters were enriched in GO terms for

"response to xenobiotic stimulus", "cellular response to lipid", "negative regulation of

small molecule metabolic process", "lung development", "cellular response to external

stimulus", "negative regulation of carbohydrate metabolic process", "neuron migration",

and "positive regulation of epithelial cell differentiation" (P-value < 10^{-3} and FDR <

0.5) (Tables S3l and S3n). Whereas, genes in both the CB1 and CB2 clusters were

enriched in GO terms for "mitotic spindle assembly checkpoint signaling" and "L-alpha-

amino acid transmembrane transport" (P-value < 10^{-3} and FDR < 0.5) (Tables S3m and

S3o). Notably, recent studies have suggested that the perturbed cerebral carbohydrate

metabolism contributes to AD pathogenesis [65, 66], whereas neuron migration ability

contributes to neurogenesis in the brain [67, 68]. These results suggested that the three

AD subgroups may differ in these functional features.

We also evaluated the chromosomal distribution of genes belonging to the CA1 or CB1

cluster in N-A1 set, and that of those belonging to the CA2 or CB2 clusters in N-A2 set

(Fig. 4). Compared with the chromosomal distribution of all human coding and

noncoding genes, both gene sets belonging to the CA1 or CB1 clusters and CA2 or CB2

clusters seemed to be accumulated at Chr. 1, 7, 9, 16, 19, and 22 (Fig. 4).

Epigenome data analysis supported the classification of patients with AD into three

groups

The data from the dorsolateral prefrontal cortex of the Rush AD database analyzed in

the present study also included information on the genome-wide distribution of

epigenomic markers such as histone modifications (H3K4me3, H3K27me3, and

H3K27ac) and CTCF binding site distribution for approximately half of all individuals

(Table S1). Hence, to classify these individuals based on features other than

transcriptome, we next evaluated the similarity of such epigenomic features among

individuals using Jaccard coefficients of the genome-wide distribution of previously

defined peak regions of epigenome status. At the first divergence of the dendrogram

according to hierarchical clustering using Jaccard coefficients, individuals were divided

into two groups: a group with a typical epigenome status distribution and another one

with an untypical distribution for the respective epigenomic markers (Figs. 5–6 and S4).

The Jaccard coefficients of peak regions of epigenome status among individuals

suggested that more AD_{HL} individuals exhibited untypical patterns compared with those

in other patients with AD (Figs. 5–6 and S4, Table 2); in particular, the AD_{HL} group had

a significantly larger number of patients with H3K4me3 modifications compared with

the AD_{HH} and AD_{HL} groups (Table 2a). Similar features were obtained for H3K27ac,

H3K27me3, and CTCF (Table 2b–2d); however, these results were not significant.

Moreover, AD_{HL} individuals tended to exhibit low epigenetic peak region density (Figs.

5–6 and S4).

Further clustering revealed that individuals exhibiting typical H3K4me3 status were

further divided into two groups, named Typical 1 and Typical 2 (Fig. 5). Based on their

specific epigenome features, more AD_{HH} individuals belonged to the Typical 2

subgroup compared with other patients with AD (Table 3e). These facts suggest that the
AD$_{HH}$, AD$_{HL}$, and AD$_{LL}$ individuals exhibited differences in the genome wide
distribution of H3K4me3 modifications.
415
No effect of *APOE* gene polymorphisms in classifying individuals
To examine the relationships between the *APOE* genotype and the subgroup to which
individuals belong, the *APOE* genotype of each individual was identified using the
RNA-seq read data from the dorsolateral prefrontal cortex of individuals in the Rush
AD database. Accordingly, no individuals were found to simultaneously possess the ε2
and ε4 alleles. Therefore, individuals were classified in three categories: APOE2 carrier,
possessing one or two ε2 alleles; APOE3 carrier, possessing two ε3 alleles; and APOE4
carrier, possessing one or two ε4 alleles. Notably, the number of APOE2, APOE3, and
APOE4 carriers was not correlated with the clinical diagnostic stage of individuals
(Tables 3a and S1a–b). In addition, the number of individuals also seemed to be
independent of the two MCI (MCI$_{Major}$ and MCI$_{NCI\_AD}$) and three AD (AD$_{HH}$, AD$_{HL}$,
and AD$_{LL}$) substages (Table 3b–c, Table S1a–b).
428
Adjacency network among substages
The data on the expression levels of genes in the M-NA set revealed differences
between patients with typical MCI, NCI, and AD, whereas those in the N-A1 or N-A2
sets revealed differences between patients with NCI and AD. To clarify the
relationships among disease substages and the pathways of disease progression in these
substages, we explored the adjacency networks among substages using M-NA and N-
A1 sets, and M-NA and N-A2 sets by performing the partition-based graph abstraction
(PAGA) method [69]. Using PAGA, we generated a graph of substages in which the
edge weights represented connection confidence. Notably, connection confidence
between NCI and AD substages often involved finite values (Table S4); however, the
direct connection between NCI and AD substages seemed unnatural in disease
progression. Therefore, we assumed that only connections with confidence values
sufficiently larger than the maximum confidence values of the connection between NCI
and AD substages (MD$_{N\_A}$) were effective for the adjacency network among substages
(Fig. S5). Subsequently, we used connections with confidence values larger than twice
that of MD$_{N\_A}$ to construct the adjacency network (Table S4).
445
Changes in gene expression patterns that characterize disease states are considered to
occur gradually as the disease progresses. Therefore, we observed that disease appeared

to progress from NCI to AD along the previously obtained adjacency network. Note that

the adjacency network obtained using the M-NA and N-A1 sets (Fig S5a) and that

constructed using the M-NA and N-A2 sets (Fig. S5b) exhibited a slightly different

structure because of differences in the contained genes in these sets. Thus, we focused

on the network constructed using common connections obtained from both networks

constructed using the M-NA and N-A1 sets (Fig S5a) and M-NA and N-A2 sets (Fig.

S5b) (Fig. 7). We did not obtain any connections from NCI to MCI_{Major}, the major group

of MCI individuals, in any of the networks (Figs. 7 and S5a–b), which seemed unnatural

for AD progression. Thus, we performed more detailed subgroup classifications of

patients with NCI, MCI, and AD to reveal the relationship between NCI and MCI_{Major}

individuals.

Detailed classification of patients with MCI by comparing between patients with

NCI and those with MCI and between patients with MCI and those with AD

To classify patients with MCI more comprehensively, we conducted comparisons

between patients with NCI and those with MCI and between patients with MCI and

those with AD. First, we performed hierarchical clustering of individuals using the

genes satisfying the following criteria: the mean expression level (TPM) was higher in

the MCI group than that in the NCI group with smaller P-values in the likelihood ratio

test than the P^D_{MN} threshold or the mean expression level (TPM) was lower in the MCI

group than that in the NCI group, with smaller P-values in the likelihood ratio test than

the P^D_{NM} threshold. We performed cluster analysis using each gene set within the range

$P^D_{NM} \leq 0.05$ and $P^D_{MN} \leq 0.05$. However, we did not identify any cases in which patients

with NCI were completely separated from those with MCI (Figs. 8a and S6a). However,

clusters with significantly more patients with NCI (NCI clusters) and MCI clusters were

formed within all cases of P^D_{NM} and P^D_{MN} except in the case of $P^D_{NM} = 0.05$ and $P^D_{MN} =$

0.01 (Fig. 8a).

When we used two-tailed t-tests instead of the likelihood ratio test to determine the P-

values, our results were qualitatively the same. P^t_{MN} and P^t_{NM} were defined instead of

P^D_{MN} and P^D_{NM}, respectively. Analysis using each gene set within the ranges $P^t_{MN} \leq 0.05$

and $P^t_{NM} \leq 0.05$ revealed no cases in which patients with NCI were completely

separated from those with MCI (Figs. 8a and S6a). However, NCI and MCI clusters

were formed within all cases of P^t_{NM} and P^t_{MN} (Fig. 8a).

Next, we performed the same analyses for patients with MCI and those with AD. In this

case, P^D_{MA}, P^D_{AM}, P^t_{MA}, and P^t_{AM} were given as values corresponding to P^D_{NM}, P^D_{MN},

P^t_{NM}, and P^t_{MN}, respectively. Similar to the above results, we did not identify any cases

in which patients with MCI were completely separated from those with AD (Fig. 8b and

S6b). However, MCI and AD clusters were formed within all cases of P^D_{MA}, P^D_{AM},

P^t_{MA}, and P^t_{AM} except in the case of $P^D_{MA} = 0.05$ and $P^D_{AM} = 0.01$ and that of $P^t_{MA} =$

0.05 and $P^t_{AM} = 0.01$ (Fig. 8b).

In this study, variations were observed in patients with MCI who deviated from the MCI

cluster depending on P^D_{NM}, P^D_{MN}, P^t_{NM}, P^t_{MN}, P^D_{MA}, P^D_{AM}, P^t_{MA}, and P^t_{AM} values. For

each patient with MCI, we counted the number of times they belonged to the NCI

cluster upon changing the P^D_{NM}, P^D_{MN}, P^t_{NM}, and P^t_{MN} values, and those belonging to

the AD cluster upon changing the P^D_{MA}, P^D_{AM}, P^t_{MA}, and P^t_{AM} values (Fig. 8c).

Consequently, individuals M3, M4, M5, M8, M12, M13, M14, M16, M28, and M29

belonged to the NCI cluster more than once, whereas individuals M3, M5, M9, M12,

M14, M16, M23, M27, and M28 belonged to the AD cluster more than once.

Notably, individuals M3, M12, M14, M16, and M28, (same as those in MCI$_{NCI\_AD}$),

clearly exhibited a greater number of deviations from patients with NCI and AD than

MCI$_{Major}$ individuals (Fig. 8c). To classify MCI$_{Major}$ individuals in more detail, we

performed k-means clustering using the number of times each individual belonged to

the NCI cluster and that belonging to the AD cluster (Figs. 8c and S6c). As a result,

MCI$_{Major}$ individuals were divided into three subtypes: MCI$_{Typical}$ individuals (17

individuals) who rarely or never belonged to other clusters; MCI$_{NCI}$ individuals (4

individuals) who may belong to NCI clusters only; and MCI$_{AD}$ individuals (3

individuals) who may belong to AD clusters only (Table S1a–S1b). Thus, MCI was

divided into four substages: MCI$_{Typical}$, MCI$_{NCI}$, MCI$_{AD}$, and MCI$_{NCI\_AD}$.

Detailed classification of patients with NCI by comparing them with patients with

MCI

Similar analysis showed that individuals N11, N12, N18, N19, N21, and N22 belonged

to the MCI cluster more than once when the P^D_{NM}, P^D_{MN}, P^t_{NM}, and P^t_{MN} values were

changed (Fig. 9a). After k-means clustering of patients with NCI using the number of

times each individual belonged to the MCI cluster, patients with NCI were divided into

two clusters: one comprising N11, N21, and N22, and another comprising those who

never belonged to the MCI cluster (Figs. 9a and S7a). Therefore, these individuals were

classified as typical patients with NCI belonging to the NCI cluster, named $NCI_{Typical}$
individuals, who were assumed to be in the $NCI_{Typical}$ substage. Individuals N12, N18,
and N19 were classified as patients with NCI close to MCI, named NCI_{MCI} individuals,
who were assumed to be in the NCI_{MCI} substage (Table S1a–S1b).
523

Detailed classification of patients with AD by comparing them with patients with
MCI
Similarly, 17 patients with AD belonged to the MCI cluster more than once, when the
P^D_{MA}, P^D_{AM}, P^t_{MA}, and P^t_{AM} values were changed (Fig. 9b). Using k-means clustering of
patients with AD based on the number of times each individual belonged to the MCI
cluster, patients with AD were divided into three clusters, with individuals A10, A14,
A15, A17, A18, A19, A22, and other patients with AD who never belonged to the MCI
cluster being included in the same cluster (Figs. 9b and S7b). Therefore, these
individuals were expected to be in one of the typical AD substages: AD_{HH}, AD_{HL}, or
AD_{LL} (Table S1a–S1b). In contrast, individuals A2, A3, A4, A6, A7, A8, A9, A11,
A20, and A21 were assumed to be at one of the AD substages close to MCI and were
named AD_{HH_MCI}, AD_{HL_MCI}, or AD_{LL_MCI} (Table S1a–S1b).
536

Adjacency network among substages obtained after detailed classifications of
individuals suggests multiple disease progression pathways from NCI to AD
Detailed classifications of individuals suggested that patients with NCI were classified
into $NCI_{Typical}$ and NCI_{MCI}, patients with MCI were classified into $MCI_{Typical}$, MCI_{NCI},
MCI_{AD}, and MCI_{NCI_AD}, and patients with AD were classified into AD_{HH}, AD_{HH_MCI},
AD_{HL}, AD_{HL_MCI}, AD_{LL}, and AD_{LL_MCI}. We constructed adjacency networks among
these 12 substages using gene expression data from the M-NA and N-A1 sets (Fig. S8a,
Table S5a) as well as from the M-NA and N-A2 sets (Fig. S8b, Table S5b) as
previously described. Subsequently, we focused on the adjacency network that
consisted of common connections between the network from the M-NA and N-A1 sets,
and that from the M-NA and N-A2 sets (Fig. 10). We identified more than four
progression paths from NCI to AD that passed through different substages, as follows:
I) from NCI ($NCI_{Typical}$ or NCI_{MCI}) to AD_{LL} through MCI_{NCI_AD}, II) from NCI to MCI_{NCI}
and from MCI_{NCI} to AD_{LL} through AD_{HH_MCI} and AD_{LL_MCI}, respectively, III) from NCI
to MCI_{NCI} and from MCI_{NCI} to AD_{HH} through $MCI_{Typical}$ or AD_{HH_MCI}, and IV) from
NCI to AD_{HL} or AD_{LL} via MCI_{NCI}, MCI_{AD}, or AD_{HL_MCI}. This suggested that at least
four disease progression pathways are active in the three classified AD substages.
554

Comparisons with the findings of recent studies

Many recent studies have performed transcriptome analyses and identified various significant differentially expressed genes (significant DEGs) between healthy (control) individuals and patients with AD, which they termed "AD risk genes" [50, 59-61]. The cluster analysis of patients with NCI, MCI, and AD and that of patients with NCI and AD was performed using these AD risk genes (Fig. S9, Table S6). However, these results differed from the individual clustering results in Figures 1, 2, and 3. Notably, most of the AD risk gene data used here were obtained from research on the hippocampus, whereas the data analyzed in the previous sections of this study were obtained from the prefrontal cortex. A recent study showed that gene expression patterns differ between the hippocampus and prefrontal cortex [39]. Hence, the observed inconsistencies were attributed to these differences.

From the dataset analyzed in the previous sections, no significant DEGs were detected between patients with NCI and those with AD, whereas only nine significant DEGs were identified among patients with NCI, MCI, and AD (Tables S1a–S1b and S7). Herein, significant DEGs were evaluated via the likelihood test using DEseq2 with FDR < 0.01. Notably, the abovementioned study reported that not many significant DEGs were yielded in areas belonging to the prefrontal cortex [39]. These facts supported the consistency between the results of the present and those of recent studies.

DISCUSSION

In this study, the analysis of data from women individuals in the Rush AD database suggested that the pathological progression pathway of AD is nonuniform, exhibiting different typical forms of deterioration. First, cluster analysis using gene expression data from individuals diagnosed with NCI, MCI, and AD suggested that NCI can be classified in at least two substages, whereas MCI and AD can be classified in four and six substages, respectively. Differences among AD substages were identified based on epigenomic features, in particular the genome-wide distribution of H3K4me3. However, the *APOE* gene polymorphisms of individuals appeared to not correlate with the disease substage. In addition, some of the genes and gene sets that were suggested in recent studies to be AD-associated contributed to the characterization of each AD substage in the present study. Second, an inference of adjacency networks among the different substages suggested the existence of four typical NCI $\rightarrow$ AD disease progression

591 pathways from different MCI to different AD substages. These findings on disease
592 progression features will facilitate further advancements in AD research. Moreover,
593 they are expected to induce changes in the strategic bases of AD prevention, treatment,
594 and studies for biomarkers [70-72].
595
Recent studies using hippocampal and frontal lobe transcriptomes have also suggested
the existence of multiple AD subtypes. However, the relationship between multiple
states, such as their connection in a series on the same disease progression path or on
different paths, has not been clarified. Furthermore, these studies did not focus on the
diversity of NCI and MCI. In contrast, in this study, both patients with NCI and MCI
were divided into multiple subgroups at different substages based on their respective
gene expression characteristics. In addition, estimating the connectivity among
substages clearly revealed the existence of multiple pathological pathways of disease
progression.
605
This study analyzed data from the dorsolateral prefrontal cortex obtained from the Rush
AD database, rather than the hippocampus, which has been the focus of many previous
studies. The reason for analyzing this dataset in this study was that, it included not only
transcriptome data but also data on the genome-wide distribution of histone
modifications (H3K4me3, H3K27me3, H3K27ac) and CTCF binding sites (Table S1).
Such datasets can potentially reveal the relationship between a wide range of
transcriptional regulatory states and disease conditions. Recently, by analyzing such
datasets, changes in CTCF binding and transcription activities were shown to be
associated with changes in pathological conditions [73]. The analysis of such
epigenome data in the present study suggests that the genome-wide distribution of
H3K4me3 differed among individuals belonging in three different AD substages.
However, the mechanism by which such differences in the epigenetic status are related
to gene expression patterns remains unclear and warrants elucidation in future studies.
619
Notably, many recent studies have performed transcriptome analyses to identify
significant DEGs between healthy individuals and patients with AD. However, by
focusing on the expression patterns of more widespread candidate genes rather than the
significant DEGs, we identified different characteristics appearing in the dorsolateral
prefrontal cortex between healthy individuals and patients with AD and that disease
substages occur in response to changes in the disease state.
626

In addition, many studies have focused on gene expression patterns in the hippocampus. One such previous research suggested the existence of multiple states in the hippocampus of patients with AD based on differences in the gene expression profiles [39]. However, as previously mentioned, the gene expression patterns differ between the hippocampus and prefrontal cortex, and these differences exhibit further variations between healthy individuals and patients with AD. Hence, different results would be obtained even from the same analysis. Importantly, the hippocampal data analyzed in previous studies can be analyzed in the same way as in this study; connections between substages can be extracted and pathological routes can be estimated, all of which are crucial issues that need to be addressed in the future.

The present study has some limitations. We only classified female individuals as we used transcriptome and epigenomic data of female patients from the Rush AD database. However, the insights of this study are expected to be useful not only for female patients but also for understanding the rapid progression of symptoms in male patients. To validate this expectation, the dataset from male individuals should also be analyzed. In this dataset, however, the number of female patients in the NCI, MCI, and AD groups was 25, 29, and 28, respectively, whereas that of men was 14, 4, and 7, respectively, that is, three times smaller than that of female individuals (seven and four times smaller than those of female patients with MCI and AD). Moreover, the number of male patients with MCI and AD in the database (four and seven) was smaller than that of individuals in each of the three AD groups obtained for female individuals. With such restricted data, performing similar analyses on male individuals or sex-independent analyses was impossible. Obtaining more data on male patients is one of the essential issues that must be resolved in the future for performing such analyses and eliminating potential sex-related bias.

The likelihood ratio test using DESeq2 and *t*-test were used to evaluate the differences in RNA-seq data between different groups when searching for gene sets to characterize and classify individuals. The *t*-test is generally better for detecting differences and was expected to identify more candidate genes in a gene set than the likelihood ratio test or other nonparametric testing methods. However, the *t*-test produces more false positives than other testing methods when data deviate from a Gaussian distribution. The gene sets used in this study were expected to contain few false-positive genes because they were chosen under imposing the additional condition that healthy individuals could be clearly distinguished from patients through cluster analysis. However, inappropriate

663 genes may have still been retained. For more precise consideration, verification is
664 necessary using various nonparametric tests.
665
In addition to the transcriptome of each individual, the database targeted for the present
analysis also contained various epigenome data; however, other information on each
individual, such as postmortem interval, brain PH, and medical history was not
included. Therefore, although this study has yielded some unprecedented findings, the
validation and characterization of each substage through comparisons with other
physiological data could not be included. However, such analysis may be achieved by
combining meta-analysis with databases focusing on other feature groups.
673

EXPERIMENTAL PROCEDURES
Data processing
Gene expression data (read count and TPM) from the dorsolateral prefrontal cortex of
patients with NCI, MCI, and AD were acquired from the Rush AD database. In the
present analysis, the data of genes with average expression levels (TPM) in patients
with NCI, MCI, and AD were smaller than 0.1, whereas those of spike-in-RNAs were
eliminated.
681

The data provided by the Rush AD study in ENCODE does not publish information on
RNA quality such as the postmortem interval of samples and RIN values. However,
according to ENCODE standards, no specific alerts regarding the quality of the RNA-
seq data used in this study were provided. Thus, the quality of the data was assumed to
be guaranteed. The set of genes whose expression changes between patients with NCI
and AD and among patients with NCI, MCI, and AD, involved many genes identified as
AD risk genes in previous studies. This further supported the lack of data bias that
would contradict previous studies.
690

Statistical analyses
Two-tailed *t*-tests and ANOVA were performed using Python (version 3.9.16). All *t*-
tests in this study were Welch's *t*-tests. The false discovery rate (FDR) was estimated
through a set of P-values using the Benjamini–Hochberg procedure [74] implemented in
MultiPy (https://puolival.github.io/multipy/) [75]. The search for DEGs via the
likelihood ratio test using DESeq2 and the calculation of P-values for the Fisher's exact
test were performed using R (version 4.1.3).
698

Cluster analyses

Cluster analyses were performed using the values of $X = \log_2(TPM + 1) - C$, where C was chosen to satisfy the average of X over individuals = 0 for each gene. iDEP.96 (http://bioinformatics.sdstate.edu/idep96/) was used for hierarchical clustering [76]. In hierarchical clustering, the group-average method, was used to create a dendrogram, using the correlation coefficient as the distance. If the cluster formed by the first branch of hierarchical clustering satisfied the following criteria for a certain group (NCI, MCI, and AD), then that cluster was regarded as a cluster including significantly more individuals belonging to that group. These criteria were as follows: i) containing more than 2/3 of the individuals belonging to that group, and ii) Fisher exact test resulting in a P-value < 0.05. Further, k-means cluster analysis was performed using the Python package scikit-learn (version 1.0.2) [77]. The number of clusters in each k-means cluster analysis was determined using the elbow method.

The construction of UMAP was performed using sc.tl.pca() [77] with default parameters, scanpy.pp.neighbors() with "n_neighbors" = 30 [78], and the scanpy.tl.umap() functions with n_components = 3 [69] of scanpy (version 1.9.3) according to the literature on basic workflows for clustering analysis using Scanpy (https://scanpy-tutorials.readthedocs.io/en/latest/pbmc3k.html). All principal components were used for clustering using the scanpy.pp.neighbors() function. The obtained results were qualitatively independent of the parameters of these applications.

Epigenomic data analysis

The Jaccard coefficient of the genome-wide distribution of epigenomic markers among individuals was estimated as follows. The bigwig formatted files of fold change over control of these epigenome markers were acquired from individuals whose ChIP-seq data of H3K4me3, H3K27ac, H3K27me3, and CTCF were provided in the Rush AD database. From each bigwig format data, bedgraph format data was obtained using bigWigToBedGraph [79]. From the bedgraph format data, bed format data of peak regions were obtained following the removal of regions with FC < 5. The Jaccard coefficients between each pair of individuals were estimated using Bedtools (version 2.31.0). The density of each epigenomic marker on each chromosome was estimated according to the following equation: [Total length of genome regions with narrow peaks in ChIP-seq data in the chromosome obtained from the bed formatted file (bp)]/[Total length of chromosome obtained from GRCh38 (bp)].

Analysis of *APOE* gene polymorphisms

RNA-seq read data (fastq format data) from the dorsolateral prefrontal cortex of patients
with NCI, MCI, and AD were acquired from the Rush AD database. Fastp
(https://github.com/OpenGene/fastp) was used to remove low-quality reads and trim
adapter sequences. Following fastp processing, the reads were mapped to the CDS
sequence of the *APOE* exon 4 of GRCh38 using Hisat2
(https://daehwankimlab.github.io/hisat2/), and the results were obtained as a bam format
file. Variant calling was performed using the obtained bam format file and CDS
sequence data of the GRCh38 *APOE* exon 4 using bcftools
(https://github.com/samtools/bcftools). After filtering out low-quality calls (QUAL<20),
genotypes were determined based on the type of mutation and allele frequency.
745

746 **Enrichment analysis**
747 Enrichment analysis was performed for the M-NA, N-A1, and N-A2 sets using
748 Metascape (v3.5.20240101) (https://metascape.org/gp/index.html#/main/step1) [80].
749 The biological process terms in Gene Ontology were the main focus of this study.
750

Construction of an adjacency network among substages
The adjacency network among substages was estimated via PAGA [69] using the gene
expression data of each individual. Each group at each substage was regarded as a
partition in PAGA. PAGA can generate a graph of substages in which the edge weights
represent connection confidence. PAGA was performed using sc.tl.pca() [77] with
default parameters, scanpy.pp.neighbors() with "n_neighbors" = 2 [78], and the
scanpy.tl.paga() functions [69] of scanpy (version 1.9.3) according to the literature for
trajectory inference using transcriptome data (https://scanpy-
tutorials.readthedocs.io/en/latest/paga-paul15.html). All principal components were
used for clustering using the scanpy.pp.neighbors() function. Additionally, to extract the
essential network among substages, the value of the parameter "n_neighbors" of the
function scanpy.pp.neighbors() was set as two, which is the minimum value in the range
of this parameter. Thus, a network with nonredundant connections among substages was
expected to be obtained.
765

766 **AUTHOR CONTRIBUTIONS**
767 Kousei Honda and Akinori Awazu conceived and designed the study. Kousei Honda
768 performed the analyses. Kousei Honda and Akinori Awazu drafted the manuscript.

770 **ACKNOWLEDGEMENTS**

This research was supported by JST Moonshot R&D (Grant Number JPMJMS2021) and a JSPS KAKENHI grant (award number 21K06124, 24K09248). Computations were partially performed on the NIG supercomputer at the ROIS National Institute of Genetics.

REFERENCES

1. Gatz, M., Reynolds, C. A., Fratiglioni, L., Johansson, B., Mortimer, J. A., Berg, S.,... & Pedersen, N. L. (2006). Role of genes and environments for explaining Alzheimer disease. *Archives of General Psychiatry*, *63*, 168–174. https://doi.org/10.1001/archpsyc.63.2.168

2. World Health Organization. (2017). Global action plan on the public health response to dementia 2017–2025.

3. Suresh, S., Singh, A., Rushendran, R., Vellapandian, C., & Prajapati, B. (2023). Alzheimer's disease: the role of extrinsic factors in its development, an investigation of the environmental enigma. *Frontiers in Neurology*, *14*, 1303111. https://doi.org/10.3389/fneur.2023.1303111

4. Hardy, J. A., & Higgins, G. A. (1992). Alzheimer's disease: the amyloid cascade hypothesis. *Science*, *256*, 184–185. https://doi.org/10.1126/science.1566067

5. Hardy, J., & Selkoe, D. J. (2002). The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science*, *297*, 353–356. https://doi.org/10.1126/science.1072994

6. Jin, M., Shepardson, N., Yang, T., Chen, G., Walsh, D., & Selkoe, D. J. (2011). Soluble amyloid β-protein dimers isolated from Alzheimer cortex directly induce Tau hyperphosphorylation and neuritic degeneration. *Proceedings of the National Academy of Sciences*, *108*, 5819–5824. https://doi.org/10.1073/pnas.1017033108

7. Selkoe, D. J., & Hardy, J. (2016). The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Molecular Medicine*, *8*(6), 595–608. https://doi.org/10.15252/emmm.201606210

806   8. Sekiya, M., Wang, M., Fujisaki, N., Sakakibara, Y., Quan, X., Ehrlich, M. E.,...
807   Iijima, K. M. (2018). Integrated biology approach reveals molecular and
808   pathological interactions among Alzheimer's Aβ42, Tau, TREM2, and TYROBP in
809   Drosophila models. *Genome Medicine*, *10*(1), 1–20. https://doi.org/10.1186/s13073-
810   018-0530-9
811
812   9. Arriagada, P. V., Growdon, J. H., Hedley-Whyte, E. T., & Hyman, B. T. (1992).
813   Neurofibrillary tangles but not senile plaques parallel duration and severity of
814   Alzheimer's disease. *Neurology*, *42*, 631–631. https://doi.org/10.1212/wnl.42.3.631
815
816   10. Strittmatter, W. J., Saunders, A. M., Schmechel, D., Pericak-Vance, M., Enghild, J.,
817   Salvesen, G. S., & Roses, A. D. (1993). Apolipoprotein E: high-avidity binding to
818   beta-amyloid and increased frequency of type 4 allele in late-onset familial
819   Alzheimer disease. *Proceedings of the National Academy of Sciences*, *90*, 1977–
820   1981. https://doi.org/10.1073/pnas.90.5.1977
821
822   11. Itzhaki, R. F., Lin, W. R., Shang, D., Wilcock, G. K., Faragher, B., & Jamieson, G.
823   A. (1997). Herpes simplex virus type 1 in brain and risk of Alzheimer's disease. *The
824   Lancet*, *349*, 241–244. https://doi.org/10.1016/s0140-6736(96)10149-5
825
826   12. Osaka, R., Kobayashi, N., Shimada, K., Ishii, A., Oka, N., & Kondo, K. (2022).
827   VP26, a herpes simplex virus type 1 capsid protein, increases DNA methylation in
828   COASY promoter region. *Brain, Behavior, & Immunity - Health*, *26*, 100545.
829   https://doi.org/10.1016/j.bbih.2022.100545
830
831   13. Readhead, B., Haure-Mirande, J. V., Funk, C. C., Richards, M. A., Shannon, P.,
832   Haroutunian, V.,... Dudley, J. T. (2018). Multiscale analysis of independent
833   Alzheimer's cohorts finds disruption of molecular, genetic, and clinical networks by
834   human herpesvirus. *Neuron*, *99*, 64–82.
835   https://doi.org/10.1016/j.neuron.2018.05.023
836
837   14. Wang, L., Davis, P. B., Volkow, N. D., Berger, N. A., Kaelber, D. C., & Xu, R.
838   (2022). Association of COVID-19 with new-onset Alzheimer's disease. *Journal of
839   Alzheimer's Disease*, *89*(2), 1–4. https://doi.org/10.3233/jad-220717
840

841   15. Kandimalla, R., Thirumala, V., & Reddy, P. H. (2017). Is Alzheimer's disease a
842   type 3 diabetes? A critical appraisal. *Biochimica et Biophysica Acta (BBA)-*
843   *Molecular Basis of Disease*, *1863*, 1078–1089.
844   https://doi.org/10.1016/j.bbadis.2016.08.018
845

846   16. Ott, A., Stolk, R. P., van Harskamp, F., Pols, H. A. P., Hofman, A., & Breteler, M.
847   M. B. (1999). Diabetes mellitus and the risk of dementia: The Rotterdam
848   Study. *Neurology*, *53*, 1937–1937. https://doi.org/10.1212/wnl.53.9.1937
849

850   17. McGeer, P. L., & McGeer, E. G. (1995). The inflammatory response system of
851   brain: implications for therapy of Alzheimer and other neurodegenerative diseases.
852   *Brain Research Reviews*, *21*, 195–218. https://doi.org/10.1016/0165-
853   0173(95)00011-9
854

855   18. Hirai, K., Aliev, G., Nunomura, A., Fujioka, H., Russell, R. L., Atwood, C. S.,...
856   Smith, M. A. (2001). Mitochondrial abnormalities in Alzheimer's disease. *Journal*
857   *of Neuroscience*, *21*, 3017–3023. https://doi.org/10.1523/jneurosci.21-09-
858   03017.2001
859

860   19. Nunomura, A., Perry, G., Aliev, G., Hirai, K., Takeda, A., Balraj, E. K.,... Smith, M.
861   A. (2001). Oxidative damage is the earliest event in Alzheimer disease. *Journal of*
862   *Neuropathology & Experimental Neurology*, *60*, 759–767.
863   https://doi.org/10.1093/jnen/60.8.759
864

865   20. Wang, X., Wang, W., Li, L., Perry, G., Lee, H. G., & Zhu, X. (2014). Oxidative
866   stress and mitochondrial dysfunction in Alzheimer's disease. *Biochimica et*
867   *Biophysica Acta (BBA)-Molecular Basis of Disease*, *1842*, 1240–1247.
868   https://doi.org/10.1016%2Fj.bbadis.2013.10.015
869

870   21. Chakravarthy, B., Gaudet, C., Ménard, M., Brown, L., Atkinson, T., LaFerla, F.
871   M.,... & Whitfield, J. (2012). Reduction of the immunostainable length of the
872   hippocampal dentate granule cells' primary cilia in 3xAD-transgenic mice
873   producing human Aβ1-42 and tau. *Biochemical and Biophysical Research*
874   *Communications*, *427*(1), 218–222. https://doi.org/10.1016/j.bbrc.2012.09.056
875

876   22. Morelli, A., Sarchielli, E., Guarnieri, G., Coppi, E., Pantano, D., Comeglio, P.,...
877   Vannelli, G. B. (2017). Young human cholinergic neurons respond to physiological
878   regulators and improve cognitive symptoms in an animal model of Alzheimer's
879   disease. *Frontiers in Cellular Neuroscience, 11*, 339.
880   https://doi.org/10.3389/fncel.2017.00339
881
882   23. Plácido, A. I., Pereira, C. M. F., Duarte, A. I., Candeias, E., Correia, S. C., Santos,
883   R. X.,... Moreira, P. I. (2014). The role of endoplasmic reticulum in amyloid
884   precursor protein processing and trafficking: implications for Alzheimer's disease.
885   *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease, 1842*, 1444–
886   1453. https://doi.org/10.1016/j.bbadis.2014.05.003
887
888   24. Vorobyeva, A. G., & Saunders, A. J. (2018). Amyloid-β interrupts canonical Sonic
889   hedgehog signaling by distorting primary cilia structure. *Cilia, 7*(1), 1–11.
890   https://doi.org/10.1186/s13630-018-0059-y
891
892   25. Bartzokis, G. (2004). Age-related myelin breakdown: a developmental model of
893   cognitive decline and Alzheimer's disease. *Neurobiology of Aging, 25*, 5–18.
894   https://doi.org/10.1016/j.neurobiolaging.2003.03.001
895
896   26. Bartzokis, G. (2011). Alzheimer's disease as homeostatic responses to age-related
897   myelin breakdown. *Neurobiology of Aging, 32*, 1341–1371.
898   https://doi.org/10.1016/j.neurobiolaging.2009.08.007
899
900   27. Cheng, Y., Saville, L., Gollen, B., Veronesi, A. A., Mohajerani, M., Joseph, J. T., &
901   Zovoilis, A. (2021). Increased Alu RNA processing in Alzheimer brains is linked to
902   gene expression changes. *EMBO Reports, 22*, e52255.
903   https://doi.org/10.15252/embr.202052255
904
905   28. Macciardi, F., Giulia Bacalini, M., Miramontes, R., Boattini, A., Taccioli, C.,
906   Modenini, G.,... Mapstone, M. (2022). A retrotransposon storm marks clinical
907   phenoconversion to late-onset Alzheimer's disease. *GeroScience, 44*, 1525–1550.
908   https://doi.org/10.1007/s11357-022-00580-w
909

910   29. Ravel-Godreuil, C., Znaidi, R., Bonnifet, T., Joshi, R. L., & Fuchs, J. (2021).
911   Transposable elements as new players in neurodegenerative diseases. *FEBS Letters*,
912   *595*, 2733–2755. https://doi.org/10.1002/1873-3468.14205
913

914   30. Duara, R., & Barker, W. (2021). Heterogeneity in Alzheimer's disease diagnosis
915   and progression rates: implications for therapeutic trials. *Neurotherapeutics*, *19*, 8–
916   25. https://doi.org/10.1007/s13311-022-01185-z
917

918   31. Komarova, N. L., & Thalhauser, C. J. (2011). High degree of heterogeneity in
919   Alzheimer's disease progression patterns. *PLoS Computational Biology*, *7*,
920   e1002251. https://doi.org/10.1371/journal.pcbi.1002251
921

922   32. Younas, N., Zafar, S., Shafiq, M., Noor, A., Siegert, A., Arora, A. S., … Zerr, I.
923   (2020). SFPQ and Tau: Critical factors contributing to rapid progression of
924   Alzheimer's disease. *Acta Neuropathologica*, *140*, 317–339.
925   https://doi.org/10.1007/s00401-020-02178-y
926

927   33. Younas, A., Younas, N., Iqbal, M. J., Ferrer, I., & Zerr, I. (2024). Comparative
928   interactome mapping of Tau-protein in classical and rapidly progressive
929   Alzheimer's disease identifies subtype-specific pathways. *Neuropathology and*
930   *Applied Neurobiology*, *50*, e12964. https://doi.org/10.1111/nan.12964
931

932   34. Jansen, I. E., Savage, J. E., Watanabe, K., Bryois, J., Williams, D. M., Steinberg,
933   S.,... Posthuma, D. (2019). Genome-wide meta-analysis identifies new loci and
934   functional pathways influencing Alzheimer's disease risk. *Nature Genetics*, *51*,
935   404–413. https://doi.org/10.1038/s41588-018-0311-9
936

937   35. Wightman, D. P., Jansen, I. E., Savage, J. E., Shadrin, A. A., Bahrami, S., Holland,
938   D.,... Posthuma, D. (2021). A genome-wide association study with 1,126,563
939   individuals identifies new risk loci for Alzheimer's disease. *Nature Genetics*, *53*,
940   1276–1282. https://doi.org/10.1038%2Fs41588-021-00921-z
941

942   36. Yuen, S. C., Zhu, H., & Leung, S. W. (2020). A systematic bioinformatics workflow
943   with meta-analytics identified potential pathogenic factors of Alzheimer's disease.
944   *Frontiers in Neuroscience*, *14*, 209. https://doi.org/10.3389/fnins.2020.00209
945

946 37. Bellenguez, C., Küçükali, F., Jansen, I. E., Kleineidam, L., Amin, N., Naj, A. C., ...
947   Lambert, J. (2022). New insights into the genetic etiology of Alzheimer's disease
948   and related dementias. *Nature Genetics*, *54*, 412–436.
949   https://doi.org/10.1038/s41588-022-01024-z
950

951 38. Johnson, E. C., Carter, E. K., Dammer, E. B., Duong, D. M., Gerasimov, E. S., Liu,
952   Y.,... Seyfried, N. T. (2022). Large-scale deep multi-layer analysis of Alzheimer's
953   disease brain reveals strong proteomic disease-related changes not observed at the
954   RNA level. *Nature Neuroscience*, *25*, 213–225. https://doi.org/10.1038/s41593-021-
955   00999-y
956

957 39. Neff, R. A., Wang, M., Vatansever, S., Guo, L., Ming, C., Wang, Q.,... Zhang, B.
958   (2021). Molecular subtyping of Alzheimer's disease using RNA sequencing data
959   reveals novel mechanisms and targets. *Science Advances*, *7*, eabb5398.
960   https://doi.org/10.1126/sciadv.abb5398
961

962 40. Tijms, B. M., Gobom, J., Reus, L., Jansen, I., Hong, S., Dobricic, V.,... Visser, P. J.
963   (2020). Pathophysiological subtypes of Alzheimer's disease based on cerebrospinal
964   fluid proteomics. *Brain*, *143*, 3776–3792. https://doi.org/10.1093/brain/awaa325
965

966 41. Tijms, B. M., Gobom, J., Teunissen, C., Dobricic, V., Tsolaki, M., Verhey, F.,...
967   Visser, P. J. (2021). CSF proteomic Alzheimer's disease-predictive subtypes in
968   cognitively intact amyloid negative individuals. *Proteomes*, *9*, 36.
969   https://doi.org/10.3390/proteomes9030036
970

971 42. ENCODE Project Consortium. (2012). An integrated encyclopedia of DNA
972   elements in the human genome. *Nature*, *489*(7414), 57.
973   https://doi.org/10.1038/nature11247
974

975 43. Luo, Y., Hitz, B. C., Gabdank, I., Hilton, J. A., Kagda, M. S., Lam, B.,... Cherry, J.
976   M. (2020). New developments on the Encyclopedia of DNA Elements (ENCODE)
977   data portal. *Nucleic Acids Research*, 48, D882–D889.
978   https://doi.org/10.1093/nar/gkz1062
979

44. Davis, E. J., Broestl, L., Abdulai-Saiku, S., Worden, K., Bonham, L. W., Miñones-
Moyano, E., ... & Dubal, D. B. (2020). A second X chromosome contributes to

resilience in a mouse model of Alzheimer's disease. *Science translational*

medicine, *12*(558), eaaz5677. https://doi.org/aaz5677

45. Lopez-Lee, C., Torres, E. R. S., Carling, G., & Gan, L. (2024). Mechanisms of sex

differences in Alzheimer's disease. *Neuron*.

https://doi.org/10.1016/j.neuron.2024.01.024

46. Anders, S., & Huber, W. (2010). Differential expression analysis for sequence count

data. *Genome Biology*, *11*, R106. https://doi.org/10.1186/gb-2010-11-10-r106

47. Shin, H., Liu, T., Duan, X., Zhang, Y., & Liu, X. S. (2013). Computational

methodology for ChIP-seq analysis. *Quantitative Biology*, *1*(1), 54-

70. https://doi.org/10.1007/s40484-013-0006-2

48. Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change

and dispersion for RNA-seq data with DESeq2. *Genome biology*, *15*, 1-21.

https://doi.org/10.1186/s13059-014-0550-8

49. Schuler, G. D., Epstein, J. A., Ohkawa, H., & Kans, J. A. (1996). Entrez: Molecular

biology database and retrieval system. In Methods in enzymology (Vol. 266, pp.

141-162). Academic Press. https://doi.org/10.1016/S0076-6879(96)66012-1

1003

1004 50. Hu, Y. S., Xin, J., Hu, Y., Zhang, L., & Wang, J. (2017). Analyzing the genes

1005   related to Alzheimer's disease via a network and pathway-based

1006   approach. *Alzheimer's research & therapy*, *9*, 1-15. https://doi.org/10.1186/s13195-

1007   017-0252-z

1008

1009 51. Singh, S., Sahu, K., Singh, C., & Singh, A. (2022). Lipopolysaccharide induced

1010   altered signaling pathways in various neurological disorders. *Naunyn-*

1011   *Schmiedeberg's Archives of Pharmacology*, *395*(3), 285-294.

1012   https://doi.org/10.1007/s00210-021-02198-9

1013

1014 52. Zhan, X., Stamova, B., & Sharp, F. R. (2018). Lipopolysaccharide associates with

1015   amyloid plaques, neurons and oligodendrocytes in Alzheimer's disease brain: a

1016   review. *Frontiers in aging neuroscience*, *10*, 42.

1017   https://doi.org/10.3389/fnagi.2018.00042

1018

1019   53. Adhikarla, S. V., Jha, N. K., Goswami, V. K., Sharma, A., Bhardwaj, A., Dey, A., ...

1020   & Jha, S. K. (2021). TLR-mediated signal transduction and neurodegenerative

1021   disorders. *Brain sciences*, *11*(11), 1373. https://doi.org/10.3390/brainsci11111373

1022

1023   54. Wu, L., Xian, X., Xu, G., Tan, Z., Dong, F., Zhang, M., & Zhang, F. (2022). Toll-

1024   like receptor 4: a promising therapeutic target for alzheimer's disease. *Mediators of*

1025   *inflammation*, *2022*. https://doi.org/10.1155/2022/7924199

1026

1027   55. Parachikova, A., Agadjanyan, M. G., Cribbs, D. H., Blurton-Jones, M., Perreau, V.,

1028   Rogers, J., ... & Cotman, C. W. (2007). Inflammatory changes parallel the early

1029   stages of Alzheimer disease. *Neurobiology of aging*, *28*(12), 1821-1833.

1030   https://doi.org/10.1016/j.neurobiolaging.2006.08.014

1031

1032   56. Grimm, M. O., Mett, J., & Hartmann, T. (2016). The impact of vitamin E and other

1033   fat-soluble vitamins on Alzheimer s disease. *International journal of molecular*

1034   *sciences*, *17*(11), 1785. https://doi.org/10.3390/ijms17111785

1035

1036   57. Lau, S. F., Wu, W., Wong, H. Y., Ouyang, L., Qiao, Y., Xu, J., ... & Ip, N. Y.

1037   (2023). The VCAM1–ApoE pathway directs microglial chemotaxis and alleviates

1038   Alzheimer's disease pathology. *Nature Aging*, *3*(10), 1219-1236.

1039   https://doi.org/10.1038/s43587-023-00491-1

1040

1041   58. Rogers, J., & Lue, L. F. (2001). Microglial chemotaxis, activation, and phagocytosis

1042   of amyloid β-peptide as linked phenomena in Alzheimer's disease. *Neurochemistry*

1043   *international*, *39*(5-6), 333-340. https://doi.org/10.1016/S0197-0186(01)00040-7

1044

1045   59. Xiang, S., Huang, Z., Wang, T., Han, Z., Yu, C. Y., Ni, D., ... & Zhang, J. (2018).

1046   Condition-specific gene co-expression network mining identifies key pathways and

1047   regulators in the brain tissue of Alzheimer's disease patients. *BMC medical*

1048   *genomics*, *11*, 39-51. https://doi.org/10.1186/s12920-018-0431-1

1049

1050   60. Rahman, M. R., Islam, T., Turanli, B., Zaman, T., Faruquee, H. M., Rahman, M.

1051   M., ... & Moni, M. A. (2019). Network-based approach to identify molecular

1052   signatures and therapeutic agents in Alzheimer's disease. *Computational biology*

1053   *and chemistry*, *78*, 431-439. https://doi.org/10.1016/j.compbiolchem.2018.12.011

1054

61. Yang, A. C., Vest, R. T., Kern, F., Lee, D. P., Agam, M., Maat, C. A., ... & Wyss-
    Coray, T. (2022). A human brain vascular atlas reveals diverse mediators of
    Alzheimer's risk. *Nature*, *603*(7903), 885-892. https://doi.org/10.1038/s41586-021-
    04369-3

62. Kaltschmidt, B., & Kaltschmidt, C. (2015). NF-KappaB in long-term memory and
    structural plasticity in the adult mammalian brain. *Frontiers in molecular
    neuroscience*, *8*, 69. https://doi.org/10.3389/fnmol.2015.00069

63. Lin, X., Kapoor, A., Gu, Y., Chow, M. J., Peng, J., Zhao, K., & Tang, D. (2020).
    Contributions of DNA damage to Alzheimer's disease. *International Journal of
    Molecular Sciences*, *21*(5), 1666. https://doi.org/10.3390/ijms21051666

64. Ranganathan, R., Sapozhnikov, G., Ni, W., Li, S., & Song, Y. (2023). Recent
    developments in the role of DNA damage response and understanding its
    implications for new therapeutic approaches in Alzheimer's disease. *Translational
    Medicine of Aging*. https://doi.org/10.1016/j.tma.2023.06.004

65. Chen, Z., & Zhong, C. (2013). Decoding Alzheimer's disease from perturbed
    cerebral glucose metabolism: implications for diagnostic and therapeutic
    strategies. *Progress in neurobiology*, *108*, 21-43.
    https://doi.org/10.1016/j.pneurobio.2013.06.004

66. Cunnane, S., Nugent, S., Roy, M., Courchesne-Loyer, A., Croteau, E., Tremblay,
    S., ... & Rapoport, S. I. (2011). Brain fuel metabolism, aging, and Alzheimer's
    disease. *Nutrition*, *27*(1), 3-20. https://doi.org/10.1016/j.nut.2010.07.021

67. Esteve, D., Molina-Navarro, M. M., Giraldo, E., Martínez-Varea, N., Blanco-
    Gandia, M. C., Rodríguez-Arias, M., ... & Lloret, A. (2022). Adult neural stem cell
    migration is impaired in a mouse model of Alzheimer's disease. *Molecular
    Neurobiology*, 1-15. https://doi.org/10.1007/s12035-021-02620-6

68. Young-Pearse, T. L., Bai, J., Chang, R., Zheng, J. B., LoTurco, J. J., & Selkoe, D. J.
    (2007). A critical function for β-amyloid precursor protein in neuronal migration

revealed by in utero RNA interference. *Journal of Neuroscience*, *27*(52), 14459-
14469. https://doi.org/10.1523/JNEUROSCI.4701-07.2007

69. Wolf, F. A., Hamey, F. K., Plass, M., Solana, J., Dahlin, J. S., Göttgens, B.,... Theis,
F. J. (2019). PAGA: graph abstraction reconciles clustering with trajectory inference
through a topology preserving map of single cells. *Genome Biology*, *20*, 1–9.
https://doi.org/10.1186%2Fs13059-019-1663-x

70. Karikari, T. K., Pascoal, T. A., Ashton, N. J., Janelidze, S., Benedet, A. L.,
Rodriguez, J. L., Chamoun, M., Savard, M., Kang, M. S., Therriault, J., Schöll, M.,
Massarweh, G., Soucy, J., Höglund, K., Brinkmalm, G., Mattsson, N., Palmqvist, S.,
Gauthier, S., Stomrud, E., . . . Blennow, K. (2020). Blood phosphorylated tau 181 as
a biomarker for Alzheimer's disease: A diagnostic performance and prediction
modelling study using data from four prospective cohorts. *The Lancet*
Neurology, *19*(5), 422-433. https://doi.org/10.1016/S1474-4422(20)30071-5

71. Therriault, J., Pascoal, T. A., Lussier, F. Z., Tissot, C., Chamoun, M., Bezgin, G.,
Servaes, S., Benedet, A. L., Ashton, N. J., Karikari, T. K., Kunach, P., Wang, Y.,
Massarweh, G., Vitali, P., Soucy, J., Blennow, K., Zetterberg, H., & Gauthier,
S. (2022). Biomarker modeling of Alzheimer's disease using PET-based Braak
staging. *Nature Aging*, *2*(6), 526-535. https://doi.org/10.1038/s43587-022-00204-0

72. Yada, Y., & Naoki, H. (2023). Few-shot prediction of amyloid β accumulation from
mainly unpaired data on biomarker candidates. *Npj Systems Biology and*
Applications, *9*(1), 1-11. https://doi.org/10.1038/s41540-023-00321-5

73. Patel, P. J., Ren, Y., & Yan, Z. (2023). Epigenomic analysis of Alzheimer's disease
brains reveals diminished CTCF binding on genes involved in synaptic organization.
Neurobiology of Disease, 184, 106192. https://doi.org/10.1016/j.nbd.2023.106192

74. Benjamini, Y., Hochberg, Y. (1995). Controlling the false discovery rate: a practical
and powerful approach to multiple testing. *Journal of the Royal Statistics Society*
Series B, *57*, 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

75. Puoliväli, T., Palva, S., & Palva, J.M. (2020). Influence of multiple hypothesis
testing on reproducibility in neuroimaging research: A simulation study and Python-

based software. *Journal of Neuroscience Methods*, *337*, 108654.

https://doi.org/10.1016/j.jneumeth.2020.108654

1127

1128   76. Ge, S.X., Son, E.W., & Yao, R. (2018). iDEP: an integrated web application for

1129   differential expression and pathway analysis of RNA-Seq data. *BMC*

1130   *Bioinformatics*, *19*, 534. https://doi.org/10.1186/s12859-018-2486-6

1131

1132   77. Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., …

1133   Duchesnay, É. (2011). Scikit-learn: machine learning in Python. *JMLR*, *12*, 2825–

1134   2830.

1135

1136   78. McInnes, L., Healy, J., & Melville, J. UMAP: Uniform manifold approximation and

1137   projection for dimension reduction. arXiv:1802.03426.

1138   https://doi.org/10.48550/arXiv.1802.03426

1139

1140   79. Kent, W. J., Zweig, A. S., Barber, G., Hinrichs, A. S., & Karolchik, D. (2010).

1141   BigWig and BigBed: enabling browsing of large distributed

1142   datasets. *Bioinformatics*, *26*(17), 2204-2207.

1143   https://doi.org/10.1093/bioinformatics/btq351

1144

1145   80. Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O.,

1146   … Chanda, S. K. (2019). Metascape provides a biologist-oriented resource for the

1147   analysis of systems-level datasets. *Nature Communications*, *10*(1),1523.

1148   https://doi.org/10.1038/s41467-019-09234-6

1149

1150 **Table 1**. Summary of the individuals included in this study (see Table S1a and S1b for
1151 more details).

Clinical stage	No Cognitive Impairment (NCI)	Mild Cognitive Impairment (MCI)	Alzheimer's Disease (AD)
# of individuals	25	29	28
Individual index	N1 – N25	M1 – M29	A1 – A28

1152

Table 2. Number of individuals exhibiting the typical genome-wide distributions of
each epigenetic state, E_T, and of those exhibiting untypical distributions, E_U, among the
three AD substages (AD_{HH}, AD_{LL}, and AD_{HL}); H3K4me3 (a), H3K27ac (b), H3K27me3
(c), and CTCF (d). The E_T and E_U in each substage were obtained from Figures 5, 6, and
S4. P-values obtained using the Fisher's exact test were 0.023843 (a), 0.193439 (b),
0.193439 (c), and 0.285294 (d). Number of individuals exhibiting E_T were divided into
E_{T1} (Typical 1) and E_{T2} (Typical 2) among the three AD substages by further clustering
for H3K4me3 obtained from Figure 5 (e). The P-value obtained using the Fisher's exact
test was 0.0420316 (e).

1162

1163 (a)

H3K4me3	AD_{HH}	AD_{LL}	AD_{HL}
# of individuals with typical (E_T)	6	4	3
# of individuals with untypical (E_U)	0	0	4

1164

1165 (b)

H3K27ac	AD_{HH}	AD_{LL}	AD_{HL}
# of individuals with typical (E_T)	5	4	3
# of individuals with untypical (E_U)	1	0	4

1166

1167 (c)

H3K27me3	AD_{HH}	AD_{LL}	AD_{HL}
# of individuals with typical (E_T)	5	4	3
# of individuals with untypical (E_U)	1	0	4

1168

1169 (d)

CTCF	AD_{HH}	AD_{LL}	AD_{HL}
# of individuals with typical (E_T)	6	3	4
# of individuals with untypical (E_U)	0	1	3

1170

1171 (e)

H3K4me3	AD_{HH}	AD_{LL}	AD_{HL}
# of individuals with Typical 1 (E_{T1})	1	2	2
# of individuals with Typical 2 (E_{T2})	5	2	1

# of individuals with untypical (E_U)	0	0	4

1172

1173

1174 **Table 3.** Number of APOE2, APOE3, and APOE4 carriers in each clinical diagnostic
1175 stage (a), in the two MCI substages (b), and in the three AD substages (c). The genotype
1176 of each individual is given in Table S1a and S1b. The P-values obtained using the
1177 Fisher's exact test were 0.8841 (a), 0.08148 (b), and 0.4379 (c).
1178
1179 (a)

Clinical Diagnostic Stage	NCI	MCI	AD
# of APOE2 carriers	5	3	4
# of APOE3 carriers	16	20	18
# of APOE4 carriers	4	6	6

1180
1181 (b)

MCI substages	MCI_{Major}	MCI_{NCI_AD}
# of APOE2 carriers	1	2
# of APOE3 carriers	18	2
# of APOE4 carriers	5	1

1182
1183 (c)

AD substages	AD_{HH}	AD_{LL}	AD_{HL}
# of APOE2 carriers	3	1	0
# of APOE3 carriers	6	7	5
# of APOE4 carriers	1	2	3

1184
1185

1186

(a)

M-NA set

■ NCI ■ MCI ■ AD

MCI$_{Major}$ individual

1187

(b)

M-NA set

■ NCI
■ MCI
■ AD

1.5

■ UP (
■ DOW

N$_G$: # of
N$_{kG}$: # of

(c)

● NCI   ● MCI$_{Major}$ ● MCI$_{NCI\_AD}$ ● AD

● NCI
● MCI Major
● MCI NCI_AD
● AD

1188

Fig. 1. Re   **ied**

using the   the

M-NA set   r

each indiv

expression levels of each gene was described using X = log2(TPM + 1) − C, where C

was chosen to satisfy the average of X over individuals = 0 for each gene. MCI$_{Major}$

individuals occupy the right region in the dendrogram. N$_G$ and N$_{kG}$ indicate the total

number of genes and number of genes annotated as protein coding or noncoding RNA

genes with higher (UP) and lower (DOWN) expression levels in MCI_{Major} individuals,

respectively. Scatter plot of individuals in 3-dimensional space constructed using U-

MAP with 1, 2, and 3 axes shown from various directions (c).

1200

(a)

1201

(b)

N-A1 set (P^D_{NA} = 0.05, P^D_{AN} = 0.02, Pt_{NA} = 0.08, Pt_{AN} = 0.10)

1202

(c)

N-A2 set (P^D_{NA} = 0.03, P^D_{AN} = 0.03, Pt_{NA} = 0.08, Pt_{AN} = 0.10)

1203

Fig. 2 Phase diagrams of the hierarchical clustering of patients with NCI and AD,

and the results of the hierarchical clustering of individuals in the N-A1 and N-A2

sets. Phase diagram of the results of hierarchical clustering when using genes where p ≤

P^D_{NA} and p ≤ P^D_{AN} (a, left), and that when using genes where p ≤ Pt_{NA} and p ≤ Pt_{AN} (a,

right). Circles indicate the complete separation of patients with NCI from those with

AD into different clusters at the first branch of the dendrogram. The dendrograms

obtained using the N-A1 (b) and N-A2 sets (c) are shown. Broken lines indicate the

boundary between the AD_{HL}, AD_{LL}, and AD_{HH} groups.

1212

1213

Fig. 3 Results of the hierarchical clustering of individuals in the N-A1 and N-A2 sets and expression level of each gene. The dendrograms and expression level of each gene in the N-A1 set for each individual (a) (see Table S3d for more details) and those in the N-A2 set for each individual (b) (See Table S3e for more details) were plotted. The heatmap of the expression levels of each gene was described using $X = \log_2(TPM + 1) - C$, where C was chosen to satisfy the average of X over individuals = 0 for each gene. Broken lines indicate the boundary between the AD_{HL}, AD_{LL}, and AD_{HH} groups.

Fig. 4 Chromosomal distribution of genes belonging to the CA1 or CB1 cluster (Fig.
3a), and that of genes belonging to the CA2 or CB2 cluster (Fig. 3b). The chromosomal
distribution of coding and non-coding genes was estimated using the human genome
data GRCh38.

H3K4me3

Fig. 5 Jaccard coefficient of the genome-wide distribution of H3K4me3 among
individuals. Individuals were divided into two groups at the first divergence of the
dendrogram. The epigenetic state exhibited by individuals belonging to the group
containing the larger number of individuals was regarded as the "typical" epigenetic
state, whereas that exhibited by individuals belonging to the group containing the fewer
individuals was regarded as the "untypical" epigenetic state. Further, individuals
belonging to the "typical" group were divided into two groups, named Typical 1 and
Typical 2, at the second divergence of the dendrogram. Dashed lines indicate the
boundaries between groups.

H3K27ac

1239 **Fig. 6** Jaccard coefficient of the genome-wide distribution of H3K27ac among
1240 individuals. Analysis was conducted similar to that described in Fig. 5.
1241

1242

1243

1244 **Fig. 7 Adjacency network among substages.** The network constructed using common
1245 connections obtained from both the network constructed using the M-NA and N-A1 sets
1246 (Fig S5a) and that constructed using the M-NA and N-A2 sets (Fig. S5b). The width of
1247 each edge indicates the connection confidence between each pair of connected nodes.
1248

1249

(a)

1250

1251

(b)

1252

(c)

1253

Fig. 8 Detailed classification of patients with MCI. Number of patients with NCI who belong to the MCI cluster (left value) and that of patients with MCI who belong to the NCI cluster (right value) as a function of P^D_{NM} and P^D_{MN} (a, left), and as a function of P^t_{NM} and P^t_{MN} (a, right). Number of patients with MCI who belong to the AD cluster (left value) and that of patients with AD who belong to the MCI cluster (right value) as

a function of P^D_{MA} and P^D_{AM} (b, left), and as a function of P^t_{MA} and P^t_{AM} (b, right).

Number of deviations of each patient with MCI from the MCI to the NCI (c, upper left)

and AD (c, lower left) cluster. Broken lines indicate boundaries between MCI_{NCI_AD} and

MCI_{Major} individuals. Scatter plot of MCI_{Major} individuals as a function of the number of

times that each individual belonged to the NCI and AD cluster (c, right). Using k-means

clustering (Fig. S6c), MCI_{Major} individuals were divided into $MCI_{Typical}$, MCI_{NCI}, and

MCI_{AD} groups.

1266

Fig. 9 Detailed classification of patients with NCI. Number of deviations of each

patient with NCI from the NCI to the MCI cluster (a), and that of each patient with AD

from the AD to the MCI cluster (b). Broken lines indicate boundaries among clusters of

individuals obtained via k-means clustering (Fig. S7a and S7b).

1274

1275

1276 **Fig. 10 Adjacency networks among substages.** A network containing common
1277 connections from both networks obtained using the data in the M-NA and N-A1 sets
1278 (Fig. S8a) and that in the M-NA and N-A2 sets (Fig. S8b). The width of each edge
1279 indicates the connection confidence between each pair of connected nodes. Arrows
1280 indicated the inferred direction of disease progression.

1281