Suboptimal refeeding compensates stunting in a mouse model of juvenile malnutrition

Jean-Louis Thoumas*, Amandine Cavaroc*, Damien Sery*, François Leulier*, Filipe De Vadder*†

* Institut de Génomique Fonctionnelle de Lyon, École Normale Supérieure de Lyon, CNRS UMR 5242, UCBL Lyon-1, F-69007 Lyon, France

† To whom correspondence should be addressed: filipe.de_vadder@ens-lyon.fr

Short title: Suboptimal refeeding after juvenile malnutrition

List of abbreviations:

GTT: Glucose tolerance test

IGF-1: Insulin-like growth factor 1

ITT: Insulin tolerance test

Lp^{WJL}: *Lactiplantibacillus plantarum* WJL

LPD: Low-protein diet

PC: Principal component

PEM: Protein energy malnutrition

MWD: Modified Western Diet

WA: weight-for-age relationship

WD: Total New Western Diet

<u>Abstract:</u>

Background: The highest rate of growth in mammals occurs in early life, particularly

after weaning. Growth is conditioned by the nutritional status. Indeed, restriction of

dietary protein in early life leads to wasting and/or stunting.

Objective: Here, we developed a mouse model of juvenile protein malnutrition

triggering stunting and studied the physiological effects of refeeding using various diets

and interventions.

Methods: In a first intervention, we refed the mice with an optimal diet (breeding diet,

rich in protein and fiber). We then treated the mice during the refeeding phase with

Lactiplantibacillus plantarum WJL (Lp^{WJL}), a previously described bacterial strain that

has the ability to stimulate growth via the somatotropic axis in early life upon chronic

malnutrition. Finally, we established a model of suboptimal refeeding, upon which the

mice were given a western diet (33% kcal from fat; 17% kcal from protein) or an

isocaloric modified western diet low in proteins (33% kcal from fat; 8% kcal from

protein) after juvenile protein malnutrition.

Results: We found that, in females, optimal growth was restored by control diet

refeeding. In males, control diet refeeding after a five-week protein restriction was not

enough to catch up growth retardation. No supplementary beneficial effect was found

associated to the microbial intervention in this context. Surprisingly, our results showed

that, in males, suboptimal refeeding with a diet rich in fat but low in protein was

sufficient to buffer the deleterious effects of protein restriction on growth. However, this

macroscopic benefit was associated to metabolic alteration. While Lp^{WJL} treatment had

no effect on growth per se, we found that bacterial treatment further impaired glycemic

control upon suboptimal refeeding.

Conclusions: Overall, we describe a novel model of juvenile protein energy

malnutrition, where growth can be caught up by suboptimal refeeding.

Keywords: Mice; protein energy malnutrition; double burden of malnutrition; stunting;

probiotic intervention

Introduction

Early postnatal malnutrition is a burden in many developing countries, and is considered

the most worrying factor of death for young infants under five years old. Consequences

of malnutrition include wasting and stunting, which are defined respectively weight-for-

height and height-for-age below two standard deviations from the median of the World

Health Organization Child Growth Standards. Wasting is indicative of acute growth

disturbance from malnutrition, whereas stunting is more symptomatic of chronic

malnutrition, with impaired long-term growth (1). In 2020, about 150 million children

under five years of age were stunted (2). Stunting is directly caused by an inadequate

intake of nutrients, especially dietary protein (3). Besides stunting, severe malnutrition

in early life is associated to metabolic disturbances that persist later in life (4,5).

Extensive epidemiological studies have demonstrated that malnutrition during early life

has detrimental effects on health, which can last throughout the life of the malnourished

person (6). This is called nutritional or metabolic programming, a process through

which metabolic and physiological adaptations in response to a poor diet persist even

after the nutritional stress is removed (7–10). The first 1,000 days of life are crucial for

growth because of the high developmental plasticity during that time, after which

stunting generally becomes irreversible and leads to growth impairment in adulthood

(11). Recently, three articles of large human cohorts have highlighted the importance of

early life interventions to prevent wasting and stunting in more than 30 cohorts of

malnourished children (12–14).

In this context, establishing animal models of early life malnutrition seems crucial to

understand the mechanisms behind the observed adverse consequences. Previously, we

established a model of mouse juvenile malnutrition, upon which mice were fed a low-

protein low-fat diet from weaning (post-natal day 21, P21) to eight weeks of age

(15,16), which is the rough physiological equivalent of 1 year to 20 years of age in

humans (17). We showed that supplementation with the probiotic *Lactiplantibacillus*

plantarum WJL (Lp^{WJL}) sustains postnatal growth of malnourished animals by

stimulating anabolic metabolism and hormonal changes, namely the somatotropic axis

via the secretion of insulin-like growth factor 1 (IGF-1) (15,16). However, knowledge

on the long-term consequences of early malnutrition after refeeding is lacking. Indeed,

the first intervention to stop malnutrition is resolving the nutritional stress associated to

the lack of protein by providing an optimized diet.

Using a mouse model of juvenile protein energy malnutrition, we sought to investigate

the consequences on growth catch-up after refeeding with either an optimal or

suboptimal diets (richer in fat and poorer in protein). While growth was not affected by

a suboptimal diet, we also explored the metabolic alterations associated to this switch.

METHODS

Animals

Male and female specific pathogen-free C57Bl/6J mice, aged 9 to 10 weeks, were

purchased from Charles River (L'Arbresle, France). After one week of acclimatization

at the conventional animal room, they were mated in our animal facility and offspring

pups were used for experiments. Mice were kept in Innorack IVC Mouse 3.5 disposable

cages (Inovive, USA), exposed to 12:12 hours light-dark cycles, supplied with tap water

and given free access to food. All experiments were performed in accordance with the

European Community Council Directive of September 22, 2010 (2010/63/EU)

regarding the protection of animals used for experimental and other scientific purposes.

The research project was approved by a local animal care and user committee

(C2EA015) and subsequently authorized by the French Ministry of Research

(APAFIS#22788-2019091713204925 v8 and APAFIS#32380-2021061809124474 v4).

Experimental design

Specific-pathogen free C57Bl/6J mice were mated. After the delivery, the litter size was

reduced to six offspring per dam. From birth to post-natal day 14 (P14), the dams were

given free access to breeding diet (control diet; Altromin 1310). At P14, the pups were

kept with their mother, but food was changed according to the experimental groups. All

diets were irradiated at 40 kGy before use. Composition of the experimental diets is

given in Table 1. At P21, mice were weaned on the diet introduced at P14. Body weight

and length were controlled once a week. For the measurement of body length, mice

were briefly anesthetized with isoflurane and length was measured using a graduated

ruler. Body length was defined as the nose-to-anus distance. At the end of the

experiments, mice were killed by cervical dislocation. Tissues were sampled, weighed

and measured, and snap-frozen in liquid nitrogen.

Experiment 1 (male and female mice)

A control group was given control diet (Altromin 1310) from P14 to P91. Experimental

groups were given low-protein diet (LPD, composition in Table 1), which is a modified

isocaloric version of AIN-93G diet. LPD was custom-made by Sciences de l'Animal et

de l'Aliment de Jouy unit (SAAJ, INRAE UE1298, Jouy-en-Josas, France). Switch to

control diet was operated in 5 different groups from P28 to P56, then mice were kept on

control diet until P91. Blood samples were taken from the submandibular vein three

times: at diet switch, two weeks after the switch, and at P91. Time-matched samples

were taken in the control group.

Experiment 2 (males only)

Three groups were formed: a control group fed control diet until P119; a low-protein

group fed LPD from P14 to P119; and a refeeding group, fed LPD from P14 to P56,

then control diet from P56 to P119. From P56 to P119, two treatment arms were formed

in each group: one receiving a placebo solution (maltodextrin) and one receiving

10^8 CFU *Lactiplantibacillus plantarum* WJL (Lp^{WJL}) as described previously (16).

Solutions were given per os 5 times a week, in the morning. At P112, mice were taken

to ANIPHY Platform (Lyon) and put in the Minispec LF90 II (Bruker, Wissembourg,

France) for non-invasive determination of body composition using NMR. Mice were

taken back to our animal facility and given the experimental diet until P119.

Experiment 3 (males only)

Groups were formed as described in Figure 5. A control group was fed control diet from

P14 to P119. Experimental diets were control diet (Altromin 1310), LPD, Total New

Western Diet (WD; Teklad TD.140203) and a modified isocaloric version of WD

(MWD), with reduced protein content. MWD was custom-made for the experiment.

WD and MWD were purchased from Envigo (Gannat, France). WD is based on Total

Western Diet, as described in (18), with the modifications described by Monsanto et al.

(19). Composition of the diets is given in Table 1. As in Experiment 2, mice were given

either a placebo or an Lp^{WJL} solution, from P35 to P119.

Glucose and insulin tolerance tests

In Experiment 3, at P91 and P105, mice underwent respectively a glucose (GTT) or

insulin tolerance test (ITT). Mice were food-deprived for 6 hours before the start of the

tests. For the GTT, mice were given an intraperitoneal (i.p.) injection of a 20% sterile

glucose solution (1 g.kg^{-1} body weight). For the ITT, mice were given an i.p. injection

of human insulin (Novorapid, Novo Nordisk) at 0.75 U.kg^{-1} body weight. Glycemia was

monitored at the tip of the tail vein at times 0, 15, 30, 45, 60, 90 and 120 minutes post-

injection using a glucometer (One Touch Verio Reflect, Life Scan).

IGF-1 measurement in serum

Blood samples were collected from the submandibular vein at the time of diet switch,

two weeks after the switch and at the end of the experiment, into a sample tube

containing a clotting activator (Micro sample tube serum gel, 1.1 mL, Sarstedt). After

centrifugation at 10,000 g (Eppendorf 5424 R) for 5 minutes, sera were collected into

sterile 1.5 mL microtubes and kept at -70°C for further use. IGF-1 in serum was

measured using Mouse/Rat IGF-1 Quantikine ELISA (R&D Systems) according to

manufacturer's instructions.

Data treatment and statistics

Statistical analysis was performed using GraphPad Prism software version 10.0

(GraphPad, San Diego, California, USA). For Experiment 1, we used one-way ANOVA

followed by Tukey's multiplicity-adjusted test. Growth catch-up slopes were compared

using ANCOVA. Principal Component (PC) Analysis was performed based on the

morphological features measured, with PC selection based on percent of total explained

variance. For Experiment 2, we used two-way ANOVA followed by multiplicity-

adjusted Šídák's test. For Experiment 3, we used one-way ANOVA followed by Tukey's

multiplicity-adjusted test when comparing to control and two-way ANOVA followed by

multiplicity-adjusted Šídák's test when comparing treated groups. Unless stated

otherwise, results are presented as mean ± SEM. *P*-values are reported after adjustment

for multiple comparison. A value of $P < 0.05$ was considered statistically significant.

<u>RESULTS</u>

Optimal refeeding after long-term juvenile malnutrition is insufficient to fully recover

growth in males, but not in females

In order to model protein energy malnutrition (PEM), we designed an AIN-93G-based

isocaloric low-protein diet (LPD), with 5.1% total kcal coming from protein (Table 1).

We first investigated how relevant our model of malnutrition was to human disease.

Indeed, although several models have been published (20,21), there is no standard

murine model of PEM. We first characterized the impact on weight gain and linear

growth in C57Bl/6J mice fed an LPD, starting at P14 and going on after weaning

(which took place at P21). We constituted 5 different groups, which corresponded to 5

different times on LPD, going from 2 weeks to 6 weeks. From P28 to P56, mice were

refed with control chow diet. To make a parallel with juvenile human malnutrition, we

defined stunting and underweight in our model as body size and weight inferior to two

standard deviations from the mean of a control group of mice fed chow diet, i.e.,

optimal diet for mouse growth (dashed lines, Figure 1A-D). As expected, LPD-fed mice

had severe growth retardation following weaning, associated to reduced body weight

(Figure 1A-D). Since definition of wasting (low weight-for-length) in humans is

complex and requires a Box-Cox power transformation (22), and growth does not occur

in the same manner in mice and humans, we plotted length vs. weight in our

experimental animals. We found that weight and length are linearly correlated during

growth (Figure 1E-H). In LPD-fed mice, the slope was significantly smaller than in age-

matched control-fed mice (males and females: ANCOVA: $P < 10^{-4}$), meaning that the

mice were wasted (Figure 1E-F). In females, when mice were refed a control chow diet

after LPD, growth and weight catch-up occurred within three weeks of diet switch

(Figure 1B, D, H). However, in males, the longer the mice stood on LPD, the slower

growth and weight catch-up occurred (Figure 1A, C). Indeed, when male mice were fed

on LPD from P14 to P56, they were still stunted at P91 (Figure 1C), despite the removal

of the nutritional stress. It is noteworthy that, despite the prevalence of stunting in male

mice at P91, mice were not wasted after refeeding (Figure 1G-H).

Using the weight-for-age (WA) relationship described by Anstead et al. (20), based on

the classification by Gómez et al. (23), we calculated WA in all our groups (Figure 2A-

B). After weaning, WA was inferior to 50% in LPD-fed male mice (Figure 2A), showing

that the mice suffered from severe malnutrition. In females, however, LPD induced

severe malnutrition, but the severity was corrected with time, reaching moderate

malnutrition at P56 (Figure 2B). Refeeding with chow diet in male mice only partially

corrected the severity of malnutrition, since RF42 mice had a WA score of $85.1 \pm 2.1\%$

(5% confidence interval), going as low as $81.6 \pm 3.0\%$ for RF56 mice, meaning they

still faced mild malnutrition. As was the case previously, refeeding malnourished female

mice completely corrected early-life malnutrition (Figure 2B). Overall, our data show a

good example of nutritional imprinting of PEM, especially in males. Growth in mice

occurs with a linear phase that lasts about two weeks after weaning followed by a

plateau phase (24). We observed that this pattern was similar after refeeding. We thus

compared growth catch-up by comparing the slopes following refeeding (or weaning, in

the case of the control group) (Figure 2C-D). In males, growth catch-up differed

significantly between the groups ($P = 0.0205$; ANCOVA). After weaning on control

diet, mice gained on average 0.11 cm.day^{-1}. This value was similar when mice were

refed after only one week on LPD (group RF28), but differed significantly with all other

groups, which had a catch-up rate of about 0.08 cm.day^{-1} (Figure 2C). Thus, in males,

growth catch-up after refeeding is impaired by long-term exposure to LPD in early life.

In females, however, growth catch-up did not differ significantly between groups

($P = 0.0504$; ANCOVA; Figure 2D), showing a sex-dependent phenotype related to

resilience to LPD feeding in early life.

Furthermore, we measured weight of liver, soleus muscle and subcutaneous and

perigonadal fat, and length of femur, tibia, small intestine and colon (Figure 3). In

males, we observed significantly reduced size of femur and tibia (Figure 3E-F),

alongside reductions in intestinal and colon sizes, which were all the more marked as

the time on LPD was long (Figure 3G-H). When we plotted the totality of the measured

parameters using Principal Component Analysis, we saw that values were mostly spread

across PC1, which accounted for 30.44% of the total variance (Figure 3I). While control

animals clustered separately from refed mice, the latter formed a cluster, showing a

biological difference between control and refed mice. In contrast, in females, we only

saw slight variations of fat deposits, but all points clustered together in a Principal

Component Analysis (Supplementary Figure 1).

Insulin-like Growth Factor 1 (IGF-1) is a hormone mainly produced by the liver as a

downstream effector of growth hormone secretion (25). Circulating levels of IGF-1

directly regulate bone growth during juvenile age (26), and IGF-1 secretion is reduced

upon chronic undernutrition (27). We thus measured circulating IGF-1 in males at three

timepoints: at diet switch, two weeks after (at the end of the linear growth phase), and at

P91 when the mice were killed (Figure 3J). We took time-matched samples in control

mice. In control mice, IGF-1 levels did not differ significantly from P28 to P91 (about

350 ng.mL^{-1}). In contrast, all mice fed LPD had an IGF-1 serum concentration around

100 ng.mL^{-1} at the time of refeeding, capturing an important signature of stunting in our

model of PEM. Two weeks after refeeding, IGF-1 levels were comparable to those of

control male mice, and did not significantly vary at P91. Thus, while refeeding with an

optimal diet stimulated IGF-1 secretion, it was not enough to compensate for the growth

retardation, suggesting irreversible effects on male growth induced by long-term

exposure to LPD after weaning.

Lp^{WJL} treatment during the refeeding fails to improve systemic growth

Given the sex-related difference we observed previously, for the reminder of the study

we focused on male mice. We chose the most severe stunting phenotype that we

observed after refeeding, i.e., mice that were refed at P56. We separated mice in three

experimental groups: mice fed control diet from P14 to P119, mice fed LPD during the

same period of time, and mice fed LPD from P14 to P56, then control diet from P56 to

P119. We decided to keep until day 119, to see if the stunting phenotype observed at day

91 was still present four weeks later.

We previously reported that the Lp^{WJL} strain stimulates growth of juvenile mice upon

chronic malnutrition (15,16). In these studies, treatment with Lp^{WJL} was performed in a

model where, besides protein, fat was also drastically reduced in the diet. Treatment was

performed after weaning, i.e., during the linear phase of growth. We wondered about the

effect of Lp^{WJL} upon PEM and hypothesized that treating our mice during the refeeding

phase, which corresponds to a linear phase of growth, would stimulate their growth to a

point where mice would totally catch up their growth after a period of protein scarcity.

As expected, mice fed LPD all the time showed severe stunting and underweight

(Figure 4A-B). Despite a significant increase in body weight and length after refeeding,

placebo-treated refed mice (RF56, Figure 4A-B) still presented signs of stunting, since

their body length was less than 2 standard deviations from the mean of the control

group. Contrary to what we had hypothesized, supplementation with Lp^{WJL} during the

refeeding phase did not have an additive effect on growth or weight, with only a slight

increase in weight in LPD-fed mice at P91. Indeed, besides fat mass (Figure 4E-F),

Lp^{WJL} treatment had no impact on size and weight of several organs (Figure 4C-K). This

was evident when running Principal Component Analysis (Figure 4L), where mice

clustered mainly along PC1 (59.01% of total variance) according to diet, but not to

treatment. Thus, treatment with Lp^{WJL} during the refeeding phase failed to further

stimulate growth after prolonged exposition to protein scarcity, highlighting the long-

lasting deleterious effects of protein malnutrition during early life.

Suboptimal refeeding with a non-obesogenic Western diet compensates early-life

associated stunting, but impairs glucose metabolism

One of the main drivers of malnutrition is suboptimal nutrition, defined as inadequate

nutrient intake that does not meet the age-specific recommendations (2). Finally, we

sought to characterize the impact on weight gain and growth of LPD feeding followed

by refeeding with a suboptimal diet. For this experience, we decided to refeed the

animals at P35, i.e., the first time when growth catch-up was impaired upon refeeding

with an optimal diet (Figure 1E). Indeed, we sought to establish a novel model of

"double burden of malnutrition", defined as the simultaneous manifestation of both

undernutrition and overweight (28). During the first burden (P14 to P35), mice were fed

control diet or LPD. During the second burden (P35 to P119), mice were fed two types

of suboptimal diet: Total New Western Diet (WD) or a modified version of WD

(MWD), with protein accounting for 8.4% of total kcal (instead of 17%). In parallel to

these groups, we also fed mice control diet, WD, or MWD from P14 to P112 (Figure

1A). Unlike diet-induced obesity diet traditionally used in nutritional studies, WD has a

variety of fat sources and is more similar to the human Western diet. It is noteworthy

that this version of WD is not obesogenic in C57Bl/6J mice (19), so the idea was to

provide a suboptimal diet without strongly affecting body weight.

During the first burden, mice on LPD gained significantly less weight and were stunted

(Figure 5B-C; G3 and G4). However, upon refeeding with WD or MWD, mice quickly

gained weight and had a significant growth catch-up (Figure 5B-C; days 35 to 119).

Indeed, at P119 all groups of mice had a similar body length (Figure 5D). Feeding with

WD after control diet induced a significant gain in body weight compared to control-fed

animals (Figure 5E), but this was not the case after LPD. In contrast, MWD did not

induce a significant increase in body weight, even when mice were fed MWD from P14

onwards.

To assess whether LPD followed by WD or MWD dysregulates glucose homeostasis,

we evaluated glucose and insulin tolerance. At P91 or P119, after 6 hours of food

deprivation, no diet had a significant impact on glycemia (Figure 5F-G). It is

noteworthy that glycemia did not differ significantly between the two time points (two-

way ANOVA: effect of time, $P = 0.81$). When we performed an intraperitoneal glucose

tolerance test (GTT), we saw that, unlike WD, MWD feeding after from P14 onwards

(group 6), induced a significant glycemic increase in the early phase of the GTT (the so-

called glycemic excursion) (Figure 5H, G6). However, this was not the case when

MWD was given after P35 (Figure 5H, G2 and G4). Interestingly, while WD per se, or

given after chow feeding, did not modify glucose tolerance (Figure 5H, G1 and G5),

priming with LPD during early life induced a transient glucose intolerance (Figure 5H,

G3). As a whole, LPD-fed mice were susceptible to develop early-phase glucose

intolerance when they were refed with MWD, i.e., a suboptimal diet rich in fat and poor

in protein.

To infer if this phenotype was due to a change in insulin sensitivity, we performed an

insulin tolerance test (ITT). We saw that insulin tolerance was not modified by any of

the different diet combinations (Figure 5I), highlighting that the aforementioned glucose

intolerance was not due to impaired insulin sensitivity. Overall, we show that, while it

does not affect growth catch-up after protein malnutrition-mediated early life stunting,

suboptimal nutrition leads to impaired glucose control later in life.

Supplementation with Lp^{WJL} in suboptimal nutritional conditions further impairs

glucose tolerance after early-life protein restriction

In the same line as previously, we investigated the effect of a supplementation with

Lp^{WJL} during the second burden of the aforementioned protocol (i.e., from P35 to P119).

We found no effect of adding Lp^{WJL} on weight gain and systemic growth

(Supplementary Figure 2). Likewise, two-way ANOVA analysis did not reveal any

significant effect of Lp^{WJL} on organ size or weight (Supplementary Figure 3). We further

assessed glucose metabolism by performing a GTT and an ITT. We previously showed

that Lp^{WJL} treatment resulted in improved glucose tolerance during the post-weaning

period in chronic malnutrition mice (16). It is noteworthy that, in our experimental

setting, treatment with Lp^{WJL} did not affect glycemia after a 6-hour-long food

deprivation (Supplementary Figure 4; effect of Lp^{WJL}, $P = 0.0869$, two-way ANOVA).

Overall, glucose tolerance was not affected in mice fed control diet after weaning (G1

and G2 mice, Figure 6), nor mice that were fed WD or MWD all long (G5 and G6

mice). Surprisingly, mice facing a double burden of malnutrition, particularly those fed

WD after LPD (G3 mice), showed significant glucose intolerance when given Lp^{WJL}

during the second burden. Furthermore, Lp^{WJL} did not significantly affect insulin

tolerance in any of the groups (Figure 6). Overall, these findings indicate that Lp^{WJL}

treatment exacerbates glucose intolerance in a mouse model of a double burden of

malnutrition manifested by early protein malnutrition followed by western diet

nutrition.

DISCUSSION

PEM early in life has deadly consequences, being one of the most important causes of

infant death in the developing world (2). Maintenance of adequate nutrition and

rehabilitation are the most effective interventions to reduce the deleterious effects of

PEM (29). In this study, we aimed at establishing a mouse model of PEM, followed by a

nutritional rehabilitation, in order to explore the long-lasting consequences of early-life

PEM.

Early-life PEM results in stunting (30). In our protein malnutrition model, slow growth

and weight gain rate occurred after weaning, with over 50% reduction after 5 weeks on

LPD. In humans, the two main clinical syndromes of the extreme forms of children with

PEM are marasmus and kwashiorkor (31). While marasmus results from adaptation to

severe depletion of calories and nutrients, kwashiorkor ("sickness of the weaning" in Ga

language) is the consequence of a diet with inadequate protein but reasonably normal

caloric intake. Thus, our mouse model seems closer to kwashiorkor than marasmus.

However, it does not exhibit classic traits of kwashiorkor such as edema and

hepatomegaly (31), probably due to the fact that our mice are not in contact with any

enteric pathogens and thus are not susceptible to infection and inflammation. Indeed,

several studies have shown that malnourished mice are more susceptible to infection,

thus mimicking one of the hallmarks of human malnutrition (20,21). Furthermore,

nutritional rehabilitation after kwashiorkor results in increased IGF-1 levels, but the

values are lower than in healthy children (32). In our mouse model, IGF-1 levels are

fully restored by nutritional rehabilitation, suggesting that mice are more resilient than

humans to early-life nutritional stress. This is confirmed by the fact that, in our model,

mice are kept under severe protein malnutrition for several weeks after weaning. While

this results in a severe phenotype regarding growth and weight, our mice did not show

any particular sign of torpor, reduced activity, or change in appetite during the

nutritional challenge. Moreover, unlike Ferreira-Paes et al. (21), who describe a model

of marasmus using weanling BALB/c mice, nutritional rehabilitation in male mice did

not fully resolve the severity of protein malnutrition. Indeed, in our experimental

setting, male mice were stunted after nutritional rehabilitation, although early-life

wasting was corrected. It is noteworthy that the earlier the rehabilitation took place, the

better the response was. Nevertheless, even after post-natal day 56 (i.e., young adult

mice), growth occurred at a rate of 0.08 cm.day^{-1} after refeeding on an optimal diet,

showing the remarkable capacity of mice to adapt to deleterious nutritional

environments.

Recent advances in the knowledge of malnutrition have shown that the gut microbiota is

a contributor to the deleterious stunting and wasting phenotype (33). Indeed,

colonization of germ-free mice with kwashiorkor-associated microbiota in combination

with a Malawian diet leads to severe weight loss (34). Our previous work has shown

that modulating the microbiota of chronically malnourished weanling mice with Lp^{WJL}

stimulates growth via activation of the somatotropic axis (15,16). We hypothesized that

Lp^{WJL} would be able to stimulate growth during the refeeding phase in our PEM model,

thus compensating the stunting phenotype we had previously observed. In our

experimental setting, we saw no improvement of growth associated to Lp^{WJL} during the

refeeding phase. Thus, it is likely that the effects of the bacterium are associated to an

overall dietary challenge, with reduced protein and fat, rather than low-protein feeding

per se. In line with that, the model we previously described exhibited a marked decrease

of weight after weaning,

Overall, our results confirm the Developmental Origins of Health and Disease

(DOHaD) paradigm, which stipulates that developmental and early postnatal insults,

such as malnutrition, can have deleterious impacts in adult life (35). In line with that,

the DOHaD paradigm is particularly appropriate in understanding the nutritional

transitions in countries that face the double burden of malnutrition. Therefore, we

sought to establish a model with suboptimal refeeding after a moderately long period of

low-protein diet. It is well known that severe acute malnutrition affects glucose

homeostasis in infants (4,36), a feature that persists later in life (37,38). Furthermore,

studies in rodents have shown that post-weaning PEM induces hepatic alterations, such

as steatosis and impaired hepatic mitochondrial turnover (39–41). Interestingly, two

independent studies using a high-fat diet (58% kcal from fat) after LPD have shown that

PEM after weaning does not potentiate fat accumulation, hepatic steatosis and insulin

resistance in adult young mice (39,42). In their mouse model of double burden of

malnutrition, Dalvi et al. described that LPD feeding for 4 weeks post-weaning induces

transient glucose intolerance after a recovery on chow feeding (39), showing that LPD

priming induces deleterious effects later in life associated to poor metabolic health.

Indeed, in our model, while we saw no differences in growth with mice on MWD after

LPD, LPD priming also induces deficiencies in glucose control later in life. Although

results on mouse models are still inconsistent and fail to fully recapitulate the metabolic

alterations observed in malnourished humans, they provide key mechanistic insights

into the different pathologies associated to stunting in early life.

Indeed, cohorts of malnourished mice and humans have shown that growth and

metabolic functions are impaired, associated to alterations in the gut microbiota, and

that this impaired phenotype could be transferred (33). Similarly, extensive research has

shown that the overnourished microbiota is associated to loss of richness of taxa and

enrichment of bacteria associated to inflammation (43,44). While most studies have

been correlative on this issue, they have given an insightful look into the alterations of

microbial ecology in cases of malnutrition. Little is known on humans facing the double

burden of malnutrition. A 2018 study on a Mexican cohort of 36 patients

(undernourished, obese or controls) showed similarities in the composition of the

microbiota between the obese and undernourished group. In line with these findings, we

hypothesized that treatment with Lp^{WJL} could have a beneficial impact in mice facing a

double burden of malnutrition. Not only did we not see any beneficial effect of Lp^{WJL} in

this context, but the treatment induced further impairment of glucose tolerance in mice

fed a suboptimal Western diet. Thus, caution should be used when developing probiotic

strategies for PEM when infants also face a double burden of malnutrition.

Overall, we provide a new model of refeeding after early-life PEM, where we show sex-

associated differences in the long-term response to refeeding. We provide data showing

that suboptimal refeeding with a moderately low-protein diet, which is also rich in fat, is

sufficient to induce growth in malnourished mice, although this is associated to

metabolic perturbations. Further investigations into the specific mechanisms driving

these features will help uncover mechanisms associated to PEM and the double burden

of malnutrition.

ACKNOWLEDGEMENTS

This work was supported by funding from Fondation pour la Recherche Médicale

(Équipe FRM EQU202203014629). FDV was a recipient of a Young Researcher Prize

from Fondation des Treilles. The authors declare no conflict of interest associated to this

study.

JLT, FL, and FDV designed research; JLT, AC and DS conducted research; JLT and

FDV analyzed data; and JLT, FL and FDV wrote the paper. FDV had primary

responsibility for final content. All authors read and approved the final manuscript.

REFERENCES

1.   Baer MT, Harris AB. Pediatric nutrition assessment: identifying children at risk.
J Am Diet Assoc 1997;97:S107-115.
2.   2021 Global Nutrition Report [Internet]. [cited 2021 Dec 9]. Available from:
https://globalnutritionreport.org/
3.   Endrinikapoulos A, Afifah DN, Mexitalia M, Andoyo R, Hatimah I, Nuryanto
N. Study of the importance of protein needs for catch-up growth in Indonesian stunted
children: a narrative review. SAGE Open Med 2023;11:20503121231165562.
4.   Spoelstra MN, Mari A, Mendel M, Senga E, van Rheenen P, van Dijk TH,

Reijngoud D-J, Zegers RGT, Heikens GT, Bandsma RHJ. Kwashiorkor and marasmus are both associated with impaired glucose clearance related to pancreatic β-cell dysfunction. Metabolism 2012;61:1224–30.

5.   Bandsma RHJ, Mendel M, Spoelstra MN, Reijngoud D-J, Boer T, Stellaard F, Brabin B, Schellekens R, Senga E, Heikens GT. Mechanisms behind decreased endogenous glucose production in malnourished children. Pediatr Res 2010;68:423–8.

6.   Victora CG, Adair L, Fall C, Hallal PC, Martorell R, Richter L, Sachdev HS, Maternal and Child Undernutrition Study Group. Maternal and child undernutrition: consequences for adult health and human capital. Lancet Lond Engl 2008;371:340–57.

7.   Petry CJ, Ozanne SE, Hales CN. Programming of intermediary metabolism. Mol Cell Endocrinol 2001;185:81–91.

8.   Lind MV, Larnkjær A, Mølgaard C, Michaelsen KF. Dietary protein intake and quality in early life: impact on growth and obesity. Curr Opin Clin Nutr Metab Care 2017;20:71–6.

9.   Bhutta ZA, Berkley JA, Bandsma RHJ, Kerac M, Trehan I, Briend A. Severe childhood malnutrition. Nat Rev Dis Primer 2017;3:17067.

10.   Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, Ezzati M, Grantham-McGregor S, Katz J, Martorell R, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. Lancet Lond Engl 2013;382:427–51.

11.   Shrimpton R, Victora CG, de Onis M, Lima RC, Blössner M, Clugston G. Worldwide timing of growth faltering: implications for nutritional interventions. Pediatrics 2001;107:E75.

12.   Benjamin-Chung J, Mertens A, Colford JM, Hubbard AE, van der Laan MJ, Coyle J, Sofrygin O, Cai W, Nguyen A, Pokpongkiat NN, et al. Early-childhood linear growth faltering in low- and middle-income countries. Nature 2023;621:550–7.

13.   Mertens A, Benjamin-Chung J, Colford JM, Coyle J, van der Laan MJ, Hubbard AE, Rosete S, Malenica I, Hejazi N, Sofrygin O, et al. Causes and consequences of child growth faltering in low-resource settings. Nature 2023;621:568–76.

14.   Mertens A, Benjamin-Chung J, Colford JM, Hubbard AE, van der Laan MJ, Coyle J, Sofrygin O, Cai W, Jilek W, Rosete S, et al. Child wasting and concurrent stunting in low- and middle-income countries. Nature 2023;621:558–67.

15.   Schwarzer M, Makki K, Storelli G, Machuca-Gayet I, Srutkova D, Hermanova P, Martino ME, Balmand S, Hudcovic T, Heddi A, et al. Lactobacillus plantarum strain maintains growth of infant mice during chronic undernutrition. Science 2016;351:854–7.

16.   Schwarzer M, Gautam UK, Makki K, Lambert A, Brabec T, Joly A, Šrůtková D, Poinsot P, Novotná T, Geoffroy S, et al. Microbe-mediated intestinal NOD2 stimulation improves linear growth of undernourished infant mice. Science 2023;379:826–33.

17.   Dutta S, Sengupta P. Men and mice: Relating their ages. Life Sci 2016;152:244–8.

18.   Hintze KJ, Benninghoff AD, Ward RE. Formulation of the Total Western Diet (TWD) as a basal diet for rodent cancer studies. J Agric Food Chem 2012;60:6736–42.

19.   Monsanto SP, Hintze KJ, Ward RE, Larson DP, Lefevre M, Benninghoff AD. The new total Western diet for rodents does not induce an overweight phenotype or alter parameters of metabolic syndrome in mice. Nutr Res N Y N 2016;36:1031–44.

20.   Anstead GM, Chandrasekar B, Zhao W, Yang J, Perez LE, Melby PC. Malnutrition alters the innate immune response and increases early visceralization

following Leishmania donovani infection. Infect Immun 2001;69:4709–18.

21. Ferreira-Paes T, Seixas-Costa P, Almeida-Amaral EE. Validation of a Feed
Protocol in a Mouse Model That Mimics Marasmic Malnutrition. Front Vet Sci
2021;8:757136.

22. Heude B, Scherdel P, Werner A, Le Guern M, Gelbert N, Walther D, Arnould
M, Bellaïche M, Chevallier B, Cheymol J, et al. A big-data approach to producing
descriptive anthropometric references: a feasibility and validation study of paediatric
growth charts. Lancet Digit Health 2019;1:e413–23.

23. Gomez F, Galvan RR, Frenk S, Munoz JC, Chavez R, Vazquez J. Mortality in
second and third degree malnutrition. J Trop Pediatr Lond Engl 1955 1956;2:77–83.

24. Body Weight Information for C57BL/6J (000664) [Internet]. The Jackson
Laboratory. [cited 2023 Jul 21]. Available from: https://www.jax.org/jax-mice-and-
services/strain-data-sheet-pages/body-weight-chart-000664

25. Yakar S, Isaksson O. Regulation of skeletal growth and mineral acquisition by
the GH/IGF-1 axis: Lessons from mouse models. Growth Horm IGF Res Off J Growth
Horm Res Soc Int IGF Res Soc 2016;28:26–42.

26. Yakar S, Rosen CJ, Beamer WG, Ackert-Bicknell CL, Wu Y, Liu J-L, Ooi GT,
Setser J, Frystyk J, Boisclair YR, et al. Circulating levels of IGF-1 directly regulate
bone growth and density. J Clin Invest 2002;110:771–81.

27. Smith WJ, Underwood LE, Clemmons DR. Effects of caloric or protein
restriction on insulin-like growth factor-I (IGF-I) and IGF-binding proteins in children
and adults. J Clin Endocrinol Metab 1995;80:443–9.

28. Popkin BM, Corvalan C, Grummer-Strawn LM. Dynamics of the double burden
of malnutrition and the changing nutrition reality. Lancet Lond Engl 2020;395:65–74.

29. Bhutta ZA, Das JK, Rizvi A, Gaffey MF, Walker N, Horton S, Webb P, Lartey
A, Black RE, Lancet Nutrition Interventions Review Group, the Maternal and Child
Nutrition Study Group. Evidence-based interventions for improvement of maternal and
child nutrition: what can be done and at what cost? Lancet Lond Engl 2013;382:452–77.

30. de Onis M, Branca F. Childhood stunting: a global perspective. Matern Child
Nutr 2016;12 Suppl 1:12–26.

31. Grover Z, Ee LC. Protein energy malnutrition. Pediatr Clin North Am
2009;56:1055–68.

32. Kouanda S, Doulougou B, De Coninck V, Habimana L, Sondo B, Tonglet R,
Ketelslegers JM, Robert A. Insulin Growth Factor-I in Protein-Energy Malnutrition
during Rehabilitation in Two Nutritional Rehabilitation Centres in Burkina Faso. J Trop
Med 2009;2009:832589.

33. Blanton LV, Barratt MJ, Charbonneau MR, Ahmed T, Gordon JI. Childhood
undernutrition, the gut microbiota, and microbiota-directed therapeutics. Science
2016;352:1533.

34. Smith MI, Yatsunenko T, Manary MJ, Trehan I, Mkakosya R, Cheng J, Kau AL,
Rich SS, Concannon P, Mychaleckyj JC, et al. Gut microbiomes of Malawian twin pairs
discordant for kwashiorkor. Science 2013;339:548–54.

35. Uauy R, Kain J, Corvalan C. How can the Developmental Origins of Health and
Disease (DOHaD) hypothesis contribute to improving health in developing countries?
Am J Clin Nutr 2011;94:1759S-1764S.

36. Bandsma RHJ, Spoelstra MN, Mari A, Mendel M, van Rheenen PF, Senga E,
van Dijk T, Heikens GT. Impaired glucose absorption in children with severe
malnutrition. J Pediatr 2011;158:282-287.e1.

37.   Ferdous F, Filteau S, Schwartz NB, Gumede-Moyo S, Cox SE. Association of
postnatal severe acute malnutrition with pancreatic exocrine and endocrine function in
children and adults: a systematic review. Br J Nutr 2022;129:1–34.
38.   Thompson DS, Bourdon C, Massara P, Boyne MS, Forrester TE, Gonzales GB,
Bandsma RHJ. Childhood severe acute malnutrition is associated with metabolic
changes in adulthood. JCI Insight 2020;5.
39.   Dalvi PS, Yang S, Swain N, Kim J, Saha S, Bourdon C, Zhang L, Chami R,
Bandsma RHJ. Long-term metabolic effects of malnutrition: Liver steatosis and insulin
resistance following early-life protein restriction. PloS One 2018;13:e0199916.
40.   van Zutphen T, Ciapaite J, Bloks VW, Ackereley C, Gerding A, Jurdzinski A, de
Moraes RA, Zhang L, Wolters JC, Bischoff R, et al. Malnutrition-associated liver
steatosis and ATP depletion is caused by peroxisomal and mitochondrial dysfunction. J
Hepatol 2016;65:1198–208.
41.   Hu G, Ling C, Chi L, Thind MK, Furse S, Koulman A, Swann JR, Lee D, Calon
MM, Bourdon C, et al. The role of the tryptophan-NAD + pathway in a mouse model
of severe malnutrition induced liver dysfunction. Nat Commun 2022;13:7576.
42.   Branco RCS, Camargo RL, Batista TM, Vettorazzi JF, Lubaczeuski C, Bomfim
LHM, Silveira LR, Boschero AC, Zoppi CC, Carneiro EM. Protein malnutrition
mitigates the effects of a high-fat diet on glucose homeostasis in mice. J Cell Physiol
2019;234:6313–23.
43.   Turnbaugh PJ, Bäckhed F, Fulton L, Gordon JI. Diet-induced obesity is linked to
marked but reversible alterations in the mouse distal gut microbiome. Cell Host
Microbe 2008;3:213–23.
44.   Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An
obesity-associated gut microbiome with increased capacity for energy harvest. Nature
2006;444:1027–31.
556

FIGURE LEGENDS

Figure 1: Chronically malnourished juvenile male and female mice differ in their

dynamics of growth.

Body length (**A, B**) and weight (**C, D**) of male and female control diet- or low-protein

diet-fed mice refed with control diet respectively at post-natal day 28 (RF28), 35

(RF35), 42 (RF42), 49 (RF49), or 56 (RF56). Arrows correspond to the day of

refeeding. (**E, F**) Length-versus-weight of control- and low-protein diet- (LPD) fed

animals. (**G, H**) Length-versus-weight of control and refed mice, after the time of

refeeding.

SD: standard deviation of control group.

Data are mean ± SEM.

Figure 2: Optimal refeeding after long-term juvenile malnutrition is insufficient to

fully recover systemic growth in males, but not in females.

(A, B) Weight-for-age relationship of of male and female control diet- or low-protein

diet-fed mice refed with control diet respectively at post-natal day 28 (RF28), 35

(RF35), 42 (RF42), 49 (RF49), or 56 (RF56). Arrows correspond to the day of

refeeding. **(C, D)** Body length evolution during the two weeks following refeeding or

from day 21 to day 35 in control mice.

Data are mean ± SEM.

Figure 3: Optimal refeeding after early-life malnutrition in males is insufficient to

fully recover organ growth.

Weight of liver **(A)**, subcutaneous fat **(B)**, perigonadal fat **(C)**, and soleus muscle **(D)**;

length of femur **(E)**, tibia **(F)**, small intestine **(G),** and colon **(H)** of male control diet- or

low-protein diet-fed mice refed with control diet respectively at post-natal day 28

(RF28), 35 (RF35), 42 (RF42), 49 (RF49), or 56 (RF56). **(I)** Principal Component

Analysis of parameters measured in **A** to **H**. **(J)** Serum concentration of IGF-1 measured

at the switch of diet, 2 weeks after switch and at the end of the experiment in

aforementioned mice.

Data are mean ± SEM.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; *** $P < 10^{-4}$, one-way ANOVA followed by

Tukey's post-hoc test.

Figure 4: Lp^{WJL} treatment during the refeeding phase fails to improve systemic

growth.

Body length **(A)** and weight **(B)**; weight of liver **(C)**; fat content **(D)** ; weight of

subcutaneous fat **(E)**, perigonadal fat **(F)**, and soleus muscle **(G)**; length of femur **(H)**,

tibia **(I)**, small intestine **(J)**, and colon **(K)** of male mice refed at post-natal day 56

(RF56) or maintained on low-protein diet (LPD) and treated with *Lactiplantibacillus*

plantarum WJL *(*Lp^{WJL}) or a placebo. **(L)** Principal Component Analysis of parameters

measured in **C** and **E** to **K**.

Data are mean ± SEM.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; *** $P < 10^{-4}$, two-way ANOVA followed by

Šídák's post-hoc test.

Figure 5: Suboptimal refeeding with a non-obesogenic Western diet compensates

early life-associated stunting, but impairs glucose metabolism.

(A) Experimental procedure . Body length **(B)** and weight **(C)** of mice described in

panel **A**. Length **(D)** and weight **(E)** at post-natal day 119. Plasma glucose after 6 hours

of food deprivation at post-natal day 91 **(F)** and 112 **(G)** of of mice described in panel

A. Intraperitoneal glucose tolerance test **(H)** and insulin tolerance test **(I)** performed at

post-natal day 91 and 112 respectively on mice described in panel **A**. Inset: area under

the curve (AUC) of the respective tests.

Data are mean ± SEM.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; *** $P < 10^{-4}$, one-way ANOVA followed by

Tukey's post-hoc test (when not indicated, value vs. control).

Figure 6: Supplementation with Lp^{WJL} in suboptimal conditions further impairs

glucose tolerance after early-life protein restriction.

Intraperitoneal glucose tolerance test (GTT) and insulin tolerance test (ITT) performed

respectively at P91 and P112 on male mice treated with *Lactiplantibacillus plantarum*

WJL *(*Lp^{WJL}*)* or a placebo from P35 to P119. Experimental groups are described in

Figure 5**A**.

Data are mean ± SEM.

* $P < 0.05$; ** $P < 0.01$, Student's *t*-test with Welch's correction.