1

Astrocytes mediate two forms of spike timing-dependent depression at entorhinal cortex-hippocampal synapses

Heriberto Coatl Cuaya*[1], Irene Martínez-Gallego*[1] and Antonio Rodríguez-Moreno[1]

[1]Laboratory of Cellular Neuroscience and Plasticity, Department of Physiology, Anatomy and Cell Biology, Universidad Pablo de Olavide, ES-41013 Seville, Spain.

*These authors contributed equally to this work

Address for correspondence: Prof. Antonio Rodríguez-Moreno, Laboratory of Cellular Neuroscience and Plasticity, Department of Physiology, Anatomy and Cell Biology, Universidad Pablo de Olavide, ES-41013 Seville, Spain. E-mail: arodmor@upo.es; Phone: +34 954977393

Running title: STDP at entorhinal cortex-dentate gyrus synapses

Key words: Plasticity, t-LTD, entorhinal cortex, dentate gyrus, hippocampus, medial and lateral perforant pathways, astrocytes, NMDARs.

Abstract

The entorhinal cortex (EC) connects to the hippocampus sending different information from cortical areas that is first processed at the dentate gyrus (DG) including spatial, limbic, and sensory and information. Excitatory afferents from lateral (LPP) and medial (MPP) perforant pathways of the EC connecting to granule cells of the DG play a role in memory encoding and information processing and are deeply affected in humans suffering Alzheimer´s disease and temporal lobe epilepsy, contributing to the dysfunctions found in these pathologies. The plasticity of these synapses is not well known yet, as are not known the forms of long-term depression (LTD) existing at those connections. We investigated whether spike timing-dependent long-term depression (t-LTD) exists at these two different EC-DG synaptic connections in mice, and whether they have different action mechanisms. We have found two different forms of t-LTD, at LPP- and MPP-GC synapses and characterised their cellular and intracellular mechanistic requirement. We found that both forms of t-LTD are expressed presynaptically and that whereas t-LTD at LPP-GC synapses does not require NMDAR, t-LTD at MPP-GC synapses requires ionotropic NMDAR containing GluN2A subunits. The two forms of t-LTD require different group I mGluR, mGluR5 LPP-GC synapses and mGluR1 MPP-GC synapses. In addition, both forms of t-LTD require postsynaptic calcium, eCB synthesis, CB1R, astrocyte activity, and glutamate. Thus, we discovered two novel forms of t-LTD that require astrocytes at EC-GC synapses.

Introduction

The entorhinal cortex (EC) conveys spatial, limbic and sensory information to the hippocampus, a structure responsible for important activities in the brain including learning, memory and spatial information coding. Excitatory projections from the EC to the dentate gyrus (DG) play a role in memory encoding, and they are affected in humans suffering from Alzheimer Disease (AD) and temporal lobe epilepsy, contributing critically to these pathologies (Paré et al., 1992; Llorens-Martín et al., 2014). In rodents, evidence suggests that the lateral perforant pathway (LPP) conveys more sensory related information, whereas the medial perforant pathway (MPP) conveys information related to spatial location and limbic signals that are involved in attention and motivation. Evidence suggests that LPP- and MPP-GC synapses have distinct electrophysiological properties, these differences potentially affecting the functional processing of information (Krueppel et al., 2011; Petersen et al., 2013; Kim et al., 2018). Indeed, these functional differences probably reflect their distinct inputs, as they contact the outer (LPP) or middle (MPP) third of the molecular layer of the DG (Paré et al., 1992; Burgess et al., 2002). Significantly, MPP-GC synapses are associated with a higher probability of release (Min et al., 1998) and the existence and/or activity of presynaptic NMDARs, which differs between the two types of fibres (Savtchouk et al., 2019). Interestingly, LPP and MPP synapses appear to influence pathological changes distinctly, with the former showing changes in release probability in models of epilepsy (Scimemi et al., 2006), preferential β-amyloid deposition (Reilly et al., 2003), earlier susceptibility to AD (Khan et al., 2014) and reduced plasticity with age (Froc et al., 2003). The mechanistic bases of these differences are virtually unknown and they pose a challenge when attempting to understand circuit-specific biological computations, as well as the susceptibility to pathological insults. In addition, the forms of plasticity at these synapses remain unclear and the limited studies into plasticity available did not distinguish between these two pathways.

Plasticity drives the organization of cortical maps during development, and it underlies learning and memory (reviewed in Malenka and Bear, 2004). The most extensively studied forms of plasticity are long-term potentiation (LTP) and long-term depression (LTD) of synaptic transmission. Spike timing-dependent plasticity (STDP) is a Hebbian form of long-term synaptic plasticity found in all known nervous systems, from insects to humans. Indeed, STDP appears to be a synaptic mechanism that

underlies circuit remodelling during development, and it is thought to be involved in learning and memory (Dan and Poo, 2004; Feldman, 2012). In STDP, the relative order and millisecond timing of pre- and postsynaptic action potentials (spikes/APs) determine the direction and magnitude of the synaptic changes. Thus, spike timing-dependent LTP (t-LTP) occurs when a postsynaptic spike follows a presynaptic spike within ms, whereas spike timing-dependent LTD (t-LTD) is induced when this order is reversed (Feldman, 2012). In addition, *in vivo* studies have suggested modifications to LTP and LTD after mice undertake behavioural tasks (Kemp and Manahan-Vaughan, 2004; Clarke et al., 2010). However, it is surprising that while plasticity of EC-DG granule cell (GC) synapses has been studied using classical protocols (Froc et al., 2003; Harney et al., 2006; Hayashi et al., 2014, Savtchouk et al., 2019), no information exists as to whether STDP or t-LTD can be induced at these synapses.

Here we studied STDP at PP-GC synapses, differentiating the lateral and medial pathways, determining whether these individual synapses are subject to t-LTD and if this form of plasticity at these synapses influences learning, memory or information processing. As a result, we identified two different forms of presynaptic t-LTD at LPP- and MPP-GC synapses. Unlike LPP-GC synapses, the t-LTD at MPP-GC synapses requires ionotropic NMDARs that contain GluN2A subunits, and the t-LTD at each type of synapse require different group I mGluRs, with LPP-GC synapses dependent on mGluR5 and MPP-GC t-LTD requiring mGluR1. In addition, we found that both forms of t-LTD require postsynaptic calcium, endocannabinoid (eCB) synthesis, CB_1 receptor (CB_1R) signalling, astrocytes and glutamate. Hence, we describe here two novel forms of presynaptic t-LTD in the brain at EC-GC synapses with different action mechanisms, probably implicated in the different aspects of information processing.

Materials and Methods

Animals and ethical approval

All animal procedures were conducted in accordance with the European Union Directive 2010/63/EU regarding the protection of animals used for scientific purposes, and they were approved by the Ethics Committee at the Universidad Pablo de Olavide and that of the Andalusian Government. C57BL/6 mice were purchased from Harlan Laboratories (Spain) and P13–21 (P, post-natal day) mice of either sex were used. Animals were kept at temperatures between 18 and 24 °C on a continuous 12 h

light/dark cycle, and at 40–60% humidity, with ad libitum access to food and water. In

some experiments, dominant-negative (dn) SNARE mice (Pascual et al., 2005) of the

same age were used. These mice were not fed with doxycycline from birth, the

supplement that drives transgene expression. In these mice, the human glial fibrillary

acidic protein (hGFAP) promoter mediates the specific expression of the tetracycline

transactivator (tTA) in astrocytes, which in turn activates the tetracycline responsive

element operator and drives the cytosolic expression of VAMP2/synaptobrevin II in

these cells, along with the enhanced green fluorescence protein (eGFP). Expression of

the dnSNARE transgene interferes with the formation of the SNARE complex, blocking

exocytosis and impairing vesicle release by astrocytes (Sultan et al., 2015).

Slice preparation. Hippocampal slices were prepared as described previously

(Andrade-Talavera et al., 2016; Pérez-Rodríguez et al., 2019; Falcón-Moya et al., 2020).

Briefly, mice were anesthetized with isoflurane (2%), decapitated, and their whole brain

was removed and placed in an ice-cold ACSF solution (pH 7.2, 300 mOsm*L^{-1})

containing (in mM): 126 NaCl, 3 KCl, 1.25 NaH$_2$PO$_4$, 2 MgSO$_4$, 2 CaCl$_2$, 26 NaHCO$_3$,

and 10 glucose. Slices (350 μm thick: Leica VT1000S Vibratome) were maintained

oxygenated (95% O$_2$/5% CO$_2$) in the same solution for at least 1 h before use.

Experiments were carried out at 33-34 °C and during the experiments; the slices were

superfused continuously with the same solution indicated above.

Electrophysiological recordings. Whole-cell patch-clamp recordings of DG

GCs were obtained under visual guidance by infrared differential interference contrast

microscopy. The identity of the neurons was verified by their characteristic voltage

response to a current step protocol applied in current-clamp configuration using a patch-

clamp amplifier (Multiclamp 700B), and acquiring the data with pCLAMP 10.2

software (Molecular Devices). Patch-clamp electrodes were pulled from borosilicate

glass tubes and had a resistance of 4-7 MΩ when filled with the intracellular solution

consisting in(in mM, pH 7.2–7.3, 290 mOsm L^{-1}): potassium gluconate, 110; HEPES,

40; NaCl, 4; ATP-Mg, 4; and GTP, 0.3. Only cells with a stable resting membrane

potential below −55 mV were assessed, and the cell recordings were excluded from the

analysis if the series resistance changed by >15%. All recordings were low-pass filtered

at 3 kHz and acquired at 10 kHz. For plasticity experiments, excitatory postsynaptic

potentials (EPSPs) were evoked alternately at 0.2 Hz in two input pathways, test and

control, by using two monopolar stimulating electrodes placed 200-400 μm from the

cell soma. Stimulating electrodes were situated on fibres contacting the distal and medial third of the dendritic arbour of GCs, corresponding to LPP- and MPP-GC synapses, respectively. Stimulation was adjusted (200 μs, 0.1-0.2 mA) to obtain an EPSP peak amplitude of 3-5 mV in control conditions. Pathway independence was ensured by the lack of cross-facilitation when the pathways were stimulated alternately at 50 ms intervals. Plasticity was assessed through the changes in the EPSP slope, measured in its rising phase as a linear fit between time points corresponding to 25-30% and 70-75% of the peak amplitude under control conditions. Miniature responses were recorded in the presence of tetrodotoxin (TTX, 300 nM).

Plasticity protocols. After establishing a stable basal EPSP over 10 min, the test input was paired 100 times with a single postsynaptic spike. The single postsynaptic spike was evoked by a brief somatic current pulse (5 ms, 0.1–0.5 pA), whereas the control pathway was left unstimulated during the pairing period. To induce t-LTD, a postsynaptic AP was evoked 18 ms before the onset of the EPSP. EPSP slopes were monitored for at least 30 min after the pairing protocol, and the presynaptic stimulation frequency remained constant throughout the experiment. In some experiments, "glutamate puffs" were applied using a Picospritzer (Parker Hannifin), expulsing glutamate dissolved in the external solution through a micropipette over an astrocyte in proximity to the LPP-GC or MPP-GC synapses at a pressure of 10 psi for 50–200 ms. In each experiment, 100 glutamate puffs were applied to the recording neuron at 0.2 Hz at baseline, 18 ms before the onset of the EPSP, which did not affect patch clamping. The EPSP slopes were then monitored for 30 min.

Drugs. The following agents were purchased from: Sigma-Aldrich - BAPTA, Zinc chloride, and the compounds used to prepare the ACSF and current clamp internal solutions; and Tocris Bioscience - MK-801 maleate, D-AP5, TTX, PPDA, Ro 25-6981 maleate, L-glutamic acid, MPEP, LY367385, LY341495, AM251, Orlistat (THL), FK506, Nimodipine, Thapsigargin and Ifenprodil. All compounds were dissolved in distilled water except for Nimodipine, Thapsigargin, PPDA, FK506, AM251 and THL, which were dissolved in dimethyl sulfoxide (DMSO).

Data analysis. The data were analysed using the Clampfit 10.2 software (Molecular Devices) and the last 5 min of the recordings were used to estimate the changes in synaptic efficacy relative to the baseline. For the paired-pulse ratio (PPR) experiments,

two EPSPs were evoked for 30 s at the baseline frequency, one at the beginning of the

baseline recording (40 ms apart) and another 30 min after the end of the pairing

protocol. The PPR was expressed as the slope of the second EPSP divided by the slope

of the first EPSP. A Coefficient of Variation (CV) analysis was carried out on the EPSP

slopes (Rodríguez-Moreno and Paulsen, 2008; Falcón-Moya et al., 2020) and the noise-

free CV of the EPSP slopes was calculated as:

$$CV = \sqrt{\frac{\sigma^2(EPSP) - \sigma^2(noise)}{EPSP\ slope}}$$

where σ^2 (EPSP) and σ^2 (noise) are the variance of the EPSP and baseline, respectively.

The plot compares the variation in the mean EPSP slope (M) to the change in response

variance of the EPSP slope ($1/CV^2$: see Brock et al., 2020 for a comprehensive

explanation). Graphs were prepared using SigmaPlot 11.0.

Statistical analysis. Before applying any statistical comparison, the data were subjected

to Shapiro-Wilk normality and equal variance tests, applying a confidence interval (CI)

of 95%. Unpaired or paired Student's t-tests were used for comparisons between two

groups. For multiple comparisons of more than two groups, one-way ANOVA was used

followed by a Holm–Sidak *post-hoc* test (when necessary). All the data are expressed as

the mean ± standard error of the mean (SEM) and P values were considered significant

when < 0.05: *p = 0.05, **p = 0.01, ***p = 0.001.

Results

Pairing presynaptic activity with single postsynaptic action potentials at low frequency

can induce t-LTD at mouse lateral and medial perforant pathway-dentate gyrus granule

cell synapses

We initially determined whether spike timing-dependent LTD can be induced at

perforant pathway-DG GC synapses by pairing presynaptic stimulation with single

postsynaptic spikes at low frequency (0.2 Hz). We used a post-pre protocol, pairing a

single postsynaptic spike followed 18 ms later by presynaptic stimulation 100 times, as

described previously at cortical and hippocampal synapses (Andrade-Talavera et al.,

2016; Martínez-Gallego et al., 2022a). We monitored the EPSPs evoked by extracellular

stimulation of the LPP- and MPP-GC synapses, which make synaptic contacts with the distal and medial third of the dendritic arbour of hippocampal DG GCs, respectively (Petersen et al., 2013; Savtchouk et al., 2019). EPSPs were recorded from GCs using the current clamp mode of the whole-cell patch-clamp configuration (Fig. 1a, k). A post-before-pre pairing protocol induced robust t-LTD at both LPP-GC (63 ± 5 %, n = 14: Fig 1b, c) and MPP-GC synapses (59 ± 5 %, n = 14: Fig. 1l, m), while unpaired control pathways remained unaffected (103 ± 4 % and 105 ± 7 %, respectively: Fig. 1b, c, l, m).

Presynaptic expression of t-LTD at lateral and medial perforant pathway-dentate gyrus granule cell synapses

We next determined the site where this t-LTD was expressed through several approaches. As we observed failures in synaptic transmission at both types of synapse, we analysed these and found that the number of failures increased consistently after t-LTD at both LPP-GC (from 43 ± 7 % at baseline to 76 ± 6 % after t-LTD, n = 8: Fig. 1d) and MPP-GC synapses (from 40 ± 5 % at baseline to 59 ± 4 % after t-LTD, n = 6: Fig. 1n), suggesting a presynaptic mechanism. We then analysed the PPR at baseline and 30 min after the pairing protocol, identifying a significant increase in the PPR after t-LTD at both LPP-GC (from 1.16 ± 0.11 at baseline to 1.67 ± 0.19 after t-LTD, n = 14: Fig. 1e) and MPP-GC synapses (from 1.17 ± 0.11 at baseline to 1.81 ± 0.15 % after t-LTD, n = 14: Fig. 1o), again suggesting a presynaptic expression of these two forms of t-LTD. Third, we estimated the noise-subtracted CV of the synaptic responses before and after t-LTD induction. A plot of CV^{-2} versus the change in the mean evoked EPSP slope (M) before and after t-LTD mainly yielded points below the diagonal line at LPP-GC and MPP-GC synapses (Fig. 1f, p), consistent with a modification of the release parameters (Rodríguez-Moreno and Paulsen, 2008; Brock et al., 2020). Finally, we recorded and analysed the miniature responses (mEPSP) in the presence of tetrodotoxin (300 nM, TTX), before and after t-LTD, adding TTX at the baseline and washing it out before performing the t-LTD experiment, and then adding TTX again after t-LTD (Falcón-Moya et al., 2020). Under these conditions, a t-LTD similar to previous experiments was evident at both types of synapse (LPP-GC synapses 55 ± 11 %, n = 6; MPP-GC synapses 67 ± 8 %, n = 7), and we found that the frequency of mEPSPs decreased at both types of synapse after t-LTD induction: LPP-GC - baseline 1.4 ± 0.13 Hz, after t-LTD induction 0.7 ± 0.09 Hz, n = 6 (Fig. 1g, i); and MPP-GC - baseline 1.2

± 0.1 H, after t-LTD induction 0.7 ± 0.07 Hz, n = 8 (Fig. 1q, s). No effect of TTX on mEPSP amplitude was evident at any of these synapses: LPP-GC - baseline 1.13 ± 0.09, after t-LTD 1.13 ± 0.02 mV, n = 5 (Fig. 1h, j); MPP-GC - baseline 1.03 ± 0.07, after t-LTD 1.00 ± 0.09 mV, n = 6 (Fig. 1r, t). These results again suggest a presynaptic locus for t-LTD expression. Hence, together these results are consistent with a decrease in the probability of neurotransmitter release and they are indicative of a presynaptic locus for these two forms of t-LTD at both types of synapse.

Presynaptic t-LTD requires NMDARs at MPP-GC but not at LPP-GC synapses

Different forms of t-LTD have been shown to require NMDARs at cortical synapses and in the hippocampus (Bender et al., 2006; Nevian and Sakman, 2006; Rodríguez-Moreno and Paulsen, 2008; Rodríguez-Moreno et al., 2011, 2013; Andrade-Talavera et al., 2016). To determine whether the two forms of t-LTD studied here require NMDARs, we treated slices with the NMDAR antagonists D-AP5 (50 μM) or MK-801 (500 μM), completely blocking NMDARs and NMDAR-mediated currents (from 57 ± 8 pA, n = 5). In these experimental conditions, t-LTD at LPP-GC synapses was not affected by these antagonists (D-AP5 66 ± 4 %, n = 8; MK-801 62 ± 9 %, n = 6; interleaved controls 60 ± 6 %, n = 8: Fig. 2a, b), whereas t-LTD at MPP-GC synapses was blocked in slices treated with either D-AP5 (90 ± 7 %, n = 10) or MK-801 (107 ± 10 %, n = 6) relative to the interleaved controls (60 ± 7 %, n = 8: Fig. 2e, f). When the experiment was repeated by loading the postsynaptic cell with MK-801 (500 μM-1 mM), t-LTD at MPP-GC synapses was not affected (54 ± 7 %, n = 9) relative to the interleaved controls (60 ± 7 %, n = 8: Fig. 2e, f). These results indicate that while t-LTD at MPP-GC synapses requires NMDARs, t-LTD at LPP-GC synapses does not. Moreover, the NMDARs required for t-LTD at MPP-GC synapses are ionotropic and not postsynaptic, consistent with data from neocortical synapses where presynaptic NMDARs are required for t-LTD (Sjöström et al., 2003; Bender et al., 2006; Nevian and Sakmann, 2006; Rodríguez-Moreno and Paulsen, 2008).

Presynaptic NMDARs containing GluN2B and GluN3A subunits have been found at MPP-GC synapses and they modulate glutamate release (Jourdain et al., 2007; Savtchouk et al., 2019). To determine the subunit composition of the NMDARs involved in t-LTD at MPP-GC synapses, we first repeated the experiments in slices treated with Zn^{2+} (300 nM), an antagonist of NMDARs containing GluN2A subunits

(Bidoret et al., 2009, Andrade-Talavera et al., 2016; Prius-Mengual et al., 2019) that prevented t-LTD (95 ± 5 %, n = 7: Fig. 2g, h). We also repeated the experiments on slices in the presence of Ro25-6981 (0.5 μM), an antagonist of NMDARs containing GluN2B subunits (Hu et al., 2009; Mares et al., 2021). This antagonist did not affect t-LTD (55 ± 11 %, n = 6: Fig. 2g, h) and neither did an additional antagonist of NMDARs containing GluN2B subunits, ifenprodil (3 μM, 62 ± 11 %, n = 6: Fig. 2g, h). Finally, t-LTD was not affected in the presence of PPDA (10 μM, 61 ± 7 %, n = 6: Fig. 2g, h) relative to the pooled interleaved controls (63 ± 6 %, n = 9), an antagonist of NMDARs that contain GluN2C/2D subunits (Chen et al., 2021; Wang et al., 2020). These results indicate that the presynaptic NMDARs involved in t-LTD induction at MPP-GC synapses probably contain GluN1/GluN2A subunits.

Presynaptic t-LTD at lateral and medial perforant pathway-dentate gyrus granule cell synapses requires mGluRs

As mGluRs have been implicated in plasticity and t-LTD in different brain regions, and at distinct synapses (Gladding et al., 2009), and they may modulate synaptic transmission at PP-DG synapses (Dietrich et al., 1997), we tested whether the presynaptic forms of t-LTD at LPP- and MPP-GC synapses require mGluRs. Exposure to the broad-spectrum mGluR antagonist LY341495 (100 μM) completely blocked t-LTD at both LPP-GC (100 ± 10 %, n = 8: Fig. 2c, d) and MPP-GC synapses (99 ± 9 %, n = 10: Fig. 2i, j). Indeed, the mGluR5 antagonist MPEP (20 μM) prevented t-LTD at LPP-GC (103 ± 10 %, n = 8: Fig. 2c, d) but not at MPP-GC synapses (61 ± 7 %, n = 6: Fig. 2i, j), whereas the specific mGluR1 antagonist LY367385 (100 μM) did not affect t-LTD at LPP-GC synapses (76 ± 3 %, n = 6: Fig. 2c, d) but it did block t-LTD at MPP-GC synapses (95 ± 7 %, n = 7: Fig. 2i, j) relative to their respective interleaved controls (LPP-GC synapses 68 ± 7 %, n = 9; MPP-GC synapses 65 ± 7 %, n = 9: Fig. 2c, d, i, j). Hence, both LPP-GC and MPP-GC synapses require mGluR activity to induce these presynaptic forms of t-LTD, but whereas t-LTD at LPP-GC synapses requires mGluR5, mGluR1 receptors are required at MPP-GC synapses.

Postsynaptic Ca^{2+} is required for t-LTD at lateral and medial perforant pathway-dentate gyrus granule cell synapses

Both t-LTP and t-LTD require postsynaptic Ca^{2+} at neocortical and hippocampal

synapses (Bender et al., 2006; Nevian and Sakmann, 2006; Rodríguez-Moreno et al.,

2013; Andrade-Talavera et al., 2016; Inglebert et al., 2021; Martínez-Gallego et al.,

2022a), which led us to investigate the postsynaptic Ca^{2+} requirements of t-LTD at PP-

DG synapses. As L-type Ca^{2+} channels have previously been implicated in plasticity

(Nevian and Sakmann, 2006; Andrade-Talavera et al., 2016; Wiera et al., 2017; Falcón-

Moya et al., 2020; Martínez-Gallego et al., 2022a), we assessed whether they are

involved in the t-LTD studied here by performing the pairing protocol after bath

application of the L-type Ca^{2+} channel blocker, nimodipine (10 μm). This channel

blocker did not impede t-LTD induction at LPP-GC (79 ± 4 %, n = 6: Fig. 3a, b) or

MPP-GC synapses (67 ± 6 %, n = 6: Fig. 3e, f), indicating that L-type Ca^{2+} channels are

not required for these forms of t-LTD. The Ca^{2+} involved in synaptic plasticity may also

be released from intracellular stores (Bender et al., 2006; Nevian and Sakmann, 2006;

Andrade-Talavera et al., 2016; Martínez-Gallego et al., 2022a) and indeed, t-LTD was

prevented at both LPP-GC and MPP-GC synapses when thapsigargin (10 μM) was

present in the superfusion fluid (LPP-GC synapses 108 ± 16 %, n = 6; MPP-GC

synapses108 ± 6 %, n = 6), a drug that depletes intracellular Ca^{2+} stores (Singh et al.,

2021). A similar effect was witnessed when thapsigargin (10 μM) was loaded into the

postsynaptic neuron (LPP-GC synapses 109 ± 12 %, n = 9; MPP-GC synapses 107 ± 5

%, n = 7) relative to untreated interleaved controls (LPP-GC synapses 66 ± 7 %, n = 9;

MPP-GC synapses 67 ± 7, n = 9: Fig. 3a, b, e, f). Hence, Ca^{2+} release from the

intracellular stores appears to be necessary for t-LTD at both LPP and MPP-GC

synapses.

Postsynaptic calcineurin is required for t-LTD induction at LPP-GC but not MPP-GC
synapses

To gain better insight into the mechanisms involved in t-LTD, we hypothesized

that a Ca^{2+}-dependent enzyme might be involved. Since the Ca^{2+}-dependent protein

phosphatase calcineurin has been implicated in synaptic plasticity in the hippocampus

(Mulkey et al., 1994) and neocortex (Torii et al., 1995; Rodríguez-Moreno et al., 2013;

Andrade-Talavera et al., 2016), we tested the effect of inhibiting calcineurin on t-LTD.

Accordingly, t-LTD induction was prevented at LPP-GC (110 ± 7 %, n = 8) but not

MPP-GC synapses (72 ± 7 %, n = 6: Fig. S1) when the calcineurin blocker FK506 (10

μM) was added to the superfusion fluid, relative to the untreated interleaved controls (LPP-GC 66 ± 8 %, n = 7; MPP-GC 71 ± 7%, n = 6: Fig. S1). Subsequently, we loaded the postsynaptic neuron at LPP-GC synapses with FK506 and as t-LTD was no longer detected at these synapses (105 ± 11 %, n = 7: Fig. S1), postsynaptic calcineurin appears to be involved in t-LTD induction at LPP-GC but not MPP-GC synapses.

Endocannabinoid signalling at lateral and medial perforant pathway-dentate gyrus granule cell synapses is required for t-LTD

Postsynaptic Ca^{2+} dynamics may affect the synthesis of eCBs like anandamide and 2-arachidonoylglycerol (2-AG), ligands of CB_1Rs (Laguerre et al., 2021) necessary for t-LTD at neocortical, striatal and hippocampal synapses in rodents (Bender et al., 2006; Andrade-Talavera et al., 2016; Cui et al., 2016; Hogrefe et al., 2022). These receptors may be expressed in neurons and/or astrocytes, where they can mediate the release of gliotransmitters like glutamate, ATP/adenosine and D-serine through exocytotic mechanisms (Covelo and Araque, 2018). The induction of t-LTD at hippocampal and neocortex synapses requires the eCB system and astrocyte signalling, which may in turn drive gliotransmitter release (Min and Nevian, 2012; Andrade-Talavera et al., 2016; Martínez-Gallego et al., 2022a). To investigate the role of eCB signalling in these forms of t-LTD, we loaded the postsynaptic neurons with tetrahydrolipstatin (THL, 5 μM) to inhibit diacylglycerol lipase activity and prevent eCB synthesis. THL prevented t-LTD induction at both LPP-GC (92 ± 7 %, n = 8 versus interleaved controls 67 ± 3% n = 8: Fig. 3c, d) and MPP-GC synapses (92 ± 5 %, n = 6 versus interleaved controls 66 ± 5%, n = 8: Fig. 3g, h). Moreover, the presence of the CB_1R antagonist AM251 (3 μM) also blocked t-LTD at both LPP-GC (102 ± 7 %, n = 14: Fig. 3c, d) and MPP-GC synapses (98 ± 5 %, n = 6: Fig. 3g, h). Thus, eCB synthesis by postsynaptic cells and CB_1R activation of is required for presynaptic t-LTD at LPP- and MPP-DG GC synapses.

Astrocyte signalling and glutamate release at lateral and medial perforant pathway-dentate gyrus granule cell synapses is required for t-LTD

Astrocytes are involved in plasticity and synaptic transmission at hippocampal and cortical synapses (Min and Nevian, 2012; Araque et al., 2014; Andrade-Talavera et

al., 2016; Covelo and Araque, 2018; Pérez-Rodríguez et al., 2019; Falcón-Moya et al., 2020; Durkee et al., 2021; Martínez-Gallego et al., 2022a), and at the synapses studied here (Jourdain et al., 2007; Savtchouk et al., 2019; Pérez-Otaño and Rodríguez-Moreno, 2019). To investigate the possible involvement of astrocytes in the induction of t-LTD at the EC-DG synapses studied here, two different approaches were used. First, individual astrocytes were loaded with the Ca^{2+} chelator BAPTA (20 mM) though the patch pipette to inhibit vesicular and Ca^{2+} dependent gliotransmitter release (Parpura and Zorec, 2010), both at LPP-GC and MPP-GC synapses, and in neurons situated at 50-100 μm from the loaded astrocytes (Fig. 4a, f). In these experimental conditions, t-LTD was impaired at both LPP-GC (103 ± 9 %, n = 6: Fig. 4b, c) and MPP-GC synapses (94 ± 6 %, n = 6: Fig. 4g, h). Second, we assessed t-LTD in transgenic mice expressing a selective dnSNARE domain (dnSNARE) in astrocytes, preventing functional vesicular gliotransmitter release from these cells (Pascual et al., 2005; Sultan et al., 2015; Sardinha et al., 2017). In contrast to the typical t-LTD observed in WT mice (LPP-GC synapses 71 ± 5 %, n = 7; MPP-GC synapses 60 ± 5 %, n = 6: Fig. 4b, c, g and h), t-LTD could not be induced in these dnSNARE mice (LPP-GC synapses 102 ± 9 %, n = 9; MPP-GC synapses 112 ± 6 %, n = 7: Fig. 4c, d, g and h). These results clearly indicate that astrocytes are required to induce t-LTD at both LPP-GC and MPP-GC synapses.

At these synapses it is thought that glutamate is released from astrocytes to modulate synaptic transmission and participate in plasticity (Jourdain et al., 2007; Savtchouk et al., 2019), as indicated for cortical synapses (Min and Nevian, 2012; Martínez-Gallego et al., 2022a). To determine whether glutamate, probably released by astrocytes, is required for t-LTD at LPP- and MPP-GC synapses, we repeated the experiments in slices from dnSNARE mutant mice but applying glutamate puffs (100 μM) in the proximity of the astrocytes and neurons under study rather than pairing. In these experimental conditions, t-LTD, which was not evident in untreated dnSNARE mice (LPP-GC synapses 95 ± 5 %, n = 7; MPP-GC synapses 109 ± 6 %, n = 6) was fully recovered at both LPP-GC (72 ± 11 %, n = 6: Fig. 4d, e) and MPP-GC synapses (70 ± 8 %, n = 6: Fig. 4i, j) relative to the control interleaved slices from WT mice (LPP-GC synapses 62 ± 8 %, n = 7; MPP-GC synapses 64 ± 5 %, n = 7). These results indicate that glutamate, probably released by astrocytes as gliotransmitters, are necessary for the induction of t-LTD at LPP- and MPP-GC synapses.

Discussion

In this study, two new forms of t-LTD were seen to be elicited at LPP and MPP synapses onto DG GCs by pairing presynaptic activity with single postsynaptic APs at low frequency (0.2 Hz). Using different approaches, we found that these two forms of t-LTD are expressed presynaptically, not least through a decrease in the glutamate release probability. Interestingly, t-LTD induction at MPP-GC synapses but not at LPP-GC synapses depends on presynaptic NMDARs containing GluN2A subunits. In addition, mGluRs are required for t-LTD at both types of PP-DG synapses, although LPP-GC synapses require mGluR5 activation and MPP-GC synapses require mGluR1 activity. These two forms of presynaptic t-LTD require postsynaptic Ca^{2+} released from postsynaptic intracellular stores and whereas t-LTD at LPP-GC synapses, requires calcineurin phosphatase activity; this is not the case at MPP-GC synapses. Both these two forms of t-LTD require eCB synthesis and CB_1R activation, and both require astrocytes and astrocyte signalling to promote gliotransmitter release, as well as glutamate that is probably released by astrocytes.

While different time intervals between the postsynaptic spike and presynaptic EPSP of a pairing protocol may induce t-LTD (Feldman, 2012), it is clear here that pairing EPSPs of 3-5 mV size 100 times with single postsynaptic spikes 18 ms later at 0.2 Hz induces robust LTD at both LPP- and MPP-GC synapses. Interestingly, a similar protocol induces t-LTD at hippocampal CA3-CA1 synapses (Andrade-Talavera et al., 2016; Pérez-Rodríguez et al., 2019, Falcón-Moya et al., 2020) and at L4-L2/3 synapses of the somatosensory (Rodríguez-Moreno and Paulsen, 2008; Banerjee et al., 2009; 2014; Martínez-Gallego et al., 2022a), visual (Larsen et al., 2014) and cingulate cortices (Hogrefe et al., 2022). As such, the results presented here extend the existence of t-LTD to LPP and MPP synapses onto hippocampal DG GCs. Although there are several uncertainties associated with quantal analysis in the CNS (see Brock et al., 2020), we used different approaches here to determine the locus of expression of these two forms of t-LTD (failure rate, PPRs, CV, and mEPSP frequency and amplitude analysis), all of which were consistent with presynaptic changes.

NMDARs are required for general synaptic plasticity and for STDP, and in
different experimental models (cell cultures, rodent brain slices and human neurons)
they are required for t-LTD and t-LTP at different synapses in the hippocampus,
neocortex, cerebellum and striatum (Bi and Poo, 1998; Bender et al., 2006; Nevian and
Sakmann, 2006; Banerjee et al., 2009, 2014; Fino et al., 2010; Verhoog and
Mansvelder, 2011; Kwag and Paulsen, 2012; Andrade-Talavera et al., 2016; Sgritta et
al., 2017; Louth et al., 2021; Martínez-Gallego et al., 2022a). NMDARs have been
implicated in synaptic plasticity at MPP-GC but not LPP-GC synapses through classical
protocols (Dahl et al., 1990; Min et al., 1998; Beck et al., 2000; Hayashi et al., 2014).
Our results are consistent with the presence of NMDARs at MPP-GC synapses, in our
case participating in LTD. Indeed, t-LTD is not present at MPP-GC synapses in the
presence of D-AP5, and the presence of presynaptic NMDARs at MPP-GC synapses
was demonstrated previously by immunogold labelling (Jourdain et al., 2007;
Savtchouk et al., 2019). At these synapses t-LTD was also blocked when MK801 was
present in the bath but not when it was loaded into the postsynaptic neuron, indicating
that the NMDARs necessary for t-LTD are not postsynaptic, as seen in L4-L2/3
synapses of the somatosensory (Bender et al., 2006; Nevian and Sakmann, 2006,
Rodríguez-Moreno and Paulsen, 2008; Rodríguez-Moreno et al., 2011, 2013) and visual
cortex (Larsen et al., 2014). Since the NMDA channel blocker MK801 prevents t-LTD,
the NMDARs required for t-LTD appear to be ionotropic, although an additional role
for a postsynaptic metabotropic NMDARs cannot be ruled-out (Nabavi et al., 2013).
Thus, the involvement of NMDARs is one important difference between LPP- and
MPP-GC synapses that may account for at least some of the physiological differences
described between these two pathways. By using antagonists with different NMDAR
subunit specificities, we also found that the NMDARs required for t-LTD at MPP-GC
synapses contain GluN2A but not GluN2B or GluN2C/D subunits. NMDARs that
enhance glutamate release or that participate in LTP at PP-GC synapses are thought to
contain GluN2B subunits in rats as their effect is blocked by ifenprodil (Jourdain et al.,
2007; Savtchouk et al., 2019). However, ifenprodil or Ro25698 had no effect on t-LTD
here, such that the preNMDARs containing GluN2B subunits that participate in LTP
and enhance glutamate release may differ from those participating in t-LTD, and there
may even be differences between the species (i.e.: rat and mouse). Thus, while both
forms of t-LTD described here appear to require group I mGluRs, they require different
ones: mGluR5 at LPP-GC synapses and mGluR1 at MPP-GC contacts. Nevertheless,

the exact location of these mGluRs is not clear at present, which may represent a further

difference between LPP- and MPP-GC synapses, and with important behavioural and

physiological consequences.

Both, t-LTD and t-LTP require postsynaptic Ca^{2+} at neocortical (Bender et al.,

2006; Nevian and Sakmann, 2006; Rodríguez-Moreno et al., 2013; Martínez-Gallego et

al., 2022a) and hippocampal synapses (Andrade-Talavera et al., 2016; Falcón-Moya et

al., 2020; Mateos-Aparicio and Rodríguez-Moreno, 2020; Ingebert et al., 2021).

Moreover, like other forms of t-LTD, that at LPP- and MPP-GC synapses requires a rise

in Ca^{2+} in the postsynaptic cell due to Ca^{2+} release from internal stores, as also seen in

other situations of hippocampal, neocortical and cerebellar t-LTD (Bender et al., 2006;

Nevian and Sakmann, 2006; Andrade-Talavera et al., 2016; Sgritta et al., 2017; Falcón-

Moya et al., 2020; Martínez-Gallego et al., 2022a). This is similar to the LTD induced

by LFS in DG GCs (Wang et al., 1997). Our data also implies an involvement of

calcineurin in t-LTD induction at LPP-GC synapses but not at MPP-GC synapses, an

additional striking difference between these two types of synapses. Protein

phosphatases, including calcineurin, may be required for different forms of LTD, both

in the hippocampus (Mulkey et al., 1994) and neocortex (Torii et al., 1995). Future

experiments will determine the exact mechanism by which calcineurin mediates t-LTD

at LPP but not at MPP synapses. What is clear for the moment is that while the

requirements for Ca^{2+} seem to be similar at both LPP-GC and MPP-GC synapses, the

Ca^{2+}-dependent protein phosphatase calcineurin is only involved in t-LTD at LPP-GC

synapses. Significantly, the functional consequences of this difference remain to be

determined. Our results indicate that eCBs and CB_1Rs are involved in the induction of

these two different forms of t-LTD, in agreement with previous field recordings

showing that the eCB system is involved in synaptic plasticity at PP-DG synapses

(Wang et al., 2016; Peñasco et al., 2019; Fontaine et al., 2020). Here we did not

elucidate the location of the CB_1Rs mediating t-LTD, such that further work will be

required to address this question. However, the demonstration that astrocytes are

required for t-LTD is suggestive that astrocytic CB_1Rs might be involved.

Through two different approaches (introducing BAPTA into astrocytes and

using mutant dnSNARE mice) we demonstrated the involvement of astrocytes in

presynaptic t-LTD, both at LPP- and MPP-GC synapses. Astrocytes participate in

synaptic potentiation at PP-GC synapses (Jourdain et al., 2007), although little is known

about the contribution of astrocytes to LTD. Thus, this is the first demonstration of the participation of astrocytes in LTD at these synapses. Interestingly, and as found in the somatosensory cortex (Min and Nevian, 2012), glutamate is required to mediate t-LTD induction at both types of PP-GC synapses, and in fact, t-LTD can be recovered by the local application of glutamate at both LPP- and MPP-GC synapses in slices from dnSNARE mice that do not develop t-LTD otherwise. These findings agree with previous studies showing that astrocytes can release glutamate that reaches the NMDARs located in the presynaptic boutons at PP-GC synapses (Bezzi et al., 2004; Jourdain et al., 2007), influencing synaptic transmission at MPP-GC synapses (Savtchouk et al., 2019). It is interesting to note that the presynaptic targets of the glutamate at least in part released by astrocytes are not the same at LPP- and MPP-GC synapses, as they are mGluRs in the case of LPP-GC synapses, and both NMDARs and mGluRs in the case of MPP-GC synapses. While it is possible that glutamate from the presynaptic neuron also activates preNMDARs and mGluRs, the amount of glutamate released by presynaptic terminals may be insufficient to drive t-LTD, and hence, glutamate released from astrocytes may be required for t-LTD.

What is the physiological role of these forms of plasticity? For the moment, the exact role of STDP in the perforant cortex or in the hippocampus is not known, and more work is necessary to specifically determine functions for t-LTP and t-LTD. STDP seems a good candidate to mediate spatial learning in the hippocampus (Bush et al., 2010) and the possible role of t-LTP and t-LTD in forms of learning involving the hippocampus and the perforant cortex will be addressed in future studies. Our studies were done in developing P13-P21 animals indicating its relevance during maturation. The functions of t-LTP and t-LTD during development are most probably related to the refinement of synaptic connections and remodelling of neuronal circuits (Andrade-Talavera et al., 2023). As a Hebbian learning rule, t-LTP should occur when the spike order is pre-before-post, strengthening those connections in which the presynaptic neuron takes part in firing the postsynaptic cell, as predicted by Hebb, whereas t-LTD occurs when the spike order is reversed, so that non-causal spiking weakens the connections involved, possibly as a first step in the elimination of those connections during development as has been suggested (see Caporale and Dan, 2008). Indeed, further studies will be necessary to determine the true influence of STDP in the perforant cortex and in the hippocampus, and the specific developmental role of t-LTD

and t-LTP in these circuits. As discussed previously (Pérez-Rodríguez et al., 2019;

Falcón-Moya et al., 2020), t-LTD most probably participates in refining synapses,

potentially weakening excitatory synapses that are underused or behaviorally irrelevant

(Buonomano and Merzenich, 1998; Feldman and Brecht, 2005).

It is interesting to note that with these new results it is clear that both forms of t-

LTD may require glutamate from astrocytes. Presynaptic plasticity may involve

structural changes and may change the short-term properties of neurotransmitter release,

participating in circuit computations, and changing the excitatory/inhibitory balance or

sensory adaptations (Monday et al., 2018). Why some synapses, like L4-L2/3 synapses

at somatosensory cortex (and as observed in the hippocampus at CA3-CA1 synapses),

show pre- and/or post-synaptic plasticity requires further study. Interestingly, in the

synapses studied here, t-LTD of LPP-GC and MPP-GC synapses have different

requirements, indicating that the presynaptic expression of plasticity is fundamental for

the correct functioning of brain circuits and it is possible that different presynaptic

forms of t-LTD from synapses onto the same neuron as pre- and postsynaptic forms of

LTD (Banerjee et al., 2009, 2014) are regulated differently. Presynaptically expressed

forms of t-LTD may in fact control the trial-to-trial reliability, and they may induce a

larger change in signal-to-noise ratio than postsynaptic changes alone, as described in

auditory cortex (Froemke et al., 2013). Broad-ranging input representations from the EC

transmitted via the PP are sparse, and the DG processes this information and converts it

into a less well-correlated output to become less similar and more specific (Borzello et

al., 2023). Whether t-LTD at LPP and MPP synapses described here are involved in this

important process require further research and the different cellular and molecular

mechanisms involved in t-LTD at LPP- and MPP-GC synapses may account for some

of the difference in their physiological influence. Notwithstanding this, the potential

behavioural influence of the presynaptic forms of LTD studied here is still an emerging

issue of particular interest in the near future.

Acknowledgements

We thank Dr Joao Oliveira for providing the dnSNARE mice. This work

received support from the Spanish Agencia Estatal de Investigación and Fondo Europeo

de Desarrollo Regional (FEDER, Grants PID2019-107677GB-I00 and PID2022-

136597NB-I00), and from the Junta de Andalucía and Fondo Europeo de Desarrollo
Regional (FEDER, Grant P20-0881) to A. R-M. H. C-C was supported by a Visiting
Scholarship from the ESN, and I. M-G was supported by a FPU Fellowship from the
Spanish Agencia Estatal de Investigación.

Author contributions

592   H. C-C and I. M-G. performed the experiments and analysed the results. A. R-M. designed and
593   supervised the research, and wrote the paper.

References

Andrade-Talavera, Y., Duque-Feria, P., Paulsen, O. & Rodríguez-Moreno, A.
Presynaptic Spike Timing-Dependent Long-Term Depression in the Mouse Hippocampus.
Cereb. Cortex **26**, 3637-3654 (2016).

Araque, A. et al. Gliotransmitters travel in time and space. *Neuron* **81**, 728-739 (2014).

Banerjee, A. et al. Double dissociation of spike timing-dependent potentiation and
depression by subunit-preferring NMDA receptor antagonists in mouse barrel cortex. *Cereb.*
Cortex **19**, 2959-2969 (2009).

Banerjee, A., González-Rueda, A., Sampaio-Baptista, C., Paulsen, O. & Rodríguez-
Moreno, A. Distinct mechanisms of spike timing-dependent LTD at vertical and horizontal
inputs onto L2/3 pyramidal neurons in mouse barrel cortex. *Physiol. Rep.* **2**, e00271 (2014).

Beck, H., Goussakov, I. V., Lie, A., Helmstaedter, C. & Elger, C. E. Synaptic plasticity
in the human dentate gyrus. *J. Neurosci.* **20**, 7080-7086 (2000).

Bender, V. A., Bender, K. J., Brasier, D. J. & Feldman, D. E. Two coincidence
detectors for spike timing-dependent plasticity in somatosensory cortex. *J. Neurosci.* **26**, 4166-
4177 (2006).

Bezzi, P. et al. Astrocytes contain a vesicular compartment that is competent for
regulated exocytosis of glutamate. *Nat. Neurosci.* **7**, 613-620 (2004).

Bi, G. Q. & Poo, M. M. Synaptic modifications in cultured hippocampal neurons:
dependence on spike timing, synaptic strength, and postsynaptic cell type. *J. Neurosci.* **18**,
10464-10472 (1998).

Bidoret, C., Ayon, A., Barbour, B. & Casado, M. Presynaptic NR2A-containing NMDA
receptors implement a high-pass filter synaptic plasticity rule. *Proc. Natl. Acad. Sci. USA* **106**,
14126-14131 (2009).

Bliss, T. V. & Lømo, T. Long-lasting potentiation of synaptic transmission in the
dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J. Physiol.*
232, 331-356 (1973).

Borzello, M. et al. Assessments of dentate gyrus function: discoveries and debates.
Nature reviews. *Neuroscience* **24**, 502-517 (2023).

Brock, J. A., Thomazeau, A., Watanabe, A., Li, S. S. Y. & Sjöström, P. J. A Practical
Guide to Using CV Analysis for Determining the Locus of Synaptic Plasticity. *Front. Syn.*
Neurosci. **12**, 11 (2020).

Burgess, N., Maguire, E. A. & O'Keefe, J. The human hippocampus and spatial and
episodic memory. *Neuron* **35**, 625-641 (2002)

Chen, Q. Y. et al. NMDA GluN2C/2D receptors contribute to synaptic regulation and
plasticity in the anterior cingulate cortex of adult mice. *Mol. Brain* **14**, 60 (2021).

Clarke, J. R., Cammarota, M., Gruart, A., Izquierdo, I. & Delgado-García, J. M. Plastic
modifications induced by object recognition memory processing. *Proc. Natl. Acad. Sci. USA*
107, 2652-2657 (2010).

Contreras, M. P., Born, J. & Inostroza, M. The expression of allocentric object-place
recognition memory during development. *Behav. Brain Res.* **372**, 112013 (2019).

Covelo, A. & Araque, A. Neuronal activity determines distinct gliotransmitter release
from a single astrocyte. *eLife* **7**, e32237 (2018).

Cui, Y. et al. Endocannabinoid dynamics gate spike-timing dependent depression and
potentiation. *eLife* **5**, e13185 (2016).

Dahl, D., Burgard, E. C. & Sarvey, J. M. NMDA receptor antagonists reduce medial,
but not lateral, perforant path-evoked EPSPs in dentate gyrus of rat hippocampal slice. *Exp.*
Brain Res. **83** (1990).

Dan, Y. & Poo, M. M. Spike timing-dependent plasticity of neural circuits. *Neuron* **44**,
23-30 (2004).

Dietrich, D. et al. Metabotropic glutamate receptors modulate synaptic transmission in
the perforant path: pharmacology and localization of two distinct receptors. *Brain Res.* **767**,
220-227 (1997).

Durkee, C., Kofuji, P., Navarrete, M. & Araque, A. Astrocyte and neuron cooperation
in long-term depression. *Trends Neurosci.* **44**, 837-848 (2021).

Falcón-Moya, R. et al. Astrocyte-mediated switch in spike timing-dependent plasticity
during hippocampal development. *Nat. Commun.* **11**, 4388, doi:10.1038/s41467-020-18024-4
(2020).

Feldman, D. E. The spike-timing dependence of plasticity. *Neuron* **75**, 556-571 (2012).

Fino, E. et al. Distinct coincidence detectors govern the corticostriatal spike timing-
dependent plasticity. *J. Physiol.* **588**, 3045-3062 (2010).

Fontaine, C. J. et al. Endocannabinoid receptors contribute significantly to multiple
forms of long-term depression in the rat dentate gyrus. *Learn Mem.* **27**, 380-389 (2020).

Froc, D. J. et al. Reduced synaptic plasticity in the lateral perforant path input to the
dentate gyrus of aged C57BL/6 mice. *J. Neurophysiol.* **90**, 32-38 (2003).

Gladding, C. M., Fitzjohn, S. M. & Molnar, E. Metabotropic glutamate receptor-
mediated long-term depression: molecular mechanisms. *Pharmacol. Rev.* **61**, 395-412 (2009).

Harney, S. C., Rowan, M. & Anwyl, R. Long-term depression of NMDA receptor-
mediated synaptic transmission is dependent on activation of metabotropic glutamate receptors
and is altered to long-term potentiation by low intracellular calcium buffering. *J. Neurosci.* **26**,
1128-1132 (2006).

Hayashi, Y. et al. Enhanced stability of hippocampal place representation caused by
reduced magnesium block of NMDA receptors in the dentate gyrus. *Mol. Brain* **7**, 44 (2014).

Hogrefe, N. et al. Long-Lasting, Pathway-Specific Impairment of a Novel Form of
Spike-Timing-Dependent Long-Term Depression by Neuropathic Pain in the Anterior Cingulate
Cortex. *J. Neurosci.* **42**, 2166-2179 (2022).

Hu, N. W., Klyubin, I., Anwyl, R. & Rowan, M. J. GluN2B subunit-containing NMDA
receptor antagonists prevent Abeta-mediated synaptic plasticity disruption in vivo. *Proc. Natl.*
Acad. Sci. USA **106**, 20504-20509 (2009).

Inglebert, Y. & Debanne, D. Calcium and Spike Timing-Dependent Plasticity. *Front.*
Cell Neurosci. **15**, 727336 (2021).

Jourdain, P. et al. Glutamate exocytosis from astrocytes controls synaptic strength. *Nat.*
Neurosci. **10**, 331-339 (2007).

Kemp, A. & Manahan-Vaughan, D. Hippocampal long-term depression and long-term
potentiation encode different aspects of novelty acquisition. *Proc. Natl. Acad. Sci. USA* **101**,
8192-8197 (2004).

Khan, U. A. et al. Molecular drivers and cortical spread of lateral entorhinal cortex
dysfunction in preclinical Alzheimer's disease. *Nat. Neurosci.* **17**, 304-311 (2014).

Kim, S., Kim, Y., Lee, S. H. & Ho, W. K. Dendritic spikes in hippocampal granule cells
are necessary for long-term potentiation at the perforant path synapse. *eLife* **7**, e35269 (2018).

Krueppel, R., Remy, S. & Beck, H. Dendritic integration in hippocampal dentate
granule cells. *Neuron* **71**, 512-528 (2011).

Kwag, J. & Paulsen, O. Gating of NMDA receptor-mediated hippocampal spike timing-
dependent potentiation by mGluR5. *Neuropharmacology* **63**, 701-709 (2012).

Laguerre, A., Keutler, K., Hauke, S. & Schultz, C. Regulation of Calcium Oscillations
in beta-Cells by Co-activated Cannabinoid Receptors. *Cell Chem. Biol.* **28**, 88-96 e83 (2021).

Larsen, R. S. et al. Synapse-specific control of experience-dependent plasticity by
presynaptic NMDA receptors. *Neuron* **83**, 879-893 (2014).

Llorens-Martín, M., et al., Selective alterations of neurons and circuits related to early
memory loss in Alzheimer's disease. *Front. Neuroanat.* **8**, 38 (2014).

Louth, E. L., Jørgensen, R. L., Korshoej, A. R., Sørensen, J. C. H. & Capogna, M.
Dopaminergic Neuromodulation of Spike Timing Dependent Plasticity in Mature Adult Rodent
and Human Cortical Neurons. *Front. Cell. Neurosci.* **15**, 668980 (2021).

Malenka, R. C. & Bear, M. F. LTP and LTD: an embarrassment of riches. *Neuron* **44**,
5-21 (2004).

Mares, P., Kozlova, L., Mikulecka, A. & Kubova, H. The GluN2B-Selective Antagonist
Ro 25-6981 Is Effective against PTZ-Induced Seizures and Safe for Further Development in
Infantile Rats. *Pharmaceutics* **13**, 1482 (2021).

Martínez-Gallego, I., Pérez-Rodríguez, M., Coatl-Cuaya, H., Flores, G. & Rodríguez-
Moreno, A. Adenosine and astrocytes determine the developmental dynamics of spike timing-
dependent plasticity in the somatosensory cortex. *J. Neurosci.* **42**, 6038-6052 (2022a).

Martínez-Gallego, I., Rodríguez-Moreno, A. & Andrade-Talavera, Y. Role of Group I
Metabotropic Glutamate Receptors in Spike Timing-Dependent Plasticity. *Int. J. Mol. Sci.* **23**,
7807 (2022b).

Mateos-Aparicio, P. & Rodríguez-Moreno, A. Calcium Dynamics and Synaptic
Plasticity. *Adv. Exp. Med. Biol.* **1131**, 965-984 (2020).

Min, M. Y., Asztely, F., Kokaia, M. & Kullmann, D. M. Long-term potentiation and
dual-component quantal signaling in the dentate gyrus. *Proc. Natl. Acad. Sci. USA* **95**, 4702-
4707 (1998).

Min, R. & Nevian, T. Astrocyte signaling controls spike timing-dependent depression at
neocortical synapses. *Nat. Neurosci.* **15**, 746-753 (2012).

Mulkey, R. M., Endo, S., Shenolikar, S. & Malenka, R. C. Involvement of a
calcineurin/inhibitor-1 phosphatase cascade in hippocampal long-term depression. *Nature* **369**,
486-488 (1994).

Nabavi, S. et al. Metabotropic NMDA receptor function is required for NMDA
receptor-dependent long-term depression. *Proc. Natl. Acad. Sci. USA* **110**, 4027-4032 (2013).

Nevian, T. & Sakmann, B. Spine Ca^{2+} signaling in spike-timing-dependent plasticity. *J.*
Neurosci. **26**, 11001-11013 (2006).

Paré, D., DeCurtis, M. & Llinás R. Role of the hippocampal-entorhinal loop in temporal
lobe epilepsy: Extra- and intracellular study in the isolated Guinea pig brain in vitro. *J.*
Neurosci. **12**, 1867–1881 (1992)

Parpura, V. & Zorec, R. Gliotransmission: Exocytotic release from astrocytes. *Brain.*
Res. Rev. **63**, 83-92 (2010).

Pascual, O. et al. Astrocytic purinergic signaling coordinates synaptic networks. *Science*
310, 113-116 (2005).

Peñasco, S. et al. Endocannabinoid long-term depression revealed at medial perforant
path excitatory synapses in the dentate gyrus. *Neuropharmacology* **153**, 32-40 (2019).

Pérez-Otano, I. & Rodríguez-Moreno, A. Presynaptic NMDARs and astrocytes ally to
control circuit-specific information flow. *Proc. Natl. Acad. Sci. USA* **116**, 13166-13168 (2019).

Pérez-Rodríguez, M. et al. Adenosine Receptor-Mediated Developmental Loss of Spike
Timing-Dependent Depression in the Hippocampus. *Cereb. Cortex.* **29**, 3266-3281 (2019).

Petersen, R. P. et al. Electrophysiological identification of medial and lateral perforant
path inputs to the dentate gyrus. *Neuroscience* **252**, 154-168 (2013).

Prius-Mengual, J., Pérez-Rodríguez, M., Andrade-Talavera, Y. & Rodríguez-Moreno,
A. NMDA Receptors Containing GluN2B/2C/2D Subunits Mediate an Increase in Glutamate
Release at Hippocampal CA3-CA1 Synapses. *Mol. Neurobiol.* **56**, 1694-1706 (2019).

Reilly, J. F. et al. Amyloid deposition in the hippocampus and entorhinal cortex:
quantitative analysis of a transgenic mouse model. *Proc. Natl. Acad. Sci. USA* **100**, 4837-4842
(2003).

Rodríguez-Moreno, A. & Paulsen, O. Spike timing-dependent long-term depression
requires presynaptic NMDA receptors. *Nat. Neurosci.* **11**, 744-745 (2008).

Rodríguez-Moreno, A. et al. Presynaptic induction and expression of timing-dependent
long-term depression demonstrated by compartment-specific photorelease of a use-dependent
NMDA receptor antagonist. *J. Neuurosci.* **31**, 8564-8569 (2011).

Rodríguez-Moreno, A. et al. Presynaptic self-depression at developing neocortical
synapses. *Neuron* **77**, 35-42 (2013).

Sardinha, V. M. et al. Astrocytic signaling supports hippocampal-prefrontal theta
synchronization and cognitive function. *Glia* **65**, 1944-1960 (2017).

Savtchouk, I. et al. Circuit-specific control of the medial entorhinal inputs to the dentate
gyrus by atypical presynaptic NMDARs activated by astrocytes. *Proc. Natl. Acad. Sci. USA*
116, 13602-13610 (2019).

Scimemi, A., Schorge, S., Kullmann, D. M. & Walker, M. C. Epileptogenesis is
associated with enhanced glutamatergic transmission in the perforant path. *J. Neurophysiol.* **95**,
1213-1220 (2006).

Sgritta, M., Locatelli, F., Soda, T., Prestori, F. & D'Angelo, E. U. Hebbian Spike-
Timing Dependent Plasticity at the Cerebellar Input Stage. *J. Neurosci.* **37**, 2809-2823 (2017).

Singh, N., Bartol, T., Levine, H., Sejnowski, T. & Nadkarni, S. Presynaptic
endoplasmic reticulum regulates short-term plasticity in hippocampal synapses. *Commun. Biol.*
4, 241 (2021).

Sjöström, P. J., Turrigiano, G. G. & Nelson, S. B. Neocortical LTD via Coincident
Activation of Presynaptic NMDA and Cannabinoid Receptors. *Neuron* **39**, 641-654 (2003).

Sultan, S. et al. Synaptic Integration of Adult-Born Hippocampal Neurons Is Locally
Controlled by Astrocytes. *Neuron* **88**, 957-972 (2015).

Torii, N., Kamishita, T., Otsu, Y. & Tsumoto, T. An inhibitor for calcineurin, FK506,
blocks induction of long-term depression in rat visual cortex. *Neurosci. Lett.* **185**, 1-4 (1995).

Verhoog, M. B. & Mansvelder, H. D. Presynaptic ionotropic receptors controlling and
modulating the rules for spike timing-dependent plasticity. *Neural plasticity* **2011**, 870763
(2011).

Wang, J. X. et al. Structural basis of subtype-selective competitive antagonism for
GluN2C/2D-containing NMDA receptors. *Nat. Commun.* **11**, 423 (2020).

Wang, J.-H. & Kelly, P. T. Postsynaptic Calcineurin Activity Downregulates Synaptic
Transmission by Weakening Intracellular Ca^{2+} Signaling Mechanisms in Hippocampal CA1
Neurons. *J. Neurosci.* **17**, 4600-4611 (1997).

Wang, W. et al. A Primary Cortical Input to Hippocampus Expresses a Pathway-
Specific and Endocannabinoid-Dependent Form of Long-Term Potentiation. *eNeuro* **3**,
ENEURO.0160-16.2016 (2016).

Wiera, G. et al. Mechanisms of NMDA Receptor- and Voltage-Gated L-Type Calcium
Channel-Dependent Hippocampal LTP Critically Rely on Proteolysis That Is Mediated by
Distinct Metalloproteinases. *J. Neurosci.* **37**, 1240-1256 (2017).

Figure legends

Figure 1. Input-specific presynaptic spike timing-dependent LTD at lateral and
medial perforant pathway-dentate gyrus granule cell synapses. (**a, k**) Schemes show
the general set-up for the electrophysiological recordings and pairing protocol (Δt = -18
ms, time between the peak of the postsynaptic spike and the onset of the EPSP) at both
LPP- and MPP-GC synapses: CA, *cornus ammonis*; DG, dentate gyrus; EC, entorhinal
cortex; S1 and S2, stimulating electrodes; R, recording electrode. A post-before-pre
pairing protocol induced t-LTD at both LPP-GC (**b**) and MPP-GC (**l**) synapses. The
EPSP slopes monitored in the paired (LPP-GC synapses black triangles, n = 14; MPP-
GC synapses open black triangles, n = 14) and unpaired (LPP-GC synapses red
symbols; MPP-GC synapses open red symbols) pathways are shown. The traces show
the EPSP before (1) and 30 minutes after (2) pairing. Depression was only observed in
the paired pathways. (**c, m**) Summary of the results. *At the lateral (**d-j**) and medial (**n-t**)*
perforant pathway synapses onto dentate gyrus granule cells t-LTD is expressed
presynaptically. The number of failures increases after t-LTD induction at both LPP- (n
= 8, **d**) and MPP-GC (n = 6, **n**) synapses. Traces show the EPSPs before (black) and 30
min after pairing (red). PPR increases after t-LTD pairing protocol at LPP-GC (**e**) and

MPP-GC **(o)** synapses. The traces show the EPSP before (black) and 30 min after t-

LTD induction (red). **(f, p)** Normalized plots of CV^{-2} versus mean EPSP slopes yield

data points mainly below the diagonal after t-LTD induction at both LPP- (n = 14) and

MPP-GC (n = 13) synapses. The traces show the EPSPs before (black) and 30 min after

t-LTD induction (red). Miniature EPSPs (mEPSPs) were monitored at baseline and after

t-LTD induction in the presence of TTX (300 nM) at LPP- (**g-j**, n = 6) and MPP-GC

synapses (**q-t**, n = 6). Cumulative graphs and histograms show that after t-LTD

induction, the frequency of mEPSPs decreases at both types of synapses, whereas the

amplitude of the mEPSPs remains constant. Scale bar for the % failures - 5 mV, 50 ms;

scale bar for PPR - 5 mV, 80 ms. Error bars indicate the S.E.M.: * $p < 0.05$, **$p < 0.01$,

*** $p < 0.001$, two-sided Student's t-test.

Figure 2. The t-LTD at lateral and medial perforant pathway-dentate gyrus

granule cell synapses has distinct requirements for glutamate receptors. *The t-LTD*

at LPP-GC synapses requires mGluR5. **(a)** The addition of D-AP5 (50 μM) or MK-801

(500 μM) to the superfusion fluid does not prevent the induction of t-LTD at LPP-GC

synapses. The EPSP slopes shown are from D-AP5-treated (red triangles, n = 8), MK-

801-treated (grey triangles, n = 6) and untreated cells (black triangles, n = 8). The traces

show EPSPs before (1) and 30 min after (2) pairing. **(b)** Summary of the results. **(c)** The

t-LTD at LPP-GC synapses requires mGluR5. The EPSP slopes are shown from control

slices (black triangles, n = 9), and slices treated with the mGluR antagonist LY341495

(100 μM, red squares, n = 8), the mGluR1 antagonist LY367385 (100 μM, grey

triangles, n = 6) or the mGluR5 antagonist MPEP (20 μM, blue squares, n = 8). The

traces show the EPSPs before (1) and 30 min after (2) pairing. **(d)** Summary of the

results. **(e)** *The t-LTD at MPP-GC synapses requires NMDARs containing GluN2A*

subunits and mGluR1. The addition of D-AP5 (50 μM) or MK-801 (500 μM) to the

superfusion fluid prevented t-LTD induction at MPP-GC synapses, whereas

postsynaptic loading of MK-801 (500 μM) does not block t-LTD induction. The EPSP

slopes are shown from D-AP5 (open red squares, n = 10) or MK-801 treated cells

(batch, open grey squares, n = 6; loaded postsynaptic cell, open blue triangles, n = 9),

and untreated cells (open black triangles, n = 8). The traces show the EPSPs before (1)

and 30 min after (2) pairing. **(f)** Summary of the results. **(g)** *The NMDARs involved in t-*

LTD at MPP-GC synapses contain GluN2A subunits. The t-LTD at MPP-GC synapses

was completely blocked in slices exposed to Zn^{2+} (300 nM), while it remained

unaffected in slices treated with Ro 25-6981 (0.5 μM), ifenprodil (3 μM) or PPDA. The

EPSP slopes shown are from control slices (open black triangles, n = 9) and slices

treated with Zn^{2+} (open red squares, n = 7), Ro-25-6981 (open grey triangles, n = 6),

ifenprodil (open blue triangles, n = 6) or PPDA (open pink triangles, n = 6). The traces

show the EPSPs before (1) and 30 min after (2) pairing. (**h**) Summary of the results. (**i**)

The t-LTD at MPP-GC synapses requires mGluR1. The EPSP slopes shown are from

control slices (open black triangles, n = 9), or slices treated with LY341495 (100 μM,

open red squares, n = 10), LY367385 (100 μM, open grey squares, n = 7) or MPEP (20

μM, open blue triangles, n = 6). The traces show the EPSPs before (1) and 30 min after

(2) pairing. (**j**) Summary of the results. The error bars indicate the S.E.M.: *$p < 0.05$, **

$p < 0.01$, ***$p < 0.001$, One-way ANOVA + Holm–Sidak test.

Figure 3. The t-LTD at lateral and medial perforant pathway-dentate gyrus

granule cell synapses requires postsynaptic calcium and eCB signalling. (**a**) The

induction of t-LTD at LPP-GC synapses in slices was unaffected by exposure to

nimodipine (10 μM) but it was prevented by treatment with thapsigargin (10 μM), either

in the superfusion fluid or when loaded into the postsynaptic cell. The EPSP slopes

shown are from control slices (black triangles, n = 9) and slices treated with nimodipine

(red triangles, n = 6), thapsigargin in the superfusion fluid (grey squares, n = 6) or

thapsigargin loaded into the postsynaptic cell (blue squares, n = 9). The traces show

EPSPs before (1) and 30 min after (2) pairing. (**b**) Summary of the results. (**c**) The t-

LTD at LPP-GC synapses requires endocannabinoids and CB1 receptors, as it was

blocked when THL was loaded into the postsynaptic neuron (pTHL, 5 μM) and in slices

treated with AM251 (3 μM). The EPSP slopes shown are from control slices (black

triangles, n = 14) and slices with THL loaded into the postsynaptic neuron (red squares,

n = 8) or treated with AM251 (grey squares, n = 14). The traces show the EPSPs before

(1) and 30 min after (2) pairing. (**d**) Summary of the results. (**e**) The induction of t-LTD

at MPP-GC synapses was unaffected in nimodipine-treated slices (10 μM) but

prevented in slices treated with thapsigargin (10 μM), either in the superfusion fluid or

when loaded into the postsynaptic cell. The EPSP slopes shown are from control slices

(open black triangles, n = 9) and slices treated with nimodipine (open red triangles, n =

6), thapsigargin in the superfusion fluid (open grey squares, n = 6) or thapsigargin

loaded into the postsynaptic cell (open blue squares, n = 7). The traces show EPSPs

before (1) and 30 min after (2) pairing. **(f)** Summary of the results. **(g)** The t-LTD at

MPP-GC synapses requires endocannabinoids and CB1 receptors, and it was blocked

when THL was loaded into the postsynaptic neuron (pTHL, 5 μM) and in slices treated

with AM251 (3 μM). The EPSP slopes shown are from control slices (open black

triangles, n = 8) and slices with THL loaded into the postsynaptic neuron (open red

squares, n = 6) or treated with AM251 (open grey squares, n = 6). The traces show

EPSPs before (1) and 30 min after (2) pairing. **(h)** Summary of the results. The error

bars indicate S.E.M. *p < 0.05, ** p < 0.01, *** p < 0.001, One-way ANOVA + Holm–

Sidak test.

Figure 4. The t-LTD at lateral and medial perforant pathway-dentate gyrus

granule cell synapses requires astrocytes and glutamate. (a, f) Schemes showing the

general experimental set-up: A, astrocyte; CA, *cornus ammonis*; DG, dentate gyrus; EC,

entorhinal cortex; GC, granule cell; S1 and S2, stimulating electrodes; R1 and R2,

recording electrodes. Inset: voltage responses of an astrocyte shown in current clamp

configuration. **(b)** The t-LTD at LPP-GC synapses requires astrocytes as it was

prevented by loading astrocytes with BAPTA (20 mM) and in slices from dnSNARE

mice. The EPSP slopes shown are from control slices (black triangles, n = 7), slices

with BAPTA-loaded astrocytes (red squares, n = 6) and slices from dnSNARE mice

(grey circles, n = 9). The traces show the EPSPs before (1) and 30 min after (2) pairing.

(c) Summary of the results. **(d)** The t-LTD at LPP-GC synapses requires glutamate as it

is absent in slices from dnSNARE mice but it is restored by glutamate puffs (100 μM).

The EPSP slopes shown are from control slices (black triangles, n = 7), slices from

dnSNARE mice (red circles, n = 7) and slices from dnSNARE mice administered

glutamate puffs (grey triangles, n = 6). The traces show the EPSPs before (1) and 30

min after (2) pairing. **(e)** Summary of the results. **(g)** The t-LTD at MPP-GC synapses

requires astrocytes as t-LTD induction at MPP-GC synapses was prevented by loading

astrocytes with BAPTA (20 mM) and in slices from dnSNARE mice. The EPSP slopes

shown are from control slices (open black triangles, n = 6), slices with BAPTA-loaded

astrocytes (open red squares, n = 6) and slices from dnSNARE mice (open grey circles,

n = 7). The traces show the EPSPs before (1) and 30 min after (2) pairing. **(h)** Summary

of the results. **(i)** The t-LTD at MPP-GC synapses requires glutamate as it is absent in

slices from dnSNARE mice but is restored by glutamate puffs (100 µM). The EPSP

slopes shown are from control slices (open black triangles, n = 7), slices from

dnSNARE mice (open red circles, n = 6) and slices from dnSNARE mice administered

to glutamate puffs (open grey triangles, n = 6). The traces show the EPSPs before (1)

and 30 min after (2) pairing. **(j)** Summary of the results. The error bars indicate the

S.E.M.: *$p < 0.05$, **$p < 0.01$, ***$P < 0.001$, One-way ANOVA + Holm–Sidak test.

Figure 1

Figure 2

Figure 3

Figure 4