PIM kinase control of CD8 T cell protein synthesis and cell trafficking

Julia M Marchingo[1,3,#], Laura Spinelli[1,2], Shalini Pathak[1,4], Doreen A Cantrell[1,#]

Affiliations:
1. Cell Signalling and Immunology Division, School of Life Sciences, University of Dundee,
United Kingdom
2. Molecular Cell and Developmental Biology Division, School of Life Sciences, University of
Dundee, United Kingdom
3. Current address: Blood Cells and Blood Cancer Division, Walter and Eliza Hall Institute of
Medical Research, Melbourne, Australia
4. Current address: Innovation Factory Core Technology Facility, University
of Manchester, UK
Correspondence: d.a.cantrell@dundee.ac.uk (D.A.C.), marchingo@wehi.edu.au (J.M.M.)

Abstract:

Integration of a large network of kinase signalling pathways co-ordinates changes in the

transcription, translation and metabolic events required for T cell activation and

differentiation. The present study explores the role of the Serine/Threonine kinases PIM1

and PIM2 in controlling murine CD8 T lymphocyte antigen receptor-mediated activation and

differentiation in response to the cytokines Interleukin 2 (IL-2) or IL-15. We show that PIM

kinases are dispensable for the differentiation programs controlled by the antigen-receptor

and IL-15. There is however a selective role for the PIM kinases in the context of IL-2

regulation of CD8 T cell fate. One key insight was that the PIM kinases controlled the

migratory capabilities of effector CD8 T cells, with *Pim1*/*Pim2*-deficient CD8 T cells unable to

fully switch off the naïve T cell chemokine and adhesion receptor program during effector

differentiation. PIM kinases were also needed for IL-2 to sustain high expression of the

glucose transporters SLC2A1 and SLC2A3 and to maintain activity of the nutrient sensing

kinase mTORc1. Strikingly, PIM kinases did not have a dominant impact on IL-2-driven

transcriptional programs but rather selectively modulated protein synthesis to shape

cytotoxic T cell proteomes. This study reveals a selective role of PIM kinases in IL-2 control

of CD8 T cells and highlights how regulated changes in protein synthesis can impact T cell

phenotypes.

Introduction:

CD8 T lymphocytes are key effector cells for adaptive immune responses to viruses, intracellular pathogens and tumours. Their clonal expansion and differentiation to cytotoxic effector or memory CD8 T cells is controlled by signalling pathways initiated by the T cell antigen receptor (TCR), co-stimulatory molecules and cytokines. Receptor engagement by immune stimuli triggers the activation of cytosolic tyrosine kinases that then direct the activity of a diverse network of Serine/Threonine kinases, disseminating signals from plasma membrane to cell interior. The Src family and ZAP70 Tyrosine kinases initiate this process downstream of the TCR, with the Janus kinases (JAK) as the main signal transducer for important cytokine regulators of T cell differentiation, IL-2 and IL-15 (Cantrell, 2015; Saravia et al., 2020). The subsequent network of Serine/Threonine kinases that is engaged beyond these initiating events coordinates phosphorylation of a diverse array of protein targets. These protein targets include chromatin regulators, transcription factors, translation machinery, cytoskeletal proteins and metabolic enzymes, which then control the changes in transcriptional, translational and metabolic programs that drive effector and memory T cell differentiation (Navarro et al., 2011; Tan et al., 2017; Locard-Paulet et al., 2020).

CD8 T cells express ∼200 Serine/Threonine protein kinases (Navarro et al., 2014; Howden et al., 2019). There is a good understanding of the roles of some Serine/Threonine kinases in CD8 T cells including the role of the diacylglycerol regulated kinases, MAP kinases ERK1 and 2, Phosphatidylinositol (3,4,5) triphosphate- controlled kinases PDK1 and AKT, and the nutrient sensing kinases mammalian Target of Rapamycin complex 1 (mTORc1) and AMP-activated protein kinase (AMPK) (Finlay et al., 2012; Rolf et al., 2013; Blagih et al., 2015; Cantrell, 2015; Hukelmann et al., 2016; Tan et al., 2017; Howden et al., 2019; Saravia et al., 2020; Damasio et al., 2021; Spinelli et al., 2021). However, beyond these core kinases there is much still to learn about how many of the ~200 Serine/Threonine kinases expressed in CD8 T cells contribute to the control of T cell function. In this context, the PIM kinases are a family of Serine/Threonine kinases that are strongly and rapidly induced in T cells responding to immune stimuli (Aho et al., 2005; Fox et al., 2005; Peperzak et al., 2010; Jackson et al., 2012). The PIM family contains 3 Serine/Threonine kinases, PIM1, PIM2 and PIM3, of which T lymphocytes predominantly express PIM1 and PIM2 (Mikkers et al., 2004; Fox et al., 2005). As PIM kinases are constitutively active, their participation in signalling

cascades is dictated by regulated changes in their protein expression levels (Warfel and

Kraft, 2015). PIM1 and PIM2 have little to no expression in naïve T cells but are induced by

T-cell receptor activation and costimulation (e.g. CD27) (Wingett et al., 1996; Fox et al.,

2005; Peperzak et al., 2010). PIM kinases are also direct transcriptional targets of JAK/STAT

signalling pathways and as such PIM1 and/or PIM2 are rapidly induced in T cells

downstream of a diverse array of cytokines, including multiple common cytokine receptor γ-

chain family cytokines, IL-6 and IL-12 (Matikainen et al., 1999; Aho et al., 2005; Fox et al.,

2005; Warfel and Kraft, 2015; Buchacher et al., 2023).

There have been some analyses of how deletion of PIM kinases impacts CD8 T cell function.

A mild reduction in *Pim1/Pim2/Pim3*-deficient CD4 and CD8 T cell proliferation has been

observed in response to sub-optimal TCR stimulation and IL-2 (Mikkers et al., 2004) and

there is evidence that PIM1 and PIM2 control survival pathways in naïve CD4 and CD8 T cells

in concert with mTORc1 (Fox et al., 2005). *Pim1/Pim2/Pim3* deletion is reported to promote

expression of stemness and memory-associated transcription factors including Bcl6 and Tcf7

(Chatterjee et al., 2019) and PIM kinase activity has been linked with maintenance of

memory CD8 T cell survival (Knudson et al., 2017). It has also been shown recently that

Pim1/Pim2 deficiency prevented mTORc1 activation and the proliferation of non-

conventional intra-epithelial gut CD8 T lymphocytes (IEL) in response to the cytokine IL-15

(James et al., 2021).

Elevated PIM kinase expression has been noted in patient T cells for a number of

autoimmune conditions, including rheumatoid arthritis, uveitis and coeliac disease, and PIM

kinase inhibition has been proposed as a possible intervention to dampen autoimmunity

(James et al., 2021; Maney et al., 2021; Li et al., 2022). PIM kinase inhibitors have also

entered clinical trials to treat some cancers and there is active interest in combining these

with immunotherapies (Clements and Warfel, 2022). Given the role of CD8 T cells in anti-

cancer immunity, it is important to understand how these kinase control CD8 T cell function.

However, there has not yet been a systematic analysis of the role of PIM kinases in

conventional CD8 T cells as they activate and differentiate to form effector and memory

cells. Accordingly, the current study, uses high resolution quantitative analysis of

Pim1/Pim2-deficient CD8 T cell proteomes and transcriptomes to obtain a comprehensive

and unbiased understanding of how PIM kinases control CD8 T cell TCR-driven activation,

and IL-2 or IL-15-driven proliferation, and differentiation. The study reveals a selective role

for PIM kinases as mediators of IL-2 signalling to control CD8 T cell trafficking and effector

functionality. We uncover that the dominant impact of PIM kinase loss was to shape

effector CD8 T cell proteomes rather than transcriptomes, reflecting PIM kinase modulation

of protein synthesis.

Results:

PIM1 and PIM2 are strongly induced by antigen receptor engagement but are dispensable

for T cell activation

T cell antigen receptor (TCR) -driven activation of naïve T cells initiates their clonal

expansion and differentiation into effector and memory cells. Quantitative mass

spectrometry-based proteomics analysis of naïve OT1 CD8 T cell stimulated with their

cognate peptide over time (Marchingo et al., 2020) showed that PIM1 protein was strongly

induced within 3 hours of TCR engagement, which was sustained at a 24 hour time point

(**Fig 1A**). PIM2 expression was not detected until 9 hours post-stimulation but was also

sustained at the 24 hour time point. 24 hour polyclonal activation of CD8 T cells with

αCD3/αCD28 agonist antibodies similarly induced expression of PIM1 and PIM2 proteins

(**Fig 1B, Table S1**). These increases in PIM kinases expression at a protein level were

underpinned by transcriptional reprograming, with both *Pim1* and P*im2* mRNA very lowly

expressed in naïve CD8 T cells and strongly upregulated by immune activation (**Fig 1C**)

(Spinelli et al., 2021). Consistent with previous reports of Pim1 and Pim2 being the major

PIM kinases in T cells (Mikkers et al., 2004; Fox et al., 2005), *Pim3* mRNA was only expressed

at very low levels in naïve and activated CD8 T cells (**Fig 1C**), with PIM3 protein not detected

at all by mass spectrometry (**Table S1**).

PIM1 and PIM2 have similar substrate selectivity and hence functional redundancy (Mikkers

et al., 2004; Bullock et al., 2005; Fox et al., 2005; Peng et al., 2007; Warfel and Kraft, 2015).

Accordingly, to explore the role of the PIM kinases in T cells, mice with a whole body

deletion of *Pim1* and *Pim2*, Pim1$^{-/-}$ (Pim1 KO) (Laird et al., 1993) and Pim2$^{-/-}$ or Pim2$^{-/Y}$ (Pim2

KO) (Mikkers et al., 2004), were backcrossed for >10 generations onto a C57BL/6

background and inter-crossed to generate *Pim1/Pim2* double knockout (Pim dKO) mice.

There were normal proportions of peripheral T cells in spleens of Pim dKO mice, though the

total number of splenocytes was lower than in age/sex matched wild-type (WT) mice (**Fig**

S1A-C). However, Pim dKO T cells showed a normal increase in cell size in response to

immune activation with αCD3/αCD28 agonist antibodies (**Fig 1D**) and 24 hour αCD3/αCD28

activated Pim dKO CD4 and CD8 T cells exhibited normal upregulation of established

markers of T cell activation including CD25, CD44 and the transferrin receptor CD71 (**Fig 1E**).

24 hour TCR-activated Pim dKO CD8 T cells also produced normal levels of the key effector

cytokine IFNγ (**Fig 1E**) and there was no discernible difference in TCR-driven proliferation

between WT and Pim dKO T cells over 3 days in culture (**Fig 1F, J**). To systematically explore

in an unbiased way if *Pim1*/*Pim2*-deficiency impacted TCR-driven activation phenotypes, we

performed high-resolution mass spectrometry to measure the proteomes of 24 hour

αCD3/αCD28 activated WT and Pim dKO CD4 and CD8 T cells. These experiments quantified

expression of >7000 proteins but found no substantial quantitative or qualitative differences

in protein content or proteome composition in activated WT versus Pim dKO CD4 and CD8 T

cells (**Fig 1G-H**) (**Table S1**), indicating a redundancy between PIM kinases and other

signalling pathways downstream of TCR activation.

One prominent study has reported that *Pim1*/*Pim2* deficiency sensitises T cells to the

mTORc1 inhibitor rapamycin (Fox et al., 2005), which suggests mTORc1 as a candidate for

mediating this redundancy. This study showed naïve Pim dKO T cells cultured in IL-7 died in

response to rapamycin treatment, whereas WT cells did not, and that rapamycin synergised

with *Pim1*/*Pim2*-deficiency to substantially reduce TCR-driven activation and proliferation

(Fox et al., 2005). However, we found no impact of *Pim1*/*Pim2*-deficiency on the ability of

naïve CD8 T cells cultured with IL-7 to survive in response to rapamycin treatment (**Fig 1I**).

Furthermore, we found no difference in the effect of rapamycin on mean division number in

Pim dKO versus WT CD8 T cells (**Fig 1J**). Collectively these data show that PIM 1 and PIM2

are rapidly and strongly expressed in response to immune activation, but they are

dispensable for the antigen receptor-driven proteome remodeling that initiates T cell clonal

expansion and differentiation. We also found no evidence that the *Pim1* and *Pim2* loss

synergised with mTORC1 inhibition.

PIM kinases do not control IL-15-driven CD8 T cell memory differentiation

The proliferative expansion and differentiation programs of CD8 T cells are shaped by cytokines. In this context, PIM1 and PIM2 kinase expression is regulated by the cytokines IL-2 and IL-15 which are key to shaping CD8 T cell effector and memory cell differentiation respectively (Weninger et al., 2001; Cornish et al., 2006; Buck et al., 2016). Both IL-2 and IL-15 control PIM1 and PIM2 expression in antigen primed CD8 T cells (**Fig 2A**). Accordingly, we assessed the contribution of PIM1 and PIM2 kinases to cytokine-driven expansion and differentiation of antigen activated CD8 T cells. In these experiments, antigen-primed CD8 T cells are expanded in IL-15 to generate memory phenotype CD8 T cells or expanded in IL-2 to generate effector cytotoxic T lymphocytes (CTL) (Weninger et al., 2001; Cornish et al., 2006; Buck et al., 2016) (**Fig 2B**).

Previous work has shown that PIM1 and PIM2 mediate IL-15 induced proliferation of gut intra-epithelial lymphocytes (IEL) (James et al., 2021). However, their role in IL-15 responses in conventional antigen receptor activated CD8 T cells is not known. We therefore first examined the impact of *Pim1/Pim2*-deficiency on IL-15-driven conventional CD8 T cell expansion. Antigen experienced WT CD8 T cells show a robust proliferative response to IL-15 such that there is an ~90-fold expansion of the CD8 T cells between days 2 and 6 of culture (**Fig 2C**). There was no impact of *Pim1/Pim2*-deficiency on the ability of IL-15 to sustain CD8 T cell viability (**Fig 2D**). Immune activated Pim dKO CD8 T cells also proliferated strongly in response to IL-15, but showed a moderate proliferative disadvantage resulting in a ~4-fold difference in total cell number by day 6 of the culture (**Fig 2C**). Antigen-primed CD8 T cells expand in IL-15 to generate memory phenotype CD8 T cells (Weninger et al., 2001). We therefore examined whether *Pim1/Pim2*-deficiency had any impact on the ability of IL-15 to induce memory differentiation. Accordingly, we performed parallel bulk RNAseq and high-resolution mass spectrometry to analyse the transcriptome and proteome of IL-15 expanded WT and Pim dKO CD8 T cells. The RNAseq analysis quantified ~14,000 RNA and using a cut off of >1.5 fold-change and q-value <0.05 we saw that the abundance of 381 RNA was modified by *Pim1/Pim2*-deficiency (**Fig 2E**) (**Table S2A**). Of the 381 RNA that were detected as differentially expressed (FC>1.5, q<0.05) only ~30% were detected with an average expression over 10 transcripts per million (TPM), of these gene ~36% were non-coding RNA, uncharacterised genes or pseudogenes, leaving only 72 substantially expressed

protein coding mRNA that were differentially expressed in Pim dKO versus WT IL-15

expanded CD8 T cells (**Table S2B**). These data indicate that Pim1/Pim2-deficiency has very

little effect on IL-15 induced protein-coding mRNA expression in differentiated T cells. We

then specifically examined if Pim dKO T cells had any differences in mRNA that are known to

be critical for memory cell differentiation. We found that *Pim1*/*Pim2*-deficiency did not

decrease expression of mRNA levels for critical memory T cell genes such as the adhesion

molecules and chemokine receptors, *Sell* (CD62L), *Ccr7* and *S1pr1* that direct the trafficking

of CD8 T cells to secondary lymphoid tissue. (**Fig 2F**). *Pim1/Pim2*-deficiency also did not

reduce expression of mRNA for the key memory cell transcription factors *Tcf7, Foxo1, Foxo3*

Klf2 and Id3. (**Fig 2G**). Indeed, if anything, there were small increases in expression of the

Klf2, *Id3* and *Ccr7* mRNA in Pim dKO versus WT IL-15 expanded CD8 T cells (**Fig 2F-G**).

Critically, when the proteomes of IL15 expanded WT and Pim dKO CD8 T cells were analysed

and expression of ~6900 total proteins quantified, there were no major changes in the

proteomes between WT and Pim dKO IL15 expanded CD8 T cells (**Fig 2H, Table S3**). Of note,

the mitochondrial proteome composition of IL15 expanded WT and Pim dKO CD8 T cells

were indistinguishable (**Fig 2I-J**). This is pertinent because one key feature of how IL-15

controls memory T cells is via mitochondrial remodelling to support energy production via

oxidative phosphorylation and fatty acid oxidation (Buck et al., 2016) and it has been

described that PIM kinases can control mitochondrial phenotypes in cancer cell lines and

cardiomyocytes (Din et al., 2013; Chauhan et al., 2020). These data show that PIM1 and

PIM2 do not drive IL-15-mediated metabolic or differentiation programs in antigen-primed

CD8 T cells.

PIM kinases selectively modulate IL-2 controlled effector CD8 T cell differentiation

IL-2 controls the expression of PIM1 and PIM2 kinases (**Fig 2A**) and in this context, we noted

that the copy numbers per cell of PIM1 protein is considerably higher in actively expanding

antigen-activated CD8 cells maintained with IL-2 compared to IL-15 (**Fig 3A**)**.** We therefore

examined the role of PIM1 and PIM2 in the IL-2 signalling pathways that control the

differentiation of effector CTL. The data show that Pim dKO antigen activated CD8 T cells

responded normally to IL-2 to induce STAT5 Y694 phosphorylation (**Fig 3B**). They also

express normal levels of the IL-2-receptor alpha chain, CD25, (**Fig 3C**), which is sustained by

STAT5, and together these data broadly indicate IL-2 receptor signal transduction is

unaltered by *Pim1/Pim2*-deficiency. Antigen activated WT CD8 T cells have a strong survival

and proliferative response to IL-2 producing an ~270-fold expansion between days 2 and 6

of culture in saturating IL-2 concentrations (**Fig 3D**). There was no impact of *Pim1/Pim2*-

deficiency on the ability of IL-2 to maintain CD8 T cell viability (**Fig 3E**) and Pim dKO cells CD8

T cells cultured in IL-2 proliferated robustly. They did however exhibit a very mild

proliferative disadvantage resulting in a ~2-fold difference in total cell number by day 6 of

the culture (**Fig 3D).** We then examined whether *Pim1/Pim2*-deficiency had any impact on

the differentiation of Pim dKO CD8 T cells into effector CTL. Expression of activation markers

CD44 and CD71 in IL-2-expanded CTL were similar between WT and Pim dKO CD8 T cells on

days 3 and 6 of co-cultures (**Fig 3F**). However, one consistent difference between IL-2

maintained WT and Pim dKO CD8 T cells is that WT CTL downregulated expression of CD62L

(L-selectin, *Sell*), whereas the Pim dKO CTL sustained higher CD62L expression (**Fig 3G**). We

also noted that IL-2 maintained Pim dKO CTL had reduced forward and side scatter

properties i.e were smaller and less granular than corresponding WT CTL (**Fig 3F).**

To explore the apparent size change between Pim dKO and WT IL-2 expanded CTL, we used

high resolution mass spectrometry to compare the proteomes of day 6 IL-2 cultured WT or

Pim dKO CTL. These analyses quantified expression of ~6,900 proteins and showed a small

decrease in the total protein mass of IL-2 maintained Pim dKO CD8 T cells compared to WT

cells (**Fig 4A, Table S4**). The data showed down-regulation of the expression of 513 proteins

(>1.5 fold-change, q<0.05) in Pim dKO CTL along with increased copy numbers of 33 proteins

(**Fig 4B**, **Table S4**). The transcription factor profile of Pim dKO T cells was consistent with

that of an effector CTL with high levels of TBX21 and low levels of TCF1 (*Tcf7*) (**Fig 4C**). Pim

dKO T cells also exhibited an effector-like mitochondrial proteome profile (**Fig S2**). However,

there were salient differences in proteins important for effector functionality between WT

and Pim dKO CD8 T cells. These included lower levels of the glucose transporters, SLC2A1

and SLC2A3 (GLUT1 and GLUT3) (**Fig 4D**) which are the major glucose transporters in CD8 T

cells (Hukelmann et al., 2016) and are key for enabling T cells to fuel high rates of glycolysis

and protein O-GlcNAcylation required to support effector functions(Chang et al., 2015;

Swamy et al., 2016). Pim dKO CTL also had decreased expression of a number of proteins

involved in fatty acid and cholesterol biosynthesis including SCD2, HMGCS1, MVK, IDI1,

FDPS, FDFT1 and SQLE (**Fig 4E**). Another striking observation was that the Pim dKO CD8 cells

had reduced levels of a number of key effector proteins including multiple granzymes,

perforin, IFNγ and TNFα (**Fig 4F**). Orthogonal flow cytometric analysis of Granzyme B and

IFNγ levels confirmed reduced levels of these effector molecule in IL-2 expanded Pim dKO

CTL, though IFNγ protein is only lowly produced in IL2 maintained CTL. (**Fig 4G).** In parallel

experiments we also examined the impact of an acute loss of PIM kinase activity on the

proteomes of IL-2 maintained CTL to see if this could recapitulate key observations from Pim

dKO T cells. When IL-2 maintained WT CTL were treated for 24 hours with the pan-PIM

kinases inhibitors PIM447 or AZD1208 we saw reductions in cell proliferation, cell size, cell

granularity, fatty acid biosynthesis proteins SCD1-3, glucose transporters SLC2A1, SLC2A3

and a reduction in Granzyme B expression (**Fig S3A-E, Table S5**). This corroborated some key

observations from the experiments with Pim dKO CTL, where *Pim1* and *Pim2* had been

deleted over the lifespan of the T cell.

To examine the molecular regulation underpinning PIM1 and PIM2 kinase control of key

metabolic and effector proteins we also performed RNAseq on IL-2 differentiated WT and

Pim dKO CTL samples collected in parallel with the proteomics analysis described in Fig 4. In

these experiments ~14,000 mRNA were quantified and using a cut off >1.5 fold-change,

q<0.05 we saw that the abundance of 223 mRNA were decreased and 155 increased by

Pim1/Pim2 deficiency (**Fig 5A, Table S6**). These data revealed that the reduced expression of

perforin and Granzymes C-G and K protein corresponded to decreases in their mRNA (**Fig**

5B-C). However, there were striking examples where a clear decrease in protein levels did

not correspond with an appreciable decrease in mRNA expression. These included the

predominant cytolytic Granzymes B and A (**Fig 5C-D**), as well as both glucose transporters

SLC2A1 and SLC2A3 (**Fig 5E**). Furthermore, contrasting differential expression of proteins

and mRNA data in Pim dKO vs WT CTL revealed that ~75 % of the proteins whose expression

was changed in Pim dKO T cells (FC>1.5, q<0.05), exhibited no strong corresponding changes

in their mRNA (FC<1.2) (**Fig 5F, Table S7**). Indeed, of the ~500 proteins whose expression

decreased at the protein level in Pim dKO CTL only 17 showed a strong decrease in

magnitude at the mRNA level as well (**Fig 5G, Table S7**).

These observations could reflect that the PIM kinases are required for IL-2 to maximally

control protein synthesis in CD8 T cells. To examine this hypothesis we first interrogated our

proteomics and transcriptomics data. Consistent with this hypothesis we observed a small

reduction in ribosome protein content in Pim dKO CTL, which scaled with the reduction in total protein content (**Fig 5H**). Moreover the translational repressor PDCD4 was increased in Pim dKO CTL (**Fig 5I**). This could be confirmed by flow cytometry (**Fig 5J**) and was underpinned at the mRNA level (**Fig 5B**). Similarly, 24 hour inhibition with pan PIM kinases inhibitors was sufficient to substantially increase PDCD4 protein expression (**Fig S3F**). We also noted that there was a small decrease in the expression of EIF4A1, a key component of the eIF4F translation initiation complex, in Pim dKO IL2 maintained CTL (**Fig 5K**). PDCD4 inhibits translation by binding EIF4A1 in a 2:1 ratio, preventing its interaction with the EIF4F complex (Suzuki et al., 2008). Together the changes in both proteins equated to an approximate halving of the ratio between EIF4A1 and PDCD4 (**Fig 5L**). These data all support that loss of PIM kinases could reduce protein synthesis in IL-2 maintained CTL. To test this hypothesis we used a single cell assay that quantifies the incorporation of an analogue of puromycin (OPP) into newly synthesized protein chains in the ribosome to assess cellular rates of protein synthesis. We found that 24 hours of treatment with pan-PIM kinase inhibitors PIM447 or AZD1208 caused a quantitative reduction in protein synthesis rates of IL-2 expanded CTL (**Fig 5M**). This data is consistent with PIM kinase effects on protein expression being via control of protein translation.

PIM1 and PIM2 regulate mTORc1 and sustain lymphoid homing in Pim1/Pim2-deficient effector CTL

The pattern of proteome remodelling in Pim dKO CTL had some phenotypic features in common with the previously reported changes that occur following inhibition of the Serine/Threonine kinase mTORc1 in CTL (Hukelmann et al., 2016; Howden et al., 2019). These included: increased expression of PDCD4 and decreased expression of cholesterol biosynthesis enzymes and effector molecules including granzymes, perforin, IFNγ and TNFα (Hukelmann et al., 2016; Howden et al., 2019). In this respect, mTORc1 activity in CTL is acutely sensitive to reduced availability of glucose (Rolf et al., 2013) and the reduced expression of glucose transporters in Pim dKO T cells might predict that these cells would have lower activity of mTORc1. We therefore assessed if PIM kinases were required for IL-2 activation of mTORc1. In these experiments we compared the phosphorylation of S6K1 (p70 S6-kinase 1) on its mTORC1 substrate sequence T389 in WT and Pim dKO CD8 T cells by western blot. These data show a modest reduction of mTORc1 activity in the Pim dKO T cells

(**Fig 6A**). We also assessed the activity of S6K1 at a single cell level by quantifying the

phosphorylation of its downstream target protein S6 ribosomal protein on S235/S236 by

flow cytometry over the course of IL-2 driven CTL expansion. These data also show that

Pim1/Pim2-deficiency reduced IL-2 induced mTORc1 dependent phosphorylation of S6 but

did not compromise the ability of the Pim dKO cells to phosphorylate S6 in response to

phorbol esters and calcium ionophores; pharmacological activators that induce RSK

mediated S6 phosphorylation (**Fig 6B**).

Previously, it has been shown that decreased glucose availability or inhibition of mTORc1

prevents the downregulation of CD62L that normally accompanies IL-2 induced CTL

differentiation (Sinclair et al., 2008; Finlay et al., 2012). Thus, the reduced expression of

glucose transporters and reduced activity of mTORc1 in Pim dKO IL-2 expanded CTL provides

a potential explanation for the striking phenotype shown in Fig 3G, that IL-2 maintained Pim

dKO CTL retain high levels of CD62L/L-selectin. The IL-2 CTL transcriptomics data showed

there is increased expression of CD62L at the mRNA level in IL-2 maintained PimdKO CTL (**Fig**

5B, 6C). The expression of *Sell* (CD62L) mRNA is controlled by the transcription factor KLF2

and the expression of *Klf2* mRNA is also increased in IL-2 maintained Pim dKO CTL (**Fig 6D**) as

are mRNA levels for other KLF2 targets: the chemokine receptors *Ccr7* and *S1pr1 (***Fig 6C***)*.

CD62L/L-selectin controls T cell adhesion to the endothelium of high endothelial venules

and is essential for lymphocyte transit from blood into secondary lymphoid tissue such as

lymph nodes. CCR7, and the $S1P_1$ receptor also direct migration by controlling T cell entry

and egress respectively into secondary lymphoid tissue. In this context, the loss of CD62L,

CCR7 and S1PR1 expression following immune activation reprograms the trafficking of

effector T cells away from lymphoid tissue toward sites of inflammation in peripheral tissues

(Sinclair et al., 2008; Nolz et al., 2011). Accordingly, the differences between Pim dKO and

WT CTL in terms of CD62L expression could indicate that in addition to having reduced

effector protein expression, Pim dKO CTL also do not switch their trafficking behaviour to

that of a fully differentiated effector CTL and rather retain the capacity to home to lymphoid

tissues. To test this possibility *in vivo*, we performed adoptive transfer experiments

comparing the ability of WT effector CTL and Pim dKO CTL to home to secondary lymphoid

tissues. In these experiments, αCD3/αCD28 activated WT or Pim dKO T cells were

differentiated in IL-2 and cells then were labelled with either CFSE or CTV dyes; mixed at a

ratio of 1:1, and adoptively transferred into C57BL/6 hosts. **Fig 6E** shows that Pim dKO CTL,

but not WT CTL, retained substantial capacity to home to secondary lymphoid organs and,

hence, accumulated in lymph nodes and spleen: PIM kinases are thus required for the

normal reprogramming of CTL trafficking.

Discussion:

The objective of the present study was to use quantitative analysis of T cell proteomes and

transcriptomes to explore how deletion of the Serine/Threonine kinases PIM1 and PIM2

impacts CD8 T cell activation and differentiation. The expression of PIM1 and PIM2 is

induced during immune activation of CD8 T cells by antigen and co-stimulation and then

sustained to varying degrees by the cytokines IL2 or IL-15. The loss of PIM1 and PIM2 was

shown to have no effect on the CD8 T cell programs controlled by antigen receptor/co-

stimulation and had very little impact on the IL-15-driven transcriptional and proteomic

programs in memory differentiated CD8 T cells. However, PIM1 and PIM2 were required for

IL-2 programming of effector CD8 T cell differentiation. IL-2 controls essential cytolytic

effector and metabolic programs in CTL, is a key cytokine for CD8 T cell anti-viral immunity

and is also used to produce cytotoxic T cells for anti-cancer immunotherapy (Kalia and

Sarkar, 2018). The understanding that PIM kinases mediate IL-2 control of the expression of

glucose transporters, lipid metabolic enzymes and key cytolytic proteins such as granzymes

A and B in effector CD8 T cells is thus fundamental information about how this key cytokine

controls T cells. It was also important to understand that PIM kinases did not control the

transcriptional programs that direct expression of glucose transporters and effector

enzymes and the hence the full consequences of PIM kinase deletion could only be assessed

by in depth analysis of the proteomes of *Pim1*/*Pim2*-deficient CTL and not from mRNA

sequencing. The loss of PIM kinase activity reduced protein synthesis rates in CTL which

explains the discordance between mRNA and protein reported herein. The reduction in

protein synthesis was modest but previous studies have shown that modest reductions in

protein synthesis and loss of mTORc1 activity impact CD8 T cell differentiation programs

(Cornish et al., 2006; Hukelmann et al., 2016; Howden et al., 2019). PIM kinase regulation of

protein translation has been reported previously in mouse B cell lymphoma and mouse

embryonic fibroblasts (Schatz et al., 2011) and some of the key examples of proteins we

found to be unaltered at the mRNA level but down-regulated at the protein level have prior

reports of sensitivity to translational regulation in immune cells (Fehniger et al., 2007;

Ricciardi et al., 2018). Reduced protein synthesis capacity in cells lacking PIM kinases could

be caused by reduced mTORc1 activity combined with increased expression at the mRNA

and protein level of the translational repressor PDCD4. Moreover, protein synthesis is highly

energetically demanding and once CTL have reduced glucose transporter expression this

would reduce glucose availability to fuel protein production. These data collectively point

towards modulation of protein translation as an important target of PIM kinase activity

downstream of IL-2 in CD8 T cells.

One other important insight from the current work is that PIM kinases are required for IL2

to fully repress expression of the adhesion molecule CD62L and the chemokine receptors

CCR7 and S1PR1. These molecules play a key role in controlling T cell homing to secondary

lymphoid tissue and are expressed at high levels in naïve and memory T cells. IL-2 induces

down regulation of CD62L, CCR7 and S1PR1 expression as part of a program that causes

effector cells to lose their capacity to home to secondary lymphoid tissues (Sinclair et al.,

2008; Nolz et al., 2011). The present data show that PIM kinases can mediate this IL-2

repression of lymph node homing receptors and the ability of IL-2 to switch off the naïve T

cell trafficking program. IL2 maintained Pim dKO CD8 effector T cells thus retain the capacity

to home to secondary lymphoid tissues *in vivo*. This lymph-node homing pattern is also a

memory CD8 T cell phenotype but here it is pertinent that the current in-depth analysis of

the transcriptomes and proteomes of *Pim1/Pim2*-deficient IL-2 differentiated CD8 T cells

show that the loss of PIM kinases does not cause effector T cells to differentiate instead into

memory cells. For example, IL-2 maintained Pim dKO CD8 T cells do not re-express key

molecules that maintain the stem-like properties of memory cells such as the transcription

factor TCF7 and IL-7 receptor (Table S2, S6). Nor do Pim dKO CTL cultured in IL-2 have the

mitochondrial proteome of a memory T cell. Hence *Pim1/Pim2*-deficiency prevents optimal

differentiation of effector CTL but does not cause these cells to fully switch to a memory

phenotype. This is worth knowing as pan-PIM kinase inhibitors are in development as anti-

cancer drugs and it has been suggested that PIM inhibitors might be used as a strategy to

generate more stem-memory-like CD8 T cells in combination with cancer immunotherapy

treatments (Chatterjee et al., 2019; Clements and Warfel, 2022). Information about how the

loss of PIM kinases will impact CD8 T cells is very relevant to informed usage of such drugs.

Finally, the selectivity of PIM kinases for IL-2 control of conventional CD8 T cell

differentiation with no evidence that PIMs were important for IL-15 responses beyond a

modest effect on cell proliferation was interesting. This failure to see a role for PIM

signalling in conventional peripheral CD8 T cell differentiation is in contrast to observations

that PIM1 and PIM2 direct IL15 control of the metabolic programming of intestinal IEL

(James et al., 2021). A notable difference between these scenarios was that in IEL, IL15 was

being used individually to drive immune activation and was sufficient to strongly induce

PIM1 and PIM2 expression. In contrast, in our conventional CD8 T cell system we uncovered

that that IL-15 does not sustain high levels of PIM kinases, which could possibly explain this

discrepancy. However, high levels of PIM expression alone are not inevitably predictive of

their importance. In this study we observed very high levels of both PIM1 and PIM2 in

antigen activated CD8 T cells yet loss of PIM kinases there had no functional impact. There

are thus clearly redundant kinase pathways that can substitute for the PIM kinases in the

immediate response of CD8 T cells to immune activation. Previously it was suggested that

mTORc1 may be one of these pathways (Fox et al., 2005), however we found no evidence

for that here, and in fact observed that PIM kinase deficiency was upstream of a reduction

in mTORc1 activity in IL2 expanded CTL. Thus, this examination of the role of PIM kinases in

multiple signalling situations reveal cell context dependent regulatory roles for PIM kinases

rather than hard wired functions that apply to every stimulatory situation in T cells.

References:

Aho, T. L., R. J. Lund, E. K. Ylikoski, S. Matikainen, R. Lahesmaa, and P. J. Koskinen. 2005.
Expression of human pim family genes is selectively up-regulated by cytokines
promoting T helper type 1, but not T helper type 2, cell differentiation. *Immunology*
116: 82-88.
Blagih, J., F. Coulombe, E. E. Vincent, F. Dupuy, G. Galicia-Vazquez, E. Yurchenko, T. C. Raissi,
et al. 2015. The energy sensor AMPK regulates T cell metabolic adaptation and
effector responses in vivo. *Immunity* 42: 41-54.
Buchacher, T., A. Shetty, S. A. Koskela, J. Smolander, R. Kaukonen, A. G. G. Sousa, S. Junttila,
et al. 2023. PIM kinases regulate early human Th17 cell differentiation. *Cell Rep* 42:
113469.
Buck, M. D., D. O'Sullivan, R. I. Klein Geltink, J. D. Curtis, C. H. Chang, D. E. Sanin, J. Qiu, et al.
2016. Mitochondrial Dynamics Controls T Cell Fate through Metabolic Programming.
Cell 166: 63-76.
Bullock, A. N., J. Debreczeni, A. L. Amos, S. Knapp, and B. E. Turk. 2005. Structure and
substrate specificity of the Pim-1 kinase. *J Biol Chem* 280: 41675-41682.
Cantrell, D. 2015. Signaling in lymphocyte activation. *Cold Spring Harb Perspect Biol* 7.
Chang, C. H., J. Qiu, D. O'Sullivan, M. D. Buck, T. Noguchi, J. D. Curtis, Q. Chen, et al. 2015.
Metabolic Competition in the Tumor Microenvironment Is a Driver of Cancer
Progression. *Cell* 162: 1229-1241.
Chatterjee, S., P. Chakraborty, A. Daenthanasanmak, S. Iamsawat, G. Andrejeva, L. A.
Luevano, M. Wolf, et al. 2019. Targeting PIM Kinase with PD1 Inhibition Improves
Immunotherapeutic Antitumor T-cell Response. *Clin Cancer Res* 25: 1036-1049.
Chauhan, S. S., R. K. Toth, C. C. Jensen, A. L. Casillas, D. F. Kashatus, and N. A. Warfel. 2020.
PIM kinases alter mitochondrial dynamics and chemosensitivity in lung cancer.
Oncogene 39: 2597-2611.
Clements, A. N., and N. A. Warfel. 2022. Targeting PIM Kinases to Improve the Efficacy of
Immunotherapy. *Cells* 11.
Cornish, G. H., L. V. Sinclair, and D. A. Cantrell. 2006. Differential regulation of T-cell growth
by IL-2 and IL-15. *Blood* 108: 600-608.
Damasio, M. P., J. M. Marchingo, L. Spinelli, J. L. Hukelmann, D. A. Cantrell, and A. J. M.
Howden. 2021. Extracellular signal-regulated kinase (ERK) pathway control of CD8+ T
cell differentiation. *Biochem J* 478: 79-98.
Din, S., M. Mason, M. Volkers, B. Johnson, C. T. Cottage, Z. Wang, A. Y. Joyo, et al. 2013. Pim-
1 preserves mitochondrial morphology by inhibiting dynamin-related protein 1
translocation. *Proc Natl Acad Sci U S A* 110: 5969-5974.
Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, et al. 2013. STAR:
ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15-21.
Ewels, P., M. Magnusson, S. Lundin, and M. Kaller. 2016. MultiQC: summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* 32: 3047-3048.
Fehniger, T. A., S. F. Cai, X. Cao, A. J. Bredemeyer, R. M. Presti, A. R. French, and T. J. Ley.
2007. Acquisition of murine NK cell cytotoxicity requires the translation of a pre-
existing pool of granzyme B and perforin mRNAs. *Immunity* 26: 798-811.
Finlay, D. K., E. Rosenzweig, L. V. Sinclair, C. Feijoo-Carnero, J. L. Hukelmann, J. Rolf, A. A.
Panteleyev, et al. 2012. PDK1 regulation of mTOR and hypoxia-inducible factor 1
integrate metabolism and migration of CD8+ T cells. *J Exp Med* 209: 2441-2453.

Fox, C. J., P. S. Hammerman, and C. B. Thompson. 2005. The Pim kinases control rapamycin-
resistant T cell survival and activation. *J Exp Med* 201: 259-266.
Howden, A. J. M., J. L. Hukelmann, A. Brenes, L. Spinelli, L. V. Sinclair, A. I. Lamond, and D. A.
Cantrell. 2019. Quantitative analysis of T cell proteomes and environmental sensors
during T cell differentiation. *Nat Immunol* 20: 1542-1554.
Hughes, C. S., S. Foehr, D. A. Garfield, E. E. Furlong, L. M. Steinmetz, and J. Krijgsveld. 2014.
Ultrasensitive proteome analysis using paramagnetic bead technology. *Mol Syst Biol*
10: 757.
Hukelmann, J. L., K. E. Anderson, L. V. Sinclair, K. M. Grzes, A. B. Murillo, P. T. Hawkins, L. R.
Stephens, et al. 2016. The cytotoxic T cell proteome and its shaping by the kinase
mTOR. *Nat Immunol* 17: 104-112.
Jackson, L. J., J. A. Pheneger, T. J. Pheneger, G. Davis, A. D. Wright, J. E. Robinson, S. Allen, et
al. 2012. The role of PIM kinases in human and mouse CD4+ T cell activation and
inflammatory bowel disease. *Cell Immunol* 272: 200-213.
James, O. J., M. Vandereyken, J. M. Marchingo, F. Singh, S. E. Bray, J. Wilson, A. G. Love, and
M. Swamy. 2021. IL-15 and PIM kinases direct the metabolic programming of
intestinal intraepithelial lymphocytes. *Nat Commun* 12: 4290.
Kalia, V., and S. Sarkar. 2018. Regulation of Effector and Memory CD8 T Cell Differentiation by
IL-2-A Balancing Act. *Front Immunol* 9: 2987.
Knudson, K. M., C. J. Pritzl, V. Saxena, A. Altman, M. A. Daniels, and E. Teixeiro. 2017.
NFkappaB-Pim-1-Eomesodermin axis is critical for maintaining CD8 T-cell memory
quality. *Proc Natl Acad Sci U S A* 114: E1659-E1667.
Laird, P. W., N. M. van der Lugt, A. Clarke, J. Domen, K. Linders, J. McWhir, A. Berns, and M.
Hooper. 1993. In vivo analysis of Pim-1 deficiency. *Nucleic Acids Res* 21: 4750-4755.
Law, C. W., M. Alhamdoosh, S. Su, X. Dong, L. Tian, G. K. Smyth, and M. E. Ritchie. 2016. RNA-
seq analysis is easy as 1-2-3 with limma, Glimma and edgeR. *F1000Res* 5.
Lawrence, M., W. Huber, H. Pages, P. Aboyoun, M. Carlson, R. Gentleman, M. T. Morgan, and
V. J. Carey. 2013. Software for computing and annotating genomic ranges. *PLoS*
Comput Biol 9: e1003118.
Li, H., L. Xie, L. Zhu, Z. Li, R. Wang, X. Liu, Z. Huang, et al. 2022. Multicellular immune
dynamics implicate PIM1 as a potential therapeutic target for uveitis. *Nat Commun*
13: 5866.
Locard-Paulet, M., G. Voisinne, C. Froment, M. Goncalves Menoita, Y. Ounoughene, L. Girard,
C. Gregoire, et al. 2020. LymphoAtlas: a dynamic and integrated phosphoproteomic
resource of TCR signaling in primary T cells reveals ITSN2 as a regulator of effector
functions. *Mol Syst Biol* 16: e9524.
Maney, N. J., H. Lemos, B. Barron-Millar, C. Carey, I. Herron, A. E. Anderson, A. L. Mellor, et
al. 2021. Pim Kinases as Therapeutic Targets in Early Rheumatoid Arthritis. *Arthritis*
Rheumatol 73: 1820-1830.
Marchingo, J. M., L. V. Sinclair, A. J. Howden, and D. A. Cantrell. 2020. Quantitative analysis of
how Myc controls T cell proteomes and metabolic pathways during T cell activation.
Elife 9.
Matikainen, S., T. Sareneva, T. Ronni, A. Lehtonen, P. J. Koskinen, and I. Julkunen. 1999.
Interferon-alpha activates multiple STAT proteins and upregulates proliferation-
associated IL-2Ralpha, c-myc, and pim-1 genes in human T cells. *Blood* 93: 1980-
1991.

McCarthy, D. J., Y. Chen, and G. K. Smyth. 2012. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res* 40: 4288-
4297.
McCarthy, D. J., K. R. Campbell, A. T. Lun, and Q. F. Wills. 2017. Scater: pre-processing, quality
control, normalization and visualization of single-cell RNA-seq data in R.
Bioinformatics 33: 1179-1186.
Mikkers, H., M. Nawijn, J. Allen, C. Brouwers, E. Verhoeven, J. Jonkers, and A. Berns. 2004.
Mice deficient for all PIM kinases display reduced body size and impaired responses
to hematopoietic growth factors. *Mol Cell Biol* 24: 6104-6115.
Navarro, M. N., J. Goebel, J. L. Hukelmann, and D. A. Cantrell. 2014. Quantitative
phosphoproteomics of cytotoxic T cells to reveal protein kinase d 2 regulated
networks. *Mol Cell Proteomics* 13: 3544-3557.
Navarro, M. N., J. Goebel, C. Feijoo-Carnero, N. Morrice, and D. A. Cantrell. 2011.
Phosphoproteomic analysis reveals an intrinsic pathway for the regulation of histone
deacetylase 7 that controls the function of cytotoxic T lymphocytes. *Nat Immunol* 12:
352-361.
Nolz, J. C., G. R. Starbeck-Miller, and J. T. Harty. 2011. Naive, effector and memory CD8 T-cell
trafficking: parallels and distinctions. *Immunotherapy* 3: 1223-1233.
Peng, C., A. Knebel, N. A. Morrice, X. Li, K. Barringer, J. Li, S. Jakes, et al. 2007. Pim kinase
substrate identification and specificity. *J Biochem* 141: 353-362.
Peperzak, V., E. A. Veraar, A. M. Keller, Y. Xiao, and J. Borst. 2010. The Pim kinase pathway
contributes to survival signaling in primed CD8+ T cells upon CD27 costimulation. *J
Immunol* 185: 6670-6678.
Ricciardi, S., N. Manfrini, R. Alfieri, P. Calamita, M. C. Crosti, S. Gallo, R. Muller, et al. 2018.
The Translational Machinery of Human CD4(+) T Cells Is Poised for Activation and
Controls the Switch from Quiescence to Metabolic Remodeling. *Cell Metab* 28: 961.
Ritchie, M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law, W. Shi, and G. K. Smyth. 2015. limma
powers differential expression analyses for RNA-sequencing and microarray studies.
Nucleic Acids Res 43: e47.
Robinson, M. D., D. J. McCarthy, and G. K. Smyth. 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* 26:
139-140.
Rolf, J., M. Zarrouk, D. K. Finlay, M. Foretz, B. Viollet, and D. A. Cantrell. 2013. AMPKalpha1: a
glucose sensor that controls CD8 T-cell memory. *Eur J Immunol* 43: 889-896.
Saravia, J., J. L. Raynor, N. M. Chapman, S. A. Lim, and H. Chi. 2020. Signaling networks in
immunometabolism. *Cell Res* 30: 328-342.
Schatz, J. H., E. Oricchio, A. L. Wolfe, M. Jiang, I. Linkov, J. Maragulia, W. Shi, et al. 2011.
Targeting cap-dependent translation blocks converging survival signals by AKT and
PIM kinases in lymphoma. *J Exp Med* 208: 1799-1807.
Sinclair, L. V., D. Finlay, C. Feijoo, G. H. Cornish, A. Gray, A. Ager, K. Okkenhaug, et al. 2008.
Phosphatidylinositol-3-OH kinase and nutrient-sensing mTOR pathways control T
lymphocyte trafficking. *Nat Immunol* 9: 513-521.
Sinclair, L. V., A. J. Howden, A. Brenes, L. Spinelli, J. L. Hukelmann, A. N. Macintyre, X. Liu, et
al. 2019. Antigen receptor control of methionine metabolism in T cells. *Elife* 8.
Sollberger, G., A. J. Brenes, J. Warner, J. S. C. Arthur, and A. J. M. Howden. 2023. Quantitative
proteomics reveals tissue-specific, infection-induced and species-specific neutrophil

580   protein signatures. *bioRxiv* 10.1101/2023.09.05.556395 DOI:
581   2023.2009.2005.556395.
582 Spinelli, L., J. M. Marchingo, A. Nomura, M. P. Damasio, and D. A. Cantrell. 2021.
583   Phosphoinositide 3-Kinase p110 Delta Differentially Restrains and Directs Naive
584   Versus Effector CD8(+) T Cell Transcriptional Programs. *Front Immunol* 12: 691997.
585 Storey, J. D., A. J. Bass, A. Dabney, and D. Robinson. 2023. qvalue: Q-value estimation for
586   false discovery rate control, version R package version 2.32.0.
587 Suzuki, C., R. G. Garces, K. A. Edmonds, S. Hiller, S. G. Hyberts, A. Marintchev, and G. Wagner.
588   2008. PDCD4 inhibits translation initiation by binding to eIF4A using both its MA3
589   domains. *Proc Natl Acad Sci U S A* 105: 3274-3279.
590 Swamy, M., S. Pathak, K. M. Grzes, S. Damerow, L. V. Sinclair, D. M. van Aalten, and D. A.
591   Cantrell. 2016. Glucose and glutamine fuel protein O-GlcNAcylation to control T cell
592   self-renewal and malignancy. *Nat Immunol* 17: 712-720.
593 Tan, H., K. Yang, Y. Li, T. I. Shaw, Y. Wang, D. B. Blanco, X. Wang, et al. 2017. Integrative
594   Proteomics and Phosphoproteomics Profiling Reveals Dynamic Signaling Networks
595   and Bioenergetics Pathways Underlying T Cell Activation. *Immunity* 46: 488-503.
596 Warfel, N. A., and A. S. Kraft. 2015. PIM kinase (and Akt) biology and signaling in tumors.
597   *Pharmacol Ther* 151: 41-49.
598 Weninger, W., M. A. Crowley, N. Manjunath, and U. H. von Andrian. 2001. Migratory
599   properties of naive, effector, and memory CD8(+) T cells. *J Exp Med* 194: 953-966.
600 Wingett, D., A. Long, D. Kelleher, and N. S. Magnuson. 1996. pim-1 proto-oncogene
601   expression in anti-CD3-mediated T cell activation is associated with protein kinase C
602   activation and is independent of Raf-1. *J Immunol* 156: 549-557.
603 Wisniewski, J. R., M. Y. Hein, J. Cox, and M. Mann. 2014. A "proteomic ruler" for protein copy
604   number and concentration estimation without spike-in standards. *Mol Cell*
605   *Proteomics* 13: 3497-3506.
606 Zougman, A., P. J. Selby, and R. E. Banks. 2014. Suspension trapping (STrap) sample
607   preparation method for bottom-up proteomics analysis. *Proteomics* 14: 1006-1000.
608

Acknowledgements:
The authors thank members of the Cantrell group for their critical discussion of the data, in
particular we thank Linda Sinclair for critical reading and comments on the manuscript. We
thank Victor Tybulewicz for providing the Pim1-/- and Pim2-/- mice, E Emslie for help with
mouse maintenance and experimental support, B Stubbs for technical assistance with
mouse genotyping, A Brenes for assistance with analytical tools, A Gardner, A Rennie and R
Clarke from the Flow Cytometry facility for cell sorting, K Beattie and A Atrih from the
Fingerprint proteomics facility for performed mass spectrometry, S Thomson for assistance
with adoptive transfer experiments, the Biological Resource Unit at the University of
Dundee and the Finnish Functional Genomics Centre, University of Turku and Abo Akademi
Biocenter Finland for performing RNA sequencing. This research was supported by a
Wellcome Trust Principal Research Fellowship to DAC (205023/Z/16/Z), a Wellcome Trust
Equipment Award to DAC (202950/Z/16/Z), an EMBO Long term Fellowship (ALTF 1543–
2015) and NHMRC CJ Martin Early Career fellowship (APP1120748) to JMM and has received
funding from the European Union's Horizon 2020 research and innovation programme
under the Marie Skłodowska-Curie grant agreement No 705984 to JMM and DAC.
625

Materials and Methods:

Mice

Pim1$^{-/-}$ (Pim1 KO)(Laird et al., 1993) and Pim2$^{-/-}$ or Pim2$^{-/Y}$ (Pim2 KO) mice(Mikkers et al.,
2004) on the FVB/N background were backcrossed for >10 generations onto a C57BL/6
background and were a generous gift from Prof Victor Tybulewicz. Pim1 KO and Pim2 KO
mice were inter-crossed to generate the Pim1/Pim2 double KO (Pim dKO) strain. Pim dKO,
C57BL/6, Ly5.1, P14 and OT1 mice were bred and maintained in the WTB/RUTG, University
of Dundee in compliance with UK Home Office Animals (Scientific Procedures) Act 1986
guidelines. Mice used for proteomics and RNAseq studies were male and 12-15 weeks of
age. Age/sex matched mice were used for all other experiments between 5–52 weeks of
age, with most being between 10 – 20 weeks.

Cell culture

Single cell suspensions were generated by mashing mouse lymph nodes (brachial, axial,
inguinal, superficial cervical, deep cervical, lumbar) or spleens through a 70 μm strainer. Red
blood cells were lysed with Ack buffer (150 mM NH$_4$Cl 10 mM KHCO$_3$110 μM Na$_2$EDTA pH
7.8). Cells were cultured in RPMI 1640 containing glutamine (Invitrogen), supplemented
with 10% FBS (Gibco), penicillin/streptomycin (Gibco) and 50 μM β-mercaptoethanol
(Sigma) at 37 °C with 5% CO$_2$.

For 24 hour TCR activation WT and Pim dKO proteomics and phenotyping, lymph node
suspension from 2 mice per biological replicate were activated with 0.5 μg/mL anti-mouse
CD3 (Biolegend) and 0.5 μg/mL anti-mouse CD28 (eBioscience) in 2 × 10 mL complete
culture medium in six well plates. Proteomics samples were generated in biological
triplicate.

For generation of memory-like or effector cytotoxic T lymphocytes (CTL) from mice with
polyclonal T cell repertoires, LN or spleen single cell suspensions were activated with 0.5
μg/mL anti-mouse CD3 (Biolegend) and 0.5 μg/mL anti-mouse CD28 (eBioscience) and
recombinant human IL-15 (20 ng/mL, Peprotech) or recombinant human IL-2 (20 ng/mL,
Proleukin, Novartis) in 10 mL complete culture medium per organ per mouse. After 2 days
of activation (~40 hours) cells were washed out of activation medium and resuspended at

0.3 million T cells/mL in fresh culture medium and cytokine 20 ng/mL of IL-15 or IL-2 for

memory-like and effector T cells respectively. T cells were subsequently split into fresh

medium and cytokine daily at a density of 0.5 million/ mL and 0.3 million/mL for memory-

like and effector T cells respectively until analysis timepoint.

For generation of effector CTL from TCR-transgenic P14 mice LN single cell suspensions were

activated with 100 ng/mL gp33 peptide, recombinant human IL-2 (20 ng/mL) and

recombinant mouse IL12 (2 ng/mL) (Peprotech) for 2 days, before washing out of activation

media and splitting cells daily into fresh media containing IL-2 (20 ng/mL) at a density of 0.3

million/mL.

For co-culture experiments Ly5.1+ WT and Pim dKO (Ly5.2+) cell suspensions were mixed in

a 50:50 ratio based on total T cell numbers before activation. A portion of the mixed naïve T

cells maintained in recombinant mouse IL-7 (5 ng/mL, Peprotech) to verify baseline T cell

ratios over time in culture.

For experiments where a pure CD8+ CTL population were required from non-TCR transgenic

mice (adoptive transfer, proteomics and RNAseq experiments), CD4 T cells were depleted

on day 3 of CTL cultures (i.e. 24 hours after being washed out of initial activation conditions)

by negative magnetic selection. T cells were resuspended in MACS buffer (PBS, 1mM EDTA,

2%FBS) at 10^8/mL, blocked with 50 μL/mL Rat Serum, incubated for 10 min with anti-CD4

biotin antibody (5 μg/mL, Biolegend), incubated for 5 min with 125 μL/mL RapidSphere

beads (StemCell Technologies), then volume topped up to 2.5 mL and placed in EasySep

magnet for 2.5 min. Supernatant containing CD8 T cells was collected and returned to

culture as per CTL protocol above. CD4 T cell depletion was confirmed by flow cytometry,

with CD4+ cells making up <0.4% of live cells.

In experiments where cytokine was withdrawn T cells were washed at least twice with warm

complete medium. Where indicated CTL were treated with 100 nM Tofacitinib

(Selleckchem), 20 nM of rapamycin (Merck), 20 ng/mL PDBu (Cell Signaling Technologies)

and ionomycin 500 ng/mL (Merck) or the pan PIM kinase inhibitors PIM447 (1 or 5 μM as

indicated) and AZD1208 (1 or 10 μM as indicated) (both Medchemexpress).

Flow cytometry

Flow cytometry data was acquired on a FACSVerse using FACSuite software, FACSCanto, or LSR II Fortessa with FACS DIVA software (BD Biosciences) or Novocyte (Acea Biosciences Inc.) with NovoExpress software (Agilent). Data was analysed using Flowjo software version 9.9.6 or version 10.6.1 and above (Treestar).

Cell surface staining antibodies conjugated to BV421, BV510, V500, FITC, PE, PerCPCy5.5, PECy7, APC, A647 and APCeF780 were obtained from BD Biosciences, eBioscience or Biolegend. Fc receptors were blocked using Fc block (BD Biosciences). Antibody staining for surface markers was performed at 1:200 in PBS 1%FBS. Antibody clones were as follows: CD4 (RM4-5), CD8 (53–6.7), CD25 (7D4 or PC61), CD44 (IM7), CD45.1 (A20), CD45.2 (104), CD62L (MEL-14), CD69 (H1.2F), CD71 (RI7217), TCRbeta (H57-597).

For IFNγ and Granzyme B intracellular staining cells were fixed and permeabilised using eBioscience Intracellular Fixation and Permeabilisation kit (eBioscience) as per manufacturer instructions. Cells were stained with anti-IFNγ (XMG1.2) and anti-Granzyme B (NGZB) at 1:100 and 1:200 respectively. For intracellular IFNγ staining GolgiPlug (BD) was added to the culture 4 hours prior to analysis.

For intracellular phospho S6 staining cells were fixed with 0.5% paraformaldehyde at 37 °C and permeabilised with ice cold 90% methanol. Cells were stained at 1:100 with anti-phospho S6 ribosomal protein (Ser235/236) Alexa Fluor 647 (D57.2.2E, Cell Signaling Technologies cat#4851S).

For intracellular phospho-STAT5 staining cells were fixed with 1% paraformaldehyde at room temperature and permeabilised with ice cold 90% methanol. Cells were stained at 1:200 with rabbit anti-phospho STAT5 (Y694) (C11C5, Cell Signaling Technologies cat#9359S) followed by anti-rabbit IgG Fab2 Alexa Fluor 647 (Cell Signaling Technologies cat#4414S).

For intracellular PDCD4 staining cells were fixed with 1% paraformaldehyde at room temperature and permeabilised with Perm Buffer (eBioscience, cat# 00-8333-56). Cells were stained at 1:100 with rabbit anti-PDCD4 (D29C6, Cell Signaling Technologies cat#9535S) followed by anti-rabbit IgG Fab2 Alexa Fluor 647 (Cell Signaling Technologies cat#4414S).

Cell sorting

Cell sorting was performed on a Sony SH800S cell sorter (Sony Biotechnology). Staining was performed in PBS 1% FBS and sorting and collection of cells for proteomics analysis was performed in RPMI 1640 containing glutamine, supplemented with 1% FBS. 24 hour TCR activated WT and Pim dKO CD4 and CD8 T cells were sorted as: DAPI-CD69+ CD8+ or CD4+. Cells were washed twice with HBSS (no calcium, no magnesium, no phenol red, GIBCO) before being snap frozen in liquid nitrogen and stored at −80°C until further processing.

CTL number estimation and viability

Absolute CD8 T cell number per well/flask and % live cells was assessed in CTL cultures by volumetric measurement of cell number from 50 µL culture medium. Cells were transferred into 450 µL PBS containing low concentration anti-CD8, anti-CD4 antibody (both 1:1500) and DAPI (1 µg/mL, Thermofisher Scientific) or propidium iodide (0.2 µg/mL, Sigma) and cell number and viability assessed on either FACSVerse or Novocyte flow cytometers. Estimated live cell counts were corrected for assay dilution ratio, total culture volume and splitting ratios for each time point. Proportions of CD8 T cells that were Ly5.1+ or Ly5.2+ from parallel surface staining were used to calculate WT and Pim dKO numbers respectively in co-culture experiments.

CTV proliferation assay

For CTV proliferation assay, lymph node suspensions at 10^7 cells per mL in PBS 0.5%FCS were labelled with 5 µM CTV (Invitrogen) for 20 min at 37°C before reaction was quenched and cells washed with cold medium. 10^5 live cells were activated with αCD3/αCD28 (both 0.5 µg/mL) or maintained in IL-7 (5 ng/mL) +/- rapamycin (20 nM) in 200 µL total volume in 96 well flat-bottomed plates. Propidium iodide (0.2 µg/mL), low dose αCD4, αCD8 antibody (both 1:3200 final concentration) and 10,000 sphero rainbow beads (BD) were added to wells prior to analysis of beads number, cell number, cell viability, CD4 and CD8 T cell composition and CTV profile of samples by flow cytometry. The ratio between the known number of beads added to the number of beads measured per well was multiplied with the number of cells measured to calculate the absolute cell number per well. Cell number per division was then corrected for the effect of expansion by dividing cell number per division

753 by 2^division number and the arithmetic mean division number calculated from these

754 values.

Protein synthesis assay

To assess protein synthesis cells were cultured with 20 µM O-propargyl puromycin (OPP,

Jena Bioscience) for 10 min. Control cells were cultured with cycloheximide (100 µg/mL,

Sigma-Aldrich) for 20 min prior to addition of OPP to assess background staining level when

cytosolic protein synthesis is completely inhibited. Cells were washed, fixed with 1% PFA,

permeabilised with 0.5% Triton x-100 and OPP incorporation into newly synthesised

peptides was measured by conjugating with Alexa647-azide via a Click-IT chemistry reaction

(2 µM Alexa-647-azide, 2 mM CuSO$_4$, 5 mM ascorbic acid in PBS) (Invitrogen) and assessing

fluorescence by flow cytometry.

Adoptive Transfer

WT and Pim dKO CD8 T cells were activated and expanded in IL-2 as described above. On

day 6 of culture WT and Pim dKO CTL were labelled with either CTV, or 5 µM CFSE

(Invitrogen) as per CTV assay protocol described above, with the exception that labelling

was performed in RPMI 1640 and CFSE labelling was for 10 min. CTV and CFSE labelling was

alternated between genotypes across biological replicates to prevent any potential

introduction of bias due to possible differences in dye toxicity. Labelled WT and Pim dKO

CD8 T cells were mixed in a 50:50 ratio and 5 million cells/mouse were adoptively

transferred via intravenous tail injection into C57BL/6 recipient mice. Recipient mice were

sacrificed and blood, spleen and lymph nodes (axillary, brachial and inguinal) removed 4 and

23 hours post-transfer and the ratio of CTV to CFSE cells assessed by flow cytometry.

Western blotting

Cell pellets were lysed in RIPA buffer (100 mM HEPES, pH 7.4, 150 mM NaCl, 1% NP40, 0.1%

SDS, 0.5% sodium deoxycholate, 10% glycerol, 1 mM EDTA, 1 mM EGTA, 1 mM TCEP, and

protease and phosphatase inhibitors (Roche)) at 20 million T cells/mL at 4 °C. Lysates were

sonicated, centrifuged 12 min, 13,000 rpm, 4°C. Supernatant was transferred to fresh tube

and NuPAGE LDS sample buffer (1×) (Life Technologies) and tris(2-carboxyethyl)phosphine

(TCEP, 25mM, Thermofisher) was added before boiling samples at 100°C for 3 min. Samples

were loaded and separated by SDS-PAGE (NuPAGE 4-12% gradient precast gels (ThermoFisher) or 12% polyacrylamide running gel), and then transferred to nitrocellulose membranes (Whatman). Equal cell numbers (~140,000 cells) were loaded per lane. Blots were probed with the following primary antibodies: S6K (5G10, Cell Signaling Technology, cat# 2217S), S6K p-T389 (108D2, Cell Signaling Technology, cat# 9234S), STAT5 p-Y694 (C11C5, Cell Signalling Technology, cat# 9359S), Pim1 (12H8, Santa Cruz, cat# SC-13513), Pim2 (1D12, Santa Cruz, cat# SC-13514). Horseradish peroxidase (HRP)-conjugated anti-rabbit or anti-mouse secondary antibodies were used to assess protein signal (Thermo Scientific, Cell Signaling Technologies). Chemiluminescence was measured using an Odyssey Fc Imaging System (Licor) and Image Studio Software (LI-COR) was used to quantify chemiluminescence.

Proteomics:

Proteomics sample preparation

Cell pellets for 24 hour TCR WT and Pim dKO CD4 and CD8 T cells were activated and sorted as described above.

IL-2 and IL-15 expanded WT and Pim dKO CD8 T cells on day 6 of culture days, were cultured in fresh warm complete medium and cytokine for 2 hours before cells were harvested, washed twice in ice-cold HBSS and cell pellets snap frozen in liquid nitrogen. Duplicate cell pellets were collected per sample for parallel processing for proteomics or RNAseq analysis.

IL-2 expanded P14 CD8 T cells on day 5 of culture were treated for 24 hours with PIM447 (5 μM) or AZD1208 (10 μM). Cells were harvested on day 6 of culture, washed twice in ice-cold HBSS and cell pellets snap frozen in liquid nitrogen.

For all proteomics samples, cell pellets were lysed in 4% SDS, 50 mM TEAB pH 8.5, 10 mM TCEP (5 min, 1200 rpm, room temperature), boiled (5 min, 500 rpm, 95 °C), then sonicated with a BioRuptor (30 s on, 30 s off x30 cycles). Protein concentration was determined using EZQ protein quantitation kit (Invitrogen) as per manufacturer instructions. Lysates were alkylated with 20 mM iodoacetamide (Sigma) for 1 hr at room temperature in the dark.

24 hour TCR WT and Pim dKO CD4 and CD8 T cell proteomics samples and Day 6, 24 hour

pan-PIM kinase inhibitor treated IL2 expanded CD8 T cell samples were cleaned up and

peptides generated by an SP3 bead protocol(Hughes et al., 2014) as described

previously(Marchingo et al., 2020). As outlined in reference(Marchingo et al., 2020):

"Briefly, 200 μg of 1:1 mixed Hydrophobic and Hydrophilic Sera-Mag SpeedBead

Carboxylate-Modified Magnetic Particles were added per protein sample then acidified

to ~pH 2.0 by addition 10:1 Acetonitrile: Formic Acid. Beads were immobilised on a

magnetic rack and proteins washed with 2 × 70% ethanol and 1 × 100% acetonitrile. Rinsed

beads were reconstituted in 0.1% SDS 50 mM TEAB pH 8.5, 1 mM CaCl2 and digested

overnight with LysC followed by overnight digestion with Trypsin, each at a 1:50 enzyme to

protein ratio. Peptide clean up was performed as per SP3 procedure(Hughes et al., 2014).

Briefly, protein-bead mixtures were resuspended and 100% acetonitrile added for 10 min

(for the last 2 min of this beads were immobilised on a magnetic rack). Acetonitrile and

digest buffer were removed, peptides were washed with acetonitrile and eluted in 2%

DMSO. Peptide concentration was quantified using CBQCA protein quantitation kit

(Invitrogen) as per manufacturer protocol. Formic acid was added to 5% final

concentration".

24 hour TCR WT and Pim dKO CD4 and CD8 T cell proteomics samples were fractionated

using high pH reverse phase liquid chromatography as described previously(Marchingo et

al., 2020). As outlined in reference(Marchingo et al., 2020): "Samples were loaded onto a

2.1 mm x 150 mm XBridge Peptide BEH C18 column with 3.5 μm particles (Waters). Using a

Dionex Ultimate3000 system, the samples were separated using a 25 min multistep gradient

of solvents A (10 mM formate at pH 9 in 2% acetonitrile) and B (10 mM ammonium formate

pH 9 in 80% acetonitrile), at a flow rate of 0.3 mL/min. Peptides were separated into 16

fractions which were consolidated into eight fractions. Fractionated peptides were dried in

vacuo then dissolved in 5% Formic Acid for analysis by LC-ES-MS/MS".

24 hour pan-PIM kinase inhibitor treated day 6 IL2 expanded CD8 T cell samples, were not

fractionated. They were dried in vacuo then dissolved in 5% Formic Acid for analysis by LC-

ES-MS/MS after SP3 cleanup.

Day 6 IL-2 and IL-15 expanded WT and PimdKO CD8 T cells samples were cleaned up and

peptides generated by an S-Trap protocol (Zougman et al., 2014) as per manufacturer

protocol (Protifi). Briefly, add phosphoric acid (to final concentration 1.2%) then S-Trap

binding buffer (90% methanol, 100 mM TEAB, pH 7.1) at a ratio 1:7 v:v lysate:binding buffer

was added to sample lysates containing 100-300 µg total protein. Acidified lysates were

loaded to S-Trap mini columns and spun (4,000 g, 30 s) until all SDS lysate/S-Trap buffer had

passed through the column. Proteins were digested on column in digest buffer (50 mM

sodium bicarbonate) containing Trypsin at a 1:20 enzyme to protein ratio (2 hrs, 47°C

without shaking). Peptides were eluted by spinning in digest buffer (1000g, 1 min), then

0.2% aqueous formic acid (1000g, 1 min), then 50% acetonitrile 0.2% formic acid (4000g, 1

min). Peptide concentration was quantified using CBQCA protein quantitation kit

(Invitrogen) as per manufacturer protocol. Peptides were dried in vacuo then dissolved in

5% Formic Acid for analysis by LC-ES-MS/MS.

Liquid chromatography electrospray tandem mass spectrometry analysis (LC-ES-MS/MS)

≤1 µg of peptide was analysed per fraction by Data-Dependent Acquisition (DDA) Mass

Spectrometry for 24 hour TCR WT and Pim dKO CD4 and CD8 T cell proteomics samples.

For label-free DDA proteomics of WT and Pim dKO CD4 and CD8 24 hour TCR activated T

cells samples were analysed as described previously(Sinclair et al., 2019). As outlined in

reference(Sinclair et al., 2019): "samples were injected onto a nanoscale C18 reverse-phase

chromatography system (UltiMate 3000 RSLC nano, Thermo Scientific) then electrosprayed

into an Orbitrap mass spectrometer (LTQ Orbitrap Velos Pro; Thermo Scientific). For

chromatography buffers were as follows: HPLC buffer A (0.1% formic acid), HPLC buffer B

(80% acetonitrile and 0.08% formic acid) and HPLC buffer C (0.1% formic acid). Peptides

were loaded onto an Acclaim PepMap100 nanoViper C18 trap column (100 µm inner

diameter, 2 cm; Thermo Scientific) in HPLC buffer C with a constant flow of 10 µl/min. After

trap enrichment, peptides were eluted onto an EASY-Spray PepMap RSLC nanoViper, C18, 2

µm, 100 Å column (75 µm, 50 cm; Thermo Scientific) using the buffer gradient: 2% B (0 to 6

min), 2% to 35% B (6 to 130 min), 35% to 98% B (130 to 132 min), 98% B (132 to 152 min),

98% to 2% B (152 to 153 min), and equilibrated in 2% B (153 to 170 min) at a flow rate of 0.3

µl/min. The eluting peptide solution was automatically electrosprayed using an EASY-Spray

nanoelectrospray ion source at 50° and a source voltage of 1.9 kV (Thermo Scientific) into

the Orbitrap mass spectrometer (LTQ Orbitrap Velos Pro; Thermo Scientific). The mass spectrometer was operated in positive ion mode. Full-scan MS survey spectra (mass/charge ratio, 335 to 1800) in profile mode were acquired in the Orbitrap with a resolution of 60,000. Data were collected using data- dependent acquisition: the 15 most intense peptide ions from the preview scan in the Orbitrap were fragmented by collision-induced dissociation (normalized collision energy, 35%; activation Q, 0.250; activation time, 10 ms) in the LTQ after the accumulation of 5000 ions. Precursor ion charge state screening was enabled, and all unassigned charge states as well as singly charged species were rejected. The lock mass option was enabled for survey scans to improve mass accuracy. (Using Lock Mass of 445.120024)".

~0.2 μg of peptide was analysed per sample by single shot DDA Mass Spectrometry for 24 hour pan-PIM kinase inhibitor treated, day 6 IL2 expanded CD8 T cell samples. Samples were injected onto a nanoscale C18 reverse-phase chromatography system (UltiMate 3000 RSLC nano, Thermo Scientific) then electrosprayed into a Q-Exactive HF (Thermo Scientific). A 2- 35% B gradient comprising of eluent A (0.1% formic acid) and eluent B (80% acetonitrile/0.1% formic acid) was used to run a 120-minute gradient per sample. With the mass spectrometer in positive mode the top 20 most intense peaks from a mass range of 335-1800 m/z in each MS1 scan with a resolution of 60,000 were then taken for MS2 analysis at a resolution of 15,000. Spectra were fragmented using Higher-energy C-trap dissociation (HCD).

2 μg of peptide was analysed per sample by Data-Independent Acquisition (DIA) Mass Spectrometry for the IL-2 and IL-15 expanded CD8 T cell proteomics samples. For label-free DIA proteomics of Day 6 IL-2 and IL-15 expanded WT and Pim dKO CD8 T cells peptide samples were analysed as described in (Sollberger et al., 2023) with some minor differences. Peptides were injected onto a nanoscale C18 reverse-phase chromatography system (UltiMate 3000 RSLC nano, Thermo Scientific) then electrosprayed into an Orbitrap Exploris 480 Mass Spectrometer (Thermo Scientific). For liquid chromatography buffers were as follows: buffer A (0.1% formic acid in Milli-Q water (v/v)) and buffer B (80% acetonitrile and 0.1% formic acid in Milli-Q water (v/v). Sample were loaded at 10 μL/min onto a trap column (100 μm × 2 cm, PepMap nanoViper C18 column, 5 μm, 100 Å, Thermo Scientific) equilibrated in 0.1%

trifluoroacetic acid (TFA). The trap column was washed for 3 min at the same flow rate with 0.1% TFA then switched in-line with a Thermo Scientific, resolving C18 column (75 μm × 50 cm, PepMap RSLC C18 column, 2 μm, 100 Å). The peptides were eluted from the column at a constant flow rate of 300 nl/min with a linear gradient from 3% buffer B to 6% buffer B in 5 min, then from 6% buffer B to 35% buffer B in 115 min, and finally to 80% buffer B within 7 min. The column was then washed with 80% buffer B for 4 min and re-equilibrated in 3% buffer B for 15 min. Two blanks were run between each sample to reduce carry-over. The column was kept at a constant temperature of 40°C.

The data was acquired using an easy spray source operated in positive mode with spray voltage at 2.1 kV and the ion transfer tube temperature at 275°C. The MS was operated in DIA mode. A scan cycle comprised a full MS scan (m/z range from 350-1650, with RF lens at 40%, AGC target set to custom, normalised AGC target at 300%, maximum injection time mode set to custom, a maximum ion injection time of 20 ms, microscan set to 1 and source fragmentation disabled. MS survey scan was followed by MS/MS DIA scan events using the following parameters: multiplex ions set to false, collision energy mode set to stepped, collision energy type set to normalized, HCD collision energies set to 25.5, 27 and 30%, orbitrap resolution 30000, first mass 200, RF lens 40%, AGC target set to custom, normalized AGC target 3000%, microscan set to 1 and maximum injection time 55 ms. Data for both MS scan and MS/MS DIA scan events were acquired in profile mode.

Proteomics data analysis

The 24 hour TCR WT and Pim dKO CD4 and CD8 T cell DDA proteomics data and 24 hour pan-PIM kinase inhibitor treated Day 6 IL2 expanded CD8 T cell DDA proteomics data were processed, searched and quantified with the MaxQuant software package, version 1.6.10.43. For the protein and peptide searches we generated a hybrid database consisting of all manually annotated mouse SwissProt entries, combined with mouse TrEMBL entries with protein level evidence available and a manually annotated homologue within the human SwissProt database as described in (Marchingo et al., 2020). For the 24 hour TCR activated data this was from the Uniprot release 2020_06, for the 24 hour PIM kinase inhibitor data this was from the Uniprot release 2019_08. Search parameters were as described in (Marchingo et al., 2020): "The following MaxQuant search parameters were used: protein N-terminal acetylation, methionine oxidation, glutamine to pyroglutamate,

and glutamine and asparagine deamidation were set as variable modifications and carbamidomethylation of cysteine residues was selected as a fixed modification; Trypsin and LysC were selected as the enzymes with up to two missed cleavages permitted; the protein and PSM false discovery rate was set to 1%; matching of peptides between runs was switched off".

Data filtering and protein copy number quantification was performed in the Perseus software package, version 1.6.6.0. Proteins were quantified from unique peptides and razor peptides (peptides assigned to a group, but not unique to that group). The data set was filtered to remove proteins categorised as 'contaminants', 'reverse' and 'only identified by site'.

To generate a point of reference between TCR activated WT and Pim dKO and inactive T cells, naïve WT CD4 and CD8 T cell data from (Marchingo et al., 2020), which was collected and processed identically, was included in proteomics search, results table and plot of protein content. At this point biological replicate 3 from Pim dKO CD4 T cells was excluded from further analysis due to poor sample quality (e.g. protein sequence coverage was an average of 13.9% compared to an average of 18.2-24.6% for all other samples and estimated total protein content per cell was ~1/3rd of that observed in the other two biological replicates for that condition, despite no obvious difference in FSC measured when FACS sorting).

The day 6 IL-2 and IL-15 expanded WT and Pim dKO CD8 T cell DIA proteomics data files were searched using Spectronaut version 14.7. Data was analysed using a library-free approach. Raw proteomics data files were searched using the Pulsar tool within Spectronaut using the following settings: 0.01 FDR at the protein and peptide level with digest rule set to 'Trypsin'P'. A maximum of two missed cleavages and minimum peptide length of 7 amino acids was selected. Carbamidomethyl of cysteine was selected as a fixed modification while protein n-terminal acetylation and methionine oxidation were selected as variable modifications. Manufacturer default settings were used for identification, with protein and precursor q-value set to 0.01 using default decoy strategy. For quantification settings, the MS-Level quantity was set to 'MS2', major group Top N and minor group Top N were set as 'False' and profiling was set as 'False'. The data was searched against the same mouse Uniprot database as used above for DDA analysis.

Data was filtered to only include proteins for which at least one condition had peptides

detected in ≥2 biological replicates for 24 hour TCR and day 6 IL2 and IL15 WT and Pim dKO

data. Data was filtered to only include proteins for which at least two conditions had

peptides detected in ≥2 biological replicates for PIM kinase inhibitor data. Mean copy

number per cell was calculated using the "proteomic ruler' plugin as described in

(Wisniewski et al., 2014).

Mass contribution of proteins (g/cell) was calculated as (protein copy number) * (molecular

weight (Daltons)) / (Avogadro's constant). Differential expression analysis of protein copy

number was performed using RStudio (version 1.2.5033). P-values and fold-change were

calculated with the Bioconductor package Limma (version 3.44.3). Q-values were calculated

using Bioconductor package qvalue (version 2.20.0).

All scripts used to perform analysis are available from authors upon request.

RNAseq:

RNA extraction and library preparation and RNA sequencing:

Biological triplicate cell pellets of day 6 IL-2 or IL-15 differentiated CD8 T cells were collected

in parallel with proteomics samples as described above. Total RNA was extracted from cell

pellets with Qiagen RNAeasy mini kit (Qiagen) as per manufacturer instructions.

2 µg of RNA was submitted per sample to the Finnish Functional Genomics Centre, University

of Turku and Åbo Akademi University for library preparation and sequencing. RNA quality was

verified using Agilent Bioanalyzer 2100 or Advanced Analytical Fragment Analyzer. 300 ng of

total RNA was used for library preparation using the Illumina TruSeq Stranded mRNA sample

preparation kit (15031047) as per manufacturer instructions. Library quality was confirmed

with Advanced Analytical Fragment Analyzer and quantified with Qubit Fluorometric

Quantitation (Life Technologies). Sample libraries were pooled and run in a single lane on an

Illumina NovaSeq 6000 with read length 2x 50 bp. Base calling was performed with bcl2fastq2

conversion software (NovaSeq 6000) and automatic adapter trimming performed for fastq

raw read files generated.

1005

1006

RNAseq data analysis

Quality assessment of raw reads was evaluated using MultiQC (Ewels et al., 2016). Raw reads were aligned to the GRCm38.p6 (Ensembl) build of the Mus musculus reference genome using the STAR aligner (Dobin et al., 2013) (version 2.7.1a), with 'outFilterType BySJout' option switched on to reduce spurious junctions. The STAR quantMode GeneCounts option was used to map aligned reads to genes to generate a gene counts matrix, using the annotation file GRCm38.101 (Ensembl).

Bioconductor package EdgeR (version 3.30.3) (Robinson et al., 2010; McCarthy et al., 2012) was used to reduce noise from low count genes with filterByExpr function and the resulting count matrix was normalised by the EdgeR Trimmed Mean of M-values (TMM) method. Differential expression analysis was performed using Limma package (version 3.44.3) (Ritchie et al., 2015; Law et al., 2016). Reads were converted to $\log_2$ counts per million and weighted using the voomWithQualityWeights function. A linear model was fitted to each gene and empirical Bayes moderated t-statistics were applied. q-values were calculated from using Bioconductor package qvalue (version 2.20.0) (Storey et al., 2023). The calculateTPM fuction from the Bioconductor package Scater (version 1.16.2) (McCarthy et al., 2017) was used to calculate TPM values for visualisation purposes, with total exon length calculated using GenomicRanges (version 1.40.0) (Lawrence et al., 2013) as the sum of the non-overlapping exonic regions for a gene.

Combining Proteomics and RNAseq data

Proteomics data was annotated during Spectronaut search and Perseus analysis with Uniprot ID, MGI symbol and MGI ID information. BioMart (version 2.47.2) was used to extract UniProt identifiers, MGI symbols and MGI IDs for corresponding ENMUSG IDs of RNAseq data. RNAseq data was first combined with proteomics data based on UniProt ID. Proteins that didn't receive an initial match, were then combined by matching MGI ID values, then if still unmatched by MGI symbol. Data combinations were manually checked for obvious mis-annotations. Proteins with no corresponding mRNA identified or vice versa were excluded from the combined data file. In a few instances multiple proteins aligned to a single mRNA measurement or multiple mRNA aligned to a single protein measurement. For the purposes

1038   of this broad examination of how differences in expression at the protein or mRNA level these

1039   results were also excluded from the combined data table and subsequent analysis.

1040

1041   **Graphing and statistics**

1042   Heatmaps were generated using Broad Institute software Morpheus

1043   (https://software.broadinstitute.org/morpheus). Statistical tests performed for experiments

1044   other than proteomics and RNAseq are indicated in figure legends and were performed in

1045   Prism (GraphPad, version 9 or 10).

1046

Data availability

All data generated or analysed during this study are included in the manuscript and

supporting files. Raw mass spec data files and MaxQuant or Spectronaut analysis files are

available on the ProteomeXchange data repository and can be accessed with the following

identifiers: PXD051198 for 24 hour TCR activated WT and Pim dKO CD4 and CD8 T cells;

Naïve T cell data used as a point of reference for plotting 24 hour TCR proteomics has

previously been published as part of (Marchingo et al., 2020) and can also be found under

the identifier PXD016105; PXD051214 for Pim inhibitor proteomics; PXD051213 for day 6 IL-

2 and IL-15 expanded WT and Pim dKO CD8 T cells. Raw RNAseq data files are available on

the NCBI GEO website and can be accessed with the identifier GSE261667.

Figure 1. Pim1 and Pim2 are strongly TCR-induced, but dispensable for T cell activation
Estimated copies per cell of PIM1 and PIM2 protein from quantitative proteomics analysis of (**A**) OT1 CD8 T cells stimulated with SIINFEKL peptide for indicated times from published dataset (Marchingo et al., 2020) or (**B**) naïve ex vivo and 24 hour αCD3/αCD28 (TCR) activated WT CD8 T cells (see Fig 1G-H) for further details. (**C**) Fragments per kilobase million (FPKM) of *Pim1*, *Pim2* and *Pim3* mRNA from published (Spinelli et al., 2021) bulk RNAseq analysis of naïve and 24 hour gp33-41 peptide stimulated P14 CD8 T cells. Lymph node cell suspensions from C57BL/6 (WT) and *Pim1*KO/*Pim2*KO (Pim dKO) mice were activated for 24 hours with αCD3/αCD28 (both 0.5 μg/mL) and CD4 and CD8 T cell (**D**) FSC-A SSC-A profiles, (**E**) expression of surface activation markers (CD25, CD44, CD71) or CD8 T cell intracellular IFNγ were measured by flow cytometry. (**F**) Lymph node single cell suspensions from WT and Pim dKO mice were labelled with CellTrace Violet (CTV), activated with αCD3/αCD28 (both 0.5 μg/mL) and CD4 and CD8 T cell CTV proliferation profiles were measured at indicated time points. (G-H) Lymph node cell suspensions from WT and Pim dKO mice were stimulated for 24 hours with αCD3/αCD28 (both 0.5 μg/mL) and activated CD4 and CD8 T cells were sorted for analysis by quantitative proteomics. Data was analysed using proteomic ruler method (Wisniewski et al., 2014) to estimate protein copy number per cell. (**G**) Total protein content (μg/million cells) (one-way ANOVA), (**H**) Volcano plots of p-value (-log$_{10}$) versus fold-change (log$_2$) in protein copy number between Pim dKO and WT. Horizontal dotted line represents multi-test correction cut-off of q=0.05, vertical dotted line shows 1.5-fold change. Lymph node single cell suspensions from WT and Pim dKO mice were labelled with CellTrace Violet (CTV) and (**I**) cells were cultured in IL-7 (5ng/mL) +/- rapamycin (20 nM) and CD8 T cell numbers measured over time or (**J**) cells were activated with αCD3/αCD28 (both 0.5 μg/mL) +/- rapamycin (20 nM) and CD8 T cell mean division number was calculated over time (two-way ANOVA). Symbols in bar charts represent biological replicates, symbols in (I) represent the mean. Error bars show mean ± S.D. Flow cytometry dot plots and histograms are representative of (D-E) n = 3, except for IFNγ staining which is n=2 or show pooled data from (I) n=3-4 and (F, J) n=5 biological replicates, with data collected over at least 2 independent experiments. Quantitative proteomics was performed on biological triplicates.

Figure 2. Pim1/Pim2 deficiency reduces IL-15-driven CD8 T cell proliferation but does not prevent memory differentiation.

(**A**) OT1 lymph node cell suspensions were SIINFEKL peptide activated for 36 hours, washed then cultured with no cytokine, IL-15 (20 ng/mL) or IL-2 (20 ng/mL) for 4 or 24 hours. Western blots of PIM1, PIM2 or pSTAT5 Y694 expression. (**B**) Schematic of cytokine driven memory and effector CD8 T cell expansion and differentiation cultures. Lymph node or spleen cell suspensions were activated for 2 days with TCR stimulus + cytokine, washed, then split daily into fresh media + cytokine. (**C**) WT (Ly5.1) and Pim dKO LN suspensions were mixed at a 50:50 ratio for T cells and cultured as outlined in (B) with TCR stimulus αCD3/αCD28 (both 0.5 μg/mL) + cytokine IL-15 (20 ng/mL), and CD8 T cell number was measured daily. (**D**) WT and Pim dKO T cells were expanded with IL-15 in separate cultures as per (B-C) and % live cells (PI-ve) were assessed on days 4 and 6 (two-way ANOVA). (E-J) WT and Pim dKO CD8 T cells were activated with TCR stimulus αCD3/αCD28 (both 0.5 μg/mL) + cytokine IL-15 (20 ng/mL), expanded with IL-15 as per (B), with an additional CD4 T cell magnetic depletion step on day 3 of culture. CD8 T cells were harvested on day 6 for parallel RNAseq and proteomic analysis. (**E**) Fold-change in mRNA expression between Pim dKO and WT versus average mRNA expression (TPM). mRNA expression (Transcripts per million, TPM) of (**F**) secondary lymphoid homing receptors Sell, Ccr7, S1pr1 and (**G**) key transcription factors involved in CD8 T cell memory differentiation and maintenance Tcf7, Klf2, Foxo1, Foxo3, Id3. (**H**) WT vs Pim dKO protein copy numbers, differentially expression proteins (FC>1.5, q<0.05) are highlighted in red (**I**) Protein copy numbers per cell for key mitochondrial proteins DRP1, OPA1 and CPT1A. (**J**) WT vs Pim dKO protein copy numbers, mitochondrial proteins (as defined in MitoCarta 3.0) are highlight in pink. Symbols in bar charts represent biological replicates, symbols in (C,H,J) represent the mean. Error bars show mean ± S.D. Data are representative of (A) n=3 or show pooled data from (C) n = 4, and (D) n=5 biological replicates with data collected over at least 2 independent experiments. Quantitative proteomics and RNAseq was performed on biological triplicates. ** q≤0.01, fold-change (FC) shown on bar graphs when q<0.05.

Figure 3. Pim dKO IL-2 differentiated effector T cells have reduced cell size and sustained expression of CD62L

(**A**) Estimated copies per cell of PIM1 and PIM2 protein from published quantitative proteomics analysis (Howden et al., 2019; Brenes et al., 2023) of CD8 T cells expanded in IL-2 or IL-15 as outlined in Fig 2B. (B-D, F-G) WT (Ly5.1) and Pim dKO lymph node or spleen single cell suspensions were mixed at a 50:50 ratio of T cells, activated for 2 days with αCD3/αCD28 (both 0.5 μg/mL) and IL-2 (20 ng/mL), washed then split into fresh medium containing IL-2 (20 ng/mL) daily (as per Fig 2B). Some of the mixed cell suspensions were also cultured in IL-7 (5 ng/mL) to sustain a naïve T cell reference. (**B**) WT and Pim dKO CTL were treated 1 hour +/- Jak1/3 inhibitor Tofacitinib (100 nM) (negative control) before pSTAT5 Y694 expression was measured on day 3 and 6 of culture, (**C**) surface CD25 expression was measured on days 3 and 6 or culture, (**D**) CD8 T cell number vs time was calculated, (**F**) CD8 T cell FSC-A, SSC-A and surface activation markers (CD44, CD71) were measured on days 3 and 6 of culture (**G**) expression of adhesion molecule CD62L was measured daily. (**E**) WT and Pim dKO T cells were activated and expanded with IL-2 as per (Fig 3B-D, F-G) except in separate cultures and % live cells (PI-ve) was assessed on days 4 and 6 (two-way ANOVA).

Symbols in bar charts represent biological replicates, symbols in (D) represent the mean. Error bars show mean ± S.D. Data are representative of (B, G) n=4, (C,F) n=6 or show pooled data from (D) n=4, (E) n=6 biological replicates with data collected over at least 2 independent experiments

Figure 4. Major glucose transporters and effector proteins are reduced in Pim dKO IL-2 expanded CTL

WT and Pim dKO CD8 T cells were activated for 2 days with αCD3/αCD28 (both 0.5 μg/mL) and IL-2 (20 ng/mL), washed then split into fresh medium containing IL-2 (20 ng/mL) daily (as per Fig 2B), with an additional CD4 T cell magnetic depletion step on day 3 of culture. CD8 T cells were harvested on day 6 of culture for high resolution mass spectrometry. (**A**) Estimated total protein content per cell (student t-test). (**B**) Volcano plots Pim dKO vs WT protein copy numbers, differentially expressed proteins (FC>1.5, q<0.05) are highlighted in red. Estimated protein copy number per cell of (**C**) transcription factor TBX21 and TCF1 (**D**) glucose transporters SLC2A1 and SLC2A3. (**E**) Volcano plots Pim dKO vs WT protein copy numbers. Proteins with KEGG term = "terpenoid backbone biosynthesis", "biosynthesis of unsaturated fatty acids" or "steroid biosynthesis" are highlighted with proteins with FC>1.5, q<0.05 shown in red and proteins with FC<1.5 and/or q>0.05 shown in pink. (**F**) Heatmap of protein copy numbers for granzymes, perforin and effector cytokines. (**G**) Granzyme B and IFNγ expression was measured by flow cytometry in day 6 IL-2 expanded WT and Pim dKO CTL.

Symbols in bar charts show biological replicates. Error bars show mean ± S.D. Data are representative of (G) n = 3-4, with data collected over at least 2 independent experiment Quantitative proteomics was performed on biological triplicates. * indicates q<0.05, fold-change (FC) shown on graph when q<0.05

Figure 5. Disconnect between protein and mRNA expression in Pim1/Pim2-deficient effector CTL corresponds with a reduction in protein synthesis

RNAseq analysis was performed in day 6 IL-2 expanded WT and Pim dKO CD8 T cells which were collected in parallel with proteomics analysis described in Fig 4A. (**A**) Volcano plot of RNAseq data, differentially expressed mRNA (FC>1.5, q<0.05) are highlighted in red. (**B**) Volcano plot of RNAseq data, Granzymes C-K, perforin, Pdcd4 and Sell are highlighted in red. (**C**) Heatmap of mRNA expression (TPM) for granzymes, perforin and effector cytokines. Bar chart of mRNA expression (TPM) of (**D**) Granzymes A and B (**E**) Glucose transporters Slc2a1 and Slc2a3. (F-G) Fold change of PimdKO/WT protein (copies) from proteomics analysis described in Fig 4 vs mRNA (TPM) (**F**) highlighting in red proteins that are differentially expressed (FC>1.5, q<0.05) where mRNA is not substantially different (FC<1.2) and (**G**) highlighting in red protein and mRNA that are both differentially expressed (FC>1.5, q<0.05). (**H**) Estimated cytosolic ribosome content per cell (left), % ribosome of total cellular protein content (right). (**I**) Estimated protein copy number per cell of translation repressor PDCD4. (**J**) PDCD4 expression measured by flow cytometry on day 3 and 6 in IL-2 expanded WT vs Pim dKO CD8 T cells. (**K**) Estimated protein copy number per cell of EIF4A1. (**L**) adjusted ratio of PDCD4 : EIF4A1 (assuming 1 PDCD4 binds 2x EIF4A1) in WT and Pim dKO proteomes. (**M**) Protein synthesis measured by OPP incorporation in day 6 IL-2-expanded WT CTL treated for 24 hours with pan PIM kinase inhibitors PIM447 (5 µM) or AZD1208 (10 µM). 30 min cycloheximide (100 µg/mL) treatment gives no protein synthesis background control.

Symbols in bar charts show biological replicates: error bars show mean ± S.D. Data are representative of (M) n=2 biological replicates collected over two independent experiments, (J) n = 2 biological replicates. Quantitative proteomics and RNAseq were performed on biological triplicates. * indicates q<0.05, fold-change (FC) shown on graph when q<0.05.

Figure 6. PIM kinases regulate mTORc1 activity and lymphoid homing in effector CTL
(**A**) Western blot of pS6K T389 and pan S6K from day 6 WT and PimdKO IL-2 CTL (paired student t-test, *= p<0.05). (**B**) WT and Pim dKO T cells were mixed in a 50:50 ratio, activated with αCD3/αCD28 (both 0.5 μg/mL) and IL-2 (20 ng/mL) and expanded in IL-2 as per Fig 2B and pS6 (Ser235/236) measured after 1 hr +/- PdBU and ionomycin (positive control), +/- rapamycin (mTORc1 inhibitor, negative control) or no additional treatment. mRNA expression (TPM) from RNAseq analysis of IL-2 expanded WT and Pim dKO CTL described in Fig 5A for (**C**) cell homing receptors S1pr1, Ccr7 and Sell and (**D**) transcription factor Klf2. (**E**) WT and Pim dKO T cells were activated and expanded with IL-2 in separate cultures as per Fig 2B. On day 6 of culture WT and Pim dKO CTL were labelled with CFSE or CTV, mixed at a 50:50 ratio and transferred into C57Bl/6 recipient mice. Values indicate percentage of transferred cells detected in blood, lymph node or spleen 4 or 23 hours post-transfer that were WT or Pim dKO (one-way ANOVA).
Symbols show biological replicates. Error bars show mean ± S.D. Data are representative of (A) n= 6, (B) n=2-4 collected across at least 2 independent experiments and (E) n= 6 recipient mice, from n=2 biological donor replicates. RNAseq was performed on biological triplicates. * q<0.05, ** q<0.01, *** q<0.001, fold-change (FC) shown on graph when q<0.05 but FC<1.5.

Figure S1. T cell counts in WT vs Pim dKO spleen

Comparison of Pim dKO and age/sex matched WT control mice for (**A**) total number of CD4 and CD8 T cells in spleens (one-way ANOVA), (**B**) total number of splenocytes (student t-test), (**C**) proportion of live splenocytes that were CD4 and CD8 (one-way ANOVA). Symbols in bar charts represent biological replicates: error bars show mean ± S.D. (A-C) n = 8 collected over 4 independent experiments. ** q<0.01, ***q<0.001.

Figure S2. Pim dKO IL2 expanded CD8 T cells exhibit an effector-like mitochondrial proteome profile

Heatmap showing protein copy number of mitochondrial proteins (as defined by inclusion on Mitocarta 3.0 list) from IL-2 expanded WT (effector), Pim dKO and IL-15 expanded WT (memory) proteomics experiments described in Fig 2 and 4. Proteins are ranked based on average expression in IL2 WT condition. Proteomics was performed on biological triplicates.

Figure S3. 24 hour treatment of IL-2 CTL with pan-PIM kinase inhibitors PIM447 or AZD1208 recapitulates many features of Pim1/Pim2-deficiency

Single cell suspension from P14 TCR-transgenic mouse lymph nodes were activated with gp33 peptide (100 ng/mL), IL-2 (20 ng/mL) and IL-12 (2 ng/mL) for 2 days, then split daily into fresh media containing IL-2 (20 ng/mL). On day 5 of culture IL-2 expanded CTL were treated with pan-PIM kinase inhibitors PIM447 (5 μM) or AZD1208 (10 μM) or DMSO vehicle control for 24 hours harvested on day 6 of culture to measure (**A**) Cell number (**B**) FSC-A SSC-A. Proteome analysis was also performed on inhibitor treated CTL to measure (**C**) protein content, (**D**) protein expression of SCD1-3, SLC2A1, SLC2A3, GZMB or (**F**) PDCD4. (**E**) Day 6 IL-2 expanded CTL from WT (C57BL/6) mice were treated for 24 hours with PIM447 or AZD1208 (both 1 μM) and GZMB expression was measured by flow cytometry.

Symbols show biological replicates. Error bars show mean ± S.D. Data are representative of (E) n=2 biological replicates collected across at least 2 independent experiments. Proteomics analysis was performed on biological triplicates, with (A-B) collected in parallel with proteomics analysis. * q<0.05.