Chilean brush tailed mouse (*Octodon degus*): a diurnal precocial rodent as a new model to study visual receptive field properties of superior colliculus neurons

N.I. Márquez[1]* [https://orcid.org/0000-0002-4200-6211], P.F. Fernández-Aburto[2] [https://orcid.org/0000-0002-5695-6234] , A. Deichler[1] [https://orcid.org/0000-0002-2048-8539], I. Perales[1] [https://orcid.org/0009-0006-0394-3049], J.-C. Letelier[1] [https://orcid.org/0000-0001-9484-026X], G. J. Marín[1,3] [https://orcid.org/0000-0003-2961-6469], J. Mpodozis[1†] [https://orcid.org/0000-0003-0786-5579], S.L. Pallas[4†*] [https://orcid.org/0000-0002-1760-7437]

[1]Depto. Biol., Univ. de Chile, Santiago, Chile; [2] Instituto de Ecología y Biodiversidad (IEB), Las Palmeras 3425, Ñuñoa, Santiago, Chile; [3]Facultad de Medicina, Universidad Finis Terrae, Santiago, Chile; [4]Dept. of Biol., Univ. of Massachusetts-Amherst, Amherst, MA; [†]These authors share senior authorship.

Abbreviated title: **Octodon degus superior colliculus receptive field properties**

*Correspondence should be addressed to:
S.L. Pallas
Department of Biology, University of Massachusetts Amherst
221 Morrill Science Center III
611 North Pleasant Street
Amherst, MA 01003 USA
spallas@umass.edu

N.I. Márquez
Departamento de Biología, Facultad de Ciencias, Universidad de Chile
Las Palmeras 3425, Ñuñoa, Santiago de Chile
CHILE
imarquez@uchile.cl

Number of figures: **7**
Number of tables: **1**

Acknowledgements: We would like to thank Solano Henríquez for his valuable assistance.

Conflict of interest: Authors report no conflict of interest.

Funding sources: Support for this work was provided by a National Science Foundation grant (IOS-1656838, IOS-2029980), a DARPA grant (HR0011-18-2-0019, TA2), and UMass startup funds awarded to S.L.P., by a Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT) grant to J.M. (FONDECYT 1210069), by a FONDECYT grant to G.M. (FONDECYT 1210169), and by a postdoctoral FONDECYT grant to A.D (FONDECYT 3220871).

ABSTRACT

Lab rodent species used to study the visual system and its development (hamsters, rats, and mice) are nocturnal, altricial, and possess simpler visual systems than carnivores and primates. To widen the spectra of studied species, here we introduce an alternative model, the Chilean degu (*Octodon degus*), a diurnal, precocial Caviomorph rodent with a cone enriched, well-structured retina, and well-developed central visual projections. To assess degu visual physiological properties, we characterized the visual responses and receptive field (RF) properties of isolated neurons in the superficial layers of the superior colliculus (sSC). To facilitate comparison with studies in other rodent species, we used four types of stimuli: (1) a moving white square, (2) sinusoidal gratings, (3) an expanding black circle (looming), and (4) a stationary black circle. We found that as in other mammalian species, RF size increases from superficial to deeper SC layers. Interestingly, compared to other lab rodents, degus have smaller RF sizes, likely indicating higher acuity. sSC neurons displayed spatial frequency tuning to grating stimuli from 0.08 to 0.24 cycles/degree. Additionally, neurons from sSC showed transient ON, OFF, or ON-OFF responses to stationary stimuli but increased their firing rates as a looming object increased in size. Our results suggests that degus have higher visual acuity, higher frequency tuning, and lower contrast sensitivity than commonly used nocturnal lab rodents, positioning degus as a well-suited model for studies of diurnal vision that are more relevant to humans.

INTRODUCTION

In studies of visual processing circuitry, the choice of animal model is often based on tradition or convenience. In contrast, animals that have been chosen for auditory processing research are species that have exceptional auditory processing ability. It could easily be argued that more has been learned about auditory signal processing from studies of bats (Razak 2011; Razak and Fuzessery 2015), owls (DeBello and Knudsen 2004), songbirds (Brainard and Doupe 2013), and chinchillas (Jones *et al.* 2015), than from traditional lab animals. The foundational studies of central visual processing were done largely in cats and monkeys (Hubel and Wiesel 1998), but mice are now the model of choice. Although mouse visual system is similar to that of other mammals in many ways (Huberman and Niell 2011; Garrett *et al.* 2014; Juavinett and Callaway 2015), there are important differences, possibly related to their nocturnal habit and poor eyesight, that have unknown consequences for reaching a general understanding of visual system development and function.

In the Pallas lab, the superior colliculus (SC) and visual cortex (V1) of Syrian hamsters were chosen for studies of visual circuit development, largely because hamsters are born earlier and in larger litters than more typically chosen rodent subjects. As in other mammals studied, receptive fields (RFs) in hamster SC and V1 are large at birth and undergo a postnatal refinement process to reach adult size. It had been assumed, based largely on studies of primary visual cortex (V1) in cats and monkeys (Hensch *et al.* 1998; Espinosa and Stryker 2012), that visual pathway development requires early visual experience during a critical period for maturation but not for maintenance of refined receptive fields and response properties. In contrast, the Pallas lab has found that

developmental refinement of receptive fields in SC and V1 occurs normally in dark reared hamsters. Continued visual deprivation in adulthood results in a gradual loss of RF refinement in both SC and V1 of adult hamsters (> postnatal day (P) 60) (Carrasco *et al.* 2005; Balmer and Pallas 2015). A brief, late juvenile exposure to light stabilizes receptive field size permanently despite continued visual deprivation, but visual experience after sexual maturity (P 60) has no effect (Carrasco and Pallas 2006; Balmer and Pallas 2015). These results raise the question of whether hamsters are unique or if other rodents, in particular those that are diurnal, would exhibit similar independence from visual stimulation during development.

Here we characterize the RF properties of SC neurons in Chilean degus (*Octodon degus*), a diurnal, precocial rodent species (Fulk 1976; Reynolds and Wright 1979; Long and Ebensperger 2010) with a cone enriched retina and a well-developed visual system (Vega-Zuniga *et al.* 2013). Unlike typical lab rodent species, degus open their eyes at P1 and exhibit visually guided behaviors shortly after birth, exposing their visual system to an early visual experience that could shape its maturation. Our results show that degus have clear diurnal characteristics and a better visual acuity in comparison to nocturnal lab rodents. Here we argue that degus are a promising model to study the role of visual experience in the maturation of the visual system and thus might be better suited to model the human visual system.

METHODS

Animals

A total of 10 animals from our colony were used in this study. Animals were housed in

metal cages (50 x 40 x 35 cm) with wood shavings as previously described by Márquez

et al. (2015) in a climate-controlled room under a 12:12 photoperiod in groups of 3-4

degus of the same sex. They were fed with rodent-chow Prolab® RMH 3000 (Purina

LabDiet®, Inc.), Mazuri chinchilla diet (Mazuri® Exotic Animal Nutrition) and provided with

water *ad libitum*. All animal procedures were approved by the *Comité Institucional de*

Cuidado y Uso Animal (CICUA) of the University of Chile and the IACUC of the University

of Massachusetts-Amherst and followed the National Institutes of Health (NIH) Guide for

the Care and Use of Laboratory Animals. All procedures met additional standards of care

established by SfN and IBRO.

Histology and neuronal tracers

Some degus (n=2) were used only for neuroanatomical data collection. The labeling of

retinal projections to the SC was performed under gaseous anesthesia using isoflurane

(4% in medical oxygen at a flow rate of 100 cc/kg/min) complemented with diazepam (5

mg/kg) administrated IP. A glass pipette (tip diameter 15–20 µm) containing 1% cholera

toxin subunit B (CTB, List Biological Laboratories, Campbell, CA) in 0.1 M phosphate

buffer was injected intraocularly. After 5–7 days survival, the degus were sedated with

gaseous isoflurane, placed in a deep state of anesthesia with ketamine/xylazine

(ketamine 40-80 mg/kg, xylazine 5-10 mg/kg), and perfused intracardially by passing 200

ml of warm saline solution (0.9% in 0.1 M PB) followed by 200 ml of cold

paraformaldehyde (PFA, 4% in 0.1 M PBS). The brains were extracted and postfixed

overnight in 4% PFA at 4°C, and then transferred to 30% sucrose solution in 0.1 M PB

until they sank, for cryoprotection. Brains were sectioned (60 μm), and a CTB

immunoreaction assay was performed as described by Deichler *et al.* (2019).

For histological confirmation of electrode positioning during electrophysiological

recording degus were given an overdose of urethane (3 mg/kg, IP) and perfused

intracardially as described above. The brains were postfixed overnight in 4% PFA at 4°C

and then transferred to 30% sucrose for cryoprotection. Finally, brains were sectioned

(60 μm), mounted, and Nissl-stained (cresyl violet, Merck, Darmstadt, Germany).

Reconstruction of electrode penetrations in several experiments verified that our

recordings were taken at intervals of 100 microns through a transect passing across

layers of the SC, from the SGS into the SGI.

In vivo recordings

Animal Preparation. Pre-op: Degus store wood shavings in their cheeks (Deichler,

pers. comm.). Thus, to avoid aspiration of the shavings during anesthesia, they were

taken from their cages and placed in a clean cage with paper towels the night before the

experiment. Degus (n = 8) were fasted overnight after receiving Dexamethasone (1

mg/kg, SQ) to prevent cerebral edema. The morning of surgery, a second dose of

Dexamethasone (1 mg/kg, SQ) was given along with atropine (0.1 mg/kg, SQ) to

counteract bradycardia and apnea and to reduce mucosal secretions.

Surgery. Isoflurane (4% in medical oxygen at a flow rate of 100 cc/kg/min) was used

for anesthesia induction, then diazepam (5 mg/kg) was administrated IP to induce

sedation before degus were anesthetized using urethane (1.25 mg/kg, IP). If necessary,

Doxapram was used (2 mg/kg SQ or sublingual) to stimulate respiration. Body

temperature was kept between 36 °C and 37 °C throughout the experiment by using a

thermo-regulated blanket (FHC, Inc.). To prevent asphyxia due to accumulation of

bronchial secretions, degus were tracheostomized after the site was prepped by shaving

the neck and chest, cleaning the area with ethanol and povidone, and inserting a tracheal

tube. The scalp was shaved, and the head stabilized in a stereotaxic head holder. Then,

the surgical site was cleaned with ethanol and povidone followed by a midline incision in

the scalp. The muscles and periosteum overlying the skull were retracted, and a small

craniotomy above the SC region was made with a surgical micro-drill. After unilaterally

aspirating the cortex to expose the SC and covering the site with saline,

electrophysiological recordings were obtained by using Epoxy-insulated tungsten

microelectrodes (1–2 MOhms; FHC, Inc.) or a 16 Channel multi- electrode array (MEA)

(~1 MOhm; Microprobes for Life Science, Inc.) and a multichannel differential amplifier

(Model 3600; AM Systems Inc.). A silver chloride wire was implanted in between the

exposed bone and muscle as a reference electrode. Physiological data were sampled

just before, during, and just after the stimulation at 10-20 kHz, with a band-pass of 3 Hz

to 10 kHz, by using a standard PC with a 16-channel A/D converter board (National

Instruments Corp). Data acquisition and offline analysis were performed using Igor Pro-8

(WaveMetrics, Inc.) or MATLAB (MathWorks, Inc) software.

Visual stimulation

Visual stimuli were presented on an LCD monitor (Alienware 25 AW2518HF, 1920 x 1080

pixels, 60 Hz refresh rate) placed 32 cm from the degu's eye. Stimulus parameters were

controlled with a lab-designed program using the Python (Python Software Foundation) -

based open-source application PsychoPy (v.3.1.3). A photodiode that detected the

stimulus produced a 5-volt TTL signal that was recorded by the amplifier and used to

synchronize stimuli and neural responses. Four types of visual stimuli were used:

Drifting squares. A white square (~3° diameter) on a dark background was swept from

the top to the bottom of the monitor over 4 s, with 1 s intervals between successive stimuli.

The position of the square changed in an orderly manner along the horizontal axis of the

screen at each sweep, covering the entire screen within 64 runs (**Fig. 1A**). The stimulation

protocol was repeated 3 times for each recording location/depth.

Drifting gratings. Stimuli consisted of full-field, vertical, sinusoidal gratings drifting from

temporal to nasal on the LCD monitor. Frequency, contrast and temporal frequency

parameters were chosen as in Niell and Stryker (2008). The sinusoidal gratings were

presented over 4 s with a 2 s interstimulus interval at a constant temporal frequency (1.5

Hz). All combinations of spatial frequency (0.01, 0.02, 0.04, 0.08, 0.16, 0.32 cycles/deg)

and contrast (0.0625, 0.125, 0.25, 0.5, 0.75, 1) were presented in a pseudo-random

manner. The stimulation protocol was repeated 5 times for each recording location/depth.

Dark looming object. We presented a looming object consisting of a small, black circle

(2° diameter) on a gray background, expanding to 20° in diameter over 250 ms, followed

by a 250 ms presentation at its maximum size (Yilmaz and Meister 2013).

Dark static object. As a control for the looming experiment, we presented a 20°

diameter static black circle on a gray background for 500 ms. The looming and static

object stimuli were both presented 15 times with 2 s inter-stimulus intervals.

Data processing

The data were band pass filtered (400 to 5000 Hz) by applying a Finite Impulse Response

(FIR) filter with a Hamming window (Choi *et al.* 2006). The background noise was

estimated using an adaptive filter algorithm (Biffi *et al.* 2010). Events that crossed a

threshold of 3.8 times the background noise were considered as spikes. Spikes were

sorted into clusters using the discrete wavelet transform for feature extraction, and

coefficients were selected to maximize the clusters' distance (Letelier and Weber 2000).

The clustering of single units was performed using a *k*-means algorithm with optimal *k*-

values obtained by the silhouette cluster analysis (Rousseeuw 1987) (see **Fig. 1B-C**).

Characterization of receptive field properties

214 *Receptive field size*

215 The RF area was assessed for each single unit by binning the data into 100 ms intervals,

216 subtracting the spontaneous firing rate, and fitting a two-dimensional (2D) Gaussian curve

217 to the average of the firing rate across trials:

$$fit = z_0 + Ae^{\left(\frac{1}{2(1-cor^2)}\right)\left[\left(\frac{x-x_0}{xWidth}\right)^2 + \left(\frac{y-y_0}{yWidth}\right)^2 - \left(\frac{2cor(x-x_0)(y-y_0)}{xWidth \cdot yWidth}\right)\right]} \qquad (1)$$

where z_0 is the matrix baseline (minimum value of firing rate after the subtraction of the

spontaneous firing rate), A is the maximal response amplitude, x_0 and y_0 are the horizontal

and vertical positions of the RF's center, *xWidth* and *yWidth* are the radii of the RF, and

cor is the correlation coefficient between two correlated Gaussian random variables. The

RF areas obtained for each single unit were subsequently categorized into depth ranges,

using the ranges defined in Table 1.

Spatial frequency tuning and contrast responses

Single unit responses were binned in 100 ms intervals and spontaneous firing rate was

subtracted. Spontaneous firing rate was obtained from one second interstimulus intervals

ranging from 0.5 -1.5 s after each stimulus presentation. Units considered in the analysis

showed a mean evoked firing rate $\geq$ 3 Hz after the spontaneous firing rate subtraction.

Tuning curves for spatial frequency (SF) were built using mean SF responses at the

contrast where the maximum response was evoked, and by fitting the data using a single

Gaussian function:

$$\mathrm{R}(sf) = R_b + \mathrm{Ae}^{\frac{(-sf - x_0)^2}{2\sigma^2}} \qquad (2)$$

where R_b is the baseline of the curve, A is the response's amplitude, x_0 is the position of

the center of the peak, and σ is the standard deviation. All variables were optimized by

finding the minimum of the unconstrained multivariable function using a derivative-free

method from MATLAB- fminsearch (Hendrikje *et al.* 2013; Camillo *et al.* 2020). Neurons

with $R^2 \leq 0.7$ in the Gaussian fitting function were not considered further for analysis. The

values for preferred spatial frequency were obtained from each single unit and were

classified as low-pass, band-pass or high-pass according to the specific spatial frequency

tuning they displayed. *Low-pass neurons* responded to low spatial frequencies (i.e., 0.01

cpd) with a peak firing rate larger than half of the peak's activity amplitude and showed

smaller (less than half of the peak's amplitude) or non-responses to higher spatial

frequencies. *Band-pass neurons* showed a bell shape response with a maximum at its

preferred SF while having low firing rates (lower than half of the peak's response

amplitude) for low *sf* (low cut-off) and for high SF (high cut-off). Finally, *high-pass neurons*

responded to higher spatial frequencies with firing rates larger than half of the peak

response (i.e., > 0.24 cpd). For those neurons showing a band-pass curve we calculated

the $\log_2$ bandwidth (high cut-off frequency/low cut-off frequency) to determine how narrow

or wide their tuning was (Heimel *et al.* 2005). We also evaluated the linearity of the

responses to the drifting grating stimuli at the preferred spatial frequency by calculating

the F1/F0 ratio, with F0 being the mean response and F1 the response modulation at the

drifting frequency. Both values were calculated by applying the discrete Fourier

transformation (Niell and Stryker 2008; Wang *et al.* 2010). Cells showing a F1/F0 $\geq$ 1

were classified as linear or simple, whereas those showing a F1/F0 < 1 were classified

as non-linear or complex. Single unit responses to different contrasts were obtained by

using the spatial frequency value exhibiting the maximum response. The mean responses

were fitted using a Naka-Rushton function (Naka and Rushton 1966):

$$R(C) = R_b + \frac{gC^n}{S^n + C^n} \qquad (3)$$

where R_b is the baseline of the curve, g the gain, S the contrast value at the mid saturation

point, n the exponent, and C the contrast. From the fitted curve we obtained: i) the contrast

value at half the peak response (C_{50}), which is a measure of contrast response linearity

(Heimel *et al.* 2005) and ii) the maximum slope of the fitted curve (maximum rate minus

the normalized spontaneous rate), which is a measure for non-linear responses to

contrast (Heimel *et al.* 2005).

Responses to dark looming objects.

Single unit responses to looming and static objects were binned in 10 ms intervals before,

during, and after the stimulus presentation. We calculated the neuronal response by first

obtaining the spontaneous firing rate for the 300 ms before the stimulus presentation,

then we averaged the responses of the 15 repetitions of each kind of stimulus (looming

and static objects) and subtracted the spontaneous firing rate. Raster-plots, however,

represent the response of a neuron (firing rate minus the spontaneous firing rate) for each

trial. To assess the onset time (latency) and peak response time to the object

presentation, z-scores were calculated for each bin taking 300 ms before and 500 ms

after each stimulus presented (t_{stim} = 0). To ensure the specific response to our visual

stimulation we chose a strict threshold; only neurons with a maximum response $\geq$ 5 z-

scores were considered in the analysis. The onset time of the response was the first bin

at which the response exceeded 2 z-score values and continued increasing until it

reached the peak response. The peak time corresponds to the time of the maximal

response between the onset of the stimulus and 250 ms (time for the looming object to

reach maximum size).

RESULTS

Cytoarchitecture of the SC

The mammalian superior colliculus is a laminar structure composed of 7 main layers,

namely, stratum zonale (SZ), stratum griseum superficiale (SGS), stratum opticum (SO),

stratum griseum intermediale (SGI), stratum album intermedium, (SAI) stratum griseum

profundus (SGP) and stratum album profundus (SAP). This latter layer separates the

SGC from the underlying periaqueductal grey (PAG). In agreement with previous findings

(Vega-Zuniga et al., 2014; Deichler et al., 2019), we found that SC in degus is a large

and highly differentiated structure, in which all layers are well-defined and readily

identifiable in Nissl preparations (**Fig. 2A, A'**). As in other mammals (e.g., mouse (Basso

and May 2017; Ito and Feldheim 2018), cat (May 2006), tree shrew (Albano *et al.* 1979)

and Galago (Balaram *et al.* 2011), but see Major *et al.* (2003)), the SGS in degus appears

to contain two subdivisions, upper (uSGS) and deep (dSGS), the former being composed

of smaller and less densely packed cells than the latter. Another striking feature is that

the SGI in degus appears composed of three subdivisions- upper (uSGI), intermediate

(iSGI) and deep (dSGI)- that are readily distinguishable by their differences in cellular

density (**Fig. 2A)**, these subdivisions have also been also observed in rats and mice

(Wiener 1986; Helms *et al.* 2004).

To determine the extent of retinal innervation of SC, we made retinal CTB injections

into the vitreous humor. We found that the SC is the largest and most densely innervated

retinal target in the degus visual system (**Fig. 2B**). Retinal projections to the degu SC

originate almost exclusively from the contralateral eye (Vega-Zuniga *et al.* 2013, 2014;

Deichler *et al.* 2020) and thus we did not quantify the ipsilateral projections. The

contralateral retinal projections enter the SC through the SO and terminate very densely

in the SZ and in both SGS subdivisions, and more sparsely in the SO. However, the

dorsal portion of the SO receives denser terminal arborizations than its ventral part (**Fig.**

2B, 2B'). Retinal projections were not present in the deeper layers of the SC, SGI, and

SGP.

Receptive field properties in the degu's SC

To characterize the RF properties in the Chilean degu (*Octodon degus*), we presented

different visual stimuli, as detailed in the methods section. Our main objective was to

characterize the RF properties within the retino-recipient layers, specifically the SGS and

SO, that are part of the superficial SC (sSC). Additionally, we assessed whether neurons

in the degu's SC exhibited the canonical organization of visual RFs characterized by

smaller RF sizes in more superficial layers compared to deeper layers, including the SGI.

Receptive field size and shape vary with depth in SC

As a first step to characterize the RF properties of SC neurons in degus, and to compare

our findings with previous work in hamsters (Carrasco *et al.* 2005) and mice (Fernández-

Aburto *et al.*, in prep), we recorded from neurons with RFs located in the frontal visual

field, and determined how their size and shape varied across the different SC layers (from

SGS to SGI). Illustrations of the size and shape of receptive fields for two single units

recorded from the SGS and SGI, respectively, are presented in Figures **3A** and **3B**.

Consistent with previous findings, degus showed the canonical organization of visual RFs

in the SC (Humphrey 1968), with RF sizes of single units increasing significantly with

depth from the SGS to the SGI layer (ANOVA $F_{[6,32]}$: 5.58, *p* = 0.00048; **Fig. 3C**). The

Tukey HSD (honestly significant difference) post-hoc test revealed that the RF areas of

single units recorded from the SGI depth range **7** (see **Table 1**) were significantly larger

than RF areas of single units from the SGS depth range **1** ($p = 0.0006$) and significantly

larger than RF areas recorded from units in the SGS depth range **2** ($p = 0.003$). RF areas

of single units from SGI depth range **7** were also significantly larger than RF areas of the

upper SO depth range **3** ($p = 0.003$; **Fig.3C**). In addition, we compared RF sizes between

the SC layers combined (SGS, SO and SGI), and found a significant difference (ANOVA

$F_{[2,36]}$: 12.46, $p < 0.0001$; **Fig. 3D).** RF areas from SGI neurons (area size range: 42.84-

162.8 deg^2) were significantly larger (Tukey HSD, $p = 0.0001$) than RF areas from SGS

neurons (area range: 19.8 – 90.97 deg^2), and significantly larger (Tukey HSD, $p = 0.0018$)

than RF areas from SO neurons (area range: 30.35 – 86.01 deg^2). RF sizes from SGS

neurons and SO neurons were similar (Tukey HSD test, $p = 0.36$).

345   Studies on rat and monkey SC have shown that there is a change in RF shape

across SC layers, going from near circular RFs in the superficial layers to RFs elongated

along the horizontal axis in the deeper layers (Humphrey 1968). To investigate whether

degus also exhibit an alteration in receptive RF shape (i.e., circular vs. oval shape), we

compared the ratio between the horizontal and vertical axes of the RFs across the SC

depth ranges. A ratio close to 1 indicates a symmetrical RF with a circular shape, whereas

deviations from this value indicate an elongation of the horizontal axis. Similar to the

findings reported by Humphrey (1968), RF symmetry changes significantly between depth

ranges; RFs from deeper layers are more elongated along the horizontal axis than RFs

from upper layers (ANOVA $F_{[6,32]}$: 4.04, $p = 0.004$; **Fig. 3E**). The RFs of neurons from SGI

depth range **7** were more elongated along the horizontal axis (i.e., asymmetrical) than the

circular RF shape of neurons from SGS depth range **1** (p = 0.004) and were also more

elongated than the RF shape of neurons recorded from the SGS depth range **2** (p = 0.01).

Likewise, neurons from SO depth range **3** showed RF shapes that are significantly more

circular than the elongated RF shapes of neurons from SGI depth range **7** (p = 0.037;

Fig. 3E). Comparisons focusing on the differences between SC layers as a whole rather

than the depth level confirmed that as recordings sampled the lower layers, RFs shapes

became more elongated along the horizontal axis (ANOVA $F_{[2,36]}$: 6.69, p = 0.003; **Fig.**

3F). The RFs of SGS neurons were significantly more symmetric (i.e., circular) than RFs

of SGI neurons (Tukey HSD, p = 0.002). The shape of the RFs from SO neurons was

statistically the same as the RF shape of single units from the SGS and SGI, suggesting

a tendency of SO neurons to have elongated RFs.

Spatial frequency tuning

Sinusoidal gratings of different spatial frequencies (SF) and contrasts (**Fig. 4A**) have

been widely used to characterize RF properties of different visual brain structures in

several animal species (Jassik-Gerschenfeld and Hardy 1979; Issa *et al.* 2000; Heimel *et*

al. 2005). In the case of degus, we found that 47% (228/485) of SGS neurons that were

recorded responded to sinusoidal grating stimuli. Of those, 58.3% (133/228) were tuned

to specific combinations of SF and contrast (see below). According to their spatial

frequency tuning, we classified neurons as low-pass, band-pass or high-pass (see

Methods). Most SC neurons were band-pass (60.9%), or low pass (36.8 %); only a few

were classified as high-pass neurons (2.3%; see Figures **4B** through **4E'**). Even though

the median value of SF preference across all three cell types (low-pass, band-pass and

high-pass) was 0.04 cycles per degree (cpd), we found a subset of band-pass neurons

(18%) tuned to a higher SF (0.24 cpd) (**Fig. 4F**).

To further assess the specificity of SF tuning in band-pass neurons, we quantified the

tuning curves' bandwidths (the range of SF to which a band-pass neuron responds).

Interestingly, 41% of band-pass neurons were tuned to their preferred SF, with

bandwidths ranging between 0.5 and 2.0 octaves (**Fig. 4G**). Additionally, we assessed

the extent to which spatial frequency modulates the activity of degus' SC units by

calculating the linearity index $F1/F0$ (where $F1$ is the modulation of the response (firing

rate) at the drift frequency of the stimulus, and $F0$ is the average firing rate of the

response, see Methods) (De Valois *et al.* 1982). We found that most SGS neurons (86%)

responded non-linearly to their preferred SF ($F1/F0 < 1$) **(Fig. 4H)**, and the rest (14%)

responded linearly ($F1/F0 > 1$) (**Fig. 4I**). The distributions of linear/non-linear responses

for the neuronal population is shown in Figure **4J**.

Contrast tuning

Overall, SGS neurons increased their response levels with increased contrast. However,

there were neurons that showed a linear response to contrast without saturating their

response (53.3%; **Fig. 5A**), and others (46.7%) that exhibited different degrees of

saturation at higher contrast (**Fig. 5B-C**). Interestingly, such saturating responses are

characterized by high contrast gain at low contrasts and low contrast gain at high

contrasts (Alitto *et al.* 2019). We further characterized linear and saturating responses by

quantifying the contrast at half of the maximum response (C_{50} value) and the relative

maximum gain (RMG), i.e., the maximum slope of the normalized contrast response

curves (Heimel *et al.* 2005). At the population level, 50% of the neurons showed that the

contrast at half of the maximum response (C_{50} value) was 48%, indicating a linear

response to contrast (**Fig. 5D**). On the other hand, neurons that show saturating

responses can have high (C_{50} < 30, 36% of neurons) or moderate (C_{50} between 30 and

40, 17% of neurons) saturation of their responses to contrast. Additionally, the median

value of RMG was 1.47 (**Fig. 5E**), indicating slight non-linearity, compared to non-linear

responses to contrast in the squirrel's V1 with a median RMG value of 2.8 (Heimel *et al.*

2005) . Only 22/135 neurons (16.3%) had a RMG of approximately 1 (between 0.8 and

1.4) that we considered as a linear behavior, whereas 28/135 showed RMG values $\geq$ 3

(20.7%) and were thus considered non-linear. Interestingly, 50% of neurons (170/477)

showed RMG values between 1.4 and 1.5, which we interpreted as weak non-linear

responses. Although both measures, C_{50} and RMG, are indicators of linearity, the RMG

is less dependent on the fitting process (Heimel *et al.* 2005). Still, both measures suggest

that the degus SGS has neuron types that respond in a linear and a nonlinear fashion.

SGS neurons respond in a continuously increasing way to looming

To evaluate the units' responses to looming stimuli, we presented 15 repetitions of a dark

disc that increased in size from 2 to 20 deg in 250 ms and stayed ON for another 250 ms.

As a control we presented a static dark disc (20°) for 500 ms to address the specificity of

the response to looming (**Fig. 6A and A'**). Neurons exposed to the looming object

continuously increased their firing rate as the object increased in size, followed by a

decrease of the firing rate when the object no longer increased in size. A small OFF

response can be observed (**Fig. 6B and 6C**). Neurons exposed to the static object, on

the other hand, showed a typical ON-OFF response (**Fig. 6B' and 6C'**). Of a total of 251

responsive neurons, 146 responded to both stimuli, 58 only to looming, and 47 only to the

static object. Interestingly, the onset time of the response to both stimuli did not differ

significantly (paired *t*-test, $p = 0.3$), showing that neurons start their response at the same

time irrespective of the visual object shown (**Fig. 7A**). The difference between the

response to the looming and to the static object can be observed in other parameters- in

peak time (**Fig. 7B**) and maximal firing rate after the initial response (**Fig. 7D**). With the

looming parameters used, neurons exposed to the looming take longer to reach their

maximal response (paired *t*-test, $p = 0.0066$) and show significantly higher firing rates

(paired *t*-test, $p < 0.0016$) than when exposed to the static object.

DISCUSSION

It is well known that mature visual properties as well as visual system development differ between

mammalian species in a way that is related to the role played by vision in their behavior.

Interestingly, most of the recent publications about visual system development and function come

from a single model species, the lab mouse *(Mus musculus)*. Lab mice are altricial, nocturnal, and

feature a poorly developed visual system when compared with diurnal rodents such as squirrels,

guinea pigs and some other New World rodents. Thus, mouse-centered studies may have biased

the current understanding of rodent visual system function and its development. As a starting

point to counterbalance this bias, we characterized the visual properties of SC neurons in the

adult *Octodon degus* in the present study. Degus, contrary to mice and other typical lab rodent

species such as rats and hamsters, are diurnal (Fulk 1976) and precocial (Reynolds and Wright

1979; Long and Ebensperger 2010). Furthermore, degus represent a phylogenetic group

(suborder Hystricomorpha) apart from old world rodents such as rats and mice (suborder

Myomorpha) or squirrels (suborder Sciuromorpha). As we predicted, we found that *O. degus'* SC

is a well-developed visual structure, featuring visual properties comparable to that of highly visual

rodents such as squirrels (Van Hooser *et al.* 2003) and gerbils (Baker and Emerson 1983). We

conclude that, given its precocial and diurnal habit, *O. degus* is especially well-suited as an animal

model for studies of visual system development.

Cytoarchitecture of the superior colliculus in *Octodon degus*

Our anatomical results readily confirmed that degus' SC features a laminar differentiation

comparable to that of other diurnal mammals, such as the squirrel (Major *et al.* 2003; Fredes *et*

al. 2012). In particular, the extensive laminar differentiation of the SGI suggests that in degus the

SC serves a stronger and more complex visuomotor role than in other rodents, a conjecture

worthy of further studies. In addition, the results of our retinal injections confirm previous studies

(Vega-Zuniga et al., 2013; Deichler et al., 2019; 2020) reporting that degu SC receives massive

and highly ordered retinal projections. As in the previous studies, we found that retinal afferents

enter the contralateral SC through the SO to form dense terminal domains in the stratum griseum

superficiale (SGS) and stratum zonale (SZ), and sparser arborizations in the stratum opticum

(SO). As expected from studies of other species, retinal projections were not present in the deeper

layers of the SC (dSC), namely the stratum griseum intermediale (SGI) and the stratum griseum

profundus (SGP).

Receptive field size

We found that the degu's SC has a laminar segregation of RF sizes, with smaller RFs located in

superficial layers, and larger RFs in deeper layers. This seems to be a canonical property of the

SC -*aka* optic tectum- across vertebrate taxa, in that it has been reported not only in other rodents,

such as mice, rats and hamsters (Humphrey 1968; Rhoades and Chalupa 1976; Ito *et al.* 2017),

but also in Carnivore species (cats and ferrets; Brecht et al., 1999; Stitt et al., 2013), primates

(Humphrey 1968; Cynader and Berman 1972) and birds (Knudsen 1982; Marín *et al.* 2005). This

change in RF size across layers correlates with the different morphological characteristics of

neurons located in each layer. In mammals, narrow field and stellate neurons located in the upper

SGS (uSGS) have narrow dendritic trees and small RFs (Mooney *et al.* 1985; May 2006; Hoy *et*

al. 2019), whereas wide field and horizontal cells, located in the deeper SGS (dSGS), have large

dendritic trees (~500-800 μm) along with very large RFs (~300-1000 deg^2) (Major *et al.* 2000;

Fredes *et al.* 2012; Gale and Murphy 2014). In the dSC (SGI and SGP), neurons that have large

dendritic trees also exhibit large RFs, most of them being premotor neurons that receive

convergent inputs from neurons located in the superficial layers (Isa and Hall 2009).

Superior colliculus RF sizes are highly variable across rodent species. In hamsters the SGS

RFs are quite large (ranging from 169 deg^2 to ~295 deg^2 [19.4 ± 0.56° diameter] (Carrasco *et al.*

2005; Balmer and Pallas 2015). In mice, on the other hand, contrasting results have been

reported. Some studies report large RF sizes (average 165 ± 21 deg^2 (Chandrasekaran *et al.*

2005; Ito *et al.* 2017) but others report smaller values (107.5 $\pm$ 14.9 deg^2, (Wang *et al.* 2010)).

These differences may depend on the visual stimulation method used. Ito *et al.* (2017) used a 10º

flashing-spot but Wang *et al.* (2010) used a 5º flashing square. In the present study we used a

small (3º), moving bright square as a visual stimulus, which is bigger than that used in previous

studies of hamsters (1º) (Carrasco *et al.* 2005; Mudd *et al.* 2019). Nevertheless, despite

methodological differences, degus seem to have the smallest RF sizes reported in rodents for the

sSC so far, with area sizes that never surpassed 100 deg^2 (ranging from 19.8 deg^2 to 74.4 deg^2).

It is noteworthy that degus are diurnal animals and possess a cone-rich retina (Chávez *et al.*

2003; Jacobs *et al.* 2003; Vega-Zuniga *et al.* 2013). These features suggest that degus would

have a visual acuity higher than that of nocturnal species. The comparatively small size of

degu SC receptive fields is indeed consistent with a greater visual acuity, reflecting the diurnal

specialization of the visual system in these rodents.

Response to grating stimuli

In cats, hooded rats, and squirrels, it has been shown that the presence of visual neurons tuned

to high spatial frequencies correlates with higher visual acuity (Dean 1981; Issa *et al.* 2000;

Heimel *et al.* 2005). In the case of degus, we found that most sSC neurons (> 50%) that responded

to gratings exhibited preferences for low/medium spatial frequencies. Interestingly, we found

another set of neurons (15%) that were tuned to higher SFs (with a peak of 0.24 cpd). Overall,

the SF tuning reported for mouse SC [peak=0.08 cpd, (Wang *et al.* 2010); median= 0.052 cpd,

(De Franceschi and Solomon 2018) is lower than what we found in degus, suggesting that, as

expected from its diurnal visual specialization, degus have a higher visual acuity than mice.

Interestingly, it has been reported that awake mice show an approximately 2-fold higher SF tuning

than anesthetized mice (De Franceschi and Solomon 2018)]. Here we only tested deeply

anesthetized animals, thus awake degus might show higher SF tuning than we report here.

Spatial frequency modulation. In degu sSC, most (86%) of the neurons recorded exhibited non-

linear (F1/F0 <1) responses to grating stimuli, thus showing low modulation by the SF presented.

However, some neurons (14%) were modulated by the SF (F1/F0 >1). A similar distribution of

linear and non-linear neurons was found in mice (Wang *et al.* 2010). Interestingly, neurons

showing low or no SF modulation were tuned to higher frequencies, a phenomenon also reported

in mice (De Franceschi and Solomon 2018).

Contrast sensitivity. We found that almost half (44%) of the neurons responded in a linear,

monotonically increasing fashion to increasing stimulus contrast (C_{50} > 45), whereas others

showed highly or moderately saturating responses to contrast. It has been proposed that

saturated responses to contrast, rather than being the result of biophysical limitations of neurons,

are due to an active mechanism controlling the gain of contrast responses (Peirce 2007). Thus,

saturating responses to contrast relate to higher sensitivity to subtle contrast differences at low

contrast values, which is an important adaptation for active nocturnal vision. In mice most SC

neurons exhibit non-linear, saturating responses to contrasts (De Franceschi and Solomon 2018),

as expected for nocturnal species. Unlike what has been observed in mice, we found no clear

bias toward saturating responses to contrast in the degus' SC, likely reflecting the diurnal

adaptations of their visual system.

Looming responses

In degus, we found many sSC neurons that showed a progressive increase in their responses to

an expanding dark disc that simulated a dark looming stimulus approaching against a lighter

background. This is similar to what has been reported in mice and hamsters, in which neurons

located in sSC respond reliably to repeated presentation of a dark looming stimulus in their RFs

(Mudd *et al.* 2019; Lee *et al.* 2020). Most of the neurons recorded in this study (> 55 %) showed

a clear difference in their response properties between looming and static stimuli, which reflects

the importance of animals responding differentially to different kinds of stimuli, i.e., a fast-

approaching object (e.g., a predator) vs. a large, stationary object. In this regard, we observed

that neurons responding to both stimulus types (looming and static objects) had significantly

higher firing rates when exposed to the looming object than when exposed to the static control,

suggesting that the SC possesses a neural mechanism that allows it to identify a fast-approaching

object. This mechanism might be involved in the execution of behaviors in this context. In rodents,

looming stimuli have been reported to unleash behaviors modulated by SC such as escaping and

freezing (Yilmaz and Meister 2013; De Franceschi and Solomon 2018). In degus, looming stimuli

projected from the upper, binocular portion of the visual field elicit escape behavior, which is

thought to be triggered by the SC/parabigeminal nucleus interaction (Deichler *et al.* 2020).

Conclusion

Degus show remarkably specialized visual properties in the superior colliculus compared to

commonly studied lab rodents. Our results suggest that degus have diurnal visual characteristics,

and a higher visual acuity than other lab rodents. Moreover, degus are precocial, diurnal and

belong to a separate clade within the Rodentia order, allowing studies to address from a

comparative perspective which visual features are conserved among rodents or other mammals,

and which might be determined by their ecological niche. Altogether, we strongly argue that degus

are a promising model to study the visual system in general and its development in particular.

BIBLIOGRAPHY

Albano JE, Norton TT, Hall WC (1979) Laminar origin of projections from the superficial layers of the superior colliculus in the tree shrew, *Tupaia glis*. Brain Res 173 (1): 1-11. doi: https://doi.org/10.1016/0006-8993(79)91090-4

Alitto HJ, Rathbun DL, Fisher TG, Alexander PC, Usrey WM (2019) Contrast gain control and retinogeniculate communication. Eur J Neurosci 49 (8): 1061-1068. doi: https://doi.org/10.1111/ejn.13904

Baker AG, Emerson VF (1983) Grating acuity of the mongolian gerbil (*Meriones unguiculatus*). Behav Brain Res 8 (2): 195-209. doi: https://doi.org/10.1016/0166-4328(83)90054-2

Balaram P, Hackett TA, Kaas JH (2011) VGLUT1 mRNA and protein expression in the visual system of prosimian galagos (*Otolemur garnetti*). Eye Brain 3: 81-98. doi: https://doi.org/10.2147/EB.S23007

Balmer TS, Pallas SL (2015) Refinement but not maintenance of visual receptive fields is independent of visual experience. Cereb Cortex 25 (4): 904-917. doi: https://doi.org/10.1093/cercor/bht281

Basso MA, May PJ (2017) Circuits for action and cognition: A view from the Superior Colliculus. Annu Rev Vis 3 (1): 197-226. doi: https://doi.org/10.1146/annurev-vision-102016-061234

Biffi E, Ghezzi D, Pedrocchi A, Ferrigno G (2010) Development and validation of a spike detection and classification algorithm aimed at implementation on hardware devices. Comput Intell Neurosci 2010: Article ID 659050, 15 pages, 2010. doi: https://doi.org/10.1155/2010/659050I

Brainard MS, Doupe AJ (2013) Translating birdsong: songbirds as a model for basic and applied medical research. Annu Rev Neurosci 36: 489-517. doi: https://doi.org/10.1146/annurev-neuro-060909-152826

Camillo D, Ahmadlou M, Heimel JA (2020) Contrast-dependence of temporal frequency tuning in mouse V1. Front Neurosci 14: 868. doi: https://doi.org/10.3389/fnins.2020.00868

Carrasco MM, Pallas SL (2006) Early visual experience prevents but cannot reverse deprivation-induced loss of refinement in adult superior colliculus. Vis Neurosci 23 (6): 845-852. doi: https://doi.org/10.1017/S0952523806230177

Carrasco MM, Razak KA, Pallas SL (2005) Visual experience is necessary for maintenance but not development of receptive fields in superior colliculus. J Neurophysiol 94 (3): 1962-1970. doi: https://doi.org/10.1152/jn.00166.2005

Chandrasekaran A, R. , Plas D, T. , Gonzalez E, Crair M, C. (2005) Evidence for an instructive role of retinal activity in retinotopic map refinement in the superior colliculus of the mouse. J Neurosci 25 (29): 6929-6938. doi: https://doi.org/10.1523/JNEUROSCI.1470-05.2005

Chávez AsE, Bozinovic F, Peichl L, Palacios AnG (2003) Retinal spectral sensitivity, fur coloration, and urine reflectance in the genus *Octodon* (rodentia): Implications for visual ecology. Invest Ophthalmol Vis Sci 44 (5): 2290-2296. doi: https://doi.org/10.1167/iovs.02-0670

Choi JH, Jung HK, Kim T (2006) A new action potential detector using the MTEO and its effects on spike sorting systems at low signal-to-noise ratios. IEEE Trans Biomed Eng 53 (4): 738-746. doi: https://doi.org/10.1109/TBME.2006.870239

Cynader M, Berman N (1972) Receptive-field organization of monkey superior colliculus. J Neurophysiol 35 (2): 187-201. doi: https://doi.org/10.1152/jn.1972.35.2.187

De Franceschi G, Solomon SG (2018) Visual response properties of neurons in the superficial layers of the superior colliculus of awake mouse. J Physiol 596 (24): 6307-6332. doi: https://doi.org/10.1113/JP276964

De Valois RL, Albrecht DG, Thorell LG (1982) Spatial frequency selectivity of cells in macaque visual cortex. Vision Res 22 (5): 545-559. doi: https://doi.org/10.1016/0042-6989(82)90113-4

Dean P (1981) Visual pathways and acuity in hooded rats. Behav Brain Res 3 (2): 239-271. doi: https://doi.org/10.1016/0166-4328(81)90050-4

DeBello WM, Knudsen EI (2004) Multiple sites of adaptive plasticity in the owl's auditory localization pathway. J Neurosci 24 (31): 6853-6861. doi: https://doi.org/10.1523/jneurosci.0480-04.2004

Deichler A, Carrasco D, Gonzalez-Cabrera C, Letelier JC, Marín G, Mpodozis J (2019) The nucleus pretectalis principalis: A pretectal structure hidden in the mammalian thalamus. J Comp Neurol 527 (2): 372-391. doi: https://doi.org/10.1002/cne.24540

Deichler A, Carrasco D, Lopez-Jury L, Vega-Zuniga T, Márquez N, Mpodozis J, Marín GJ (2020) A specialized reciprocal connectivity suggests a link between the mechanisms by which the superior colliculus and parabigeminal nucleus produce defensive behaviors in rodents. Sci Rep 10 (1): 16220. doi: https://doi.org/10.1038/s41598-020-72848-0

Espinosa JS, Stryker Michael P (2012) Development and plasticity of the primary visual cortex. Neuron 75 (2): 230-249. doi: https://doi.org/10.1016/j.neuron.2012.06.009

Fredes F, Vega-Zuniga T, Karten H, Mpodozis J (2012) Bilateral and ipsilateral ascending tectopulvinar pathways in mammals: A study in the squirrel (*Spermophilus beecheyi*). J Comp Neurol 520 (8): 1800-1818. doi: https://doi.org/10.1002/cne.23014

Fulk G (1976) Notes on the activity, reproduction and social behaviour of *Octodon degus.* J of Mammalogy 57: 495-505. doi: https://doi.org/10.2307/1379298

Gale SD, Murphy GJ (2014) Distinct representation and distribution of visual information by specific cell types in mouse superficial superior colliculus. J Neurosci 34 (40): 13458-13471. doi: https://doi.org/10.1523/jneurosci.2768-14.2014

Garrett ME, Nauhaus I, Marshel JH, Callaway EM (2014) Topography and areal organization of mouse visual cortex. J Neurosci 34 (37): 12587-12600. doi: https://doi.org/10.1523/jneurosci.1124-14.2014

Heimel JA, Van Hooser SD, Nelson SB (2005) Laminar organization of response properties in primary visual cortex of the gray squirrel (*Sciurus carolinensis*). J Neurophysiol 94 (5): 3538-3554. doi: https://doi.org/10.1152/jn.00106.2005

Helms MC, Özen G, Hall WC (2004) Organization of the intermediate gray layer of the superior colliculus. I. Intrinsic vertical connections. J Neurophysiol 91 (4): 1706-1715. doi: https://doi.org/10.1152/jn.00705.2003

Hendrikje N, Andrea H, Ian N, Hiroki T, Huang ZJ, Edward MC (2013) Contrast dependence and differential contributions from somatostatin- and parvalbumin-expressing neurons to spatial integration in mouse V1. J Neurosci 33 (27): 11145-11154. doi: https://doi.org/10.1523/jneurosci.5320-12.2013

Hensch TK, Fagiolini M, Mataga N, Stryker MP, Baekkeskov S, Kash SF (1998) Local GABA circuit control of experience-dependent plasticity in developing visual cortex. Science 282 (5393): 1504-1508. doi: https://doi.org/10.1126/science.282.5393.1504

Hoy JL, Bishop HI, Niell CM (2019) Defined cell types in superior colliculus make distinct contributions to prey capture behavior in the mouse. Curr Biol 29 (23): 4130-4138. doi: https://doi.org/10.1016/j.cub.2019.10.017

Hubel DH, Wiesel TN (1998) Early exploration of the visual cortex. Neuron 20 (3): 401-412. doi: https://doi.org/10.1016/s0896-6273(00)80984-8

Huberman AD, Niell CM (2011) What can mice tell us about how vision works? Trends Neurosci 34 (9): 464-473. doi: https://doi.org/10.1016/j.tins.2011.07.002

Humphrey NK (1968) Responses to visual stimuli of units in the superior colliculus of rats and monkeys. Exp Neurol 20 (3): 312-340. doi: https://doi.org/10.1016/0014-4886(68)90076-9

Isa T, Hall WC (2009) Exploring the superior colliculus *in vitro*. J Neurophysiol 102 (5): 2581-2593. doi: https://doi.org/10.1152/jn.00498.2009

Issa NP, Trepel C, Stryker MP (2000) Spatial frequency maps in cat visual cortex. J Neurosci 20 (22): 8504-8514. doi: https://doi.org/10.1523/jneurosci.20-22-08504.2000

Ito S, Feldheim DA (2018) The mouse superior colliculus: An emerging model for studying circuit formation and function. Front Neural Circuits 12: 10. doi: https://doi.org/10.3389/fncir.2018.00010

Ito S, Feldheim DA, Litke AM (2017) Segregation of visual response properties in the mouse superior colliculus and their modulation during locomotion. J Neurosci 37 (35): 8428-8443. doi: https://doi.org/10.1523/jneurosci.3689-16.2017

Jacobs GH, Calderone JB, Fenwick JA, Krogh K, Williams GA (2003) Visual adaptations in a diurnal rodent, Octodon degus. J Comp Physiol A 189 (5): 347-361. doi: https://doi.org/10.1007/s00359-003-0408-0

Jassik-Gerschenfeld D, Hardy O (1979) Single-neuron responses to moving sine-wave gratings in the pigeon optic tectum. Vision Res 19 (9): 993-999. doi: https://doi.org/10.1016/0042-6989(79)90224-4

Jones HG, Brown AD, Koka K, Thornton JL, Tollin DJ (2015) Sound frequency-invariant neural coding of a frequency-dependent cue to sound source location. J Neurophysiol 114 (1): 531-539. doi: https://doi.org/10.1152/jn.00062.2015

Juavinett Ashley L, Callaway Edward M (2015) Pattern and component motion responses in mouse visual cortical areas. Curr Biol 25 (13): 1759-1764. doi: https://doi.org/10.1016/j.cub.2015.05.028

Knudsen EI (1982) Auditory and visual maps of space in the optic tectum of the owl. J Neurosci 2 (9): 1177-1194. doi: https://doi.org/10.1523/jneurosci.02-09-01177.1982

Lee KH, Tran A, Turan Z, Meister M (2020) The sifting of visual information in the superior colliculus. eLife 9: e50678. doi: https://doi.org/10.7554/eLife.50678

Letelier JC, Weber PP (2000) Spike sorting based on discrete wavelet transform coefficients. J Neurosci Methods 101 (2): 93-106. doi: https://doi.org/10.1016/S0165-0270(00)00250-8

Long CV, Ebensperger LA (2010) Pup growth rates and breeding female weight changes in two populations of captive bred degus (*Octodon degus*), a precocial caviomorph rodent. Reprod Domest Anim 45 (6): 975-982. doi: https://doi.org/10.1111/j.1439-0531.2009.01470.x

Major DE, Luksch H, Karten HJ (2000) Bottlebrush dendritic endings and large dendritic fields: Motion-detecting neurons in the mammalian tectum. J Comp Neurol 423 (2): 243-260. doi: https://doi.org/10.1002/1096-9861(20000724)423:2<243::AID-CNE5>3.0.CO;2-5

Major DE, Rodman HR, Libedinsky C, Karten HJ (2003) Pattern of retinal projections in the California ground squirrel (*Spermophilus beecheyi*): Anterograde tracing study using cholera toxin. J Comp Neurol 463: 317-340. doi: https://doi.org/10.1002/cne.10764

Marín G, Mpdozis J, Sentis E, Ossandón T, Letelier JC (2005) Oscillatory bursts in the optic tectum of birds represent re-entrant signals from the nucleus isthmi pars parvocellularis. J Neurosci 25 (30): 7081-7089. doi: https://doi.org/10.1523/jneurosci.1379-05.2005

Márquez N, Martínez-Harms J, Vásquez RA, Mpodozis J (2015) Early olfactory environment influences social behaviour in adult *Octodon degus*. PLOS ONE 10 (2): e0118018. doi: https://doi.org/10.1371/journal.pone.0118018

May PJ (2006) The mammalian superior colliculus: laminar structure and connections. Prog Brain Res 151: 321-378. doi: https://doi.org/10.1016/S0079-6123(05)51011-2

Mooney RD, Klein BG, Rhoades RW (1985) Correlations between the structural and functional characteristics of neurons in the superficial laminae and the hamster's superior colliculus. J Neurosci 5 (11): 2989-3009. doi: https://doi.org/10.1523/jneurosci.05-11-02989.1985

Mudd DB, Balmer TS, Kim SY, Machhour N, Pallas SL (2019) Trkb activation during a critical period mimics the protective effects of early visual experience on perception and the stability of receptive fields in adult superior colliculus. J Neurosci 39 (23): 4475-4488. doi: https://doi.org/10.1523/jneurosci.2598-18.2019

Naka KI, Rushton WA (1966) S-potentials from colour units in the retina of fish (Cyprinidae). J Physiol 185 (3): 536-555. doi: https://doi.org/10.1113/jphysiol.1966.sp008001

Niell CM, Stryker MP (2008) Highly selective receptive fields in mouse visual cortex. J Neurosci 28 (30): 7520-7536. doi: https://doi.org/10.1523/jneurosci.0623-08.2008

Peirce JW (2007) The potential importance of saturating and supersaturating contrast response functions in visual cortex. J Vis 7 (6): 1-10. doi: https://doi.org/10.1167/7.6.13

Razak KA (2011) Systematic representation of sound locations in the primary auditory cortex. J Neurosci 31 (39): 13848-13859. doi: https://doi.org/10.1523/jneurosci.1937-11.2011

Razak KA, Fuzessery ZM (2015) Development of echolocation calls and neural selectivity for echolocation calls in the pallid bat. Dev Neurobiol 75 (10): 1125-1139. doi: https://doi.org/10.1002/dneu.22226

Reynolds TJ, Wright JW (1979) Early postnatal physical and behavioural development of degus (*Octodon degus*). Lab Anim 13 (2): 93-100. doi: https://doi.org/10.1258/002367779780943576

Rhoades RW, Chalupa LM (1976) Directional selectivity in the superior colliculus of the golden hamster. Brain Res 118 (2): 334-338. doi: https://doi.org/10.1016/0006-8993(76)90721-6

Rousseeuw PJ (1987) Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. J Comput Appl Math 20: 53-65. doi: https://doi.org/10.1016/0377-0427(87)90125-7

Van Hooser SD, Heimel JAF, Nelson SB (2003) Receptive field properties and laminar organization of lateral geniculate nucleus in the gray squirrel (*Sciurus carolinensis*). J Neurophysiol 90 (5): 3398-3418. doi: https://doi.org/10.1152/jn.00474.2003

Vega-Zuniga T, Medina FS, Fredes F, Zuniga C, Severín D, Palacios AG, Karten HJ, Mpodozis J (2013) Does nocturnality drive binocular vision? Octodontine rodents as a case study. PLOS ONE 8 (12): e84199. doi: https://doi.org/10.1371/journal.pone.0084199

Wang L, Sarnaik R, Rangarajan K, Liu X, Cang J (2010) Visual receptive field properties of neurons in the superficial superior colliculus of the mouse. J Neurosci 30 (49): 16573-16584. doi: https://doi.org/10.1523/jneurosci.3305-10.2010

Wiener SI (1986) Laminar distribution and patchiness of cytochrome oxidase in mouse superior colliculus. J Comp Neurol 244 (2): 137-148. doi: https://doi.org/10.1002/cne.902440202

Yilmaz M, Meister M (2013) Rapid innate defensive responses of mice to looming visual stimuli. Curr Biol 23 (20): 2011-2015. doi: https://doi.org/10.1016/j.cub.2013.08.015

771 **TABLES**

772 **Table 1.**

Depth range (µm)*	SC layer	Depth range
< 400	SGS	1
(400, 500)	SGS	2
(501, 600)	SO	3
(601, 750)	SO	4
(751, 950)	SGI	5
(951, 1100)	SGI	6
(1101, 1300)	SGI	7

774 **Table 1. Categorization of depth ranges.** Visual responses were systematically
775 recorded from neurons with RFs located within the frontal visual field, at intervals of 100
776 µm along transects across the SC layers. Neuronal responses were categorized into
777 seven distinct depth ranges according to the depth at which they were recorded.

Figure 1. Visual stimulation, extracellular recording, and spike sorting. (A) Visual stimuli
used to characterize RF size and shape. A small white square (~3° diameter) on a black
background was swept across the screen monitor from top to bottom with each vertical
sweep (27°/s, 4 sec), displaced horizontally by 3°, at 1s intervals, to cover the entire
screen. (B) Top trace, extracellular responses in the SC. Bottom trace, the visual
stimulus on/off pattern as described in A. (C) Example of 3 different single units isolated
from recordings.

790
791
Figure 2. (A) Photomicrograph of a Nissl-stained coronal section showing the
subdivisions of the degus' SC: stratum griseum superficiale (SGS) upper (uSGS)
and deeper (dSGS), stratum opticum (SO), stratum griseum intermediale (SGI) upper (uSGI),
intermediate (iSGI) and deeper (dSGI), the stratum griseum profundus (SGP), and PAG,
periacueductal gray. **(A')** Detail of a Nissl-stained coronal section of SC laminae. **(B)**
Anterograde labeling of retinotectal projections after a CTB injection into the contralateral
eye. **(B')** Detail of the retinotectal projection into the different layers of the degus' SC.
799

Figure 3. Receptive field size and shape change across SC layers in degus. (A) Heat

map showing the RF of a single unit isolated at 320 µm depth in the SGS mapped by

visual stimulation as in Fig. 1A. **(B)** Heat map showing the RF of a unit at 1100 (µm) in

the SGI mapped with the same stimulation, at the same recording site (stereotaxic

coordinates) as in **A**, but deeper. **(C)** Boxplot of RF areas (in deg^2) across depths ranges

from the SGS to the SGI. Boxplots show standard interquartile range (IQR) and median

value (black central line), yellow dots are the individual data points. RFs increase in size

with depth (ANOVA: $p < 0.01$). **(D)** Boxplot of RF areas across SC layers (SGS, SO and

SGI). **(E)** Boxplot of RF shape expressed as the width/height ratio at different SC depth

ranges. RFs widen along the horizontal axis with depth (ANOVA: $p < 0.01$). **(F)** Boxplot of

RF shape at different SC layers. Different letters above each group in B, C, D, and F

represent statistically significant differences (Tukey post-hoc test, see text for exact *p*-

values).

Figure 4. Single units show selectivity to spatial frequency. sSC neurons have a

narrow bandwidth, and a small proportion of them (14%) respond linearly to sinusoidal

gratings. **(A)** Illustration of the drifting grating stimulus types. Spatial frequency and

contrast were varied in a pseudorandom manner. **(B)** Percentage of neurons with low-

pass, band-pass or high-pass selectivity to different spatial frequencies (n = 128). **(C-E)**

Response matrix (instantaneous firing rate, in Hz) of three different neurons to all contrast

and spatial frequency combinations. **(C′-E′)** Examples of low-pass **(C′)**, band-pass **(D′)**,

and high pass **(E′)** tuning of the same single neurons as in **C-E**. **(F)** Bar plot of spatial

frequency preference for the population. Most units are tuned to 0.04 or 0.08 (cpd), and

a smaller group is tuned to higher spatial frequency (0.24 cpd). **(G)** Bar plot of bandwidth

showing that most neurons (>50%) are tuned to a certain spatial frequency with narrow

bandwidths. **(H-I)** Solid lines are the mean firing rate (100 ms bins) and shaded regions

represent mean $\pm$ sd. **(H)** Example of a single unit responding non-linearly to 4 seconds

of stimulation with a sinusoidal grating (linearity index F1/F0 < 1, where *F1* is the response

at the drift frequency, and *F0* is the average firing rate). **(I)** Example of a different single

unit responding in a linear manner to sinusoidal gratings (F1/F0 >1). **(J)** Histogram of the

linearity index across the entire population of single units (n = 232) showing neurons

responding linearly to sinusoidal gratings (red bars) and neurons with non-linear

responses to sinusoidal gratings (blue bars).

Figure 5. SC Single units show linear or non-linear responses to contrast. Example of three different single units' tuning to contrast, ranging from linear **(A)** to non-linear **(B and C).** All three curves are fitted with a Naka-Rushton function (black line). The dotted line is the spontaneous firing rate (Hz) and the dashed line is the C_{50} (the contrast at half the peak response). Note that **(B)** and **(C)** show saturating responses and differ in their C_{50} values (~40 and ~20, respectively). **(D)** Cumulative histogram of C_{50} distribution (N=135), with the median C_{50} = 48. **(E)** Cumulative histogram (N=135) of the relative maximum gain (see Methods).

849
850

Figure 6. Single units show a continuous buildup of response level to looming stimuli but an ON-OFF response to static stimuli. (A) Diagram of visual stimulation. The looming object increased in size from 2° to 20° over 250 ms, followed by 250 ms at the 20°size. **(A')** The static object was 20° in diameter and was presented for 500 ms. **(B-B')** Heatmap showing instantaneous firing rates of a single unit responding to 15 consecutive presentations of a looming object **(B)** or of a static object **(B'). (C-C')** PSTH

857   of the same single unit showing the mean firing rate for responses to the looming **(C)** and

858   static objects **(C')** across the 15 trials (mean $\pm$ SE).

Figure 7. Neurons respond differentially to looming and static objects.

(A) Cumulative probability for onset time comparing both visual stimulus types. Single units responded with similar onset time to both types (static object (red), looming object (black)). **(B)** Cumulative probability for the timing of peak activity comparing responses to looming (black) and static visual stimuli (red). Single units reach their peak activity significantly faster with the static object than with the looming object (p ≥ 0.05). **(C)** Time at maximal firing rate of neurons responding to both the looming and to the static object; note that most neurons fall above the identity line (slope = 1.0; red) indicating that their maximal firing occurs later in time when exposed to looming compared to the static object.

872   **(D)** Maximum firing rate to each visual stimulus of neurons that responded to both stimuli.

873   Interestingly, neurons show higher firing rates when exposed to looming stimuli.

 FIGURE LEGENDS

Figure 1. Visual stimulation, extracellular recording, and spike sorting. (A) Visual

stimuli used to characterize RF size and shape. A small white square (~3° diameter) on

a black background was swept across the screen monitor from top to bottom with each

vertical sweep (27°/s, 4 sec), displaced horizontally by 3°, at 1s intervals, to cover the

entire screen. **(B)** Top trace, extracellular responses in the SC. Bottom trace, the visual

stimulus on/off pattern as described in **A**. **(C)** Example of 3 different single units isolated

from recordings.

Figure 2. (A) Photomicrograph of a Nissl-stained coronal section showing the

subdivisions of the degus' SC: stratum griseum superficiale (SGS) upper (uSGS) and

deeper (dSGS), stratum opticum (SO), stratum griseum intermediale (SGI) upper (uSGI),

intermediate (iSGI) and deeper (dSGI), the stratum griseum profundus (SGP), and PAG,

periacueductal gray. **(A')** Detail of a Nissl-stained coronal section of SC laminae. **(B)**

Anterograde labeling of retinotectal projections after a CTB injection into the contralateral

eye. **(B')** Detail of the retinotectal projection into the different layers of the degus' SC.

Figure 3. Receptive field size and shape change across SC layers in degus. (A) Heat

map showing the RF of a single unit isolated at 320 μm depth in the SGS mapped by

visual stimulation as in Fig. 1A. **(B)** Heat map showing the RF of a unit at 1100 (μm) in

the SGI mapped with the same stimulation, at the same recording site (stereotaxic

coordinates) as in **A**, but deeper. **(C)** Boxplot of RF areas (in deg^2) across depths ranges

from the SGS to the SGI. Boxplots show standard interquartile range (IQR) and median

value (black central line), yellow dots are the individual data points. RFs increase in size with depth (ANOVA: *p* < 0.01). **(D)** Boxplot of RF areas across SC layers (SGS, SO and SGI). **(E)** Boxplot of RF shape expressed as the width/height ratio at different SC depth ranges. RFs widen along the horizontal axis with depth (ANOVA: *p* <0.01). **(F)** Boxplot of RF shape at different SC layers. Different letters above each group in B, C, D, and F represent statistically significant differences (Tukey post-hoc test, see text for exact *p*-values).

Figure 4. Single units show selectivity to spatial frequency. sSC neurons have a narrow bandwidth, and a small proportion of them (14%) respond linearly to sinusoidal gratings. **(A)** Illustration of the drifting grating stimulus types. Spatial frequency and contrast were varied in a pseudorandom manner. **(B)** Percentage of neurons with low-pass, band-pass or high-pass selectivity to different spatial frequencies (n = 128). **(C-E)** Response matrix (instantaneous firing rate, in Hz) of three different neurons to all contrast and spatial frequency combinations. **(C′-E′)** Examples of low-pass **(C′)**, band-pass **(D′)**, and high pass **(E′)** tuning of the same single neurons as in **C-E**. **(F)** Bar plot of spatial frequency preference for the population. Most units are tuned to 0.04 or 0.08 (cpd), and a smaller group is tuned to higher spatial frequency (0.24 cpd). **(G)** Bar plot of bandwidth showing that most neurons (>50%) are tuned to a certain spatial frequency with narrow bandwidths. **(H-I)** Solid lines are the mean firing rate (100 ms bins) and shaded regions represent mean ± sd. **(H)** Example of a single unit responding non-linearly to 4 seconds of stimulation with a sinusoidal grating (linearity index F1/F0 < 1, where *F1* is the response

at the drift frequency, and *F0* is the average firing rate). **(I)** Example of a different single

unit responding in a linear manner to sinusoidal gratings (F1/F0 >1). **(J)** Histogram of the

linearity index across the entire population of single units (n = 232) showing neurons

responding linearly to sinusoidal gratings (red bars) and neurons with non-linear

responses to sinusoidal gratings (blue bars).

Figure 5. SC Single units show linear or non-linear responses to contrast. Example

of three different single units' tuning to contrast, ranging from linear **(A)** to non-linear **(B**

and C). All three curves are fitted with a Naka-Rushton function (black line). The dotted

line is the spontaneous firing rate (Hz) and the dashed line is the C_{50} (the contrast at half

the peak response)**.** Note that **(B)** and **(C)** show saturating responses and differ in their

C_{50} values (~40 and ~20, respectively). **(D)** Cumulative histogram of C_{50} distribution

(N=135), with the median C_{50} = 48. **(E)** Cumulative histogram (N=135) of the relative

maximum gain (see Methods).

Figure 6. Single units show a continuous buildup of response level to looming

stimuli but an ON-OFF response to static stimuli. (A) Diagram of visual stimulation.

The looming object increased in size from 2° to 20° over 250 ms, followed by 250 ms at

the 20°size. **(A')** The static object was 20° in diameter and was presented for 500 ms. **(B-**

B') Heatmap showing instantaneous firing rates of a single unit responding to 15

consecutive presentations of a looming object **(B)** or of a static object **(B'). (C-C')** PSTH

of the same single unit showing the mean firing rate for responses to the looming **(C)** and

static objects **(C')** across the 15 trials (mean $\pm$ SE).

Figure 7. Neurons respond differentially to looming and static objects.

(A) Cumulative probability for onset time comparing both visual stimulus types. Single units responded with similar onset time to both types (static object (red), looming object (black)). **(B)** Cumulative probability for the timing of peak activity comparing responses to looming (black) and static visual stimuli (red). Single units reach their peak activity significantly faster with the static object than with the looming object ($p \geq 0.05$). **(C)** Time at maximal firing rate of neurons responding to both the looming and to the static object; note that most neurons fall above the identity line (slope = 1.0; red) indicating that their maximal firing occurs later in time when exposed to looming compared to the static object. **(D)** Maximum firing rate to each visual stimulus of neurons that responded to both stimuli. Interestingly, neurons show higher firing rates when exposed to looming stimuli.