Skull evolution and lineage diversification in endemic Malagasy carnivorans

Chris J. Law[1,2], Tate J. Linden[2], and John J. Flynn[3]

[1]Department of Integrative Biology, University of Texas

[2]Burke Museum and Department of Biology, University of Washington

[3]Division of Paleontology, American Museum of Natural History

cjlaw@uw.edu; ORCID ID: 0000-0003-1575-7746

jflynn@amnh.org; ORCID ID 0000-0003-4705-3591

Abstract

Madagascar is one of the world's foremost biodiversity hotspots with more than 90% of

its species endemic to the island. Malagasy carnivorans are one of only four extant terrestrial

mammalian clades endemic to Madagascar. Although there are only eight extant species, these

carnivorans exhibit remarkable phenotypic and ecological diversity that is often hypothesized to

have diversified through an adaptive radiation. Here, we investigated the evolution of skull

diversity in Malagasy carnivorans and tested if they exhibited characteristics of convergence and

an adaptive radiation. We found that their skull disparity exceeds that of any other feliform

family, as their skulls vary widely and strikingly capture a large amount of the morphological

variation found across all feliforms. We also found evidence of shared adaptive zones in cranial

shape between euplerid subclades and felids, herpestids, and viverrids. Lastly, contrary to

predictions of adaptive radiation, we found that Malagasy carnivorans do not exhibit rapid

lineage diversification and only marginally faster rates of mandibular shape evolution, and to a

lesser extent cranial shape evolution, compared to other feliforms. These results reveal that

exceptional diversification rates are not necessary to generate the striking phenotypic diversity

that evolved in carnivorans after their dispersal to and isolation on Madagascar.

Key terms: Adaptive radiation; Carnivora; craniomandibular; Madagascar; phenotypic

evolution; phylogenetic comparative methods

Introduction

Many islands are considered as biodiversity hotspots [1] and contain species that are not

found anywhere else in the world [2–4]. High rates of endemism could be spurred by adaptive

radiations [5]. Islands in particular, which are often geographically isolated, provide novel

ecological opportunities for founding species to rapidly diversify to a variety of species and

phenotypes to fill previously unoccupied ecological niches [6–8]. Unsurprisingly, the best-

studied adaptive radiations are found among insular island groups such as Galápagos finches,

Hawaiian honeycreepers, and Caribbean anoles [9–11]. One of the world's foremost biodiversity

hotspots is Madagascar, with more than 90% of its species endemic to the island [12]. As

Madagascar has been geographically isolated for over 88 million years, most endemic Malagasy

vertebrates are a result of several independent successful dispersal events from Africa or Asia

[13]. Upon arrival on Madagascar, adaptive radiations are hypothesized to have facilitated the

diversification of many Malagasy clades, such as beetles [14,15], frogs [16,17], vangas [18], and

lemurs [19–21]. In this study, we investigate the diversity of Malagasy euplerid carnivorans, one

of four extant terrestrial mammalian clades (i.e., Lemuroidea, Eupleridae, Tenrecidae and

Nesomyinae) endemic to Madagascar and hypothesized to have diversified through an adaptive

radiation [22].

Although there are only eight extant species, Malagasy carnivorans exhibit remarkable

phenotypic and ecological diversity [22–25], so much so that no anatomical character can be

used to define them. As a result, Malagasy carnivorans have traditionally been thought to belong

to or originate from multiple feliform families (i.e., Herpestidae, Viverridae, and Felidae) and

thus presumably dispersed to Madagascar through multiple independent dispersal events [26–

29]. Molecular data has since revealed that all Malagasy carnivorans originated from a single

ancestor, thus forming a monophyletic clade and most likely exhibited a single dispersal event to

Madagascar [30–32]. Eupleridae consists of three major clades: Galidiinae, Euplerinae, and the

fossa (*Cryptoprocta ferox*) (Fig. 1). Galidiines are comprised of five species of vontsira. Their

superficial resemblance to mongooses (Herpestidae) have led to their "Malagasy mongoose"

alias and even to their classification as herpestids [26,27]. Galidiines exhibit diverse ecologies

such as in diets that range from insectivory, similarly found in herpestids, to more carnivorous

diets in *Galidictis* species [22]. Euplerines superficially resemble civets (Viverridae), leading to

them having been described as "Malagasy civets" and classified as viverrids [28]. Euplerines

also exhibit diverse ecologies ranging from the omnivorous fanaloka (*Fossa fossana*) that feeds

on a variety of invertebrate and vertebrate prey, similar to viverrids, to the Eastern falanouc

(*Eupleres goudotii*) that specialize on soft-bodied invertebrates such as worms and slugs [22].

The last euplerid clade consists of the fossa, the largest and most carnivorous of the extant

euplerids [33]. Its large-bodied civet-like and carnivorous cat-like resemblances have historically

led its placement either within Viverridae or Felidae [28,29].

Here, we investigated the morphological disparity among Malagasy euplerids and in

comparison to taxa in other feliform carnivoran groups with which they have been historically

classified. We used skull shape in our examination of phenotypic evolution because of its strong

associations with dietary ecology across carnivorans [34–37]. Because of their superficial

resemblances, we also investigated convergence among euplerid and feliform clades with which

they have been classified historically. We predict that euplerids will exhibit greater disparity

compared to other feliform clades and that the three euplerid clades—galidiines, euplerines, and

the fossa—will converge towards herpestid-, viverrid-, and felid-like morphotypes in skull

morphology, respectively. The observed phenotypic and ecological diversity found in euplerids

suggest that they exhibited an adaptive radiation. Therefore, we also test if rates of their lineage

diversification and phenotypic evolution are significantly accelerated relative to other clades as

predicted under adaptive radiation. We predict that euplerids will exhibit shifts towards faster

rates of lineage diversification and phenotypic evolution compared to other feliform clades.

Methods

Morphological data

We quantified the shapes of 110 crania across 57 feliforms (~50% of species diversity)

and the shapes of 315 mandibles across 90 feliforms (~79% of species diversity) using 3D

geometric morphometrics [38,39]. Our dataset encompasses all eight euplerid species. Whether

Eupleres major and *Salanoia durrelli* are true species remains debated [40,41]. Shape data were

collected from 3D scans of feliform skulls obtained from surface scanning with an EinScan Pro

HD scanner at various museums, an Artec Spider structured light 3D scanner at the American

Museum of Natural History, or from previously published work [35] (see Table S1 for list of

specimens and museums). All specimens were fully mature, determined by the closure of

exoccipital–basioccipital and basisphenoid–basioccipital sutures on the cranium and full tooth

eruption. Although carnivorans exhibit sexual dimorphism [42,43], a combination of females,

males, and sex unknown individuals was used because just one sex cannot be used without

compromising the inclusion of as many euplerid species as possible. We used 35 landmarks and

seven curves with 134 semi-landmarks for the cranium, and 21 landmarks and four curves with

24 semi-landmarks for the mandible (Fig. S1). Landmarks were digitized using Slicer, and

curves were digitized by oversampling semi-landmarks in Slicer. Landmarks were superimposed

by Generalized Procrustes analysis, and semi-landmarks on the curves were allowed to slide

along their tangent vectors until their positions minimized bending energy [39,44]. As part of the

superimposition procedure, bilaterally homologous landmarks and semi-landmarks were

reflected across the median plane and averaged using the geomorph function bilat.symmetry. All

Procrustes superimpositions were performed in the R package geomorph v4.0.6 [45]. We used

centroid size as our metric of cranial and mandibular size.

Craniomandibular shape analyses

We performed all analyses under a phylogenetic framework using the most recent

phylogeny of mammals, pruned to include just feliform carnivorans [46]. All analyses were

performed in R v4.1.1 (R Core Team 2021). Because allometry has been shown to facilitate or

constrain skull shape evolution [47], we first tested for evolutionary allometry on cranial and

mandibular shape by performing a phylogenetic Procrustes regression [48] with a random

residual permutation procedure (1,000 iterations) using the geomorph function procD.pgls. Both

cranial shape (SS = 0.022, MS = 0.022, R^2 = 0.20, F = 14.127, Z = 3.45, P < 0.001) and

mandibular shape (SS = 0.044, MS = 0.044, R^2 = 0.25, F = 29.360, Z = 5.07, P = 0.001)

exhibited significant evolutionary allometry. We, therefore, extracted allometry-free shape as the

shape residuals from the phylogenetic Procrustes regressions and used them in all subsequent

analyses.

Craniomandibular shape disparity. We visualized the phylomorphospace of cranial and

mandibular shape by performing principal component analyses (PCA) using the geomorph

function gm.prcomp. We then examined if cranial and mandibular disparity (i.e., Procrustes

variance) differed among the feliform clades using the geomorph functions procD.plgs and

morphol.disparity.

Testing for convergence towards different feliform morphotypes. Euplerids are often cited

to superficially resemble and were historically classified together with other feliforms,

specifically galidiines with mongooses (Herpestidae), euplerines with civets (Viverridae), and

the fossa with cats (Felidae). Therefore, we tested whether there was statistical evidence for

convergence in these various euplerid-feliform comparisons. We assessed convergence using the

Ct-measure in the R package convevol [49,50], which tests whether the geometric distance

between focal lineages in phylomorphospace are shorter than the distances between ancestral

nodes at a specific time point. Second, we used a frequency-based measure of convergence (C_5),

which determines whether the number of lineages that crosses into a particular region of

morphospace is greater than expected. We conducted 100 simulations under Brownian motion to

assess statistical significance of Ct and C_5 measures.

We also tested whether each of the three groups exhibited shared adaptive zones in skull

shapes by fitting five multivariate evolutionary models [51,52] on the first three PCs of the

cranial shape dataset (accounting for 74.44% of total cranial shape variation) and mandibular

shape dataset (87.4% of total mandibular shape variation) using the R package mvMORPH

version 1.1.7 [53] to incorporate covariances between axes. These models included single-rate

Brownian motion model (mvBM1), single-peak Ornstein–Uhlenbeck model (mvOU1), and three

two-peak Ornstein–Uhlenbeck models (mvOUM). Each of these mvOUM models designated

two optima, one optimum for the focal comparison and a second optimum for the remaining

feliforms. All models were fit across 250 mapped trees to account for uncertainty in phylogenetic

topology and stochastic character maps of group designations. Models were assessed with small

sample corrected Akaike weights (AICcW), and all models with a ∆AICc < 2 were considered

best-fitting models. Support for a mvOUM model as a best-fitting model would suggest a shared

adaptive zone between the focal clades of interest. These results, however, also would provide

more nuanced interpretations of the convergent evolution because they are unable to distinguish

between potential convergence or conservatism [49,50].

Craniomandibular shape evolutionary rates. We investigated rates of cranial shape and

mandibular shape evolution using two approaches. First, we tested if evolutionary rates in cranial

shape and mandibular shape differed among the feliform clades using the geomorph function

compare.evol.rates. Second, we examined branch-specific evolutionary rates and rate shifts of

cranial shape and mandibular shape using a variable rates model under a reversible-jump Markov

Chain Monte Carlo (rjMCMC) framework in BayesTraits v4.1.1

(http://www.evolution.rdg.ac.uk/). To reduce the dimensions of the shape data for BayesTraits

analysis, we used only the phylogenetic principal component scores (pPC) scores that represent

>95% of the shape variation in the cranium (first 18 pPCs) and mandible (first 11 pPCs). We ran

two independent chains, each with 200,000,000 iterations sampled every 20,000 iterations. After

the first 25,000,000 iterations were discarded as burn-in, we assessed convergence of the two

chains by checking the trace plots and then the effective sample size (ESS) with Gelman and

Rubin's diagnostics in the R package coda v0.19-4. We plotted rates shifts and branch-specific

rates across the feliform phylogeny and constructed density plots to compare evolutionary rates

among feliform families using custom functions from [54].

Rates of lineage diversification

169   We estimated speciation and extinction rates through time and across the feliform

phylogeny using two methods, Bayesian Analysis of Macroevolutionary Mixtures (BAMM) [55]

and the Lineage Specific Birth–Death–Shift (LSBDS) model ([56]. We performed both

diversification analyses using the most recent phylogeny of mammals, pruned to include just

feliform carnivorans [46]. BAMM applies a reversible jump Markov Chain Monte Carlo

(RJMCMC) to explore candidate models of lineage diversification as well as quantify

heterogeneity in evolutionary rates. We performed two independent BAMM runs of five million

generations on feliform phylogeny, sampling every 1,000 generations and with priors chosen

using the R package BAMMtools v2.0 [57]. We assessed the convergence of each BAMM run

using the R package BAMMtools v2.1.11 (Rabosky et al. 2014). The LSBDS model samples rate

regimes from a prior distribution, discretized into a fixed number of rate categories. We

implemented the LSBDS model using seven categories (one for each of the seven extant feliform

families) for speciation and extinction, and ran two MCMC chains for 5,000 generations in

RevBayes [58]. We merged the posteriors, retaining the last 4,000 generations from the MCMC

in the R package RevGadgets [59]. Convergence for both methods was assessed by checking that

the ESS values for all model parameters in the log files were greater than 200, using the R

package coda [60].

Results

Craniomandibular shape morphospace and disparity

PCs 1–3 explain 74.44% of the cranial shape variation across feliforms (Fig. 2). PC 1

distinguishes hypercarnivorous felids and hyaenids (negative PC 1) from the remaining, less

carnivorous feliforms including euplerids (positive PC 1). Positive PC 1 is characterized by

lateral narrowing of the cranium at the zygomatic arches, expansion of the braincase through

widening of the nuchal crests, and reduction of the sagittal crest resulting in dorsoventral

flattening. The diversity of euplerid cranial shapes is best captured by PC 2, as euplerid species

occupy the full range of PC 2. Positive PC 2 describes shortening of the rostrum and broadening

of the cranium at the zygomatic arches and nuchal crests. Galidiines are associated with positive

PC 2 and are encompassed within the range of variation of herpestids (and felids), whereas

euplerines are associated with negative PC 2 with *Fossa fossana* falling within the range of

viverrids (and hyaenids) but *Eupleres goudotii* distinct from all other feliforms. In PC2,

Cryptoprocta is intermediate between galidiines and euplerines (closer to the former), and falls

within the broad range of values of felids. In the total PC1/2 cranial morphospace, galidiines

cluster with herpestids, euplerines are closest to and overlap with viverrids, and *Cryptoprocta* is

distinct from other feliforms. Euplerids show little variation along cranial PC 3, falling entirely

within the range of felids and overlapping with the lower range of viverrids and upper range of

herpestids. All other non-monotypic feliform families show a greater range of variance than

and/or do not overlap with euplerids in PC3. Positive PC 3 describes slight elongation and

ventrodorsal flexion of the rostrum and slight broadening of the zygomatic arches. Ancestral

node reconstruction suggests that euplerids exhibit a cranial shape between viverrids and

herpestids (Fig. 3). Overall, euplerids exhibit significantly greater cranial shape disparity

(Procrustes variance = 0.0078) compared to felids (0.0047, P = 0.026) and viverrids (0.0033, P =

0.004), but not hyaenids (0.0066, P = 0.595) or herpestids (0.0061, P = 0.299).

Euplerids occupy greater regions of mandibular morphospace when compared to cranial

morphospace. PCs 1–3 explain 87.4% of the mandibular shape variation across feliforms (Fig.

2). Like in the crania, PC 1 of mandibular shape largely distinguishes hypercarnivorous felids,

and to a lesser extent hyaenids (both with negative PC 1 values), from the remaining, less

carnivorous feliforms (positive PC 1), except for overlap with hyaenids of two species of

Galidictis (galidiine euplerids) and one herpestid. In mandibular PC1, euplerines fall within the

range of viverrids and a few herpestids at the extreme positive end of their range of; galidiines

have a wide range of mandibular PC1 values, overlapping with herpestids at the most negative

end of their range of variation, hyaenids, and one species each of felid and viverrid;

Cryptoprocta is very close to hyaenids in PC1 value. Positive PC 1 describes dorsoventral

mandibular flexure and elongation, decrease in coronoid height, and lateral compression of the

mandibular body. However, unlike in the crania, euplerids exhibit greater variation along PC 1

than in any other feliform family (although viverrids are similar in range of variation, due to one

extreme outlier, the binturong [*Arctictis binturong*]), and spans negative and positive PC 1.

Euplerids also approach both the negative and positive ends of variation in both PCs 2

and 3, with *Eupleres goudotii* again having the most extreme negative value for PC2 of any

feliform (as for the cranial PCA). Positive PC 2 describes anteroposterior shortening of the

mandibular body, increases in coronoid height, and lateral broadening of the mandibular body;

positive PC 3 describes anteroposterior elongation of the mandibular body, dorsoventral

compression of the mandibular body, and reduction in coronoid height. Ancestral node

reconstruction suggests that euplerids exhibit a cranial shape between viverrids and herpestids,

with closer resemblance to the latter (Fig. 3). Euplerids exhibit significantly greater mandibular

shape disparity (Procrustes variance = 0.0100) compared to felids (0.0035, P = 0.001), viverrids

(0.0054, P = 0.020), and herpestids (0.0040, P = 0.003), but not hyaenids (0.0050, P = 0.692).

Convergence towards different feliform morphotypes

We did not find statistical support for the distance-based measure of convergence

between galidiines and herpestids (cranial Ct_1 = -0.19, P = 0.96; mandibular Ct_1 = -0.03, P =

XX), between euplerines and viverrids (cranial Ct_1 = -0.24, P = 0.84; mandibular Ct_1 = -0.25, P =

XX), or between the fossa (*Cryptoprocta ferox*) and felids (cranial Ct_1 = 0.05, P = 0.08;

mandibular Ct_1 = -0.05, P = 0.48). Similarly, the frequency-based measure of convergence

indicated that the number of transitions into regions of phylomorphospace occupied by

herpestids, viverrids, and felids was not significantly greater than expected under Brownian

motion (all P > 0.380).

Using evolutionary modeling, we found evidence of shared adaptive zones in cranial

shape between galidiines and herpestids (best model = mvOUM, AICcW = 0.98) and partial

evidence of shared adaptive zones in cranial shape between euplerines and viverrids (best models

= mvOUM [AICcW = 0.52] and mvBM [AICcW = 0.47; ΔAICc = 0.20]) and between the fossa

and felids (best models = mvBM [AICcW = 0.66] and mvOUM [AICcW = 0.32; ΔAICc = 1.47])

(Table S2). In contrast, we found no evidence of shared adaptive zones in mandibular shape in

all three comparisons (best model for all = mvOU1 [AICcW > 081]; Table S2).

Rates of craniomandibular shape evolution

Rates of cranial shape evolution in euplerids (ln σ^2_{multi} = -12.86) do not significantly

differ from most feliform clades (felid ln σ^2_{multi} = -12.94, P = 0.592; hyaenid ln σ^2_{multi} = -12.47, P

= 0.100; herpestid ln σ^2_{multi} = -13.14, P = 0.105), except for viverrids (ln σ^2_{multi} = -13.72, P =

0.001), in which euplerids exhibited significantly faster rates of cranial shape evolution.

Although we found six significant rate shifts (posterior probability (pp) > 0.60) across feliforms,

none of these occurred within euplerids (Fig. 3). Instead, rate shifts occurred towards or within

the branches of the other major feliform clades including Hyaenidae, Felidae (pantherines and

the *Puma* lineage), Viverridae (crownward *Genetta* species), Herpestidae (subclade of

herpestines), and the overall feliform clade excluding Nandiniidae (Fig. 3).

Euplerids exhibited greater rate heterogeneity in evolution of mandibular shape compared

to cranial shape. Euplerids exhibited significantly faster rates of mandibular shape evolution (ln

σ^2_{multi} = -12.29) compared to viverrids (ln σ^2_{multi} = -13.08, P = 0.001) and herpestids (ln σ^2_{multi} = -

12.79, P = 0.001) but significantly slower rates compared to felids (ln σ^2_{multi} = -11.80, P = 0.023).

There was no difference in mandibular shape evolutionary rates between euplerids and hyaenids

(ln σ^2_{multi} = -12.23, P = 0.889). Furthermore, we found that two of the five significant rate shifts

in mandibular shape (pp > 0.70) occurred within euplerid clades: *Galidictis* species and

euplerines (Fig. 3). The remaining rate shifts occurred within the same subclade of herpestines

that exhibited a rate shift in cranial shape evolution, within *Leopardus* cats, and for the overall

feliform clade excluding Nandiniidae (Fig. 3).

Diversification rate analyses

Euplerids do not exhibit exceptional rates of diversification, as both BAMM and the

LSBDS model are united in revealing that there are no increased species diversification rates on

the branches leading towards Eupleridae (Fig. 4; Fig. S1). Furthermore, we found no significant

shifts in diversification rate within feliforms, albeit there are slightly elevated rates on the

branches leading towards to the viverrid subfamily Genettinae and felids (Fig. 4; Fig. S1).

Discussion

Skull diversity in Malagasy carnivorans

Euplerids exhibit greater disparity in their skulls compared to most feliform families, for

both cranial and mandibular shapes, and particularly so in their mandibles. Euplerids overlap

with all extant feliform families in mandibular shape morphospace, whereas they only overlap

with non-hyercarnivorous families (i.e., Herpestidae, Nandiniidae, Prionodontidae, and

Viverridae) in cranial shape morphospace (Fig. 2). The only euplerid to approach

hypercarnivorous feliforms (Felidae, Hyaenidae) in both cranial and mandibular morphospace is

the fossa (*Cryptoprocta ferox*), the most carnivorous of the euplerids [22]. The fossa exhibits

cranial and mandibular morphologies that appear intermediate between the average euplerid and

average felid (Fig. 5A, 6A). Compared to all other euplerids, the fossa exhibits relatively broader

zygomatic arches and more pronounced sagittal and nuchal crests in the cranium (Fig. 5A) and

relatively broader coronoid processes in the mandible (Fig. 6A). These adaptations suggest that

the fossa exhibits larger temporalis and masseter jaw muscles for increased biting ability [61–64]

and are thus better adapted for their hypercarnivorous diets [22]. There is some evidence that the

fossa and felids share a similar adaptive zone in cranial shape (ΔAIC = 1.47; Table S2), but there

is no statistical support for convergence in skull shape between the fossa and felids. These results

suggest that cranial adaptations towards hypercarnivory in the fossa do not quite reach the felid

morphotype (Fig. 5A, 6A) because the fossa and felids share an adaptive zone characterized by

broad adaptive slopes rather than distinct adaptive peaks with steep slopes [24,50,65,66].

Euplerines occupy overlapping regions of cranial and mandibular morphospace with

viverrids, the family to which they have been compared or classified with historically (Fig. 2).

The fanaloka (*Fossa fossana*) exhibits only slight differences in skull shape (i.e., narrower and

more elongate crania and mandibles) compared to the average viverrid, which is consistent with

their similar diets of invertebrate and vertebrate prey, whereas the falanouc (*Eupleres goudotii*)

exhibits a more specialized skull that includes relatively elongate rostrum and mandible and less

pronounced zygomatic arches, nuchal crest, sagittal crests, coronoid processes compared to the

average viverrid (Fig. 5B) and all other feliforms. These adaptations in the skull, along with

reduced dentition, enable the falanouc to specialize in feeding on soft-bodied invertebrates such

as worms and slugs [22]. We found evidence that euplerines and viverrids share an adaptive zone

in cranial shape but not mandibular shape (Table S2), but neither skull component provided

statistical support for convergence between the two groups. Therefore, like for the fossa and

felids, our results suggest that euplerines and viverrids may share an adaptive zone with broad

adaptive slopes instead of exhibiting convergence characterized by distinct similar adaptive

peaks with steep slopes.

Galidiines exhibit distinct patterns of cranial and mandibular shape distribution in

morphospace. Galidiines occupy overlapping regions of cranial morphospace with herpestids

(Fig. 2), and there appear to be few cranial shape differences between the galidiine species

closest to the average cranial shape of all galidiines and the herpestid species closest to the

average cranial shape of all herpestids (Fig. 5C). Evolutionary modeling also strongly supports a

shared adaptive zone in cranial shape between galidiines and herpestids (Table S2) but no

statistical support for convergence. However, galidiines approach regions of mandibular

morphospace occupied by hypercarnivorous feliforms (Fig. 2) and exhibit mandibular

morphologies that appear intermediate between herpestids and felids (Fig. 6C, D): galidiine

mandibles tend to be relatively broader and more robust compared to the herpestid species

closest to the average mandibular shape of all herpestids, but are not as broad or robust as the

felid species closest to the average mandibular shape of all felids (Fig. 6C, D). Of the galidiines,

Galidictis species appear most similar to felids in mandibular shape, providing morphological

evidence that carnivory likely is an important component of their diet. Previous work surmised

that in addition to insects, *Galidictis* feed on rodents, reptiles, amphibians, and even small

lemurs; however, their dietary ecologies remain poorly known, as they are some of the rarest

carnivorans in the world [22]. That *Galidictis* exhibits a rate shift towards faster mandibular

evolution (Fig. 3) leads us to postulate that shifts towards more carnivorous diets may be due to

more relatively recent (< 5 million years ago) selection. The distinct patterns between the

cranium and mandible in galidiines follow an overall trend of decoupled modes of evolution

between the cranium and mandible across carnivorans, where cranial evolution tends to follow

clade-based shifts whereas mandibular evolution is linked instead to broad dietary regimes

[35,67]. Our results suggest that galidiines may have retained cranial morphology from euplerid-

herpestid ancestors through conservatism rather than convergence whereas the more evolutionary

labile mandible adapted towards a more carnivorous diet than the typical insectivorous diets

found in herpestids, most euplerids, and most likely their most recent common ancestors (of the

euplerid + herpestid clade).

Are Malagasy carnivorans an adaptive radiation?

Like many endemic Malagasy clades, euplerids are often considered to represent an

adaptive radiation after dispersal to a long-isolated island. We found that euplerids show extreme

ranges of skull shape variation and many distinct cranial and mandibular shapes in

phylomorphospace and geometric morphometric analyses. Contrary to predictions of adaptive

radiation, we found that euplerids do not exhibit exceptional lineage diversification and have

only marginally faster rates of mandibular shape evolution and, to a lesser extent, cranial shape

evolution compared to other feliform families (Fig. 3, 4). Thus, it is tempting to postulate that

euplerids may have evolved via an "adaptive non-radiation" [8], where ecomorphological

divergence within a clade fails to produce rapid species diversification. Rapid phenotypic

evolution without rapid taxonomic diversification rates have been found in a variety of vertebrate

clades [68–70] including other carnivorans such as mustelids [71] as well as other endemic

Malagasy vertebrates such as mantellid frogs [8]. Those studies cited possible factors that may

mask a signal of rapid diversification [8,68–71] including the absence of fossil data in

diversification analyses, which may have erased signatures of early rapid diversification [72–74];

continental radiations, in which many clades are simply unable to rapidly radiate spatially across

entire continents as soon as they arise [69,71,75–77]; and clade age, where younger clades

diversify faster than older clades [78]. These factors also may influence our analyses of euplerid

diversification. First, euplerids are not an exceptionally young clade, having diverged from

herpestids ~24–18 million years ago [30] with a crown clade age of 15–14 million years old [46].

Despite their long evolutionary history, euplerids are extremely poorly represented in the fossil

record, known only from Holocene subfossil remains of the extinct giant fossa (*Cryptoprocta*

spelea) [79]. Therefore, unaccounted cladogenesis of extinct euplerids may lead to

underestimation of taxonomic diversification rates in euplerids. Second, unaccounted speciation

of modern euplerids also may lead to underestimation of diversification rates. At least two

euplerids—*Eupleres goudotii* and *Salanoia concolor*—are thought, though with uncertainty, to

each consist of two species (i.e., *E. goudotii* and *E. major* and *S. concolor* and *S. durrelli*)

[40,41,80]; if valid, this would lead to increased diversification rates, albeit at more crownward

tips of the tree. Lastly, euplerid diversification is perhaps better characterized as a continental

radiation rather than as an insular radiation. Researchers often suggest that Madagascar, as the 4th

largest island globally, resembles a continent more than an island, with a geological record that

spans more than three billion years and several vicariance and dispersal events over the past ~88-

90 million years, after its final separation from other major Gondwanan landmasses [81–83].

Therefore, the diversification of Malagasy clades with long residence times in Madagascar may

be better characterized as multiple bursts of speciation events throughout their evolutionary

history [19,84] rather than as a single early burst of rapid diversification that has been found in

some other insular clades [9–11]. That diversification rates do not differ statistically between

euplerids and continental feliforms support the hypothesis that the diversification dynamics of

other endemic Malagasy clades follow a pattern more similar to diversification on a continent

than on an island [8]. Therefore, while found only on Madagascar, there is no indication that

insular processes spurred a rapid burst of taxonomic diversification of euplerids, as often has

been hypothesized.

Conclusion

Euplerids exhibit great disparity in cranial and mandibular morphology that corresponds

to their wide dietary diversity ranging from worm-eating falanoucs to lemur-eating fossas. We

document for the first time that their skull morphologies, particularly their mandibles, show

significantly greater shape disparity compared to other feliform clades, and this single small

clade of Malagasy endemic carnivorans captures a large amount of the morphological variation

that is found across the entire clade of feliforms. We also found evidence that the three euplerid

clades—galidiines, euplerines, and the fossa—exhibited shared adaptive zones in cranial shape

morphology with herpestids, viverrids, and felids, respectively. We did not find statistical

evidence, however, that these three euplerid clades—galidiines, euplerines, and the fossa—

converged towards herpestid-, viverrid-, and felid-morphotypes in skull morphology,

respectively, as had been implicit in classifications or directly hypothesized historically. Despite

demonstrating great skull disparity, we found that euplerids do not clearly exhibit an adaptive

radiation, as we found no evidence of rapid lineage diversification rates and only marginally

faster rates of mandibular shape evolution compared to other feliform families. This work

supports an increasing number of studies demonstrating that adaptive radiations in endemic

Malagasy clades may not as common as previously thought [8,19] and diversification dynamics

on Madagascar may resemble those found on continents rather than on islands.

Acknowledgements

We thank C. Merrill for scanning of euplerid specimens at AMNH, and R. Andrade Luna, E.

Blackwell, F. Levy, and J. Padro for helping scan and landmark specimens. We are grateful to

the staff of and collections at the American Museum of Natural History, California Academy of

Sciences, Field Museum of Natural History, Natural History Museum of Los Angeles County,

Museum of Vertebrate Zoology, Natural History Museum London, National Museum of Natural

History, and Burke Museum of Natural History and Culture.

Funding

Funding was supported by the National Science Foundation (DBI–2128146), a University of

Texas Early Career Provost Fellowship, and Stengl-Wyer Endowment Grant (SWG-22-02) to

CJL.

Ethics

This work did not require ethical approval from a human subject or animal welfare committee.

Data accessibility

All data and original code are available from the Dryad Digital Repository: XXXX

Declaration of AI use

We have not used AI-assisted technologies in creat- ing this article.

Conflict of interest declaration

We declare we have no competing interests

Figures (with Captions)

Fig. 1. Phylogeny of Eupleridae and other extant feliform Carnivora families. Oligo = Oligocene;

Pli = Pliocene; Ple = Pleistocene. Phylogeny based on [46].

Fig. 2. Morphospace of allometry-free cranial (left) and mandibular (right) shape defined by

principal component (PC) axes 1–3. Taxa illustrated for the cranial morphospace: (a) Brown

hyena (*Hyaena brunnea* [Hyaenidae]), –PC1; (b) Javan mongoose (*Urva javanicus*

[Herpestidae]), +PC1; (c) Pallas's cat (*Otocolobus manul* [Felidae]), +PC2; (d) Eastern falanouc

(*Eupleres goudotii* [Eupleridae]), –PC2; (e) Asian palm civet (*Paradoxurus hermaphroditus*

[Eupleridae]), +PC3; and (f) Egyptian mongoose (*Herpestes ichneumon* [Herpestidae]), –PC3.

Taxa illustrated for the mandibular morphospace: (g) tiger (*Panthera tigris* [Felidae]), –PC1; (h)

small Indian civet (*Viverricula indica* [Viverridae]), +PC1; (i) Ethiopian dwarf mongoose

446   (*Helogale hirtula* [Herpestidae]), +PC2; (j) Eastern falanouc (*Eupleres goudotii* [Eupleridae]), –

447   PC2; (k) Large Indian civet (*Viverra zibetha* [Viverridae]), +PC3; and (l) Binturong (*Arctictis*

448   *binturong* [Viverridae]), –PC3. Grey circles indicate estimated ancestral shapes at nodes, and the

449   green star indicates estimated ancestral shape of the ancestral euplerid node.

452
453 **Fig. 3.** Phylorate plots of cranial and mandibular shape evolutionary rates across feliform

454 Carnivora. Warmer colors indicate faster rates and cooler colors indicate slower rates. Significant

455 shifts in evolutionary rate are shown as grey circles with sizes proportional to the posterior

456 probability. All rates were log-transformed.

Fig. 4. Phylorate plot of lineage diversification rates across feliform Carnivora using BAMM.

Colors at each point in time along the branches of the phylorate plot denote instantaneous rate of

diversification. Warmer colors indicate faster rates and cooler colors indicate slower rates. There

are no significant shifts in diversification rates across the phylogeny. Analyses of lineage

diversification rates using LSBDS resulted in similar results (Fig. S1).

Fig. 5. Thin-plate spline (TPS) deformation grids depicting cranial shape differences between euplerid clades and other feliform families. A) Cranial shape of the fossa (*Crytoprocta ferox*) appears intermediate between felids and other euplerids. B) Cranial shape of euplerines: the fanaloka (*Fossa fossana*) resembles viverrids whereas the Eastern falanouc (*Eupleres goudotii*) exhibits a more specialized morphology. C) Cranial shapes of most galidiines resemble herpestids.

Fig. 6. Thin-plate spline (TPS) deformation grids depicting mandibular shape differences

euplerid clades and other feliform families. A) Mandibular shape of the fossa (*Crytoprocta ferox*)

appears intermediate between felids and euplerids. B) Mandibular shape of euplerines: the

fanaloka (*Fossa fossana*) resemble viverrids whereas the Eastern falanouc (*Eupleres goudotii*)

exhibits a more specialized morphology. C) Mandibular shapes of most galidiines resemble

herpestids, although the mandibular shape (D) of Grandidier's vontsira (*Galidictis grandidieri*)

appears intermediate between felids and other galidiines.

References

1. Myers N, Mittermeier RA, Mittermeier CG, Fonseca GAB da, Kent J. 2000 Biodiversity hotspots for conservation priorities. *Nature* **403**, 853–858. (doi:10.1038/35002501)

2. Kier G, Kreft H, Lee TM, Jetz W, Ibisch PL, Nowicki C, Mutke J, Barthlott W. 2009 A global assessment of endemism and species richness across island and mainland regions. *Proc. Natl. Acad. Sci.* **106**, 9322–9327. (doi:10.1073/pnas.0810306106)

3. Losos JB, Ricklefs RE. 2009 Adaptation and diversification on islands. *Nature* **457**, 830–836. (doi:10.1038/nature07893)

4. Veron S, Haevermans T, Govaerts R, Mouchet M, Pellens R. 2019 Distribution and relative age of endemism across islands worldwide. *Sci. Rep.* **9**, 11693. (doi:10.1038/s41598-019-47951-6)

5. Stroud JT, Losos JB. 2016 Ecological Opportunity and Adaptive Radiation. *Annu. Rev. Ecol., Evol., Syst.* **47**, 507–532. (doi:10.1146/annurev-ecolsys-121415-032254)

6. Simpson GG. 1955 *Major features of evolution*. Columbia University Press. See http://krishikosh.egranth.ac.in/handle/1/2034793.

7. Schluter D. 2000 *The Ecology of Adaptive Radiation*. Oxford University Press. See http://books.google.com/books/content?id=Q1wxNmLAL10C&printsec=frontcover&img=1&zoom=1&edge=curl&source=gbs_api.

8. Moen DS, Ravelojaona RN, Hutter CR, Wiens JJ. 2021 Testing for adaptive radiation: A new approach applied to Madagascar frogs*. *Evolution* **75**, 3008–3025. (doi:10.1111/evo.14328)

9. Grant PR, Grant BR. 2007 *How and why species multiply: the radiation of Darwin's finches*. Princeton University Press.

10. Losos JB. 2011 *Lizards in an evolutionary tree: ecology and adaptive radiation of anoles*. University of California Press.

11. Lerner HRL, Meyer M, James HF, Hofreiter M, Fleischer RC. 2011 Multilocus Resolution of Phylogeny and Timescale in the Extant Adaptive Radiation of Hawaiian Honeycreepers. *Curr. Biol.* **21**, 1838–1844. (doi:10.1016/j.cub.2011.09.039)

12. Ralimanana H *et al.* 2022 Madagascar's extraordinary biodiversity: Threats and opportunities. *Science* **378**, eadf1466. (doi:10.1126/science.adf1466)

13. Yoder AD, Nowak MD. 2006 Has Vicariance or Dispersal Been the Predominant Biogeographic Force in Madagascar? Only Time Will Tell. *Annu. Rev. Ecol., Evol., Syst.* **37**, 405–431. (doi:10.1146/annurev.ecolsys.37.091305.110239)

14. Wirta H, Orsini L, Hanski I. 2008 An old adaptive radiation of forest dung beetles in
Madagascar. *Mol. Phylogenetics Evol.* **47**, 1076–1089. (doi:10.1016/j.ympev.2008.03.010)

15. Moore W, Robertson JA. 2014 Explosive Adaptive Radiation and Extreme Phenotypic
Diversity within Ant-Nest Beetles. *Curr. Biol.* **24**, 2435–2439. (doi:10.1016/j.cub.2014.09.022)

16. Valero KCW *et al.* 2017 Transcriptomic and macroevolutionary evidence for phenotypic
uncoupling between frog life history phases. *Nat. Commun.* **8**, 15213.
(doi:10.1038/ncomms15213)

17. Bossuyt F, Milinkovitch MC. 2000 Convergent adaptive radiations in Madagascan and Asian
ranid frogs reveal covariation between larval and adult traits. *Proc. Natl. Acad. Sci.* **97**, 6585–
6590. (doi:10.1073/pnas.97.12.6585)

18. Reddy S, Driskell A, Rabosky DL, Hackett SJ, Schulenberg TS. 2012 Diversification and the
adaptive radiation of the vangas of Madagascar. *Proc. R. Soc. B: Biol. Sci.* **279**, 2062–2071.
(doi:10.1098/rspb.2011.2380)

19. Everson KM *et al.* 2023 Not one, but multiple radiations underlie the biodiversity of
Madagascar's endangered lemurs. *bioRxiv* , 2023.04.26.537867.
(doi:10.1101/2023.04.26.537867)

20. Godfrey LR, Samonds KE, Baldwin JW, Sutherland MR, Kamilar JM, Allfisher KL. 2020
Mid-Cenozoic climate change, extinction, and faunal turnover in Madagascar, and their bearing
on the evolution of lemurs. *BMC Evol. Biol.* **20**, 97. (doi:10.1186/s12862-020-01628-1)

21. Herrera JP. 2016 Testing the adaptive radiation hypothesis for the lemurs of Madagascar. *R.*
Soc. Open Sci. **4**, 161014. (doi:10.1098/rsos.161014)

22. Goodman SM. 2009 Family Eupleridae. Madagascar carnivores. In *Handbook of the*
mammals of the world, Volume 1, Carnivores. (eds D Wilson, RA Mittermeier), pp. 330–351.
Barcelona, Spain: Lynx edicions.

23. Michaud M, Veron G, Peigné S, Blin A, Fabre A-C. 2018 Are phenotypic disparity and rate
of morphological evolution correlated with ecological diversity in Carnivora? *Biological Journal*
of the Linnean Society **124**, 294–307. (doi:10.1093/biolinnean/bly047/4994739)

24. Law CJ. 2022 Different Evolutionary Pathways Lead to Incomplete Convergence of Elongate
Body Shapes in Carnivoran Mammals. *Systematic Biol* **71**, 788–796.
(doi:10.1093/sysbio/syab091)

25. Law CJ. 2021 Evolutionary and morphological patterns underlying carnivoran body shape
diversity. *Evolution* **75**, 365–375. (doi:10.1111/evo.14143)

26. Lydekker R. 1894 *A hand-book to the Carnivora.. Part 1,, Cats, civets, and mungooses /.*
London: Allen. (doi:10.5962/bhl.title.9520)

27. Pocock RI. 1915 XLIV.—On some external characters of Galidia, Galidictis, and related
genera. *J. Nat. Hist. Ser. 8* **16**, 351–356. (doi:10.1080/00222931508693726)

28. Gregory WK, Hellman M. 1939 On the evolution and major classification of the civets
(Viverridae) and allied fossil and recent Carnivora: a phylogenetic study of the skull and
dentition. *Proceedings of the American Philosophical Society* , 309–392.

29. Bennett ET. 1833 Notice of a new genus of Viverridous Mammalia from Madagascar.
Proceedings of the Zoological Society of London **46**.

30. Yoder AD, Burns MM, Zehr S, Delefosse T, Veron G, Goodman SM, Flynn JJ. 2003 Single
origin of Malagasy Carnivora from an African ancestor. *Nature* **421**, 734–737.
(doi:10.1038/nature01303)

31. Yoder AD, Flynn JJ. 2003 Origin of Malagasy carnivora. In *The natural history of*
Madagascar, pp. 1253–1256.

32. Flynn JJ, Finarelli JA, Zehr S, Hsu J, Nedbal MA. 2005 Molecular Phylogeny of the
Carnivora (Mammalia): Assessing the Impact of Increased Sampling on Resolving Enigmatic
Relationships. *Syst. Biol.* **54**, 317–337. (doi:10.1080/10635150590923326)

33. Wilson DE, Mittermeier RA. 2009 *Handbook of the mammals of the world, Volume 1,*
Carnivores. Barcelona, Spain: Lynx Edicions. See http://www.sidalc.net/cgi-
bin/wxis.exe/?IsisScript=sibe01.xis&method=post&formato=2&cantidad=1&expresion=mfn=03
4105.

34. Law CJ, Duran E, Hung N, Richards E, Santillan I, Mehta RS. 2018 Effects of diet on cranial
morphology and biting ability in musteloid mammals. *J. Evol. Biol.* **31**, 1918–1931.
(doi:10.1111/jeb.13385)

35. Law CJ, Blackwell EA, Curtis AA, Dickinson E, Hartstone☐Rose A, Santana SE. 2022
Decoupled evolution of the cranium and mandible in carnivoran mammals. *Evolution* **76**, 2959–
2974. (doi:10.1111/evo.14578)

36. Figueirido B, Tseng ZJ, Martín-Serra A. 2013 Skull shape evolution in durophagous
carnivorans. *Evolution* **67**, 1975–1993. (doi:10.1111/evo.12059)

37. Meloro C, Clauss M, Raia P. 2015 Ecomorphology of Carnivora challenges convergent
evolution. *Org Divers Evol* , 1–10. (doi:10.1007/s13127-015-0227-5)

38. Rohlf FJ, Slice D. 1990 Extensions of the Procrustes method for the optimal superimposition
of landmarks. *Syst. Zool.* **39**, 40–21. (doi:10.2307/2992207)

39. Zelditch ML, Swiderski DL, Sheets HD. 2012 *Geometric morphometrics for biologists: a*
primer. Academic Press.

40. Veron G, Goodman SM. 2018 One or two species of the rare Malagasy carnivoran Eupleres
(Eupleridae)? New insights from molecular data. *Mammalia* **82**, 107–112.
(doi:10.1515/mammalia-2016-0182)

41. Durbin J, Funk SM, Hawkins F, Hills DM, Jenkins PD, Moncrieff CB, Ralainasolo FB. 2010
Investigations into the status of a new taxon of Salanoia (Mammalia: Carnivora: Eupleridae)
from the marshes of Lac Alaotra, Madagascar. *Syst. Biodivers.* **8**, 341–355.
(doi:10.1080/14772001003756751)

42. Law CJ. 2019 Solitary meat-eaters: solitary, carnivorous carnivorans exhibit the highest
degree of sexual size dimorphism. *Sci Rep-uk* **9**, 15344. (doi:10.1038/s41598-019-51943-x)

43. Law CJ, Mehta RS. 2018 Carnivory maintains cranial dimorphism between males and
females: evidence for niche divergence in extant Musteloidea. *Evolution* **72**, 1950–1961.
(doi:10.1111/evo.13514)

44. Bookstein FL. 1997 Landmark methods for forms without landmarks: morphometrics of
group differences in outline shape. *Medical Image Analysis* **1**, 225–243.

45. Baken EK, Collyer ML, Kaliontzopoulou A, Adams DC. 2021 geomorph v4.0 and gmShiny:
Enhanced analytics and a new graphical interface for a comprehensive morphometric experience.
Methods Ecol Evol **12**, 2355–2363. (doi:10.1111/2041-210x.13723)

46. Upham NS, Esselstyn JA, Jetz W. 2019 Inferring the mammal tree: Species-level sets of
phylogenies for questions in ecology, evolution, and conservation. *PLoS Biol* **17**, e3000494-44.
(doi:10.1371/journal.pbio.3000494)

47. Mitchell DR, Sherratt E, Weisbecker V. 2023 Facing the facts: adaptive trade□offs along
body size ranges determine mammalian craniofacial scaling. *Biol. Rev.* (doi:10.1111/brv.13032)

48. Adams DC. 2014 A method for assessing phylogenetic least squares models for shape and
other high-dimensional multivariate data. *Evolution* **68**, 2675–2688. (doi:10.1111/evo.12463)

49. Stayton CT. 2015 The definition, recognition, and interpretation of convergent evolution, and
two new measures for quantifying and assessing the significance of convergence. *Evolution* **69**,
2140–2153. (doi:10.1111/evo.12729)

50. Grossnickle DM, Brightly WH, Weaver LN, Stanchak KE, Roston RA, Pevsner SK, Stayton
CT, Polly PD, Law CJ. 2023 Challenges and advances in measuring phenotypic convergence.
bioRxiv , 2022.10.18.512739. (doi:10.1101/2022.10.18.512739)

51. Hansen TF. 1997 Stabilizing selection and the comparative analysis of adaptation. *Evolution*
51, 1341–1351. (doi:10.1111/j.1558-5646.1997.tb01457.x)

52. Butler MA, King AA. 2004 Phylogenetic comparative analysis: a modeling approach for
adaptive evolution. *Am Nat* **164**, 683–695. (doi:10.1086/426002)

53. Clavel J, Escarguel G, Merceron G. 2015 mv morph: an R package for fitting multivariate
evolutionary models to morphometric data. *Methods Ecol Evol* **6**, 1311–1319.
(doi:10.1111/2041-210x.12420)

54. Coombs EJ, Felice RN, Clavel J, Park T, Bennion RF, Churchill M, Geisler JH, Beatty B,
Goswami A. 2022 The tempo of cetacean cranial evolution. *Curr Biol* **32**, 2233-2247.e4.
(doi:10.1016/j.cub.2022.04.060)

55. Rabosky DL. 2014 Automatic detection of key innovations, rate shifts, and diversity-
dependence on phylogenetic trees. *PLoS ONE* **9**, e89543-15. (doi:10.1371/journal.pone.0089543)

56. Höhna S, Freyman WA, Nolen Z, Huelsenbeck JP, May MR, Moore BR. 2019 A Bayesian
Approach for Estimating Branch-Specific Speciation and Extinction Rates. *bioRxiv* , 555805.
(doi:10.1101/555805)

57. Rabosky DL, Grundler M, Anderson C, Title P, Shi JJ, Brown JW, Huang H, Larson JG.
2014 BAMMtools: an R package for the analysis of evolutionary dynamics on phylogenetic
trees. *Methods Ecol Evol* **5**, 701–707. (doi:10.1111/2041-210x.12199)

58. Höhna S, Landis MJ, Heath TA, Boussau B, Lartillot N, Moore BR, Huelsenbeck JP,
Ronquist F. 2016 RevBayes: Bayesian Phylogenetic Inference Using Graphical Models and an
Interactive Model-Specification Language. *Syst. Biol.* **65**, 726–736. (doi:10.1093/sysbio/syw021)

59. Tribble CM, Freyman WA, Landis MJ, Lim JY, Barido☐Sottani J, Kopperud BT, H☐hna S,
May MR. 2022 RevGadgets: An R package for visualizing Bayesian phylogenetic analyses from
RevBayes. *Methods Ecol. Evol.* **13**, 314–323. (doi:10.1111/2041-210x.13750)

60. Plummer M, Best N, Cowles K, Vines K. 2006 CODA: Convergence Diagnosis and Output
Analysis for MCMC. *R news* **6**, 7–11.

61. Radinsky LB. 1981 Evolution of skull shape in Carnivores .1. Representative modern
Carnivores. *Biol J Linn Soc* **15**, 369–388. (doi:10.1111/j.1095-8312.1981.tb00770.x)

62. Law CJ, Young C, Mehta RS. 2016 Ontogenetic scaling of theoretical bite force in southern
sea otters (Enhydra lutris nereis). *Physiol Biochem Zool* **89**, 347–363. (doi:10.1086/688313)

63. Dickinson E *et al.* 2021 Evaluating bony predictors of bite force across the order Carnivora. *J
Morpho* **282**, 1499–1513. (doi:10.1002/jmor.21400)

64. Hartstone-Rose A, Dickinson E, Deutsch AR, Worden N, Hirschkorn GA. 2021 Masticatory
muscle architectural correlates of dietary diversity in Canidae, Ursidae, and across the order
Carnivora. *Anat. Rec.* (doi:10.1002/ar.24748)

65. Phillips PC, Arnold SJ. 1989 Visualizing multivariate selection. *Evolution* **43**, 1209–1222.
(doi:10.1111/j.1558-5646.1989.tb02569.x)

66. Polly PD. 2004 On the simulation of the evolution of morphological shape: multivariate
shape under selection and drift. *Palaeontologia Electornica* **7**.

67. Law CJ, Hlusko LJ, Tseng ZJ. 2024 Uncovering the mosaic evolution of the carnivoran
skeletal system. *Biol. Lett.* **20**, 20230526. (doi:10.1098/rsbl.2023.0526)

68. Adams DC, Berns CM, Kozak KH, Wiens JJ. 2009 Are rates of species diversification
correlated with rates of morphological evolution? *Proc R Soc B* **276**, 2729–2738.
(doi:10.1098/rspb.2009.0543)

69. Derryberry EP, Claramunt S, Derryberry G, Chesser RT, Cracraft J, Aleixo A, Pérez-Emán J,
Jr JVR, Brumfield RT. 2011 Lineage diversification and morphological evolution in a
large□scale continental radiation: the neotropical ovenbirds and woodcreepers (Aves:
Furnariidae). *Evolution* **65**, 2973–2986. (doi:10.1111/j.1558-5646.2011.01374.x)

70. Aristide L, Rosenberger AL, Tejedor MF, Perez SI. 2015 Modeling lineage and phenotypic
diversification in the New World monkey (Platyrrhini, Primates) radiation. *Mol Biol Evol* **82**,
375–385. (doi:10.1016/j.ympev.2013.11.008)

71. Law CJ, Slater GJ, Mehta RS. 2018 Lineage diversity and size disparity in Musteloidea:
testing patterns of adaptive radiation using molecular and fossil-based methods. *Syst. Biol.* **67**,
127–144. (doi:10.1093/sysbio/syx047)

72. Rabosky DL, Lovette IJ. 2008 Explosive evolutionary radiations: decreasing speciation or
increasing extinction through time? *Evolution* **62**, 1866–1875. (doi:10.1111/j.1558-
5646.2008.00409.x)

73. Quental TB, Marshall CR. 2009 Extinction during evolutionary radiations: reconciling the
fossil record with molecular phylogenies. *Evolution* **63**, 3158–3167. (doi:10.1111/j.1558-
5646.2009.00794.x)

74. Liow LH, Quental TB, Marshall CR. 2010 When can decreasing diversification rates be
detected with molecular phylogenies and the fossil record? *Syst. Biol.* **59**, 646–659.
(doi:10.1093/sysbio/syq052)

75. Day JJ, Peart CR, Brown KJ, Friel JP, Bills R, Moritz T. 2013 Continental diversification of
an African catfish radiation (Mochokidae: Synodontis). *Syst. Biol.* **62**, 351–365.
(doi:10.1093/sysbio/syt001)

76. Kozak KM, Wahlberg N, Neild AFE, Dasmahapatra KK, Mallet J, Jiggins CD. 2015
Multilocus species trees show the recent adaptive radiation of the mimetic heliconius butterflies.
Systematic Biology **64**, 505–524. (doi:10.1093/sysbio/syv007)

77. Liedtke HC *et al.* 2016 No ecological opportunity signal on a continental scale?
Diversification and life-history evolution of African true toads (Anura: Bufonidae). *Evolution* **70**,
1717–1733. (doi:10.1111/evo.12985)

78. Diaz LFH, Harmon LJ, Sugawara MTC, Miller ET, Pennell MW. 2019 Macroevolutionary
diversification rates show time dependency. *Proc. Natl. Acad. Sci.* **116**, 7403–7408.
(doi:10.1073/pnas.1818058116)

79. Goodman SM, Rasoloarison RM, Ganzhorn JU. 2004 On the specific identification of
subfossil Cryptoprocta (Mammalia, Carnivora) from Madagascar. *Zoosystema* **26**, 129–143.

80. Goodman SM, Helgen KM. 2010 Species limits and distribution of the Malagasy carnivoran
genus Eupleres (Family Eupleridae). *mamm* **74**, 177–185. (doi:10.1515/mamm.2010.018)

81. Wit MJ de. 2003 MADAGASCAR: Heads It's a Continent, Tails It's an Island. *Earth Planet.*
Sci. **31**, 213–248. (doi:10.1146/annurev.earth.31.100901.141337)

82. Vences M, Wollenberg KC, Vieites DR, Lees DC. 2009 Madagascar as a model region of
species diversification. *Trends Ecol. Evol.* **24**, 456–465. (doi:10.1016/j.tree.2009.03.011)

83. Krause DW, O'Connor PM, Sertich JJW, Rogers KC, Rogers RR, Rakotozafy B. 2022 The
New Natural History of Madagascar. , 59–68. (doi:10.2307/j.ctv2ks6tbb.13)

84. Raxworthy CJ, Pearson RG, Zimkus BM, Reddy S, Deo AJ, Nussbaum RA, Ingram CM.
2008 Continental speciation in the tropics: contrasting biogeographic patterns of divergence in
the Uroplatus leaf□tailed gecko radiation of Madagascar. *J. Zoo□l.* **275**, 423–440.
(doi:10.1111/j.1469-7998.2008.00460.x)

697