The Effects of Acute Coordinative vs. Acute Endurance Exercise on the Cortisol Concentration

Henning Budde*[1], Mirko Wegner[2], Christiane Ahrens[1], Bruna Velasques[3,4], Pedro Ribeiro[3,4], Sergio Machado[5], Anett Mueller-Alcazar[6]

[1] Institute for Systems Medicine (ISM), MSH Medical School Hamburg University of Applied Sciences and Medical University Hamburg, Germany.

[2] Department of Health Science, Institute of Human Movement Science, University of Hamburg, Germany.

[3] Bioscience Department, School of Physical Education of the Federal University of Rio de Janeiro, Rio de Janeiro, Brazil; Institute of Applied Neuroscience, Rio de Janeiro, Brazil.

[4] Brain Mapping and Sensory Motor Integration, Institute of Psychiatry, Rio de Janeiro, Brazil.

[5] Panic and Respiration, Institute of Psychiatry, Federal University of Rio de Janeiro, Brazil and Laboratory of Physical Activity Neuroscience, Neurodiversity Institute, Queimados-RJ, Brazil

[6] Institute for Cognitive and Affective Neuroscience, MSH Medical School Hamburg University of Applied Sciences and Medical University Hamburg, Germany.

Running Title: Cortisol due to Coordinative or Endurance Exercise

***Correspondence**:

Henning Budde, E-Mail address: henning.budde@medicalschool-hamburg.de

ABSTRACT

Objective: Physical exercise interventions can cause neuroendocrine activation,

which in turn increases salivary cortisol concentrations. Until now there have been

mostly studies focusing on endurance as one possible mode of exercise. The

purpose of this study was to examine the effects of these different interventions

with an intraindividual comparison. Methods: 61 students between 18 and 30

years of age were included and first completed a coordinative exercise and seven

days later an endurance exercise of the same intensity and length which was self-

set on the first day, with a maximum heart rate of 64 - 76% (HRmax) over a period

of 15 min. To measure changes in the hypothalamic-pituitary-adrenal axis (HPA-

axis) activity, saliva samples were collected before (t1) and after exercise (t2 and

t3). Results: Baseline values of cortisol (t0) did not differ significantly between

coordinative and endurance exercise, $t(55) = .233$, $p = .816$. Post hoc tests

revealed that the cortisol values of the second coordinative vs. endurance time

point, $t(55) = 2.097$, $p = .040$, $d = .741$, and third coordinative vs. endurance time

point, $t(55) = 3.004$, $p = .004$, $d = .735$, differed significantly with large effect sizes.

Conclusion: The results show that coordinative exercise produced a higher cortisol

release than endurance exercise. Interventions such as coordinative exercise

require higher cognitive components resulting in stronger cortisol release than

endurance exercise of the same intensity and length. Thus, the type of acute

physical stress is a psychophysiological factor in determining the neuroendocrine

stress response.

57 Keywords: Acute exercise, Coordinative exercise, Cortisol concentration,

58 Endurance exercise, Physical stress, Psychological stress

Introduction

Acute physical exercise influences neuroendocrine responses and can lead to an increase in cortisol concentration if the intensity of the intervention exceeds a certain threshold [1]. In addition, studies suggest that enhanced chronical physical activity (PA) facilitates neuroplasticity of certain brain structures and is associated with an improvement in neurogenesis, synaptogenesis, the release of neurotrophins and neuroendocrinological changes that are linked to benefits for cognitive and affective functions [2]. In the long term, this may be reflected in a reduction in psychosocial stress symptoms [3] and in short term lead to an improvement in cognitive performance [4]. Different modes of exercise have gained increased attention in research in recent years [5].

In the event of acute physical or psychological stress, our body's stress response essentially occurs via two axes: the sympatho-adrenomedullary system puts the body into an alarm state when it is exposed to stress and ensures an increase in heart and respiratory rate by releasing adrenaline and noradrenaline [6]. The hypothalamic-pituitary-adrenal axis (HPA-axis) forms the second stress axis, which is primarily characterized by the release of the stress hormone cortisol [7]. Cortisol has multiple effects on metabolism [8], the immune system and inflammatory processes [9], and is the most important hormone of the HPA-axis alongside corticotropin-releasing hormone (CRH) and adrenocorticotropic hormone (ACTH). Acute physical exercise is acknowledged as an immediate stressor for the body and affects the secretion processes of several endocrine tissues and the subsequent release of hormones like cortisol [10].

The circadian rhythm of cortisol is stable and is only slowly influenced by external

conditions. During acute physical stress, a peak secretion of cortisol can be

observed approximately 20 to 30 minutes after the onset of the stimulus [11]. If the

organism is unable to terminate the stress response, e.g. in the case of chronic

stress exposure, stress-adaptive mechanisms can lead to pathological changes.

The role of the endocrine stress system can be seen in the pathophysiology of

various mental illnesses such as depression [3], burnout [12], or PTSD [13]. Also

physical exercise can induce changes on the functioning of the HPA-axis, provided

that a certain intensity threshold is exceeded [14] [3].

Physical exercise refers to targeted, planned and structured forms of PA [15].

Physical exercise should be differentiated into acute physical exercise (single

bout) and chronic physical exercise (repeated bouts of physical exercise or

exercise training). The intensity and duration of the activity is considered the

primary physiological factor for determining the neuroendocrine stress response [10].

Adult subjects showed a linear increase in salivary cortisol concentration over a

period of 15 minutes at a maximum exercise intensity of 70-85% maximum Heart

rate (HRmax), whereas no significant changes were observed at lower exercise

intensities [14]. In connection with psychosocial stress, the cortisol response

appears to be even more pronounced [16]. It can therefore be assumed that, in

addition to the intensity of the stressors, the type of stressor itself also has an

influence on cortisol levels and that psychosocial stress can increase cortisol

levels significantly more than purely physical stress of moderate intensity (65-75 %

HRmax) [17].

Salivary cortisol responses show great intra- and inter-individual variability and are determined by various factors [18]. Personality traits, age, gender, endogenous and exogenous sex steroid levels, genetic predispositions or critical life events and in particular negative childhood experiences are worth mentioning in this context [19]. Several studies have shown that chronic endurance training (which is defined as planned, structured, repetitive, and purposeful PA leading to a change in fitness [20]). can effect cortisol concentration in the long term [21] and thus has both a preventive and therapeutic effect in reducing the experience of stress [22]. This shows that high levels of PA are related to a lower physiological stress reactivity. The immediate neuroendocrine stress responses induced by exercise are not yet fully understood. Results of studies comparing different types of *chronic* interventions showed that coordinative exercise led to a greater reduction in cortisol concentrations and stress levels compared to chronic endurance exercise [21]. However, the immediate stress responses produced by *acute* exercise interventions are still not clear specially regarding how the acute responses differ depending on the type of exercise performed.

Based on current research, the question of whether there are direct effects of different acute interventions of the same intensity (i.e. with the same cardiovascular load) on the cortisol concentration will therefore be investigated. A distinction is made between acute physical exercise of a coordinative nature and acute physical endurance exercise. As higher cognitive abilities will be required during the coordinative exercise (e.g. the ability to react and concentrate)[23], both physiological and psychological stressors are at work here. For this reason, it is assumed that stress load will be higher. Thus, the type of acute physical stress could be a physiological factor in determining the neuroendocrine stress response.

To our knowledge, there have been no studies on whether cortisol concentration

differ between an acute coordinative exercise and an acute endurance exercise of

the same intensity and duration. The reaction to physical interventions is

demonstrably individual and depends on many influencing factors [24]. For this

reason, the intra- and inter-individual variability should be given special

consideration here [5] and influencing factors such as gender, PA and will be

investigated exploratively.

Material and Methods

Participants

The G*Power analysis (α = 5%, 1-β= 95%, d = 0.95) resulted in a required sample size of 60 participants. The participants were informed on the characteristics of their participation, agreed to participate and signed the informed consent. The study was approved by the ethics committee of the Medical School Hamburg (MSH-2021/131) and is conducted in accordance with the latest revision of the Declaration of Helsinki (2013). Further inclusion criteria for participation in the study were the absence of mental or physical impairment, drug or alcohol dependence, acute psychotic illness and a BMI of < 30 (overweight). Based on these criteria, no participant had to be excluded. Both of the exercise sessions had to be accomplished with a moderate intensity between 64 – 76% HRmax [25]. In addition, we asked the participants about their perceived social support and their satisfaction with their everyday social relationships and measured their self-perceived stress. Motor fitness and PA were also assessed. A total of 61 people between the ages of 18 and 29 took part in the study. 32 were female. The average age was 21.8 years (SD = 3.09, N = 61).

Study design

For the present study, a moderate exercise intensity just before the vigorous range was aimed for, based on Garber's classification of exercise intensity [25]. The intensity of the different exercise sessions should be between 64 - 76% HRmax. According to Tanaka's formula: HRmax = 208 - 0.7 x age [26], this corresponds to approximately 140 to 145 beats per minute (bpm). Additionally we assessed the

Rate of Perceived Exertion (RPE) using the Borg Scale, a numerical scale between 6 and 20 to record the subjective feeling of exertion [27], which should correspond to a score between 13 and 14 (somewhat strenuous to strenuous). The HR (bpm) was measured before and during each intervention. The standardized individual target HR and the RPE of the coordinative intervention (Co) determined the intensity of the endurance intervention (En) and were continuously recorded and monitored with a HR monitor (Polar M430, Kempele, Finland).

After registering, the participants were instructed not to exercise 24 hours and not to eat for two hours before the start of the exercises. Each participant took part in two interventions, which took place between 2 p.m. and 4 p.m. seven days apart. In the first session, questionnaires were used to collect demographic data PA as well as the current state of loneliness which took about 20 minutes. During this time, participants were able to calm down. A coordination exercise (Co) lasting 15 minutes was initially performed. Using a coordination ladder 10 cm above the floor, the subjects were asked to complete various step and jump sequences After a washout phase of seven days, the second intervention was carried out at the same time in form of endurance exercise (En) with the same intensity for the cardiovascular system. Again, the subjects first completed a questionnaire in order to acclimatize for about 20 minutes. After putting on the HR measuring devices, the subjects had to run for 15 minutes, while getting acoustic feedback about the HR level they should achieve (second session). The rate the RPE was again assessed using the Borg scale [27].

The following questionnaires were used:

A demographic questionnaire contained physiological parameters (height, age and

weight) as well as questions on the subject's personal circumstances.

The Godin Leisure-Time Exercise Questionnaire (GLTQ) [28] was used to assess

PA. The GLTQ determines the level of PA and has been shown to correlate with

the level of physical fitness [29].

FFB-mot [30] was used to assess motor fitness. The subscales of the FFB-mot are

the components strength, endurance, flexibility and coordination.

In addition, before the second intervention (En), the subjects' subjective

experience of loneliness was assessed using the UCLA Loneliness Scale [31].

The planned study has a quasi-experimental, „pre-post-intervention" design with

repeated measurements [32]. The intraindividual variability of the cortisol

concentration describes the different responses of the same subject to two

different interventions at three different measurement times. The timing of both

interventions was exactly the same. In order to standardize the intensity of the

endurance exercise the coordinative exercise had to be performed first. A

randomization of the subjects was therefore not possible. See Figure 1.

Insert Figure 1 here

The first sample was taken immediately before the intervention (t0), the second

sample five minutes after completion (t1) and the third sample 30 minutes after the

end of the exercise (t2). Saliva samples were collected using the SaliCap system

(IBL Hamburg) and stored at minus 20°C until laboratory analysis. The saliva

analysis method is considered a reliable non-invasive way of determining stress-

related adrenal hormone levels [33]. The concentration of cortisol in the saliva

correlates closely with the concentration of free cortisol in the blood, and it has

been confirmed that the measurement in the saliva is a reliable tool for

investigations of HPA activities [34]. The biochemical analysis was carried out at the

University of Technology Dresden using immunoassay analysis. Intra- and inter-

assay coefficients of variation were 1.8 and 2.9 %, respectively.

Data Analysis

Due to non-evaluable cortisol samples, two participants had to be excluded. Three

others did not complete the exercise with the required intensity and were also

excluded. The sample for the calculation comprises 56 participants (25 male and

31 female).

For two of the six measurement points a total of 5 samples were missing. Overall

this was 1.5% missing data which allowed for imputation techniques. Multiple

imputation techniques were employed using SPSS to complete the data set using

the mean of five different random models. As a result complete cortisol values for

all six measurement points and all 56 participants could be analyzed. For further

analysis the cortisol values were log-transformed because they were not normally

distributed.

The main analysis of the changes in cortisol release after both interventions (acute

coordination and endurance exercise) was carried out using an ANOVA with

repeated measures. This was carried out here as a special form of analysis of

variance due to the presence of linked measurements or data with repeated

measurements. Data were tested for sphericity and if present corrected is first

tested using the Mauchly test. Effect sizes were calculated in the ANOVAS,

significant results are reported by Eta-square (η^2).

In addition, the influence of the variables gender, motor fitness (surveyed by

questionnaire FFBmot) and PA (questionnaire GLTQ) on the differences between

the individual measurements was exploratively examined and considered as

covariates in the context of the ANOVA. The influence of gender, $F_{(1,50)} = 0.959$,

p = .332, η² = 0.019, motor fitness (FFBmot), $F_{(1,50)} = 1.91$, *p = 0.172*, η² =

0.037, and PA $F_{(1,50)} = 0.029$, *p = .866*, η² = 0.001, were not significant.

However, the measured cortisol concentrations were not influenced by any of

these variables. The subjective feeling of loneliness (questionnaire UCLA) (F

$_{(1,50)} = 0.022$, *p = .883*, η² = 0.001) also showed no influence on the

measurement results. Therefore, in the following, analyses for Cortisol were

reported and analysed without any covariate.

T-tests (two-sided) for dependent samples were calculated to test whether the

cortisol concentration differed between the two interventions (Co/En). The

prerequisite for the application of this method is normal distribution of the samples.

The effect sizes are reported as Cohen's d. In some cases, more than 10% of the

data points for the variables under consideration are below the 5th percentile or

above the 95th percentile.

Insert Table 1 here

Results

Preliminary analyses suggest that none of the demographic variables (age, gender, BMI, physical activity, mood) affected the results significantly and were thus not included as covariates in the ANOVA. Furthermore, baseline values of cortisol (t0) did not differ significantly between coordinative and endurance exercise, t (55) = .233, p = .816.

Mean values and standard deviations are shown in Table 1. Repeated measure ANOVA revealed a main effect of the measurement point, F (2) = 22.472, $p < .001$, η^2 = .290, and the type of exercise, F (1) = 5.587, p = .022, η^2 = .092. The interaction of measurement points and type of exercise was also significant, F (2, 110) = 4.322, p = .016, η^2 = .073 (see Figure 2).

Post hoc tests revealed that the cortisol values of the second coordinative vs. endurance time point, t (55) = 2.097, p = .040, d = .741, and third coordinative vs. endurance time point, t (55) = 3.004, p = .004, d = .735, differed significantly with large effect sizes. Moreover, within the coordinative exercise condition, cortisol levels at time point three were significantly higher than in time point one, t (55) = -3.858, $p < .001$, d = .684, and two, t (55) = -6.814, $p < .001$, d = .484. In the coordinative condition, cortisol levels at time point one and two did not differ significantly.

In the endurance exercise condition, cortisol levels significantly differed between time point one and two, t (55) = 4.000, $p < .001$, d = .519, and between time point

two and three, *t* (55) = -5.631, *p* < .001, *d* = .469. Cortisol levels did not differ

significantly between time points one and three. The within conditions showed

medium effect sizes.

Insert Figure 2 here

During the coordinative exercise session, participant's mean HR was not

significantly different from the predefined HR target (144 bpm; 142 bpm). The

mean HR during the endurance exercise was 143 bpm.

Discussion

Our main result was already hypothesized before the study. We saw that more

cortisol is released in the aftermath of acute coordinative exercise of moderate

intensity than due to an acute endurance exercise of the same intensity and

duration. The endurance intervention was carried out at the same time seven days

after the coordinative exercise, between 64 and 76% HRmax [27] lasting 15 minutes.

We explain our acute results with the effect we observed when comparing different

types of *chronic* interventions (Cortisol Awakening Response before and after a

training 3 times a week over 10 weeks). Showing that coordinative training led to a

greater reduction in cortisol concentration compared to an endurance training [21].

According to Athanasiou et al. (2023) regular acute exercise leads to adaptations

that are reflected in a reduced physiological stress response [35]. It would therefore

be expected that enhanced cortisol concentration acutely results in a reduced

concentration after a chronic treatment. If this homeostatic threat appears

chronical it might lead to a lower response compared to the pretraining levels.

Rehfeld et al., (2018) observed a significant elevation in both white and gray

matter, mainly in the cerebellum after a coordinative compared to a noncoordinative training [36].

It is possible that different neurobiological signaling pathways are activated by different exercise interventions [21]. The coordinative intervention places more demanding cognitive requirements on movement accuracy than endurance exercise[23]. Budde et al. (2008) observed significantly higher attention after 10 min of coordinatively demanding exercise compared to endurance exercise [23]. Coordinative exercise requires perceptual and higher-level cognitive processes, such as attention, that are essential for mapping sensation to action and ensuring anticipatory and adaptive aspects of coordination [37]. This may show that coordinative interventions lead to a pre-activation of cognitive neuronal networks [23], which is reflected in endocrine responses.

After the interventions, cortisol concentration was significantly increased in the third measurement (t3) only after the Co condition, which was performed 30 min after the end of the intervention, compared to t1. This result is not completely in line with those of comparable studies. After acute physical running exercise of a higher intensity (70-85% HRmax) over a period of 12 min in younger students, an increase in cortisol concentration could be observed in between 5 minutes after the exercise [1] whereby the intensity and duration of the exercise play a significant role in the cortisol secretion. Cortisol is first secreted a few minutes after the stressor before it then rises continuously until the peak of secretion is reached after approximately 20 to 30 minutes [38]. However, individuals respond differently to certain exercise stimuli.

In addition to the environmental influencing factors already listed, particularly

pronounced personality traits such as implicit affiliation motive or power motive

[19,39] could also be attributed to affect the cortisol response.

With regard to gender, there was no difference to be found in cortisol secretion.

However which is not surprising, since the available data is very heterogeneous [18].

Due to our research question, we were unable to use an randomized controlled

trial design [40]. We first had to determine the HR during the coordinative exercise in

order to determine the intensity of the second exercise. This is more difficult with a

coordinative exercise than with an endurance exercise. It therefore needed to be

determined first. The lack of random assignment is the major weakness of the

quasi-experimental study design [41]. However, this was unavoidable. The

participants were exercising in an interindividual comparable intensity according to

Garber et al. (2011) [25], we calculated their HRmax using a formula [26] and did not

explicitly test their HRmax for example, with the shuttle run test [42]. However, the

used questionnaire generates a reliable measurement of the motor fitness

including the subscales cardiorespiratory fitness/endurance and gross motor

coordination [30]. The PA-questionnaire (GLTQ) [28] we used is significantly

associated with fitness of the participants [29].

This intraindividual study was able to show that after an acute exercise

intervention of moderate intensity (64-76% HRmax) over a period of 15 min,

cortisol levels increased more under the coordinative conditions than under the

endurance conditions. These results indicate that these different modes of acute

exercise cause different psychophysiological HPA activities, which could be of

interest for future stress research. These results might be extrapolated to other

clinical areas focused on stress responses (like exercising with a motor handicap).

References

1.   Budde H, Pietrassyk-Kendziorra S, Bohm S, Voelcker-Rehage C. Hormonal responses to

physical and cognitive stress in a school setting. *Neurosci Lett*. 05/2010a;474(3):131-134.

doi:10.1016/j.neulet.2010.03.015

2.   Budde H, Velasques B, Ribeiro P, Soya H. Editorial: Neuromodulation of Exercise: Impact on

Different Kinds of Behavior. *Front Neurosci*. 2020;14:455. doi:10.3389/fnins.2020.00455

3.   Helmich I, Latini A, Sigwalt A, et al. Neurobiological Alterations Induced by Exercise and Their

Impact on Depressive Disorders. *Clin Pract Epidemiol Ment Health CP EMH*. 2010;6:115-125.

doi:10.2174/1745017901006010115

4.   Budde H, Voelcker-Rehage C, Pietrassyk-Kendziorra S, Machado S, Ribeiro P, Arafat AM.

Steroid hormones in the saliva of adolescents after different exercise intensities and their

influence on working memory in a school setting. *Psychoneuroendocrinology*.

4/2010b;35(3):382-391. doi:10.1016/j.psyneuen.2009.07.015

5.   Herold F, Törpel A, Hamacher D, et al. Causes and Consequences of Interindividual Response

Variability: A Call to Apply a More Rigorous Research Design in Acute Exercise-Cognition

Studies. *Front Physiol*. 2021;12. Accessed September 28, 2023.

https://www.frontiersin.org/articles/10.3389/fphys.2021.682891

6.   Olsson EMG, Norlund F, Rondung E, et al. The e-mental health treatment in Stockholm

myocardial infarction with non-obstructive coronaries or Takotsubo syndrome study (E-

SMINC): a study protocol for a randomised controlled trial. *Trials*. 2022;23(1):597.

doi:10.1186/s13063-022-06530-3

7.   Godoy LD, Rossignoli MT, Delfino-Pereira P, Garcia-Cairasco N, de Lima Umeoka EH. A

Comprehensive Overview on Stress Neurobiology: Basic Concepts and Clinical Implications.

Front Behav Neurosci. 2018;12. Accessed December 29, 2023.

https://www.frontiersin.org/articles/10.3389/fnbeh.2018.00127

8.   Stimson RH, Anderson AJ, Ramage LE, et al. Acute physiological effects of glucocorticoids on

fuel metabolism in humans are permissive but not direct. *Diabetes Obes Metab*.

2017;19(6):883-891. doi:10.1111/dom.12899

9.   Oppong E, Cato ACB. Effects of Glucocorticoids in the Immune System. *Adv Exp Med Biol*.

2015;872:217-233. doi:10.1007/978-1-4939-2895-8_9

10. Hackney AC. Exercise as a stressor to the human neuroendocrine system. *Med Kaunas Lith*.

2006;42(10):788-797.

11. Kirschbaum C, Hellhammer DH. Salivary cortisol in psychoneuroendocrine research: Recent

developments and applications. *Psychoneuroendocrinology*. 1994;19(4):313-333.

doi:10.1016/0306-4530(94)90013-2

12. Monfared SS, Lebeau JC, Mason J, et al. A Bio-Physio-Psychological Investigation of Athletes'

Burnout. *Res Q Exerc Sport*. 2021;92(1):189-198.

13. Budde H, Akko DP, Ainamani HE, Murillo-Rodríguez E, Weierstall R. The impact of an exercise

training intervention on cortisol levels and post-traumatic stress disorder in juveniles from an

Ugandan refugee settlement: study protocol for a randomized control trial. *Trials*.

2018;19(1):364. doi:10.1186/s13063-018-2753-x

14. O'Connor PJ, Corrigan DL. Influence of short-term cycling on salivary cortisol levels. *Med Sci*

Sports Exerc. 1987;19(3):224-228.

15. Budde H, Schwarz R, Velasques B, et al. The need for differentiating between exercise, physical

activity, and training. *Autoimmun Rev*. 2016;15(1):110-111. doi:10.1016/j.autrev.2015.09.004

16. Clemente-Suárez VJ, Robles-Pérez JJ, Fernández-Lucas J. Psycho-physiological response in an

automatic parachute jump. *J Sports Sci*. 2017;35(19):1872-1878.

doi:10.1080/02640414.2016.1240878

17. Wegner M, Müller-Alcazar A, Jäger A, Machado S, Arias-Carrión O, Budde H. Psychosocial

Stress But Not Exercise Increases Cortisol and Reduces State Anxiety Levels in School Classes -

Results from a Stressor Applicable in Large Group Settings. *CNS Neurol Disord Drug Targets*.

2014;13:1015-20. doi:10.2174/1871527313666140612103425

18. Kudielka BM, Hellhammer DH, Wüst S. Why do we respond so differently? Reviewing

determinants of human salivary cortisol responses to challenge. *Psychoneuroendocrinology*.

2009;34(1):2-18. doi:10.1016/j.psyneuen.2008.10.004

19. Wegner M, Schüler J, Budde H. The implicit affiliation motive moderates cortisol responses to

acute psychosocial stress in high school students. *Psychoneuroendocrinology*. 2014;48:162-

168. doi:10.1016/j.psyneuen.2014.06.013

20. Wegner M, Amatriain-Fernández S, Kaulitzky A, Murillo-Rodriguez E, Machado S, Budde H.

Systematic Review of Meta-Analyses: Exercise Effects on Depression in Children and

Adolescents. *Front Psychiatry*. 2020;11. Accessed August 29, 2023.

https://www.frontiersin.org/articles/10.3389/fpsyt.2020.00081

21. Wegner M, Koutsandréou F, Müller-Alcazar A, Lautenbach F, Budde H. Effects of Different

Types of Exercise Training on the Cortisol Awakening Response in Children. *Front Endocrinol*.

2019;10. Accessed August 29, 2023.

https://www.frontiersin.org/articles/10.3389/fendo.2019.00463

22. Chovanec L, Gröpel P. Effects of 8-week endurance and resistance training programmes on

cardiovascular stress responses, life stress and coping. *J Sports Sci*. 2020;38(15):1699-1707.

doi:10.1080/02640414.2020.1756672

23. Budde H, Voelcker-Rehage C, Pietraßyk-Kendziorra S, Ribeiro P, Tidow G. Acute coordinative

exercise improves attentional performance in adolescents. *Neurosci Lett*. 2008;441(2):219-

223. doi:10.1016/j.neulet.2008.06.024

24. Gronwald T, Törpel A, Herold F, Budde H. Perspective of Dose and Response for Individualized

Physical Exercise and Training Prescription. *J Funct Morphol Kinesiol*. 2020;5.

doi:10.3390/jfmk5030048

25. Garber C, Blissmer B, Deschenes M, et al. Quantity and Quality of Exercise for Developing and

Maintaining Cardiorespiratory, Musculoskeletal, and Neuromotor Fitness in Apparently

Healthy Adults: Guidance for Prescribing Exercise. *Med Sci Sports Exerc*. 2011;43:1334-1359.

doi:10.1249/MSS.0b013e318213fefb

26. Tanaka H, Monahan K, Seals D. Age-Predicted Maximal Heart Rate Revisited. *J Am Coll Cardiol*.

2001;37:153-156. doi:10.1016/S0735-1097(00)01054-8

27. Borg G, Loellgen H. Borg's Perceived Exertion and Pain Scales. 2001;52.

28. Godin G. The Godin-Shephard Leisure-Time Physical Activity Questionnaire. *Health Fit J Can*.

2011;4(1):18-22. doi:10.14288/hfjc.v4i1.82

29. Milton K, Bull F, Bauman A. Reliability and validity of a single-item physical activity measure.

Br J Sports Med. 2011;45:203-208. doi:10.1136/bjsm.2009.068395

30. Bös K, Abel T, Woll A, Niemann S, Tittlbach S, Schott N. Der Fragebogen zur Erfassung des

motorischen Funktionsstatus (FFB-MOT). *Diagnostica*. 2002;48:101-111. doi:10.1026//0012-

1924.48.2.101

31. Russell DW. UCLA Loneliness Scale (Version 3): Reliability, validity, and factor structure. *J Pers

Assess*. 1996;66(1):20-40. doi:10.1207/s15327752jpa6601_2

32. Harris AD, McGregor JC, Perencevich EN, et al. The Use and Interpretation of Quasi-

Experimental Studies in Medical Informatics. *J Am Med Inform Assoc JAMIA*. 2006;13(1):16.

doi:10.1197/jamia.M1749

33. Jollin L, Thomasson R, Panse B, et al. Saliva DHEA and cortisol responses following short-term

corticosteroid intake. *Eur J Clin Invest*. 2009;40:183-186. doi:10.1111/j.1365-

2362.2009.02219.x

34. Kirschbaum C, Hellhammer D. Salivary Cortisol in Psychobiological Research: An Overview.

Neuropsychobiology. 1989;22:150-169. doi:10.1159/000118611

35. Athanasiou N, Bogdanis GC, Mastorakos G. Endocrine responses of the stress system to

different types of exercise. *Rev Endocr Metab Disord*. 2023;24(2):251-266.

doi:10.1007/s11154-022-09758-1

36. Rehfeld K, Lüders A, Hökelmann A, et al. Dance training is superior to repetitive physical

exercise in inducing brain plasticity in the elderly. *PloS One*. 2018;13(7):e0196636.

doi:10.1371/journal.pone.0196636

37. Voelcker-Rehage C, Niemann C. Structural and functional brain changes related to different

types of physical activity across the life span. *Neurosci Biobehav Rev*. 2013;37(9, Part B):2268-

2295. doi:10.1016/j.neubiorev.2013.01.028

38. Dickerson S, Kemeny M. Acute Stressors and Cortisol Responses: A Theoretical Integration and

Synthesis of Laboratory Research. *Psychol Bull*. 2004;130:355-391. doi:10.1037/0033-

2909.130.3.355

39. Wegner M, Schüler J, Scheuermann K, Machado S, Budde H. The Implicit Power Motive

andAdolescents' Salivary Cortisol Responses to Acute Psychosocial Stress and Exercise in

School. *CNS Neurol Disord - Drug Targets Former Curr Drug Targets - CNS Neurol Disord*.

Published online November 11, 2015. doi:10.2174/1871527315666151111123122

40. Lim CY, In J. Considerations for crossover design in clinical study. *Korean J Anesthesiol*.

2021;74(4):293-299. doi:10.4097/kja.21165

41. Iwahori M, Oshiyama C, Matsuzaki H. A quasi-experimental controlled study of a school-based

mental health programme to improve the self-esteem of primary school children. *Humanit*

Soc Sci Commun. 2022;9(1):1-10. doi:10.1057/s41599-022-01156-x

42. Léger LA, Mercier D, Gadoury C, Lambert J. The multistage 20 metre shuttle run test for

aerobic fitness. *J Sports Sci*. 1988;6(2):93-101. doi:10.1080/02640418808729800

Measurement	Co (A, B, C)		En (D, E, F)	
	M	*SD*	*M*	*SD*
Cortisol (nmol/l)				
pre (t0)	3.52	2.69	3.24	1.98
post I (t1)	3.24	2.42	2.54	1.45
post II (t2)	5.41	5.73	3.70	2.53

Table 1 *Means (M) and standard deviations (SD) for cortisol (nmol/l) at pre- and post-test*

I and II as well as for change in cortisol concentration and change in cortisol mean value

between pre- and post-test II (n = 56).

Figure 1 *Explanation of the study design (d1/d2 = day of intervention, Co = coordinative*

exercise, En = endurance exercise, t0,1,2 = time of measurement.)

Figure 2 *Changes in cortisol concentration after the interventions coordinative (Co)*

compared to endurance (En).