Research Article

Glutamine catabolism supports amino acid biosynthesis and suppresses the integrated stress response to promote photoreceptor survival

Moloy T. Goswami[1,6], Eric Weh[1,6], Shubha Subramanya[1], Katherine M. Weh[1], Hima Bindu Durumutla[1,2], Heather Hager[1], Nicholas Miller[1], Sraboni Chaudhury[1], Anthony Andren[3], Peter Sajjakulnukit[3], Li Zhang[3], Cagri G. Besirli[1], Costas A. Lyssiotis[3,4,5], Thomas J. Wubben[1,7]

[1]Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, MI, USA; [2]Molecular and Developmental Biology Graduate Program, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA; [3]Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, MI, USA; [4]Department of Internal Medicine, Division of Gastroenterology and Hepatology, University of Michigan, Ann Arbor, MI, USA; [5]Rogel Cancer Center, University of Michigan, Ann Arbor, MI, USA; [6]equal contribution

[7]Corresponding author and lead contact: Thomas J. Wubben, MD, PhD, Kellogg Eye Center, University of Michigan, 1000 Wall Street, Ann Arbor, MI, USA, Phone: 734-936-8072, Email: twubben@med.umich.edu

Impact statement (one sentence, typically 15-30 words): For the first time, this work reveals the metabolic dependency of photoreceptors on glutamine catabolism *in vivo* and further demonstrates the flexibility of photoreceptors to utilize fuel sources beyond glucose.

Abstract

Photoreceptor loss results in vision loss in many blinding diseases, and metabolic dysfunction underlies photoreceptor degeneration. So, exploiting photoreceptor metabolism is an attractive strategy to prevent vision loss. Yet, the metabolic pathways that maintain photoreceptor health remain largely unknown. Here, we investigated the dependence of photoreceptors on glutamine (Gln) catabolism. Gln is converted to glutamate via glutaminase (GLS), so mice lacking GLS in rod photoreceptors were generated to inhibit Gln catabolism. Loss of GLS produced rapid rod photoreceptor degeneration. In vivo metabolomic methodologies and metabolic supplementation identified Gln catabolism as critical for glutamate and aspartate biosynthesis. Concordant with this amino acid deprivation, the integrated stress response (ISR) was activated with protein synthesis attenuation, and inhibiting the ISR delayed photoreceptor loss. Furthermore, supplementing asparagine, which is synthesized from aspartate, delayed photoreceptor degeneration. Hence, Gln catabolism is integral to photoreceptor health, and these data reveal a novel metabolic axis in these metabolically-demanding neurons.

Keywords (10 or less words)

Photoreceptor, glutaminase, metabolism, neurodegeneration

Introduction

Photoreceptor (PR) death is the cause of vision loss in many retinal diseases. A paucity of effective therapies exist that prevent PR death, so there is an unmet need for therapeutics that improve or prolong PR survival. The retina has a significant energetic demand, driven in large part by PRs (Pan et al., 2021). The prodigious bioenergetic requirements of photoreceptors (PRs) are due to the need to conduct phototransduction and neurotransmission as well as manufacture the lipid- and protein-rich outer segment, which is shed daily and phagocytosed by the retinal pigment epithelium (RPE) (Ng et al., 2015; Young, 1967). PRs have little reserve capacity to generate adenosine triphosphate (ATP) and as a result, are adversely affected by small changes in energy homeostasis. As such, metabolic dysfunction has been shown to underlie PR cell death (Bowne et al., 2006; Du et al., 2015; Hartong et al., 2008; Kooragayala et al., 2015; Pan et al., 2021), and exploiting PR metabolism to make these cells more robust to stress is an attractive neuroprotective strategy (Pan et al., 2021). Yet, beyond glucose, the metabolic pathways integral to PR health remain largely unknown. This is a critical knowledge gap as identification of these pathways is likely to reveal new strategies for therapeutic intervention (Duncan et al., 2018).

Glucose is central to PR metabolism as these cells utilize aerobic glycolysis, or the conversion of glucose to lactate despite the presence of oxygen, for the production of both energy and anabolic building blocks, similar to cancer cells (Ait-Ali et al., 2015; Chinchore et al., 2017; Kanow et al., 2017; Petit et al., 2018; Swarup et al., 2019). Previous studies have shown that genetic knockdown of enzymes key to aerobic glycolysis leads to PR dysfunction and death (Chinchore et al., 2017; Petit et al., 2018; Rajala, 2020; Weh et al., 2020; Wubben et al., 2017). However, like other metabolically-demanding cells, recent work has demonstrated that PRs have the flexibility to utilize fuel sources beyond glucose to meet their metabolic needs (Adler et al., 2014; Daniele et al., 2022; Du, Cleghorn, Contreras, Linton, et al., 2013; Grenell et al., 2019; Joyal et al., 2016; Xu et al., 2020).

Glutamine (Gln) is the most abundant amino acid found in the body and circulating within the blood (Yang et al., 2017), and many rapidly dividing cells, including cancer cells, which utilize aerobic glycolysis, depend on Gln for their survival and proliferation (Yang et

al., 2017). Gln can serve as a substrate for multiple pathways, providing a carbon and
nitrogen source for biosynthesis, energetics, and cellular reactive oxygen species (ROS)
homeostasis (Adler et al., 2014; Xu et al., 2020; Yang et al., 2017). These features
potentially make Gln an ideal alternative fuel source for PRs, whose bioenergetic demand
rivals that of cancer cells despite being terminally differentiated (Ng et al., 2015). To this
end, over half of the Gln in the retina is found in the outer retina, which is primarily
composed of PRs (Du, Cleghorn, Contreras, Linton, et al., 2013; Macosko et al., 2015;
Voigt et al., 2020). *Ex* vivo experiments have shown that Gln can be a source of carbons
for TCA cycle intermediates in the retina and contribute to amino acid biosynthesis (Du,
Cleghorn, Contreras, Linton, et al., 2013; Grenell et al., 2019; Tsantilas et al., 2021; Xu et
al., 2020). Additionally, Gln was demonstrated to support nicotinamide adenine
dinucleotide phosphate (NADPH) generation in the absence of glucose *in vitro* in isolated
PRs (Adler et al., 2014). Finally, in a genetically altered mouse model that disrupts glucose
transport to the PRs, both Gln and its transporter were upregulated in the retina, implying
that metabolism of this amino acid may be supporting PR survival when glucose is limiting
(Swarup et al., 2019).
Glutaminolysis is the process by which Gln is metabolized into TCA cycle
intermediates for critical biosynthetic precursors. This process is initiated by the
catabolism of Gln to glutamate (Glu) via one of two glutaminase enzymes: kidney-type
glutaminase (GLS) or liver-type glutaminase (GLS2) (Yang et al., 2017). Single-cell RNA
sequencing data (GSE63473 and GSE142449) has demonstrated that GLS is the
predominant isoform in the retina and PRs (Macosko et al., 2015; Voigt et al., 2020).
Additionally, it has been shown that glutaminase activity is at least two-fold higher in
mitochondria-rich PR inner segments (Ross et al., 1987). Since the inner segments of PRs
are responsible for supporting the majority of energy production and metabolism, GLS
activity is likely critical for metabolic functions in PRs.
Glutaminolysis via GLS is indispensable to the metabolism of many cancer cells
(Altman et al., 2016; Yang et al., 2017), and a variety of data show the similarities between
cancer cell metabolism and retinal metabolism, and specifically, metabolism of the PRs in
the outer retina (Chinchore et al., 2017; Du, Cleghorn, Contreras, Linton, et al., 2013; Ng
et al., 2015; Rajala, 2020). Therefore, we hypothesized that GLS-initiated Gln catabolism
may also be essential to PR metabolism, function, and survival. Previous studies utilizing
in vitro or *ex vivo* methods with whole retinas from wild-type mice or mice with genetic
perturbations not confined to PRs (Adler et al., 2014; Du, Cleghorn, Contreras, Linton, et
al., 2013; Grenell et al., 2019; Xu et al., 2020) have provided a foundation for this
governing hypothesis, but none have examined the role of Gln catabolism specifically in
PRs *in vivo.* In this study, we generated a rod photoreceptor-specific knockout of GLS to
comprehensively study the importance of GLS-driven Gln catabolism in PRs.

Results
116
Generation of a rod photoreceptor-specific, *Gls* knockout mouse
To confirm the single-cell RNA sequencing data (Macosko et al., 2015; Voigt et al., 2020),
we conducted real-time reverse transcription PCR (qRT-PCR) with primers specific for
either *Gls* or *Gls2* (Supplementary File 1). These data show that *Gls* expression is 14
times greater than that of *Gls2* in the mouse retina (Figure 1 – figure supplement 1A),
indicating that GLS is indeed the predominant isoform in murine retina. Furthermore,
retinal sections stained for GLS using immunofluorescence showed that GLS is expressed
throughout the retina with enrichment in the PR inner segments (Figure 1 – figure
supplement 1B). It has been shown that GLS activity is at least two-fold higher in the PR
inner segments, which are rich in mitochondria (Ross et al., 1987). Accordingly, GLS

segregated to the mitochondrial-enriched fraction more than the cytosolic fraction in the
retina (Figure 1 – figure supplement 1C).
129
We generated a rod PR-specific, *Gls* conditional knockout mouse to determine the role of
GLS activity in PR survival and function. Animals homozygous for a floxed *Gls* allele and
expressing a Cre-recombinase under the control of the rhodopsin promoter (*Gls$^{fl/fl}$;Rho-*
Cre$^+$, cKO) as well as animals expressing only the Cre-recombinase (*Gls$^{wt/wt}$;Rho-Cre$^+$*,
WT) were generated. Total retinal lysate was collected from cKO and WT animals at
postnatal day 14 (P14) and GLS expression was measured (Figure 1A). These data show
that cKO animals have significantly less GLS expression compared to WT animals. Whole
eyes were collected at P14 and stained for GLS using immunofluorescence. Figure 1B
shows significant loss of GLS expression in PR inner segments of cKO animals compared
to WT animals. The remaining GLS in the inner segment layer is found within cone PRs,
as shown in Figure 1B (arrows), confirming the knockout is specific to rod PRs. A lack of
compensatory upregulation of *Gls2* in the retina was confirmed via qRT-PCR and in PRs
using immunofluorescence (Figure 1 – figure supplement 1D-E).
143
Loss of GLS causes rapid PR degeneration
These data clearly show that GLS is significantly downregulated in cKO animals. To
determine if loss of GLS effects PR survival, optical coherence tomography (OCT) was
used to measure the *in vivo* thickness of each retinal layer at various timepoints in cKO
and WT animals (Figure 1C). At P14, cKO and WT animals are indistinguishable in total
retinal and outer nuclear layer (ONL) thickness (Figures 1D and 1E). However, by P21 a
significant loss in total retinal, ONL, and IS/OS thickness is observed (Figures 1D-1F) and
cKO animals continue to experience loss of retinal, ONL and IS/OS thickness out to P84.
Considering mice open their eyes around P14, we assessed if light exposure was
contributing to PR degeneration. cKO mice were reared in the dark, but no change in the
rate of ONL degeneration was observed when comparing dark-reared mice to those
reared in 12-hour light/12-hour dark cyclic lighting conditions (Figure 1 – figure supplement
2). Histology at P14, P21, and P42 confirmed loss of PR cell bodies (Figures 1G and 1H).
As has been seen in other models of PR degeneration, glial fibrillary acidic protein (GFAP),
which is a marker of stress-induced Müller glial cell activation, was increased in the retina
of the cKO mouse at P21 and P42 (Figure 1 – figure supplement 3) (Grenell et al., 2019).
160
Next, to assess the cell death pathways contributing to PR degeneration in the cKO
mouse, retinal sections from P21 animals were stained for TUNEL (Figure 1I). At P21,
there is a significant increase in TUNEL positive outer retinal cells in cKO mice compared
to WT mice (Figure 1J). qRT-PCR analysis of cell death-related gene expression at P14
(Figure 1K) demonstrated an increase in genes involved in apoptosis, necroptosis, and
ferroptosis. These data coincide with the thought that apoptosis is the predominant
mechanism of PR death in many retinal diseases, considering *Casp8* showed the greatest
increase in expression in the cKO retina, but also that other cell death mechanisms can
contribute to PR cell death (Wubben et al., 2016).
170
Retinas were stained for RHO expression using immunofluorescence, which indicated rod
OSs were shorter after *Gls* knockout (Figure 1 – figure supplement 4A). Because cKO
mice demonstrated shorter OSs, transmission electron microscopy (TEM) was used to
investigate the ultrastructure of cKO rod OSs (Figures 1 – figure supplement 4B-C). The
rod OSs in the cKO mouse appeared largely normal with well-organized, stacked disc

membranes that maintained interdigitation with the RPE similar to the WT mouse, just
shorter than WT rod OSs.
178
A deficiency in GLS activity could lead to a significant decrease in Glu, the main
neurotransmitter used by PRs. TEM was utilized to assess the ultrastructure of rod PR
ribbon synapses, which appear structurally intact in the remaining rod PRs of the cKO
retina at P21 (Figure 1 – figure supplement 4D). To further evaluate the synaptic
connectivity between PRs and the inner retina, retinal sections from cKO and WT mice at
P14 were labeled with wheat germ agglutinin (WGA), a plant lectin that binds N-
acetylglucosamine and sialic acid residues, to highlight PR synaptic membranes and non-
synaptic membranes (McLaughlin et al., 1980). Retinal sections from P14 WT and cKO
mice stained with WGA did not demonstrate significant differences in the labeling of PR
synaptic membranes in the outer plexiform layer (OPL) (Figure 1 – figure supplement 5A).
Staining of P14 retinal sections from WT and cKO mice with an antibody against Bassoon,
which labels the ribbon synapses of rods and cones in the OPL (Kutsyr et al., 2021), did
not demonstrate differences in the synaptic connectivity between PRs and second-order
neurons (Figure 1 – figure supplement 5B). Additionally, retinal sections at P14, P21, and
P42 were stained with antibodies against major cell-types in the inner retina, such as
amacrine, ganglion, and bipolar cells, and no significant alterations in immunofluorescent
patterns to suggest an inner retinal developmental abnormality were observed between
cKO and WT animals (Figure 1 – figure supplement 5C). Accordingly, inner retinal thinning
was not observed until P42 (Figure 1 – figure supplement 5D), when approximately 25%
of cells remained in the ONL (Figure 1H). This thinning of the inner retinal area is likely
secondary to the primary PR degeneration similar to that seen in other mouse models of
retinal degeneration (Ueta et al., 2012).
201
Loss of *Gls* in rod photoreceptors impairs retinal function
Loss of IS/OS thickness, shorter OSs, and loss of PR cell bodies can result in functional
loss. Thus, electroretinography (ERG) analysis was performed on cKO and WT animals
at P21 and P42 (Figure 2). As expected, cKO animals show a significant loss in rod-driven
scotopic a- and b-wave amplitudes at P21, which are further decreased by P42 (Figure
2A-2C). Interestingly, a significant loss in photopic b-wave amplitude was also found at
P42, suggesting a cone PR defect (Figure 2D). Immunofluorescent staining found that
cone outer segments shorten as rod degeneration progresses from P14 to P42 (Figure
2E) supporting secondary cone degeneration, which is a common phenotype associated
with rod-mediated retinal degeneration (Caruso et al., 2020).
212
GLS is necessary for maintenance of mature photoreceptors
In mice expressing a Cre-recombinase under the control of the rhodopsin promoter, Cre-
mediated excision of floxed genomic DNA has been observed as early as P7 (Le et al.,
2006). To ensure the function of GLS is not restricted to this early phase of PR
development and maturation and that it is critical to fully developed PRs as well, mice
homozygous for a floxed *Gls* allele (*Gls^{fl/fl}*) and expressing an inducible Cre-recombinase
under the control of the *Pde6g* promoter (*Gls^{fl/fl};Pde6g-Cre^{ERT2}*) (Koch et al., 2015) as well
as animals expressing only the inducible Cre-recombinase (*Gls^{wt/wt};Pde6g-Cre^{ERT2}*) were
generated. This allowed for induction of cre-recombinase activity in fully-mature PRs,
specifically, by the administration of tamoxifen (TAM). Animals heterozygous for *Pde6g-*
Cre^{ERT2} and homozygous for either the WT or floxed *Gls* allele (IND-cKO) were generated,
and TAM was administered intraperitoneally for 5 consecutive days starting at P22 (Figure
3) as retinal development is typically considered complete by P21 (Zhou et al., 2021).
Significant reduction of GLS in the retina of IND-cKO mice after TAM induction was

confirmed via Western blot and in PRs with immunofluorescence analysis (Figures 3A and
3B). Ten days after TAM induction, IND- cKO mice began to show a thinning of the IS/OS
layer that preceded and then paralleled the rapid ONL degeneration observed on
longitudinal OCT and histology (Figures 3C-3G). Hence, GLS is also critical for the survival
of fully developed PRs.
232
Beyond survival, GLS is also critical for PR function. ERG analyses performed 10 days
after TAM induction, a time prior to major structural changes in the outer retina,
demonstrated statistically significant reductions in the IND-cKO scotopic a- and b-waves
as compared to the WT (Figure 3H). Similarly, photopic ERG demonstrated statistically
significant decreases in the b-wave of the IND-cKO retina (Figure 3I). These data suggest
that GLS-driven Gln catabolism plays a significant role not only in rod PR survival but their
function as well. Additionally, *Pde6g* is expressed by rods to a significant degree but also
by cones (Voigt et al., 2020). Therefore, the IND-cKO likely knocks out GLS from both
rods and cones, which is in accordance with the immunofluorescence image in Figure 3B
where GLS is not observed in rod or cone inner segments unlike in Figure 1B where GLS
remains in cones. Hence, the reduction in IND-cKO photopic b-wave may suggest that
GLS-driven Gln catabolism in cones impairs their synaptic transmission.
245
246
GLS knockout does not alter nucleotide metabolism
Gln, and its breakdown product, Glu, have several fates in cellular metabolism (Figure
4A), which may underly the rapid and significant PR degeneration observed when GLS is
knocked out of rod PRs. PRs require nucleotides to support phototransduction as well as
transcriptional efforts for the continued synthesis of OSs. The catabolism of Gln to Glu by
GLS produces an ammonium ion that can be used for the synthesis of nucleotides (Yoo
et al., 2020). Additionally, inhibition of GLS has previously been demonstrated to inhibit
pyrimidine and purine biosynthesis (Alkan et al., 2018; Okazaki et al., 2017). To explore
the role of GLS-initiated Gln catabolism in nucleotide metabolism, *in vivo* targeted liquid
chromatography-tandem mass spectrometry (LC-MS/MS) metabolomics were performed
on P14 retina from WT and cKO animals (Supplementary File 2). No differences in key
purine or pyrimidine metabolism intermediates, such as ribose 5-phosphate (R5P), inosine
monophosphate (IMP), and uridine monophosphate (UMP), were observed between the
WT and cKO retina (Figure 4B), suggesting that GLS-mediated catabolism of Gln is not a
critical pathway for nucleotide synthesis in rod PRs.
262
Redox balance is altered upon knockout of GLS in rod photoreceptors
Gln can also be used for the generation of the antioxidant molecule glutathione as well as
in a non-canonical NADPH generating pathway (Son et al., 2013; Yoo et al., 2020).
Glutathione is a tripeptide of glutamate, glycine, and cysteine. Glutamate is a product of
the GLS reaction, and previous studies have demonstrated that inhibiting GLS reduces
glutathione levels and increases reactive oxygen species (Li et al., 2015). As such, the
NADP$^+$/NADPH ratio was statistically significantly increased by 16% in the P14 cKO retina
as compared to WT (Figure 4C), and the relative abundance of oxidized glutathione
(GSSG) was decreased in the P14 cKO retina as compared to WT (Figure 4D). While an
increased NADP$^+$/NADPH ratio in the cKO retina may imply that the abundance of GSSG
should be increased in the cKO retina since reduced glutathione can be recovered from
the oxidized form by the conversion of NADPH to NADP$^+$, the levels of GSSG have been
seen to be reduced in other models where GLS is pharmacologically inhibited or
genetically knocked down (Daemen et al., 2018). The lower levels of GSSG may signal
an overall reduction in the biosynthesis of glutathione considering the product of the GLS

reaction, glutamate, is directly necessary for its synthesis as well as indirectly responsible
for cysteine via the SLC7A11 (xCT) cystine/glutamate antiporter (Yoo et al., 2020). The
expression of *Slc7a11* was reduced in the P14 cKO retina (Figure 4E). Hence, the
biosynthesis of glutathione in rod PRs may be affected directly and indirectly when GLS
is knocked out. In accordance with the antioxidants NADPH and glutathione potentially
being reduced, the expression of the antioxidant enzymes *Sod1* and *Sod2*, which detoxify
superoxide radicals, were increased in the P14 cKO retina (Figure 4E). These data
suggest that GLS-mediated Gln catabolism regulates redox balance in rod PRs, and the
altered redox balance that is a consequence of its knockout may be one factor contributing
to PR degeneration in the GLS cKO mouse.
288
Relative abundance of TCA cycle intermediates and mitochondrial function mostly
unchanged in *Gls* cKO retina
While the changes in certain redox elements were statistically significant, they were
modest alterations and may not account in full for the significant and rapid PR
degeneration noted in the GLS cKO mouse. Many cancer cells that utilize aerobic
glycolysis rely on Gln to replenish TCA cycle intermediates, which maintain oxidative
metabolism and provide biosynthetic precursors. Considering PRs utilize aerobic
glycolysis, similar to cancer cells, and have significant bioenergetic demands, we next
sought to determine if rod PRs also depend on GLS-initiated glutaminolysis for TCA cycle
metabolites and oxidative metabolism. Interestingly, LC-MS/MS-based targeted
metabolomics demonstrated very few changes in the relative pool sizes of TCA cycle
metabolites in the cKO compared to WT retina at P14 with only malate showing a
statistically significant decrease (Figure 4F). Since GLS is enriched in the PR mitochondria
(Figure 1 – figure supplement 1B) and previous studies demonstrated that inhibiting GLS
in certain cell lines reduces mitochondrial function (Alkan et al., 2018), mitochondrial
stress tests were also performed on P14 cKO and WT retina, prior to PR loss, using the
BaroFuse (Kamat et al., 2023). The basal oxygen consumption rate (OCR) as well as the
changes in OCR in response to oligomycin or carbonyl cyanide 4-
(trifluoromethoxy)phenylhydrazone (FCCP) were not statistically significantly different
between WT and cKO retina (Figures 4G and 4H). The expression of complexes involved
in oxidative phosphorylation were also unchanged between the WT and cKO retina at P14
(Figure 4I).
311
As stated earlier, a Cre-recombinase under the control of the rhodopsin has been shown
to be activated as early as P7 (Li et al., 2005). To circumvent this early knockout of GLS,
rod PRs may rewire their metabolism, as has been seen in other conditional knockout
mouse models with this Cre-recombinase system (Subramanya et al., 2023; Wubben et
al., 2017), to utilize different fuel sources for the maintenance of the TCA cycle and
mitochondrial function. To this end, the expression of genes involved in glycolysis,
pyruvate metabolism, and the TCA cycle were examined in the P14 cKO and WT retina
before PR degeneration. Numerous genes were significantly altered in the cKO retina
across the metabolic pathways (Figure 4 – figure supplement 1). The expression of
multiple genes in glycolysis and pyruvate metabolism were increased (Figure 4 – figure
supplement 1A), possibly suggesting that rod PRs are stimulating glucose oxidation to
maintain the TCA cycle and mitochondrial function similar to that seen in certain cancer
cells when GLS is inhibited (Okazaki et al., 2017). In contrast, the only significant change
in expression of these genes in eyecup tissue was *Pdk4,* with a significant decrease in
cKO vs WT mice (Figure 4 – figure supplement 2). However, targeted metabolomics
demonstrated only a minor change in glycolytic intermediates between the cKO and WT
retina at P14 (Figure 4F). As targeted metabolomics at the P14 timepoint provides only a

snapshot of the pool size, additional analyses with stable isotope tracing metabolomics are necessary to assess nutrient utilization, in this case glucose. Uniformly-labeled, $^{13}C_6$-glucose was intraperitoneally injected in P14 WT and cKO mice, and the retina harvested 45 minutes later for metabolomic analysis via LC-MS/MS (Yuan et al., 2019). No change in the fractional labeling of glycolytic intermediates was observed between the WT and cKO retina, and more so, an increase in the fractional labeling of TCA cycle intermediates was not observed in the cKO retina (Figures 5A, 5B and Figure 5 – figure supplement 1). Hence, it is unlikely rod PRs are increasing glucose oxidation to maintain the levels of TCA cycle intermediates and mitochondrial function.

Gls cKO in rod photoreceptors decreases the utilization of Gln in the TCA cycle

We initially anticipated that GLS knockout in rod PRs would significantly alter the relative TCA cycle metabolite pools in the cKO retina, similar to what is seen in Gln-dependent cancer cells (Daemen et al., 2018; Yang et al., 2017), as PRs and cancer cells have common metabolic relationships (Du, Cleghorn, Contreras, Linton, et al., 2013; Ng et al., 2015; Rajala, 2020), and previous *ex vivo* studies demonstrated Gln can supplement the TCA cycle in the retina (Grenell et al., 2019; Tsantilas et al., 2021). However, as observed in Figure 4F, very few changes were observed in the relative pool sizes of TCA cycle metabolites in the cKO compared to WT retina. To delineate how Gln is metabolized, we used LC-MS/MS to trace the metabolic fate of uniformly-labeled $^{13}C_5$-Gln in the WT and cKO retina *in vivo* at P14. Uniformly-labeled $^{13}C_5$-Gln was intraperitoneally injected into the mice and the retina harvested 45 minutes later. In accordance with GLS being the enzyme that initiates glutaminolysis (Yang et al., 2017), the fractional labeling of glutamate, TCA cycle intermediates, and pyruvate was decreased in the P14 cKO retina as compared to WT (Figures 5C, 5D and Figure 5 – figure supplement 2). Previous *ex vivo* studies have also observed that isotopically labeled $^{13}C_5$-Gln can contribute to m+3 pyruvate in the retina possibly via the decarboxylation of m+4 malate by malic enzyme (Grenell et al., 2019).

a-ketoglutarate (α-KG) is the main entry point for Gln into the TCA cycle (Figure 5C) and has been shown to be a key metabolite in Gln metabolism (Yang et al., 2017). *In vivo* stable isotope tracing demonstrated decreased incorporation of Gln carbons into α-KG in the cKO retina compared to the WT (Figure 5D). Furthermore, supplementation with α-KG has been shown to improve PR survival in mouse models of inherited retinal disease (Wert et al., 2020). So, we attempted to rescue the PR degeneration phenotype in the GLS cKO mouse by supplementing cKO animals with 10 mg/mL of α-KG in the drinking water starting at P4 (Rowe et al., 2021; Wert et al., 2020). A small, but significant increase in ONL thickness was identified in α-KG-treated animals at P22 using OCT (Figure 5E). These data further suggest that Gln's role in supporting the TCA cycle is not the major mechanism by which PRs utilize Gln to suppress PR apoptosis.

Gls cKO retina has reduced NEAA levels and an upregulated integrated stress response

Gln and GLS-derived Glu play a central role in the biosynthesis of several NEAAs (Yoo et al., 2020). So, the disruption of GLS-initiated Gln catabolism may be causing a disruption in available NEAAs for biomass production. To this end, targeted metabolomics on the cKO retina at P14 showed a significant increase in the substrate of GLS, Gln, and a significant decrease in its NEAA product Glu, further illustrating loss of GLS function (Figures 6A and 6B). The NEAA Asp was also significantly reduced (Figure 6A), which is consistent with previous studies that inhibited GLS in cancer cells (Alkan et al., 2018; Okazaki et al., 2017). Notably, similar decreases in Glu and Asp were noted in IND-cKO

compared to WT retina at a point prior to PR degeneration (Figure 6 – figure supplement 1).

Deprivation of amino acids can activate the integrated stress response (ISR), which responds to an array of stressors to maintain cellular homeostasis and particularly, protein homeostasis. Acutely, ISR activation can protect cells by temporarily halting protein synthesis. However, chronic ISR activation and global protein synthesis inhibition can trigger apoptosis (Pakos-Zebrucka et al., 2016). ISR activation leads to phosphorylation of the alpha subunit of eukaryotic translation initiation factor 2 (eIF2α), which reduces global protein synthesis while preferentially allowing for the translation of activating transcription factor 4 (ATF4) (Pakos-Zebrucka et al., 2016). Considering the reduced levels of NEAAs Glu and Asp, retinas from P14 GLS cKO animals were assayed for these ISR components using Western blotting (Figures 6C and 6D), which demonstrated increased levels of both phosphorylated eIF2α and total ATF4, suggesting ISR activation. To determine if global protein synthesis was affected in GLS cKO retinas, we applied the SUnSET (SUrface SEnsing of Translation) method to quantify protein synthesis in WT and cKO retinas at P14 (Fort et al., 2022). These data show a significant decrease of puromycin incorporation into nascent polypeptide chains, indicating a decrease in global protein synthesis consistent with ISR activation (Figures 6E and 6F). To determine if PR degeneration could be rescued by inhibiting the ISR, ISRIB (2.5 mg/kg) was administered systemically daily starting at P5. In WT mice, ISRIB treatment did not impact retinal anatomy as assessed by OCT at P21 (Figure 6 – figure supplement 2A). Notably, at P21, ISRIB treatment in the cKO mouse significantly increased ONL thickness compared to vehicle using OCT (Figure 6G) and histology (Figure 6 – figure supplement 3A), and this effect was sustained at P28 (Figure 6 – figure supplement 2B). These results suggest that ISR activation is downstream of Gln catabolism and contributes to PR degeneration in the cKO mouse retina.

Previous work has demonstrated that when certain cancer cell lines are faced with Gln deprivation, α-KG alone does not restore cell proliferation (Pavlova et al., 2018; Son et al., 2013). Rather, NEAA supplementation is necessary to restore proliferation in these reports. Similarly, α-KG supplementation had an incomplete survival impact on PR survival, showing approximately 10% increase in ONL thickness in cKO animals (Figure 5E). Since systemic or intravitreal glutamate supplementation has demonstrated retinal toxicity (Sisk & Kuwabara, 1985), and Asp concentration is normally low in plasma with poor cellular membrane permeability (Sullivan et al., 2018), we investigated other NEAA approaches to rescue the PR degeneration observed in the GLS cKO mouse. The potential of asparagine (Asn) to rescue PR degeneration was chosen due to the growing list of Asn-mediated cellular processes that improve cancer cell survival and growth under metabolic stress, such as Gln deprivation. These Asn-mediated cellular processes include inhibiting ISR-induced apoptosis (Zhang et al., 2014),(1) promoting protein synthesis (Pavlova et al., 2018), and maintaining Asp pools (Halbrook et al., 2022). All these processes may be relevant to the PR degeneration observed in the cKO mouse. Following administration of systemic Asn (200 mg/kg) daily starting at P5, *in vivo* analysis of retinal structure via OCT and histology at P21 demonstrated a significant improvement in ONL thickness compared to animals treated with vehicle, with a 22-26% increase in ONL thickness in some retinal locations (Figure 6H and Figure 6 – figure supplement 3B).

Discussion

In this study, we demonstrate for the first time the importance of Gln catabolism in PR metabolism, function, and survival *in vivo*. Rod PRs lacking GLS demonstrated rapid

and complete degeneration with concomitant functional loss. Mechanistically, lack of GLS-
initiated Gln catabolism in rod PRs, specifically, resulted in changes in redox homeostasis
and a decrease in the fractional labeling of TCA cycle intermediates from stable isotope-
labeled Gln *in vivo*. However, Gln anaplerosis did not appear to be driving the TCA cycle
as targeted metabolomics showed few changes in the relative levels of TCA cycle
intermediates, and supplementation with αKG showed a modest rescue effect in PR
degeneration. The NEAAs Glu and Asp were found to be significantly decreased in the
cKO mouse retina, and in accordance with this AA deprivation, the ISR was upregulated
with a reduction in global protein synthesis suggesting Gln catabolism in rod PRs plays a
significant role in supporting biomass production via NEAA synthesis. Interestingly,
supplementing cKO mice with Asn significantly rescued PR degeneration, revealing a
novel link between Gln and Asn metabolism in PRs. This work further demonstrates that
PRs have the flexibility to utilize fuel sources other than glucose to meet their metabolic
needs and that Gln is a critical amino acid that supports PR cell biomass, redox balance,
and survival.
The rapid degenerative phenotype observed with conditional deletion of *Gls* in rod
PRs was unexpected considering glucose is viewed as the primary substrate for the retina
and PRs (Swarup et al., 2019). The *Gls* cKO mouse demonstrated comparable ONL
thickness to WT controls at P14 with approximately 50% loss of PR cell bodies by P21
(Figure 1 – figure supplement 3A-B) and near complete PR degeneration by P84 (Figure
1C). In contrast, conditional deletion of numerous enzymes within central glucose
metabolism, including hexokinase 2 (HK2) (Weh et al., 2020), pyruvate kinase M2 (PKM2)
(Wubben et al., 2017), glucose transporter 1 (GLUT1) (Daniele et al., 2022), mitochondrial
pyruvate carrier 1 (MPC1) (Grenell et al., 2019), and lactate dehydrogenase A (LDHA)
(Rajala et al., 2023), results in a slower, age-related degeneration. For example, loss of
GLUT1 in the retina and rod PRs does not demonstrate 50% thinning of the ONL until
approximately 4 months of age (Daniele et al., 2022), and the others noted above
demonstrate even slower outer retinal degeneration. It has also been shown that PRs can
utilize fatty acid β-oxidation for energy (Joyal et al., 2016). Interestingly, in retinas lacking
the very-low-density lipoprotein receptor (VLDLR), which facilitates fatty acid uptake, the
PRs remained largely intact. In any of these transgenic mouse models, PRs may use other
transporters to take up fatty acids or glucose, rewire their metabolism, or utilize metabolic
redundancies to maintain metabolic homeostasis and stave off degeneration
(Subramanya et al., 2023; Wubben et al., 2017). Our data show that any metabolic
reprogramming that is occurring in the cKO mouse retina appears unable to significantly
circumvent the significant and rapid PR degeneration suggesting the importance of Gln
catabolism in rod PRs. Furthermore, inducing GLS knockdown in mature PRs also
demonstrated rapid PR degeneration (Figure 3).
Traditionally, it has been thought that Gln maintains cell survival through Gln-
derived α-KG and maintenance of the TCA cycle (Zhang et al., 2014).(1) While stable
isotope tracing with uniformly labeled Gln demonstrated a reduction in fractional labeling
of the TCA cycle intermediates (Figure 5D), targeted metabolomics showed few changes
in the relative levels of the intermediates (Figure 4F) and supplementation with αKG did
not rescue PR degeneration to a large degree (Figure 5E). Gln is also required for the
biosynthesis of NEAAs, and the NEAAs Glu and Asp were reduced nearly two-fold in the
retina of cKO mouse (Figure 6A). Glu is the product of the GLS reaction so it is not
surprising that Glu was substantially reduced. Glu is involved in glutathione biosynthesis
directly and indirectly, can be converted into α-KG to enter the TCA cycle, and is crucial
for the biosynthesis of NEAAs including Asp (Yoo et al., 2020). Alterations in redox
homeostasis were observed in the retina of the cKO mouse in accordance with the
reduced level of Glu (Figures 4C-4E). While these redox imbalances likely contribute to

the PR degeneration observed, it is unlikely these small but statistically significant
changes fully account for the rapid and complete PR degeneration. Future rescue studies
with antioxidants, such as N-acetylcysteine, are needed to shed light on the role of redox
imbalance in this novel transgenic mouse model.
Previous *ex vivo* studies demonstrated that Gln can contribute carbons to Asp
synthesis via the TCA cycle and/or contribute a nitrogen via glutamate-oxaloacetate
transaminases (Du, Cleghorn, Contreras, Lindsay, et al., 2013; Xu et al., 2020), the latter
of which have been shown to be critical in the Gln metabolic rewiring of cancer cells (Son
et al., 2013) and more recently, in rod PR metabolism, function, and survival (Subramanya
et al., 2023). Our data shows that Gln-derived carbons via Glu are entering the TCA cycle
in vivo and contributing to the synthesis of Asp (Figure 5), and this contribution is reduced
in the cKO mouse retina. Based on the previous studies noted above, Gln may also be
contributing to Asp synthesis via the glutamate-oxaloacetate transaminase catalyzed
transfer of a nitrogen from Glu. Future studies using amine-labeled Gln are needed to
dissect the contributions of Gln-derived carbon versus nitrogen to the synthesis of Asp in
the retina *in vivo*.
Asp is a proteinogenic amino acid that has many biosynthetic roles (Alkan et al.,
2018), and *ex vivo* neural retina studies have suggested that the retina needs Asp to
maintain its metabolic homeostasis (Li et al., 2020). In accordance with this data, Asp was
one of the few metabolites that was significantly reduced in the retina of cKO mouse prior
to rapid PR degeneration (Figure 6A). Based on *ex vivo* studies, it has been postulated
that Asp utilization in the retina is necessary to maintain aerobic glycolysis and
mitochondrial metabolism through the recycling of NADH to NAD^+, shuttling electrons into
the mitochondria via the malate-aspartate shuttle, replenishing oxaloacetate for
biosynthesis and/or producing pyruvate via malic enzyme (Li et al., 2020). Our *in vivo*
metabolomics data further support this last point as uniformly labeled Gln contributed to
m+3 pyruvate in the retina possibly via the decarboxylation of m+4 malate by malic
enzyme (Figure 6D). The m+4 isotopologue of malate could arise from m+4 Asp via
glutamate-oxaloacetate transaminase (GOT) and malate dehydrogenase (MDH) as part
of the malate-aspartate shuttle. Furthermore, the decarboxylation of malate by malic
enzyme produces pyruvate and also NADPH. A previous study revealed high malic
enzyme activity in the retina and suggested it is capable of producing NADPH in the retina
(Winkler et al., 1986). The observation that malate was reduced (Figure 4F) and the
$NADP^+$/NADPH ratio was increased (Figure 4C) in the cKO retina further supports a
metabolic pathway in PRs where the Gln carbon skeleton is converted into Asp, then OAA,
malate, and finally, pyruvate via the cytosolic enzymes of the malate-aspartate shuttle and
malic enzyme. This non-canonical Gln metabolism pathway is also necessary to sustain
cancer cell growth (Son et al., 2013).
The m+4 isotopologue of oxaloacetate (OAA) can also be converted to m+3
pyruvate by phosphoenolpyruvate carboxykinase (PEPCK). A recent *ex vivo* study
postulated that PRs can utilize different metabolic cycles, such as the Cahill cycle or mini-
Krebs cycle, to uncouple glycolysis from oxidative phosphorylation (Chen et al., 2024).
These cycles are fueled by Gln and require PEPCK to replenish pyruvate from OAA. While
the lack of m+4 citrate in both the P14 WT and cKO retina and decrease in m+2 citrate in
the cKO retina (Figure 5 – figure supplement 2) may further support the existence of these
previously postulated metabolic cycles, which avoid the citrate synthase step of the TCA
cycle as well as others, further studies are needed to confirm the activity of these pathways
in the retina and specifically, the PRs.
Interestingly, we did not observe any significant changes in glycolysis or
mitochondrial function in the cKO retina despite a reduction in the relative level of Asp. It
is hypothesized that Asp catabolism sustains glycolysis and oxidative phosphorylation in

the retina by recycling NADH to NAD$^+$ and shuttling electrons into the mitochondria via the
malate-aspartate shuttle (Li et al., 2020). However, in the GLS cKO retina, the
NAD$^+$/NADH ratio ($P = 0.125$) and its oft-used proxy, the pyruvate/lactate ratio ($P = 0.192$),
were not statistically significantly altered. Compensatory metabolic rewiring, which has
been seen in other conditional knockout mouse models using the same Cre-recombinase
system (Subramanya et al., 2023; Wubben et al., 2017), may be maintaining the
NAD$^+$/NADH ratio in the cKO retina at P14 when there are equal numbers of PRs in the
WT and cKO retina. To this end, the expression of the *Mdh1* gene, which is a component
of the malate-aspartate shuttle that recycles NADH to NAD$^+$ in the cytoplasm, was
upregulated in the cKO retina (Figure 4- figure supplement 1). The conversion of
dihydroxyacetone phosphate (DHAP) to glycerol 3-phosphate via glycerol 3-phosphate
dehydrogenase is an alternative pathway to regenerate NAD$^+$. DHAP levels were
significantly reduced in the cKO retina possibly suggesting increased activity of this
pathway (Figure 4F). Yet, the gene expression and activity of glycerol 3-phosphate
dehydrogenase in the retina has been shown to be low (Adler & Klucznik, 1982; Voigt et
al., 2020).
Asp is also the immediate precursor to Asn. Intracellular levels of Asn are the
lowest among the NEAAs (Zhang et al., 2014), so it was not surprising that Asn was below
the level of confident detection in our targeted metabolomics analyses on the WT or cKO
retina. Asn had been shown to rescue cancer cells from ISR-induced apoptosis, increase
protein synthesis during Gln deprivation, and protect the Asp pool (Halbrook et al., 2022;
Pavlova et al., 2018; Zhang et al., 2014). As metabolic similarities exist between PRs and
cancer cells (Du, Cleghorn, Contreras, Linton, et al., 2013; Ng et al., 2015), and the *Gls*
cKO mouse retina demonstrated ISR activation and apoptosis (Figures 1G-1I and 6C-6D),
decreased protein synthesis (Figure 6E), and decreased Asp (Figure 6A), Asn
supplementation was explored and proved effective as observed in the rescue of PR
degeneration in the cKO mouse (Figure 6H). These results offer the first evidence for a
role of Asn downstream of Gln metabolism in PR survival, but further studies are
necessary to define which of the Asn-mediated processes is crucial for PR neuroprotection
following targeted *Gls* knockout. Of note, while Asn supplementation provided a greater
PR neuroprotective effect than α-KG, the two supplements had different routes of
administration with Asn being provided intraperitoneally as previously described (Xu et al.,
2021) and α-KG being provided in the drinking water as had previously improved PR
survival in a mouse model of inherited retinal disease (Wert et al., 2020). It is unclear if
supplementing α-KG to GLS cKO animals intraperitoneally would further boost retinal
protection. Yet, the relative abundance of α-KG and most other TCA cycle intermediates
were unchanged between WT and cKO retinas suggesting Gln may not be driving the
TCA cycle in PRs.
The ISR is activated with reduced global protein synthesis in the *Gls* cKO mouse
retina (Figures 6C-6F) and inhibiting the ISR with ISRIB delayed PR degeneration (Figure
6G). ISR activation is a hallmark of neurodegenerative diseases including retinal
degenerative diseases (Gorbatyuk et al., 2020). Chronic activation of the ISR and protein
synthesis attenuation has been observed in a multitude of preclinical models of retinal
degeneration and shown to contribute to PR degeneration (Gorbatyuk et al., 2020).
Metabolic dysfunction and deprivation of key nutrients, such as glucose and amino acids,
are not only known stressors that activate the ISR (Pakos-Zebrucka et al., 2016), but also
underlying mechanisms of PR degeneration (Ait-Ali et al., 2015; Caruso et al., 2020; Pan
et al., 2021). There is a paucity of studies that examine the link between metabolism and
the ISR in retinal degenerative disease, which is a critical gap in our knowledge since
identification of molecular and metabolic pathways triggering PR death is likely to reveal
novel targets for therapeutic intervention. A recent study demonstrated that imbalanced

pro-apoptotic ISR signaling contributes to deoxysphingolipid-mediated toxicity in retinal
disease and identified potential therapeutic strategies, such as pharmacologic
enhancement of ATF6 activity or treatment with ATF6-regulated neurotrophic factor
MANF, that attenuate the associated retinal degeneration (Rosarda et al., 2023). The
novel transgenic mouse model of retinal degeneration described here provides a unique
tool to obtain further insight on the nexus of metabolism, ISR activation, and protein
synthesis attenuation in PR degeneration to identify pharmacologically and metabolically
tractable nodes for therapeutic intervention.
Collectively, our results indicate that Gln is critical for maintaining the pools of key
biosynthetic precursors, Glu and Asp, in rod PRs and disrupting Gln catabolism results in
profound loss of PR function and survival in part secondary to an imbalance in ISR
activation and protein synthesis attenuation. Glucose remains central in PR metabolism,
but improving our understanding of other metabolic pathways that support PR function
and survival and how these metabolic pathways connect with cell death mechanisms
could be transformative for preventing PR degeneration and vision loss in a multitude of
retinal degenerative diseases.
600
Ideas and Speculation
Beyond glucose, the metabolic pathways integral to photoreceptor health remain
largely unknown. This is a critical knowledge gap as identification of these pathways is
likely to reveal new strategies for therapeutic intervention. This work demonstrates that
rod photoreceptors depend on glutamine catabolism and suggests a metabolic axis where
glutamine catabolism in rod photoreceptors supports the production of aspartate and
asparagine to promote anabolism and prevent signaling through the pro-apoptotic
integrated stress response pathway. Considering activation of the integrated stress
response is a hallmark of neurodegenerative diseases and metabolic dysfunction
underlies photoreceptor degeneration, defining the pathways by which glutamine
catabolism contributes to photoreceptor health is likely to identify nodes that may be
targeted to make photoreceptors less vulnerable to stress.
613
614

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Genetic reagent (*Mus musculus*)	*Gls$^{fl/fl}$*	PMID: 26778975		
Genetic reagent (*Mus musculus*)	Rho-cre	PMID: 16636658		
Genetic reagent (*Mus musculus*)	*Gls$^{fl/fl}$;Rho-Cre^{+}*	This study		
Genetic reagent (*Mus musculus*)	*Pde6g-Cre^{ERT2}*	PMID: 26301813		
Genetic reagent (*Mus musculus*)	*Gls$^{wt/wt}$;Pde6g-Cre^{ERT2}*	This study		

Genetic reagent (*Mus musculus*)	*Gls^fl/fl;Pde6g-Cre^ERT2*	This study		
Antibody	(Mouse monoclonal) anti-GLS	Proteintech	66265-1-Ig	1:200; immunofluorescence
Antibody	(Rabbit polyclonal) anti-GLS2	Abcam	Ab113509	1:200; immunofluorescence
Antibody	(Mouse monoclonal) anti-Rhodopsin	Abcam	Ab5417	1:1,000; immunofluorescence
Antibody	(Goat polyclonal) anti-OPN1MW/LW	Santa Cruz Biotechnology	Sc-22117	1:200; immunofluorescence
Antibody	(Mouse monoclonal) anti-BRN3A	Santa Cruz Biotechnology	Sc-8429	1:100; immunofluorescence
Antibody	(Rabbit polyclonal) anti-calretinin	Millipore Sigma	C7479	1:100; immunofluorescence
Antibody	(Rat monoclonal) anti-GFAP	Thermo Fisher	13-300	1:200; immunofluorescence
Antibody	(Mouse monoclonal) anti-Chx10	Santa Cruz Biotechnology	Sc-365519	1:200; immunofluorescence
Antibody	(Mouse monoclonal) anti-Bassoon	Enzo	SAP7F407	1:1,000; immunofluorescence
Antibody	Goat-anti-mouse Alexa 488	Invitrogen	A11001	1:1,000; immunofluorescence
Antibody	Donkey-anti-mouse Alexa 594	Jackson ImmunoResearch Laboratories	715-585-151	1:500, immunofluorescence
Antibody	Donkey-anti-rabbit Alexa 594	Jackson ImmunoResearch Laboratories	711-585-152	1:500; immunofluorescence
Antibody	Donkey-anti-goat Alexa 647	Invitrogen	A21447	1,2000; immunofluorescence

Antibody	Donkey-anti-rat Alexa 594	Invitrogen	A21209	1:500; immunofluorescence
Antibody	PNA Alexa 594 conjugate	Invitrogen	L32459	1:200; immunofluorescence
Antibody	WGA Alexa 594 conjugate	Invitrogen	W11262	1:1,000; immunofluorescence
Antibody	(Rabbit polyclonal) anti-KGA-specific GLS	Proteintech	20170-1-AP	1:1,000; western
Antibody	(Rabbit polyclonal) anti-GAC-specific GLS	Proteintech	19959-1-AP	1:1,000; western
Antibody	(Rabbit polyclonal) anti-GLS	Proteintech	12855-1-AP	1:1,000; western
Antibody	(Rabbit polyclonal) anti-VDAC	Cell Signaling Technology	4866	1:1,000; western
Antibody	(Mouse monoclonal) Anti-TIM23	BD Biosciences	611223	1:1,000; western
Antibody	(Mouse monoclonal) anti-HSP90	Cell Signaling Technology	4877	1:2,000; western
Antibody	Horse-anti-mouse-HRP	Cell Signaling Technology	7076	1:5,000; western
Antibody	Goat-anti-rabbit-HRP	Cell Signaling Technology	7074	1:5,000; western
Antibody	(Rabbit monoclonal) anti-eIF2α	Cell Signaling Technology	5324	1:1,000; western
Antibody	(Rabbit monoclonal) anti-Phospho-eIF2α	Cell Signaling Technology	3398	1:5,000; western

Antibody	(Rabbit polyclonal) anti-ATF4	Invitrogen	PA5-27576	1:1,000; western
Antibody	(Mouse monoclonal) anti-puromycin	Biolegend	381502	1:1,000; western
Antibody	(Mouse monoclonal) Total OXPHOS rodent antibody	Abcam	Ab110413	1:1,000; western

615
Methods and Materials
617

Animals
All animals were treated in accordance with the Association for Research in Vision and
Ophthalmology Statement for the Use of Animals in Ophthalmic and Vision Research. The
protocol was approved by the University Committee on Use and Care of Animals of the
University of Michigan (PRO00011133). All animals were housed under standard
husbandry conditions at room temperature in 12-hour light/12-hour dark cycles unless
explicitly stated in the text. Both male and female mice were used for all experiments. A
transgenic mouse where *Gls* is selectively deleted from rod photoreceptors was created
by crossing mice with Lox-P sites flanking exon 1 of the *Gls* gene (*Gls^{fl/fl}*, courtesy of Dr.
Stephen Rayport, Columbia University) with Rho-Cre mice, in which Cre-recombinase
expression is driven specifically in rod PRs. *Gls^{fl/fl}* and Rho-Cre mice have been previously
described (Le et al., 2006; Mingote et al., 2015). Animals were screened for the *rd8*
mutation (Mattapallil et al., 2012). Alpha-ketoglutarate (α-KG, Millipore-Sigma, St. Louis,
MO, USA, Cat # K1128) was provided to mice in their drinking water (10 mg/mL) starting
at P4. Asparagine (200 mg/kg, Millipore-Sigma, Cat # A4159) or vehicle (PBS) was
injected IP starting at P5. ISRIB (2.5 mg/kg, Caymen Chemical, Ann Arbor, MI, USA, Cat
16258) or vehicle (50% PEG 400, 43.4% saline, 6.6% DMSO) (Halliday et al., 2015) was
injected IP starting at P5. For inducible deletion of *Gls*, *Gls^{fl/fl}* mice were crossed to *Pde6g-*
Cre^{ERT2} mice (courtesy of Dr. Stephen Tsang, Columbia University) (Koch et al., 2015).
The Cre-recombinase was activated via IP injection of tamoxifen (Millipore-Sigma, Cat #
T5648) at a concentration of 100 mg/kg bodyweight for 5 consecutive days. Whole retinas
were extracted from animals using the "cut-and-pick" method as previously described
(Winkler, 1981), being careful to remove any adherent ciliary body or RPE before
processing. Total retina was then either snap frozen on dry ice (Western blotting,
metabolomics), immersed in RNAlater (Qiagen, Hilden, Germany, Cat # 76104) for qRT-
PCR, or used immediately (BaroFuse or NADP$^+$/NADPH bioluminescent assay).
644

In vivo functional and structural assessment
Visual function was assessed as previously described (Weh et al., 2020; Wubben et al.,
2017). Mice were anesthetized using a mixture of ketamine/xylazine (90/10 mg/kg) and
their pupils were dilated using 1% tropicamide and 2.5% phenylephrine ophthalmic drops.
Retinal function was determined using a Diagnosys Celeris ERG system (Diagnosys LLC,
Lowell, MA, USA) following overnight dark adaptation. *In vivo* retinal thickness was

measured using the Envisu-R SD-OCT imager (Leica Microsystems Inc., Buffalo Grove,
IL, USA). A 1.5 mm horizontal B-scan (1000 A-scans x 100 frames) and a 1.5 mm x 1.5
mm rectangular volume (1000 A-scans x 36 B-scans x 6 frames) were captured, registered
and averaged using the built-in software, and analyzed using the Diver 1.0 software suite
(Leica Microsystems). Images were segmented manually to determine total retinal, outer
nuclear layer, and combined inner segment/outer segment thickness. Measurements
were taken at 15 points on a 9x9 grid and averaged as previously described (Weh et al.,
2022).
659
Immunofluorescence
Mouse eyes were enucleated and immersed in 4% paraformaldehyde overnight before
embedding in paraffin and sectioned at 6 mm thickness. Following standard protocols,
sections were de-paraffinized and antigen retrieval performed as previously described
(Weh et al., 2020; Wubben et al., 2017). Sections were blocked with 1% bovine serum
albumin (BSA, Millipore-Sigma, Cat # A9647) in 1X phosphate buffered saline (PBS,
Thermo Fisher Scientific, Waltham, MA, USA, Cat # BP399) with 0.125% Tween 20
(Thermo Fisher Scientific, Cat # BP337) and 10% normal goat serum prior to incubating
with primary antibody in 1% BSA and 1% normal goat serum overnight at 4°C. Slides were
then washed, secondary antibody applied for 1 hour at room temperature before washing,
and finally, counterstained with DAPI (Thermo Fisher Scientific, Cat # P36930). Images
were obtained on a Leica DM6000 microscope with a 40X objective. The antibodies used
for immunofluorescence are found in the key resources table.
673
TUNEL staining and cell counts
TUNEL staining was performed as previously described using the DeadEnd kit (Promega,
Madison, WI, USA, Cat # G3250) (Wubben et al., 2017; Wubben et al., 2020). TUNEL-
positive cells were counted in a masked fashion and normalized to the total number of
nuclei using a custom ImageJ macro (Busov & Besirli, 2014; Wubben et al., 2017; Wubben
et al., 2020). Tissue sections through the plane of the optic nerve were also stained with
hematoxylin and eosin and the total number of nuclei in the ONL were determined after
normalization to inner retinal area (Wubben et al., 2020).
682
Sub-cellular fractionation
Whole retina was fractionated into cytosolic and post-cytosolic (mitochondrial enriched)
fractions as previously described (Weh et al., 2020) using the Subcellular Protein
Fractionation Kit for Tissues (Thermo Fisher Scientific, Cat # 87790). Both retinas from a
single animal were pooled and homogenized with a Dounce homogenizer in cytoplasmic
extraction buffer supplemented with protease (Halt™ Protease Inhibitor Cocktail, Thermo
Fisher Scientific, Cat # 87786) and phosphatase (Halt™ Phosphatase Inhibitor Cocktail,
Thermo Fisher Scientific, Cat # 78420) inhibitors. The retinal lysate was then centrifuged
at 10,000 x relative centrifugal force (RCF) for 10 minutes at 4°C. The resulting
supernatant was saved as the cytosolic fraction, and the resulting pellet was resuspended
in RIPA lysis buffer (Thermo Fisher Scientific, Cat # 89900) that included protease and
phosphatase inhibitors (Cell Signaling Technology, Danvers, MA, USA, Cat # 5872) and
sonicated at 20% amplitude with 1s on/off pulse for 10s. The lysate was centrifuged for 10
minutes at 10,000 x RCF at 4°C. The resulting supernatant was saved as the
mitochondrial enriched fraction. The percentage of GLS in each fraction was determined
using Western blotting.
699
Western blotting

Immunoblots were performed as previously described (Weh et al., 2020). Protein
estimation was performed using the Pierce™ BCA kit (Thermo Fisher Scientific, Cat #
23225). Equivalent micrograms of protein from each sample were diluted using 4X
Laemmli buffer (Bio-Rad, Hercules, CA, USA, Cat # 1610747) supplemented with β-
mercaptoethanol (Millipore-Sigma, Cat # M6250) before heating at 95°C for 5 minutes and
finally loaded onto 4-20% polyacrylamide gel (Bio-Rad, Cat # 4561094). Samples were
then transferred to a PVDF membrane using the Trans-Blot® Turbo™ Transfer System
(Bio-Rad, Cat # 1704150). Membranes were blocked using 5% non-fat milk powder diluted
in TBST (Tris-buffered Saline, Bio-Rad, Cat # 1706435, supplemented with 0.1% Tween-
20, Thermo Fisher Scientific, Cat # 28320) for 4 hours at room temperature. Primary
antibodies were diluted in 5% BSA and added to blots before incubating overnight at 4°C.
Blots were then washed and appropriate secondary antibody was added for 1 hour at
room temperature. Chemiluminescence was performed using the SuperSignal™ West
Dura/Femto Extended Duration Substrate (Thermo Fisher Scientific, Cat # 34075 and
34094) and the immunoblots were imaged with an Azure 600 imaging system (Azure
Biosystems; Dublin, CA USA). All antibodies and dilutions used are found in the key
resources table.
718
Quantitative real-time PCR
Total RNA was extracted from whole retina using the RNeasy Mini Kit (Qiagen, Cat #
74104) following the manufacturer's protocol. Isolated RNA was assayed for quantity and
quality with a Nanodrop 1000 (Thermo Fisher Scientific) and 1 µg of RNA was used as
input for cDNA synthesis using the RNA QuantiTect transcription kit (Qiagen, Cat #
205311). Approximately 100 ng of cDNA was used as a template for each qRT-PCR
reaction using the PowerTrack SYBR Green supermix (Thermo Fisher Scientific, Cat #
46109) as previously described (Subramanya et al., 2023). The Ct values for *Actb* were
used to determine relative transcript expression levels using the $2^{-\Delta\Delta Ct}$ method with a cycle
threshold cutoff of 35 cycles for the presence of transcript. The geometric mean was used
to normalize samples. Custom qRT-PCR primers (Supplementary File 1) designed to
specifically detect spliced transcripts were used to determine transcript levels following
the above protocol.
732
Metabolomics
For unlabeled targeted metabolomics, both retinas from a single animal were rinsed in
PBS, combined, and snap-frozen on dry ice. Metabolites were extracted using 80%
methanol at -80°C and an OMNI Bead Ruptor (OMNI International, Kennesaw, GA, USA,
Cat # 19-050A). Lysates were centrifuged at 14,000 x RCF for 10 minutes at 4°C, and the
supernatant stored at -80°C until being processed in the SpeedVac. The pellet from each
sample was saved for protein estimation. To determine the protein concentration for each
sample, 150 µL of 0.1 M NaOH was added to the pellet for 24 hours at 37°C. The sample
was then vortexed and centrifuged at 5000 x RCF at room temperature. The protein
estimation was performed as described above. The protein concentration of each sample
was used to normalize amount of sample for lyophilization with a SpeedVac concentrator
(Thermo Fisher Scientific, Cat # 13875355). Dried metabolite pellets were resuspended
for liquid chromatography-coupled mass spectrometry (LC/MS) analysis using an Agilent
Technologies Triple Quad 6470 instrument (Santa Clara, CA, USA) as previously
described (Wubben et al., 2020). Previously published parameters were used for data
collection (Yuan et al., 2012). Agilent MassHunter Workstation Quantitative Analysis
Software (B0900) was used to process raw data. Additional statistical analyses were
performed in Microsoft Excel. Each sample was normalized by the total intensity of all
metabolites to reflect sample protein content. To obtain relative metabolites, each

metabolite abundance level in each sample was divided by the mean of the abundance
levels across all control samples.
754
To analyze the incorporation of non-radioactive stable isotope carbon-13 (^{13}C) into
metabolites in central carbon metabolism and related pathways, mice were
intraperitoneally injected with 2g/kg of uniformly labeled ^{13}C$_6$-glucose (Cambridge Isotope
Laboratories, Cat # CLM-1396) or 300 mg/kg of uniformly labeled ^{13}C$_5$-glutamine
(Cambridge Isotope Laboratories, Cat # CLM-1822-H) and retinas were harvested 45
minutes later and snap-frozen as described above. Metabolites were extracted as
described above and data collected according to previously published protocols (Yuan et
al., 2019).
763
SUrface SEnsing of Translation (SUnSET) Method
In vivo protein synthesis in whole retina from P14 WT and cKO mice was measured using
the SUnSET protocol as previously described (Fort et al., 2022). Briefly, a stock solution
of 40 mg/mL puromycin hydrochloride (Millipore-Sigma, Cat # P7255) was prepared in
sterile 0.9% sodium chloride and injected intraperitoneally into mice at a final
concentration of 200 mg/kg body weight. Mice were sacrificed after 30 min and fresh
retinas were harvested and lysed in RIPA buffer (Thermo Fisher Scientific, Cat # 89900)
with protease and phosphatase inhibitors (Cell Signaling Technology, Cat # 5872). Protein
quantitation was conducted with the BCA assay as described above and 10 µg of protein
was analyzed by western blot analysis using an anti-puromycin antibody.
774
BaroFuse
Oxygen consumption rate (OCR) was determine as previously described (Kamat et al.,
2023). Briefly, a BaroFuse (Entox Sciences, Mercer Island, WA) was used to determine
the OCR of freshly isolated retinas. Single, whole retinas were dissected into Hank's
Balanced salt solution supplemented with 0.1 g/ 100 mL BSA (HBSS, Cytiva,
Marlborough, MA, USA, Cat # SH30031FS). Perifusion media consisted of commercial
Krebs-Ringer Bicarbonate buffer (KRB, Thermo Fisher Scientific, Cat # J67795.K2)
supplemented with 0.1g /100 mL fatty-acid free BSA (Millipore-Sigma, Cat # A9647) and
4.4 mM glucose, for a final concentration of 5.5 mM glucose (Millipore-Sigma, Cat #
G8270). The oxygen and CO_2 concentration of the perifusion media is maintained by
saturating the solution in a 21% oxygen, 5% CO_2 atmosphere, and the temperature was
maintained at 37°C throughout the experiment. At various times throughout the
experiment Oligomycin-A (Cayman Chemical, Ann Arbor, MI, USA, Cat # 11342), FCCP
(Trifluoromethoxy carbonylcyanide phenylhydrazone, Cayman Chemical, Cat# 15218),
and KCN (Potassium Cyanide, Thermo Fisher, Cat # 012136) were added to the perifusion
media through an injection port. A chamber without tissue was used as a negative control.
791
NAD$^+$/NADH and NADP$^+$/NADPH Measurements
The NAD$^+$/NADH and NADP$^+$/NADPH measurements were conducted using the
NAD$^+$/NADH-Glo Assay (Promega, Cat # G9071) and NADP$^+$/NADPH-Glo Assay
(Promega, Cat # G9081), respectively, following manufacturer's instructions in whole
retina from P14 WT and cKO mice. Briefly, two fresh retina per mouse were harvested for
each sample in 150 µL of PBS/Bicarbonate/0.5%dodecyltrimethylammonimum bromide
(DTAB) buffer. DTAB is used in the buffer to preserve dinucleotide stability. The samples
were sonicated (20% strength and 1sec/10 iterations) to create a uniform suspension and
diluted with 150 µL of PBS/Bicarbonate/0.5% DTAB buffer (lysate). For measuring
NAD$^+$/NADP$^+$, 150 µL of lysate was mixed with 75 µL of 0.4N HCl (Thermo Fisher
Scientific, Cat # A144) and heated to 60°C for 15 minutes. The lysate mixture was cooled

to room temperature for 10 min and then neutralized by adding 150 µL 0.5M Trizma base (Millipore-Sigma, Cat # T1503). For measuring NADH/NADPH, the remaining 150 µL of lysate was heated to 60°C for 15 minutes, cooled to room temperature for 10 min and 150 µL 0.5M Trizma base was added to the sample. Following lysate preparation, 50 µL of lysate was incubated at room temperature for 30 min with 50 µL of either the NAD+/NADP+-Glo or NADH/NAPDH-Glo detection reagent in a 96-well white walled tissue culture plate (Thermo Fisher Scientific, Cat # 3610). Luminescence was recorded using the Omega plate reader (BMG Labtech) and data are reported as a ratio of $NAD^+/NADH$ or $NADP^+/NADPH$ of N=3 animals in triplicate technical measurements per animal sample.

Statistical analysis. All data is presented as mean ± SEM. The significance of the difference between means was determined using a two-tailed student's *t*-test or one-way ANOVA in Excel or Prism 9.0. Results with a *p*-value ≤ 0.05 were considered significant.

Resource Availability. Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Thomas Wubben (twubben@med.umich.edu).

References

Adler, A. J., & Klucznik, K. M. (1982). Glycerol phosphate dehydrogenase in developing chick retina and brain. *J Neurochem*, *38*(4), 909-915. https://doi.org/10.1111/j.1471-4159.1982.tb05329.x

Adler, L. t., Chen, C., & Koutalos, Y. (2014). Mitochondria contribute to NADPH generation in mouse rod photoreceptors. *J Biol Chem*, *289*(3), 1519-1528. https://doi.org/10.1074/jbc.M113.511295

Ait-Ali, N., Fridlich, R., Millet-Puel, G., Clerin, E., Delalande, F., Jaillard, C., Blond, F., Perrocheau, L., Reichman, S., Byrne, L. C., Olivier-Bandini, A., Bellalou, J., Moyse, E., Bouillaud, F., Nicol, X., Dalkara, D., van Dorsselaer, A., Sahel, J. A., & Leveillard, T. (2015). Rod-derived cone viability factor promotes cone survival by stimulating aerobic glycolysis. *Cell*, *161*(4), 817-832. https://doi.org/10.1016/j.cell.2015.03.023

Alkan, H. F., Walter, K. E., Luengo, A., Madreiter-Sokolowski, C. T., Stryeck, S., Lau, A. N., Al-Zoughbi, W., Lewis, C. A., Thomas, C. J., Hoefler, G., Graier, W. F., Madl, T., Vander Heiden, M. G., & Bogner-Strauss, J. G. (2018). Cytosolic Aspartate Availability Determines Cell Survival When Glutamine Is Limiting. *Cell Metab*, *28*(5), 706-720 e706. https://doi.org/10.1016/j.cmet.2018.07.021

Altman, B. J., Stine, Z. E., & Dang, C. V. (2016). From Krebs to clinic: glutamine metabolism to cancer therapy. *Nat Rev Cancer*, *16*(10), 619-634. https://doi.org/10.1038/nrc.2016.71

Bowne, S. J., Liu, Q., Sullivan, L. S., Zhu, J., Spellicy, C. J., Rickman, C. B., Pierce, E. A., & Daiger, S. P. (2006). Why do mutations in the ubiquitously expressed housekeeping gene IMPDH1 cause retina-specific photoreceptor degeneration? *Invest Ophthalmol Vis Sci*, *47*(9), 3754-3765. https://doi.org/10.1167/iovs.06-0207

Busov, B., & Besirli, C. G. (2014). Optimization of ImageJ for automated image analysis to assess for photoreceptor cell death in retinal tissue sections. *Investigative Ophthalmology & Visual Science*, *55*(13), 4365-4365.

Campos-Sandoval, J. A., Martin-Rufian, M., Cardona, C., Lobo, C., Penalver, A., & Marquez, J. (2015). Glutaminases in brain: Multiple isoforms for many purposes. *Neurochem Int*, 88, 1-5. https://doi.org/10.1016/j.neuint.2015.03.006

Caruso, S. M., Ryu, J., Quinn, P. M., & Tsang, S. H. (2020). Precision metabolome reprogramming for imprecision therapeutics in retinitis pigmentosa. *J Clin Invest*, *130*(8), 3971-3973. https://doi.org/10.1172/JCI139239

Chen, Y., Zizmare, L., Calbiague, V., Wang, L., Yu, S., Herberg, F. W., Schmachtenberg, O., Paquet-Durand, F., & Trautwein, T. (2024). The retina uncouples glycolysis and oxidative phosphorylation via Cori-, Cahill-, and mini-Krebs-cycle. *Elife*, *12*, RP91141. https://doi.org/https://doi.org/10.7554/eLife.91141.1

Chinchore, Y., Begaj, T., Wu, D., Drokhlyansky, E., & Cepko, C. L. (2017). Glycolytic reliance promotes anabolism in photoreceptors. *Elife*, *6*. https://doi.org/10.7554/eLife.25946

Daemen, A., Liu, B., Song, K., Kwong, M., Gao, M., Hong, R., Nannini, M., Peterson, D., Liederer, B. M., de la Cruz, C., Sangaraju, D., Jaochico, A., Zhao, X., Sandoval, W., Hunsaker, T., Firestein, R., Latham, S., Sampath, D., Evangelista, M., & Hatzivassiliou, G. (2018). Pan-Cancer Metabolic Signature Predicts Co-Dependency on Glutaminase and De Novo Glutathione Synthesis Linked to a High-Mesenchymal Cell State. *Cell Metab*, *28*(3), 383-399 e389. https://doi.org/10.1016/j.cmet.2018.06.003

Daniele, L. L., Han, J. Y. S., Samuels, I. S., Komirisetty, R., Mehta, N., McCord, J. L., Yu, M., Wang, Y., Boesze-Battaglia, K., Bell, B. A., Du, J., Peachey, N. S., & Philp, N. J. (2022). Glucose uptake by GLUT1 in photoreceptors is essential for outer segment renewal and rod photoreceptor survival. *FASEB J*, *36*(8), e22428. https://doi.org/10.1096/fj.202200369R

Du, J., Cleghorn, W., Contreras, L., Linton, J. D., Chan, G. C., Chertov, A. O., Saheki, T., Govindaraju, V., Sadilek, M., Satrustegui, J., & Hurley, J. B. (2013). Cytosolic reducing power preserves glutamate in retina. *Proc Natl Acad Sci U S A*, *110*(46), 18501-18506. https://doi.org/10.1073/pnas.1311193110

Du, J., Cleghorn, W. M., Contreras, L., Lindsay, K., Rountree, A. M., Chertov, A. O., Turner, S. J., Sahaboglu, A., Linton, J., Sadilek, M., Satrustegui, J., Sweet, I. R., Paquet-Durand, F., & Hurley, J. B. (2013). Inhibition of mitochondrial pyruvate transport by zaprinast causes massive accumulation of aspartate at the expense of glutamate in the retina. *J Biol Chem*, *288*(50), 36129-36140. https://doi.org/10.1074/jbc.M113.507285

Du, J., Linton, J. D., & Hurley, J. B. (2015). Probing Metabolism in the Intact Retina Using Stable Isotope Tracers. *Methods Enzymol*, *561*, 149-170. https://doi.org/10.1016/bs.mie.2015.04.002

Duncan, J. L., Pierce, E. A., Laster, A. M., Daiger, S. P., Birch, D. G., Ash, J. D., Iannaccone, A., Flannery, J. G., Sahel, J. A., Zack, D. J., Zarbin, M. A., & and the Foundation Fighting Blindness Scientific Advisory, B. (2018). Inherited Retinal Degenerations: Current Landscape and Knowledge Gaps. *Transl Vis Sci Technol*, *7*(4), 6. https://doi.org/10.1167/tvst.7.4.6

Fort, P. E., Losiewicz, M. K., Elghazi, L., Kong, D., Cras-Meneur, C., Fingar, D. C., Kimball, S. R., Rajala, R. V. S., Smith, A. J., Ali, R. R., Abcouwer, S. F., & Gardner, T. W. (2022). mTORC1 regulates high levels of protein synthesis in retinal ganglion cells of adult mice. *J Biol Chem*, *298*(6), 101944. https://doi.org/10.1016/j.jbc.2022.101944

Gorbatyuk, M. S., Starr, C. R., & Gorbatyuk, O. S. (2020). Endoplasmic reticulum stress: New insights into the pathogenesis and treatment of retinal degenerative

diseases. *Prog Retin Eye Res*, *79*, 100860.
https://doi.org/10.1016/j.preteyeres.2020.100860
Grenell, A., Wang, Y., Yam, M., Swarup, A., Dilan, T. L., Hauer, A., Linton, J. D., Philp,
N. J., Gregor, E., Zhu, S., Shi, Q., Murphy, J., Guan, T., Lohner, D.,
Kolandaivelu, S., Ramamurthy, V., Goldberg, A. F. X., Hurley, J. B., & Du, J.
(2019). Loss of MPC1 reprograms retinal metabolism to impair visual function.
Proc Natl Acad Sci U S A, *116*(9), 3530-3535.
https://doi.org/10.1073/pnas.1812941116
Halbrook, C. J., Thurston, G., Boyer, S., Anaraki, C., Jimenez, J. A., McCarthy, A.,
Steele, N. G., Kerk, S. A., Hong, H. S., Lin, L., Law, F. V., Felton, C., Scipioni, L.,
Sajjakulnukit, P., Andren, A., Beutel, A. K., Singh, R., Nelson, B. S., Van Den
Bergh, F., Krall, A. S., Mullen, P. J., Zhang, L., Batra, S., Morton, J. P., Stanger,
B. Z., Christofk, H. R., Digman, M. A., Beard, D. A., Viale, A., Zhang, J.,
Crawford, H. C., Pasca di Magliano, M., Jorgensen, C., & Lyssiotis, C. A. (2022).
Differential integrated stress response and asparagine production drive
symbiosis and therapy resistance of pancreatic adenocarcinoma cells. *Nat
Cancer*, *3*(11), 1386-1403. https://doi.org/10.1038/s43018-022-00463-1
Halliday, M., Radford, H., Sekine, Y., Moreno, J., Verity, N., le Quesne, J., Ortori, C. A.,
Barrett, D. A., Fromont, C., Fischer, P. M., Harding, H. P., Ron, D., & Mallucci, G.
R. (2015). Partial restoration of protein synthesis rates by the small molecule
ISRIB prevents neurodegeneration without pancreatic toxicity. *Cell Death Dis*,
6(3), e1672. https://doi.org/10.1038/cddis.2015.49
Hartong, D. T., Dange, M., McGee, T. L., Berson, E. L., Dryja, T. P., & Colman, R. F.
(2008). Insights from retinitis pigmentosa into the roles of isocitrate
dehydrogenases in the Krebs cycle. *Nat Genet*, *40*(10), 1230-1234.
https://doi.org/10.1038/ng.223
Joyal, J. S., Sun, Y., Gantner, M. L., Shao, Z., Evans, L. P., Saba, N., Fredrick, T.,
Burnim, S., Kim, J. S., Patel, G., Juan, A. M., Hurst, C. G., Hatton, C. J., Cui, Z.,
Pierce, K. A., Bherer, P., Aguilar, E., Powner, M. B., Vevis, K., Boisvert, M., Fu,
Z., Levy, E., Fruttiger, M., Packard, A., Rezende, F. A., Maranda, B., Sapieha, P.,
Chen, J., Friedlander, M., Clish, C. B., & Smith, L. E. (2016). Retinal lipid and
glucose metabolism dictates angiogenesis through the lipid sensor Ffar1. *Nat
Med*, *22*(4), 439-445. https://doi.org/10.1038/nm.4059
Kamat, V., Grumbine, M. K., Bao, K., Mokate, K., Khalil, G., Cook, D., Clearwater, B.,
Hirst, R., Harman, J., Boeck, M., Fu, Z., Smith, L. E. H., Goswami, M., Wubben,
T. J., Walker, E. M., Zhu, J., Soleimanpour, S. A., Scarlett, J. M., Robbings, B.
M., Hass, D., Hurley, J. B., & Sweet, I. R. (2023). A versatile pumpless multi-
channel fluidics system for maintenance and real-time functional assessment of
tissue and cells. *Cell Rep Methods*, *3*(11), 100642.
https://doi.org/10.1016/j.crmeth.2023.100642
Kanow, M. A., Giarmarco, M. M., Jankowski, C. S., Tsantilas, K., Engel, A. L., Du, J.,
Linton, J. D., Farnsworth, C. C., Sloat, S. R., Rountree, A., Sweet, I. R., Lindsay,
K. J., Parker, E. D., Brockerhoff, S. E., Sadilek, M., Chao, J. R., & Hurley, J. B.
(2017). Biochemical adaptations of the retina and retinal pigment epithelium
support a metabolic ecosystem in the vertebrate eye. *Elife*, *6*.
https://doi.org/10.7554/eLife.28899
Koch, S. F., Tsai, Y. T., Duong, J. K., Wu, W. H., Hsu, C. W., Wu, W. P., Bonet-Ponce,
L., Lin, C. S., & Tsang, S. H. (2015). Halting progressive neurodegeneration in
advanced retinitis pigmentosa. *J Clin Invest*, *125*(9), 3704-3713.
https://doi.org/10.1172/JCI82462

Kooragayala, K., Gotoh, N., Cogliati, T., Nellissery, J., Kaden, T. R., French, S.,
Balaban, R., Li, W., Covian, R., & Swaroop, A. (2015). Quantification of Oxygen
Consumption in Retina Ex Vivo Demonstrates Limited Reserve Capacity of
Photoreceptor Mitochondria. *Invest Ophthalmol Vis Sci*, *56*(13), 8428-8436.
https://doi.org/10.1167/iovs.15-17901
Kutsyr, O., Noailles, A., Martinez-Gil, N., Maestre-Carballa, L., Martinez-Garcia, M.,
Maneu, V., Cuenca, N., & Lax, P. (2021). Short-term high-fat feeding
exacerbates degeneration in retinitis pigmentosa by promoting retinal oxidative
stress and inflammation. *Proc Natl Acad Sci U S A*, *118*(43).
https://doi.org/10.1073/pnas.2100566118
Le, Y. Z., Zheng, L., Zheng, W., Ash, J. D., Agbaga, M. P., Zhu, M., & Anderson, R. E.
(2006). Mouse opsin promoter-directed Cre recombinase expression in
transgenic mice. *Mol Vis*, *12*, 389-398.
https://www.ncbi.nlm.nih.gov/pubmed/16636658
Li, B., Zhang, T., Liu, W., Wang, Y., Xu, R., Zeng, S., Zhang, R., Zhu, S., Gillies, M. C.,
Zhu, L., & Du, J. (2020). Metabolic Features of Mouse and Human Retinas: Rods
versus Cones, Macula versus Periphery, Retina versus RPE. *iScience*, *23*(11),
101672. https://doi.org/10.1016/j.isci.2020.101672
Li, J., Csibi, A., Yang, S., Hoffman, G. R., Li, C., Zhang, E., Yu, J. J., & Blenis, J. (2015).
Synthetic lethality of combined glutaminase and Hsp90 inhibition in mTORC1-
driven tumor cells. *Proc Natl Acad Sci U S A*, *112*(1), E21-29.
https://doi.org/10.1073/pnas.1417015112
Li, S., Chen, D., Sauve, Y., McCandless, J., Chen, Y. J., & Chen, C. K. (2005).
Rhodopsin-iCre transgenic mouse line for Cre-mediated rod-specific gene
targeting. *Genesis*, *41*(2), 73-80. https://doi.org/10.1002/gene.20097
Macosko, E. Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M., Tirosh, I.,
Bialas, A. R., Kamitaki, N., Martersteck, E. M., Trombetta, J. J., Weitz, D. A.,
Sanes, J. R., Shalek, A. K., Regev, A., & McCarroll, S. A. (2015). Highly Parallel
Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets.
Cell, *161*(5), 1202-1214. https://doi.org/10.1016/j.cell.2015.05.002
Mattapallil, M. J., Wawrousek, E. F., Chan, C. C., Zhao, H., Roychoudhury, J., Ferguson,
T. A., & Caspi, R. R. (2012). The Rd8 mutation of the Crb1 gene is present in
vendor lines of C57BL/6N mice and embryonic stem cells, and confounds ocular
induced mutant phenotypes. *Invest Ophthalmol Vis Sci*, *53*(6), 2921-2927.
https://doi.org/10.1167/iovs.12-9662
McLaughlin, B. J., Wood, J. G., & Gurd, J. W. (1980). The localization of lectin binding
sites during photoreceptor synaptogenesis in the chick retina. *Brain Res*, *191*(2),
345-357. https://doi.org/10.1016/0006-8993(80)91286-x
Mingote, S., Masson, J., Gellman, C., Thomsen, G. M., Lin, C. S., Merker, R. J., Gaisler-
Salomon, I., Wang, Y., Ernst, R., Hen, R., & Rayport, S. (2015). Genetic
Pharmacotherapy as an Early CNS Drug Development Strategy: Testing
Glutaminase Inhibition for Schizophrenia Treatment in Adult Mice. *Front Syst*
Neurosci, *9*, 165. https://doi.org/10.3389/fnsys.2015.00165
Ng, S. K., Wood, J. P., Chidlow, G., Han, G., Kittipassorn, T., Peet, D. J., & Casson, R.
J. (2015). Cancer-like metabolism of the mammalian retina. *Clin Exp Ophthalmol*,
43(4), 367-376. https://doi.org/10.1111/ceo.12462
Okazaki, A., Gameiro, P. A., Christodoulou, D., Laviollette, L., Schneider, M., Chaves,
F., Stemmer-Rachamimov, A., Yazinski, S. A., Lee, R., Stephanopoulos, G., Zou,
L., & Iliopoulos, O. (2017). Glutaminase and poly(ADP-ribose) polymerase
inhibitors suppress pyrimidine synthesis and VHL-deficient renal cancers. *J Clin*
Invest, *127*(5), 1631-1645. https://doi.org/10.1172/JCI87800

Pakos-Zebrucka, K., Koryga, I., Mnich, K., Ljujic, M., Samali, A., & Gorman, A. M. (2016). The integrated stress response. *EMBO Rep, 17*(10), 1374-1395. https://doi.org/10.15252/embr.201642195

Pan, W. W., Wubben, T. J., & Besirli, C. G. (2021). Photoreceptor metabolic reprogramming: current understanding and therapeutic implications. *Commun Biol, 4*(1), 245. https://doi.org/10.1038/s42003-021-01765-3

Pavlova, N. N., Hui, S., Ghergurovich, J. M., Fan, J., Intlekofer, A. M., White, R. M., Rabinowitz, J. D., Thompson, C. B., & Zhang, J. (2018). As Extracellular Glutamine Levels Decline, Asparagine Becomes an Essential Amino Acid. *Cell Metab, 27*(2), 428-438 e425. https://doi.org/10.1016/j.cmet.2017.12.006

Petit, L., Ma, S., Cipi, J., Cheng, S. Y., Zieger, M., Hay, N., & Punzo, C. (2018). Aerobic Glycolysis Is Essential for Normal Rod Function and Controls Secondary Cone Death in Retinitis Pigmentosa. *Cell Rep, 23*(9), 2629-2642. https://doi.org/10.1016/j.celrep.2018.04.111

Rajala, A., Bhat, M. A., Teel, K., Gopinadhan Nair, G. K., Purcell, L., & Rajala, R. V. S. (2023). The function of lactate dehydrogenase A in retinal neurons: implications to retinal degenerative diseases. *PNAS Nexus, 2*(3), pgad038. https://doi.org/10.1093/pnasnexus/pgad038

Rajala, R. V. S. (2020). Aerobic Glycolysis in the Retina: Functional Roles of Pyruvate Kinase Isoforms. *Front Cell Dev Biol, 8*, 266. https://doi.org/10.3389/fcell.2020.00266

Rosarda, J. D., Giles, S., Harkins-Perry, S., Mills, E. A., Friedlander, M., Wiseman, R. L., & Eade, K. T. (2023). Imbalanced unfolded protein response signaling contributes to 1-deoxysphingolipid retinal toxicity. *Nat Commun, 14*(1), 4119. https://doi.org/10.1038/s41467-023-39775-w

Ross, C. D., Bowers, M., & Godfrey, D. A. (1987). Distribution of glutaminase activity in retinal layers of rat and guinea pig. *Brain Res, 401*(1), 168-172. https://doi.org/10.1016/0006-8993(87)91178-4

Rowe, A. A., Patel, P. D., Gordillo, R., & Wert, K. J. (2021). Replenishment of TCA cycle intermediates provides photoreceptor resilience against neurodegeneration during progression of retinitis pigmentosa. *JCI Insight, 6*(17). https://doi.org/10.1172/jci.insight.150898

Sisk, D. R., & Kuwabara, T. (1985). Histologic changes in the inner retina of albino rats following intravitreal injection of monosodium L-glutamate. *Graefes Arch Clin Exp Ophthalmol, 223*(5), 250-258. https://doi.org/10.1007/BF02153655

Son, J., Lyssiotis, C. A., Ying, H., Wang, X., Hua, S., Ligorio, M., Perera, R. M., Ferrone, C. R., Mullarky, E., Shyh-Chang, N., Kang, Y., Fleming, J. B., Bardeesy, N., Asara, J. M., Haigis, M. C., DePinho, R. A., Cantley, L. C., & Kimmelman, A. C. (2013). Glutamine supports pancreatic cancer growth through a KRAS-regulated metabolic pathway. *Nature, 496*(7443), 101-105. https://doi.org/10.1038/nature12040

Subramanya, S., Goswami, M. T., Miller, N., Weh, E., Chaudhury, S., Zhang, L., Andren, A., Hager, H., Weh, K. M., Lyssiotis, C. A., Besirli, C. G., & Wubben, T. J. (2023). Rod photoreceptor-specific deletion of cytosolic aspartate aminotransferase, GOT1, causes retinal degeneration [Original Research]. *Frontiers in Ophthalmology, 3*. https://doi.org/10.3389/fopht.2023.1306019

Sullivan, L. B., Luengo, A., Danai, L. V., Bush, L. N., Diehl, F. F., Hosios, A. M., Lau, A. N., Elmiligy, S., Malstrom, S., Lewis, C. A., & Vander Heiden, M. G. (2018). Aspartate is an endogenous metabolic limitation for tumour growth. *Nat Cell Biol, 20*(7), 782-788. https://doi.org/10.1038/s41556-018-0125-0

Swarup, A., Samuels, I. S., Bell, B. A., Han, J. Y. S., Du, J., Massenzio, E., Abel, E. D., Boesze-Battaglia, K., Peachey, N. S., & Philp, N. J. (2019). Modulating GLUT1 expression in retinal pigment epithelium decreases glucose levels in the retina: impact on photoreceptors and Muller glial cells. *Am J Physiol Cell Physiol, 316*(1), C121-C133. https://doi.org/10.1152/ajpcell.00410.2018

Tsantilas, K. A., Cleghorn, W. M., Bisbach, C. M., Whitson, J. A., Hass, D. T., Robbings, B. M., Sadilek, M., Linton, J. D., Rountree, A. M., Valencia, A. P., Sweetwyne, M. T., Campbell, M. D., Zhang, H., Jankowski, C. S. R., Sweet, I. R., Marcinek, D. J., Rabinovitch, P. S., & Hurley, J. B. (2021). An Analysis of Metabolic Changes in the Retina and Retinal Pigment Epithelium of Aging Mice. *Invest Ophthalmol Vis Sci, 62*(14), 20. https://doi.org/10.1167/iovs.62.14.20

Ueta, T., Inoue, T., Furukawa, T., Tamaki, Y., Nakagawa, Y., Imai, H., & Yanagi, Y. (2012). Glutathione peroxidase 4 is required for maturation of photoreceptor cells. *J Biol Chem, 287*(10), 7675-7682. https://doi.org/10.1074/jbc.M111.335174

Voigt, A. P., Whitmore, S. S., Lessing, N. D., DeLuca, A. P., Tucker, B. A., Stone, E. M., Mullins, R. F., & Scheetz, T. E. (2020). Spectacle: An interactive resource for ocular single-cell RNA sequencing data analysis. *Exp Eye Res, 200*, 108204. https://doi.org/10.1016/j.exer.2020.108204

Weh, E., Lutrzykowska, Z., Smith, A., Hager, H., Pawar, M., Wubben, T. J., & Besirli, C. G. (2020). Hexokinase 2 is dispensable for photoreceptor development but is required for survival during aging and outer retinal stress. *Cell Death Dis, 11*(6), 422. https://doi.org/10.1038/s41419-020-2638-2

Weh, E., Scott, K., Wubben, T. J., & Besirli, C. G. (2022). Dark-reared rd10 mice experience rapid photoreceptor degeneration with short exposure to room-light during in vivo retinal imaging. *Exp Eye Res, 215*, 108913. https://doi.org/10.1016/j.exer.2021.108913

Wert, K. J., Velez, G., Kanchustambham, V. L., Shankar, V., Evans, L. P., Sengillo, J. D., Zare, R. N., Bassuk, A. G., Tsang, S. H., & Mahajan, V. B. (2020). Metabolite therapy guided by liquid biopsy proteomics delays retinal neurodegeneration. *EBioMedicine, 52*, 102636. https://doi.org/10.1016/j.ebiom.2020.102636

Winkler, B. S. (1981). Glycolytic and oxidative metabolism in relation to retinal function. *J Gen Physiol, 77*(6), 667-692. https://doi.org/10.1085/jgp.77.6.667

Winkler, B. S., DeSantis, N., & Solomon, F. (1986). Multiple NADPH-producing pathways control glutathione (GSH) content in retina. *Exp Eye Res, 43*(5), 829-847. https://doi.org/10.1016/s0014-4835(86)80013-6

Wubben, T. J., Besirli, C. G., & Zacks, D. N. (2016). Pharmacotherapies for Retinal Detachment. *Ophthalmology, 123*(7), 1553-1562. https://doi.org/10.1016/j.ophtha.2016.02.040

Wubben, T. J., Pawar, M., Smith, A., Toolan, K., Hager, H., & Besirli, C. G. (2017). Photoreceptor metabolic reprogramming provides survival advantage in acute stress while causing chronic degeneration. *Sci Rep, 7*(1), 17863. https://doi.org/10.1038/s41598-017-18098-z

Wubben, T. J., Pawar, M., Weh, E., Smith, A., Sajjakulnukit, P., Zhang, L., Dai, L., Hager, H., Pai, M. P., Lyssiotis, C. A., & Besirli, C. G. (2020). Small molecule activation of metabolic enzyme pyruvate kinase muscle isozyme 2, PKM2, circumvents photoreceptor apoptosis. *Sci Rep, 10*(1), 2990. https://doi.org/10.1038/s41598-020-59999-w

Xu, R., Ritz, B. K., Wang, Y., Huang, J., Zhao, C., Gong, K., Liu, X., & Du, J. (2020). The retina and retinal pigment epithelium differ in nitrogen metabolism and are metabolically connected. *J Biol Chem, 295*(8), 2324-2335. https://doi.org/10.1074/jbc.RA119.011727

Xu, Y., Shi, T., Cui, X., Yan, L., Wang, Q., Xu, X., Zhao, Q., Xu, X., Tang, Q. Q., Tang,
H., & Pan, D. (2021). Asparagine reinforces mTORC1 signaling to boost
thermogenesis and glycolysis in adipose tissues. *EMBO J*, *40*(24), e108069.
https://doi.org/10.15252/embj.2021108069
Yang, L., Venneti, S., & Nagrath, D. (2017). Glutaminolysis: A Hallmark of Cancer
Metabolism. *Annu Rev Biomed Eng*, *19*, 163-194.
https://doi.org/10.1146/annurev-bioeng-071516-044546
Yoo, H. C., Yu, Y. C., Sung, Y., & Han, J. M. (2020). Glutamine reliance in cell
metabolism. *Exp Mol Med*, *52*(9), 1496-1516. https://doi.org/10.1038/s12276-
020-00504-8
Young, R. W. (1967). The renewal of photoreceptor cell outer segments. *J Cell Biol*,
33(1), 61-72. https://doi.org/10.1083/jcb.33.1.61
Yuan, M., Breitkopf, S. B., Yang, X., & Asara, J. M. (2012). A positive/negative ion-
switching, targeted mass spectrometry-based metabolomics platform for bodily
fluids, cells, and fresh and fixed tissue. *Nat Protoc*, *7*(5), 872-881.
https://doi.org/10.1038/nprot.2012.024
Yuan, M., Kremer, D. M., Huang, H., Breitkopf, S. B., Ben-Sahra, I., Manning, B. D.,
Lyssiotis, C. A., & Asara, J. M. (2019). Ex vivo and in vivo stable isotope labelling
of central carbon metabolism and related pathways with analysis by LC-MS/MS.
Nat Protoc, *14*(2), 313-330. https://doi.org/10.1038/s41596-018-0102-x
Zhang, J., Fan, J., Venneti, S., Cross, J. R., Takagi, T., Bhinder, B., Djaballah, H., Kanai,
M., Cheng, E. H., Judkins, A. R., Pawel, B., Baggs, J., Cherry, S., Rabinowitz, J.
D., & Thompson, C. B. (2014). Asparagine plays a critical role in regulating
cellular adaptation to glutamine depletion. *Mol Cell*, *56*(2), 205-218.
https://doi.org/10.1016/j.molcel.2014.08.018
Zhou, M., Liu, Y., & Ma, C. (2021). Distinct Nuclear Architecture of Photoreceptors and
Light-Induced Behaviors in Different Strains of Mice. *Transl Vis Sci Technol*,
10(2), 37. https://doi.org/10.1167/tvst.10.2.37
1134
1135

Figures and Figure Legends

1137

1138
1139
Figure 1. Rod photoreceptor-specific knockout of glutaminase (GLS) displays rapid
retinal degeneration and increased markers of cell death. (A) Western blot analysis
showing decreased GLS protein levels in the retina of WT and cKO mice at post-natal day
14 (P14). Quantitation of Western blot results for N=3-4 animals per group. (B)
Representative images for immunofluorescence of P14 mouse retinas (N=3 animals per
group) stained for GLS (red), cone opsin (green) and nuclei (DAPI, blue) in WT and cKO
mice. White arrows indicate remaining GLS expression in cone photoreceptors. Left and
middle panel scale bars are 40 μm. Right image scale bar is 20 μm. (C) OCT images

detailing outer retinal changes in cKO mice over time compared to WT. Retinal structures
are comparable to WT mice at P14 but rapidly thin with age. (D) Total retinal thickness,
(E) ONL thickness and (F) IS/OS thickness as determined by OCT in WT and cKO mice
over time. N=4-9 eyes per group. (G) Representative hematoxylin and eosin stained retinal
sections from rod photoreceptor-specific *Gls* conditional knockout (cKO) mice compared
to wild-type (WT) mice at P14, P21 and P42. ONL, outer nuclear layer; INL, inner nuclear
layer; GCL, ganglion cell layer. Scale bar is 40 μm. H) ONL cell counts as a percent of WT
retinas at P14, P21 and P42. N=5-10 eyes per group. (I) Representative images of WT
and cKO retinas stained to detect TUNEL-positive cells (green) at P21. Scale bar is 40
μm. N=3-5 animals per group. (J) Quantitation of percent TUNEL-positive cells at P21
showing an increase in TUNEL-positive cells in cKO animals. N=3-5 animals per group.
(K) qRT-PCR of genes related to cell death pathways including apoptosis (Apop.),
necroptosis (Nec.), autophagy (Aut.) and ferroptosis (Ferrop) in WT and cKO mice at P14.
N=6 animals per group. Statistical differences in (A), (D), (E), (F), (H), (J) and (K) are
based on an unpaired two-tailed Student's t-test where *P <0.05, **P<0.01 and
***P<0.001. Data are presented as mean ± standard error of the mean. OCT: optical
coherence tomography, OS: outer segment, IS: inner segment, ONL: outer nuclear layer,
INL: inner nuclear layer, GCL: ganglion cell layer, TUNEL: terminal deoxynucleotidyl
transferase dUTP nick and labeling.

1167

1168

Figure 1 – figure supplement 1. GLS is the predominant isoform in the mouse retina and the rod photoreceptor-specific *Gls* conditional knockout mouse does not demonstrate compensatory upregulation of the *Gls2* isoform. (A) qRT-PCR analysis of *Gls* and *Gls2* in P14 mouse retina. Relative expression was analyzed by comparative threshold cycle (2^-ΔΔCT) method. Expression values were represented as fold change over *Gls* after normalization with β-actin. N=9 animals per group. (B) Representative GLS immunofluorescence in P21 WT animals. Scale bar is 40 μm. (C) Western blot analysis and quantitation of GLS in fractionated wild-type (WT) mouse retinas. TIM23 was used as a mitochondrial fraction marker and HSP90 as a cytosolic fraction marker. WR whole retina extract; C, cytosolic fraction; M, mitochondrial fraction. N=3 animals. (D) qRT-PCR analysis of *Gls2* mRNA expression in the retina at P14 demonstrates no change between the rod photoreceptor-specific *Gls* conditional knockout mouse (cKO) and the wild-type (WT) mouse where both *Gls* alleles are present. Data are normalized to β-actin. N=5-6 animals per group. (E) Representative GLS2 immunofluorescence (red) images in cKO mice at P21 shows no increase in the expression of GLS2 as compared WT mice. Scale bar is 40 μm. Statistical differences in (A), (B) and (D) are based on an unpaired two-tailed Student's t-test where **P<0.01 and ***P<0.001. Data are presented as mean ± standard error of the mean.

1187

Figure 1 – figure supplement 2. *Gls* **conditional knockout mouse demonstrates**
similar degeneration under cyclic- and dark-reared conditions. (A) Representative
OCT images from the rod photoreceptor-specific *Gls* conditional knockout (cKO) mouse
compared to wild-type (WT) mice at P21 and P28 housed with 12-hour light/12-hour dark
cycles (cyclic-reared) or dark-reared. The white vertical bar represents the thickness of
the outer nuclear layer (ONL). (B) Quantitation of ONL thickness as determined by OCT
in cKO mice compared to WT mice at P21 and P28 under the different housing conditions.
N=4-6 animals per group. Statistical differences in (B) are based on a One-way ANOVA
with Tukey's post hoc test for multiple comparisons where *** P<0.001. Data are presented
as mean ± standard error of the mean.

1198

1199

Figure 1 – figure supplement 3. Histology confirms increased Müller glial cell
activation in rod photoreceptor-specific *Gls* **conditional knockout mice.**
Representative images from GFAP staining indicate increased Müller glial cell activation
in cKO compared to WT mice at P21 and P42. Scale bar is 40 µm.

1204

1205
Figure 1 – figure supplement 4. Rod photoreceptor-specific *Gls* conditional
knockout mice have shorter outer segments. (A) Representative immunofluorescence
images of rhodopsin (RHO) staining (green) in rod photoreceptor-specific *Gls* conditional
knockout (cKO) mice compared to wild-type (WT) mice at P21. N=3 animals per group.
Scale bar is 40 µm. (B) Representative transmission electron microscopy (TEM) images
of retinal sections from cKO mice compared to WT mice at P21. Images taken at 250X
magnification and scale bar is 5 µm. (C) TEM images of retinal sections from cKO mice
compared to WT mice at P21 taken at 500X magnification and scale bar is 3 µm. (D) TEM
images of retinal sections from cKO and WT mice at P21 taken at 8000X and 6000X
magnification, respectively, to show ribbon synapses. Intact ribbon synapses in WT rods
are shown in yellow boxes. A structurally intact ribbon synapse in cKO rods is denoted by
the yellow arrow. Scale bars are 500 nm.

1218

1219
Figure 1 – figure supplement 5. Rod photoreceptor-specific *Gls* knockout does not
alter synaptic connectivity between photoreceptors and second-order neurons. (A)
Representative retinal sections stained with wheat germ agglutinin (WGA) to label
photoreceptor synaptic membranes and non-synaptic membranes in the rod
photoreceptor-specific *Gls* conditional knockout (cKO) mice compared to wild-type (WT)
mice at P14. OPL, outer plexiform layer; IPL, inner plexiform layer. N=3 animals per group.
Scale bar is 40 µm. (B) Representative retinal sections stained with an antibody against
Bassoon to label ribbon synapses of rods and cones in the OPL in cKO and WT mice at

P14. N=3 animals per group. Scale bar is 40 µm. (C) Representative retinal sections from
WT and cKO animals at P14, P21 and P42 are immunostained with antibodies against
major cell-type specific markers (red): calretinin (amacrine cells), BRN3A (ganglion cells),
CHX10 (bipolar cells). ONL, outer nuclear layer; INL, inner nuclear layer; GCL, ganglion
cell layer. N=3 animals per group. Scale bars are 40 µm. (D) Inner retinal area as a percent
of WT retinas at P14, P21 and P42. N=5-10 eyes per group. Statistical differences in (D)
are based on an unpaired two-tailed Student's t-test where **$P<0.01$. Data are presented
as mean ± standard error of the mean.

1236

Figure 2. Loss of *Gls* in rod photoreceptors impairs retinal function. (A)
Representative scotopic and photopic electroretinography (ERG) tracings for the rod
photoreceptor-specific *Gls* conditional knockout mice (cKO) compared wild-type (WT)
mice at P21 and P42. (B) ERG scotopic a-wave and (C) b-wave amplitudes in cKO mice
compared to WT mice at P21 and P42. A flash intensity of 32 cd*s/m^2 was utilized. N=3-
6 animals per group. (D) ERG photopic b-wave amplitudes in cKO mice compared to WT
mice at P21 and P42. A flash intensity of 100 cd*s/m^2 was used. N=3-6 animals per
group. (E) Representative images from staining of the cone-specific marker peanut
agglutinin (PNA, green) and nuclei (DAPI, blue) in retinal sections from cKO mice
compared to WT mice at P14, P21 and P42. Scale bars are 40 μm. Statistical
differences in (B), (C) and (D) are based on an unpaired two-tailed Student's t-test
where *P <0.05, **P<0.01 and ***P<0.001. Data are presented as mean ± standard error
of the mean. ONL, outer nuclear layer; IS, inner segments; OS, outer segments. N=3 per
group.

1253
Figure 3. Rod photoreceptors require GLS for maintenance and maturation. *Gls^fl/fl*
mice carrying a tamoxifen (TAM)-inducible Cre-recombinase (IND-cKO) under the
control of *Pde6g* (*Gls^fl/fl;Pde6g-Cre^ERT2*) compared to mice expressing only the inducible
Cre-recombinase (*Gls^wt/wt;Pde6g-Cre^ERT2*; WT). Both IND-cKO and WT were
administered tamoxifen for 5 days starting at P22. (A) Quantitation of Western blot
results showing decreased GLS protein levels in the retina of IND-cKO animals at 10
days after tamoxifen induction compared to WT mice. HSP90 was used as a loading
control. N=3-4 animals per group. (B) Representative GLS immunofluorescence (red) in
IND-cKO mice compared to the wild-type (WT) mouse 10 days after tamoxifen induction.
N=3 animals per group. Scale bar is 40 µm. (C) OCT images detailing outer retinal
changes in WT and IND-cKO animals over time. Total retinal thickness (D), outer

nuclear layer (ONL) thickness (E) and inner segment/outer segment (IS/OS) thickness
(F) as determined by OCT for 38 days post tamoxifen. N= 5-10 eyes per group. (G)
Representative hematoxylin and eosin-stained retinal sections from IND-cKO mice
compared to WT mice at 10 and 38 days after tamoxifen induction. N=3 animals per
group. Scale bar is 40 μm. ONL, outer nuclear layer; INL, inner nuclear layer; GCL,
ganglion cell layer. Quantitation of scotopic (H) and photopic (I) ERG a- and b-waves in
WT and IND-cKO retina 10 days post TAM. N>12 eyes per genotype. Statistical
differences in (A), (D), (E), (F), (H), and (I) are based on an unpaired two-tailed
Student's t-test where *P<0.05, **P<0.01 and ***P<0.001. Data are presented as mean ±
standard error of the mean.

**Figure 4. GLS cKO mice maintain levels of nucleotide, glycolytic and TCA cycle
metabolites and mitochondrial function but demonstrate altered redox balance.** (A)
Schematic summarizing the biosynthetic and bioenergetic roles of glutamine. (B) Relative
abundance of key intermediates in nucleotide metabolism in the retina of WT and cKO
mice at P14 as determined by targeted metabolomics. N=6-7 animals per group. Relative
abundance is the ion intensity normalized to the WT. (C) The NADP+/NADPH ratio, as
determined by bioluminescence assay, is significantly increased in the cKO as compared
to WT retina at P14. N=5-6 animals per group. (D) Relative abundance of GSSG in the
retina of WT and cKO mice at P14, prior to PR degeneration, as determined by targeted
metabolomics. N=6-7 animals per group. (E) qRT-PCR of genes related to redox
homeostasis are significantly altered in cKO compared to WT mice. N=6 animals per
group. (F) Relative abundance of metabolites in glycolysis and the TCA cycle in WT and
cKO retina at P14. N=6-7 animals per group. (G) Mitochondrial stress test carried out on
isolated WT and cKO retina at P14 using the BaroFuse. The baseline was established by
perifusing the tissue for 90 min and then oligomycin, FCCP, and KCN were injected into

the perifusate sequentially as indicated. (H) Comparison of the effects of oligomycin and FCCP on OCR in P14 WT and cKO retina. N=6-8 animals per group. (I) Western blot analysis and quantitation of the mitochondrial electron transport chain complexes show no differences between WT and cKO retina. N=3-4 animals per group. Fold change is in relation to WT. Statistical differences in (B-F), (H) and (I) are based on an unpaired two-tailed Student's t-test where *P<0.05. Data are presented as mean ± standard error of the mean. R5P: ribose 5-phosphate, IMP: inosine monophosphate, UMP: uridine monophosphate, GSSG: glutathione disulfide, F6P: fructose 6-phosphate, DHAP: dihydroxyacetone phosphate, 2PG: 2-phospho-D-glycerate, PEP: phosphoenolpyruvate, Pyr: pyruvate, Lac: lactate, Cit: citrate, α-KG: alpha-ketoglutarate, Suc: succinate, Mal: malate, Oligo: oligomycin, FCCP: carbonyl cyanide p-trifluoromethoxyphenylhydrazone, KCN: potassium cyanide, CI-NDUFB8: complex 1, NADH:ubiquinone oxidoreductase subunit B8, CII-SDHB: complex 2, succinate dehydrogenase complex iron sulfur subunit B, CIII-UQCRC2: complex 3, ubiquinol-cytochrome c reductase core protein 2, CV-ATP5A: complex 5, ATP synthase F1 subunit alpha, HSP90: heat shock protein 90.

Figure 4 – figure supplement 1. Loss of *Gls* in rod photoreceptors alters the expression of retinal metabolism-related genes. qRT-PCR analysis of genes related to (A) glycolysis and pyruvate metabolism as well as (B) the TCA cycle in rod photoreceptor-specific *Gls* conditional knockout mice (cKO) compared to wild-type (WT) mice retina at P14. Expression values are represented as fold change over WT after normalization with β-actin. N=6 animals per group. Statistical differences in (A) and (B) are based on an unpaired two-tailed Student's t-test where *P<0.05, **P<0.01 and ***P<0.001. Data are presented as mean ± standard error of the mean.

Figure 4 – figure supplement 2. Loss of *Gls* in rod photoreceptors does not alter the expression of metabolism-related genes in eyecups. qRT-PCR analysis of genes related to (A) glycolysis and pyruvate metabolism as well as (B) the TCA cycle in rod photoreceptor-specific *Gls* conditional knockout mice (cKO) eyecups compared to wild-type (WT) mice eyecups at P14. Expression values are represented as fold change over WT after normalization with β-actin. N=3-4 eyecups per group. Statistical differences in (A) and (B) are based on an unpaired two-tailed Student's t-test where **$P<0.01$. Data are presented as mean ± standard error of the mean.

1329

Figure 5. Loss of GLS in rod photoreceptors has significant effects on TCA cycle metabolism with only partial rescue upon α-KG supplementation. (A) Schematic summarizing $^{13}C_6$-glucose labeling in glycolytic and TCA cycle intermediates. (B) Fractional labeling of glycolytic and TCA cycle metabolites in the retina following intraperitoneal injection of $^{13}C_6$-glucose in WT and cKO mice at P14. N=5-6 animals per group. (C) Schematic summarizing $^{13}C_5$-Gln labelling in the TCA cycle. (D) Fractional labeling of TCA cycle metabolites in the retina following intraperitoneal injection of $^{13}C_5$-Gln in WT and cKO mice at P14. N=6-11 animals per group. (E) ONL thickness in cKO mice at P22 as assessed by OCT following α-KG supplementation (10 mg/mL) or vehicle (water) in the drinking water from P4-P22. N=5 animals per group. Statistical differences in (B), (D) and (E) are based on an unpaired two-tailed Student's t-test where *P <0.05, **P<0.01 and ***P<0.001. Data are presented as mean ± standard error of the mean. F6P: fructose 6-phosphate, FBP: fructose 1,6-bisphosphate, DHAP: dihydroxyacetone phosphate, 2PG: 2-phosphoglycolate, PEP: phosphoenolpyruvate, Pyr: pyruvate, Lac: lactate, Gln: glutamine, Glu: glutamate, α-KG: alpha-ketoglutarate, Suc: succinate, Fum: fumarate, Mal: malate, OAA: oxaloacetate, Cit: citrate, Asp: aspartate.

1346

1347
Figure 5 – figure supplement 1. Mass isotopologue distribution of $^{13}C_6$-glucose in
glycolysis and TCA cycle intermediates in rod photoreceptor-specific *Gls*
conditional knockout (cKO) mice compared to wild-type (WT) mice. Fractional
labeling of glycolytic and TCA cycle intermediates with uniformly-labeled, $^{13}C_6$-glucose in
retina from rod photoreceptor-specific *Gls* cKO mice compared to WT mice at P14.
Statistical differences are based on an unpaired two-tailed Student's t-test where *P<0.05
and **P<0.01. Data are presented as mean ± standard error of the mean.

1355

1356
Figure 5 – figure supplement 2. Mass isotopologue distribution of $^{13}C_5$-Gln in TCA
cycle intermediates in rod photoreceptor-specific *Gls* conditional knockout mice
(cKO) compared to wild-type (WT) mice. Fractional labeling of TCA cycle intermediates
with uniformly-labeled, $^{13}C_5$-Gln in retina from rod photoreceptor-specific *Gls* cKO mice
compared to WT mice at P14. Statistical differences are based on an unpaired two-tailed
Student's t-test where *$P<0.05$, **$P<0.01$ and ***$P<0.001$. Data are presented as mean ±
standard error of the mean.

1364

1365

Figure 6. GLS cKO retina demonstrates decreased nonessential amino acids, ISR activation, and decreased global protein synthesis. (A) Amino acids significantly altered in the *Gls* cKO mouse retina at P14. Relative abundance is the ion intensity relative to WT retina. N=6-7 animals per group. (B) Ratio of glutamine to glutamate in WT and cKO retina. N=6-7 animals per group. (C) Western blot of ISR proteins phospho-eIF1α^{S51} (p-eIF2α), total eIF2α and ATF4 in WT and cKO mice. N=3-4 animals per group. (D) Quantitation of Western blot in Panel C. (E) Western blot of protein puromycinylation in the WT and cKO mouse retina at P14 harvested 30 min after systemic puromycin administration. (F) Quantitation of puromycin incorporation in WT and cKO retina. N=4-5 animals per group. (G) ONL thickness at P21 in cKO mice as assessed by OCT following intraperitoneal injection of ISRIB (2.5 mg/kg) or vehicle (50% PEG 400, 43.4% saline, 6.6% DMSO) from P5-P21. N=3-6 animals per group. (H) ONL thickness at P21 in cKO mice as assessed by OCT following intraperitoneal injection of Asn (200 mg/kg) or vehicle (PBS) from P5-P21. N=3-5 animals per group. Statistical differences in (A), (B), (D), (F), (G), and (H) are based on an unpaired two-tailed Student's t-test where *P <0.05, **P<0.01 and ***P<0.001. Data are presented as mean ± standard error of the mean. Gln: glutamine, Glu: glutamate, Asp: aspartate, ONL: outer nuclear layer.

1383

1384
Figure 6 – figure supplement 1. Amino acid levels are altered in the IND-cKO
retina. Relative abundance of glutamine (Gln), glutamate (Glu), aspartate (Asp) and
asparagine (Asn) in retinal tissue from WT and IND-cKO 10 days post tamoxifen (TAM),
which is prior to PR degeneration. N=5 animals per group. Statistical differences are
based on an unpaired Student's t-test where *P<0.05. Data are presented as mean ±
standard error of the mean.
1391

1392
Figure 6 – figure supplement 2. ISRIB does not impact retinal anatomy in WT mice
and prolongs PR survival in the cKO retina. (A) ISRIB (2.5 mg/kg) or vehicle (50% PEG
400, 43.4% saline and 6.6% DMSO) was delivered intraperitoneally in WT mice starting
at P5. Total retinal (TRT), outer nuclear layer (ONL) and the inner segment/outer segment
thickness were measured at P21 by OCT. N=3 animals per group. (B) ONL thickness at
P28 in cKO mice as assessed by OCT following intraperitoneal injection of ISRIB (2.5
mg/kg) or vehicle (50% PEG 400, 43.4% saline and 6.6% DMSO) from P5-P28. N=4-5
animals per group. Statistical differences in (A) and (B) are based on an unpaired two-tail
Student's t-test where *P <0.05 and **P<0.01. Data are presented as mean ± standard
error of the mean.
1403

Figure 6 – figure supplement 3. Histology confirms ISRIB and Asn prolong photoreceptor survival in the GLS cKO retina. Representative hematoxylin and eosin stained retinal sections from rod photoreceptor-specific *Gls* conditional knockout (cKO) mice at P21 following intraperitoneal injection with A) ISRIB (2.5 mg/kg) or vehicle (50% PEG 400, 43.4% saline, 6.6% DMSO) or B) Asn (200 mg/kg) or vehicle (PBS) from P5-P21. ONL, outer nuclear layer; INL, inner nuclear layer. Asn, asparagine. Scale bar is 50 µm.

Acknowledgments
Funding for this research was supported by a NEI K08EY031757, a RPB Unrestricted Grant, and the Global Ophthalmology Awards Program (GOAP), a Bayer-sponsored initiative committed to supporting ophthalmic research across the world. This work utilized the Vision Research Core funded by P30 EY007003 from the National Eye Institute. C.A.L. was supported by the following grants: R37CA237421, R01CA248160, R01CA244931, and P01HL149633. C.G.B was supported by R01EY029675. We thank Dr. Stephen Rayport in the Department of Psychiatry, Columbia University, New York, NY, for providing the *Gls^{flox/flox}* mice. We kindly thank Dr. Stephen Tsang in the Department of Ophthalmology, Columbia University, New York, NY, for the photoreceptor-specific, tamoxifen inducible mouse model (*Pde6g-Cre^{ERT2}*).

Author Contributions

Moloy T. Goswami, Conceptualization, Formal Analysis, Investigation, Writing – review and editing; Eric Weh, Conceptualization, Formal Analysis, Investigation, Writing – original draft preparation, Writing – review and editing, Visualization, Supervision; Shubha Subramanya, Formal Analysis, Investigation, Writing – review and editing, Visualization; Katherine M. Weh, Writing – review and editing, Visualization; Hima Bindu Durumutla, Investigation; Heather Hager, Investigation; Nicholas Miller, Investigation; Sraboni Chaudhury, Investigation; Anthony Andren, Resources; Peter Sajjakulnukit, Resources; Li Zhang, Resources; Cagri G. Besirli, Writing – review and editing, Supervision; Costas A. Lyssiotis, Conceptualization, Methodology, Resources, Supervision; Thomas J. Wubben, Conceptualization, Methodology, Formal Analysis, Writing – original draft preparation, Writing – review and editing, Visualization, Supervision, Funding acquisition.

Declaration of Interests

Moloy T. Goswami declares no competing interests.
Eric Weh declares no completing interests.
Shubha Subramanya declares no competing interests.
Katherine M. Weh declares no competing interests.

Hima Bindu Durumutla declares no competing interests.
Heather Hager declares no competing interests.
Nicholas Miller declares no competing interests.
Sraboni Chaudhury declares no competing interests.
Anthony Andren declares no competing interests.
Peter Sajjakulnukit declares no competing interests.
Li Zhang declares no competing interests.
Cagri G. Besirli owns Johnson & Johnson stock and has equity interest in Ocutheia and
iRenix Medical.
Costas A. Lyssiotis has consulted for Astellas Pharmaceuticals, Odyssey Therapeutics,
Third Rock Ventures, and T-Knife Therapeutics. C.A.L. is an inventor on patents pertaining
to Kras regulated metabolic pathways, redox control pathways in pancreatic cancer, and
targeting the GOT1-ME1 pathway as a therapeutic approach (US Patent No: 2015126580-
A1, 05/07/2015; US Patent No: 20190136238, 05/09/2019; International Patent No:
WO2013177426-A2, 04/23/2015).
Thomas J. Wubben has equity interest in Ocutheia.

1461

1462   **Supplemental Information**
1463   Supplementary File 1. Primers utilized in qRT-PCR experiments to measure gene
1464   expression.
1465

Gene	Forward Primer	Reverse Primer
Gls	CGTTCCATGTTGGTCTTCCT	ATCACCGACTTCACCCTTTG
Gls2	GACCGTGGTGAACCTGCTAT	ACCTCCAGGTGGTTGAACTG
Actb	AGCCATGTACGTAGCCATCC	CTCTCAGCTGTGGTGGTGAA
Aco2	CAACATGGGTGCAGAAATTG	GTGAGCCAAGTCAGGGGTAA
Idh3a	GAGGTTTTGCTGGTGGTGTT	TCCTCCTGGTCCTTGAATTG
Idh3b	ATCTGAGCGAGGTGCAGAAT	TACGTTGGCAAACAAATCCA
Idh3g	TGTAAGCTCCAACGCTGATG	CACTCCTGGCAGGCTCTTAC
Ogdh	ACATGGCACAGTGCATCATT	ACATCTGCAGAAACCGCTCT
Suclg1	AGATTCCCTTGGTTGTGTGC	GGGTTGTTTGGTGAACTGCT
Suclg2	CTTTGGTGGGATCGTCAACT	AACAGCTTTCTTGGCTGCAT
Sdha	ACACAGACCTGGTGGAGACC	GGATGGGCTTGGAGTAATCA
Sdhb	ACTGGTGGAACGGAGACAAG	TTAAGCCAATGCTCGCTTCT
Sdhc	GGAGGGGTCTCTCTTTTTGG	AAGTGTCGGATCCCATTCAG
Sdhd	GATCCCTGCTGGGTACTTGA	AAGTAGCAAAGCCCAGCAAA
Fh1	AGCAATGCATATTGCTGCTG	CGCATACTGGACTTGCTGAA
Mdh1	GAAGCCCTGAAAGACGACAG	TCGACACGAACTCTCCCTCT
Mdh1b	CATAGCCACCACCCTGAAGT	TGAATCAGATCGCCTGTGAG
Mdh2	GCTTTGTCTTCTCCCTCGTG	CAAAGTCCTCGCCTTTCTTG
Cs	TGCCTAAGGATCCCATGTTC	TTCATCTCCGTCATGCCATA
Acly	GAAGCTGACCTTGCTGAACC	CTGCCTCCAATGATGAGGAT
Pdha1	GGGACGTCTGTTGAGAGAGC	TGTGTCCATGGTAGCGGTAA
Pdhb	TCGAAGCCATAGAAGCCAGT	AGGCATAGGGACATCAGCAC
Hk2	GGGACGACGGTACACTCAAT	GCCAGTGGTAAGGAGCTCTG
Pdk1	GGCGGCTTTGTGATTTGTAT	ACCTGAATCGGGGGATAAAC

Pdk2	AGGAAGTCAATGCCACCAAC	GAGGGCCACCATAATCTTGA
Pdk3	GTGGAGTCCCACTTCGAAAA	AAAACTGGCAGCCTCTCAAA
Pdk4	CCTTTGGCTGGTTTTGGTTA	CCTGCTTGGGATACACCAGT
Pcx	ATGTTGTGGACGTGGCAGTA	AATCGAAGGCTGCGTACAGT
Bcat1	TAGAATGTGCCGATCTGCTG	CTTTGGAAGGCTTCTTGACG
Bcat2	GGACCCATGAAGACGGAGTA	CCCGCTTCAATTCCTTCATA
Me1	GGGATTGCTCACTTGGTTGT	GTTCATGGGCAAACACCTCT
Me2	TTGTGTTCCCTGCATGGTTA	ATACAGACGGGCAGACCAAC
Sod1	CCAGTGCAGGACCTCATTTT	CACCTTTGCCCAAGTCATCT
Sod2	CCGAGGAGAAGTACCACGAG	GCTTGATAGCCTCCAGCAAC
Slc7a11	CCCAGATATGCATCGTCCTT	CGTCTGAACCACTTGGGTTT
Gch1	CACCAAGGGATACCAGGAGA	AGCCAATATGGACCCTTCCT
Chac1	GTACGGCTCCCTAGTGTGGA	GTCTTCAAGGAGGGTCACCA
Ptgs2	AGAAGGAAATGGCTGCAGAA	GCTCGGCTTCCAGTATTGAG
Atg5	AGATGGACAGCTGCACACAC	GCTGGGGGACAATGCTAATA
Sqstm1	GCTCAGGAGGAGACGATGAC	AGAAACCCATGGACAGCATC
Ripk1	CCTGCTGGAGAAGACAGACC	CATCATCTTCCCCTCTTCCA
Ripk3	ACACGGCACTCCTTGGTATC	CCGAACTGTGCTTGGTCATA
Casp3	GGGCCTGTTGAACTGAAAAA	CCGTCCTTTGAATTTCTCCA
Casp8	CCTAGACTGCAACCGAGAGG	GCAGGCTCAAGTCATCTTCC
Casp9	GATGCTGTCCCCTATCAGGA	GGGACTGCAGGTCTTCAGAG

1466
1467 Supplementary File 2. List of metabolites and their parameters detected by LC-MS/MS.
1468

Compound Name	Precursor Ion (Da)	Product Ion (Quantifier) (Da)	Product Ion (Qualifier) (Da)	Retention Time (min)	Collision Energy (Volts)
±-Mevalonolactone	189.1	59.1		2.91	6
2-2-Dimethyl Succinic acid	145.1	101.1		14	11
	145.1		127.1	14	8
2-3-Dihydroxybenzoic acid	153	109		14.5	15
	153		53.2	14.5	25
2-3-Dihydroxyisovalerate	133.1	75.1		10.8	10
	133.1		57.2	10.8	15
2-3-Pyridinedicarboxylic acid	166	122		14.9	6
	166		78.1	14.9	13
2-4-Quinolinediol	160	117.9		15.2	19
	160		42.2	15.2	29
2-Deoxyadenosine	250	134.05		6.5	12
	310.1	250		6.5	4
2-Deoxyadenosine 5-diphosphate	410	79.1		15.7	44

	410		158.7	15.7	24
2-Deoxyadenosine 5-monophosphate	330	79		12.7	48
	330		134.05	12.7	28
2-Deoxycytidine	286.1	226		2	4
	286.1		93	2	20
2-Deoxycytidine 5-diphosphate	386	79		14.7	36
	386		97	14.7	20
2-Deoxycytidine 5-monophosphate	306	79		10.4	44
	306		194.9	10.4	14
2-Deoxy-D-glucose 6-phosphate	243	79.1		9.1	45
	243		97	9.1	14
2-Deoxy-D-ribose	193.1	59.2		1.4	16
	193.1		133	1.4	0
2-Deoxyguanosine	266.1	150		4	12
	266.1		108	4	36
2-Deoxyguanosine 5-diphosphate	426	79.1		15.4	48
	426		158.9	15.4	28
2-Deoxyguanosine 5-monophosphate	346	79.1		12.6	48
	346		97	12.6	36
2-Deoxyinosine	251.1	135		3.7	20
	251.1		108	3.7	42
2-Deoxyribose 5-phosphate	213	79.1		9.23	40
	213		96.9	9.23	12
2-Deoxyuridine	227.1	184		2.7	6
	227.1		42.2	2.7	20
2-Deoxyuridine 5-triphosphate	467	158.8		16.8	36
	467		368.9	16.8	20
2-Isopropylmalic acid	175.1	113		15.5	13
	175.1		115	15.5	13
2-Ketobutyrate	101	57.2		12.4	4
2-Methyl-1-butanol	87.1	43.3		9.6	5
2-Phosphoglyceric acid	185	97		14.8	14
	185		79	14.8	37
3-2-Hydroxyethylindole	160.1	130.06		15.4	14
	160.1		142.07	15.4	16
3-Dehydroshikimic acid	171	127		7.4	8
	171		109	7.4	17
3-Hydroxyanthranilic acid	152	108		12.1	12
	152		107	12.1	23

Compound					
3-Hydroxy-DL-kynurenine	223.1	205.9		2.3	4
	223.1		162	2.3	10
3-Hydroxyphenylacetic acid	151	107.1		14.2	4
	151		65.2	14.2	26
3-Indoleacetic acid	174.1	130		16.5	7
	174.1		128	16.5	19
3-Methylglutaric acid	145.1	101.1		14.1	11
	145.1		41.3	14.1	52
4-Aminobenzoic acid	136	92		9.8	8
4-Guanidobutyric acid	144.1	102		1.4	8
	144.1		41.2	1.4	32
4-Hydroxybenzoic acid	137	93		12.1	14
	137		65.2	12.1	34
4-Hydroxy-L-glutamic acid	162	144		5.3	6
	162		72.1	5.3	16
4-Hydroxyphenyl-pyruvic acid	179	107		14.7	4
4-Methyl-2-oxovaleric acid	129.1	85.1		13.7	7
4-Pyridoxic acid	182	138		15.3	12
	182		108.1	15.3	20
4-Quinolinol	144	66.1		9.2	44
	144		116	9.2	28
5-Deoxy-5-(methylthio)adenosine	356.1	134		11.8	12
	356.1		296	11.8	4
5-Hydroxy-3-indoleacetic acid	190.1	144		13	21
	190.1		116	13	46
5-Methoxytryptamine	189.1	144		2.4	28
	189.1		174	2.4	12
6-Hydroxynicotinic acid	138	94.1		9	10
	138		42.3	9	27
Adenine	134	107		2.8	18
	134		92.1	2.8	20
Adenosine	266	134		6.4	10
	326.1	134		6.4	20
Adenosine 3-5-cyclic monophosphate	328	134		13.4	24
	328		79	13.4	48
Adenosine 5-diphosphate	426	159		15.4	28
	426		328	15.4	16
Adenosine 5-monophosphate	346	79		11.6	38
	346		97	11.6	24

Compound					
Adenosine 5-triphosphate	506	408.1		17.2	22
	506		159	17.2	38
Adenylosuccinic acid	462	97		16.4	24
	462		134.05	16.4	48
Adipic acid	145.1	101.1		13.9	11
	145.1		83.1	13.9	12
AICAR	257.1	125		2.3	10
	257.1		42.2	2.3	44
Allantoin	157	42.2		1.3	16
	157		97	1.3	12
alpha-D(+)Mannose 1-phosphate	259	79		7.6	48
	259		97	7.6	14
alpha-D-Glucose-1-phosphate	259	79		7.6	28
	259		240.9	7.6	9
alpha-Ketoglutaric acid	145	101		14.3	5
	145		57.2	14.3	9
Arabinose-5-phosphate	229	79		8.4	36
	229		97	8.4	8
Argininosuccinic acid	289	88		4.6	36
	289		131.1	4.6	27
beta-Nicotinamide adenine dinucleotide	662.1	540		9.5	12
	662.1		327.9	9.5	36
beta-Nicotinamide mononucleotide	333	251.1		3.7	12
	333		135.1	3.7	36
Cellobiose	341.1	161		1.3	5
Chorismic acid	225	189		6.7	6
cis-Aconitic acid	173	129		15.5	4
	173		85.1	15.5	8
Citramalic acid	147	85.1		13.8	12
	147		87	13.8	14
Citric acid	191	111.01		14.9	10
	191		87.01	14.9	16
CoA	766.1	408.1		17.8	40
	766.1		686.2	17.8	40
Creatine	130.1	88.1		1.3	8
	130.1		41.2	1.3	36
Creatine phosphate	210	79		7	16
	210		96.9	7	4
Creatinine	112	41.2		1.35	24
	112		88.1	1.35	16
Cysteine	120	33.2		1.31	10
Cytidine	242.1	109		1.8	8

	242.1		42.2	1.8	16
Cytidine 5-diphosphate	402.01	79		14.4	48
	402.01		158.92	14.4	24
Cytidine 5-triphosphate	482	158.8		17	40
	482		79	17	40
Cytidine-5-monophosphate	322	79		9.7	44
	322		97	9.7	22
Cytosine	110	67.1		1.4	8
	110		42.2	1.4	12
D-+-Galactosamine	238.1	159.9		1.1	8
	238.1		100	1.1	6
Deoxyadenosine 5-triphosphate	490	391.9		17.2	24
	490		158.9	17.2	32
Deoxycytidine 5-triphosphate	466	158.9		16.8	28
	466		367.9	16.8	20
Deoxyguanosine 5-triphosphate	506	407.9		17.2	20
	506		158.8	17.2	32
Deoxythymidine 5-triphosphate	481	158.8		17.2	36
	481		383.1	17.2	20
D-erythro-Dihydrosphingosine	300.3	199		8.9	8
	300.3		282	8.9	6
D-Fructose 1,6-biphosphate	338.9	97		14.8	22
	338.9		241.01	14.8	12
D-Fructose 6-phosphate	259	79		7.6	48
	259		97	7.6	14
D-Gluconic acid	195.1	75.2		5.9	18
	195.1		129	5.9	10
D-Glucosamine 6-phosphate	258	79		1.6	48
	258		97	1.6	16
D-Glucose 6-phosphate	259	79		7.6	48
	259		97	7.6	14
Dihydroxyacetone phosphate	169	79		10.5	32
	169		97	10.5	6
DL-2-Aminoadipic acid	160.1	142		4.3	9
	160.1		116	4.3	12
DL-Glyceraldehyde 3-phosphate	169	79		11.7	28
	169		97	11.7	4
DL-Isocitric acid	191	111.02		15.1	11
	191		173	15.1	6

Compound					
D-Maltose	341.1	161		1.3	5
	341.1		179	1.3	4
D-Mannose	179.1	89		1.3	4
	179.1		59.2	1.3	16
D-pantothenic acid	218	88		12.1	10
	218		146.08	12.1	14
D-Ribose 5-phosphate	229	79		8.5	48
	229		97	8.5	10
D-Ribulose 1,5-biphosphate	308.9	79		14.7	46
	308.9		97	14.7	22
D-Sedoheptulose-7-phosphate	289	96.9		8.2	18
	289		79	8.2	48
D-Xylose	149.05	89.02		1.2	4
	149.05		71.01	1.2	4
D-Xylulose-5-phosphate	229	79		8.5	36
	229		138.98	8.5	8
Epicatechin	289.1	245		12.2	11
	289.1		109	12.2	26
Flavin adenine dinucleotide	784.1	346		17	35
	784.1		437	17	30
Folinic acid	472.2	315		14.8	28
	472.2		343	14.8	20
Galactonic acid	195.1	75.1		5.9	18
	195.1		129	5.9	11
gamma-Aminobutyric acid	102	84.1		1.2	8
gamma-Glu-Cys	249.1	128		8.1	8
	249.1		171	8.1	9
GlcNAC	220	119		1.4	2
Glucoheptonic acid	225.1	129		5.8	11
	225.1		75.1	5.8	18
Glutathione Reduced	306.1	143		8.43	17
	306.1		128.1	8.43	17
Glyceric acid	105	75.1		5.9	8
	105		57.2	5.9	14
Glyoxylic acid	73	45.2		5.9	6
Guanine	150	133.02		2	12
	150		66.01	2	36
Guanosine	282.1	150		3.9	17
	282.1		132.9	3.9	32
Guanosine 3,5-cyclic monophosphate	344	150.042		12.2	24
	344		133.02	12.2	44

Guanosine 5-diphosphate	442	150		14.8	31
	442		158.9	14.8	20
Guanosine 5-triphosphate	522.3	158.9		17	36
	522.3		424	17	20
Homocitrate	205	125		15.4	10
	205		145	15.4	10
Hypoxanthine	135	92		2	17
	135		65.1	2	28
Indoline-2-carboxylate	162.1	116		16.1	16
	162.1		118	16.1	11
Inosine	267.1	135		3.3	22
	267.1		108	3.3	44
Inosine 5-diphosphate	427	158.9		14.5	28
	427		135	14.5	24
Inosine 5-monophosphate	347	79		11.1	44
	347		97	11.1	22
Inosine 5-triphosphate	507	409		17.2	19
	507		159	17.2	40
Isopentenyl pyrophosphate	245	209		12.8	4
Isopentyl acetate	129.1	85.2		14.6	6
	129.1		41.2	14.6	12
Itaconic acid	129	85.1		13.6	6
	129		41.3	13.6	12
Ketoisovaleric acid	115	71.2		16.56	4
Ketovaleric acid	115	71.2		14.5	4
Lactic acid	89	43.3		7.4	10
	89		45.3	7.4	9
L-Arabinose	149	89.1		1.4	4
	149		59.2	1.4	12
L-Arabitol	151	89		1.3	12
	151		71.02	1.3	16
L-Arginine	173.1	131		1.1	11
	173.1		156	1.1	8
L-asparagine	131	113		1.2	4
	131		70	1.2	12
L-Aspartic Acid	132	88		4.9	10
	132		71	4.9	14
L-Canavanine	175.1	74.04		1.1	10
	175.1		118	1.1	8
L-Carnitine	220.1	146		1.2	4
L-Citrulline	174.1	131		1.3	8
	174.1		42.2	1.3	48

Compound	Q1	Product 1	Product 2	RT	CE
L-Cystathionine	221.1	134		1.3	8
	221.1		120	1.3	9
L-Cystine	239	74.1		1.2	20
	239		120	1.2	4
L-Dihydroorotic acid	157	113		7.3	5
	157		42.2	7.3	30
L-Glutamic acid	146	102		4.5	11
	146		128	4.5	8
L-Glutamine	145.1	127		1.3	7
	145.1		109	1.3	10
L-Glutathione (oxidized)	611.1	306.07		13.6	24
	611.1		272.09	13.6	28
L-Histidine	154.1	137		1.1	12
	154.1		93.1	1.1	16
L-Homocysteine	134	116.9		1.4	8
L-Homocystine	267	132		1.5	8
	267		115	1.5	16
L-Homoserine	118	72.1		1.3	10
	118		55	1.3	16
L-Hydroxyglutaric acid	147	128.9		13.8	7
	147		85.1	13.8	13
Lipoamide	204.1	158		16.8	4
	204.1		64.1	16.8	20
L-Isoleucine	130	45		2.1	16
L-Kynurenine	207.1	190		3.9	4
	207.1		144	3.9	20
L-Leucine	130	45		2.1	16
L-Malic acid	133	115		13.8	8
	133		71.1	13.8	14
L-Methionine	148	47.2		1.8	12
	148		100	1.8	6
L-Phenylalanine	164.1	147		4.5	9
	164.1		103	4.5	15
L-Proline	114	68.1		1.3	12
	114		45.2	1.3	12
L-Serine	104	74.1		1.2	8
L-Sorbose	179.1	89		1.3	4
	179.1		71.02	1.3	14
L-Threonine	118	74.1		1.3	9
L-Tryptophan	203.1	116		7.9	14
	203.1		74.1	7.9	14
L-Tyrosine	180	119		2.3	15
	180		93	2.3	12
Maleic acid	115	71.2		12.8	7

Name					
	115		27.3	12.8	15
Malonic acid	103	59.2		12.8	7
	103		41.3	12.8	32
Melibiose	341.1	179		1.3	4
	341.1		89	1.3	13
Mevalonic acid	147.1	128.9		9.8	8
	147.1		103.1	9.8	11
Mevalonic acid 5-phosphate	227	79		14.8	40
	227		97	14.8	12
m-Hydroxybenzoic acid	137	93.1		13.4	10
	137		65.2	13.4	28
myo-Inositol	179	71.02		1.3	12
	239.1	179		1.3	0
N-Acetyl D-galactosamine	220.1	119		1.2	0
N-Acetyl-alpha-D-glucosamine 1-phosphate	300	97		8.7	16
	300		79	8.7	36
N-acetylaspartate	174	88.1		14	14
	174		58.1	14	22
N-acetylaspartylglutamate	303.1	285		16.1	6
	303.1		128	16.1	14
N-Acetyl-D-glucosamine 6-phosphate	300	97		8.8	18
	300		79	8.8	48
N-Acetylglutamic acid	188.1	128.04		14.1	10
	188.1		102	14.1	16
N-Acetylneuraminic acid	308.1	87		6.2	14
	308.1		169.9	6.2	10
NADH	664.1	397		15.4	32
NADPH	744.1	426.1		17.2	33
N-Carbamoyl-DL-aspartic acid	175	132		13.5	7
	175		88.1	13.5	19
N-Carbamyl-L-glutamic acid	189.1	146		13.6	7
	189.1		102.1	13.6	20
N-Formyl-L-Tyrosine	208.1	164		12.8	7
	208.1		107	12.8	12
Nicotinic acid	122	78.1		11.9	11
	122		51.2	11.9	32
Nicotinic acid mononucleotide	334	289.9		8.6	4
	334		79	8.6	40
o-Hydroxy hippuric acid	194	150		16.5	11
	194		121	16.5	22

Compound					
o-Phospho-L-Serine	184	79		10.2	36
	184		97	10.2	13
O-Phosphorylethanolamine	140	79		1.7	14
	140		63	1.7	38
Orotic acid	155	42		9.1	26
	155		40	9.1	44
O-Succinyl-L-homoserine	218.1	117		6.5	7
	218.1		118	6.5	0
Oxamic acid	88	44.3		6.5	6
	88		42.3	6.5	12
Phenylpyruvic acid	163	91.1		17.2	6
Phosphoenolpyruvic acid	167	79		15.3	12
Prephenic acid	225	101		12.4	9
Pyridoxal 5 phosphate	246	79		14.8	30
Pyridoxal hydrochloride	166.1	138		2.4	12
	166.1		108	2.4	18
Pyridoxamine	167.1	121		1.1	22
	167.1		122	1.1	18
Pyridoxine	168.1	122.1		2.4	17
	168.1		166	2.4	9
Pyruvic acid	87	43.2		9.5	4
Quinic acid	191.1	85.1		6	25
	191.1		93.1	6	24
Riboflavin	375.1	255		12.6	14
	375.1		212	12.6	28
Ribonic acid gamma lactone	147	59.2		1.5	16
	147		99	1.5	8
S-2-Aminoethyl-L-cysteine	163.1	76.1		1.1	10
	163.1		33.2	1.1	28
S-5-Adenosyl-L-homocysteine	383.1	134		5.3	24
	383.1		248	5.3	13
Salicylic acid	137	93.1		16.8	17
	137		65.2	16.8	33
Shikimic acid	173	111		5.8	7
	173		93.1	5.8	16
Succinic acid	117	73.1		13	10
	117		99	13	8
Succinic semialdehyde	101	57.2		7.1	7
Taurine	124	80		1.3	24
Taurocholic acid	514	514		20.7	50
Thiamine	264.1	234		1.1	6

	264.1		148	1.1	18
Thymidine	241.1	42.2		5.5	16
	241.1		151	5.5	6
Thymidine 5-diphosphate	401.01	97		15.4	25
	401.01		79	15.4	44
Thymine	125	42.2		2.7	14
trans-4-Hydroxy-L-proline	130.1	128		1.3	12
	130.1		66.03	1.3	16
trans-Aconitic acid	173	129		15.55	4
	173		85.1	15.55	10
trans-trans Muconic acid	141	53.3		14.7	8
Trehalose	341	89.01		1.3	12
	401.1	341		1.3	10
Trehalose 6-phosphate	421.1	241		7.9	27
Uracil	111	42.2		1.7	14
Uric acid	167	124		5.5	13
	167		42.3	5.5	26
Uridine	243.1	200		2.3	6
	243.1		110	2.3	12
Uridine 5-diphosphate	403	158.9		14.8	28
	403		79	14.8	48
Uridine 5'-diphosphogalactose	565	322.9		14.3	24
	565		384.9	14.3	29
Uridine 5-diphosphoglucose	565	322.9		14.3	24
	565		384.9	14.3	28
Uridine 5-monophosphate	323	79.1		10.8	48
	323		97	10.8	24
Uridine 5-triphosphate	483	158.9		17	36
	483		385	17	20
Vanillic acid	167	123		12.6	9
	167		108	12.6	18
Xanthine	151	108		2.1	16
	151		42.2	2.1	24
Xanthosine	283.1	151		8.2	18
	283.1		108	8.2	40
Xylitol	151.1	89		1.3	8
	151.1		71	1.3	8

1469