Hyperspectral imaging for chloroplast movement detection

Paweł Hermanowicz, Justyna Łabuz[*]

[*] Corresponding author: justyna.sojka@uj.edu.pl

Malopolska Centre of Biotechnology, Jagiellonian University, Gronostajowa 7A, 30-387 Kraków, Poland

Paweł Hermanowicz ORCID: 0000-0002-4708-442X

Justyna Łabuz ORCID: 0000-0002-2612-1658

Keywords: *Arabidopsis thaliana,* chloroplast movements, hyperspectral imaging, leaf reflectance, *Nicotiana benthamiana*, vegetation indices

Summary:

We employed hyperspectral imaging to detect chloroplast positioning in *Nicotiana benthamiana* and *Arabidopsis thaliana* leaves and assess its influence on commonly used vegetation indices. In low light, chloroplasts move to cell walls perpendicular to the direction of the incident light. In high light, they move to cell walls parallel to the light direction. Chloroplast movements result in significant changes in leaf transmittance and reflectance. The changes in leaf reflectance offer a way to examine chloroplast positioning in a non-contact way. At the same time, they may confound remote sensing of other physiological traits. The shape of reflectance spectra recorded on irradiated and non-irradiated parts of *N. benthamiana* and *A. thaliana* leaves indicated the specific position of chloroplasts. Low blue light resulted in a decrease in leaf reflectance in the green-yellow region of the spectrum. High blue light irradiation caused an increase in leaf reflectance in the visible range. The differential spectra, showing the effect of high light on leaf reflectance, exhibited a characteristic saddle in the green-yellow region and a peak at around 695 nm. Results obtained for *A. thaliana* mutants with disrupted chloroplast movements suggest that the observed spectral changes are mostly due to the chloroplast relocations. The reflectance spectra were used to train machine learning methods in the classification of leaves according to the chloroplast positioning. The convolutional network showed low levels of misclassification of leaves irradiated with high light even when different species were used for training and testing. This suggests that reflectance spectra may be used to detect the chloroplast avoidance response in heterogeneous

patches of vegetation. We also examined the correlation between chloroplast positioning and

values of indices of normalized-difference type for various combinations of wavelengths and

proposed a chloroplast movement index for validation of chloroplast positions in leaves. The

analysis of commonly used vegetation indices showed that their values may be altered due to

chloroplast rearrangements. Our work indicates that changes in leaf reflectance due to

chloroplast movements may be substantial and should be taken into account in remote sensing

studies.

Abbreviations:

ARVI - Atmospherically Resistant Vegetation Index

CAR1 - Carotenoid Reflectance Index 1

CAR2 - Carotenoid Reflectance Index 2

EVI - Enhanced Vegetation Index

NDVI - Normalized Difference Vegetation Index

mRENDVI - Modified Red Edge NDVI

mRESR - Modified Red Edge Simple Ratio Index

PAR - photosynthetically active radiation

PRI - Photochemical Reflectance Index

PSRI - Plant Senescence Reflectance Index

RENDVI - Red Edge NDVI

RGRI - Red Green Ratio Index

SG - Sum Green Index

SIPI - Structure Insensitive Pigment Index

SR - Simple Ratio Index

VOG1 - Vogelmann Red Edge Index 1

VOG2 - Vogelmann Red Edge Index 2

VOG3 - Vogelmann Red Edge Index 3

Introduction

Plants show a remarkable ability of chloroplast relocation in changing light conditions, for review see (Łabuz et al., 2022). In low light, chloroplasts move to cell walls perpendicular to the direction of incident light. This response, called chloroplast accumulation, increases light capture for photosynthesis (Gotoh et al., 2018). In high light, chloroplasts gather at cell walls parallel to the direction of incident light. Chloroplast avoidance protects the photosynthetic apparatus from excess light (Kasahara et al., 2002; Sztatelman et al., 2010), (Davis & Hangarter, 2012) or shapes light gradients within the leaf tissue (Brugnoli & Bjorkman, 1992), (Wilson & Ruban, 2020). Seed plants move chloroplasts in response to blue light, however, red light is also active in some ferns, mosses, and green algae (Suetsugu, et al., 2005). In the model plant *Arabidopsis thaliana*, blue light-induced chloroplast movements are controlled by photoreceptor serine-threonine kinases called phototropins (Christie, 2007). Two phototropins, phot1 and phot2, mediate the chloroplast accumulation response (Sakai et al., 2001). Sustained chloroplast avoidance can be elicited only by phot2 (Jarillo et al., 2001), (Kagawa et al., 2001). Thus, no substantial directional chloroplast movements are observed in the *phot1phot2* double phototropin mutant. The *phot1* mutant is less sensitive to very low light irradiances than wild-type plants, but it shows both accumulation and avoidance response (Luesse et al., 2010). In the *phot2* mutant, light elicits chloroplast accumulation regardless of irradiance, though it is preceded by a weak, transient avoidance response in high light (Luesse et al., 2010), (Łabuz et al., 2015). Chloroplast avoidance is observed in both low and high light in *jac1,* a signaling mutant, lacking JAC1, the J-domain protein required for chloroplast accumulation response 1 (Suetsugu, et al., 2005).

First microscopic chloroplast movement observations were performed by Senn (Senn, 2008), revealing a plethora of different arrangements within plant cells (Kataoka, 2015). Microscopic observations still play a pivotal role in assessing chloroplast positions within leaf cells, but they are laborious (Kagawa & Wada, 2000) and provide information for individual cells. Interpretation of microscopic observations requires care due to the steep fluence rate gradient within the mesophyll tissue. In some species, the effects of chloroplast movements on the optical properties of the leaves are so pronounced that they may be observed with the naked eye. Whereas chloroplast avoidance results in paler leaf blades, chloroplast accumulation causes the darkening of leaves. This effect is the basis for the so-called slit assay (Kagawa & Wada, 2000), and is even utilized by artists (Dolinsky & Hangarter, 2013). Changes in leaf transmittance correspond to averaged chloroplast positions within the

mesophyll and are used to assess chloroplast movements quantitatively. The development of
photometric devices for measuring blue-light-induced changes in leaf transmittance offered a
convenient approach, however, limited in its throughput and restricted mainly to laboratory
use (Shibata, 1958; Walczak & Gabrys, 1980; Yorinao & Kazuo, 1974). Leaf reflectance has
been also used to monitor chloroplast movements (Park et al., 1996), (Gorton et al., 1999). A
method using measurements of red light reflected from the leaf surface was developed to
monitor movements during the growth of *Arabidopsis thaliana* in a growth chamber (Dutta et
al., 2015). Analysis of leaf optical properties for several species indicated an important role of
the cell shape in the ability of chloroplast to move and subsequently alter the leaf absorptance
and scattering (Davis et al., 2011). A detailed study of leaf reflectance has also been
performed using *Nicotiana tabacum.* Chloroplast positioning resulted in the highest changes
in the leaf reflectance spectrum in the region between 500 and 650 nm. The changes in leaf
reflectance were about 2% in the blue and red regions and about 5% in the green region of the
light spectrum (Baránková et al., 2016). The works of (Davis et al., 2011), (Baránková et al.,
2016) used an integrating sphere to measure changes in diffuse reflectance. While this
approach yields invaluable data on the effects of chloroplast movements on the optical
properties of the leaves, it shares the limitations of the transmittance-based method in its low
throughput and requirement of direct contact with the leaf.

The aims of our work were twofold - to establish a system for high throughput
chloroplast relocation monitoring using spectra of white light reflected from leaves and to
examine whether changes in chloroplast positioning can confound the assessment of leaf traits
from commonly used vegetation indices. We have used multiple mutant lines that exhibit
disrupted chloroplast movements to distinguish between changes in leaf spectra that can be
ascribed to chloroplast relocations from changes due to other light-induced processes. To
acquire reflectance spectra and examine their spatial variability within the leaf, we have used
hyperspectral imaging, which is a high-throughput, contactless method.

Vegetation indices are a convenient approach to reduce the possibly redundant
information in spectra, focusing on the most relevant wavelengths or bands. Hyperspectral or
narrow-band vegetation indices are single values calculated using a combination of
wavelengths, which are empirically shown to correlate with the traits of interest. Vegetation
indices were developed to examine plant or canopy structure (e.g. fractional cover, green
biomass, leaf area index, senesced biomass), as well as biochemical (e.g. the content of water,
nitrogen, lignin, cellulose, and pigments: chlorophyll, carotenoids, anthocyanins) and

physiological traits (e.g. stress, changes in xanthophyll cycle pigments, fluorescence, leaf moisture) (Thenkabail et al., 2018). In particular, a large group of vegetation indices based on the visible or NIR reflectance (formulae in Supplementary Table S1) is used to assess the chlorophyll content, consisting of the classical NDVI (Rouse et al., 1974), but also SR (Rouse et al., 1974), EVI (Huete et al., 1997), ARVI (Kaufman & Tanre, 1995), RENDVI (Gitelson & Merzlyak, 1994), mRESR (Sims & Gamon, 2002), and a sum green index, (Behmann et al., 2014). VOG1, 2, and 3 link the chlorophyll and water contents (Vogelmann et al., 1993). PRI (Gamon et al., 1992), SIPI (Penuelas et al., 1995), CAR1, and CAR2 (Gitelson et al., 2002) represent carotenoid proportions, while RGRI (Gamon & Surfus, 1999), the anthocyanin-chlorophyll ratio. mRENDEVI (Sims & Gamon, 2002) and PSRI (Merzlyak et al., 1999) are used for stress and senescence estimation. The equations specifying vegetation indices are based on the visible and NIR infrared range of the solar spectrum. In the visible and near-infrared range, several bands were found particularly useful for remote sensing, including 660 – 690 nm (red-absorption maxima), 900 – 925 nm (near-infrared reflection peak), 700 – 720 nm (a portion of red-edge), and 540 – 555 nm (green reflectance maxima), 490 nm centered band (rapid change in slope of the spectra), 520 nm and 575 nm bands (the most rapid positive or negative change in reflectance), 845 nm centered band (the NIR shoulder), and 975 nm centered band (biomass/moisture sensitive) (Thenkabail et al., 2002). The estimates of plant traits based on vegetation indices may be biased due to the effects of other, nonspecific physiological responses on the leaf reflectance, such as senescence (Idso et al., 1980); as well as changes in the chlorophyll and carotenoid content (Tucker, 1977). The values of vegetation indices may be also altered by topography (Chen et al., 2020) or snow cover (Wang et al., 2023). In this work, we have investigated the impact of chloroplast positioning on commonly used vegetation indices and discussed the magnitude of their changes caused by chloroplast movements compared to the scale of changes due to the variation of other traits, reported in the literature. We have analyzed the accuracy of vegetation indices for the detection of chloroplast positioning and identified an index of the normalized difference type, tentatively referred to as a Chloroplast Movement Index, that correlates well with the chloroplast arrangement.

To examine whether a more accurate classification of leaves according to chloroplast arrangement can be obtained using whole spectra of reflectance in the visible range, we have examined the performance of three supervised machine learning procedures in the task of classifying leaves according to the chloroplast positioning based on the reflectance spectra.

We employed the support vector machine and two types of neural feed-forward network
architectures, the multilayer perceptron and convolutional networks. A mathematically
rigorous treatment of both of these types of neural networks can be found in (Caterini &
Chang, 2018) and (Berner et al., 2022). The Multilayer Perceptron (Pal & Mitra, 1992) is a
classical architecture, consisting of a variable number of densely connected layers of neurons.
It has been used already in the early works applying machine learning for the analysis of
reflectance spectra and remotely sensed data, as reviewed in (Mas & Flores, 2008). In
particular, it has been successfully applied to the detection of fungal infections from leaf
spectra (Moshou et al., 2004) and to estimate carotenoid content through inversion of the
PROSPECT model (Hao et al., 2023). In convolutional neural networks, densely connected
layers resembling the multilayer perceptron are preceded by additional layers that convolve
their input with an arbitrary number of kernels. The benefits of convolutional networks
include their good generalization (Alzubaidi et al., 2021) so that the network performance is
usually not substantially degraded when applied to an unseen dataset. They have recently
become a popular tool for analysis of spectral data. Their applications include the prediction
of the concentration of photosynthetic pigment through the inversion of leaf optical models
(Annala et al., 2020), (Shi et al., 2022) as well as the estimation of nitrogen (Pullanagari et al.,
2021) and water content (Zhang et al., 2023) from the leaf reflectance spectra. However, an
architecture based on the multilayer perceptron was found to be superior to a convolutional
network for the prediction of photosynthetic performance in Japanese beaches based on leaf
reflectance (Song & Wang, 2021).

Materials and methods

Plants and Growth Conditions

Seeds of *Nicotiana benthamiana, Arabidopsis thaliana* wild-type Col-0, *phot1*:
SALK_088841 (Lehmann et al., 2011), the *phot2* (*npl1-1*) mutant (Jarillo et al., 2001),
phot1phot2 (SALK_088841 crossed with *npl1-1*), (Hermanowicz et al., 2019), *jac1-3:*
WiscDsLox457-460P9 (Hermanowicz et al., 2019) were sown in Jiffy-7 pots (Jiffy Products
International AS) and vernalized at 4°C for two days. All plants were transferred to a growth
chamber (Sanyo MLR 350H) at 23°C, 80% relative humidity, with a photoperiod of 10 h light
and 14 h darkness. The irradiance at the plane of leaf rosettes was 70 $\mu mol \cdot m^{-2} \cdot s^{-1}$. Light was
supplied by fluorescent lamps (FL40SS.W/37, Sanyo, Japan). After 4 weeks *N. benthamiana*
were transferred to pots containing soil and moved to a walk-in growth chamber with the

187 photoperiod of 10 h light and 14 h darkness, equipped with white LEDs supplying irradiance

188 of ca. 100 μmol·m^{-2}·s^{-1}.4-week-old *Arabidopsis* and 6-week-old *N. benthamiana* plants were

189 used for the experiment. Plants were dark-adapted for at least 16 h for all kinds of

190 measurements of chloroplast movements. For microscopic observations, detached leaves were

191 either kept in darkness or irradiated with blue light of 1.6 μmol·m^{-2}·s^{-1} or 120 μmol·m^{-2}·s^{-1} for

192 1 h (EPILED, 1 W, 460 nm). For hyperspectral imaging half of the leaf was covered with

193 aluminum foil and the other half was irradiated with the same light irradiance as for confocal

194 microscopy.

195 *Microscopic observations*

196 Microscopic observations were performed with the Axio Observer.Z1 inverted

197 microscope (Carl Zeiss, Jena, Germany) and the LSM 880 confocal module. The long-

198 distance LCI Apochromat 25x, NA 0.8, objective was used with glycerol immersion.

199 Chloroplast positioning was observed in the palisade parenchyma of wild-type *Nicotiana*

200 *benthamiana* and *Arabidopsis thaliana* leaves as well as *phot1, phot2, phot1phot2,* and *jac1*

201 mutants. The 633 nm He-Ne laser was used to excite chlorophyll, and emission in the range of

202 661 - 721 nm was recorded as the magenta channel. Stacks were collected on the upper

203 surface of the leaves. Projection images were calculated from slices corresponding to the first

204 ca. 90 μm of stacks, starting from the upper surface of the epidermis. This range included the

205 epidermis and upper parts of palisade cells.

206 *Photometric method*

207 Leaf transmittance changes due to chloroplast movements were assessed with the

208 photometric method (Gabryś et al., 2017), using a custom-built photometric setup.

209 Chloroplast responses were induced with blue (peak at 455 nm, M455L4 LED, Thorlabs)

210 actinic light of 1.6 or 120 μmol m^{-2} s^{-1}. The red measuring beam was produced by a 660 nm

211 LED (M660L4, Thorlabs) and modulated at 1033 Hz. The beams were collimated and

212 combined with a dichroic mirror (DMLP550, Thorlabs). They were then directed toward a

213 detached leaf, mounted in front of a port of an integrating sphere (IS200-4, Thorlabs). The

214 signal was detected with a photodiode detector (DET100A2, Thorlabs) mounted at another

215 port of the sphere. Leaf transmittance curves were processed with a custom-written

216 Mathematica (Wolfram Research) script. Irradiance was measured at the sample plane with

217 the LI-190R sensor (LI-COR) and Keithley 6485 picoampere meter.

218 *Hyperspectral imaging*

Immediately after blue irradiation, the aluminum foil was removed from the covered leaf

half, and hyperspectral images of whole leaves were collected using a Headwall Nano

hyperspectral Camera (Nano-Hyperspec VNIR 400-1000 nm) equipped with a 12 mm

objective (Cinegon F/1.4/12-0906, C-Mount, Schneider Kreuznach, 400-1000 nm). The

camera was mounted on a rotating platform (Sevenoak SK-ebh01 Pro), 30 cm from the table.

A halogen lamp (Hedler), located 58 cm from the table, was used as a white light source

delivering irradiance of 280 $\mu mol \cdot m^{-2} \cdot s^{-1}$ in the leaf plane for collecting the reflected light in

the range of 400 - 1000 nm. The angle of incidence of light with respect to the leaf normal

was ca. 10°. All images were taken on a black velvet background. White, diffusely-reflecting

PTFE disks (SM05CP2C, Thorlabs) were used as reflectance standards. The extraction of

spectra and calculation of the mean difference in reflectance spectra between irradiated and

darkened halves was performed using a custom script written in Mathematica 13.3 (Wolfram

Research).

Machine learning procedures for leaf classification

The performance of three types of machine learning procedures, the support vector

machine, the multilayer perceptron, and the convolutional network, in the classification of

leaves based on reflectance spectra were examined using Mathematica 13.3 (Wolfram

Research). Spectra were normalized, truncated to the 400 – 750 nm range, and divided into

three classes, corresponding to dark-adapted, low-light, and high-light irradiated leaves. As a

data augmentation procedure, spectra were extracted from square blocks (60 by 60 pixels),

randomly selected within the leaf area. Training and validation were performed using spectra

recorded on *N. benthamiana* leaves. To assess the generalization capability of the trained

classifiers, final testing was performed on spectra recorded from *N. benthamiana*, as well as

on spectra from *A. thaliana*. The *N. benthamiana* dataset was first split into the test set (25%)

and the set used for building classifiers (75%). The latter one was further split into the training

set and the validation (held-out) set, using balanced five-fold cross-validation. Cross-

validation was repeated ten times, with different random partitioning of the dataset into the

folds. The performance of the trained classifiers on the validation set guided the selection of

model hyperparameters. For the support vector machine, five kernel types were examined

(linear, polynomial, sigmoid, and radial Gaussian function). Multilayer perceptron

architectures with three or fewer densely connected layers were examined, with up to 30

neurons per layer and a decreasing number of neurons in the input-output direction. Each

layer was followed by a batch normalization layer and a rectified linear unit. For the

convolution networks, we examined several combinations of the number of convolution
layers (1 to 4), the width of the convolution and max pooling kernels, the number of feature
maps, as well as number of densely connected layers. Successive convolutional layers had
decreasing kernel width and increasing number of feature maps, which is a common
approach. Rectified linear units were always used for activation functions and batch
normalization layers were inserted after each densely connected layer to regularize the trained
network. The final layer of neurons in the multilayer perceptron and convolutional networks
was connected to a softmax layer. As the modeled response variable was categorical, with
three classes, the training consisted of minimizing the negative log-likelihood of the
multinomial distribution, corresponding to the cross-entropy function. The training was
carried out in 2000 epochs, using the Adam optimizer, with a batch size of 64.

Results

Hyperspectral imaging of chloroplast movements in Nicotiana benthamiana

To assess the accuracy of hyperspectral imaging for detecting chloroplast movements,
we chose a plant with leaf blades of a considerable size, *Nicotiana benthamiana,* which
exhibited substantial chloroplast movements (Davis et al., 2011). Blue light of 1.6 $\mu mol \cdot m^{-2} \cdot s^{-1}$
induced a decrease in leaf transmittance corresponding to chloroplast accumulation, whereas
blue light of 120 $\mu mol \cdot m^{-2} \cdot s^{-1}$ increased leaf transmittance in line with the chloroplast
avoidance response (Fig. 1. A, B). The changes in chloroplast positions due to blue light
irradiation observed under the microscope (Fig. 1. C), were visible in the hyperspectral
images of partly irradiated leaves (Fig. 1. F, G). The leaf part irradiated with blue light of
1.6 $\mu mol \cdot m^{-2} \cdot s^{-1}$ for 1 h was darker than the part kept simultaneously in darkness by covering
it with an aluminum foil. The leaf part irradiated with blue light of 120 $\mu mol \cdot m^{-2} \cdot s^{-1}$ for 1 h
appeared paler than the part kept under the aluminum foil (Fig. 1. F, G). The mean reflectance
of dark-adapted leaves was ca. 11% in visible light (400-700 nm). The average difference of
reflectance in this range between leaves with chloroplasts in avoidance and accumulation
positions was ca. 2.3% (Fig. 1. D, E). Differential leaf reflectance curves were calculated
from the spectra recorded for non-irradiated and blue light-irradiated leaf parts (Fig. S1,
Fig. 1. D, E). After irradiation with 1.6 $\mu mol \cdot m^{-2} \cdot s^{-1}$, a decrease in differential leaf reflectance
curves induced by blue light was observed in the green-yellow region (500-600 nm).
Irradiation with blue light of 120 $\mu mol \cdot m^{-2} \cdot s^{-1}$ resulted in a characteristic 'saddle' in the green-
yellow region (500-600 nm) of differential reflectance curves. In addition, a trough at ca.

684 nm and a peak at ca. 695 nm were visible. One-hour irradiation with high blue light

induces quenching of chlorophyll fluorescence (Pfündel et al., 2018). Thus, we may expect

that the maxima of the chlorophyll emission spectrum (measured in intact leaves)

corresponded to troughs in the high-minus-low-light differential curves. Accordingly, the

trough at ca. 684 nm is near a small peak at ca. 685 nm visible in the fluorescence emission

spectrum of several angiosperm species (Chappelle et al., 1985), (Agati, 1998). The peak at

695 nm in the differential reflectance spectra appears to correspond to the inflection point

between the overlapping peaks in the chlorophyll emission spectrum. As the main peak of the

emission spectrum from intact leaves is located at ca. 740 nm, fluorescence quenching likely

contributes to the steep decrease in the differential reflectance spectrum visible between

700 nm and 745 nm.

Monitoring chloroplast movements in Arabidopsis thaliana wild type and chloroplast

movement mutants

To better understand the relationship between chloroplast positioning and leaf

reflectance, we examined light-induced changes in reflectance spectra of several *Arabidopsis*

thaliana mutants with disrupted chloroplast movements. The use of mutants was necessary to

distinguish between spectral changes due to chloroplast movements and other processes of

blue-light-induced responses. Microscopic observations (Fig. 2. A) were performed to enable

direct comparisons of the movement aberrations characteristic of mutants. The recorded

spectra for *Arabidopsis* wild-type and all chloroplast movement mutants are shown in

Fig. S2 & S3. For *Arabidopsis* wild-type plants, patterns of differential leaf reflectance curves

were similar to *N. benthamiana* (compare Fig 1. D, E and Fig. 2. B). A decrease in the green-

yellow region (500-600 nm) after blue light of 1.6 μmol·m^{-2}·s^{-1} and a saddle with a peak at ca.

695 nm after blue light of 120 μmol·m^{-2}·s^{-1} were observed. Spectra acquired for the *phot1*

mutant resembled those of the wild-type, as this mutant differs from the wild-type only at

very low light intensities, not investigated in this study. The *phot2* mutant exhibits chloroplast

accumulation regardless of the light intensity used, thus the differential leaf reflectance curves

were similar for both used light intensities and resembled those of wild-type after blue light

treatment of 1.6 μmol·m^{-2}·s^{-1}. Curves that were representative of the *phot1phot2* mutant did

not show specific differences for leaves irradiated with 1.6 μmol·m^{-2}·s^{-1} and 120 μmol·m^{-2}·s^{-1}.

The *jac1* mutant, which is characterized by chloroplast avoidance at both investigated light

intensities, exhibited curves showing characteristic features for chloroplast avoidance. It is

worth noting that the dark positioning in the *jac1* mutant was distinct from the wild-type and
phot1, which differed from the *phot2* and *phot1phot2* mutants.

Performance of machine learning procedures for classification of leaves according to the
position of chloroplast movements

Our next goal was to automatically assess chloroplast positioning using classification
algorithms on the spectra extracted from hyperspectral images. Two types of neural networks
were considered: a multilayer perceptron and a convolutional neural network (Malek et al.,
2018). The multilayer perceptron architecture selected based on the validation set
performance consisted of three layers, with 21, 10, and 3 neurons. The structure of the
convolutional neural network, selected based on the cross-validation results for multiple trial
architectures, is shown in Fig. 3 A. The network consisted of four convolutional layers,
interspersed with max-pooling layers and ReLU activation functions. Feature maps generated
by the last convolutional layer were concatenated into a single array that acted as an input for
densely connected parts of the network, composed of four layers.

All three machine learning methods exhibited very high (91% - 100%) accuracy in
distinguishing between spectra recorded on leaves with the chloroplast avoidance positioning
and other types of chloroplast positioning (Fig. 3 B). High accuracy was observed both when
the trained and test sets came from the same species (*N. benthamiana*) and in the heterologous
setup, in which the spectra recorded for *A. thaliana* were used as the test set. The performance
of the tested procedures in distinguishing between spectra recorded on leaves exhibiting
accumulation and dark positioning was substantially worse than that observed for the
avoidance positioning. The best results were obtained using the convolutional network. For
the *N. benthamiana* test dataset, 70% of spectra recorded on dark-adapted leaf parts and 92%
of spectra recorded on leaves exhibiting the accumulation response were correctly classified.

Chloroplast Movement Index

To identify wavelengths that can be used to formulate a vegetation index that could be
used to distinguish leaves of different chloroplast arrangements, we employed the approach of
the correlation matrix, proposed in (Thenkabail et al., 2000). We used the formula of the
normalized difference type

$$ND(\lambda_1, \lambda_2) = \frac{R_{\lambda 1} - R_{\lambda 2}}{R_{\lambda 1} + R_{\lambda 2}}$$

where $R_{\lambda 1}$ and $R_{\lambda 2}$ are relative reflectance values in narrow bands centered at two wavelengths λ_1 and λ_2 in the range of 400 – 750 nm (Fig. 4). For every pair of λ_1 and λ_2, the values of the index ND(λ_1,λ_2) from raw spectra and its biserial correlation with the chloroplast positioning (binomial variable with two levels) was calculated. The plots were similar for both investigated species, *N. benthamiana* (Fig. 4 A) and *A. thaliana* (Fig. 4 B). We chose the region between 550 nm to 670 nm, in which correlation was high, and identified the local maxima. We additionally took into account that illumination-induced changes in leaf reflectance in the range of 500 nm to 550 nm are likely partly due to changes in chloroplast thylakoid pH gradient and the de-epoxidation of violaxanthin to zeaxanthin (Gamon et al., 1990), thus indices using wavelengths in this spectral region may be biased. Our final formula for the Chloroplast Movement Index, using wavelength marked with a cross in Fig. 4 A and Fig. 4 B, is:

$$CMI = \frac{R_{635} - R_{555}}{R_{635} + R_{555}}$$

We calculated the values of this index for *Nicotiana* (Fig. S4 A) and *Arabidopsis* (Fig. S4 B) spectra and analyzed its performance to distinguish chloroplast accumulation from avoidance. The index took negative values for both species in the range of -0.1 to -0.5. CMI proved to be very effective for *Nicotiana* leaves (Fig. 4 C) irradiated with low and high blue light. For *Arabidopsis* wild type and chloroplast movement mutants (Fig. 4 D), this index seemed to be able to distinguish between chloroplasts in the accumulation and avoidance positions depending on light irradiation regimes, but also the type of mutants. Wild-type plants, *phot1,* and *phot2* mutants in low light and the *phot2* mutant in high light showed chloroplast accumulation. The *phot1phot2* mutant exhibited differences not dependent on chloroplast movements. Wild type, *phot1* and *jac1* mutants in high light demonstrated chloroplast avoidance. We also performed pixel-wise calculations of our Chloroplast Movement Index for whole leaves (Fig.4. E, F, G). Venation and petioles gave high values to the index thus in practice pre-processing of data may be necessary to exclude such areas before analysis, using for example NDVI. In lower-resolution remote sensing spectral unmixing may potentially be used to exclude the contribution of light reflected from tissues other than mesophyll.

The impact of chloroplast movements on commonly used vegetation indices

We also calculated how chloroplast positioning affects vegetation indices calculated

for *Nicotiana benthamiana* leaves. We used a set of indices that are sensitive to pigments and

cell structure, as summarized in (Behmann et al., 2014), Supplementary Table S1. Chloroplast

positioning after high light treatment reduced the values of several indices, especially NDVI

(Fig. 5 A, Fig. S5 A), SR (Fig. 5 B, Fig. S5 B), RENDVI (Fig. 5 F, Fig. S5 F), mRENDEVI

(Fig. 5 G, Fig. S5 G), mRESR (Fig. 5 H, Fig. S5 H0, VOG1 (Fig. 5 I, Fig. S5 I), PRI

(Fig. 5 L, Fig. S5 L), SIPI (Fig. 5 M, Fig. S5 M), RGRI (Fig. 5 N, Fig. S5 N), PSRI (Fig. 5 O,

Fig. S5 O) CAR1 (Fig. 5 P, Fig. S5 P), and CAR2 (Fig. 5 Q, Fig. S5 Q). Chloroplast in the

avoidance position elevated SG (Fig. 5 E, Fig. S5 E), VOG2 (Fig. 5 J, Fig. S5 J), and VOG3

(Fig. 5 K, Fig. S5 K) values. Chloroplast accumulation affected vegetation indices to a lesser

extent, with diminished SG values and enhanced SR, RENDEVI, mRENDEVI, RGRI, CAR1,

and CAR2. EVI (Fig. 5 C, Fig. S5 C) and ARVI (Fig. 5 D, Fig. S5 D) were not dependent on

the position of chloroplasts.

In wild-type *Arabidopsis,* the direction of change (either elevation or reduction of the

index value) was consistent with the results obtained for *Nicotiana* for all investigated indices

apart from EVI and PRI (Fig. S5, Fig. S6, Fig. S7, Fig. 5, Fig. 6). The use of *Arabidopsis*

chloroplast movement mutants enabled us to determine if the observed changes in the

vegetative indices are related to differences in chloroplast arrangements. Values of several

indices decreased in leaves showing the avoidance positioning, including NDVI (Fig. 6 A,

Fig. S6 A), SR (Fig. 6 B, Fig. S6 B), RENDEVI (Fig. 6 F, Fig. S6 F), mRENDEVI (Fig. 6 G,

Fig. S6 G), mRESR (Fig. 6 H, Fig. S6 H), VOG1 (Fig. 6 I, Fig. S7 I), SIPI (Fig. 6 M,

Fig. S7 M), RGRI (Fig. 6 N, Fig. S7 N), PSRI (Fig. 6 O, Fig. S7 O), CAR1 (Fig. 6 P,

Fig. S7 P) and CAR2 (Fig. 6 Q, Fig. S7 Q). Such a decrease was not observed in mutants

deficient in chloroplast avoidance, namely *phot2* and *phot1phot2*. The change in the value of

the index mimicked the impact of low light of 1.6 $\mu mol \cdot m^{-2} \cdot s^{-1}$ in those mutants. For indices

in which chloroplast avoidance led to an increase in their value, such as SG (Fig. 6 E,

Fig. S6 E), VOG2 (Fig. 6 J, Fig. S7 J), and VOG3 (Fig. 6 K, Fig. S7 K), in both mutants,

phot2, and *phot1phot2* diminished values of those indices were observed, similarly to those

observed under low light. RENDEVI, mRENDEVI, and VOG1 seemed to be sensitive to

chloroplast accumulation. Changes in EVI (Fig. 6 C, Fig. S6 C), ARVI (Fig. 6 D, Fig. S6 D),

and PRI (Fig. 6 L, Fig. S7 L) did not correspond to chloroplast positioning in cells, as no

differences were observed between the *Arabidopsis* wild-type and chloroplast movement

mutants.

Discussion

Assessment of chloroplast movements through analysis of hyperspectral images

Here we propose a new method of chloroplast movement assessment based on broad-spectrum reflectance imaging and extraction of information about chloroplast positioning contained in the reflectance spectrum, using either vegetation indices or machine-learning procedures. This method appears to be more suited for high throughput detection of chloroplast movements than methods employing hemispherical reflectance (Davis et al., 2011), (Baránková et al., 2016), which require direct contact of the measuring equipment with the leaf. In the work of (Dutta et al., 2015), directional leaf reflectance of red light was used to visualize chloroplast movements in whole *Arabidopsis* rosettes. The use of white light as the measuring light allowed us to examine the effects of chloroplast positioning not only on the absolute reflectance but also on the shape of the spectrum and ratios between reflectance in selected bands, properties that are often more robust against artifacts or confounding factors (see e.g. (Gamon & Surfus, 1999) in the context of pigment content estimation). These issues are relevant for field conditions when sunlight is used as measuring light.

Averaged hyperspectral images of partly irradiated leaves show a clear difference in the magnitude of visible light reflectance between the irradiated and non-irradiated parts (Fig. 1 F). A work on maize suggested that an increase in PAR reflectance resulting from water stress might stem from the chloroplast avoidance movement (Zygielbaum et al., 2012). Irradiation with blue light affects the shape of the reflectance spectrum (Fig. 1 D, E). The measurements performed on *Arabidopsis* mutants with disrupted chloroplast movements indicate that the observed blue-light-induced changes in the spectrum are to a large extent caused by chloroplast relocations (Fig 2). This suggests that it might be possible to use the information contained in the reflectance spectrum to classify leaves according to the chloroplast positioning. We found that machine learning procedures, in particular the convolution network, exhibited good overall performance in the classification of reflectance spectra of leaves according to the chloroplast positioning (Fig. 3). The support vector machine and neural networks can reliably distinguish leaves exhibiting chloroplast avoidance positioning from the dark-adapted and low-light treated leaves, even when data used for training and testing come from different species (*N. benthamiana* and *A. thaliana*). This indicates that it may be possible to detect chloroplast avoidance even in spectra recorded on heterogeneous vegetation patches.

The Chloroplast Movement Index proposed in this work is the first effort to establish

vegetation indices relevant for monitoring chloroplast movements in field conditions (Fig. 4).

The differences between values of the index for dark-adapted and irradiated leaves for both

investigated species *Arabidopsis thaliana* and *Nicotiana benthamiana* are substantial. The

differences of smaller magnitude are observed between dark-adapted and low-light-irradiated

leaves. The index characterizes correctly chloroplast positions in *Arabidopsis* mutants with

disrupted chloroplast movements and positioning. This suggests that the index is sensitive to

chloroplast movements and not to other blue-light-dependent processes. As phototropin

mutations affect the leaf morphology and the mesophyll architecture (Kozuka et al., 2011),

good performance of the index for the *Arabidopsis* mutants suggests that it may be relatively

robust against changes in the leaf structure.

The impact of chloroplast movements on commonly used vegetation indices

Our results indicate that chloroplast movements may affect several vegetation indices

commonly used for remote sensing of physiological traits. To assess the practical significance

of this phenomenon, it is necessary to compare the observed changes caused by chloroplast

relocations with the magnitude of variation of the indices due to changes in the traits that are

usually monitored with them. Analysis of beach forests indicates that vegetation indices are

affected by the light conditions in which leaves develop. They generally perform well with

sun-lit leaves, but poorly with shaded ones (Sonobe & Wang, 2017). In this work, we show

that chlorophyll-dependent indices, characterized by simple formulae, such as Red Edge

NDVI (RENDVI), Modified Red Edge NDVI (mRENDVI), Modified Red Edge Simple Ratio

Index (mRESR), Simple Ratio Index (SR), as well as Sum Green Index (SG), are affected by

chloroplast positioning. The values for RENDEVI or mRENDVI fall in the range of 0 - 0.7

for various leaf types (Sims & Gamon, 2002), and thus may be significantly influenced by the

chloroplast relocations. This is especially prominent when differences between dark and lit

leaf halves of investigated *Arabidopsis* mutants are analyzed. mRESR values are between 0 -

4 for a variety of leaf types and species. Low values of mRESR, comparable to those

calculated in this study, are observed for leaves that are tough or pubescent (e.g. from

drought-deciduous species) and are characterized by high surface reflectance (Sims &

Gamon, 2002). The Red Green Ratio Index representing the anthocyanin chlorophyll ratio is

also substantially affected by chloroplast movements, with ΔRGRI values of about 0.1. RGRI

values measured for *Quercus agrifolia* depend on leaf age in the range of 0.5 - 1.5. SR values

mainly fall into the range of 0 - 20 as observed for paddy rice fields in Italy and 10 - 20 for

clonal *Populus* (Thenkabail et al., 2018). By contrast, the influence of chloroplast positioning is negligible for chlorophyll-dependent indices characterized by more complex formulae, such as the Enhanced Vegetation Index (EVI) and Atmospherically Resistant Vegetation Index (ARVI). The values of EVI, ARVI, SIPI, and PRI seem not to depend on chloroplast positioning, as changes in their values did not correlate in *Arabidopsis* phototropin mutants (Fig. 6). Structure Insensitive Pigment Index (SIPI) values around 1 are shared for a wide variety of plant species, except for evergreen species (Sims & Gamon, 2002). Photochemical Reflectance Index (PRI) values fall in the range of -0.1 to 0.1 except for winter deciduous and evergreen species (Sims & Gamon, 2002). The values of VOG1 and NDVI are very similar between groups both for *Nicotiana* and *Arabidopsis* wild-type and chloroplast movement mutants. The observed differences due to chloroplast positioning are relatively small, although statistically significant. However, NDVI values representing active vegetation are in the range of 0.7 - 0.9 with a calculated uncertainty value of about 0.02 (Borgogno-Mondino et al., 2016). The difference in NDVI between chloroplast dark and avoidance positions is about 0.05, which is significantly above the uncertainty levels for this index.

The accuracy of methods for pigment content estimation from vegetation reflectance spectra is usually lower for carotenoids than chlorophylls (Hill et al., 2019), both at the leaf (Gitelson et al., 2006) and canopy (Asner et al., 2015) levels. This lower accuracy reported for carotenoids is at least partly due to their lower content. However, carotenoid-dependent indices, CAR1 and CAR2, are substantially influenced by chloroplast relocations. For tree species (maple, chestnut, beech) ΔCAR2 of 1 corresponds to a difference in carotenoid content of 1.8 nmol/cm^2 (0.018 mmol/m^2). A similar, although nonlinear relationship between carotenoid content and ΔCAR1 is observed for maple leaves (Gitelson et al., 2002). For a survey investigating leaves from a wide range of species representing different functional types, carotenoid content fell in the range of 0 - 0.5 mmol/m^2 (Sims & Gamon, 2002). Plant Senescence Reflectance Index (PSRI), useful for monitoring stress and senescence, is also significantly influenced by chloroplasts in the avoidance position. The difference that stems from chloroplast positioning may affect PSRI readings, as ΔPSRI values are in the range of 0.02. Values between -0.05 and 0.05 are observed for several annual, deciduous, and evergreen species (Sims & Gamon, 2002).

Conclusions

We present a contactless method of chloroplast movement detection through hyperspectral imaging. It serves as a proof of concept for a high throughput, remote technique applicable to field conditions. We also propose a new vegetation index of a normalized difference type, useful for assessing the positions of chloroplasts and distinguishing between the accumulation and avoidance responses. Finally, we draw the attention of the community to the impact that chloroplast movement may exert on remote sensing studies based on leaf reflectance. Several commonly used vegetation indices might be affected by the chloroplast positioning in the mesophyll cells.

Acknowledgements

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest

The authors have no competing interests to declare.

Author contributions

P.H. - Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Software; Validation; Visualization; Writing – original draft, review & editing. J.L. - Conceptualization; Data curation, Validation; Methodology; Investigation; Resources; Supervision; Writing – original draft, review & editing.

Data availability

The data that supports the findings of this study are available in the supplementary material of this article. Raw data are available from the corresponding author upon reasonable request.

Supporting Information (brief legends)

Table S1: Formulae of Vegetation Indices calculated from the spectra obtained on *Arabidopsis thaliana* and *Nicotiana benthamiana*

Fig. S1. Leaf reflectance spectra recorded for dark-adapted and illuminated leaf halves of *N. benthamiana.* Detached leaves were irradiated with either (A) 1.6 or (B) 120 mmol m^{-2} s^{-1} of continuous blue light (455 nm) for 1 h, with half of the blade covered with aluminum foil.

Fig. S2. Leaf reflectance spectra recorded for dark-adapted and illuminated leaf halves of

A thaliana wild-type and chloroplast movement mutants. Detached leaves were irradiated

with either 1.6 mmol m^{-2} s^{-1} of continuous blue light (455 nm) for 1 h, with half of the blade

covered with aluminum foil.

Fig. S3. Leaf reflectance spectra recorded for dark-adapted and illuminated leaf halves of

A thaliana wild-type and chloroplast movement mutants. Detached leaves were irradiated

with either 120 mmol m^{-2} s^{-1} of continuous blue light (455 nm) for 1 h, with half of the blade

covered with aluminum foil.

Fig. S4. Chloroplast Movement Index calculated value for leaf halves of (A) *Nicotiana*

benthamiana or (B) *Arabidopsis thaliana* wild type and chloroplast movement mutants

(*phot1, phot2, phot1phot2, jac1*) either kept in darkness (grey) or irradiated (blue). The

irradiance to induce chloroplast accumulation was 1.6 µmol·m^{-2}·s^{-1} and to trigger chloroplast

avoidance was 120 µmol·m^{-2}·s^{-1}.

Fig. S5. Vegetation indices calculated for leaf halves of *Nicotiana benthamiana* either kept in

the dark (grey) or irradiated (blue). The formulae for the indices are collected in the

Supplementary Table S1. The irradiance to induce chloroplast accumulation was

1.6 µmol·m^{-2}·s^{-1} and to trigger chloroplast avoidance was 120 µmol·m^{-2}·s^{-1}. (A) NDVI, (B)

SR, (C) EVI, (D) ARVI, (E) SG, (F) RENDVI, (G) mRENDEVI, (H) mRESR, (I) VOG1, (J)

VOG2, (K) VOG3, (L) PRI, (M) SIPI, (N) RGRI, (O) PSRI, (P) CAR1, and (Q) CAR2.

Fig. S6. Vegetation indices calculated for leaf halves of *Arabidopsis thaliana* and chloroplast

movement mutants (*phot1, phot2, phot1phot2, jac1*) either kept in the dark (grey) or irradiated

(blue). The formulae for the indices are collected in the Supplementary Table S1. The

irradiance to induce chloroplast accumulation was 1.6 µmol·m^{-2}·s^{-1} and to trigger chloroplast

avoidance was 120 µmol·m^{-2}·s^{-1}. (A) NDVI, (B) SR, (C) EVI, (D) ARVI, (E) SG, (F)

RENDVI, (G) mRENDEVI, (H) mRESR.

Fig. S7. Vegetation indices calculated for leaf halves of *Arabidopsis thaliana* and chloroplast

movement mutants (*phot1, phot2, phot1phot2, jac1*) either kept in the dark (grey) or irradiated

(blue). The formulae for the indices are collected in the Supplementary Table S1. The

irradiance to induce chloroplast accumulation was 1.6 µmol·m^{-2}·s^{-1} and to trigger chloroplast

avoidance was 120 µmol·m^{-2}·s^{-1}. (I) VOG1, (J) VOG2, (K) VOG3, (L) PRI, (M) SIPI, (N)

RGRI, (O) PSRI, (P) CAR1, and (Q) CAR2.

References

Agati, G. (1998). Response of the in vivo chlorophyll fluorescence spectrum to environmental
564   factors and laser excitation wavelength. *Pure Appl. Opt*, *7*, 797–807.

565   Alzubaidi, L., Zhang, J., Humaidi, A. J., Al-Dujaili, A., Duan, Y., Al-Shamma, O.,
566   Santamaría, J., Fadhel, M. A., Al-Amidie, M., & Farhan, L. (2021). Review of deep
567   learning: concepts, CNN architectures, challenges, applications, future directions.
568   *Journal of Big Data*, *8*(1). https://doi.org/10.1186/s40537-021-00444-8

569   Annala, L., Honkavaara, E., Tuominen, S., & Pölönen, I. (2020). Chlorophyll concentration
570   retrieval by training convolutional neural network for stochastic model of leaf optical
571   properties (SLOP) inversion. *Remote Sensing*, *12*(2). https://doi.org/10.3390/rs12020283

572   Asner, G. P., Martin, R. E., Anderson, C. B., & Knapp, D. E. (2015). Quantifying forest
573   canopy traits: Imaging spectroscopy versus field survey. *Remote Sensing of*
574   *Environment*, *158*, 15–27. https://doi.org/10.1016/j.rse.2014.11.011

575   Baránková, B., Lazár, D., & Nauš, J. (2016). Analysis of the effect of chloroplast arrangement
576   on optical properties of green tobacco leaves. *Remote Sensing of Environment*, *174*, 181–
577   196. https://doi.org/10.1016/j.rse.2015.12.011

578   Behmann, J., Steinrücken, J., & Plümer, L. (2014). Detection of early plant stress responses in
579   hyperspectral images. *ISPRS Journal of Photogrammetry and Remote Sensing*, *93*, 98–
580   111. https://doi.org/10.1016/j.isprsjprs.2014.03.016

581   Berner J, Grohs P, Kutyniok G, & Petersen P. (2023). The Modern Mathematics of Deep
582   Learning. In P. Grohs & G. Kutyniok (Eds.), *Mathematical Aspects of Deep Learning*
583   (Vol. 1, pp. 1–82). Cambridge University Press.

584   Borgogno-Mondino, E., Lessio, A., & Gomarasca, M. A. (2016). A fast operative method for
585   NDVI uncertainty estimation and its role in vegetation analysis. *European Journal of*
586   *Remote Sensing*, *49*, 137–156. https://doi.org/10.5721/EuJRS20164908

587   Brugnoli, E., & Bjorkman, O. (1992). Chloroplast movements in leaves: Influence on
588   chlorophyll fluorescence and measurements of light-induced absorbance changes related
589   to ΔpH and zeaxanthin formation. *Photosynthesis Research*, *32*(1), 23–35.
590   https://doi.org/10.1007/BF00028795

Caterini, A. L., & Chang, D. E. (2018). *Deep Neural Networks in a Mathematical*
Framework.   Springer   Briefs   in   Computer   Science.
http://www.springer.com/series/10028

Chappelle, E. W., Wood, F. M., Newcomb, W. W., & Mcmurtrey, J. E. (1985). Laser-induced
fluorescence of green plants. 3: LIF spectral signatures of five major plant types. *Applied*
Optics, *24*(1), 74–80.

Chen, R., Yin, G., Liu, G., Li, J., & Verger, A. (2020). Evaluation and normalization of
topographic   effects   on   vegetation   indices.   *Remote   Sensing*,   *12*(14),   2290.
https://doi.org/10.3390/rs12142290

Christie, J. M. (2007). Phototropin Blue-Light Receptors. *Annual Review of Plant Biology*,
58(1), 21–45. https://doi.org/10.1146/annurev.arplant.58.032806.103951

Davis, P. A., Caylor, S., Whippo, C. W., & Hangarter, R. P. (2011). Changes in leaf optical
properties associated with light-dependent chloroplast movements. *Plant, Cell and*
Environment, *34*(12), 2047–2059. https://doi.org/10.1111/j.1365-3040.2011.02402.x

Davis, P. A., & Hangarter, R. P. (2012). Chloroplast movement provides photoprotection to
plants by redistributing PSII damage within leaves. *Photosynthesis Research*, *112*, 153–
161. https://doi.org/10.1007/s11120-012-9755-4

Dolinsky Margaret, & Hangarter, R. P. (2013). The Living Canvas: Interactive chloroplasts.
Leonardo, *46*(5), 488–489. https://doi.org/10.1162/Leon

Dutta, S., Cruz, J. A., Jiao, Y., Chen, J., Kramer, D. M., & Osteryoung, K. W. (2015). Non-
invasive, whole-plant imaging of chloroplast movement and chlorophyll fluorescence
reveals photosynthetic phenotypes independent of chloroplast photorelocation defects in
chloroplast   division   mutants.   *Plant   Journal*,   *84*(2),   428–442.
https://doi.org/10.1111/tpj.13009

Gabryś, H., Banaś, A. K., Hermanowicz, P., Krzeszowiec, W., Leśniewski, S., Łabuz, J., &
Sztatelman, O. (2017). Photometric Assays for Chloroplast Movement Responses to
Blue Light. *Bio-Protocol*, *7*(11), 1–11. https://doi.org/10.21769/BioProtoc.2310

Gamon, J. A., Field, C. B., Bilger, W., Bjiirkman, O., Fredeen, A. L., & Pefiuelas, J. (1990).
Remote sensing of the xanthophyll cycle and chlorophyll fluorescence in sunflower
leaves and canopies. *Oecologia*, *85*, 1–7.

Gamon, J. A., Penuelas, J., & Field, C. B. (1992). A Narrow-Waveband Spectral Index That

Tracks Diurnal Changes in Photosynthetic Efficiency*. *Remote Sensing of Environment*,

41, 35–44.

Gamon, J. A., & Surfus, J. S. (1999). Assessing leaf pigment content and activity with a

reflectometer. *New Phytologist*, *143*(1), 105–117. https://doi.org/10.1046/j.1469-

8137.1999.00424.x

Gitelson, A. A., Keydan, G. P., & Merzlyak, M. N. (2006). Three-band model for noninvasive

estimation of chlorophyll, carotenoids, and anthocyanin contents in higher plant leaves.

Geophysical Research Letters, *33*(11), L11402. https://doi.org/10.1029/2006GL026457

Gitelson, A. A., & Merzlyak, M. N. (1994). Spectral Reflectance Changes Associated with

Autumn Senescence of Aesculus hippocastanum L. and Acer platanoides L. Leaves.

Spectral Features and Relation to Chlorophyll Estimation. *Journal of Plant Physiology*,

143(3), 286–292. https://doi.org/10.1016/S0176-1617(11)81633-0

Gitelson, A. A., Zur, Y., Chivkunova, O. B., & Merzlyak, M. N. (2002). Assessing

Carotenoid Content in Plant Leaves with Reflectance Spectroscopy. *Photochemistry and*

Photobiology, *75*(3), 272. https://doi.org/10.1562/0031-

8655(2002)075<0272:accipl>2.0.co;2

Gorton, H. L., Williams, W. E., & Vogelmann, T. C. (1999). Chloroplast movement in

Alocasia macrorrhiza. *Physiologia Plantarum*, *106*, 421–428.

Gotoh, E., Suetsugu, N., Yamori, W., Ishishita, K., Kiyabu, R., Fukuda, M., Higa, T.,

Shirouchi, B., & Wada, M. (2018). Chloroplast accumulation response enhances leaf

photosynthesis and plant biomass production. *Plant Physiology*, *178*(3), 1358–1369.

https://doi.org/10.1104/pp.18.00484

Hao, W., Sun, J., Zhang, Z., Zhang, K., Qiu, F., & Xu, J. (2023). Novel Hybrid Model to

Estimate Leaf Carotenoids Using Multilayer Perceptron and PROSPECT Simulations.

Remote Sensing, *15*(20), 4997. https://doi.org/10.3390/rs15204997

Hermanowicz, P., Banaś, A. K., Sztatelman, O., Gabryś, H., & Łabuz, J. (2019). UV-B

Induces Chloroplast Movements in a Phototropin-Dependent Manner. *Frontiers in Plant*

Science, *10*. https://doi.org/10.3389/fpls.2019.01279

Hill, J., Buddenbaum, H., & Townsend, P. A. (2019). Imaging Spectroscopy of Forest
Ecosystems: Perspectives for the Use of Space-borne Hyperspectral Earth Observation
Systems. In *Surveys in Geophysics* (Vol. 40, Issue 3, pp. 553–588). Springer Science and
Business Media B.V. https://doi.org/10.1007/s10712-019-09514-2

Huete, A. R., Liu, H. Q., Batchily, K, & Van Leeuwen, W. (1997). A Comparison of
Vegetation Indices over a Global Set of TM Images for EOS-MODIS. *Remote Sensing of*
Environment, *5*, 440–451.

Idso, S. B., Pinter, P. J., Jackson, J. D., & Reginato, R. J. (1980). Estimation of Grain Yields
By Remote Sensing of Crop Senescence Rates. *Remote Sensing of Environment*, *9*, 87–
91.

Jarillo, J. A., Gabrys, H., Capel, J., Alonso, J. M., Ecker, J. R., & Cashmore, A. R. (2001).
Phototropin-related NPL1 controls chloroplast relocation induced by blue light. *Nature*,
410(6831), 952–954. https://doi.org/10.1038/35073622

Kagawa, T., Sakai, T., Suetsugu, N., Oikawa, K., Ishiguro, S., Kato, T., Tabata, S., Okada, K.,
& Wada, M. (2001). Arabidopsis NPL1: A Phototropin Homolog Controlling the
Chloroplast High-Light Avoidance Response. *Science*, *291*(5511), 2138–2141.
https://doi.org/10.1126/science.291.5511.2138

Kagawa, T., & Wada, M. (2000). Blue Light-Induced Chloroplast Relocation in Arabidopsis
thaliana as Analyzed by Microbeam Irradiation. *Plant and Cell Physiology*, *41*(1), 84–
93. https://doi.org/10.1093/pcp/41.1.84

Kasahara, M., Kagawa, T., Oikawa, K., Suetsugu, N., Miyao, M., & Wada, M. (2002).
Chloroplast avoidance movement reduces photodamage in plants. *Nature*,
420(December), 829–832. https://doi.org/10.1038/nature01202.1.

Kataoka, H. (2015). Gustav Senn (1875-1945): The pioneer of chloroplast movement
research. *Journal of Integrative Plant Biology*, *57*(1), 4–13.
https://doi.org/10.1111/jipb.12311

Kaufman, Y. J., & Tanre, D. (1995). Strategy for Direct and Indirect Methods Correcting the
Aerosol Effect on Remote Sensing: From AVHRR to EOS-MODIS. *Remote Sensing of*
Environment, *55*, 65–79.

Kozuka, T., Kong, S. G., Doi, M., Shimazaki, K. I., & Nagatani, A. (2011). Tissue-autonomous promotion of palisade cell development by phototropin 2 in Arabidopsis. *Plant Cell*, *23*(10), 3684–3695. https://doi.org/10.1105/tpc.111.085852

Łabuz, J., Hermanowicz, P., & Gabryś, H. (2015). The impact of temperature on blue light induced chloroplast movements in Arabidopsis thaliana. *Plant Science*, *239*, 238–249. https://doi.org/10.1016/j.plantsci.2015.07.013

Łabuz, J., Sztatelman, O., & Hermanowicz, P. (2022). Molecular insights into the phototropin control of chloroplast movements. *Journal of Experimental Botany*, 1–18. https://doi.org/10.1093/jxb/erac271

Lehmann, P., Nöthen, J., Schmidt Von Braun, S., Bohnsack, M. T., Mirus, O., & Schleiff, E. (2011). Transitions of gene expression induced by short-term blue light. *Plant Biology*, *13*(2), 349–361. https://doi.org/10.1111/j.1438-8677.2010.00377.x

Luesse, D. R., Deblasio, S. L., & Hangarter, R. P. (2010). Integration of phot1, phot2, and PhyB signalling in light-induced chloroplast movements. *Journal of Experimental Botany*, *61*(15), 4387–4397. https://doi.org/10.1093/jxb/erq242

Malek, S., Melgani, F., & Bazi, Y. (2018). One-dimensional convolutional neural networks for spectroscopic signal regression. *Journal of Chemometrics*, *32*(5). https://doi.org/10.1002/cem.2977

Mas, J. F., & Flores, J. J. (2008). The application of artificial neural networks to the analysis of remotely sensed data. In *International Journal of Remote Sensing* (Vol. 29, Issue 3, pp. 617–663). Taylor and Francis Ltd. https://doi.org/10.1080/01431160701352154

Merzlyak, M. N., Gitelson, A. A., Chivkunova, O. B., & Rakitin, V. Y. (1999). Non-destructive optical detection of pigment changes during leaf senescence and fruit ripening. *Physiologia Plantarum*, *106*, 135–141.

Moshou, D., Bravo, C., West, J., Wahlen, S., McCartney, A., & Ramon, H. (2004). Automatic detection of "yellow rust" in wheat using reflectance measurements and neural networks. *Computers and Electronics in Agriculture*, *44*(3), 173–188. https://doi.org/10.1016/j.compag.2004.04.003

Pal, S. K., & Mitra, S. (1992). Multilayer Perceptron, Fuzzy Sets, and Classification. *IEEE TRANSACTIONS ON NEURAL NETWORKS*, *3*(5), 683.

Park, Y., Chow, W. S., & Anderson, J. M. (1996). Chloroplast Movement in the Shade Plant
Tradescantia albiflora Helps Protect Photosystem II against Light Stress. *Plant*
Physiology, *111*, 867–875.

Penuelas, J., Frederic, B., & Filella, I. (1995). Semi-Empirical Indices to Assess
Carotenoids/Chlorophyll-a Ratio from Leaf Spectral Reflectance. *Photosynthetica*, *31*(2),
221–230. https://www.researchgate.net/publication/229084513

Pfündel, E. E., Latouche, G., Meister, A., & Cerovic, Z. G. (2018). Linking chloroplast
relocation to different responses of photosynthesis to blue and red radiation in low and
high light-acclimated leaves of Arabidopsis thaliana (L.). *Photosynthesis Research*,
137(1), 105–128. https://doi.org/10.1007/s11120-018-0482-3

Pullanagari, R. R., Dehghan-Shoar, M., Yule, I. J., & Bhatia, N. (2021). Field spectroscopy of
canopy nitrogen concentration in temperate grasslands using a convolutional neural
network. *Remote Sensing of Environment*, *257*, 112353.
https://doi.org/10.1016/j.rse.2021.112353

Rouse, J. W., Haas, R. H., Schell, J. A., & Deering, D. W. (1974). Monitoring vegetation
systems in the Great Plains with ERTS. *NASA Special Publication*, *351*, 309–317.
https://doi.org/10.1021/jf60203a024

Sakai, T., Kagawa, T., Kasahara, M., Swartz, T. E., Christie, J. M., Briggs, W. R., Wada, M.,
& Okada, K. (2001). Arabidopsis nph1 and npl1: Blue light receptors that mediate both
phototropism and chloroplast relocation. *Proceedings of the National Academy of*
Sciences, *98*(12), 6969–6974. https://doi.org/10.1073/pnas.101137598

Senn, G. (1908). *Die Gestalts- und Lageveränderung der Pflanzen-Chromatophoren: mit*
einer Beilage: Die Lichtbrechung der lebenden Pflanzenzelle. W. Engelmann, W.

Shi, S., Xu, L., Gong, W., Chen, B., Chen, B., Qu, F., Tang, X., Sun, J., & Yang, J. (2022). A
convolution neural network for forest leaf chlorophyll and carotenoid estimation using
hyperspectral reflectance. *International Journal of Applied Earth Observation and*
Geoinformation, *108*. https://doi.org/10.1016/j.jag.2022.102719

Shibata, K. (1958). Spectrophotometry of intact biological materials. *The Journal of*
Biochemistry, *45*(8), 599–623. https://sci-hub.tw/10.1002/iroh.19620470121

Sims, D. A., & Gamon, J. A. (2002). Relationships between leaf pigment content and spectral
reflectance across a wide range of species, leaf structures and developmental stages.
Remote Sensing of Environment, *81*, 337–354. www.elsevier.com/locate/rse

Song, G., & Wang, Q. (2021). Including leaf traits improves a deep neural network model for
predicting photosynthetic capacity from reflectance. *Remote Sensing*, *13*(21).
https://doi.org/10.3390/rs13214467

Sonobe, R., & Wang, Q. (2017). Hyperspectral indices for quantifying leaf chlorophyll
concentrations performed differently with different leaf types in deciduous forests.
Ecological Informatics, *37*, 1–9. https://doi.org/10.1016/j.ecoinf.2016.11.007

Suetsugu, N., Kagawa, T., & Wada, M. (2005). An Auxilin-Like J-Domain Protein, JAC1,
Regulates Phototropin-Mediated Chloroplast Movement. *Plant Physiology*,
139(September), 151–162. https://doi.org/10.1104/pp.105.067371.

Suetsugu, N., Mittmann, F., Wagner, G., Hughes, J., & Wada, M. (2005). A chimeric
photoreceptor gene, NEOCHROME, has arisen twice during plant evolution.
Proceedings of the National Academy of Sciences, *102*(38), 13705–13709.
https://doi.org/10.1073/pnas.0504734102

Sztatelman, O., Waloszek, A., Banaś, A. K., & Gabryś, H. (2010). Photoprotective function of
chloroplast avoidance movement: In vivo chlorophyll fluorescence study. *Journal of
Plant Physiology*, *167*(9), 709–716. https://doi.org/10.1016/j.jplph.2009.12.015

Thenkabail, P. S., Lyon, J. G., & Huete, A. (2018). *Hyperspectral Remote Sensing of
Vegetation, Hyperspectral Indices and Image Classifications for Agriculture and
Vegetation* (P. S. Thenkabail, J. G. Lyon, & A. Huete, Eds.; Second Edition). CRC Press
Taylor & Francis.

Thenkabail, P. S., Smith, R. B., & De Pauw, E. (2000). Hyperspectral Vegetation Indices and
Their Relationships with Agricultural Crop Characteristics. *Remote Sensing of the
Environment*, *71*, 158–182.

Thenkabail, P. S., Smith, R. B., & De Pauw, E. (2002). Evaluation of Narrowband and
Broadband Vegetation Indices for Determining Optimal Hyperspectral Wavebands for
Agricultural Crop Characterization. *Photogrammetric Engineering & Remote Sensing*,
68(6), 607–621.

Tucker, C. J. (1977). Spectral Estimation of Grass Canopy Variables. *Remote Sensing of*
Environment, *6*, 1.

Vogelmann, J. E., Rock, B. N., & Moss, D. M. (1993). Red edge spectral measurements from
sugar maple leaves. *International Journal of Remote Sensing*, *14*(8), 1563–1575.
https://doi.org/10.1080/01431169308953986

Walczak, T., & Gabrys, H. (1980). New type of photometer for measurements of transmission
changes corresponding to chloroplast movements in leaves. *Photosynthetica*, *14*, 65–72.

Wang, R., Springer, K. R., & Gamon, J. A. (2023). Confounding effects of snow cover on
remotely sensed vegetation indices of evergreen and deciduous trees: An experimental
study. *Global Change Biology*, *29*(21), 6120–6138. https://doi.org/10.1111/gcb.16916

Wilson, S., & Ruban, A. v. (2020). Rethinking the influence of chloroplast movements on
non-photochemical quenching and photoprotection. *Plant Physiology*, *183*(3), 1213–
1223. https://doi.org/10.1104/pp.20.00549

Yorinao, I., & Kazuo, S. (1974). Comparative examination of terrestrial plant leaves in terms
of light-induced absorption changes due to chloroplast rearrangements. *Plant & Cell*
Physiology, *15*, 717–721.

Zhang, C., Li, C., He, M., Cai, Z., Feng, Z., Qi, H., & Zhou, L. (2023). Leaf water content
determination of oilseed rape using near-infrared hyperspectral imaging with deep
learning regression methods. *Infrared Physics and Technology*, *134*.
https://doi.org/10.1016/j.infrared.2023.104921

Zygielbaum, A. I., Arkebauer, T. J., Walter-Shea, E. A., & Scoby, D. L. (2012). Detection and
measurement of vegetation photoprotection stress response using PAR reflectance. *Israel*
Journal of Plant Sciences, *60*(1–2), 37–47. https://doi.org/10.1560/IJPS.60.1-2.37

Figure captions

Fig. 1. Chloroplast movements in *Nicotiana benthamiana* leaves assessed with (A, B)
transmittance measurements, (C) microscopic observations, and (D, E, F, G) leaf reflectance
measurements. (A, B) Changes in leaf transmittance at 660 nm, induced by blue light of (A)
1.6 or (B) 120 μmol m^{-2}·s^{-1}. Each curve represents the mean of 10 measurements. The
standard deviation is shown in light grey, the standard error is in dark grey. A decrease in leaf

transmittance corresponds to chloroplast accumulation and an increase - to chloroplast

avoidance. (C) Chloroplast arrangements in palisade cells of *N. benthamiana*. Leaves

detached from 6-week-old dark-adapted plants were either kept in darkness (mock irradiation)

or irradiated for 1 h with blue light of 1.6 or 120 μmol·m^{-2}·s^{-1}. Chloroplast arrangements were

imaged with a laser-scanning confocal microscope, using chlorophyll autofluorescence.

Maximum intensity projections were calculated from Z-stacks, recorded for ca. 90 μm,

starting from the leaf's upper surface. (D, E, F, G) Effect of blue light irradiation on the

reflectance spectrum of *N. benthamiana* leaves. Detached leaves were irradiated with either

1.6 or 120 mmol m^{-2} s^{-1} of continuous blue light (455 nm) for 1 h, with half of the blade

covered with aluminum foil. (D, E) The mean difference in reflectance spectra between

irradiated and darkened leaf halves calculated from hyperspectral images. The sample size

was 23 for low and 31 leaves for high light irradiation. The standard deviation is shown in

light grey, the standard error is in dark grey. (F, G) Reflectance images of tobacco leaves,

calculated as an average of the hyperspectral images in the visible (400 – 700 nm) range.

Fig. 2. Blue-light-induced chloroplast movements, examined with (A) the optical microscopy

and (B) leaf reflectance spectra in *Arabidopsis thaliana* rosette leaves of WT, *phot1*, *phot2*,

phot1phot2, and *jac1* mutant plants. (A) Chloroplast arrangements in palisade cells of

detached leaves from 4-week-old dark-adapted plants were either kept in darkness (mock

irradiation) or irradiated for 1 h with blue light of 1.6 or 120 μmol·m^{-2}·s^{-1}. Chloroplast

arrangements were imaged with a laser-scanning confocal microscope, using chlorophyll

autofluorescence. Maximum intensity projections were calculated from Z-stacks, recorded for

ca. 90 μm, starting from the leaf's upper surface. (B) Effect of blue light irradiation on the

reflectance spectrum of the adaxial surface of *Arabidopsis* wild-type and phototropin mutant

leaves. Detached leaves were irradiated with either 1.6 or 120 μmol m^{-2} s^{-1} blue light for 1 h,

with half of the blade covered with aluminum foil. The mean difference in reflectance spectra

between irradiated and darkened halves was calculated from hyperspectral images. The

sample size is 30 leaves for each combination of the plant line and irradiation conditions. The

standard deviation is shown in light grey, the standard error is in dark grey.

Fig. 3. (A) The architecture of the convolutional neuron network for classification of leaf

reflectance spectra in the range 400 – 750 nm according to the chloroplast arrangement. The

network consists of the initial convolutional part, four densely connected layers, and a

softmax layer. The convolutional part consisted of four blocks, each with a 1-D convolution

layer of variable kernel size and count, followed by a rectified linear unit (ReLU) and a max

pooling layer, with a kernel size of 5 and stride of 2. Each of the first three densely connected layers was followed by a batch normalization (BN) and a ReLU layer. The final softmax layer yields estimated probabilities of the input falling into each chloroplast arrangement class. (B) The confusion matrices showing the mean performance of the support vector machine with a linear kernel, multilayer perceptron, and convolutional network, trained on spectra recorded on *Nicotiana benthamiana* leaves and used to classify spectra recorded for the same species (upper row) or *Arabidopsis thaliana* (bottom row). Training was performed 20 times, and the standard deviation is shown.

Fig. 4. Classification of chloroplast positioning using a vegetation index of a normalized difference type. (A - B) Contour plots showing the biserial correlation between chloroplast positioning and the narrow band vegetation index of the general form specified in the plot, calculated from the leaf reflectance spectra for every combination of wavelengths λ_1 (x-axis) and λ_2 (y-axis) in the range of 400 nm to 750 nm. Spectra recorded for (A) *Nicotiana benthamiana* and (B) *Arabidopsis thaliana* leaves were used. Chloroplast accumulation was induced with 1.6 µmol·m^{-2}·s^{-1}, and avoidance with 120 µmol·m^{-2}·s^{-1} of blue light. Blue and red areas correspond to indices that assume high values for leaves exhibiting accumulation and avoidance response, respectively. The pair of wavelengths (635 and 555 nm), used to calculate the ND index for chloroplast movements (CMI), positively correlated with the avoidance response, and was marked with a cross. (C) The difference in the Chloroplast Movement Index between the dark-adapted and either low- or high-light irradiated *N. benthamiana* leaf halves. Each dot corresponds to one leaf. Statistical analysis was performed with the Welch *t*-test. (D) The difference in Chloroplast Movement Index between the dark-adapted and either low- or high-light-irradiated leaf halves of *Arabidopsis thaliana* and chloroplast movement mutants (WT- green, *phot1* - blue, *phot2* – pink, *phot1phot2* - purple, *jac1* - yellow). Boxes that do not share any letter represent groups for which the means of transformed values differ at the 0.05 level (Tukey's test, adjusted for multiple comparisons). (E, F, G) The Chloroplast Movement Index calculated for each pixel of hyperspectral images of (E) *Nicotiana benthamiana,* (F) *Arabidopsis thaliana* wild-type, and (G) the *phot2* mutant. In each leaf, one half was darkened, while the other was irradiated with either 1.6 µmol·m^{-2}·s^{-1} or 120 µmol·m^{-2}·s^{-1} of blue light.

Fig. 5. The difference between values of vegetation indices calculated from the spectra from darkened and irradiated parts of the *Nicotiana benthamiana* leaves. Each dot corresponds to one leaf. Irradiation with low blue light of 1.6 µmol·m^{-2}·s^{-1} was used to induce chloroplast

accumulation (dark blue boxes), with high blue light of 120 µmol·m^{-2}·s^{-1} to trigger the

avoidance response (light blue boxes). The index formulae are in Supplementary Table 1. The

statistical significance of differences between means was examined using the Welch t-test.

Fig. 6. The difference between values of vegetation indices calculated from the spectra from

dark-adapted and irradiated parts of *Arabidopsis thaliana* wild-type and chloroplast

movement mutants (WT - green, *phot1* - blue, *phot2* - pink, *phot1phot2* - purple, *jac1* -

yellow). Each dot corresponds to one leaf. The irradiance to induce chloroplast accumulation

was 1.6 µmol·m^{-2}·s^{-1} and to trigger chloroplast avoidance was 120 µmol·m^{-2}·s^{-1} of blue light.

The index formulae are in Supplementary Table 1. Boxes that do not share any letter represent

groups for which the means of transformed values differ at the 0.05 level (Tukey's test,

adjusted for multiple comparisons).

A) 1.6 µmol·m⁻²·s⁻¹ BL

B) 120 µmol·m⁻²·s⁻¹ BL

C) Darkened | 1.6 µmol·m⁻²·s⁻¹ | 120 µmol·m⁻²·s⁻¹

50 µm

D) 1.6 µmol·m⁻²·s⁻¹ BL

E) 120 µmol·m⁻²·s⁻¹ BL

F) 1.6 µmol·m⁻²·s⁻¹ | Darkened

G) 120 µmol·m⁻²·s⁻¹ | Darkened

VIS Reflectance

0   0.1   0.2   0.3

Nicotiana benthamiana

A) $ND(\lambda_1,\lambda_2) = \dfrac{R_{\lambda_1} - R_{\lambda_2}}{R_{\lambda_1} + R_{\lambda_2}}$

Arabidopsis thaliana

B) $ND(\lambda_1,\lambda_2) = \dfrac{R_{\lambda_1} - R_{\lambda_2}}{R_{\lambda_1} + R_{\lambda_2}}$

C) ($\mu mol \cdot m^{-2} \cdot s^{-1}$)

1.6   120

$\Delta ND\ (635\ nm, 555\ nm)$

***

D)

1.6 $\mu mol \cdot m^{-2} \cdot s^{-1}$

WT   *phot1*   *phot2*   *phot1 phot2*   *jac1*

120 $\mu mol \cdot m^{-2} \cdot s^{-1}$

WT   *phot1*   *phot2*   *phot1 phot2*   *jac1*

$\Delta ND\ (635\ nm, 555\ nm)$

a   abc   a   bc   c   d   e   a   ab   de

E) 1.6 $\mu mol \cdot m^{-2} \cdot s^{-1}$ Darkened   120 $\mu mol \cdot m^{-2} \cdot s^{-1}$ Darkened

ND (635 nm, 555 nm)

-0.4   -0.2   0.0   0.1

F) $\mu mol \cdot m^{-2} \cdot s^{-1}$

1.6   Darkened   120   Darkened

G) 1.6   Darkened   120   Darkened