1

3   Effects of early life adversity and adolescent basolateral amygdala activity on

4   corticolimbic connectivity and anxiety behaviors

10   Caitlyn R. Cody (MS) [1], Emilce Artur de la Villarmois (PhD) [2], Anabel Miguelez

11   Fernandez (PhD) [2], Janelle Lardizabal [1], Chaney McKnight [1], Kuei Tseng (PhD) [2]

12   Heather C. Brenhouse (PhD) [1]

14   1. Psychology Department, Northeastern University, Boston MA 02115

15   2. Department of Anatomy and Cell Biology, University of Illinois, Chicago IL 60612

Corresponding Author: Heather Brenhouse
360 Huntington Ave
125 Nightingale Hall
Boston MA 02115
Phone: 617-549-4549
Fax: 617.373.8714
h.brenhouse@northeastern.edu
29

Abstract

Early postnatal development of corticolimbic circuitry is shaped by the environment and

is vulnerable to early life challenges. Prior work has shown that early life adversity (ELA)

leads to hyperinnervation of glutamatergic basolateral amygdala (BLA) projections to the

prefrontal cortex (PFC) in adolescence. While hyperinnervation is associated with later-

life anxiety behaviors, the physiological changes underpinning corticolimbic and

behavioral impacts of ELA are not understood. We tested whether postsynaptic BLA-

driven PFC activity is enhanced in ELA-exposed animals, using the maternal separation

(MS) model of ELA. PFC local-field potential following BLA stimulation was facilitated in

MS-exposed adolescents. Since ELA increases activity of the early-developing BLA,

while the PFC exhibits protracted development, we further examined impacts of

glutamatergic BLA activity during early adolescence on later-life PFC innervation and

heightened anxiety. In early adolescence, MS-exposed animals exhibited decreased

anxiety-like behavior, and acute adolescent BLA inhibition induced behaviors that

resembled those of MS animals. To examine long-lasting impacts of adolescent BLA

activity on innervation, BLA-originating axonal boutons in the PFC were quantified in late

adolescence after early adolescent BLA inhibition. We further tested whether late

adolescent BLA-PFC changes were associated with anxious reactivity expressed as

heightened acoustic startle responses. MS rearing increased BLA-PFC innervation and

threat reactivity in late adolescence, however early adolescent BLA inhibition was

insufficient to prevent MS effects, suggesting that earlier BLA activity or post-synaptic

receptor rearrangement in the PFC drives altered innervation. Taken together, these

findings highlight both pre- and postsynaptic changes in the adolescent BLA-PFC circuit

following ELA.

1. Introduction

Early postnatal life is a dynamic period of brain development that involves a

coordinated symphony of circuit maturation. As many regions within the brain continue to

develop and mature postnatally, external stimuli impact this development and regulate

the degree of connectivity to other brain areas. Since subcortical limbic regions undergo

substantial synapse formation and pruning during early development while higher-order

areas such as the prefrontal cortex (PFC) mature throughout adolescence,[1–5] early

postnatal life and adolescence are two sensitive periods when limbic and prefrontal

regions form functional connections in response to external stimuli and neuronal activity.

Thus, activity within limbic regions over the course of development can affect the strength

and quantity of efferent connections to cortical areas.

The basolateral amygdala (BLA) has unique integration with aversion- and reward-

related circuitry and serves as the primary input nucleus of the amygdala.[6] Monosynaptic

BLA connectivity with the PFC[7–10] increases throughout adolescence in healthy rats,

facilitating the integration of emotional information with cognitive appraisals of the

environment.[11] Challenges to the early amygdala that alter local activity may therefore

induce activity changes within efferent projections, which may result in changes to

behavior when the circuit is recruited during adulthood.

Adolescence is a dynamic period when synapses undergo significant refinement

in the frontal cortex.[12,13] These reported changes are thought to involve glutamatergic

afferents with local remodeling of inhibitory-excitatory synaptic balance,[14,15] and coincide

with heightened emotional regulation.[16–18] Importantly, neuronal activity during sensitive

periods directly impacts pre- and post-synaptic plasticity within the PFC.[19] The present

study therefore investigated the impacts of early life environmental inputs and adolescent

BLA activity on the development of BLA-PFC circuitry and corresponding behaviors.

Theoretical frameworks regarding environmental influences on development posit

that postnatal brain circuitry has evolved to "expect" an appropriate quantity and quality

of caregiving stimuli in order to develop typically.[20] Disruptions to typical development can

be detrimental to the postnatal brain and can exert cascading effects on subsequent

maturation.[21–24] One such form of disruption includes early life adversity (ELA), where the

species-expectant environment is violated.[25,26] ELA can lead to several affective

consequences such as anxiety and major depressive disorder, which often emerge in

adolescence, and continue through adulthood.[13,27–34] The maternal separation (MS)

model of ELA in rodents induces corticolimbic and anxiety-related changes that resemble

those reported in ELA-exposed humans. Particularly, MS induces hyperinnervation of the

PFC from glutamatergic BLA projections,[18] as well as enhanced anxiety-like behavior as

measured by the acoustic startle response (ASR).[35] These effects were seen in

adolescent rodents as early as postnatal day (p)40 in male rats, and p30 in females, which

corresponds to early-mid adolescence. Therefore, we hypothesized that ELA leads to

pre- and post-synaptic changes to BLA-PFC connectivity via hyperactivity within the

developing BLA, and that preventing heightened presynaptic BLA activity during early-

mid adolescence would interrupt hyperinnervation and result in reduced anxiety

behaviors.

We first determined whether MS exposure led to facilitation of BLA-evoked local-field potentials (LFP) within the PFC in early-mid adolescence. Next, to test the role that BLA activity plays in the developing innervation to the PFC, BLA activity was manipulated via an inhibitory DREADD to reduce excitatory activity during early-mid adolescence. In an exploratory analysis, an open field test was performed following acute DREADD inhibition to test whether early adolescent BLA activity regulated anxiety-like behaviors differentially after MS exposure. Histology was performed in late adolescence to determine if earlier BLA inhibition prevented the establishment of long-range glutamatergic PFC hyperinnervation. Animals were also tested in late adolescence for enhanced acoustic startle response (ASR) at baseline or following an anxiety-provoking social threat cue (playback of 22-kHz ultrasonic vocalizations), to determine whether prevention of BLA-PFC hyperinnervation was associated with prevention of enhanced ASR.

2. Materials and Methods

All experimental procedures were conducted in accordance with the 1996 Guide for the Care and Use of Laboratory Animals (NIH) with approval from the Institutional Animal Care and Use Committee at Northeastern University and the University of Illinois Chicago.

2.1 Subjects and Maternal Separation (MS)

All experimental subjects were bred in house (details in Supplemental Materials). Whole litters were randomly assigned to control (CON) or maternal separation (MS) rearing conditions. All separations lasted 4 hours each day beginning at 0900h with pups kept in

a separate room from the dam. Upon completion of MS pups were reunited with their

dams and littermates and left undisturbed until the next day. On p21, pups were weaned

from the dam into standard rat cages with an age, sex, and condition matched partner.

2.2 Local Field Potential (LFP) recordings of prefrontal response to basolateral amygdala

(BLA) stimulation

Early-mid adolescent (p30-45) rats were anesthetized and a concentric bipolar electrode

was used to record within the prelimbic region while stimulating the BLA through a

computer-controlled pulse generator (details in Supplemental Materials). Changes in the

input-output curve of PFC responses were estimated by comparing the amplitude of BLA-

evoked LFP at different stimulating intensities (0, 0.25, 0.5, 0.75, and 1.0 mA) using single

(300µS duration) square pulses delivered every 15s in both control and MS rats.

2.3 Viral Injections and Chemogenetic Inhibition

In a separate cohort of male and female rats, on p26 subjects underwent a stereotaxic

surgical injection of the inhibitory DREADD AAV-CAMKIIa-HM4DGi-mCherry (Addgene,

Watertown, MA) to the BLA (A/P: -1.9mm, L/M: ±4.5mm, D/V: -6.75mm). Viral injections

are detailed in Supplemental Materials. Rats were allowed to recover for 1 week before

behavioral testing. Daily from p33-39, rats were weighed and given an intraperitoneal (i.p)

injection of the DREADD agonist clozapine-N-oxide dihydrochloride (CNO, 2mg/kg, Bio-

Techne Tocris, Minneapolis, MN) or vehicle saline (SAL). CNO was prepared fresh prior

to the start of injections and was dissolved in sterile saline. Functional validation methods

are detailed in Supplemental Materials.

2.4 Behavioral Assays

For each behavioral assay, rats were brought into the behavioral suite and allowed to

habituate for 15 minutes.

2.4.1 Open Field Test (OFT)

For early adolescent testing (p33), 30 minutes after the first CNO or SAL injection, rats

were placed into the open field (100 cm x 100 cm black acrylic arena). Video monitoring

of behavior was done via an overhead CCTV camera (Panasonic WV-CP500, Secaucus,

NJ), and behavioral scoring was done via Noldus Ethovision 9.0 Behavioral Tracking

software. The central zone of the arena was defined as the inner 30x30 cm portion of the

arena (Figure 4A).

2.4.2 Acoustic Startle Response (ASR)

On p54, rats were placed into a sound attenuated startle chamber (MedAssociates Inc,

St. Albens, VT) and played a series of 30 105dB white noise bursts for an assessment of

baseline startle. Twenty-four hours later, rats underwent a second ASR test, during which

ASR was measured following exposure to a five-minute 22 kHz rat ultrasonic vocalization

(USV) recording. For more details see Supplemental Materials.

2.5 Microscopy and Analysis

2.5.1 Viral Expression

Histological procedures are detailed in Supplemental Materials. In order to quantify the

amount of DREADD expression within and outside of the BLA, slices were imaged on a

Zeiss Imager M.2 microscope for bolus verification. Slices with dual hemispheric

expression in the BLA underwent volumetric analysis via the Cavalieri probe within

StereoInvestigator. Sections containing DREADD expression were analyzed at 2.5X

magnification to determine the extent of bolus infection within and spread outside of the

BLA. Average BLA volume was previously measured in a subset of animals, and the

volume of DREADD expression was quantified as a percent of the total BLA volume.

Animals with a total bolus volume that filled <50% of the BLA were excluded from

analysis.

2.5.2 Axonal Innervation

The corresponding PFC tissue of all included animals was processed using

immunohistochemistry (IHC) to identify mCherry expression. Sections containing the

infralimbic (IL) and preimbic (PL) regions were prepared for immunohistochemistry as

detailed in Falcy et al. 2020[36] (details in Supplemental Materials). Z-stacks were analyzed

for axonal bouton count and intensity.

2.6 Statistical Analysis

GraphPad Prism 10.0 Software was used for all data analyses and graphical

representation. Sample sizes were determined by power analyses using effect sizes from

prior experiments. Sex x rearing x treatment 3-way ANOVAs were conducted (with

frequency as repeated measures in BLA-evoked PFC LFP analyses), and when main

effects of sex or interactions with sex were observed, 2-way ANOVAs with factors of

treatment and rearing were conducted for each sex with Post-hoc Tukey's multiple

comparison's tests when appropriate. For simplicity, full statistical details are only

highlighted for significant comparisons in the Results section. For all statistical details,

see Supplemental Materials. Of note, three animals were removed from ASR analysis

due to data file corruption.

3. Results

3.1 MS leads to facilitation of BLA-evoked local field potential in the early adult PFC.

3-way ANOVA revealed a main effect of rearing ($F_{1,20}$ = 102.0; p < 0.0001; partial η^2 =

0.732) and a rearing x intensity interaction ($F_{4,80}$ = 38.23; p < 0.0001; partial η^2 = 0.563)

on PFC local field potential (Figure 1A-C; n=6/group), with MS exposure facilitating BLA-

evoked activity in both males (Figure 1B) and females (Figure 1C). These findings

indicate that exposure to MS results in increased BLA-evoked activity within the PFC in

early-mid adolescence.

3.2 BLA inhibition during mid-adolescence does not prevent increased BLA-PFC

connectivity after MS

Figure 2A summarizes the DREADD strategy used to inhibit the BLA. Ex-vivo recordings

of BLA pyramidal neurons show the effect of the inhibitory DREADD at 8-9 days post viral

injection (Figure 2B-D).

A subset of rats (n=6-8) were sacrificed following behavioral testing (p56-60) to

quantify BLA innervation to the PFC via axonal bouton count and fluorescent intensity.

Three-way ANOVA revealed a sex x rearing interaction with a moderate effect size on

bouton count in the PL ($F_{1,57}$ = 3.245; p=0.0770; partial η^2 = 0.054) and the IL ($F_{1,57}$ =

3.617; p=0.0623; partial η^2 = 0.06). Due to the presence of a moderate effect size for the

sex x rearing interaction, each sex was analyzed separately, with two-way ANOVAs

assessing effects of rearing condition and treatment (CNO or SAL). In males, MS

increased PL bouton count ($F_{1,25}$ = 4.661; p=0.0407; partial η^2 = 0.157; Figure 3E) and IL

bouton count ($F_{1,25}$ = 5.098; p=0.0329; partial η^2 = 0.169; Figure 3I). However, no effects

were observed in females.

Three-way ANOVA also revealed a main effect of rearing on bouton intensity in the

PL ($F_{1,57}$ = 10.87; p=0.0017; partial η^2 = 0.160) and IL ($F_{1,57}$ = 15.58; p=0.0002; partial η^2

= 0.215), as well as a sex x rearing interaction in bouton intensity within the PL ($F_{1,57}$ =

15.00; p=0.0003; partial η^2 = 0.208) and IL ($F_{1,57}$ = 13.19; p=0.0006; partial η^2 = 0.160).

Due to the presence of a sex x rearing interaction, each sex was analyzed separately,

revealing another male-specific main effect of rearing on PL bouton intensity ($F_{1,25}$ =

36.09; p=<0.0001; partial η^2 = 0.591, Figure 3G), and IL bouton intensity ($F_{1,25}$ = 29.53;

p=<0.0001; partial η^2 = 0.541; Figure 3K).

Treatment with CNO had no effects on either measure (count or fluorescent

intensity). Results of all statistical comparisons are summarized in Supplementary Tables

5-7. These findings indicate that MS rearing induces hyperinnervation of the PFC from

the BLA in late adolescent male animals only, and that BLA inhibition during early-mid

adolescence is insufficient to reverse these changes.

3.3 Exposure to MS and acute BLA inhibition alters adolescent behavior in a sex

dependent manner

Rats with inhibitory DREADD expression in the BLA were given an i.p injection of

CNO (2 mg/kg) or SAL 30 minutes before the open field test on p33. Animals (n=10-12

per condition) were assessed for time spent, latency to enter, and number of entries into

the center zone of the arena (Figure 4D-K).

A three-way ANOVA revealed a moderate effect size sex x treatment interaction

($F_{1,79}$ = 3.061; p=0.081; partial η^2 = 0.037) on distance moved. Two-way ANOVAs within

each sex assessed effects of rearing condition and treatment (CNO or SAL). No effects

were seen in male animals (Figure 4D). In female animals, a main effect of treatment

($F_{1,41}$= 4.526; p=0.0394; partial η^2 = 0.099) was found indicating that females treated with

CNO had increased locomotion (Figure 4E).

Three-way ANOVAs revealed main effects of sex ($F_{1,79}$ = 7.019; p=0.0098; partial

η^2 = 0.084) and treatment ($F_{1,79}$ = 7.148; p=0.0092; partial η^2 = 0.085) for the latency to

enter the center. Analyses also revealed a rearing x treatment interaction ($F_{1,79}$ = 5.098;

p=0.0268; partial η^2 = 0.062). Two-way ANOVAs assessed effects within each sex of

rearing condition and treatment. In male animals, a main effect of treatment ($F_{1,38}$ = 4.475;

p=0.0414; partial η^2 = 0.11) and a rearing x treatment interaction ($F_{1,38}$ = 4.235; p=0.0469;

partial η^2 = 0.105) were found for the latency to enter the center (Figure 4F). Post-hoc

analyses revealed differences between control SAL and control CNO treated animals

(p=0.0129) as well as control SAL and MS SAL animals (p=0.0258) indicating that control

animals treated with CNO as well as MS animals treated with SAL exhibited a decrease

in the latency to enter the center compared to control SAL animals. In female animals, a

significant main effect of treatment ($F_{1,41}$ = 4.185; p=0.0472; partial η^2 = 0.093) was found

in the latency to enter the center (Figure 4G), with CNO treated animals showing a

decreased latency to enter the center.

Three-way ANOVAs revealed a rearing x treatment interaction for center duration

($F_{1,79}$ = 4.912; p=0.0296; partial η^2 = 0.059). Two-way ANOVAs further assessed effects

of rearing condition and treatment within each sex. No effects were seen in male animals

(Figure 4H). In female animals, analyses revealed a main effect of rearing ($F_{1,41}$ = 4.392;

p=0.0427; partial η^2 = 0.101) and a rearing x treatment interaction ($F_{1,41}$ = 4.412;

p=0.0422; partial η^2 = 0.101) on center duration (Figure 4I). Post-hoc analyses revealed

differences between control and MS SAL treated females (p=0.0045) such that MS

animals spent more time in the center compared to controls, and that this was not

mediated by CNO treatment.

A three-way ANOVA revealed a main effect of rearing ($F_{1,79}$ = 5.353; p=0.0234;

partial η^2 = 0.065) and a rearing x treatment interaction ($F_{1,79}$ = 6.176; p=0.0151; partial

η^2 = 0.074) for center frequency. Two-way ANOVAs within each sex assessed effects of

rearing condition and treatment. In male animals, a moderate effect sized rearing x

treatment interaction was observed ($F_{1,38}$ = 3.561; p=0.0677; partial η^2 = 0.087; Figure

4J). Female animals displayed a significant main effect of rearing ($F_{1,41}$ = 11.24;

p=0.0018; partial η^2 = 0.219; Figure 4K), such that MS females entered the center more

than control-reared females, mirroring the increased duration. These findings indicate that

MS rearing results in decreased anxiety-like behavior in the OFT, and that BLA inhibition

in early adolescence results in an MS-like phenotype in control animals, while having no

effect on MS animals.

3.4 Early adolescent BLA inhibition does not prevent ASR changes in MS-exposed late

adolescence

On p54 (late adolescence), rats were tested for baseline startle levels to 30 white

noise bursts. Latency to startle, peak startle time, and peak startle value were assessed

with three-way ANOVAs. Previously established differences between males and females

were observed and are likely related to animal size; main effects of sex were observed

on baseline latency to startle ($F_{1,76}$ = 7.821; p=0.0065; partial η^2= 0.093), and baseline

peak value ($F_{1,76}$ = 5.598; p=0.0205; partial η^2= 0.069). No effects of rearing or treatment

were observed on baseline ASR in late adolescence (Figure 5A-C). On p55, rats were

presented with a five-minute playback of a 22 kHz USV recording and then immediately

tested for startle levels to 30 white noise bursts. Latency to startle, peak startle time, and

peak startle value were assessed with three-way ANOVAs. Main effects of rearing were

found for latency to startle ($F_{1,76}$ = 4.855; p=0.0306; partial η^2= 0.060; Figure 5D), and

peak time ($F_{1,76}$ = 5.210; p=0.0252; partial η^2 = 0.064; Figure 5E), with MS-exposed

animals startling more quickly than controls following USV playback. Main effects of sex

were found for latency to startle on Day 2 ($F_{1,76}$ = 4.082; p=0.047; partial η^2= 0.051) as

well as peak value on Day 2 ($F_{1,76}$ = 10.54; p=0.0017; partial η^2= 0.122). No effects of

treatment were observed on USV mediated ASR. Rats were also evaluated for the

percent change from baseline following USV presentation. A main effect of rearing was

again found in latency to startle ($F_{1,76}$ = 4.081; p=0.0469; partial η^2 = 0.051; Figure 5G)

and peak time ($F_{1,76}$ = 5.195; p=0.0255; partial η^2 = 0.064; Figure 5H), indicating that MS

animals startled quicker than control animals following USV playback. A main effect of

rearing was also observed for percent change in peak value ($F_{1,76}$ = 4.856; p=0.0306;

partial η ^{2}= 0.060) No main effects of treatment with CNO were found in any analyses,

and no interactions of rearing, sex, or treatment were present, therefore further analyses

were not conducted. These findings indicate that BLA inhibition via CNO in early-mid

adolescence had no overall effect on ASR in late adolescence.

4. Discussion

The current study identifies dissociable pre- and post-synaptic contributions to sex-

and age-specific effects of ELA on corticolimbic connectivity and behavior. We sought to

investigate if an adverse rearing environment altered BLA-PFC connectivity via aberrantly

high adolescent BLA activity, and if these changes drove anxiety-like behavioral

responses in early and late adolescence. We observed that MS rearing yielded

postsynaptic facilitation of PFC responsivity to BLA activation by mid-adolescence. We

hypothesized that presynaptic inhibition of the BLA during this period would prevent MS-

driven hyperinnervation from the BLA to the PFC, as well as MS-driven changes to

anxiety-like behaviors.

As previously observed by quantifying BLA axon terminals,[18] here, MS resulted in

hyperinnervation of the PFC as assessed by a more functionally relevant measure,

namely the number of axonal boutons. However, this increase was not attenuated by BLA

inhibition from p33-p39, suggesting that sustained adolescent activity of the BLA does not

regulate the development of presynaptic connectivity with the PFC. Additionally, no

increases in innervation were seen in females, indicating that by late adolescence, any

previously observed increases in BLA-PFC connectivity may have pruned back to control

levels. Both male and female rats exhibit significant pruning within the PFC between

adolescence and adulthood, however females display greater losses than males.[37] The

age at which peak synaptic pruning of the PFC occurs in males and females remains

unknown,[38] which allows for the possibility that inhibition of BLA activity in early-mid

adolescence had differential effects on BLA-PFC innervation across sexes. It is therefore

possible that the period of accelerated BLA-PFC innervation induced by MS rearing

occurred at an earlier timepoint in females, and thus by the time histology was performed

(p56), both control and MS females had reached adult innervation levels. Additionally,

previous work from our group showed increases in female BLA-PFC innervation that were

already apparent by p28; therefore, if BLA activity does provoke innervation to the PFC

in females, inhibition from p33-39 may have been too late to affect any MS-evoked

changes. Thus, it will be important to repeat this study with an earlier intervention time

point as well as earlier ages of assessment to capture any sex-specific effects of juvenile

BLA inhibition on adolescent innervation.

Since inhibiting presynaptic activity was not sufficient to reverse long-range

hyperinnervation, it is possible that postsynaptic plasticity within the PFC precedes and

drives these connectivity changes. For example, MS results in heightened expression of

GluN2A-containing NMDA receptors within the PL[39] as well as reduced GluA2-positive

AMPA receptors.[40] While the PFC displays late-maturing connectivity with regions such

as the BLA,[11,18] the postnatal environment is capable of directly impacting PFC activity

and receptivity,[41,42] and the nature of glutamate receptivity can influence development of

inputs.[43] Importantly, post synaptic NMDAR increases induced by MS have been shown

to correlate with anxiety-like behaviors,[44] and thus in concert with our findings here

showing a lack of structural changes following presynaptic inhibition, we posit that

postsynaptic mechanisms may be more critical for ELA induced outcomes in the PFC.

We observed anxiety-like ASR in late adolescence such that MS animals

responded to anxiogenic USV playback with elevated threat responsivity. Contrastingly,

MS-exposed animals tested in early adolescence exhibited faster center entry (males) or

increased center time (females), suggesting a decrease in anxiety-like behavior. Our

findings are largely consistent with other assessments of MS-reared animals in early

adolescence,[45–47] where by late adolescence, MS-reared animals display reduced center

time.[48] Importantly, the open field task has also been used as a measure of risk-seeking,[49]

and ELA is known to increase risk-seeking and impulsivity in adolescence.[50–59] It is

possible therefore that the anxiolytic behaviors seen during the OFT in early adolescence

may be attributable to increases in risk-seeking behaviors, and that this behavioral

phenotype shifts as animals continue to mature.   Notably, acute BLA inhibition in

adolescent controls resulted in open field behavior that mirrored MS-exposed animals,

such that control animals treated with CNO had shorter latencies to enter the center of

the open field arena. Further work will tease apart the discrete mechanisms linking MS to

risk-seeking, and how developing BLA activity differentially drives these behaviors during

adolescence.

MS does not appear to alter later-life ASR via hyperactivity of the developing BLA,

as early-mid adolescent BLA inhibition did not prevent MS-induced changes. The

amygdala regulates anxiety-related enhancement of ASR reflex, with the BLA particularly

sensitive to ambiguous or distal threats from the environment.[60,61] Here we tested the

impacts of MS and early BLA manipulation on potentiation of ASR by a distal social threat,

namely playback of 22kHz USV.[57] Previous work has shown enhanced BLA activity

following presentation of a 22 kHz USV,[62] indicating a role of the BLA in the processing

of this fearful stimuli. Control-reared animals display a potentiated ASR following USV

playback, while MS alters this potentiated response.[35] Here we observed that while MS-

exposed animals exhibited increased potentiation of the ASR, early-mid adolescent BLA

inhibition was not sufficient to rescue the heightened startle response induced by MS.

These findings should be considered with the limitation that surgery and chronic

injections, which are stressful experiences in and of themselves, were conducted during

juvenility and early adolescence. This may be considered a second hit of stress later

following MS exposure, which can alter typical stress responses.[63] We are also unable to

causally link altered BLA-PFC innervation or functional connectivity with ASR. Taken

together, we observed that postnatal ELA impacts development of both pre- and post-

synaptic BLA-PFC connectivity as well as anxiety-related behaviors across adolescence.

However, the insufficiency of pre-synaptic BLA inhibition to alter later BLA-PFC

connectivity or anxiety-related behavior highlights the importance of investigating early

post-synaptic changes in the PFC that may be driving MS-induced corticolimbic functional

connectivity changes.

382
383
384
385

Acknowledgments
The authors would like to thank the Institute for Chemical Imaging of Living Systems at Northeastern University for consultation and imaging support.

Author Contributions
Heather C. Brenhouse and Kuei Y. Tseng and designed the study. Caitlyn R. Cody performed and analyzed the behavioral and axonal innervation data under the supervision of Heather C. Brenhouse. Janelle Lardizabal and Chaney McKnight assisted Caitlyn R. Cody with behavioral experiments, histology, and data collection. Emilce Artur de la Villarmois and Anabel Miguelez Fernandez performed all electrophysical experiments and data analysis under the supervision of Kuei Y. Tseng. Caitlyn R. Cody, Heather C. Brenhouse, and Kuei Y. Tseng wrote the manuscript. Caitlyn R. Cody, Emilce Artur de la Villarmois, and Kuei Y. Tseng prepared the figures.

Funding
Supported by NIH Grant R01MH127850-01 (KYT and HCB)

Competing Interests
The authors have nothing to disclose

References

1. Drzewiecki, C. M., Willing, J. & Juraska, J. M. Synaptic number changes in the medial prefrontal cortex across adolescence in male and female rats: A role for pubertal onset. *Synap. N. Y. N* **70**, 361–368 (2016).

2. Huttenlocher, P. R. Synapse elimination and plasticity in developing human cerebral cortex. *Am. J. Ment. Defic.* **88**, 488–496 (1984).

3. Huttenlocher, P. R. Morphometric study of human cerebral cortex development. *Neuropsychologia* **28**, 517–527 (1990).

4. Huttenlocher, P. R. & Dabholkar, A. S. Regional differences in synaptogenesis in human cerebral cortex. *J. Comp. Neurol.* **387**, 167–178 (1997).

5. Konrad, K., Firk, C. & Uhlhaas, P. J. Brain development during adolescence: neuroscientific insights into this developmental period. *Dtsch. Arzteblatt Int.* **110**, 425–431 (2013).

6. Janak, P. H. & Tye, K. M. From circuits to behaviour in the amygdala. *Nature* **517**, 284–292 (2015).

7. Bacon, S. J., Headlam, A. J., Gabbott, P. L. & Smith, A. D. Amygdala input to medial prefrontal cortex (mPFC) in the rat: a light and electron microscope study. *Brain Res.* **720**, 211–219 (1996).

8. Krettek, J. E. & Price, J. L. Projections from the amygdaloid complex to the cerebral cortex and thalamus in the rat and cat. *J. Comp. Neurol.* **172**, 687–722 (1977).

9. Mcdonald, A. J. Glutamate and aspartate immunoreactive neurons of the rat basolateral amygdala: colocalization of excitatory amino acids and projections to the limbic circuit. *J. Comp. Neurol.* **365**, 367–379 (1996).

10. Verwer, R. W., Van Vulpen, E. H. & Van Uum, J. F. Postnatal development of

amygdaloid projections to the prefrontal cortex in the rat studied with retrograde and

anterograde tracers. *J. Comp. Neurol.* **376**, 75–96 (1996).

11. Cunningham, M. G., Bhattacharyya, S. & Benes, F. M. Amygdalo-cortical sprouting

continues into early adulthood: implications for the development of normal and

abnormal function during adolescence. *J. Comp. Neurol.* **453**, 116–130 (2002).

12. Andersen, S. L., Thompson, A. T., Rutstein, M., Hostetter, J. C. & Teicher, M. H.

Dopamine receptor pruning in prefrontal cortex during the periadolescent period in

rats. *Synap. N. Y. N* **37**, 167–169 (2000).

13. Andersen, S. L. & Teicher, M. H. Stress, sensitive periods and maturational events in

adolescent depression. *Trends Neurosci.* **31**, 183–191 (2008).

14. Caballero, A., Granberg, R. & Tseng, K. Y. Mechanisms contributing to prefrontal

cortex maturation during adolescence. *Neurosci. Biobehav. Rev.* **70**, 4–12 (2016).

15. Caballero, A., Orozco, A. & Tseng, K. Y. Developmental regulation of excitatory-

inhibitory synaptic balance in the prefrontal cortex during adolescence. *Semin. Cell*

Dev. Biol. **118**, 60–63 (2021).

16. Gee, D. G. *et al.* Early developmental emergence of human amygdala-prefrontal

connectivity after maternal deprivation. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 15638–

15643 (2013).

17. Swartz, J. R., Carrasco, M., Wiggins, J. L., Thomason, M. E. & Monk, C. S. Age-

related changes in the structure and function of prefrontal cortex-amygdala circuitry

in children and adolescents: a multi-modal imaging approach. *NeuroImage* **86**, 212–

220 (2014).

18. Honeycutt, J. A. *et al.* Altered corticolimbic connectivity reveals sex-specific

adolescent outcomes in a rat model of early life adversity. *eLife* **9**, e52651 (2020).

19. Guadagno, A., Belliveau, C., Mechawar, N. & Walker, C.-D. Effects of Early Life

Stress on the Developing Basolateral Amygdala-Prefrontal Cortex Circuit: The

Emerging Role of Local Inhibition and Perineuronal Nets. *Front. Hum. Neurosci.* **15**,

(2021).

20. Zeanah, C. H., Gunnar, M. R., McCall, R. B., Kreppner, J. M. & Fox, N. A. Sensitive

Periods. *Monogr. Soc. Res. Child Dev.* **76**, 147–162 (2011).

21. Knudsen, E. I. Sensitive periods in the development of the brain and behavior. *J.

Cogn. Neurosci.* **16**, 1412–1425 (2004).

22. Hensch, T. K. Critical period regulation. *Annu. Rev. Neurosci.* **27**, 549–579 (2004).

23. Tseng, K. Y., Chambers, R. A. & Lipska, B. K. The neonatal ventral hippocampal

lesion as a heuristic neurodevelopmental model of schizophrenia. *Behav. Brain Res.*

204, 295–305 (2009).

24. Tottenham, N. & Gabard-Durnam, L. J. The developing amygdala: a student of the

world and a teacher of the cortex. *Curr. Opin. Psychol.* **17**, 55–60 (2017).

25. Nelson, C. A. & Gabard-Durnam, L. J. Early Adversity and Critical Periods:

Neurodevelopmental Consequences of Violating the Expectable Environment.

Trends Neurosci. **43**, 133–143 (2020).

26. Lupien, S. J., McEwen, B. S., Gunnar, M. R. & Heim, C. Effects of stress throughout

the lifespan on the brain, behaviour and cognition. *Nat. Rev. Neurosci.* **10**, 434–445

(2009).

27. Kessler, R. C., Avenevoli, S. & Ries Merikangas, K. Mood disorders in children and adolescents: an epidemiologic perspective. *Biol. Psychiatry* **49**, 1002–1014 (2001).

28. Davey, C. G., Yücel, M. & Allen, N. B. The emergence of depression in adolescence: development of the prefrontal cortex and the representation of reward. *Neurosci. Biobehav. Rev.* **32**, 1–19 (2008).

29. Smyke, A. T. *et al.* The caregiving context in institution-reared and family-reared infants and toddlers in Romania. *J. Child Psychol. Psychiatry* **48**, 210–218 (2007).

30. McEwen, B. S. Understanding the potency of stressful early life experiences on brain and body function. *Metabolism.* **57**, S11–S15 (2008).

31. Maccari, S., Krugers, H. J., Morley-Fletcher, S., Szyf, M. & Brunton, P. J. The consequences of early-life adversity: neurobiological, behavioural and epigenetic adaptations. *J. Neuroendocrinol.* **26**, 707–723 (2014).

32. Callaghan, B. L. & Tottenham, N. The Stress Acceleration Hypothesis: Effects of early-life adversity on emotion circuits and behavior. *Curr. Opin. Behav. Sci.* **7**, 76–81 (2016).

33. Hane, A. A. & Fox, N. A. Early Caregiving and Human Biobehavioral Development: A Comparative Physiology Approach. *Curr. Opin. Behav. Sci.* **7**, 82–90 (2016).

34. Krugers, H. J. *et al.* Early life adversity: Lasting consequences for emotional learning. *Neurobiol. Stress* **6**, 14–21 (2016).

35. Granata, L., Parakoyi, A. & Brenhouse, H. C. Age- and sex-specific effects of maternal separation on the acoustic startle reflex in rats: early baseline enhancement in females and blunted response to ambiguous threat. *Front. Behav. Neurosci.* **16**, (2022).

36. Falcy, B. A., Mohr, M. A. & Micevych, P. E. Immunohistochemical amplification of

mCherry fusion protein is necessary for proper visualization. *MethodsX* **7**, 100946

(2020).

37. Koss, W. A., Belden, C. E., Hristov, A. D. & Juraska, J. M. Dendritic remodeling in

the adolescent medial prefrontal cortex and the basolateral amygdala of male and

female rats. *Synap. N. Y. N* **68**, 61–72 (2014).

38. Juraska, J. M. The last stage of development: The restructuring and plasticity of the

cortex during adolescence especially at puberty. *Dev. Psychobiol.* **66**, e22468

(2024).

39. Wieck, A., Andersen, S. L. & Brenhouse, H. C. Evidence for a neuroinflammatory

mechanism in delayed effects of early life adversity in rats: Relationship to cortical

NMDA receptor expression. *Brain. Behav. Immun.* **28**, 218–226 (2013).

40. Ganguly, P., Honeycutt, J. A., Rowe, J. R., Demaestri, C. & Brenhouse, H. C. Effects

of early life stress on cocaine conditioning and AMPA receptor composition are sex-

specific and driven by TNF. *Brain. Behav. Immun.* **78**, 41–51 (2019).

41. Stern, J. A., Kelsey, C. M., Yancey, H. & Grossmann, T. Love on the developing

brain: Maternal sensitivity and infants' neural responses to emotion in the

dorsolateral prefrontal cortex. *Dev. Sci.* e13497 (2024) doi:10.1111/desc.13497.

42. Ayala-Rodríguez, J. D. & García-Colunga, J. Maternal separation modifies

spontaneous synaptic activity in the infralimbic cortex of stress-resilient male rats.

PloS One **18**, e0294151 (2023).

43. Kelsch, W. *et al.* GluN2B-containing NMDA receptors promote glutamate synapse

development in hippocampal interneurons. *J. Neurosci. Off. J. Soc. Neurosci.* **34**,

16022–16030 (2014).

44. Ganguly, P., Holland, F. H. & Brenhouse, H. C. Functional Uncoupling NMDAR

NR2A Subunit from PSD-95 in the Prefrontal Cortex: Effects on Behavioral

Dysfunction and Parvalbumin Loss after Early-Life Stress.

Neuropsychopharmacology **40**, 2666–2675 (2015).

45. Wang, Q., Li, M., Du, W., Shao, F. & Wang, W. The different effects of maternal

separation on spatial learning and reversal learning in rats. *Behav. Brain Res.* **280**,

16–23 (2015).

46. Kundakovic, M., Lim, S., Gudsnuk, K. & Champagne, F. A. Sex-specific and strain-

dependent effects of early life adversity on behavioral and epigenetic outcomes.

Front. Psychiatry **4**, 78 (2013).

47. Wang, X. *et al.* Maternal separation affects anxiety-like behavior beginning in

adolescence and continuing through adulthood and related to Dnmt3a expression. *J.*

Neurophysiol. **128**, 611–618 (2022).

48. Jin, S. *et al.* Anxiety-like behaviour assessments of adolescent rats after repeated

maternal separation during early life. *Neuroreport* **29**, 643–649 (2018).

49. Palm, S., Momeni, S., Lundberg, S., Nylander, I. & Roman, E. Risk-assessment and

risk-taking behavior predict potassium- and amphetamine-induced dopamine

response in the dorsal striatum of rats. *Front. Behav. Neurosci.* **8**, 236 (2014).

50. Rincón-Cortés, M. Sex differences in addiction-relevant behavioral outcomes in

rodents following early life stress. *Addict. Neurosci.* **6**, 100067 (2023).

51. Kim, S. T. *et al.* Multidimensional impulsivity as a mediator of early life stress and

alcohol dependence. *Sci. Rep.* **8**, 4104 (2018).

52. Duckworth, A., Kim, B. & Tsukayama, E. Life Stress Impairs Self-Control in Early

Adolescence. *Front. Psychol.* **3**, (2013).

53. Hostinar, C. E., Stellern, S. A., Schaefer, C., Carlson, S. M. & Gunnar, M. R.

Associations between early life adversity and executive function in children adopted

internationally from orphanages. *Proc. Natl. Acad. Sci. U. S. A.* **109 Suppl 2**, 17208–

17212 (2012).

54. Lovallo, W. R. Early life adversity reduces stress reactivity and enhances impulsive

behavior: implications for health behaviors. *Int. J. Psychophysiol. Off. J. Int. Organ.*

Psychophysiol. **90**, 8–16 (2013).

55. Oshri, A. *et al.* Impulsivity as a mechanism linking child abuse and neglect with

substance use in adolescence and adulthood. *Dev. Psychopathol.* **30**, 417–435

(2018).

56. Lovic, V., Keen, D., Fletcher, P. J. & Fleming, A. S. Early-life maternal separation and

social isolation produce an increase in impulsive action but not impulsive choice.

Behav. Neurosci. **125**, 481–491 (2011).

57. Roman, E., Gustafsson, L., Berg, M. & Nylander, I. Behavioral profiles and stress-

induced corticosteroid secretion in male Wistar rats subjected to short and

prolonged periods of maternal separation. *Horm. Behav.* **50**, 736–747 (2006).

58. Marmendal, M., Eriksson, C. J. P. & Fahlke, C. Early deprivation increases

exploration and locomotion in adult male Wistar offspring. *Pharmacol. Biochem.*

Behav. **85**, 535–544 (2006).

59. Lundberg, S., Abelson, K. S. P., Nylander, I. & Roman, E. Few long-term

consequences after prolonged maternal separation in female Wistar rats. *PLOS*

ONE **12**, e0190042 (2017).

60. Koch, M. & Schnitzler, H. U. The acoustic startle response in rats--circuits mediating

evocation, inhibition and potentiation. *Behav. Brain Res.* **89**, 35–49 (1997).

61. Davis, M., Falls, W. A., Campeau, S. & Kim, M. Fear-potentiated startle: A neural

and pharmacological analysis. *Behav. Brain Res.* **58**, 175–198 (1993).

62. Demaestri, C., Brenhouse, H. C. & Honeycutt, J. A. 22 kHz and 55 kHz ultrasonic

vocalizations differentially influence neural and behavioral outcomes: Implications for

modeling anxiety via auditory stimuli in the rat. *Behav. Brain Res.* **360**, 134–145

(2019).

63. Kuhlman, K. R. Pitfalls and potential: Translating the two-hit model of early life stress

from pre-clinical non-human experiments to human samples. *Brain Behav. Immun. -*

Health **35**, 100711 (2024).

  Figure Legends

Figure 1. Effects of ELA on BLA-evoked PFC LFP. (**A**) Illustration of BLA stimulation
and PFC recording sites. (B) Prelimbic PFC response to BLA stimulation at increasing
intensities is enhanced by p30 as shown by local field potential (LFP) recordings in
males (****p<0.0001, **p<0.01 difference between control and MS animals, after
correction for multiple comparisons; N=6/group). (**C**) The prelimbic PFC LFP response
to BLA stimulation is also enhanced by p30 in females (****p<0.0001, **p<0.01
difference between control and MS animals, after correction for multiple comparisons;
N=6/group).

Figure 2. DREADD validation. (**A**) Timeline of DREADD delivery and ex-vivo recordings
of BLA pyramidal neurons. (**B**) Bath application of CNO (10μM, 10 min) markedly
hyperpolarized the resting membrane potential of all pyramidal neuron tested (**p<0.01,
unpaired t-test, t=3.114, df=14). (**C**) Accordingly, a significant increase in the rheobase
(minimal somatic current (pA) to elicit an action potential) was observed all neurons
following bath application of CNO (**p<0.01, unpaired t-test, t=4.124, df=14). (**D**)
Example traces of a BLA pyramidal neuron showing the inhibitory effect of CNO as
revealed by the reduced current evoked action potential and hyperpolarized resting
membrane potential.
641
Figure 3. Effects of ELA with or without BLA inactivation during early-mid adolescence
on late adolescent PFC innervation. (**A**) Timeline of DREADD experiment. (**B**)
Representative DREADD injections in the BLA (**C-D**) BLA-PFC terminals in the PFC. (**E**)
Male PL bouton count. (**F**) Female PL bouton count. (**G**) Male PL bouton intensity. (**H**)
Female PL bouton intensity. (**I**) Male IL bouton count. (**J**) Female IL bouton count. (**K**)
Male IL bouton intensity. (**K**) Female IL bouton intensity. Means ± SEM with individual
data points are shown. (*p < 0.05 ****p<0.0001 rearing effect). n=5-12/group.
649
Figure 4. Effects of ELA with or without acute BLA inactivation on open field behavior
during early adolescence. (**A**) Schematic of open field arena. (**B-C**) Representative
heat maps depicting control and MS animal movement within the open field arena. OFT
results for male (**D**) Distance Moved (**H**) Center Duration (**F**) Center latency to enter (**J**)
Center frequency and female (**E**) Distance Moved (**I**) Center Duration (**G**) Center latency
to enter (**K**) Center frequency. Means ± SEM with individual data points are shown. (*p
< 0.05; **p<0.005; dashed line denotes main effect of rearing). n=10-13/group.
657
658
Figure 5. Effects of ELA with or without BLA inactivation during early-mid adolescence
on late adolescent acoustic startle. (**A**) Day 1 latency to startle. (**B**) Day 1 latency to
peak startle value. (**C**) Day 1 peak startle value. (**D**) Day 2 latency to startle. (**E**) Day 2
latency to peak startle value. (**F**) Day 2 peak startle value. (**G**) Percent change latency
to startle. (**H**) Percent change latency to peak startle value. (**C**) Percent change peak
startle value. Means ± SEM with individual data points are shown. (*p < 0.05 rearing
effect). n=9-12/group.

A.

postnatal days
25   33-34

BLA
CaMKIIa
hM4D-Gi

BLA
recordings

B.

RMP (mV)

**

aCSF   +CNO

C.

rheobase (pA)

**

aCSF   +CNO

D.

20 mV
0.2 s

aCSF

+CNO

−68 mV   −73 mV

A. Control or MS Rearing — p2-p20 | DREADD Surgery — p26 | Open Field Test — p33-p34 | CNO Injections — p33-p39 | ASR — p54-p55 | Placement and Innervation — p60+

B. Representative BLA DREADD Injection — 500 μm

C. Representative BLA Terminals in PFC — ACC, PL, IL, VLO, AI, LO, AId, GI, DI — 1000 um

D. Representtive BLA-PFC Boutons — 10 um

Prelimbic Cortex

E. ♂ PL Count — Count — * — CON, MS

F. ♀ PL Count — Count — CON, MS

G. ♂ PL Intensity — Intensity — **** — CON, MS

H. ♀ PL Intensity — Intensity — CON, MS

Infralimbic Cortex

I. ♂ IL Count — Count — * — CON, MS

J. ♀ IL Count — Count — CON, MS

K. ♂ IL Intensity — Intensity — **** — CON, MS

L. ♀ IL Intensity — Intensity — CON, MS

● CON SAL ● CON CNO ○ MS SAL ○ MS CNO

A. Open Field **B.** Control **C.** MS

D. ♂ Distance Moved **E.** ♀ Distance Moved **F.** ♂ Center Latency to Enter **G.** ♀ Center Latency to Enter

H. ♂ Center Duration **I.** ♀ Center Duration **J.** ♂ Center Frequency **K.** ♀ Center Frequency

● CON SAL ● CON CNO ○ MS SAL ○ MS CNO

Day 1-Baseline Startle Response

A. Latency to Startle

B. Latency to Peak

C. Peak Value

Day 2- USV Induced Startle Response

D. Latency to Startle

E. Latency to Peak

F. Peak Value

Percent Change- Day 1 to Day 2

G. % Change Latency to Startle

H. % Change Latency to Peak

I. % Change Peak Value

CON SAL   CON CNO   MS SAL   MS CNO