Contributions of mirror-image hair cell orientation to mouse otolith organ and zebrafish neuromast function

Kazuya Ono[#1], Amandine Jarysta[#2], Natasha Hughes[#3], Alma Jukic[#4], Vanessa Chang[3], Michael R. Deans[5,6], Ruth Anne Eatock[*1], Kathleen Cullen[*3,7,8,9], Katie Kindt[*4] and Basile Tarchini[*2,10]

[1] Department of Neurobiology, University of Chicago, Chicago, IL, 60637, USA.

[2] The Jackson Laboratory, Bar Harbor, ME 04609, USA.

[3] Dept. of Biomedical Engineering, Johns Hopkins University, Baltimore, 21205 MD, USA

[4] Section on Sensory Cell Development and Function, National Institute on Deafness and Other Communication Disorders, National Institutes of Health, Bethesda, MA, USA

[5] Department of Neurobiology, Spencer Fox Eccles School of Medicine, University of Utah, Salt Lake City, Utah, USA.

[6] Department of Otolaryngology - Head & Neck Surgery, Spencer Fox Eccles School of Medicine at the University of Utah, Salt Lake City, Utah, USA.

[7] Department of Otolaryngology-Head and Neck Surgery, Johns Hopkins University, Baltimore 21205 MD, USA.

[8] Department of Neuroscience, Johns Hopkins University, Baltimore 21205 MD, USA.

[9] Kavli Neuroscience Discovery Institute, Johns Hopkins University, Baltimore 21205 MD, USA.

[10] Tufts University School of Medicine, Boston, MA 02111, USA.

[#] co-first authors

[*] Corresponding authors

SUMMARY/ABSTRACT

Otolith organs in the inner ear and neuromasts in the fish lateral-line harbor two populations of hair cells oriented to detect stimuli in opposing directions. The underlying mechanism is highly conserved: the transcription factor EMX2 is regionally expressed in just one hair cell population and acts through the receptor GPR156 to reverse cell orientation relative to the other population. In mouse and zebrafish, loss of Emx2 results in sensory organs that harbor only one hair cell orientation and are not innervated properly. In zebrafish, Emx2 also confers hair cells with reduced mechanosensory properties. Here, we leverage mouse and zebrafish models lacking GPR156 to determine how detecting stimuli of opposing directions serves vestibular function, and whether GPR156 has other roles besides orienting hair cells. We find that otolith organs in *Gpr156* mouse mutants have normal zonal organization and normal type I-II hair cell distribution and mechano-electrical transduction properties. In contrast, *gpr156* zebrafish mutants lack the smaller mechanically-evoked signals that characterize Emx2-positive hair cells. Loss of GPR156 does not affect orientation-selectivity of afferents in mouse utricle or zebrafish neuromasts. Consistent with normal otolith organ anatomy and afferent selectivity, *Gpr156* mutant mice do not show overt vestibular dysfunction. Instead, performance on two tests that engage otolith organs is significantly altered – swimming and off-vertical-axis rotation. We conclude that GPR156 relays hair cell orientation and transduction information downstream of EMX2, but not selectivity for direction-specific afferents. These results clarify how molecular mechanisms that confer bi-directionality to sensory organs contribute to function, from single hair cell physiology to animal behavior.

INTRODUCTION

The vestibular system provides essential information about the head's motion and orientation relative to gravity and plays a vital role in everyday life. Fundamental gaps exist in our understanding of vestibular development and function. For example, we do not fully understand how the abrupt reversal of hair cell (HC) orientation in otolith organs affects vestibular function, and whether the reversal also impacts HC mechano-electrical transduction properties or afferent neuron contacts. We explore these questions further in mouse and in the zebrafish lateral line, a sensory system where external fluid flow is detected by neuromast organs that also feature HC reversal.

Vestibular HCs reside in five sensory organs per inner ear in human, mouse and larval zebrafish: three cristae ampullaris and two otolith (or macular) organs. Within each of these organs, HCs are precisely oriented to detect mechanical stimuli based on direction. HCs are sensitive to the proportion of a stimulus aligned with the mechanosensory hair bundle's orientation: a vector from the bundle's short edge to its tall (kinociliary) edge. In cristae, which detect angular head motions, HCs are oriented uniformly to detect fluid flow in the plane of the attached semicircular canal. In contrast, HCs in otolith organs are oriented to respond to a range of head positions and linear motions in an approximately horizontal (utricle) or vertical (saccule) plane (Fig. 1A). Within these planes, HCs show two opposing orientations across a virtual "line of polarity reversal" (LPR), a conserved anatomical feature in all otolith organs (Denman-Johnson and Forge, 1999; Flock, 1964; Lindeman, 1969). HCs are also aligned in two opposing orientations in fish and amphibian neuromasts in the lateral-line system. Orientation reversal confers on neighboring HCs of opposing orientations opposite responses to the same local stimulus: depolarization and increased afferent spike rate in one population, but hyperpolarization and decreased spike rate in the other. This functional prediction and its utility are well established in neuromasts as the ability to sensitively detect water movements in two directions (Chitnis et al., 2012; Lopez-Schier et al., 2004). Beyond increasing the range of stimulus orientations detected, it remains unclear how HCs with opposing orientations and responses serve otolith organ function.

In the utricle, the LPR also segregates distinct zones, for example the lateral extrastriola (LES) from the more medial striola (Fig. 1A). Zones are strikingly different regions found in all vestibular epithelia, with 'central' and 'peripheral' zones of cristae corresponding to striolar and extrastriolar zones of maculae, respectively. Afferents that innervate striolar HCs have irregular spike timing and adapting responses in contrast to afferents that innervate extrastriolar HCs which have more regular and tonic spikes (Goldberg, 1991; Goldberg, 2000). Striolar afferents transmit more information about the time course of translational self-motion, while extrastriolar afferents better discriminate between different static head orientations relative to gravity (Jamali et al., 2019). Correlated with these physiological differences between zones are differences in HCs and afferent morphologies. Both the striola and extrastriola are populated by type I HCs contacted by *calyx afferent endings* and type II HCs contacted by *bouton afferent endings.* Most individual afferents in both zones contact both type I and type II HCs ("*dimorphic afferents*"). However, pure-calyx and pure-bouton individual afferents are respectively only found in the striola and extrastriola (Eatock and Songer, 2011; Goldberg, 2000). In amniotes there is no obligatory relationship between HC orientation reversal and zones. While the utricular striola is entirely medial to the LPR, the saccular striola is bisected by the LPR (Fig. 1A) (Deans, 2013; Li et al., 2008; Xue and Peterson, 2006). Moreover, central and peripheral zones of cristae bear many anatomical and physiological similarities to the striolar and extrastriolar zones, but cristae have no LPR and no abrupt variation in HC orientation, as in the organ of Corti.

Interestingly, mammalian otolith afferents have two distinct anatomical properties that segregate with the LPR. First, individual afferents are maximally excited by stimuli of one orientation and not the opposite (Fernandez et al., 1972; Goldberg et al., 1990), indicating that individual afferents selectively contact HCs sharing the same general orientation, as also observed in the zebrafish lateral line (Ji et al., 2018). Second, the central projections of afferent neurons innervating HCs on each side of the LPR are also distinct; afferents that innervate one HC orientation largely project to the cerebellum, while the opposite HC orientation projects to the brainstem (Ji et al., 2022; Maklad et al., 2010). The LPR thus

correlates with a striking segregation of both the peripheral and central projections of

afferent neurons.

Recent work has illuminated how HCs with opposing orientations are formed during

development (reviewed in (Tarchini, 2021)). Early in development all organs are polarized

by core PCP proteins that are enriched asymmetrically at cell-cell junctions before HCs

are born (Deans, 2013). In cristae, asymmetric core PCP proteins set up a default,

uniform HC orientation across the epithelium. Because the asymmetric pattern of core

PCP protein enrichment is unchanged across the LPR, a distinct, regional mechanism

defines a HC population with an opposite orientation (Deans, 2013; Deans et al., 2007;

Jones et al., 2014; Mirkovic et al., 2012). Regional expression of the transcription factor

EMX2 in the lateral utricle and posterior saccule drives the 180°-reversed orientation of

HCs compared to neighboring *Emx2*-negative HCs (Holley et al., 2010; Jiang et al., 2017).

Although regionally restricted, *Emx2* is expressed by all cell types in the sensory

epithelium. Reversal is achieved cell-autonomously in HCs by an orphan GPCR, GPR156

(Kindt et al., 2021). Unlike *Emx2, Gpr156* is transcribed throughout the sensory epithelia

but is limited to HCs. In *Emx2*-positive HCs only, the GPR156 protein is trafficked and

planar-polarized at apical junctions, where it signals through inhibitory G proteins to

trigger a reversal in HC orientation. The EMX2>GPR156 reversal cascade is conserved

in zebrafish neuromasts and loss of either of these proteins produces neuromasts where

all HCs adopt the same, uniform orientation (Jacobo et al., 2019; Jiang et al., 2017; Kindt

et al., 2021; Kozak et al., 2020; Lozano-Ortega et al., 2018).

Importantly, in addition to reversing HC orientation, Emx2 is required for afferent

selectivity. In zebrafish *emx2* mutants, individual afferent fibers either innervate nearly all

or very few HCs instead of innervating ~50% of HCs based on their shared orientation (Ji

et al., 2018). In mouse otolith organs, regional *Emx2* expression is required for local

afferents that innervate HC of one orientation to project centrally to the cerebellum and

not to the brainstem (Ji et al., 2022). In zebrafish, Emx2 was also recently shown to affect

HC mechanotransduction, thus defining an orientation-based functional asymmetry in

lateral-line neuromasts (Chou et al., 2017; Kindig et al., 2023). It remains unknown

whether, similar to Emx2, GPR156 is required for orientation-based selectivity of afferent contacts in mouse otolith organs or zebrafish neuromasts, and whether Gpr156 participates in direction-based asymmetry of mechanosensitive response in neuromasts.

Here, using animal models where *Gpr156* is inactivated, we first established that abolishing the LPR in mouse otolith organs did not produce obvious changes in HC patterning and mechanotransduction properties. In contrast, we found that similar to Emx2, Gpr156 is required for asymmetric mechanotransduction in zebrafish neuromasts. Normal afferent segregation was maintained in both mice and zebrafish lacking GPR156 despite one HC population failing to reverse its orientation. Unchanged afferent organization in the mouse allowed us to specifically interrogate how the LPR serves vestibular function. We found that although *Gpr156* mutant mice did not show overt vestibular dysfunction, performance on two tests that engage otolith organs was significantly altered. This study clarifies how molecular mechanisms that orient HCs to achieve bi-directional sensitivity contribute to vestibular and lateral-line function.

RESULTS

Normal hair cell numbers and organization in *Gpr156* mutant adult otolith organs

In previous work (Kindt et al., 2021), we established that in sensory organs of the inner

ear and lateral-line, *Gpr156* expression is limited to HCs. In neonate otolith organs, we

demonstrated that GPR156 is required for EMX2-positive HCs to adopt a reversed

orientation compared to EMX2-negative HCs. We further showed that EMX2-positive HCs

lacking GPR156 do not correct their orientation with time and remain non-reversed in

young adults (P21-P28). Focusing on the utricular sensory epithelium, we showed normal

macular surface area and number of HCs in all three zones (LES, striola and MES; Fig.

1A) in adult *Gpr156*$^{del/del}$ mutants. Here we begin by extending our morphological

characterization of *Gpr156*$^{del/del}$ adult macular organs to include the saccule as well as

zonal and HC type profiles, before proceeding to functional and behavioral assessments.

We first verified that absence of GPR156 did not obviously interfere with striolar patterning

in adult animals. We used oncomodulin (OCM) to immunolabel striolar HCs (Simmons et

al., 2010) at 4-week of age (P28) and confirmed that striolar surface area and striolar HC

density were unaffected in constitutive *Gpr156*$^{del/del}$ mutant utricles and saccules

compared to controls (*Gpr156*$^{del/+}$; Fig. 1B-C). Each macular organ contains two main HC

types, type I and type II. Therefore, we next quantified type I HCs in the LES, striola and

MES zones of the adult utricle and in the anterior (ANT), striolar and posterior (POST)

zones of the adult saccule. We used SPP1/Osteopontin, a marker that labels the neck

region of approximately 90% of type I HCs (Fig. 1D-E) (McInturff et al., 2018). We

observed no difference in SPP1-positive type I HC numbers between mutants and

controls in any zone or organ (Fig. 1F). We next used a SOX2 antibody to label type II

HCs in the same zones (Fig. 1G-H). Again, we observed no difference in type II HC

numbers between mutants and controls (Fig. 1I). In summary, loss of GPR156 prevents

Emx2-positive HCs from reversing their orientation but does not affect macular patterning,

including division into clear zones and appropriate numbers of type I and type II HCs

throughout.

No defects in mechano-electrical transduction were detected in individual mouse utricular hair cells of *Gpr156* mutants

After establishing normal numbers and types of mouse vestibular HCs, we assessed whether HCs respond normally to hair bundle deflections in the absence of GPR156. For this assessment, we recorded whole-cell mechano-electrical transduction (MET) currents of HCs in the utricle of *Gpr156$^{del/del}$* mutants. To stimulate transduction, we deflected individual mechanosensory hair bundles of HCs from *Gpr156$^{del/del}$* mutants and heterozygous *Gpr156$^{del/+}$* control utricles with a rigid probe coupled to the back (short edge) of the bundle (see Methods, Fig. 2A). All bundles were positively deflected along their orientation axes. I_{MET} recordings were taken between P8 and P100 (median age for comparisons: ~P20 for both null and heterozygous genotypes; see Tables).

We first compared properties of the evoked MET current (I_{MET}) across genotypes in the LES, the zone in which hair bundles fail to reverse in *Gpr156* mutants. ~70% (93/135) of LES HCs responded with detectable I_{MET} (within 20 pA) to hair bundle deflection, regardless of HC type and genotype (Chi-squared test, Table 1). We attribute the insensitivity of the remaining cells to damage during dissection of the tissue, given that damaged bundles can be seen in every preparation, zone, and bundle type, independent of genotype.

I_{MET} responses are shown for families of iterated 400-ms steps of displacement; subsets of these families are shown in Fig. 2B-E. During bundle displacement, I_{MET} typically peaked at step onset. As noted before (Vollrath and Eatock, 2003), traces evoked by stimuli in the middle of the operating range adapted with a time course fit by 1-2 decaying exponential terms. Because type I and II HCs were voltage-clamped at different holding potentials (V_{hold} = −94 and −84 mV, respectively, see Methods), their MET currents experienced different driving forces. We normalized for driving force by converting current to conductance (G) and plotted G as a function of hair bundle displacement (X) for each HC type and genotype (Fig. 2F-G). G(X) curves were fit with a simple Boltzmann function

(Eq. 1, Methods), yielding values for maximum conductance (G_{max}), the midpoint of the G(X) curve ($X_{1/2}$, displacement at half-maximal I_{MET}) and dx (slope factor), the displacement range corresponding to an e-fold increase in G for small displacements. Among the LES HCs tested, no significant differences were detected between genotypes for any Boltzmann parameter (G_{max}, $X_{1/2}$, or dx) (Fig. 2F-G, Table 2). G_{max} values were in the range of 2.5-3.0 nS, $X_{1/2}$ was 200-300 nm and dx was 150-200 nm (Table 2). From G(X) relations we estimated the operating range (OR; displacement range yielding 10-90% growth of G_{MET}) to be ~800-900 nm for all LES HCs (Table 2). In summary, the peak (non-adapted) displacement sensitivity of I_{MET} was not detectably affected in LES hair bundles of mutants.

Another salient feature of vestibular HC transduction is adaptation (response decay) during a steady hair bundle deflection. We fit the adaptation with one or more exponential decays, and also calculated its extent (% decay) (Eq. 4, Methods; Fig. 2B-E). As reported previously for immature mouse utricular HCs (Vollrath and Eatock, 2003), these mature mouse utricular HCs had fast and slow adaptation components. The fast component was more prominent at small displacements and the slow component dominated at large displacements; often, both were evident in the middle of the operating range. We therefore compared adaptation time constants, τ_{fast} and τ_{slow}, at the midpoint of the operating range ($X_{1/2}$). For 400-ms steps, mean values are in the range of 4 - 10 ms and 70 – 140 ms, respectively, and the two components are responsible for comparable extents of the total adaptation. These results are consistent with previous reports with similar methods from immature mouse utricular HCs (Vollrath and Eatock, 2003) and immature rat saccular HCs (Songer and Eatock, 2013. In the current sample, however, adaptation at $X_{1/2}$ included a significant third, faster adaptation component ("very fast"; τ_{VF} < 1 ms) in most type I HCs (12/13) and 2/10 type II HCs. In the LES, we detected no significant differences with genotype in adaptation rates or extent (Table 2).

No changes were detected in voltage-gated conductances of individual mouse utricular hair cells of *Gpr156* mutants

MET current initiates the receptor potential, which in turn modulates substantial voltage-
gated currents in the HC's basolateral membrane, especially K^+ currents, which further
shape the receptor potential. For example, type I HCs make relatively fast, small receptor
potentials compared to type II HCs because type I HCs have many more K channels that
are open around resting potential. To test for effects of *Gpr156* deletion on voltage-
dependent currents, we applied voltage step protocols to whole-cell clamped type I and
type II HCs in the LES of *Gpr156^(del/del)* and *Gpr156^(del/+)* utricles (Fig. 3A-D).
270
The time course and voltage dependence of outwardly rectifying voltage-gated current
from LES type I cells of *Gpr156^(del/del)* utricles resembled those of *Gpr156^(del/+)* utricles (Fig.
3A, C). Most of the current in type I HCs (Fig. 3A, C) flows through a low-voltage-activated
K^+ conductance ($g_{K,L}$) that dominates the type I HC voltage response. As described in
Methods, we used the current data to generate conductance density vs. voltage (G/Cm -
V) curves (Fig. 3E) and fitted them with the Boltzmann function (Eq. 5, Methods) to show
how the open probability of $g_{K,L}$ varies with voltage. In both genotypes, approximately half
of $g_{K,L}$ was activated at resting membrane potential, V_{rest} (~–85 mV). As a consequence
of their large K^+ conductance at V_{rest}, type I cells had low input resistances (R_{in} ~50 MΩ).
As is typical of type I HCs, the maximal K^+ conductance density (G_{max}/C_m), which is
proportional to the maximal number of open channels per unit membrane area, was much
larger than for type II HCs (Fig. 3E vs. 3F, note difference in conductance scale; Table 3).
283
Similarly, we detected no effect of genotype on the voltage-dependent outward currents
of LES type II HCs (Fig. 3B, D, F). The currents activated positive to resting potential (~–
72 mV) and had activation midpoints ($V_{1/2}$ values) of –30 to –40 mV. Because of the small
voltage-gated conductance at rest, input resistance was ~10-fold higher than in type I
HCs (~500 MΩ vs. ~50 MΩ). The outward currents inactivated significantly within 400 ms,
indicating a substantial A-current component. The maximum conductance density
(G_{max}/C_m) at 400 ms (~steady state), which is proportional to the number of open channels
per unit membrane area, was much smaller than in type I HCs. Again, no effect of
genotype was detected on the differences between type I and II HCs of the LES (Table
3).

Together these results are consistent with previous reports on key physiological properties of type I and II HCs in amniote vestibular organs. The mature physiological status of each HC type appeared to be preserved in LES HCs that failed to reverse their orientation because GPR156 was inactivated.

Hair cell physiology beyond the lateral extrastriolar domain: Variations in properties were not related to loss of GPR156

Loss of GPR156 does not impact HC orientation in the striola and MES, but *Gpr156* is transcribed in all HCs across the maculae (Kindt et al., 2021). We thus investigated transduction and adaptation in those zones as well (Fig. 4). While there was heterogeneity across HCs, we saw no striking systematic differences with genotype in the voltage-gated or mechanotransduction currents from control *vs. Gpr156* mutants (but note that the statistical power of the comparisons is low, Table 4).

Gpr156 impacted the mechanosensitive properties of lateral-line hair cells in zebrafish

Similar to otolith organs of mammals, each neuromast within the posterior lateral-line system of zebrafish has two populations of HCs that enable the detection of anterior- or posterior-directed fluid flow (Fig. 5A). To allow bi-directional detection, both Emx2 and Gpr156 are required to reverse the orientation of one HC population so that it can detect posterior flow (Fig. 5A) (Jiang et al., 2017; Kindt et al., 2021). Previous work has shown that in wild-type animals, the mechanosensitive properties of HCs that detect anterior flow (Emx2$^-$) are larger compared to those that detect posterior flow (Emx2$^+$) (Kindig et al., 2023). Further, this work demonstrated that loss or gain of Emx2 in all lateral-line HCs was able to not only alter HC orientation, but also increase and decrease mechanosensitive responses respectively (Kindig et al., 2023). Our current results in the mouse utricle did not detect significant differences in the mechanotransduction properties

of HCs lacking GPR156 (Fig. 2). Whether loss of Gpr156 alters the mechanosensitive

properties of lateral-line HCs is not known.

To address this question, we used a membrane-localized GCaMP6s reporter in HCs

(memGCaMP6s) (Kindt et al., 2021) to perform *in vivo* calcium imaging (Lukasz and Kindt,

2018) and. By imaging GCaMP6s signals in an apical plane containing the hair bundles,

we measured evoked, mechanosensation-dependent calcium signals during fluid-jet

stimulation. We used this approach to stimulate in both the anterior and posterior direction

and compared the magnitude of the mechanosensitive responses in lateral-line HCs in

both control (*gpr156+/-*) and *gpr156* mutants. As shown previously, we could reliably

identify the orientation (ANT vs POST) among control HCs using this approach (eg: Fig.

5B). We first verified that in control animals, the magnitude of the mechanosensitive

responses of HCs that detect anterior flow was greater than those that detect posterior

flow (eg: Fig. 5B and Fig. 5D, F). Consistent with previous work, in *gpr156* mutants, we

found that the majority of HCs only responded to anterior flow (eg: Fig. 5C) (Kindt et al.,

2021). Importantly, we found that the magnitude of the mechanosensitive responses in

gpr156 mutants was not different when compared to control HCs that respond to anterior

flow (eg: Fig. 5C and Fig. 5E-F). Further, the mechanosensitive responses in *gpr156*

mutants were significantly larger than control HCs that respond to posterior flow (Fig. 5F).

Our analysis thus indicates that in *gpr156* mutants, lateral-line HCs show larger

mechanosensitive responses. These responses are similar to HCs in control animals that

respond to anterior flow (Emx2-). Together, this data suggests that the EMX2-GPR156

mechanism driving HC orientation reversal also affects HC transduction in zebrafish

neuromasts.

Segregated afferent receptive fields were preserved in mouse utricles lacking

GPR156

After characterizing the functional properties of HCs in mouse and zebrafish *Gpr156*

mutants, we examined whether GPR156 plays a role in orientation-selective afferent

innervation. Previous work has demonstrated that *Emx2* expression in the LES of the mouse utricle and in HCs that sense posterior flow in the lateral line is required for afferents to innervate HCs of similar (rather than opposing) orientation (Ji et al., 2022; Ji et al., 2018). Whether GPR156 plays a similar role in direction-selective afferent innervation is unknown.

To investigate whether afferent organization in the LES is disrupted by loss of GPR156, we first compared the terminal fields of individual dye-filled calyx-bearing afferents in *Gpr156del/+ and Gpr156del/del* utricles. All LES calyces belong to dimorphic afferents, which form both calyces on type I HCs and boutons on type II HCs. To examine terminal fields, a fluorescent internal solution was injected into a calyx terminal within the LES during whole-cell recordings. The fluorescent dye was then allowed to diffuse into the calyx and from there throughout the terminal arbor.

Using this approach, we found that terminal arbors (receptive fields) in *Gpr156del/+* control utricles (n = 12) comprised 1 or 2 calyces and as many as 65 boutons (median: 29) (Fig. 6A-C). The labeled afferent terminals were restricted to LES in all but one case (Fig. 6A). In this single case, a terminal branch crossed the LPR, innervating 16 striolar HCs with opposite orientation to the 12 LES HCs also innervated. In some filled LES afferents, a thin branch originated below the epithelium (Fig. 6C, Supplemental video 1), as previously described for chinchilla extrastriola (Fernandez et al., 1990). We also labeled a single striolar calyx from a control utricle near the striolar/MES boundary; its terminal field was excluded from the calbindin-negative MES (Fig. 6D).

When we performed dye labeling in *Gpr156del/del* utricles, the innervation patterns and the size of terminal fields appeared similar to *Gpr156del/+* controls. In *Gpr156del/del* utricles, the labeled LES afferents (n = 8) were also all dimorphic, terminating in up to three calyces and 56 boutons (median: 23). The labeled receptive fields did not cross the striola/LES border (Fig. 6E-G). We also did not observe fiber branches extending across the border from parent axons (Fig. 6E-G; Supplemental video 2). As observed in the lone *Gpr156del/+* striolar afferent (Fig. 6D), *Gpr156del/del* afferents with the main dendritic fiber terminating

in the striola (n = 4) contacted relatively few HCs, and receptive fields were restricted to

the calbindin-positive region (Figure 6H).

To confirm and complement dye-filling results at the whole organ level, we used a new

genetic approach to specifically label afferents innervating EMX2$^+$ HCs in the LES.

Interestingly, when breeding the *Advillin-Cre* strain (Zhou et al., 2010) with the *Tigre Ai140*

reporter (Daigle et al., 2018), we only observed Cre recombination and EGFP expression

in vestibular ganglion neurons that contacted HCs in the lateral utricle region (Fig. 7A).

Close examination revealed clear labeling of HC calyces in the LES, and occasionally in

OCM$^+$ HCs in the lateral striola (Fig. 7B, arrowheads). We confirmed that similar to OCM$^-$

EGFP$^+$ HCs in the LES, these OCM$^+$ EGFP$^+$ HCs were invariably oriented towards the

medial edge (Fig. 7C). This demonstrated that *Advillin* expression defines a population of

neurons whose peripheral processes closely segregate with HC orientation, irrespective

of striolar (OCM$^+$) or extrastriolar (OCM$^-$) HC identity. Importantly, HC innervation by

Advillin-positive neurons was unchanged in *Gpr156$^{del/del}$* mutants (Fig. 7A-B). We

observed equivalent numbers of OCM$^+$ HCs innervated in the lateral striola in *Gpr156$^{del/del}$*

mutants compared to controls (Mann-Whitney p=0.74; average OCM$^+$/EGFP$^+$ HCs per

utricle: *Gpr156$^{del/+}$*: 20.25, *Gpr156$^{del/del}$*:19.25; 3 animals and 4 utricles per genotype; no

OCM$^+$/EGFP$^+$ HCs oriented towards the lateral edge were recorded in either genotype).

In summary, we detected no anomaly in afferent receptive fields in utricles lacking

GPR156 and thus lacking a LPR, showing that the tendency of afferents to innervate HCs

on one side of the LPR is not controlled by GPR156 or HC orientation.

Afferent firing patterns were preserved in mouse utricles lacking GPR156

Our innervation data suggested that HCs in zones demarcated by the LPR are still

innervated by distinct populations of afferent neurons in *Gpr156$^{del/del}$* mutants. We next

investigated whether the LES afferent population retains normal physiology. Striolar

afferents have highly irregular inter-spike intervals, while extrastriolar afferents in both

LES and MES have much more regular inter-spike intervals (Goldberg, 1991; Goldberg,

2000; Goldberg et al., 1990). Associated with the difference in spike timing regularity are
zonal differences in response dynamics and information encoding capacity (e.g., (Jamali
et al., 2019)). The spike timing difference is considered to reflect differences in afferents'
excitability as conferred by their intrinsic ion channels, with regular afferents being more
excitable (Kalluri et al., 2010; Ventura and Kalluri, 2019). To test for effects of *Gpr156*
inactivation on afferent excitability, we injected small depolarizing current steps into LES
calyces through the whole-cell recording pipette and measured current threshold for
spiking, I_{thresh}, and the numbers of spikes evoked by each current step.

As reported (Gonzalez-Garrido et al., 2021; Ono et al., 2020), I_{thresh} evoked 1-to-several
spikes but increments beyond I_{thresh} evoked more sustained firing (Fig. 8a-b). No
difference was observed across genotypes (Table 5): in both *Gpr156$^{del/+}$* afferents (n=18)
and *Gpr156$^{del/del}$* afferents (n=20), mean I_{thresh} was ~100 pA and ~95% of afferents
produced sustained spiking at $3 \times I_{thresh}$. These results suggest that afferent firing patterns,
and likely the underlying ion channel expression, are unaltered in the LES of *Gpr156$^{del/del}$*
utricles, despite abnormal HC orientation.

Gpr156 was not required for afferent selectivity or synaptic pairing in zebrafish
lateral-line

We next expanded our analysis to determine whether loss of Gpr156 impacts afferent
selectivity in the lateral-line. For this analysis we used mosaic expression of
neuroD:tdTomato to label single lateral-line afferent neurons. After identifying *gpr156*
mutant and control *(gpr156$^{+/-}$)* animals with labeled fibers, we immunolabeled to visualize
orientation (phalloidin), HC bodies (MYO7), and cell type (presence or absence of Emx2).
We then used confocal microscopy to image the innervation pattern of single afferents.

Consistent with previous results, some afferent fibers selectively contacted Emx2$^-$ HCs
with hair bundles oriented to respond to anterior flow in control animals (Fig. 9A-H). Other
afferent fibers selectively contacted Emx2$^+$ HCs with hair bundles oriented to respond to
posterior flow (Supplemental fig. 1A-H). We then examined the innervation patterns of

afferent fibers in *gpr156* mutants. Similar to controls we were able to identify fibers with a

clear preference for Emx2$^-$ or Emx2$^+$ HCs (Fig. 9I-P, Supplemental fig. 1I-P). Importantly,

in *gpr156* mutants, afferent fibers innervated HCs based on the presence or absence of

Emx2, despite all HCs responding to anterior flow.

Following this initial assessment, we quantified the percentage of HCs per neuromast that

were innervated by each afferent fiber. We found that *gpr156* mutant fibers innervated a

similar percentage of HCs per neuromast compared to controls (Fig. 9Q, control: 38.90 %

± 2.40, n = 18, *gpr156* ko: 41.31 % ± 2.33, n = 23, p = 0.48). We also quantified how

selective each afferent fiber was for Emx2$^+$ or Emx2$^-$ HC. For each afferent fiber, we

calculated the percentage of either Emx2$^+$ or Emx2$^-$ HCs innervated out of the total

number of innervated HCs. In both control and *gpr156* mutants, afferent fibers showed a

high selectivity (Fig. 9R, control: 85.06 % ± 4.32, n = 18, *gpr156* ko: 81.29 % ± 3.17, n =

23, p = 0.48). We split this dataset and examined the selectivity of afferent fibers that

preferentially contacted either Emx2$^-$ or Emx2$^+$ HCs (Fig. 9S-T). This analysis revealed

that in control and *gpr156* mutants, both afferent fiber types were highly selective

(afferents contacting Emx2$^-$ cells, control: 81.41 % ± 7.90, n = 8, *gpr156* ko: 79.37 % ±

4.73, n = 10, p = 0.82; afferents contacting Emx2$^+$ cells, control: 87.99 % ± 4.75, n = 10,

gpr156 ko: 82.80 % ± 4.38, n = 13, p = 0.43). Overall, our single fiber labeling revealed

that Gpr156 is not required for afferent fibers to selectively innervate HCs based on the

presence or absence of Emx2.

To extend this analysis, we examined synaptic pairing in *gpr156* mutants. Previous

research found that along with inappropriate innervation, there is an increase in the

amount of unpaired pre- and post-synapses in *emx2* mutants (Ji et al., 2018). We

immunostained control along with *gpr156* and *emx2* mutant HCs with Ribeye b to label

presynapses and pan-MAGUK to label postsynapses (Fig. 10A-C). We then quantified

the number of complete immunolabeled synapses per HC. Among all genotypes, the

number of complete synapses was unaltered (Fig. 10D, control 3.56 ± 0.06, n = 9

neuromasts, *gpr156* ko: 3.61 ± 0.10, n = 16 neuromasts, p = 0.74; control 3.51 ± 0.27, n

= 9 neuromasts, *emx2* ko: 3.67± 0.08 n = 9 neuromasts, p = 0.50). We also quantified

unpaired pre- and post-synapses. Similar to previous results, there were more unpaired

pre- and post-synapses per neuromast in *emx2* mutants compared to controls (Fig. 9E-

F, presynapses, *emx2*$^{+/+}$ or *emx2*$^{+/-}$ control 2.00 ± 0.53, n = 9 neuromasts, *emx2* ko: 4.22

± 0.70, n = 9 neuromasts, p = 0.02; postsynapses, control 1.44 ± 0.56, n = 9 neuromasts,

emx2 ko: 6.56 ± 0.78, n = 9 neuromasts, p < 0.0001). In contrast, in *gpr156* mutants,

there was no increase in unpaired pre- or post-synapses per neuromast compared to

controls (Fig. 10E-F, presynapses, *gpr156*$^{+/-}$ control 2.11 ± 0.31, n = 9 neuromasts,

gpr156 ko: 1.68 ± 0.23, n = 16 neuromasts, p = 0.29; postsynapses, control 0.33 ± 0.17,

n = 9 neuromasts, *gpr156* ko: 0.75 ± 0.17, n = 16 neuromasts, p = 0.12). A lack of unpaired

pre- and post-synapses in *gpr156* mutants further confirms that Emx2 but not Gpr156

impacts afferent innervation in lateral-line HCs.

In summary, *Gpr156* inactivation does not appear to impact afferent selectivity in either

mouse or zebrafish. This is in sharp contrast with results in *Emx2* mutants, where

orientation-specific HC afferent innervation is profoundly altered in mouse and zebrafish

(Ji et al., 2022; Ji et al., 2018). It follows that HC orientation per se does not dictate afferent

selectivity, with EMX2 likely using a different effector(s) than GPR156 in this context. Thus,

afferent studies confirm that the *Gpr156*^{del} model is well-suited to investigate specifically

how the reversal of HC orientation in otolith and lateral line organs serves organ function.

Gpr156 mutant mice had defects in otolith-driven, but not canal-driven, vestibular

behaviors

In mammals, only otolith organs (utricle, saccule) have a LPR. Therefore, the absence of

a LPR in *Gpr156* mutants should only affect otolith-driven reflexes and behaviors. In order

to test this prediction we compared *Gpr156* control and mutant mice on a number of tests

that can discriminate between otolith and canal inputs.

Normal posture, balance beam but impaired swimming performance in *Gpr156* mutants

To assess how loss of the LPR affects vestibular functions, we first determined whether

Gpr156 mutant mice showed postural/balance impairments. Visual and quantitative

assessments did not reveal circling or head tremor behaviors in *Gpr156^{del/del}* mice at rest

(Supplemental fig. 2). Additionally, we conducted basic standard behavioral tests that

involve subjective scoring: (1) tail hanging, (2) air righting, and (3) contact inhibition of

righting. Both control and *Gpr156^{del/del}* mice scored 0 (where 0=normal, see Methods) on

each of these behavioral tests (Supplemental fig. 3A-C).

After this initial assessment, we next subjected mice to more challenging balance tests

including walking along a narrow balance beam and swimming. On the balance beam,

both *Gpr156^{del/del}* and *Gpr156^{del/+}* mice showed normal coordination (Fig. 11A). However,

when *Gpr156^{del/del}* mice were placed in water, they demonstrated severely impaired

swimming compared to *Gpr156^{del/+}* controls. This impairment was characterized by

underwater tumbling and the inability to maintain normal upright posture in the water. All

mice were rescued within 60 seconds and the average time of rescue was significantly

lower in *Gpr156^{del/del}* compared to control mice (Fig. 11B; *Gpr156^{del/+}* controls= 49.9±3.0s;

Gpr156^{-del/del}= 17.4±2.7s; *P*<0.0001). Overall, these results indicated that GPR156 is

required for vestibular function important for swimming behavior.

We hypothesized that markedly impaired swimming performance could be due to deficits

in the otolith rather than semicircular canal dependent pathways. To test this proposal,

we quantified eye movements generated by two different vestibulo-ocular reflex

behaviors, one driven by activation of the semicircular canals (angular vestibular-ocular

reflex, or angular VOR) and the other driven by activation of the otoliths (off-axis vertical

rotation, or OVAR).

<u>Normal aVOR and OKR responses in *Gpr156* mutants</u>

We first quantified the angular vestibulo-ocular reflex in *Gpr156^{del/del}* and *Gpr156^{del/+}*

control mice in darkness (VORd). The angular VOR was evoked by rotating the animal

sinusoidally around earth vertical axis. To quantify the angular VORd we computed the

gain and phase for each testing frequency (0.2, 0.4, 0.8, 1, and 2Hz, ±16°/s). During

VORd, both control and *Gpr156^{del/del}* mice displayed robust compensatory eye

movements, which increased as a function of frequency (Fig. 12A-B). Further, to confirm

that there was no visual deficit in *Gpr156*^{del/del} mice, we also tested their optokinetic reflex

(OKR) by rotating a patterned visual surround sinusoidally about earth vertical axis at

the same testing frequencies, while the animal remained stationary (see Methods). The

OKR responses of *Gpr156*^{del/del} and control mice were comparable, confirming that there

was no visual deficit in *Gpr156* mutants (Fig. 12C-D). For the sake of completeness, we

quantified the angular VOR evoked in the light (VORl) by rotating the animal sinusoidally

around earth vertical axis, while the patterned visual surround remained stationary.

Again, we found no difference in the VORl response of *Gpr156*^{del/del} versus control mice

in this lit environment (Figure 12E-F). Finally, we tested whether *Gpr156*^{del/del} mice

demonstrated normal ability to adapt their angular VOR. We used a standard angular

VOR motor learning protocol (see Methods) in which we carried out 30-minute-long VOR

gain-down training for both control and *Gpr156*^{del/del} mice. Head restrained mice were

rotated with the visual surround in phase. The training stimulus was 2 Hz with a peak

velocity of 16°/s. The learning efficacy was assessed by quantifying change in the VOR

gain after training. In both control and *Gpr156*^{del/del} mice, this VOR motor learning

protocol produced a significant reduction in VOR gain after training (Supplemental fig.

4A-B). Overall, *Gpr156*^{del/del} and control mice demonstrated a comparable percent

change of gain decrease at each testing frequencies (Supplemental fig. 4C). Together,

our VOR and OKR results indicated normal semicircular canals-driven responses in

Gpr156^{del/del} mice.

<u>Altered OVAR responses in *Gpr156* mutants</u>

As no defects were apparent in semicircular canal-mediated behavior, we next tested

whether otolith-mediated vestibulo-ocular reflex behaviors were impaired in *Gpr156*^{del/del}

mice. For this assessment, we recorded eye movements while the mouse was tilted 17°

off-the vertical axis (i.e., off-vertical axis rotation (OVAR)), and then rotated at a constant

velocity (50°/s for 72 sec). The eye velocity evoked by this paradigm comprised two

different responses: (1) a transient canal-mediated response that decayed over the first

10-15 sec and (2) an otolith-mediated steady-state response in which eye velocity

oscillated around a constant bias with a sinusoidal waveform (Figure 12G). Overall, our

analysis revealed that transient OVAR responses did not differ between control and

Gpr156$^{del/del}$ mice, supporting the proposal that absence of GPR156 did not affect canal-
related ocular responses. In contrast, steady-state OVAR responses demonstrated a
significantly impaired bias in *Gpr156$^{del/del}$* mice as compared to controls (Figure 12G-H).
The OVAR of *Gpr156$^{del/del}$* mice was characterized by a significantly reduced bias relative
to controls (*Gpr156$^{del/del}$*: bias=0.039 ± 0.26 versus control: bias=3.85 ± 0.54, P = 4.03E-
5). Thus taken together, these results show that the absence of HC reversal in
Gpr156$^{del/del}$ alters otolith-mediated OVAR responses. This in turn may explain the
swimming deficits observed (Fig. 11B).
580
581
582

DISCUSSION

Hair bundle orientation vs. zonal identity

Taken together, our observations in utricles lacking GPR156 agree with other evidence

that in mouse vestibular epithelia, HC orientation is controlled separately from other

zonally differentiated properties of either HCs or afferents. Prior evidence includes data

from *Emx2* knockouts also exhibiting loss of the LPR without altering zoning (Jiang et al.,

2017) and data showing that inactivating a retinoic acid dehydrogenase deletes striolar

identity without affecting HC orientation (Ono et al., 2020). This conclusion stands in

contrast to zebrafish lateral line results, where transduction properties vary with HC

orientation depending on Emx2 (Kindig et al., 2023) and Gpr156 (this study).

Asymmetrical mechano-electrical transduction

Zebrafish lateral-line HCs use a combination of 2 mechanosensitive channels, Tmc2a

and Tmc2b (Chou et al., 2017). In the posterior lateral line, Emx2$^+$ HCs that sense

posterior flow rely on primarily on Tmc2b, while Emx2$^-$ HCs that sense anterior flow rely

on both Tmc2a and Tmc2b. Recently, a functional asymmetry was discovered between

anterior- and posterior-sensitive HCs: Emx2$^+$ HCs that rely primarily on Tmc2b have

smaller mechanosensitive responses compared to Emx2$^-$ HCs that rely on both Tmc2a

and Tmc2b. (Kindig et al., 2023). This work raises the possibility that the transcription

factor Emx2 inhibits *tmc2a* transcription to reduce mechanosensitive responses in HCs

sensing posterior flow. In our present study we find that functional asymmetry was also

lost in *gpr156* mutants (Fig. 5). In the context of mouse HC orientation, EMX2 is necessary

and sufficient to enrich and polarize GPR156 at HC apical junctions and reverse HC

orientation (Kindt et al., 2021). As the Gpr156 receptor is unlikely to act transcriptionally,

zebrafish Emx2 may act through Gpr156 to impact Tmc2a/2b function. For example,

polarization of Gpr156 at apical junctions may limit scaffolding or transport of Tmc

proteins to homomeric channels composed of Tmc2b. Overall, our work in zebrafish

indicates that both Emx2 and Gpr156 are important to dictate HC orientation and the

mechanosensitive properties of HCs with opposing orientations.

We did not see a clear impact of GPR156 loss on the transduction properties of *Emx2*[+]

HCs in the mouse utricle (Fig. 2-4). Subtle effects remain possible given the natural

variance in the data and low power of the negative results. Nevertheless, current results

suggest normal HC function in the *Gpr156* mouse mutant, a prerequisite to interrogate

how non-reversed HCs affects vestibular behavior.

Importance of the LPR for vestibular function

The brain combines information from the maculae and cristae with inputs from other

modalities, including the somatosensory and visual systems, to compute a representation

of our self-motion and spatial orientation (reviewed in (Cullen, 2019). Our present findings

demonstrate that GPR156 is *not* required for macular HCs to selectively contact different

afferents across the LPR. Accordingly, we predicted that HCs that fail to reverse in the

lateral utricle of *Gpr156* mutants would lead to deficits in paradigms that test macular

contributions to behavioral responses. Indeed, *Gpr156* mutant mice demonstrated

impaired performance on two tests that engage the otolith organs: swimming and off-

vertical-axis rotation (OVAR). First, during swimming, impaired spatial orientation results

from the combined altered otolith input and reduced somatosensory feedback due to the

aqueous environment. Second, *Gpr156* mutant mice displayed impaired sustained

responses to off-vertical axis rotation (OVAR). While initial eye movement response to

OVAR is mediated by stimulation of semicircular canal HCs, the canal response decays

such that sustained eye movement is solely driven by the otolith stimulation (Beraneck et

al., 2012; Ono et al., 2020). Quantification of the sustained responses revealed a marked

reduction in this otolith-mediated reflex response. Confirming that these deficits are

specific to otolith function, the *Gpr156* mutant mice did not show overt vestibular

dysfunction such as spinning or head tilt that can arise with global dysfunction of

vestibular epithelia. Further, they displayed normal performance during testing that

selectively engaged superior canal organs, namely angular VOR in the light and dark.

This latter finding was predicted; because HC orientation is unchanged in *Gpr156* mutant

cristae, the coding properties of semicircular canal afferents should be unchanged and

would therefore produce a normal angular VOR upon specific stimulation of canal

ampullae.

Comparing vestibular phenotypes of *Gpr156* and *Emx2* mutants

Gpr156 does not appear to be significantly expressed in the central nervous system according to the Allen Brain Atlas (Lein et al., 2007). In addition, *Advillin-Cre* tracing and dye-filling experiments (Fig. 6-7) showed that vestibular ganglion neurons still selectively innervated LES HCs even when lacking GPR156. We thus speculate that HC signals in the LES zone of *Gpr156* mutants are relayed to the cerebellum (Ji et al., 2022; Maklad et al., 2010) via the central projection of *Advillin*[+] neurons as usual, but with abnormal (non-reversed) response polarity. Overall, it appears unlikely that vestibular deficits observed in our constitutive *Gpr156* mouse mutants reflect defective neuronal activity in addition to a missing LPR. That said, we cannot entirely rule out another role of GPR156 beside its regulation of HC orientation. This could be addressed in a follow-up study by limiting *Gpr156* inactivation to HCs.

It is interesting to compare vestibular-related behaviors between the *Gpr156* and *Emx2* mouse models in light of established similarities and differences in protein distribution and function (Ji et al., 2022; Jiang et al., 2017; Kindt et al., 2021). Regarding behavior, this study and work by Ji et al., 2022 agree that vestibular deficits in each model are relatively mild (normal balance beam, VOR and OKR). *Emx2* inactivation in HCs (*Gfi1-Cre*) showed impaired swimming ability, but defects were less severe and different in nature compared to constitutive *Gpr156* mutants. *Gpr156* mutants displayed substantial difficulty in maintaining their orientation in water, and frequently rolled over such that they needed to be rescued. In contrast, *Emx2* mutants did not need rescue and were described as demonstrating "frantic" swimming behavior, spending more time trying to climb out of the water as compared to their littermate controls. OVAR highlighted otolith-specific defects in *Gpr156* mutants but was not performed in *Emx2* mutants.

GPR156 acts downstream of EMX2 to regulate hair bundle orientation, and perhaps mechanosensitivity in zebrafish. Another significant new result in the current study is to show that afferent projections segregate with HC orientation and not with zonal identity (Fig. 7; Advillin neuron labeling), yet segregation does not depend on HC orientation since

it is unchanged in mouse and zebrafish *Gpr156* mutants (Fig. 6-10). It follows that EMX2

must influence afferent segregation (Ji et al., 2022; Ji et al., 2018) via other effectors

unrelated to HC orientation. Accordingly, Ji and colleagues concluded that EMX2

regulates HC orientation and afferent patterning independently, with *Emx2* expression in

support cells critical for proper afferent contacts with HCs (Ji et al., 2022). This aligns well

with previous evidence that *Gpr156* is only expressed in HCs (Kindt et al., 2021).

Ji and colleagues also used specific inactivation of the transduction protein TMIE in the

LES (*Emx2-Cre*) to silence the *Emx2*[+] HC population. Interestingly, as with *Emx2* mutants,

they did not observe severe behavioral defects, and swimming ability in particular was

not obviously compromised in contrast to *Gpr156* mutants. It should be noted however

that while the *Gpr156 KO* and *Emx2-Cre; Tmie cKO* models both lack a normal

complement of bidirectional responses, they provide different forms of aberrant linear

acceleration signals. In the *Tmie* cKO model, LES HC inputs to the cerebellum are

missing but in the *Gpr156* model they are retained and of opposite nature since LES HCs

fail to reverse their orientation.

Several considerations could potentially explain why the *Gpr156* model appears to display

more severe behavioral defects compared to the *Emx2* models. First, HC or afferent

properties were not tested in *Emx2* mutants (Ji et al., 2022) and may be defective. Second,

compounded HC orientation and afferent defects in the *Emx2* mutants could dampen the

deleterious consequence of having LES HCs drive a neuronal response of reversed

polarity. Afferent defects in *Emx2* mutants were reported during embryogenesis (E16.5)

and innervation could further degenerate in adults, further dampening the consequence

of non-reversed HC activity in the LES. It is worth noting that aberrant vestibular afferent

signals are clinically more deleterious than absent signals, which perhaps explains why

properly relayed yet aberrant signals from LES HCs in *Gpr156* mutants result in

behavioral defects worse than observed in *Emx2* or *Emx2-Cre; Tmie cKO* models.

In conclusion, the *Gpr156* mutant mouse model is well-suited to specifically interrogate

how HC orientation reversal and the resulting LPR influence vestibular function because

it appears to lack confounding afferent patterning defects. In future studies, this model

could be improved by limiting *Gpr156* inactivation to HCs.

METHODS

Mouse strains and husbandry

The *Gpr156^del* strain (*B6N(Cg)-Gpr156tm1.1(KOMP)Vlcg/J*; MGI:5608696) was produced by the Knockout Mouse Project (KOMP) and studied previously in (Kindt et al., 2021). The *Advillin-Cre* strain is *B6.129P2-Avil^tm2(cre)Fawa/J* (MGI: 6196038) (Zhou et al., 2010). The *Tigre Ai140* strain is *B6.Cg-Igs7^tm140.1(tetO-EGFP,CAG-tTA2)Hze* (MGI: 5904001) (Daigle et al., 2018). Some whole-cell patch clamp recordings were obtained from CD1 mice (Charles River Laboratories USA). All animals were maintained in standard housing and all animal work was reviewed for compliance and approved by the Animal Care and Use Committees of The Jackson Laboratory (anatomical experiments), the University of Chicago (cellular electrophysiology and afferent labeling experiments), and the Johns Hopkins University School of Medicine (behavioral experiments).

Mouse immunofluorescence and imaging (Figure 1, Figure 7)

Temporal bones of adult animals were isolated, oval windows were cleared and the cochlear apex was punctured to allow better access to the fixative. Samples were then fixed in paraformaldehyde (PFA 4%) overnight at 4°C, rinsed in PBS and incubated overnight in 4% Ethylenediaminetetraacetic acid (EDTA) at room temperature for decalcification. Both the trio (utricle and crista) and the saccule were dissected from the bone and their epithelium exposed. After dissection, samples were permeabilized and blocked in PBS with 0.5% Triton-X100 and bovine serum albumin (1%) for at least at 1 hour at room temperature. For SPP1/Osteonpontin staining, temporal bones were immediately dissected to isolate the trio and the saccule and expose their epithelium, and an eyelash was used to scrape off the otoconia, Samples were then fixed in PFA 4% for 1h at room temperature before being permeabilized and blocked as described previously. Primary and secondary antibodies were each incubated overnight at 4°C in PBS, and fluorescent dye-conjugated phalloidin was added to secondary antibodies. Samples were washed 3 times in PBS + 0.05% Triton-X100 after each antibody incubation before being finally post fixed in PFA 4% for at least 1 hour at room temperature. Primary antibodies used were:

Mouse anti-MYO7A (Developmental Studies Hybridoma Bank/DSHB 138-1, 1:1000)

Goat anti-OCM (Invitrogen, PA547832, 1:200)

Goat anti-SPP1 (R&D Systems, AF808, 1:200)

Rabbit anti-PCNT (Biolegend, PRB432C, 1:400)

Goat anti-SOX2 (R&D Systems, AF2018, 1:200)

Secondary antibodies used were raised in donkey and coupled with AlexaFluor (AF) 488,

555, or 647 (donkey anti-rabbit 488 (A-21206), donkey anti-mouse 647 (A-31571), donkey

anti-goat 555 (A-21432); Thermofisher Scientific). Fluorescent conjugated phalloidin was

used to reveal F-actin (CF405 (89138-126); Biotum).

Confocal images were captured on a line scanning confocal microscope (LSM8000) using

Zen 2.6 software, the Airyscan detector in confocal mode, and either a 20× or a 63× 1.4

NA oil objective lens (Carl Zeiss AG). Images show a single optical plane unless stated

otherwise in the figure legend.

Mouse immunostain quantification (Figure 1, Figure 7)

All images were processed using Adobe Photoshop (CC2020), and the same image

treatment was applied across genotypes for each experiment. All quantifications included

at least three animals of each genotypes, and all values plotted in the study, as well as

animal cohort size (N), HC number and stereocilia number (n) are detailed in legends. To

quantify striolar surface, striolar density, as well as HC Type I and II density by region,

images were captured using a DM5500B fluorescence microscope using the Leica

Application Suite (LASX) and a 20x objective (Leica). Striolar surface was measured

using the polygon selection and area tools in Fiji/ImageJ to define the striola based on

the Oncomodulin staining. Striolar density was measured by using the cell counter tool in

Fiji/ImageJ to obtain the total number of Oconmodulin positive HCs. That number was

then normalized by the total striolar surface previously measured to obtain a final number

of HCs per $1000\mu m^2$. To measure Type I and Type II HC density by region, three regions

of interest (ROI) were defined centrally of either 130x50μm (utricle) or 150x40μm

(saccule). The lateral extrastriolar (LES) domain was placed 20μm inside the macula

starting from the lateral edge of the utricle, encompassing most of the region lateral to the

line of polarity reversal (LPR). The striolar domain was defined immediately adjacent to

the LES domain based on the LPR and oncomodulin signal in controls. The medial

extrastriolar (MES) domain started 50µm medial to the striolar domain. The anterior

extrastriolar domain (ANT) was placed 10µm from the anterior edge of the saccule. The

striolar domain was defined 20µm posterior to the ANT domain based on oncomodulin

staining in controls, and the posterior extrastriolar domain (POST) was defined 50µm

posterior to the striolar domain. The location of these domains is illustrated in Fig.1A. The

same fields and spacing were next used for the $Gpr156^{del/del}$ mutants. The cell counter

plugin in Fiji/ImageJ was used to count HCs in each ROI.

Data analysis and statistics – mouse anatomy (Figure 1, Figure 7)

All data were plotted in Prism 9 (GraphPad). Striolar surface, HC density and number of

Type I and type II HC density in each region of the maculae were plotted individually.

Their distribution was framed by 25-75% whisker boxes where exterior lines show the

minimum and the maximum, the middle line represents the median, and + represents the

mean. Difference of data distribution was tested using Mann-Whitney U (non-parametric,

unpaired t-test) for striolar surface and HC density, and 2-way ANOVA with Sidak's

multiple comparison for type I and type II HCs.

Mouse hair cell and afferent electrophysiology (Figures 2-4, Figure 8)

We followed procedures described in (Gonzalez-Garrido et al., 2021) to prepare and

record from HCs and calyceal afferent terminals in semi-intact utricles comprising the

sensory epithelium and attached distal vestibular nerve including the vestibular ganglion.

Recordings were analyzed for 83 cells from 57 $Gpr156$ heterozygotes and 93 cells from

73 $Gpr156$-null animals.

Preparation. Following protocols approved by the University of Chicago Animal Care and

Use Committee, mice were deeply anesthetized with gaseous isoflurane and then

decapitated to allow dissection of the utricles and attached nerve and ganglion. The

preparations were dissected in our *standard bath (external) solution*: cold Leibovitz's-L15

medium (L15, Gibco, #41300-021) supplemented with 10 mM HEPES (4-(2-

hydroxyethyl)-1-piperazineethanesulfonic acid) (~315 mmol/kg, pH 7.4). The utricles

were then treated with L15 containing proteinase XXIV (100 mg/ml, Sigma, St. Louis, MO

#P8038) for 11 min at room temperature to facilitate mechanical removal of the otolithic

membrane. The epithelium was mounted on a cover slip with glued glass fibers and

placed in a recording chamber on a microscope (Zeiss, Axio Examiner A1) equipped with

Nomarski and fluorescence optics.

Recording. For whole-cell recording, pipettes (R6, King Precision Glass) were heat-pulled

(PC100, Narishige) and parafilm-wrapped to reduce capacitance. Pipette resistances

were 4-5 MΩ for the standard internal and external solutions. Standard internal solution

contained (in mM): 135 KCl, 0.1 $CaCl_2$, 3.5 $MgCl_2$, 3 Na_2ATP, 5 creatine phosphate (Na

salt), 0.1 Na-cAMP, 0.1 Li-GTP, 5 EGTA, and 10 HEPES, plus ~28 mM KOH to bring pH

to 7.3 and osmolality to ~300 mmol/kg. Internal solution was supplemented with

sulforhodamine 101 (1 mg/100 ml; Thermo Fisher Scientific) for visualization of recorded

hair cells. The bath (external solution) was modified L15 described above and perfused

during the experiment.

We were able to locate LES, striola or MES despite loss of the LPR because anatomical

features of the striola are maintained in the otolith organs of $Gpr156^{del/del}$ mice (Kindt et

al., 2021), and some of these (hair bundle size, spacing of HCs, abundance of complex

calyces) are visible in live tissue with Nomarski optics. Usually we aimed for HCs or

calyces >50 μm away from zonal boundaries to avoid confusion over zone. In one

experimental series, however, we focused on the zonal boundary to see whether afferents

respected zonal boundaries in Gpr156-deleted mice.

To record in whole-cell mode from a specific HC or afferent calyx, we cleaned the cell

membrane by the outflow of pipette solution and applied gentle suction to promote giga-

seal formation and membrane rupture. Stimulus protocols and data acquisition were

implemented by the patch clamp amplifier (model EPC-10, HEKA Elektronik) controlled

by Patchmaster software. Data were digitized at sampling intervals of 10 or 20

microseconds. Pipette and membrane capacitive currents were nulled and series

resistance (R_s) was compensated by 80% on-line with the amplifier's controls. Voltages

were corrected offline for the liquid junction potential (–4 mV) and the voltage error due
to residual R_s (20% of total R_s, 8.1 ± 0.1, n = 207).

<u>Mechanical stimulation of mouse hair cells.</u> Hair bundles were deflected with a rigid probe
(pulled glass pipette, BF120-60-10, Sutter instrument Company) glued to a piezoelectric
bimorph ceramic and brought into contact with the staircase of stereocilia ('back' of the
bundle) at ~half-height. Displacement of the probe to driving voltage was calibrated with
a CCD camera, yielding a scale factor 330-410 nm/V, depending on the probe. Probe
motion recorded with a photodiode (PIN-6D, United Detector Technology) revealed a
primary resonance at 1900-2000 Hz. To attenuate this ringing, the voltage input to the
bimorph was low-pass filtered (8-pole Bessel filter, Model 900, Frequency Devices) set
at 1 kHz, for a step rise time (10-90%) of ~300 μs.

Mechanotransduction current was recorded in voltage-clamp mode with the holding
potential (HP) of –94 mV and –84 mV for type I and type II HCs, respectively. The larger
negative potential for type I HCs was chosen to reduce $g_{K,L}$ and so improve the voltage
clamp. Responses to three identical displacement step protocols were averaged and low-
pass filtered off line (corner frequency, 2 kHz, 8-pole Bessel filter, as above). Step
protocols comprised either 20 or 40 displacement steps iterated by 100 nm or 35 nm,
respectively, and beginning slightly negative to the resting bundle position. Transduction
sensitivity and adaptation was analyzed in 49 HCs with I_{max} > 150 pA (23 HCs from
heterozygotes and 26 HCs from null mutants), as described next.

Quantification of mouse electrophysiological data. Analysis of I_{MET} properties is
described elsewhere (Songer and Eatock, 2013; Vollrath and Eatock, 2003). Briefly,
conductance-displacement [G(X)] activation curves were calculated by dividing I_{MET}
(averaged across 2-4 presentations) by the driving force (HP – I_{MET} reversal potential, 0.2
mV) (Corns et al., 2014) and plotting G_{MET} against probe displacement (*X*). Probe
displacement as a function of driving voltage was calibrated off-line with a photodiode
(Songer and Eatock, 2013). G(X) curves were averaged across HCs and fitted with a
Boltzmann function (Eq. 1):

$$G_{MET}(X) = \frac{G_{max} - G_{min}}{1 + e^{(\frac{X_{1/2}-X}{S})}} + G_{min} \quad (1)$$

where G_{max} and G_{min} are maximum and minimum MET conductance, respectively, $X_{1/2}$ is

displacement that evokes half-maximal G_{MET} and S is displacement corresponding to an

e-fold rise in G_{MET}. We used the Boltzmann fits to calculate operating range (OR):

displacement range corresponding to 10 - 90% of the response range, $G_{max} - G_{min}$.

We characterized transducer adaptation for responses at $X_{1/2}$, shown previously to reveal

both fast and slow components of adaptation in mouse utricular hair cells (Vollrath and

Eatock 2003). To reduce noise, current responses were averaged (*n* usually 3) and low-

pass filtered at 1.5 kHz. Fast and slow adaptation components were estimated from

double-exponential fits of the first 300 ms of the step response (Eq. 2). Deviation of those

fits at the onset response signaled a very fast transducer adaptation component, which

was separately estimated with a single-exponential fit of the first 1.5 ms of the response

(Eq. 3).

$$I(t) = A_{f}e^{-\frac{t}{\tau_f}} + A_{s}e^{-\frac{t}{\tau_s}} + I_{ss} \quad (2)$$

$$I(t) = A_{vf}e^{-\frac{t}{\tau_{vf}}} + I_{ss} \quad (3)$$

Here τ_{vf}, τ_f and τ_s are the very short (1 ms or less), short (from 1 to 10 ms) and long (>10

ms) time constants, respectively; A_{vf}, A_f and A_s are the amplitudes of the corresponding

exponential terms; and I_{SS} is the steady-state current. All HCs had a very fast or fast

component or both; some lacked a slow component that was detectable in the 400-ms

step.

We calculated extent of adaptation (% decay) at $X_{1/2}$ as follows (Eq. 4):

$$900 \qquad Extent\ of\ adaptation = \left(\frac{I_{peak} - I_{ss}}{I_{peak} - I_0}\right) \times 100 \quad (4)$$

where I_0, I_{peak}, and I_{SS} are pre-step, steady-state, and peak currents, respectively.

In mouse utricular HCs, the dominant voltage-gated currents are carried by outwardly

rectifying K_V channels (Martin et al., 2023). By the end of the first postnatal week, these

currents provide distinctive electrophysiological signatures in type I and type II HCs, which

include more negatively activating K^+ currents and more negative resting potentials for

type I HCs (Rusch and Eatock, 1996; Rusch et al., 1998). To characterize the voltage

dependence of the currents, we applied families of iterated test voltage steps (400 ms

duration) from holding potentials of –74 mV (type I HCs) and –64 mV (type II HCs). The

different holding potentials reflect the difference in resting potentials of the two HC types.

Steady-state voltage dependence was measured by constructing conductance-voltage

[G(V)] relations from tail currents, measured 1 ms after the 400-ms test voltage steps, at

a common membrane voltage of –39 mV (chosen to reduce contamination of tail currents

by HCN currents). Tail current was converted to "tail conductance" by dividing by driving

force ($V_m - V_{rev}$), where V_{rev} is approximated by the K^+ equilibrium potential (–84 mV in

our solutions). Tail conductance was plotted against the iterated test step voltage and fit

with the Boltzmann function:

$$920 \qquad G(V) = \frac{G_{max} - G_{min}}{1 + e^{(V_{1/2} - V)/S}} + G_{min} \quad (5)$$

where $V_{1/2}$ is the voltage of half-maximal activation and S is the voltage change over

which G changes e-fold at voltages negative to $V_{1/2}$. Voltage was corrected for voltage

errors caused by residual R_S and for the liquid junction potential by subtracting 4 mV.

Input resistance (R_{in}) was estimated from current clamp data as the slope of a linear

regression of the $V_m(I)$ relation for small currents around resting potential.

Imaging of afferent terminal arbors in mouse utricle (Figure 6)

To investigate whether afferent receptive fields (terminal arbors) were disrupted by the
lack of bundle reversal in the lateral extrastriola (LES), we recorded in whole-cell mode
from calyces near the LES-striolar border. To label the terminal arbors for imaging, we
included Alexa Fluor 594 (#A10438 Thermo Fisher Scientific) instead of sulforhodamine
in the pipette solution (50 micromolar). After whole-cell recording was achieved, we
waited 20 min as the fluorescent pipette solution diffused into the calyx and throughout
the terminal arbor, then removed the pipette and fixed the utricular epithelium with 4%
paraformaldehyde for 10 min at room temperature (~22°). Fixed utricles were immersed
in blocking buffer (PBS with 4% normal donkey serum and 0.2% triton) then treated with
primary antibodies diluted with blocking buffer overnight at 4°C. Primary antibodies were:
mouse monoclonal anti-βII spectrin (1:500; BD Transduction, #612562) , to label hair
bundles and show the location of the kinocilium by absence of label; rabbit polyclonal
anti-calbindin (1:250; Invitrogen, #711443) to label the striola and juxtastriola (Leonard
and Kevetter, 2002). Specimens were then washed extensively and incubated with
secondary antibodies conjugated with fluorescent proteins (donkey anti-mouse or rabbit
IgG (H + L) antibody, ThermoFisher Scientific) for 1 hr at 4°C. Stained specimens were
mounted whole in mounting medium (Vector Laboratories, #H-1500), imaged with
spinning-disc confocal microscopy (Marianas SDC 3i), and analyzed with Adobe
Photoshop 2021 and FluoRender (ver. 2.26.3).

Measurement of excitability from mouse utricular afferents (Figure 8)
To document excitability of afferents in the LES, we recorded from calyces in whole-cell
current clamp mode, injected current steps and noted the thresholds for spiking and the
firing patterns. Most LES afferents are dimorphic, forming both calyces and boutons on
type I and II HCs, respectively. We rejected calyces with resting potentials less negative
than −50 mV. We evoked spikes with current steps to determine if afferents had step-
evoked firing patterns consistent with expected zonal patterns of spiking regularity in
mammalian vestibular epithelia (Goldberg et al., 1990; Iwasaki et al., 2008; Kalluri et al.,
2010). Families of 500-ms current steps were applied from zero-current potential,
beginning at −200 pA and incremented by 50 pA. Threshold current (I_{thresh}) was taken as
the minimum current (±50 pA) required to evoke any spikes.

We classified neuronal firing patterns according to the number of spikes evoked by a 400-
ms step at $I_m = 3 \times I_{thresh}$ (Ono et al., 2020): a pattern of 1-2 spikes at step onset was
considered 'transient'; if more spikes were evoked, the pattern was 'sustained'. These
patterns have been associated with the mechanisms that give rise to well-known
differences in the regularity of firing across zones. The transient step-evoked pattern is
associated with striolar, irregular afferents and the sustained step-evoked pattern with
extrastriolar, regular afferents. The difference in step-evoked firing patterns is attributed
to zonal differences in expression of K_V channels that influence spike regularity (Iwasaki
et al., 2008; Kalluri et al., 2010) and, more importantly, encoding strategy of afferent fibers
(Jamali et al., 2019).

Data analysis and statistics – mouse electrophysiology (Figures 2-4, Figure 8)
Data analysis, statistical tests, curve fitting and graphing were done with OriginPro (2022;
OriginLab, Northampton, MA), Clampfit (Molecular Probes) and Matlab (Mathworks,
Natick, MA). For samples that had homogeneous variance, significance was tested with
either a two-tailed t-test or a one-way ANOVA with post-hoc Tukey test. When samples
had unequal variance, we used either Welch's t-test or Kruskal-Wallis ANOVA with post-
hoc Mann-Whitney U test. For significant differences ($p < 0.05$), we calculated effect size
with Hedge's g equation. For non-significant (NS) differences, we calculated statistical
power *post hoc* (OriginPro 2022) and these values are entered in tables along with p-
values.

Quantification of vestibular reflexes in mice (Figures 11-12)
We tested the angular vestibulo-ocular reflex (aVOR), the optokinetic reflex (OKR), and
the otolith vestibulo-ocular reflex (off-vertical axis rotation, OVAR). Specifically, aVOR
tests semicircular canal function, whereas OVAR tests otolith function. For completeness
we also tested each animal's OKR as a control of the integrity of visuo-motor function.

aVOR and OKR. Surgical techniques and experimental setup have been previously
described (Beraneck and Cullen, 2007). Eye movement data were collected using an

infrared video system (ETL-200, ISCAN system). The rotational velocity of the turntable

(head velocity) was measured using a MEMS sensor (MPU-9250, SparkFun Electronics).

Eye movements during the OKR were evoked by sinusoidal rotations of a visual surround

(vertical black and white stripes, visual angle width of 5°) placed around the turntable at

frequencies 0.2, 0.4, 0.8, 1, 2, and 3 Hz with peak velocities of ±16°/s. To record VOR

responses, the turntable was rotated at sinusoidal frequencies 0.2, 0.4, 0.8, 1, 2, and 3

Hz with peak velocities of ±16°/s in both light and dark. In the light condition, the visual

surround remained stationary, whereas, in the dark condition, both the visual surround

and turntable rotated in phase and the surroundings was made dark. Head and eye

movement signals were low-passed filtered at 125 Hz and sampled at 1 kHz. Eye position

data were differentiated to obtain velocity traces. Cycles of data with quick phases were

excluded from the analysis. Least-square optimization determined the VOR and OKR

gains, and phases plotted as mean ± standard error of the mean (SEM) against all

frequencies for all mice.

1005

OVAR. Techniques employed for measurement of eye movements during OVAR in alert

mice were described elsewhere (Beraneck et al., 2012). Briefly, recordings made after

fixating mice on a rotating platform, which was tilted 17° with respect to the ground. The

platform speed was increased from 0 to 50°/s in 500 ms and maintained its constant

velocity for 72 s (10 complete rounds) before being stopped. Eye movements were

measured using the same Infrared video system (ETL-200, ISCAN system) used for

angular VOR and OKR recordings. Quick phases were identified as previously described

and excluded from subsequent analysis. We then estimated the time constant of the

OVAR slow-phase eye velocity response decay, as well as the amplitude and phase of

its sinusoidal modulation using a linear regression approach.

1016

Mouse swimming assessment

A large container (26.25 × 16.25 × 14.38′) was filled with water (24–26°C) at the height

of at least 15 cm. The mouse was placed into the water and observed for up to 1 min. Its

performance was rated using the following scale (Hardisty-Hughes et al., 2010):

0 = swims, body elongated, and tail propels in flagella-like motion.

1 = immobile floating.

2 = underwater tumbling.

In a time-to-rescue trial, if an animal exhibited distressed swimming (underwater

tumbling) before 60 seconds, the animal was rescued and time of rescue was marked. If

not, the trial was marked as lasting 60 seconds.

1027

1028   **Mouse balance beam assessment**

1029   A 6-mm-wide and 40-cm-long beam was used for balance beam testing. The mice

1030   traversed 40 cm to a dark box. Walking speed was measured by recording the time the

1031   mouse took to reach the goal box from the opposite end of the beam. Mice were scored

1032   "time out" if they failed to reach the endpoint in 2 min.

1033

Air Righting, Tailing Hanging, and Contact Inhibition of Righting Tests

Air Righting Test

The mouse was picked up by the tail and lowered into a container (size of container?) so

that all four feet touched the bottom. The container was quickly inverted at a height of

30–40 cm so mouse fell supine onto a foam cushion below. How the mouse landed was

observed and rated as below (Rabbath et al., 2001):

0 = the animal lands on its feet (normal).

1 = the animal lands on its side (mild deficit).

2 = the animal lands on its back (severe deficit).

1043

Tail Hanging Test

The animal was picked up by the tail and lowered to an even surface, and its posture

rated using the following scale (Rabbath et al., 2001):

0 = straight body posture with extension of forelimbs toward the earth (normal).

1 = slightly bending the body ventrally (intermediate response).

2 = persistently bending the body (severe response).

1050

Contact Inhibition of Righting Test

The mouse was picked up by the tail and lowered into a small container so that all four

feet were in contact with the bottom, then the top of the container closed in contact with

the mouse's back. The container was then quickly inverted so that the mouse became

supine while the surface remains in contact with the soles of the mouse's feet. The

mouse's reflex was rated using the following scale (Rabbath et al., 2001):

0 = animal rights successfully (normal).

1 = partial righting (intermediate response).

2 = complete loss of righting (severe response).

1060

Quantification of Tremor during Resting Head Movement (Supplemental fig. 2)

Mice were placed in a cylinder (9 cm diameter and 21.5 cm height) that limited their

motion, so mice maintained their steady posture. Head movements in six dimensions

were recorded for 2 minutes using a miniature head motion sensor (MPU-9250,

SparkFun Electronics, Niwot, CO, United States) affixed on the top of the skull, which

comprises a three-dimensional (3D) accelerometer (measures linear acceleration;

fore/aft, lateral, and vertical) and 3D gyroscope (measures angular velocity: roll, pitch,

and yaw). Data was acquired at 200 Hz using windows-based CoolTerm software. We

then computed the power spectral densities (pwelch function, MATLAB, MathWorks)

using Welch's averaged periodogram with nfft = 4,096 and a Bartlett window (4,096 ms

duration) for all six dimensions of movement.

1072

Data analysis and statistics – mouse behavior (Figures 11-12)

Data are reported as the mean ± SEM. Nonparametric Mann-Whitney U-test was

performed to test significance for time to traverse (balance beam) and time to rescue

(swimming) between two groups. Two-way repeated-measures ANOVA followed by

Bonferroni post hoc comparison tests was used for the VOR data across frequencies. For

power spectra analysis we used an independent sample permutation test to test

significant differences between the two groups. Prism 9 (GraphPad) or MATLAB was

used for statistical analyses. The same independent sample permutation test was also

used to test significance for bias and gain of OVAR responses.

1082

Zebrafish strains and husbandry

Zebrafish (*Danio rerio*) were grown at 30 °C using a 14 hr light, 10 hr dark cycle. Larvae

were raised in E3 embryo medium (5 mM NaCl, 0.17 mM KCl, 0.33 mM $CaCl_2$, and

0.33 mM $MgSO_4$, pH 7.2). Larvae were examined at 5-6 days post fertilization (dpf). The

following previously established mutant and transgenic zebrafish strains were used in this

study: *gpr156[idc15]*, *emx2[idc5]* (Jiang et al., 2017; Kindt et al., 2021),

Tg(myo6b:memGCaMP6s[idc1] (Kindt et al., 2021). For comparisons, *emx2* mutants were

compared to wild-type or *emx2[+/-]* animals. *Gpr156* mutants were compared to *gpr156[+/-]*

animals. Zebrafish work was performed at the National Institute of Health (NIH) and

approved by the Animal Use Committee at the NIH under animal study protocol #1362-

13.

1094

Zebrafish immunofluorescence and imaging (Figures 5, Figures 9-10)

Immunohistochemistry was performed on whole larvae at 5 dpf. Whole larvae were fixed

with 4% paraformaldehyde in PBS at 4°C for 3.5 hr. For pre- and post-synaptic labeling

all wash, block and antibody solutions were prepared with 0.1% Tween in PBS (PBST).

For Emx2 labeling performed on sparse afferent labeling (see below) all wash, block and

antibody solutions were prepared with PBS + 1% DMSO, 0.5% Triton-X100, 0.1% Tween-

20 (PBDTT). After fixation, larvae were washed 4 × 5 min in PBST or PBDTT. For synaptic

labeling, larvae were permeabilized with Acetone. For this permeabilization larvae were

washed for 5 min with H_2O. The H_2O was removed and replaced with ice-cold acetone

and placed at −20°C for 5 min, followed by a 5 min H_2O wash. The larvae were then

washed for 4 × 5 min in PBST. For all immunostains larvae were blocked overnight at 4°C

in blocking solution (2% goat serum, 1% bovine serum albumin, 2% fish skin gelatin in

PBST or PBDTT). Larvae were then incubated in primary antibodies in antibody solution

(1% bovine serum albumin in PBST or PBDTT) overnight, nutating at 4°C. The next day,

the larvae were washed for 4 × 5 min in PBST or PBDTT to remove the primary antibodies.

Secondary antibodies in antibody solution were added and larvae were incubated for 2

hrs at room temperature, with minimal exposure to light. Secondary antibodies were

washed out with PBST or PBDTT for 4 × 5 min. Larvae were mounted on glass slides

with Prolong Gold (ThermoFisher Scientific) using No. 1.5 coverslips. Primary antibodies

used were:

Rabbit anti-Myo7a (Proteus 25-6790; 1:1000)

Mouse anti-Ribeye b (IgG2a) (Sheets et al., 2011)

Mouse anti-pan-MAGUK (IgG1) (Millipore MABN7; 1:500)

Mouse anti-Myo7a (DSHB 138-1; 1:500)

Rabbit anti-Emx2 (Trans Genic KO609; 1:250).

The following secondaries were used at 1:1000 for synaptic labeling: goat anti-rabbit

Alexa 488, goat anti-mouse IgG2a Alexa 546, goat anti-mouse IgG1 Alexa 647, along

with Alexa 488 Phalloidin (Thermofischer; #A-11008, #A-21133, #A-21240, #A12379).

For Emx2 co-labeling the following secondaries were used at 1:1000: goat anti-mouse

Alexa 488, and goat ant-rabbit Alexa 647, along with Alexa 488 Phalloidin (Thermofischer;

#A12379, #A28175, #A27040).

Fixed samples were imaged on an inverted Zeiss LSM 780 laser-scanning confocal

microscope with an Airyscan attachment (Carl Zeiss AG, Oberkochen, Germany) using

an 63 × 1.4 NA oil objective lens. Airyscan z-stacks were acquired every 0.18 µm. The

Airyscan Z-stacks were processed with Zen Black software v2.1 using 2D filter setting of

6.0. Experiments were imaged with the same acquisition settings to maintain consistency

between comparisons. Processed imaged were further processed using Fiji.

1132

Zebrafish immunostain quantification (Figure 5, Figures 9-10)

Images were processed in ImageJ. Researcher was not blinded to genotype. Hair bundle

orientation was scored relative to the midline of the muscle somites. HC number per

neuromast were quantified based on Myo7a labeling and presence of a paired/complete

synapse. For quantification of Emx2 labeling, HCs were scored as Emx2 positive if they

labeled with both Emx2 and Myo7a. To quality as a ribbon or presynapse, the following

minimum size filters were applied to images: Ribeye b: 0.025 μm^2, MAGUK: 0.04 μm^2. A

complete synapse was comprised of both a Ribeye b and MAGUK puntca. An unpaired

presynapse consisted of only a Ribeye b puntca, while an unpaired postsynapse

consisted of only a MAGUK puncta. In each neuromast all HCs (~15 HC per neuromast)

were examined for our quantifications.

1144

Zebrafish functional calcium imaging in hair bundles (Figure 5)

GCaMP6s-based calcium imaging in zebrafish hair bundles has been previously
described (Lukasz and Kindt, 2018). Briefly, individual 5-6 dpf larvae were first
anesthetized with tricaine (0.03% Ethyl 3-aminobenzoate methanesulfonate salt,
SigmaAldrich). To restrain larvae, they were then pinned to a Sylgard-filled recording
chamber. To suppress the movement, alpha-bungarotoxin (125 µM, Tocris) was injected
into the heart. Larvae were then rinsed and immersed in extracellular imaging solution (in
mM: 140 NaCl, 2 KCl, 2 CaCl$_2$, 1 MgCl$_2$ and 10 HEPES, pH 7.3, OSM 310 +/- 10) without
tricaine. A fluid jet was used to mechanically stimulate the apical bundles of HCs of the
A-P neuromasts of the primary posterior lateral-line. To stimulate the two orientations of
HCs (A>P and P>A) a 500 ms 'push' was delivered. Larvae were rotated 180° to deliver
a comparable 'push' stimulus to both the A>P and P>A HCs.

1157

To image calcium-dependent mechanosensitive responses in apical hair bundles, a
Bruker Swept-field confocal system was used. The Bruker Swept-field confocal system
was equipped with a Rolera EM-C2 CCD camera (QImaging) and a Nikon CFI Fluor
60✕ 1.0 NA water immersion objective. Images were acquired in 5 planes along the Z-
axis at 0.5 µm intervals (hair bundles) at a 50 Hz frame rate (10 Hz volume rate). The 5
plane Z-stacks were projected into one plane for image processing and quantification.
The method to create spatial ΔF heatmaps has been described (Lukasz and Kindt, 2018).
For GCaMP6s measurements, a circular ROI with a ~1.5 µm (hair bundles) diameter was
placed on the center of each individual bundle. The mean intensity ($\Delta F/F_0$) within each
ROI was quantified. F_0 represents the GCaMP6s intensity prior to stimulation. We
examined the GCaMP6s signal in each hair bundle to determine its orientation. The
GCaMP6s responses for each neuromast were averaged to quantify the magnitude of the
A>P and P>A responses.

1171

Zebrafish sparse labeling of single afferents in the lateral-line (Figure 9)
To visualize the innervation pattern of single afferent neurons, a *neuroD1:tdTomato*
plasmid was injected into zebrafish embryos at the 1-cell stage. This plasmid consists of
a 5kb minimal promoter, *neurod1*, that drives tdTomato expression in lateral-line afferents
(Ji et al., 2018). This plasmid was pressure at a concentration of 30 ng/µl. At 3 dpf larvae

were anesthetized with 0.03% ethyl 3-aminobenzoate methanesulfonate (Tricaine), to

screen for tdTomato expression. Larvae were screened for mosaic expression of

tdTomato expression in the lateral-line afferents using a Zeiss SteREO Discovery V20

microscope (Carl Zeiss) with an X-Cite 120 external fluorescent light source (EXFO

Photonic Solutions Inc). After selecting larvae with tdTomato expression, larvae were

prepared for immunostaining at 5 dpf, and imaged as outlined above.

1183

1184 Zebrafish data analysis and statistics

Statistical analyses and data plots were performed with Prism 9 (Graphpad, San Diego,

CA). Values of data with error bars on graphs and in text are expressed as mean ± SEM

unless indicated otherwise. All experiments were performed on a minimum of 3 animals,

6 neuromasts (primary posterior lateral-line neuromasts with A-P orientations: L1-L5). For

5-6 dpf larvae, each neuromast represents analysis from 12 to 20 HCs and 41–68

synapses. All replicates are biological. No animals or samples were excluded from our

analyses unless control experiments failed–in these cases all samples were excluded.

No randomization or blinding was used for our animal studies. Where appropriate, data

was confirmed for normality using a D'Agostino-Pearson normality test. Statistical

significance between conditions was determined by an unpaired t-test.

1195

1196

REFERENCES

Beraneck, M., Bojados, M., Le Seac'h, A., Jamon, M. and Vidal, P. P. (2012). Ontogeny of mouse vestibulo-ocular reflex following genetic or environmental alteration of gravity sensing. *PLoS One* **7**, e40414.

Chitnis, A. B., Nogare, D. D. and Matsuda, M. (2012). Building the posterior lateral line system in zebrafish. *Dev Neurobiol* **72**, 234-255.

Chou, S. W., Chen, Z., Zhu, S., Davis, R. W., Hu, J., Liu, L., Fernando, C. A., Kindig, K., Brown, W. C., Stepanyan, R., et al. (2017). A molecular basis for water motion detection by the mechanosensory lateral line of zebrafish. *Nature communications* **8**, 2234.

Corns, L. F., Johnson, S. L., Kros, C. J. and Marcotti, W. (2014). Calcium entry into stereocilia drives adaptation of the mechanoelectrical transducer current of mammalian cochlear hair cells. *PNAS* **111**, 14918-14923.

Cullen, K. E. (2019). Vestibular processing during natural self-motion: implications for perception and action. *Nat Rev Neurosci* **20**, 346-363.

Daigle, T. L., Madisen, L., Hage, T. A., Valley, M. T., Knoblich, U., Larsen, R. S., Takeno, M. M., Huang, L., Gu, H., Larsen, R., et al. (2018). A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced Brain-Cell-Type Targeting and Functionality. *Cell* **174**, 465-480 e422.

Deans, M. R. (2013). A balance of form and function: planar polarity and development of the vestibular maculae. *Seminars in cell & developmental biology* **24**, 490-498.

Deans, M. R., Antic, D., Suyama, K., Scott, M. P., Axelrod, J. D. and Goodrich, L. V. (2007). Asymmetric distribution of prickle-like 2 reveals an early underlying polarization of vestibular sensory epithelia in the inner ear. *J Neurosci* **27**, 3139-3147.

Denman-Johnson, K. and Forge, A. (1999). Establishment of hair bundle polarity and orientation in the developing vestibular system of the mouse. *J Neurocytol* **28**, 821-835.

Eatock, R. A. and Songer, J. E. (2011). Vestibular hair cells and afferents: two channels for head motion signals. *Annu Rev Neurosci* **34**, 501-534.

Fernandez, C., Goldberg, J. M. and Abend, W. K. (1972). Response to static tilts of peripheral neurons innervating otolith organs of the squirrel monkey. *J Neurophysiol* **35**, 978-987.

Fernandez, C., Goldberg, J. M. and Baird, R. A. (1990). The vestibular nerve of the chinchilla. III. Peripheral innervation patterns in the utricular macula. *J Neurophysiol* **63**, 767-780.

Flock, A. (1964). Structure of the Macula Utriculi with Special Reference to Directional Interplay of Sensory Responses as Revealed by Morphological Polarization. *J Cell Biol* **22**, 413-431.

Goldberg, J. M. (1991). The vestibular end organs: morphological and physiological diversity of afferents. *Current opinion in neurobiology* **1**, 229-235.

---- (2000). Afferent diversity and the organization of central vestibular pathways. *Exp Brain Res* **130**, 277-297.

Goldberg, J. M., Desmadryl, G., Baird, R. A. and Fernandez, C. (1990). The vestibular nerve of the chinchilla. V. Relation between afferent discharge properties and peripheral innervation patterns in the utricular macula. *J Neurophysiol* **63**, 791-804.

Gonzalez-Garrido, A., Pujol, R., Lopez-Ramirez, O., Finkbeiner, C., Eatock, R. A. and Stone, J. S.
(2021). The Differentiation Status of Hair Cells That Regenerate Naturally in the
Vestibular Inner Ear of the Adult Mouse. *J Neurosci* **41**, 7779-7796.
Holley, M., Rhodes, C., Kneebone, A., Herde, M. K., Fleming, M. and Steel, K. P. (2010). Emx2
and early hair cell development in the mouse inner ear. *Dev Biol* **340**, 547-556.
Iwasaki, S., Chihara, Y., Komuta, Y., Ito, K. and Sahara, Y. (2008). Low-voltage-activated
potassium channels underlie the regulation of intrinsic firing properties of rat vestibular
ganglion cells. *J Neurophysiol* **100**, 2192-2204.
Jacobo, A., Dasgupta, A., Erzberger, A., Siletti, K. and Hudspeth, A. J. (2019). Notch-Mediated
Determination of Hair-Bundle Polarity in Mechanosensory Hair Cells of the Zebrafish
Lateral Line. *Current biology : CB* **29**, 3579-3587 e3577.
Jamali, M., Carriot, J., Chacron, M. J. and Cullen, K. E. (2019). Coding strategies in the otolith
system differ for translational head motion vs. static orientation relative to gravity. *eLife*
8.
Ji, Y. R., Tona, Y., Wafa, T., Christman, M. E., Tourney, E. D., Jiang, T., Ohta, S., Cheng, H.,
Fitzgerald, T., Fritzsch, B., et al. (2022). Function of bidirectional sensitivity in the otolith
organs established by transcription factor Emx2. *Nature communications* **13**, 6330.
Ji, Y. R., Warrier, S., Jiang, T., Wu, D. K. and Kindt, K. S. (2018). Directional selectivity of
afferent neurons in zebrafish neuromasts is regulated by Emx2 in presynaptic hair cells.
eLife **7**.
Jiang, T., Kindt, K. and Wu, D. K. (2017). Transcription factor Emx2 controls stereociliary bundle
orientation of sensory hair cells. *eLife* **6**.
Jones, C., Qian, D., Kim, S. M., Li, S., Ren, D., Knapp, L., Sprinzak, D., Avraham, K. B.,
Matsuzaki, F., Chi, F., et al. (2014). Ankrd6 is a mammalian functional homolog of
Drosophila planar cell polarity gene diego and regulates coordinated cellular orientation
in the mouse inner ear. *Dev Biol* **395**, 62-72.
Kalluri, R., Xue, J. and Eatock, R. A. (2010). Ion channels set spike timing regularity of
mammalian vestibular afferent neurons. *J Neurophysiol* **104**, 2034-2051.
Kindig, K., Stepanyan, R., Kindt, K. S. and McDermott, B. M., Jr. (2023). Asymmetric
mechanotransduction by hair cells of the zebrafish lateral line. *Current biology : CB* **33**,
1295-1307 e1293.
Kindt, K. S., Akturk, A., Jarysta, A., Day, M., Beirl, A., Flonard, M. and Tarchini, B. (2021).
EMX2-GPR156-Galphai reverses hair cell orientation in mechanosensory epithelia.
Nature communications **12**, 2861.
Kozak, E. L., Palit, S., Miranda-Rodriguez, J. R., Janjic, A., Bottcher, A., Lickert, H., Enard, W.,
Theis, F. J. and Lopez-Schier, H. (2020). Epithelial Planar Bipolarity Emerges from Notch-
Mediated Asymmetric Inhibition of Emx2. *Current biology : CB* **30**, 1142-1151 e1146.
Lein, E. S., Hawrylycz, M. J., Ao, N., Ayres, M., Bensinger, A., Bernard, A., Boe, A. F., Boguski,
M. S., Brockway, K. S., Byrnes, E. J., et al. (2007). Genome-wide atlas of gene expression
in the adult mouse brain. *Nature* **445**, 168-176.
Leonard, R. B. and Kevetter, G. A. (2002). Molecular probes of the vestibular nerve. I.
Peripheral termination patterns of calretinin, calbindin and peripherin containing fibers.
Brain Res **928**, 8-17.

Li, A., Xue, J. and Peterson, E. H. (2008). Architecture of the mouse utricle: macular
organization and hair bundle heights. *J Neurophysiol* **99**, 718-733.
Lindeman, H. H. (1969). Regional differences in structure of the vestibular sensory regions. *The*
Journal of laryngology and otology **83**, 1-17.
Lopez-Schier, H., Starr, C. J., Kappler, J. A., Kollmar, R. and Hudspeth, A. J. (2004). Directional
cell migration establishes the axes of planar polarity in the posterior lateral-line organ of
the zebrafish. *Dev Cell* **7**, 401-412.
Lozano-Ortega, M., Valera, G., Xiao, Y., Faucherre, A. and Lopez-Schier, H. (2018). Hair cell
identity establishes labeled lines of directional mechanosensation. *PLoS biology* **16**,
e2004404.
Maklad, A., Kamel, S., Wong, E. and Fritzsch, B. (2010). Development and organization of
polarity-specific segregation of primary vestibular afferent fibers in mice. *Cell and tissue*
research **340**, 303-321.
Martin, H. R., Lysakowski, A. and Eatock, R. A. (2023). The potassium channel subunit K(V)1.8
(Kcna10) is essential for the distinctive outwardly rectifying conductances of type I and II
vestibular hair cells. *bioRxiv*.
McInturff, S., Burns, J. C. and Kelley, M. W. (2018). Characterization of spatial and temporal
development of Type I and Type II hair cells in the mouse utricle using new cell-type-
specific markers. *Biol Open* **7**.
Mirkovic, I., Pylawka, S. and Hudspeth, A. J. (2012). Rearrangements between differentiating
hair cells coordinate planar polarity and the establishment of mirror symmetry in
lateral-line neuromasts. *Biol Open* **1**, 498-505.
Ono, K., Keller, J., Lopez Ramirez, O., Gonzalez Garrido, A., Zobeiri, O. A., Chang, H. H. V.,
Vijayakumar, S., Ayiotis, A., Duester, G., Della Santina, C. C., et al. (2020). Retinoic acid
degradation shapes zonal development of vestibular organs and sensitivity to transient
linear accelerations. *Nature communications* **11**, 63.
Rusch, A. and Eatock, R. A. (1996). A delayed rectifier conductance in type I hair cells of the
mouse utricle. *J Neurophysiol* **76**, 995-1004.
Rusch, A., Lysakowski, A. and Eatock, R. A. (1998). Postnatal development of type I and type II
hair cells in the mouse utricle: acquisition of voltage-gated conductances and
differentiated morphology. *J Neurosci* **18**, 7487-7501.
Sheets, L., Trapani, J. G., Mo, W., Obholzer, N. and Nicolson, T. (2011). Ribeye is required for
presynaptic Ca(V)1.3a channel localization and afferent innervation of sensory hair cells.
Development **138**, 1309-1319.
Simmons, D. D., Tong, B., Schrader, A. D. and Hornak, A. J. (2010). Oncomodulin identifies
different hair cell types in the mammalian inner ear. *J Comp Neurol* **518**, 3785-3802.
Songer, J. E. and Eatock, R. A. (2013). Tuning and timing in mammalian type I hair cells and
calyceal synapses. *J Neurosci* **33**, 3706-3724.
Tarchini, B. (2021). A Reversal in Hair Cell Orientation Organizes Both the Auditory and
Vestibular Organs. *Front Neurosci* **15**, 695914.
Ventura, C. M. and Kalluri, R. (2019). Enhanced Activation of HCN Channels Reduces Excitability
and Spike-Timing Regularity in Maturing Vestibular Afferent Neurons. *J Neurosci* **39**,
2860-2876.

Vollrath, M. A. and Eatock, R. A. (2003). Time course and extent of mechanotransducer
adaptation in mouse utricular hair cells: comparison with frog saccular hair cells. *J*
Neurophysiol **90**, 2676-2689.
Xue, J. and Peterson, E. H. (2006). Hair bundle heights in the utricle: differences between
macular locations and hair cell types. *J Neurophysiol* **95**, 171-186.
Zhou, X., Wang, L., Hasegawa, H., Amin, P., Han, B. X., Kaneko, S., He, Y. and Wang, F. (2010).
Deletion of PIK3C3/Vps34 in sensory neurons causes rapid neurodegeneration by
disrupting the endosomal but not the autophagic pathway. *PNAS* **107**, 9424-9429.
1332
1333

Figure 1. Normal striola and hair cell type organization in *Gpr156* mutants. A)
Diagrams showing general macular organization and the regions analyzed in **B-I** (LES,
MES: lateral and medial extrastriola, respectively; ANT: anterior, POST: posterior). Each
region is 130x50 µm in the utricle and 150x40 µm in the saccule. White chevrons indicate
HC orientation. **B-C**) P28 utricles (**B**) and saccules (**C**) immunolabeled with oncomodulin
(OCM) to reveal the striolar region and phalloidin to reveal F-actin (F-ACT)-rich hair
bundles. Graphs report striolar surface area and striolar HC density based on OCM
labeling and show no change in mutants compared to heterozygote littermate controls.
All points are graphed along with 25-75% boxplots (external bars: minimum and maximum,
internal bar: median, cross: mean). Mann-Whitney U test. **D-F**) P28 maculae
immunolabeled with SPP1 to reveal type I HCs. Low magnification of the utricle (**D**), high
magnification view showing SPP1 at the utricle HC neck (**E**), and quantification per region
in utricle and saccule (**F**). **G-I**) P28 maculae immunolabeled with SOX2 to reveal type II
HCs. Low magnification of the utricle (**G**), high magnification view showing SOX2 in the
utricle HC nucleus (**H**), and quantification per region in utricle and saccule (**I**). **F**, **I:** 2-way
ANOVA with Sidak's multiple comparisons. N, n: number of animals and HCs,
respectively. UTR, utricle. Scale bars: 100 µm (**B-C**, **D**, **G**), 10 µm (**E**, **H**).

Figure 2. No effects of *Gpr156* deletion on mechanosensitivity were detected in LES hair cells. A) *Left,* Schematic showing hair cell types with normal reversed bundle polarity at the LPR. *Arrows*, Directions of hair bundle motion that produce a depolarizing receptor potential. *Right,* Maps of recordings from LES zone, lateral to the LPR in heterozygotes, and similar location in null mutants, but without bundle reversal (arrows). **B-E**) Exemplar I_{MET} recordings from voltage-clamped LES HCs of different type (I vs. II, columns) and genotype (het vs. null, rows), and ranging in age from P10 to P31. XY scales for all 4 are in **D**. Each panel shows I_{MET} current (average of 3 traces) evoked by 3 of many iterated bundle displacements (bottom of **D,E**): negatively and positively saturating stimuli (X_{min},

X_{max}) plus a step to ~ $X_{1/2}$ (midpoint) of the operating range. *Red traces*, fits of current

decay for step to $X_{1/2}$ with Eq. 2 or 3 (Methods). Voltage was clamped with the indicated

holding currents at –94 mV (type I) and –84 mV (type II). **F,G**) *G(X)* relations from type I

(**F**) and type II (**G**) HCs. G, was peak MET conductance taken at the onset of each of 20

or 40 displacement steps (*X*), iterated by 100 nm or 35 nm, respectively, and averaged

across 3 repeated stimulus families. *Mean ± SEM; n = 5-8 cells.* Fitted *G(X)* parameters

are statistically similar for *Gpr156^{del/+}* and *Gpr156^{del/del}* HCs (see Table 2). *Filled circles,*

$X_{1/2}$ values from fits. X_{max}: 1.1 to 1.3 micrometers. X_{min}; –0.2 to –0.45 micrometers.

1373

1374

1375

Figure 3. No effects of *Gpr156* deletion on basolateral voltage-dependent conductances were detected in LES HCs. A-D) Exemplar voltage-dependent currents from lateral extrastriolar (LES) type I (a, c) and type II (b, d) HCs of *Gpr156^{del/+}* (**A**, **B**) and *Gpr156^{del/del}* (**C**, **D**) utricles. There are significant differences between HC types, but these differences are seen in both *Gpr156^{del/+}* (**A,B**) and *Gpr156^{del/del}* utricles (**C,D**). A pre-pulse from holding potential to –124 mV shows the usual type I-type II difference in K⁺ conductances at or near resting potential: the prepulse deactivates $g_{K,L}$ in type I cells (**A,C**), and activates a small inwardly rectifying current in type II cells (**B,D**). Following the pre-pulse, test steps were iterated in 5-mV increments from –124 mV to –54 mV (type I)

1385 or from −124 mV to +6 mV (type II), activating currents with very different time course and

1386 voltage dependence. **E,F**) G(V) relations measured from tail currents at the end of the

1387 iterated voltage steps show that $g_{K,L}$ activated positive to −114 mV (**E**) while type II

1388 delayed rectifying currents activated positive to −64 mV (**F**).

1389

Figure 4. Displacement sensitivity and time course of step-evoked transduction currents in each zone were not strongly affected by Gpr156 deletion. A-F) Exemplar I_{MET} recordings from voltage-clamped HCs of different genotype, HC type (columns), and epithelial zone (rows). I_{MET} current families (averages of 3) evoked by 3 of many iterated bundle displacements (**E,F**, bottom): negatively and positively saturating stimuli plus the step closest to the midpoint ($X_{1/2}$) of the G(X) relation. X_{max}: 1.08 to 1.32 μm. X_{min}: –0.2 to –0.45 μm. *Red traces*, fits of current decay with Eq. 2 or 3 (Methods). Holding potentials,

–94 mV (type I) and –84 mV (type II). **G-H**) Averaged peak (onset) $G_{MET}(X)$ relations for

type I (**G**) and type II (**H**) HCs. Data from 5-9 HCs of each type were fitted with the

Boltzmann functions, which were averaged to produce mean curves shown (± SEM). $X_{1/2}$

values (filled circles) are given. **I**) Mean maximum G_{MET} of type I HCs was larger in MES

than in LES or striola, but there was not a significant effect of genotype. *Black circles*, 2

striolar type I HCs from CD1 mice.

1403

Figure 5. Gpr156 impacted the mechanosensitive properties of lateral-line HCs. A) Schematic on the left shows a lateral-line neuromast from the side. HCs that detect anterior and posterior flow are color coded in blue and orange respectively. Posterior HCs do not express Emx2 and rely primarily on the MET channel subunits Tmc2a and Tmc2b. In contrast, anterior HCs express Emx2 and rely primary on Tmc2b for mechanosensation. Schematic on the right shows a plane taken through the apical hair bundle plane, viewed top-down. The directional sensitivity of each HC is dictated by the location of the

kinocilium, which is indicated by the circle on the side of each hair bundle. This apical

plane is one used to monitor mechanosensitive-calcium signals in lateral-line hair bundles.

The pipettes on each side of the hair bundles show the direction of flow delivered to

stimulate the two populations of HCs. **B1**-**B5**) Representative example of evoked-

mechanosensitive calcium signals in hair bundles of a control neuromast during a 500-

ms ANT (**B2**) or a 500-ms POST (**B3**) directed stimulus. Spatial patterns of GCaMP6s

signals during stimulation (**B2**-**B3**) are colorized according to the ΔF heat map and

superimposed onto a baseline GCaMP6s image. In **B1**, the colored circles indicate the

respective hair bundle orientation observed from calcium imaging (ANT, orange; POST,

blue; no response, gray). ROIs were placed on each hair bundle to generate the ΔF/F

GCaMP6s traces from individual ANT (**B4**) and POST (**B5**) responding hair bundles. **C1**-

C5) Representative example of evoked-mechanosensitive calcium signals in hair bundles

of a *gpr156* mutant neuromast during a 500-ms ANT (**C2**) or a 500-ms POST (**C3**)

directed stimulus. Spatial patterns of GCaMP6s signals during stimulation (**C2**-**C3**) are

colorized according to the ΔF heat map in **B2** and superimposed onto a baseline

GCaMP6s image. In **C1**, the colored circles indicate the respective hair bundle orientation

observed from calcium imaging (POST, blue). ROIs were placed on all hair bundle to

generate the ΔF/F GCaMP6s traces from individual hair bundles during ANT (**C4**) and

POST (**C5**) stimuli. **D-F)** Quantification of the average increase in GCaMP6s per

neuromast for ANT and POST hair bundles. Traces in **D** and **E** show the average

GCaMP6s response per neuromast in ANT and POST hair-bundle populations (mean ±

SEM is shown in **D-E**, n = 9 control and 10 *gpr156* neuromasts). The magnitude of the

GCaMP6s data in **D-E** is plotted to compare the average GCaMP6s increase for ANT and

POST hair bundles for each neuromast in **F**. In control neuromasts, the GCaMP6s

increase in the hair bundles ANT cells were larger compared to POST cells. In *gpr156*

mutants the ANT cells were significantly larger than control POST cells, but not different

than control ANT cells. **P = 0.0043 and P = 0.561, paired and unpaired t-test respectively.

Scale bar is 5 μm in **B3**.

1440

Figure 6. Afferent innervation patterns near the LES/S zone boundary were not substantially disturbed by *Gpr156* deletion. Afferent receptive fields (*green*) were labeled by diffusion of fluorescent dye (AlexaFluor) from whole-cell recording pipettes into calyces (*asterisks*) and throughout the terminal arbor, for (**A-D**) *Gpr156del/+* controls and (**E-H**) *Gpr156del/del* mutants. Counterstained with anti-calbindin (Calb1) antibody to show the striola (*magenta*). *Top right,* Schematic of the utricle with magenta striola; *black dots,* approximate location of each labelled calyx *shown*. All labeled afferents had a thick, medial-projecting neurite that branched to form up to 2 calyces and many bouton contacts. Anti-βII-spectrin labeling (**A**, **A'**) leaves an unlabeled hole where the kinocilium is, allowing determination of bundle orientation (*black arrows outlined in white, A*) and, in *Gpr156del/+* controls, the LPR (*dotted white line*). **A**) In one control afferent, the receptive field straddled the LPR (**A'**, **A''**), with 1 calyx on a type I HC in the LES (*white arrow*) and some boutons contacting type II HCs, as terminals or en passant, in the calbindin+ striola (*white arrowheads*). **B, C**) In all other fills, both control and mutant, the labeled LES arbors innervated only LES HCs. **C,** *arrow*, A thin branch extended from the fiber below the epithelium (Supplemental video). **D**) A receptive field labeled by filling a striolar calyx

included 2 calyces and some bouton endings, all restricted to the calbindin$^+$ striola (this

afferent is white because of the merge of AlexaFluor and calbindin stains). **E-G**) Afferent

terminal fields of LES calyces from *Gpr156$^{del/del}$* utricles largely remained in the calbindin$^-$

region (LES). **H**) A striolar (calbindin$^+$) calyx in a *Gpr156$^{del/del}$* mouse made multiple

boutons entirely in the calbindin$^+$ area (striola). Scale bar: 20 micrometers L, lateral; P,

posterior.

1465

1466

1467

Figure 7. LES afferent innervation pattern was comparable for *Gpr156*^{*del/+*} and
***Gpr156*^{*del/del*} utricles. A**) P30 utricles where oncomodulin (OCM) labels striolar HCs and
Advillin-positive afferents are labeled by EGFP in *Advillin-Cre; TIGRE; Gpr156* animals.
Advillin-Cre neurons specifically innervated lateral HCs and orientation-based
segregation was not altered in *Gpr156*^{*del/del*} mutants. **B**) Close-up views at the LES-striola
border (XY and YZ views). The vast majority of EGFP-positive HCs were in the the LES
but *Advillin*-positive neurons also innervated OCM⁺ HCs in the lateral striola in both
control and *Gpr156* mutants. **C**) Close-up view where basal bodies are labeled with
pericentrin (PCNT) to reveal HC orientation (arrowheads). Lateral striolar HCs innervated
by *Advillin* neurons were consistently oriented medially, showing that innervation from

Advillin neurons strictly segregates with HC orientation, and not with striolar

characteristics (OCM expression). Yellow dotted lines in A mark the outline of the utricle.

Scale bars: 100 µm (A) 10 µm (B-C).

1481

1482

1483

Figure 8. The excitability of LES afferents was not significantly affected by Gpr156

deletion. Exemplary voltage responses of control (**A**) and null (**B**) afferents to injected

steps of current. We recorded from the large calyceal endings as a way to access these

dimorphic afferents, which, like most LES afferents, made contact with both type I and

type II HCs. For current steps (bottom) at $3 \times I_{thresh}$ ($\times 3$, green), both genotypes produced

sustained firing, considered typical of regular afferents innervating LES and MES. Here

I_{thresh} (blue trace) was 100 pA for Gpr156$^{del/+}$ afferent and 250 pA for Gpr156$^{del/del}$ afferent,

but overall there was no significant difference in current threshold with genotype.

1492

1493

Phalloidin Myo7 tdTomato Emx2

control

Selected optical sections

Phalloidin Myo7 tdTomato Emx2

gpr156 ko

Selected optical sections

1494

Figure 9. Gpr156 was not required for lateral-line afferents to select Emx2$^+$ or Emx2$^-$ hair cells. A-H) Images of a control neuromast innervated by a single afferent fiber that contacts Emx2(-) cells at 5 dpf. **A**) Phalloidin label reveals 16 hair bundles (8 A>P and 8 P>A hair bundles). Z-stack projections show all HCs labeled with Myo7a (**B**) a single afferent fiber expressing tdTomato (**C)** and Emx2(+ and -) HCs (**D**). **E-H**) Selected optical sections of (**B**–**D**) highlight the single afferent contacting individual Emx2(-) HCs. **I-P**) Images of a *gpr156* mutant neuromast innervated by a single afferent fiber that contacts Emx2(-) cells. (**I**) Phalloidin label revealed 14 hair bundles (13 A>P and 1 P>A hair bundles). Z-stack projections show all HCs labeled with Myo7a (**J**) a single afferent fiber expressing tdTomato (**K**) and Emx2(+ and -) HCs (**L**). **M-O**) Selected optical sections of (**J**–**L**) highlight a single afferent in a *gpr156* mutants contacting individual Emx2(-) HCs. **Q**) In both controls and *gpr156* mutants, each afferent fiber contacted the same number of HCs per neuromast. **R**) The overall selectivity of afferent fiber for Emx2(+ or -) HCs were similarly high in *gpr156* mutant and controls. **S**-**T**) The selectivity of afferent fibers for Emx2(+) or Emx2(-) HCs was similarly high in both *gpr156* mutants and controls. Arrows in **A** and **I** indicate the orientation of the hair bundles in each example. SEM is shown in Q-T. An unpaired *t*-test was used for comparisons. Scale bar = 5 µm.

1513

1514

1515

Figure 10. Grp156 was not required for pairing of pre- and post-synapses in lateral-
line hair cells. A-**C**) Images neuromasts immunolabeled with Maguk and Ribeye b to
stain HC post- and pre-synapses, respectively at 5 dpf. **A**) Example of an immunostain of
a control neuromast showing 3 complete synapses to the right. **B**) Image of an *emx2*
mutant neuromast that shows complete synapses, as well as unpaired pre- and
postsynapses, shown to the right. **C**) Image of a *gpr156* mutant neuromast, with complete
synapses shown to the right. **D**) Quantification revealed the same number of complete
synapses per HC in *emx2* and *gpr156* mutants compared to controls. **E**-**F**) In *gpr156*
mutants, quantification revealed there is no difference in the number of unpaired pre- or
post-synapses per neuromast compared to controls. In *emx2* mutants, there were
significantly more pre- or post-synapses per neuromast compared to controls. SEM is
shown in **D**-**F**. Scale bar = 5 µm. Unpaired t-test were used to make comparisons. *p <
0.05, ****p< 0.0001.

1529

1530

1531

1532

1533 **Figure 11. *Gpr156* mutant mice failed to swim and maintain their balance in water.**

1534 **A**) Time taken to traverse the balance beam for control and *Gpr156del/del* mice. N is 33 and

1535 30 for control (green) and *Gpr156del/del* (red) mice respectively. **B**) Time taken to rescue

1536 an animal immediately after the animal exhibited underwater tumbling. N is 14 and 20 for

1537 control (green) and *Gpr156del/del* (red) mice respectively. ***P<0.001.

1538

VOR in dark

OKR

VOR in light

Figure 12. VOR and OKR responses were normal but OVAR responses were

altered in *Gpr156del/del* **mice. A**-**B**) VORd gain and phase (mean±SEM) plotted as a

function of frequency for control and *Gpr156del/del* mice. N is 8 and 7 for control (green)

and *Gpr156del/del* KO (red) mice respectively. **C**-**D**) OKR gain and phase (mean±SEM)

plotted as a function of frequency for control and *Gpr156del/del* mice. **E-F)** VORl gain and

phase (mean±SEM) plotted as a function of frequency for control and *Gpr156*^{del/del} mice.

(**G**) Average eye velocities (mean±SEM) during 72s-long off-vertical axis rotation for

control and *Gpr156*^{del/del} mice. N is 8 and 7 for control (green) and *Gpr156*^{-/-} (red) mice

respectively. **H)** OVAR bias during the steady state (mean±SEM) for control and

Gpr156^{del/del} mice. ***P<0.001.

1551

1552

1553

1554

1555

1556

1557

1558

1559

1560

1561

1562

1563

1564

 TABLES

1566

1567

1568

1569 **Table 1. Percent of LES HCs with transduction**

1570

	Type I	Type II	ALL
Gpr156^{del/+}	16/22 (73%)	23/38 (60%)	39/60 (65%)
Gpr156^{del/del}	20/29 (69%)	34/46 (73%)	54/75 (72%)
Age range (median)	P11-44 (P19)	P11-100 (P19)	P11-P100 (P18)
p value	0.99	0.63	0.89

1571

1572

1573

1574 **Table 2. Genotype comparisons of transduction and adaptation in LES HCs**

Genotype	n	Age (median)	$G(X)$ Boltzmann parameters				Adaptation component tau at X-1/2			Adaptation component extent at at X-1/2			
			G_{max}, nS	10-90%, nm	$X_{1/2}$, nm	dx, nm	τ_{VF}, ms (n)	τ_F, ms	τ_S, ms	% decay	% A_{VF}	% A_F	% A_S
Gpr156^{del/+}	23	P8-32 (P18)	3.0 ± 0.2	0.66 ± 0.04	0.20 ± 0.01	0.16 ± 0.01	0.3 ± 0.1 (14) n.d. (9)	10.5 ± 1.8	152 ± 31	64 ± 2	24 ± 5 (14) n.d. (9)	39 ± 4	37 ± 3
Gpr156^{del/del}	26	P8-32 (P20)	2.9 ± 0.2	0.72 ± 0.04	0.25 ± 0.02	0.16 ± 0.01	0.3 ± 0.3 (14) n.d. (12)	7.8 ± 1.0	140 ± 21	68 ± 3	24 ± 5 (14) n.d (12)	36 ± 4	40 ± 4

Statistics													
p value			0.91	0.26	0.11	0.9	0.44	0.18	0.75	0.3	0.95	0.67	0.62
NS power			0.05	0.2	0.36	0.05	0.12	0.27	0.06	0.18	0.05	0.07	0.08

Includes both cell types and all zones
*n.d. = very fast component not detected

1575

1576

1577 **Table 3. Genotype comparisons of electrical properties of Type I and II HCs in**

1578 **LES**

1579

HC type	Genotype	n	Age (median)	C_m, pF	V_{rest}, mV	$V_{1/2}$, mV	Slope factor, mV	R_{in} (MΩ)	G_{max}/C_m, nS/pF
Type I	$Gpr156^{del/+}$	9-13	P15-34 (P19)	5.8 ± 0.2	-85 ± 1	-83 ± 1	4.5 ± 0.3	46 ± 3	19 ± 2
	$Gpr156^{del/del}$	11-18	P12-44 (P18)	5.8 ± 0.3	-86 ± 1	-84 ± 1	4.8 ± 0.3	51 ± 3	24 ± 4
	p value			0.84	0.84	0.57	0.53	0.37	0.22
	*NS power			0.05	0.05	0.08	0.09	0.14	0.2
Type II	$Gpr156^{del/+}$	24-29	P11-47 (P18)	4.7 ± 0.1	-72 ± 1	-31 ± 1	8.9 ± 0.5	598 ± 35	2.7 ± 0.2
	$Gpr156^{del/del}$	28-29	P12-44 (P17)	4.4 ± 0.1	-73 ± 2	-30 ± 1	9.0 ± 0.6	575 ± 41	3.2 ± 0.2
	p value			0.14	0.71	0.48	0.85	0.67	0.11
	*NS power			0.31	0.07	0.1	0.05	0.07	0.34

1580

1581 *estimated power of non-significant result

1582

1583 **Table 4. Genotype comparison of G(X) and adaptation parameters for all hair cells**

Genotype	n	Age (median)	G(X) Boltzmann parameters				Adaptation component tau at X-1/2			Adaptation component extent at at X-1/2			
			G_{max}, nS	10-90%, nm	$X_{1/2}$, nm	dx, nm	τ_{VF}, ms (n)	τ_F, ms	τ_S, ms	% decay	% A_{VF}	% A_F	% A_S
$Gpr156^{del/+}$	23	P8-32 (P18)	3.0 ± 0.2	0.66 ± 0.04	0.20 ± 0.01	0.16 ± 0.01	0.3 ± 0.1 (14) n.d. (9)	10.5 ± 1.8	152 ± 31	64 ± 2	24 ± 5 (14) n.d. (9)	39 ± 4	37 ± 3
$Gpr156^{del/del}$	26	P8-32 (P20)	2.9 ± 0.2	0.72 ± 0.04	0.25 ± 0.02	0.16 ± 0.01	0.3 ± 0.3 (14) n.d. (12)	7.8 ± 1.0	140 ± 21	68 ± 3	24 ± 5 (14) n.d (12)	36 ± 4	40 ± 4

Statistics													
p value			0.91	0.26	0.11	0.9	0.44	0.18	0.75	0.3	0.95	0.67	0.62
NS power			0.05	0.2	0.36	0.05	0.12	0.27	0.06	0.18	0.05	0.07	0.08

Includes both cell types and all zones
*n.d. = very fast component not detected

1584

1585

1586

1587 **Table 5. Genotype comparison of excitability in LES afferents**

1588

Genotype	n	Age range (median)	# Transient	# Sustained	Vrest, mV	Rin, MΩ	I-threshold, pA
Gpr156[del/+]	18	P11-35 (P19)	1 (5%)	17 (94.5%)	-66 ± 1	614 ± 145	111 ± 27
Gpr156[del/del]	20	P13-35 (P16)	1 (5.0%)	19 (95.0%)	-67 ± 1	323 ± 67	105 ± 20
Statistics							
p value				0.99	0.20	0.09	0.85
***NS power**					0.25	0.43	0.05

1589

1590 *estimated power of non-significant result

1591

1592   **Supplemental Figures**

Phalloidin — **Myo7** — **tdTomato** — **Emx2**

Selected optical sections

Phalloidin — **Myo7** — **tdTomato** — **Emx2**

Selected optical sections

1593

1594

Supplemental figure 1. Gpr156 was not required for lateral-line afferents to select Emx2+ HCs. A-H) Images of a control neuromast innervated by a single afferent fiber that contacts Emx2+ cells. (A) Phalloidin label reveals 14 hair bundles (7 A>P and 7 P>A hair bundles). Z-stack projections show all HCs labeled with Myo7a (**B**) a single afferent fiber expressing tdTomato (**C**) and Emx2+ and Emx2- HCs (**D**). (**E-H**) Selected optical sections

of (**B**–**D**) highlight the single afferent contacting individual Emx2$^+$ HCs. (**I-P**) Images of a

gpr156 mutant neuromast innervated by a single afferent fiber that contacts Emx2$^+$ cells.

I) Phalloidin label revealed 14 hair bundles (12 A>P and 1 P>A hair bundles, 1 at 90°). Z-

stack projections show all HCs labeled with Myo7a (**J**) a single afferent fiber expressing

tdTomato (**K**) and Emx2$^+$ and Emx2$^-$ HCs (**L**). **M-O**) Selected optical sections of (**J**–**L**)

highlight a single afferent in a *gpr156* mutants contacting individual Emx2$^+$ HCs. Arrows

in **A** and **I** indicate the orientation of the hair bundles in each example. Scale bar: 5 µm.

1607

1608

Supplemental figure 2. No resting head tremor was observed in *Gpr156del/del* mice. Comparison of power spectra density of head movements at rest in translational axes and rotational axes between control and *Gpr156del/del*. N is 8 and 7 for control and *Gpr156del/del* mice respectively. Error bars: SEM.

1613

1621 **Supplemental figure 3. *Gpr156^{del/del}* mice scored normally on several basic standard**

1622 **behavioral tests**. Comparison of assigned scores between *Gpr156^{del/del}* and control mice

1623 show no impairment in (**A**) air righting, (**B**) contact inhibition, (**C**) tail hanging. N is 9 and

1624 7 for control and *Gpr156^{del/del}* mice respectively. Error bars: SEM.

1625

1626

1627

1628 **Supplemental figure 4. VOR learning was unimpaired in *Gpr156^(del/del)* mice. A**) VORd

1629 gain (mean ± SEM) before and after the VOR gain-down training plotted as a function of

1630 frequency for control mice (N=7). **B**) VORd gain (mean ± SEM) before and after the VOR

1631 gain-down training plotted as a function of frequency for *Gpr156^(del/del)* mice (N=7). **C**)

1632 Percent change in VOR gain for control and *Gpr156^(del/del)*. *P<0.05, ***P<0.001.

1633

1634

1635

1636

1637

1638

1639