Micron-resolution fiber mapping in histology independent of sample preparation

Marios Georgiadis[1]✉*, Franca auf der Heiden[2], Hamed Abbasi[3,4], Loes Ettema[3], Jeffrey Nirschl[5], Hossein Moein Taghavi[1], Moe Wakatsuki[1], Andy Liu[1], William Hai Dang Ho[1], Mackenzie Carlson[1,6], Michail Doukas[7], Sjors A. Koppes[7], Stijn Keereweer[4], Raymond A. Sobel[5], Kawin Setsompop[1], Congyu Liao[1], Katrin Amunts[2,8], Markus Axer[2,9], Michael Zeineh[1], Miriam Menzel[3,2]✉*

[1]Department of Radiology, Stanford University, Stanford, CA 94305, USA.
[2]Institute of Neuroscience and Medicine (INM-1), Forschungszentrum Jülich GmbH, Jülich, Germany.
[3]Department of Imaging Physics, Faculty of Applied Sciences, Delft University of Technology, Delft, the Netherlands.
[4]Department of Otorhinolaryngology and Head and Neck Surgery, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the Netherlands.
[5]Department of Pathology, Stanford University, Stanford, CA 94305, USA.
[6]Department of Neurology and Neurological Sciences, Stanford University, Stanford, CA 94305, USA.
[7]Department of Pathology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the Netherlands.
[8]C. and O. Vogt Institute for Brain Research, University Hospital Düsseldorf, Medical Faculty, University Düsseldorf, Germany.
[9]Department of Physics, School of Mathematics and Natural Sciences, University of Wuppertal, Wuppertal, Germany.

*These authors contributed equally: Marios Georgiadis, Miriam Menzel.
✉Correspondence and requests for materials should be addressed to Marios Georgiadis (email: marios-g@stanford.edu) and Miriam Menzel (email: m.menzel@tudelft.nl).

Abstract

Detailed knowledge of the brain's nerve fiber network is crucial for understanding its function in health and disease. However, mapping fibers with high resolution remains prohibitive in most histological sections because state-of-the-art techniques are incompatible with their preparation. Here, we present a micron-resolution light-scattering-based technique that reveals intricate fiber networks independent of sample preparation for extended fields of view. We uncover fiber structures in both label-free and stained, paraffin-embedded and deparaffinized, newly-prepared and archived, animal and human brain tissues – including whole-brain sections from the BigBrain atlas. We identify altered microstructures in demyelination and hippocampal neurodegeneration, and show key advantages over diffusion magnetic resonance imaging, polarization microscopy, and structure tensor analysis. We also reveal structures in non-brain tissues – including muscle, bone, and blood vessels. Our cost-effective, versatile technique enables studies of intricate fiber networks in any type of histological tissue section, offering a new dimension to neuroscientific and biomedical research.

Introduction

The brain contains billions of nerve fibers that form a highly complex and densely interwoven network. Detailed mapping of the neuronal trajectories is a major goal of neuroscience[1,2]. However, disentangling the fiber architecture with high resolution over large fields of view remains prohibitive: diffusion magnetic resonance imaging (dMRI) maps nerve fiber pathways in whole human brains *in vivo*, but even with longer scan times in post-mortem brains the resolution is in the tens of micrometers, not sufficient to resolve trajectories of individual nerve axons[3,4]. Electron microscopy resolves nerve cells at nanometer resolution[5,6], but tracking the fiber pathways is limited to small tissue volumes[7,8]. Small-angle X-ray scattering (SAXS) can directly probe multi-directional fiber pathways[9,10], but cannot provide whole-brain micrometer resolution and requires expensive equipment such as synchrotrons. Various light microscopy techniques visualize nerve fibers at high resolution and over larger fields of view, using staining[11,12], fluorescence[13], or other contrast mechanisms. However, they rely on structure tensor analysis to derive fiber trajectories[14,15], which decreases effective resolution and is limited in white matter regions with crossing or densely packed fibers. Polarization-based techniques directly visualize fiber trajectories by measuring their birefringence[16,17], with the current gold standard for analyzing nerve fiber architecture in whole human brain sections being 3D-polarized light imaging (3D-PLI)[18–20]. However, 3D-PLI analysis is limited in areas of crossing fibers, and section preparation needs to preserve birefringence, e.g. by avoiding dehydration or lipid removal. Scattered light imaging (SLI) has recently emerged as a method to reveal interwoven fiber networks in unstained cryo-sections with micrometer resolution[21–23].

At the same time, the most common tissue preparation method is paraffin embedding and microtome sectioning followed by histological staining[24]. These so-called formalin-fixed paraffin-embedded (FFPE) sections, which are abundantly available in research and clinical laboratories, cannot be analyzed by polarization-based techniques because the birefringence is significantly reduced during tissue preparation. One prominent example is the BigBrain atlas[11], containing thousands of FFPE sections of a whole human brain. Nissl-based structure tensor analysis (Nissl-ST) can be used to track white matter fibers in FFPE sections by analyzing the arrangement of glial cells to derive fiber directions[25]. However, Nissl-ST can only be applied to sections with Nissl-staining, has effective resolution of tens of micrometers, and –similar to polarization microscopy– does not resolve crossing fibers which are abundant in the brain.

Here, we present computational scattered light imaging (ComSLI) – a fast and cost-effective optical microscopy method that uses a rotating LED light and a high-resolution camera to reveal microscopic fiber orientations of virtually any new or archived tissue section, also in regions with densely interwoven fibers. Using ComSLI, we recover orientation maps from animal and human brain tissues prepared with different protocols, stains, thicknesses, and storage periods – including whole-brain sections from the BigBrain atlas. We show that our technique reconstructs

fiber microarchitecture in health and disease, discerning myelin from axonal loss in neurodegenerative disorders, and identifying degeneration of the perforant pathway in diseased hippocampus tissue. By comparing it to diffusion MRI, 3D-PLI and Nissl-ST, we demonstrate its key advantages in revealing highly detailed and interwoven fiber networks. Moreover, we show that ComSLI can also be applied to reveal directed structures in non-brain tissues, e.g. muscle and collagen fibers.

Our approach enables retrospective investigations of histology slides, combining information from any stain with fiber orientations, and can help reveal new micro-organization patterns as well as relationships between the microstructure and the underlying biology in healthy and diseased, animal and human tissues.

Results

Revealing fiber architecture in whole human brain sections

Oriented structures such as nerve fiber bundles scatter light predominantly perpendicular to their main axis[26], yielding characteristic light intensity profiles, $I(\varphi)$, where the mid-position of a peak pair indicates the fiber orientation (Fig. 1A). ComSLI reverses the light path, which is physically equivalent, to determine the orientations of fibers contained in each micrometer-sized image pixel. The setup (Fig. 1B) consists of a high-resolution camera with small-acceptance-angle lens, and a rotating LED-light source illuminating the tissue section from below at ~45° elevation. The camera captures images of the sample at different light source rotation angles φ, yielding an image series in which each pixel contains a light intensity profile across different illumination angles, $I(\varphi)$, according to the orientation of its contained fibers (Supplementary Movie 1). Analyzing the peaks in each intensity profile yields multiple fiber orientations for each image pixel.

First, we analyzed whether ComSLI can retrieve fiber orientations with micrometer resolution from extended field-of-view sections, such as entire FFPE human brain sections (*Jülich BigBrain*[11]), Fig. 1C. As can be seen, orientation information was recovered from single and crossing fiber bundles in each pixel, in both white and gray matter (Fig. 1D). Microscopic details become apparent: U-fibers connecting the precentral gyrus to neighboring gyri, Fig. 1E, with subjacent deeper fibers crossing the main tracts (small arrows). Fibers in the corpus callosum (cc) are relatively homogeneously decussating between the two hemispheres with small angular variations (Fig. 1F, small arrows), while fibers of the fornix (f) course superomedially, and complex fiber crossings in dense corona radiata white matter are discerned (Fig. 1G).

Such investigations at whole-brain level unveil microscopic fiber orientations, enabling detailed investigation of anatomy, correlations with cellular information, or pathology.

Fig. 1 | Retrieving fiber orientations from a whole human brain section. (**A**) Light beam scattered by in-plane fiber bundle, predominantly perpendicular to the fibers, resulting in paired peaks in the azimuthal intensity profile, $I(\varphi)$, whose midline indicates the fiber orientation. (**B**) Schematic of ComSLI measurement setup, allowing to measure the azimuthal intensity profile by reversing the light path. The tissue section is illuminated at ~45° elevation angle by a rotating light source while a camera captures a high-resolution image of the section at each (azimuthal) rotation angle φ. (**C**) Brightfield microscopy image of a coronal human brain section (second *BigBrain* dataset, section no. 3452, FFPE, silver-stained), 1μm/pixel. (**D**) Fiber orientation map of the brain section, 7μm/pixel. The fiber orientations are shown as multi-colored pixels according to the color wheel in the bottom right; multiple colors per pixel indicate multiple fiber orientations, such as the zoomed-in green-purple pixel of the right corona radiata. (**E-G**) Fiber orientations for boxes marked in (D). Orientations are encoded both by the line direction and its color. (**E**) U-fiber. (**F**) Corpus callosum (cc) and fornix (f). (**G**) Right corona radiata. For better visualization, the orientation-depicting lines of sets of 15x15 pixels (E-F) and 75x75 pixels (G) were overlaid.

ComSLI works across stains, preparation protocols, and species

Next, we tested whether ComSLI can retrieve information from histological sections independent of stain, sample preparation, and species, since the orientation of scattered light is affected solely by the orientation of scattering fibers.

We scanned consecutive FFPE sections of a human hippocampus with various stains, colors, intensities, antibody concentrations, and counter-stains (see Methods). Figure 2 and Supplementary Fig. 1 show the brightfield images (A), fiber orientation maps (B), and zoomed-in views of the cornu ammonis and multiple fiber tracts (C). Despite the different staining protocols, targets, intensities, and counter-stains, the retrieved fiber orientations are almost identical across hippocampal sections, with small anatomical differences between sections.

We also investigated a section embedded in celloidin, myelin-stained approximately 120 years ago, from the brain collection of the Cécile and Oskar Vogt Institute for Brain Research (Fig. 2D-E): Despite the section age and the low light transmission in parts of the section (Fig. 2D), the individual nerve fiber orientations were retrieved (Fig. 2E), e.g., in the corpus callosum and the Muratoff bundle (subcallosal fasciculus) running subependymal to the caudate nucleus surface[27] (arrowhead).

This led us to ask (i) if we can also retrieve orientations from sections at different steps of sample preparation, and (ii) if the sample processing affects the method's sensitivity. We scanned two sections in paraffin just after sectioning, deparaffinization, and staining (Supplementary Fig. 2). Excitingly, fiber orientations could be visualized from sections still in paraffin (Supplementary Fig. 2A-B), as well as from completely unstained sections (Supplementary Fig. 2C-D), showing similar results as the stained sections (Supplementary Fig. 2E-F). The derivation and concordance of white and gray matter orientations across sample preparation stages highlight the wide range of histology sections that can be analyzed with ComSLI.

To confirm that the technique works across species, we measured multiple animal brain sections with different stains, including from mouse (Fig. 2F-I) and pig (Fig. 2J-M). Tracts were retrieved with micrometer resolution, such as the intricate mouse cerebellar connectivity (Fig. 2I) and multiple crossing and U-fibers in the pig corona radiata (Fig. 2M).

Analyzing disturbed microarchitecture in pathologic tissues

To see if ComSLI can unveil disturbed microscopic fiber orientations and provide unique information in pathologic human tissue, we studied human brain samples with neuropathology.

First, we investigated a brain of a subject with multiple sclerosis (MS) – a neurological autoimmune disease damaging the insulating myelin sheaths of nerve fibers and impairing signal transmission. We imaged a section with focal demyelination (Fig. 3A, red-annotated lesions). To fully characterize the lesion pathologically, we performed additional myelin (Luxol fast blue - LFB) and neurofilament stains in consecutive sections (Supplementary Fig. 3).

Fig. 2 | ComSLI detects fiber orientations independent of tissue preparation and species. (**A-C**) **Human Hippocampi**, FFPE sections with different stains. (**A**) Brightfield microscopy images, 0.5μm/pixel. (**B**) ComSLI fiber orientation maps, 7μm/pixel, orientations encoded by color wheel. (**C**) Zoomed-in fiber orientation vector maps (white box in (B)), including parts of cornu ammonis and multiple tracts: fornix (f), endfolial pathway (ef), alveus (al), Schaffer collaterals (Sch). Vectors of 15x15 pixels are overlaid. (**D-E**) **Myelin-stained human brain section** (~120-year-old). (**D**) Section photograph. (**E**) ComSLI color-coded fiber orientation map (3μm/pixel) of marked region in (D), cc: corpus callosum, Cd: Caudate nucleus, MB: Muratoff bundle (arrowhead). (**F-I**) **Sagittal mouse brain section**, FFPE, microglia-stained. (**F**) Brightfield image, 0.5μm/pixel. (**G**) ComSLI average scattering signal, 4.25μm/pixel. (**H**) Fiber orientation map. (**I**) Zoom-in of box in (G) (cerebellar area), including cerebellar peduncle (cp), cerebellar nuclei (cn), lobes (inferior-il, posterior-pl, central-cl, anterodorsal-adl, anterobasal-abl), and gray matter including oriented molecular layer (ml) fibers. Vectors overlaid for 15x15 pixels. (**J-M**) **Pig hemisphere-brain section**, FFPE, H&E-stained. (**J**) Brightfield image, 0.5μm/pixel. (**K**) ComSLI average scattering signal, 9μm/pixel. (**L**) Fiber orientation map. (**M**) Fiber orientations from box in (K), overlaid for 15x15 pixels, including U-fiber tracts (curved arrows) and multiple crossing fibers (straight arrows); external capsule (ec), internal capsule (ic), corpus callosum (cc).

The light scattering signal (Fig. 3B) was not affected by myelin loss within the red-outlined lesion. The axonal orientation could be retrieved even in regions of extensive demyelination (Fig. 3C, arrowheads). In regions with a transition between normal-appearing white matter (nawm) and demyelinated white matter (dwm) axons (Fig. 3D-F), the ComSLI signal intensity is preserved and the fiber orientations are continuing their course despite the demyelination. This highlights the robustness of the method, which is not dependent on the presence of myelin, but rather of axonal orientations. Although the ComSLI signal is not significantly affected by myelin, it detects changes in tissue density; the lower ComSLI signal periventricularly (Fig. 3B, upper arrowhead) is due to the lower neuropil density in the region (Fig. 3G,I), with less corresponding fiber organization present (Fig. 3H). Even then, ComSLI retrieves fiber orientations from the remaining axons (Fig. 3H).

Next, we studied a brain sample with necrotizing leukoencephalopathy (LE) which is characterized by white matter necrosis with loss of neurons and myelin. The pathology-confirmed lesions (red-contoured areas in Fig. 3J) show a lower scattering signal (Fig. 3K), and are clearly delineated in both images. The lesions are located within the corona radiata (cr) where multiple nerve fiber crossings occur; the fiber orientation map (Fig. 3L) reveals the intricate fiber organization in this region, with grayish-appearing color showing crossing fibers of multiple orientations. A closer look (Fig. 3M,N,P,Q) shows the multiple crossings of the fibers surrounding the lesion but more heterogeneously distributed fibers inside the lesion, reflecting the disorganization associated with local tissue destruction. This partial loss of axons was confirmed with LFB and neurofilament stains respectively on consecutive sections (Fig. 3O,R), showing patchy axon loss and spongiosis as well as axonal spheroids and dystrophic axons (Supplementary Fig. 3I,J).

Looking at more microscopic fiber changes in pathology, we compared the hippocampal pathways from a subject with epilepsy-associated hippocampal sclerosis to those from a healthy control, both for coronal sections at the level of the lateral geniculate nucleus, Supplementary Fig. 4. At a macroscopic level, the diminished volume of the sclerotic hippocampus, Supplementary Fig. 4A, was accompanied by a strong degeneration of pathways, with only a few crossings across the hippocampus, Supplementary Fig. 4C. The perforant pathway (p), the principal source of entorhinal input to the hippocampus, overlying the medial subiculum, was almost completely degenerated, Supplementary Fig. 4D. On the other hand, the control hippocampus had multiple interwoven fibers across all subfields, Supplementary Fig. 4G, with an especially strong perforant pathway connecting the entorhinal cortex to CA1, CA3, and dentate gyrus, Supplementary Fig. 4H. Overall, ComSLI can reveal the organization of intact axonal fibers, despite demyelination, but does show loss of signal and loss of distributed axonal networks associated with neurodegeneration.

Fig. 3 | Human brain pathology measured with ComSLI (FFPE sections, H&E-, LFB-, and neurofilament-stained). Brightfield microscopy: 0.5µm/pixel size, ComSLI: 9µm/pixel. (**A-I**) **Multiple sclerosis (MS) brain**. (**A**) Brightfield image; white matter lesions (red contours) indicated by arrowheads. (**B**) Average scattering signal from ComSLI (v: ventricle, gm: gray matter, wm: white matter). (**C**) Fiber orientation map. (**D**) Zoom-in of brightfield H&E image corresponding to orange box in (B), at the interface of normal appearing white matter (nawm), demyelinated white matter (dwm), and gray matter (gm). (**E**) Fiber orientations in the same area. 15x15 pixel orientations are overlaid for visual clarity. (**F**) The same region in a consecutive LFB+H&E-stained section. (**G**) Zoom-in of brightfield H&E image corresponding to orange box in (B). (**H**) Fiber orientations in the same area. (**I**) The same region in a consecutive neurofilament-stained section. (**J-R**) **Leukoencephalopathy brain**. (**J**) Brightfield image; lesions are annotated with red contour, larger lesions indicated by arrowheads. (**K**) Average scattering signal. cc: corpus callosum, cg: cingulum, cr: corona radiata. (**L**) Fiber orientation map, lesions annotated with white contour. (**M**) Zoom-in of a large lesion in the H&E image corresponding to upper orange box in (K). (**N**) ComSLI

fiber orientations in the same area, showing some less aligned fibers (arrowheads). (**O**) Zoom-in of a lesion in a myelin-stained LFB+H&E consecutive section, showing the dramatic loss of myelin in the lesion as well as axonal loss and spongiosis, causing the reduced ComSLI signal. (**P**) Zoom-in of the H&E image corresponding to bottom orange box in (K). (**Q**) ComSLI fiber orientations in the same area, showing some areas of less aligned fibers (arrows). (**R**) Zoom-in of a lesion in a neurofilament-stained consecutive section.

Comparison to other fiber orientation techniques

We compared ComSLI to previously developed methods that yield fiber orientations using different mechanisms and physical phenomena.

We first contrasted our results on whole human brain sections (Fig. 4, left) to those obtained using Nissl-based Structure Tensor (Nissl-ST) on the same section (Fig. 4, middle). Nissl-ST uses 2D structure tensor analysis to retrieve fiber orientation information in most white matter areas based on the anisotropic shape of Nissl-stained glial cells[25]. We measured brain sections Nissl-stained with Cresyl violet (Fig. 4 and Supplementary Fig. 7), using the code and settings by Schurr & Mezer[25] to derive in-plane fiber orientations. Supplementary Fig. 5 shows the resulting fiber orientations for different kernel sizes, from 200x200µm to 50x50µm, showing the 100µm kernel as the best balance between data quality and resolution. The fiber orientations are in overall agreement between Nissl-ST and ComSLI (Fig. 4A,A'), with ComSLI achieving better visualization of small features, such as fiber bundles in the putamen (Pu) and globus pallidus (GPe, GPi), the medial and lateral medullary laminae (mml, lml), and the external capsule (ec), Fig. 4B,B'. Moreover, ComSLI resolves multiple crossing fibers per 3µm pixel in a label-free manner, for example in the corona radiata region (Fig. 4C,D), with Nissl-ST providing single fiber orientation per 100µm pixel (Fig. 4D').

We also compared our results to diffusion MRI-derived fiber orientations (Fig. 4, right) obtained from a high-resolution *in vivo* 18-hour diffusion MRI scan on the brain of a healthy volunteer[28]. ComSLI fiber orientations were compared with those from the MR plane that best matched the measured brain sections (see Methods). The main nerve fiber orientation patterns of ComSLI and dMRI are mostly in agreement (Fig. 4A,A''), with ComSLI unveiling distinct microscopic fiber orientation patterns, like the individual fiber bundles in the putamen and global pallidus or the medullary laminae (Fig. 4B), not resolved by dMRI *in vivo* (Fig. 4B'') or *ex vivo*[29].

ComSLI, Nissl-ST and dMRI comparison on another brain section, the cell-body silver-stained *Jülich BigBrain* section from Fig. 1C, yields similar conclusions (Supplementary Fig. 6).

Finally, 3D-Polarized Light Imaging (3D-PLI) provides micron-resolution fiber orientation maps of whole-brain sections[18–20]. Our tests on the studied whole-brain FFPE sections showed significantly reduced sensitivity (Supplementary Fig. 7, right), compared to the high-sensitivity high-resolution maps it provides in 3D-PLI sections. This is possibly because an important part of myelin lipids resolves during alcohol dehydration[30], and water removal also distorts myelin periodicity[31], resulting in loss of its birefringence.

Fig. 4 | Comparison of ComSLI, Nissl-ST, and dMRI fiber orientations. ComSLI and Nissl-ST were performed on the same brain section (human coronal FFPE section, Nissl-stained with Cresyl violet, no. 3301) – the ComSLI measurement with 3μm pixel size, and Nissl-ST on the brightfield microscopy image with 1μm pixel size. Diffusion MRI was performed *in vivo* on a healthy volunteer with 0.76mm isotropic voxel size, and the orientation distribution functions were evaluated at a plane similar to the evaluated brain section (see Methods). (**A**) Color-coded fiber orientations depicted for each image pixel. The orientations were computed with *SLIX*[32] for ComSLI and dMRI, and with the provided code for Nissl-ST[25] using 15μm blur radius and a kernel size of 200x200μm. (**B**) Enlarged views of the fiber orientation maps for the bottom rectangular area marked in (A), featuring individual fiber bundles in the globus pallidus and putamen; annotated are internal capsule (ic), external capsule (ec), optic tract (ot), globus pallidus external segment (GPe), globus pallidus internal segment (GPi), putamen (Pu), medial and lateral medullary laminae (mml, lml), and claustrum (Cl). Nissl-ST orientations were computed with 15μm blur radius and a kernel size of 100x100μm. (**C**) Enlarged views of the fiber orientations for the top rectangular area in (A), featuring crossing fibers in the corona radiata. Nissl-ST orientations were computed with 15μm blur radius and a kernel size of 100x100μm. ComSLI fiber orientations were visualized as colored lines and overlaid on 66x66 pixels for better visualization and comparison; Nissl-ST orientations were overlaid on 2x2 pixels; dMRI orientation distribution functions were visualized with *MRtrix3*'s *mrview*[33]. (**D**) Fiber orientations and distribution functions of 2x6 selected pixels, marked by the rectangular regions in (C), with orientation distributions displayed at the resolution of each method (3μm *vs.* 100μm *vs.* 760μm).

Extension to tissues beyond brain

Finally, we tested whether ComSLI can retrieve fiber orientations from non-brain human tissues, given the multiple fiber types present, often in close proximity.

We performed measurements on FFPE sections from different tissue types (Fig. 5): human tongue muscle, colorectal tissue, lower jaw bone (mandible), and arterial wall.

In the tongue, ComSLI reveals layers of differently oriented structures (Fig. 5A-D), like outer epithelium (i), underlying stroma (ii) with collagen fibers running parallel to the surface, and

inner muscular layer (iii), which contains interwoven muscle fibers that enable tongue reshaping and multi-directional movement.

In colorectal tissue, Fig. 5E-I, orientations of multiple fiber types are retrieved: epithelial lamina propria (iv) is typically oriented normal/radial to the surface, while the basal membrane

microstructure is oriented parallel to the surface (arrowheads in Fig. 5G); the mucosal and submucosal fibers (v) have a complex architecture with various orientations; circular muscle

fibers of the muscularis mucosa (vi) run from top to bottom (vertical bidirectional arrow in Fig. 5G and lower arrowhead in Fig. 5H); longitudinal fibers of the muscularis propria (vii) run

perpendicular to the circular muscle fibers (horizontal bidirectional arrow in Fig. 5G and upper arrowhead in Fig. 5H). Blood vessels yield high scattering signal (arrowheads in Fig. 5F), with

collagen fibers wrapping around vessel walls (arrowheads in Fig. 5I).

In mandible bone (Fig. 5J-L), the inner spongy bone (trabecular bone, ix) shows fibers mostly

following trabecular microstructure[34]. The U-shaped cortical bone (xiii) shows a dominant fiber orientation in the direction of the force applied during mastication (Fig. 5L, bidirectional arrow),

while the superior region (x) exhibits more variable fiber orientations, distributing incoming loads to the rest of the jaw.

In artery wall, Fig. 5M-O, ComSLI reveals microscopic arrangements of fibers in different layers. While the brightfield image (Fig. 5M) indicates the fiber type (pinkish area dominated by

collagen fibers, dark/black area by elastin fibers, grayish-greenish area by muscle fibers, indicated by arrowheads in corresponding color), the fiber orientation map (Fig. 5O) reveals

distinct fiber directions in each layer, critical for artery mechanical properties[35]: fibers run parallel to the surface in the inner, middle, and outer layers; in the layers in between, the fibers

run perpendicular to the surface (bidirectional arrows).

ComSLI revealed directed structures of non-brain tissues both for stained and unstained sections

314 (Supplementary Fig. 8), and yielded similar results across adjacent sections, for different section thicknesses (from 4μm to 15μm) and different steps of sample preparation, also for sections still

316 in paraffin (Supplementary Fig. 8A,D,G).

Fig. 5 | Non-brain samples measured with ComSLI. (**A-D**) **Human tongue muscle**. (**A**) Brightfield H&E image,
320 0.46μm/pixel; (i) epithelium, (ii) stroma, (iii) inner muscle layer. (**B**) Average scattering signal, 3μm/pixel. (**C**)
ComSLI fiber orientation map, color-coded according to color wheel in (D). (**D**) Zoomed-in area from (C) including
322 different fiber layers; fiber orientations are displayed as colored lines for every 25th pixel. (**E-I**) **Human colorectal**

tissue. (**E**) Brightfield H&E image, 0.46µm/pixel; (iv) epithelium, (v) mucosa and submucosa, (vi) circular muscle, (vii) longitudinal muscle. (**F**) Average scattering signal, 3µm/pixel, with two blood vessels indicated by arrowheads. (**G**) ComSLI fiber orientation map, color-coded according to color wheel in (H). Bidirectional arrows indicate main fiber orientation, arrowheads indicate the basal membrane. (**H,I**) Upper and lower zoomed-in areas from (G); fiber orientations are displayed as colored lines for every 40th image pixel. Arrowheads indicate vertical and horizontal fiber orientations from circular and longitudinal muscle respectively (H), and blood vessels with fibers running circumferential (I). (**J-L**) **Human mandible**. (**J**) Brightfield H&E image, 0.46µm/pixel; (xiii) cortical bone on anterior side, (ix) trabecular (cancellous) bone, (x) cortical bone on tooth side. (**K**) Average scattering signal, 3µm/pixel. (**L**) ComSLI fiber orientation map, color-coded according to color wheel. (**M-O**) **Elastic artery wall**. (**M**) Brightfield elastin-stained image, 0.46µm/pixel. (**N**) Average scattering signal, 3µm/pixel. (**O**) ComSLI fiber orientation map. Highlighted are fiber layers with different directions going from lumenal to adventitial surfaces. All samples are 4µm-thin FFPE sections and stained with either H&E (tongue, colorectal, mandible) or elastin staining (artery wall). Main fiber orientations in panels are indicated by bidirectional arrows.

Discussion

Nerve fibers form a dense network in the brain, determining its connectivity and function in health and disease. Here, we show that microscopic fiber orientations can be retrieved using ComSLI from both new and archived paraffin-treated tissue sections prepared with different stains and protocols.

Fiber orientation maps were retrieved from human whole-brain, hippocampus, isocortex, pathology, as well as pig and mouse brain sections. ComSLI provided microscopic fiber maps, invisible with brightfield microscopy and inaccessible with other methods.

The presented approach has the potential to make fiber mapping accessible to every laboratory: ComSLI is i) relatively low-cost, requiring only a light source (such as an LED light), a sample stage, and a micron-resolution camera, ii) fast, with a full measurement performed within seconds, iii) compatible with sections at virtually any stage of sample preparation. In miniaturized form, using an LED-ring instead of a rotating LED, ComSLI can be an add-on to existing microscopes and easily be integrated into clinical workflows. This enables microscopic exploration of fiber orientations in health and disease on sections readily available in thousands of laboratories and clinics worldwide.

Compared to Nissl-ST, ComSLI provides significantly higher resolution in a label-free manner, including multiple crossing fiber orientations per image pixel, and the possibility for out-of-plane orientation information[21–23]. Diffusion MRI can image living human subjects tomographically but lacks cellular and microscopic details. Compared to diffusion MRI, ComSLI provides much higher resolution leading to visualization of small fiber tracts invisible to MRI. Also, ComSLI can be applied to any section, new or archived, at any sample preparation stage, even for sections still in paraffin.

Given its independence from myelin birefringence, ComSLI can yield neuropathologic insights when combined with myelin-sensitive microscopic methods such as 3D-PLI[18–20]. polarization-sensitive optical coherence tomography[16,17], and X-ray scattering[9,10,36,37]. Especially compared to

3D-PLI, which is the method-of-choice for micron-resolution fiber mapping on sections, ComSLI can be applied to virtually any section independent of sample preparation or birefringence, including FFPE sections, and can resolve crossing fibers per pixel, with an easy-to-implement and cost-effective setup.

ComSLI enables retrospective studies that can shed light to changes in microscopic fiber orientations. In neuroscience, it can be applied to the multiple neurologic and psychiatric disorders with structural connectivity changes, such as Alzheimer's disease[38], Parkinson's disease[39], traumatic brain injury[40], stroke[41], multiple sclerosis[42], schizophrenia[43], epilepsy[44], or autism[45]. It can be used across any brain circuitry and can validate *ex vivo* diffusion MRI orientations[4], especially in areas with small or highly interwoven fiber tracts such as the hippocampus[20] or U-fibers[46], where sensitivity and high-resolution is key. Finally, it can be used to create reference maps of microscopic fiber architecture for entire brains, e.g., applied to and extending datasets such as the BigBrain atlas[11].

Our study shows that ComSLI can not only reveal nerve fiber architectures, but also other directed structures in non-brain tissues, like muscle, collagen, and elastin fibers. These form intricate biological fiber networks which control mechanical and functional properties, and are altered in disease in muscle[50], connective[47,48,51], and epithelial[35,49] tissues. ComSLI can be used to reveal collagen[52], actin/myosin[53], elastin[54], or the collagen fiber organization around tumors[55,56], which is related to staging and malignancy, potentially allowing more accurate diagnosis and tailored cancer treatment.

In this study, we only show in-plane fiber orientations, but the method also provides information of fiber inclination[21–23], and our future work will focus on quantifying out-of-plane angles. While measurements were performed in transmission mode through thin sections, we aim to extend measurements to back-scattering mode to analyze fiber orientations on surfaces without sectioning. Finally, ComSLI provides fiber orientations without distinguishing between nerve, muscle, collagen, or other fibers. Future studies will investigate the different scattering signatures from distinct fiber types to enable label-free, fiber-specific analyses.

In conclusion, ComSLI reveals the underlying fiber microarchitecture of tissues including paraffin-embedded formalin-fixed (FFPE) sections, independent of staining and birefringence. This fast, simple to implement, and cost-effective technique can become a routine tool to provide micrometer fiber orientation mapping of histological tissue sections, opening new avenues in the investigation of the microarchitecture of any tissue, in health or disease.

Methods

Sample preparation

Whole human brain (BigBrain) sections

Two human brains, obtained within 24 hours after death and without neurological disorders, were fixed in 4% formaldehyde, dehydrated in increasing alcohol series (80%, 90%, 96%, 100% ethanol for at least one week each), embedded in a solvent (chloroform) for one week, and embedded in 57-60°C paraffin solution for two months. After hardening, the brains were coronally cut into 20 µm-thin sections from anterior to posterior with a large-scale microtome (Leica SM2500 Microtome) and mounted on a 37°C heat plate in gelatin solution. The sections were placed in a decreasing alcohol series to remove the paraffin and then placed in a staining solution to highlight neuronal cell bodies: the sections of the first brain (30 years old, male) were stained with silver following the protocol of Merker[57]; the sections of the second brain (71 years old, male) were stained with Cresyl-violet. After staining, the sections were again dehydrated in increasing alcohol series, mounted/sealed on glass slides, and archived. The silver-stained sections were from the second BigBrain data set[58], 3D-reconstructed with the same spatial resolution of 20µm isotropic such as the original Jülich BigBrain (*24*), archived for 32 years, and section no. 3452 was selected for evaluation (Fig. 1 and Supplementary Fig. 6). The Cresyl-violet sections were archived for one and a half years, and sections no. 3301 (Fig. 4, left) and 2520 (Supplementary Fig. 7) were selected for evaluation. Body donors gave written informed consent for the general use of postmortem tissue used in this study for aims of research and education. The usage is covered by a vote of the ethics committee of the medical faculty of the Heinrich Heine University Düsseldorf, Germany (#4863).

120-year-old myelin-stained brain section

The myelin-stained human brain section (Fig. 2B) comes from the brain collection of the Cécile and Oskar Vogt Institute for Brain Research, Heinrich Heine University Düsseldorf, Germany. The brain of a 25-year-old male was embedded in celloidin and stained according to Weigert's iron hematoxylin myelin staining in 1904[59].

Human hippocampus, cortex, and pathology brain sections

Four-millimeter thick specimens of formalin-fixed human brains were dehydrated in increasing ethanol steps (70% x2, 95% x2, 100% x3, 3.5hrs each step), cleared in xylene (3.5hrs x2), paraffin-embedded (3.5hrs x2), and sectioned into 5µm-thin sections. The sections were de-waxed and stained with different agents as indicated. The hippocampal sections in Fig. 2A and Supplementary Fig. 1 were from a 89-year-old male with Alzheimer's pathology, stained against microglia (CD163), Perl's iron with Diaminobenzidine (DAB) enhancement, tau, and amyloid, with hematoxylin counterstain where indicated. Sections from brains with multiple sclerosis (80 years old, male, from temporal periventricular white matter and cortex) and leukoencephalopathy

(43 years old, male, from periventricular white matter and cingulum) were stained with hematoxylin and eosin, luxol fast blue plus hematoxylin and eosin, and neurofilament (2F11) (Fig. 3 and Supplementary Fig. 3). Hippocampal and visual cortex sections in Supplementary Fig. 2 were from a 60-year-old male stained with hematoxylin & eosin and a 67-year-old female stained with luxol fast blue respectively. The sclerotic hippocampal section in Supplementary Fig. 4 was from a 69-year-old female with epilepsy, the control was from a 66-year-old female with no neuropathologic abnormality. Specimens were acquired under Stanford ADRC IRB (Assurance nr. FWA00000935).

Mouse brain section

A female ~10-week-old C57BL/6 mouse (Jackson Laboratories) was housed in a temperature-controlled environment, with a 12-hour light/dark schedule and *ad libitum* food/water access. It was euthanized for the purposes of a different study (APLAC #32577) under anesthesia with 2-3% isoflurane followed by cardiac puncture and perfusion with 20mL phosphate buffered saline (PBS). The brain was harvested, kept in 4% paraformaldehyde (PFA) in PBS for 24 hours at 4°C, transferred to 10%, 20%, and 30% sucrose in PBS, embedded in Tissue-Tek O.C.T. in dry ice for 1 hour, and cut sagittally into 10μm sections using a cryotome (Leica CM1860). The sections were subsequently washed, mounted on a glass slide, intubated with Iba1 antibody (dilution 1:200), secondary antibody (goat anti-rabbit Cy3 1:200), and cover-slipped. A mid-sagittal section was selected for evaluation (Fig. 2F-I).

Pig brain section

A 4-week female Yorkshire pig (#2) was euthanized for a different study (Stanford APLAC protocol nr 33684), the brain was harvested, cut into 5-mm coronal slabs using a brain slicer, and a mid-frontal slab (#5) was paraffin-embedded, similar to the human pathologic specimen preparation above. The slab was cut in 10μm sections using a Leica HistoCore AUTOCUT microtome. After deparaffinization, a section (#127) was stained with hematoxylin and eosin and cover-slipped (Fig. 2J-M).

Human tongue, colorectal, bone, and artery wall sections

The non-brain tissue sections (Fig. 5 and Supplementary Fig. 8) were obtained from a tissue archive at Erasmus Medical Center, Rotterdam, the Netherlands, approved by the Medisch Ethische Toetsing Commissie (METC) under number MEC-2023-0587. The tissue samples were obtained from patients during surgery. The bone sample was decalcified first using DecalMATE by Milestone Medical. Afterwards, all samples were fixed in 4% formaldehyde for 24 hours, dehydrated in increasing alcohol series (70%, 80%, 90%, 96%, 100% ethanol), treated with xylene, embedded in paraffin, and cut with a microtome (Leica RM2165) into 4μm-thin sections. The sections were placed in a decreasing alcohol series to remove the paraffin, mounted on glass slides, stained with hematoxylin and eosin (artery wall with Verhoeff-Van Gieson elastin staining), and then cover-slipped.

Brightfield microscopy

The whole human brain sections were scanned with the TissueScope LE120 Slide Scanner by Huron Digital Pathology, Huron Technologies International Inc. The device measures in brightfield mode with 20X magnification and 0.74 NA, providing a pixel size of 0.4µm. The final images were stored with a pixel size of 1µm.

The hippocampus, cortex, pathology, and animal brain sections were scanned using an Aperio AT2 whole slide scanner with the ImageScope software and a 20X magnification, resulting in brightfield images with a pixel size of 0.5µm.

The stained non-brain microscopy slides were scanned using the Nanozoomer 2.0 HT digital slide scanner by Hamamatsu Photonics K.K., offering a 20X magnification and a pixel size of 0.46µm. The unstained non-brain microscopy slides were scanned using the Keyence VHX-6000 Digital Microscope (with VH-ZST objective, 20X), with a pixel size of 10µm.

ComSLI

Whole human brain (silver-stained), hippocampus, cortex, pathology, and animal brain sections

Measurements were performed with a rotating light source and camera (cf. Fig. 1A), using a Flexacam C3 12 MP microscope camera (Leica) and a Navitar 12X Zoom Lens with a 0.67X Standard Adapter and a 0.5X Lens Attachment, with 4.25-9µm pixel size, as indicated in the figure captions. As light source, an ADJ Pinspot LED II was used, with 5.1cm diameter and 3.5o full-angle of divergence, oriented at ~45° with respect to the sample plane. A motorized specimen stage enabled the whole-human-brain section scanning in 8x5 tiles, all other brain sections were scanned at single tile. Images were acquired at 10o rotation steps (36 images/sample) with 125ms exposure time, except from the sections in paraffin that gave very strong scattering and were imaged with 7ms exposure time. Prior to the measurement, a 100mm diameter diffuser plate (Thorlabs) was measured for calibration (see below for calibration details).

Whole human brain (Cresyl-violet) and non-brain tissue sections

Measurements were similarly performed with a rotating light source and camera (cf. Fig. 1A), using a fiber-coupled LED light source consisting of an ultra-high power LED (UHP-FB-W50 by Prizmatix) with 400-750nm wavelength (peak at 443nm), 2-meter long step-index multimode silica (low OH) fiber patch cord (Thorlabs), a 25.4 mm diameter collimating optics (FCM1-0.5-CN by Prizmatix), and a 25.4 mm diameter engineered diffuser (beam shaper) for homogenizing the illumination (ED1-S20-MD by Thorlabs), yielding top-hat beam with 20° full-angle of divergence. The exposure time was adjusted manually per sample for maximizing the dynamic range of the captured signal while avoiding saturation (range: 50-100 ms). The light source was oriented at ~45° with respect to the sample and rotated with a motorized specimen stage

504 (ZABER X-RSB060AD-E01-KX14A) in steps of 15° (24 images/sample). Images were taken with a 20 MP monochromatic CMOS camera (BASLER acA5472-17um) and a Rodenstock
506 Apo-Rodagon-D120 Lens, yielding a pixel size of 3µm (4µm optical resolution) and a field-of-view of 16x11mm². A motorized specimen stage was used to perform whole-slide scanning.
508 Prior to the measurement, a diffuser plate (DG100x100 N-BK7 ground glass diffuser, 1500 grit, Thorlabs) was measured for calibration.

510 *120-year-old myelin-stained brain section*

The measurement was performed with a similar camera and lens as for the non-brain tissue
512 sections (BASLER acA5472-17uc and Rodenstock Apo-Rodagon-D120), using an LED display instead of a focused light source (50x50cm², 128x128 RGB-LEDs, Absen Polaris 3.9pro
514 In/Outdoor LED Cabinet). The sample was illuminated by a green circle segment (9° azimuthal and polar widths, respectively) with an effective illumination angle of 47°, which was rotated in
516 15° steps. Images were taken with 10 seconds exposure time and a gain of 10, and 4 images were averaged per illumination angle to increase signal to noise.

518 *Calibration*

Prior to each measurement session, a diffuser plate was measured under similar conditions. A
520 100-pixel Gaussian blur was applied to resulting images to homogenize diffuser defects. Subsequently, the blurred images of all angles were divided by the average of their maxima for
522 normalization. These normalized diffuser images were used to calibrate the measured tissue images, aiming to account for the different light intensities across the field of view for each
524 image: Each tissue image was divided by its corresponding normalized diffuser image of the same illumination angle.

526 *Generation of fiber orientation and vector maps*

Each calibrated image series from a ComSLI measurement was evaluated with the open-source
528 software SLIX[32], which analyzes the position of scattering peaks to compute the fiber orientations and visualize them in in color-encoded maps, using multi-colored pixels and colored
530 vector lines. Measurements with 15° azimuthal steps were processed without filtering. Measurements with 10° azimuthal steps were processed with Fourier low pass filter (40% cutoff
532 frequency, 0.225 window width) before generating the parameter maps, as described in [22].

Nissl-ST

534 The fiber orientation maps were computed in Matlab following the procedure described and code shared by Schurr & Mezer[25], using default settings and 100µm as kernel to compute the structure
536 tensor (effective resolution).

3D-PLI

538 The 3D-PLI measurements were performed using the LMP3D microscope (Taorad GmbH, Germany), containing an evo4070MFLGEC (2048x2048) camera and a Nikon 4x (NA 0.2) lens,

540   which achieves a pixel size of 1.85µm and an in-plane optical resolution of 2.2µm (determined by US-Airforce target). The sample was illuminated by linearly polarized light in 20° rotation
542   angles and analyzed by a circular analyzer as described by Axer et al.[19]. 3D-PLI FOM was computed on the supercomputer JURECA at Forschungszentrum Jülich (grant no. 28954).

544   **Diffusion MRI**

The diffusion MRI dataset[28] is from a 30 year-old male who underwent 18 hours of diffusion
546   MRI scanning in the MGH-USC 3T Connectom scanner using gSlider-SMS[60]. After manually identifying the MR plane that most closely matched the BigBrain histology sections, the entire
548   dataset was rotated using FreeSurfer's *freeview* and the b-vectors were rotated at the same angles (rotation angles for the Silver-stained section were -34º sagittal and 1.5º axial, for the Cresyl-
550   violet -30.4º sagittal). Fiber responses and orientation distributions were computed using the *dwi2response* and *dwi2fod* functions in *MRtrix3*[33], using the multi-tissue, multi-shell algorithm[61],
552   and visualized in *mrview*. To generate whole-brain colormaps in Fig. 4 and Supplementary Fig. 6, *MRtrix3*'s *sh2amp* function was used to probe fiber orientations at the coronal plane at 5º
554   intervals, and colormaps were generated using SLIX[32].

Data availability

558   All data underlying the study and the original image files are available in the data repository Dryad: https://doi.org/10.5061/dryad.02v6wwqb2.

Code availability

562   The ComSLI fiber orientation and vector maps were generated with the open-source software *SLIX* which is available on GitHub (https://github.com/3d-pli/SLIX/). The Nissl-ST fiber
564   orientation maps were generated in Matlab using code shared by Schurr & Mezer[25]. The dMRI orientation distribution functions were computed with the open-source software *MRtrix3* which
566   is available on GitHub (https://github.com/MRtrix3/) and visualized with the tool *mrview*.

References

568   1. O. Sporns, G. Tononi, R. Kötter, The human connectome: a structural description of the human brain. *PLoS Comput Biol*. **1**, e42 (2005).

570   2. M. Axer, K. Amunts, Scale matters: The nested human connectome. *Science*. **378**, 500–504 (2022).

572   3. B. Jeurissen, M. Descoteaux, S. Mori, A. Leemans, Diffusion MRI fiber tractography of the brain. *NMR Biomed.* **32**, e3785 (2019).

574   4. A. Yendiki, M. Aggarwal, M. Axer, A. F. D. Howard, A.-M. van C. van Walsum, S. N.
576   Haber, Post mortem mapping of connectional anatomy for the validation of diffusion MRI. *Neuroimage*. **256**, 119146 (2022).

        5. S. Loomba, J. Straehle, V. Gangadharan, N. Heike, A. Khalifa, A. Motta, N. Ju, M. Siev-
578   ers, J. Gempt, H. S. Meyer, M. Helmstaedter, Connectomic comparison of mouse and human cortex. *Science*. **377**, eabo0924 (2023).

580   6. C. S. Xu, S. Pang, G. Shtengel, A. Müller, A. T. Ritter, H. K. Hoffman, S. Takemura, Z.
            Lu, H. A. Pasolli, N. Iyer, J. Chung, D. Bennett, A. V Weigel, M. Freeman, S. B. van En-
582   gelenburg, T. C. Walther, R. V Farese, J. Lippincott-Schwartz, I. Mellman, M. Solimena,
            H. F. Hess, An open-access volume electron microscopy atlas of whole cells and tissues.
584   *Nature*. **599**, 147–151 (2021).

        7. A.L. Eberle, D. Zeidler, Multi-beam scanning electron microscopy for high-throughput
586   imaging in connectomic research. *Front. Neuroanat.* **12**, 112 (2018).

        8. R.A. Salo, I. Belevich, E. Manninen, E. Jokitalo, O. Gröhn, A. Sierra, Quantification of
588   anisotropy and orientation in 3D electron microscopy and diffusion tensor imaging in in-
            jured rat brain. *NeuroImage.* **172**, 404–414 (2018).

590   9. M. Georgiadis, A. Schroeter, Z. Gao, M. Guizar-Sicairos, M. Liebi, C. Leuze, J. A. Mc-
            Nab, A. Balolia, J. Veraart, B. Ades-Aron, S. Kim, T. M. Shepherd, C. H. Lee, P. Wal-
592   czak, S. Chodankar, P. DiGiacomo, G. David, M. Augath, V. Zerbi, S. Sommer, I. Ra-
            jkovic, T. Weiss, O. Bunk, L. Yang, J. Zhang, D. S. Novikov, M. Zeineh, E. Fieremans,
594   M. Rudin, Nanostructure-specific X-ray tomography reveals myelin levels, integrity and
            axon orientations in mouse and human nervous tissue. *Nat. Commun.* **12** (2021).

596   10. M. Georgiadis, M. Menzel, J.A. Reuter, D.E. Born, S.R. Kovacevich, D. Alvarez, H.M.
            Taghavi, A. Schroeter, M. Rudin, Z. Gao, M. Guizar-Sicairos, T.M. Weiss, M. Axer, I.
598   Rajkovic, M.M. Zeineh, Imaging crossing fibers in mouse, pig, monkey, and human brain
            using small-angle X-ray scattering. *Acta Biomaterialia.* **164**, 317–331 (2023).

600   11. K. Amunts, C. Lepage, L. Borgeat, H. Mohlberg, T. Dickscheid, M.-É. Rousseau, S. Blu-
            dau, P.-L. Bazin, L. B. Lewis, A.-M. Oros-Peusquens, N. J. Shah, T. Lippert, K. Zilles,
602   A. C. Evans, BigBrain: An Ultrahigh-Resolution 3D Human Brain Model. *Science*. **340**,
            1472–1475 (2013).

604   12. V. Carriel, A. Campos, M. Alaminos, S. Raimondo, S. Geuna, Staining methods for
            normal and regenerative myelin in the nervous system. C. Pellicciari, E. Biggiogera
606   (Eds.) *Histochemistry of single molecules: methods and protocols*. Edited by New York:
            Springer. p. 207-218 (2017).

608   13. I. Costantini, E. Baria, M. Sorelli, F. Matuschke, F. Giardini, M. Menzel, G. Mazzamuto, L. Silvestri, R. Cicchi, K. Amunts, M. Axer, F.S. Pavone, Autofluorescence enhancement

610   for label-free imaging of myelinated fibers in mammalian brains. *Sci. Rep*. **11**, 8038 (2021).

612   14. A.R. Khan, A. Cornea, L.A. Leigland, S.G. Kohama, S.N. Jespersen, C.D. Kroenke, 3D structure tensor analysis of light microscopy data for validating diffusion MRI. *Neuro-*

614   *Image*. **111**, 192–203 (2015).

15. H. Wang, C. Lenglet, T. Akkin, Structure tensor analysis of serial optical coherence scan-

ner images for mapping fiber orientations and tractography in the brain. *J. Biomed. Opt*. **20**, 036003 (2015).

16. H. Wang, C. Magnain, R. Wang, J. Dubb, A. Varjabedian, L. S. Tirrell, A. Stevens, J. C. Augustinack, E. Konukoglu, I. Aganj, M. P. Frosch, J. D. Schmahmann, B. Fischl, D. A.

Boas, as-PSOCT: Volumetric microscopic imaging of human brain architecture and connectivity. *Neuroimage*. **165**, 56–68 (2018).

17. C. J. Liu, W. Ammon, R. J. Jones, J. Nolan, R. Wang, S. Chang, M. P. Frosch, A. Yendiki, D. A. Boas, C. Magnain, B. Fischl, H. Wang, Refractive-index matching en-

hanced polarization sensitive optical coherence tomography quantification in human brain tissue. *Biomed. Opt. Express*. **13**, 358–372 (2022).

18. M. Axer, K. Amunts, D. Grässel, C. Palm, J. Dammers, H. Axer, U. Pietrzyk, K. Zilles, A novel approach to the human connectome: Ultra-high resolution mapping of fiber tracts

in the brain. *Neuroimage*. **54**, 1091–1101 (2011).

19. M. Axer, D. Grässel, M. Kleiner, J. Dammers, T. Dickscheid, J. Reckfort, T. Hütz, B.

630   Eiben, U. Pietrzyk, K. Zilles, K. Amunts, High-resolution fiber tract reconstruction in the human brain by means of three-dimensional polarized light imaging. *Front. Neuroinform*.

632   **5**, 34 (2011).

20. M. M. Zeineh, N. Palomero-Gallagher, M. Axer, D. Grassel, M. Goubran, A. Wree, R.

634   Woods, K. Amunts, K. Zilles, Direct Visualization and Mapping of the Spatial Course of Fiber Tracts at Microscopic Resolution in the Human Hippocampus. *Cereb Cortex*. **27**,

636   1779–1794 (2017).

21. M. Menzel, J. A. Reuter, D. Gräßel, M. Huwer, P. Schlömer, K. Amunts, M. Axer, Scat-

638   tered Light Imaging: Resolving the substructure of nerve fiber crossings in whole brain sections with micrometer resolution. *Neuroimage*. **233**, 117952 (2021).

640   22. M. Menzel, M. Ritzkowski, J. A. Reuter, D. Gräßel, K. Amunts, M. Axer, Scatterometry Measurements With Scattered Light Imaging Enable New Insights Into the Nerve Fiber

642   Architecture of the Brain. *Front. Neuroanat*. **15** (2021).

23. M. Menzel, D. Gräßel, I. Rajkovic, M. M. Zeineh, M. Georgiadis, Using light and X-ray

644   scattering to untangle complex neuronal orientations and validate diffusion MRI. *Elife*. **12** (2023).

646   24. T. Sanderson, G. Wild, A. M. Cull, J. Marston, G. Zardin, 19—Immunohistochemical and immunofluorescent techniques. *Bancroft's Theory Pract. Histol. Tech. 8th ed.; Su-*

648   *varna, SK, Layton, C., Bancroft, JD, Eds*, 337–394 (2018).

25. R. Schurr, A. A. Mezer, The glial framework reveals white matter fiber architecture in

650   human and primate brains. *Science*. **374**, 762–767 (2021).

26. M. Menzel, M. Axer, H. De Raedt, I. Costantini, L. Silvestri, F. S. Pavone, K. Amunts,
652   K. Michielsen, Toward a High-Resolution Reconstruction of 3D Nerve Fiber Architec-
       tures and Crossings in the Brain Using Light Scattering Measurements and Finite-Differ-
654   ence Time-Domain Simulations. *Phys. Rev. X*. **10**, 21002 (2020).

27. J. D. Schmahmann, D. N. Pandya, *Fiber pathways of the brain* (Oxford University Press,
656   New York, 2006).

28. F. Wang, Z. Dong, Q. Tian, C. Liao, Q. Fan, W. S. Hoge, B. Keil, J. R. Polimeni, L. L.
658   Wald, S. Y. Huang, K. Setsompop, In vivo human whole-brain Connectom diffusion
       MRI dataset at 760 µm isotropic resolution. *Sci. data*. **8**, 122 (2021).

29. B. R. Plantinga, A. Roebroeck, V. G. Kemper, K. Uludağ, M. Melse, J. Mai, M. L. Kuijf,
660   A. Herrler, A. Jahanshahi, B. M. ter Haar Romeny, Y. Temel, Ultra-High Field MRI Post
662   Mortem Structural Connectivity of the Human Subthalamic Nucleus, Substantia Nigra,
       and Globus Pallidus. *Front. Neuroanat.* **10** (2016).

30. D. P. Leist, G. S. Nettleton, R. C. Feldhoff, Determination of lipid loss during aqueous
664   and phase partition fixation using formalin and glutaraldehyde. *J. Histochem. Cytochem.*
666   *Off. J. Histochem. Soc.* **34**, 437–441 (1986).

31. D. A. Kirschner, C. J. Hollingshead, Processing for electron microscopy alters membrane
668   structure and packing in myelin. *J. Ultrastruct. Res.* **73**, 211–232 (1980).

32. J. A. Reuter, M. Menzel, SLIX: A Python package for fully automated evaluation of
670   Scattered Light Imaging measurements on brain tissue. *J. Open Source Softw.* **5**, 2675
       (2020).

33. J.-D. Tournier, R. Smith, D. Raffelt, R. Tabbara, T. Dhollander, M. Pietsch, D. Christi-
672   aens, B. Jeurissen, C.-H. Yeh, A. Connelly, MRtrix3: A fast, flexible and open software
674   framework for medical image processing and visualisation. *Neuroimage*. **202**, 116137
       (2019).

34. M. Georgiadis, M. Guizar-Sicairos, O. Gschwend, P. Hangartner, O. Bunk, R. Muller, P.
676   Schneider, Ultrastructure organization of human trabeculae assessed by 3D sSAXS and
678   relation to bone microarchitecture. *PLoS One*. **11**, e0159838 (2016).

35. T. C. Gasser, R. W. Ogden, G. A. Holzapfel, Hyperelastic modelling of arterial layers
680   with distributed collagen fibre orientations. *J. R. Soc. Interface*. **3**, 15–35 (2005).

36. M. Georgiadis, A. Schroeter, Z. Gao, M. Guizar-Sicairos, D. S. Novikov, E. Fieremans,
682   M. Rudin, Retrieving neuronal orientations using 3D scanning SAXS and comparison
       with diffusion MRI. *Neuroimage*. **204**, 116214 (2020).

37. M. Georgiadis, M. Guizar-Sicairos, A. Zwahlen, A. J. Trüssel, O. Bunk, R. Müller, P.
684   Schneider, 3D scanning SAXS: A novel method for the assessment of bone ultrastructure
686   orientation. *Bone*. **71**, 42–52 (2015).

38. M. Filippi, S. Basaia, E. Canu, F. Imperiale, G. Magnani, M. Falautano, G. Comi, A.
688   Falini, F. Agosta, Changes in functional and structural brain connectome along the
       Alzheimer's disease continuum. *Mol. Psychiatry*. **25**, 230–239 (2020).

39. Y. Yang, C. Ye, J. Sun, L. Liang, H. Lv, L. Gao, J. Fang, T. Ma, T. Wu, Alteration of
690

brain structural connectivity in progression of Parkinson's disease: A connectome-wide network analysis. *NeuroImage Clin.* **31**, 102715 (2021).

40. R. Raizman, I. Tavor, A. Biegon, S. Harnof, C. Hoffmann, G. Tsarfaty, E. Fruchter, L. Tatsa-Laur, M. Weiser, A. Livny, Traumatic Brain Injury Severity in a Network Perspective: A Diffusion MRI Based Connectome Study. *Sci. Rep.* **10**, 9121 (2020).

41. T. Paul, V. M. Wiemer, L. Hensel, M. Cieslak, C. Tscherpel, C. Grefkes, S. T. Grafton, G. R. Fink, L. J. Volz, Interhemispheric Structural Connectivity Underlies Motor Recovery after Stroke. *Ann. Neurol.* **94**, 785–797 (2023).

42. E. Pagani, M. A. Rocca, E. De Meo, M. A. Horsfield, B. Colombo, M. Rodegher, G. Comi, M. Filippi, Structural connectivity in multiple sclerosis and modeling of disconnection. *Mult. Scler. J.* **26**, 220–232 (2019).

43. B. Yao, S. F. W. Neggers, R. S. Kahn, K. N. Thakkar, Altered thalamocortical structural connectivity in persons with schizophrenia and healthy siblings. *NeuroImage Clin.* **28**, 102370 (2020).

44. J. Royer, B. C. Bernhardt, S. Larivière, E. Gleichgerrcht, B. J. Vorderwülbecke, S. Vulliémoz, L. Bonilha, Epilepsy and brain network hubs. *Epilepsia*. **63**, 537–550 (2022).

45. M. Valenti, M. C. Pino, M. Mazza, G. Panzarino, C. Di Paolantonio, A. Verrotti, Abnormal Structural and Functional Connectivity of the Corpus Callosum in Autism Spectrum Disorders: a Review. *Rev. J. Autism Dev. Disord.* **7**, 46–62 (2020).

46. F. Movahedian Attar, E. Kirilina, D. Haenelt, K. J. Pine, R. Trampel, L. J. Edwards, N. Weiskopf, Mapping Short Association Fibers in the Early Cortical Visual Processing Stream Using In Vivo Diffusion Tractography. *Cereb. Cortex*. **30**, 4496–4514 (2020).

47. J.-Y. Rho, L. Kuhn-Spearing, P. Zioupos, Mechanical properties and the hierarchical structure of bone. *Med. Eng. Phys.* **20**, 92–102 (1998).

48. K. E. Stockhausen, M. Qwamizadeh, E. M. Wölfel, H. Hemmatian, I. A. K. Fiedler, S. Flenner, E. Longo, M. Amling, I. Greving, R. O. Ritchie, F. N. Schmidt, B. Busse, Collagen Fiber Orientation Is Coupled with Specific Nano-Compositional Patterns in Dark and Bright Osteons Modulating Their Biomechanical Properties. *ACS Nano*. **15**, 455–467 (2021).

49. D. T. Corr, D. A. Hart, Biomechanics of Scar Tissue and Uninjured Skin. *Adv. Wound Care*. **2**, 37–43 (2012).

50. P. P. Purslow, J. A. Trotter, The morphology and mechanical properties of endomysium in series-fibred muscles: variations with muscle length. *J. Muscle Res. Cell Motil.* **15**, 299–308 (1994).

51. M. Liebi, M. Georgiadis, A. Menzel, P. Schneider, J. Kohlbrecher, O. Bunk, M. Guizar-Sicairos, Nanostructure surveys of macroscopic specimens by small-angle scattering tensor tomography. *Nature*. **527**, 349–352 (2015).

52. J. Lin, Y. Shi, Y. Men, X. Wang, J. Ye, C. Zhang, Mechanical Roles in Formation of Oriented Collagen Fibers. *Tissue Eng. Part B Rev.* **26**, 116–128 (2019).

53. K. L. Weirich, S. Stam, E. Munro, M. L. Gardel, Actin bundle architecture and mechanics regulate myosin II force generation. *Biophys. J.* **120**, 1957–1970 (2021).

732   54. L. Baumann, E. F. Bernstein, A. S. Weiss, D. Bates, S. Humphrey, M. Silberberg, R. Daniels, Clinical Relevance of Elastin in the Structure and Function of Skin. *Aesthetic*
734   *Surg. J. Open Forum*. **3**, ojab019 (2021).

55. W. Han, S. Chen, W. Yuan, Q. Fan, J. Tian, X. Wang, L. Chen, X. Zhang, W. Wei, R.
Liu, J. Qu, Y. Jiao, R. H. Austin, L. Liu, Oriented collagen fibers direct tumor cell intravasation. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 11208–11213 (2016).

56. S. Z. Despotović, Đ. N. Milićević, A. J. Krmpot, A. M. Pavlović, V. D. Živanović, Z. Krivokapić, V. B. Pavlović, S. Lević, G. Nikolić, M. D. Rabasović, Altered organization
of collagen fibers in the uninvolved human colon mucosa 10 cm and 20 cm away from the malignant tumor. *Sci. Rep.* **10**, 6359 (2020).

57. B. Merker, Silver staining of cell bodies by means of physical development. *J. Neurosci. Methods*. **9**, 235–241 (1983).

58. H. Mohlberg, B. Tweddell, T. Lippert, K. Amunts, Workflows for Ultra-High Resolution 3D Models of the Human Brain on Massively Parallel Supercomputers, in *International*
Workshop on Brain-Inspired Computing (2015).

59. O. Vogt, Die myelogenetische Gliederung des Cortex cerebelli. *J. für Psychol. und Neu-*
748   *rol.* **5**, 235–258 (1905).

60. K. Setsompop, Q. Fan, J. Stockmann, B. Bilgic, S. Huang, S. F. Cauley, A. Nummenmaa,
750   F. Wang, Y. Rathi, T. Witzel, L. L. Wald, High-resolution in vivo diffusion imaging of the human brain with generalized slice dithered enhanced resolution: Simultaneous mul-
752   tislice (gSlider-SMS). *Magn. Reson. Med.* **79**, 141–151 (2018).

61. B. Jeurissen, J.-D. Tournier, T. Dhollander, A. Connelly, J. Sijbers, Multi-tissue con-
754   strained spherical deconvolution for improved analysis of multi-shell diffusion MRI data. *Neuroimage*. **103**, 411–426 (2014).

Acknowledgments

We thank Markus Cremer and the laboratory team at Forschungzentrum Jülich (INM-1), Germany, for preparing the whole human brain (BigBrain) sections, and Philipp Schlömer for performing and processing the 3D-PLI measurement. We thank David Camarillo for his help with the pig project. This work was supported by the National Institutes of Health grant nos. R03AG083702 (M.G.), R01MH092311 (K.A.), R01AG061120 (M.Z.), and R21AG072675 (M.Z.). F.H., K.A., M.A. and M.M. received funding from the European Union's Horizon 2020 Research and Innovation Programme grant no. 945539 ("Human Brain Project" SGA3) and no. 101147319 ("EBRAINS 2.0 Project"). F.H., L.E. and M.M. received funding from the Deutsche Forschungsgemeinschaft (DFG) Project no. 498596755. M.M. received funding from the Klaus Tschira Boost Fund and from the Helmholtz Association's Initiative and Networking Fund through the Helmholtz International BigBrain Analytics and Learning Laboratory (HIBALL) under the Helmholtz International Lab grant agreement InterLabs-0015. H.A. received funding from the Convergence Imaging Facility and Innovation Center (CIFIC) of TU Delft, Erasmus MC, and Erasmus UR. M.C. received funding from the National Institutes of Health Ruth L. Kirschstein F31 award number 1F31AG077905. M.G. received funding from the Alzheimer's Association Research Fellowship no. 24AARF-1241479.

Author contributions

M.G. and M.M. co-designed the study, developed the LED-spot setup, supervised the research, analyzed and interpreted the data, prepared the figures, and wrote the first version of the manuscript. M.G. performed the measurements of the whole human brain (silver-stained) section, the hippocampus, pathology, and animal brain sections. F.H. optimized the measurement parameters and hardware of the LED-display setup, performed first measurements of the FFPE whole brain sections, measured and analyzed the myelin-stained human brain section, and contributed to a figure. H.A. measured and analyzed the non-brain tissue samples and prepared the corresponding figures. L.E. measured and analyzed the Cresyl-violet stained, whole human brain sections. J.N. contributed to the design of the pathology study, provided anatomical analysis and interpretation of the brain pathology samples. H.M.T. helped with the Nissl-ST analysis, M.W. performed initial measurements of FFPE sections. A.L. contributed to scanning of the silver-stained human brain section, W.H.D.H. helped with setting up the measurements with the LED-spot, and M.C. contributed to the mouse brain study. M.D. provided anatomical analysis and interpretation of tongue, jar bone, and blood vessel samples, S.A.K. of colorectal samples, S.K. of oral tissue samples, and R.A.S. of pathological human brain samples. K.S. and C.L. helped design the comparison to dMRI, and C.L with analyzing the dMRI data set. K.A. provided the BigBrain sections, contributed to the anatomical content, and participated in the design of the study. M.A. and M.Z. co-designed the study and contributed to data interpretation and supervision. All authors contributed to revision and editing of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information accompanies this paper.

Correspondence and requests for materials should be addressed to M.G. or M.M.

Supplementary Information

802   **Supplementary Movie 1 | ComSLI dataset of whole human brain BigBrain section** (second *BigBrain* dataset, section no. 3452, FFPE, silver-stained, fiber orientations in Fig. 1). Each frame shows pixel intensities after
804   calibration (see Methods) for the entire brain at each illumination angle φ (in degrees). The movie, which has been 5x downsampled to reduce its size of many gigabytes, can be downloaded from the data repository Dryad:
806   https://doi.org/10.5061/dryad.02v6wwqb2 [SupplementaryMovie1.gif].

808 **Supplementary Fig. 1 | Differently stained FFPE sections of a human hippocampus measured with ComSLI** (same block as in Fig. 2A-C). (**A**) Brightfield microscopy images, 0.5μm/pixel. (**B**) Color-coded fiber orientation
810 maps from ComSLI, 7μm/pixel. (**C**) Zoomed-in fiber orientation vectors (rectangular region marked by white box in (B)). Vectors of sets of 15x15 pixels are overlaid for visual clarity.

Supplementary Fig. 2 | ComSLI at different steps of FFPE sample preparation. Left: human hippocampus, right: human isocortex and subcortical white matter. Each panel contains brightfield images, 0.5µm/pixel (left side) and the respective ComSLI color-coded fiber orientations, 8µm/pixel (right side). (**A,B**) Sections still in paraffin. The fiber orientations are clearly visible in white (w) and gray (g) matter, despite evident paraffin structures in the photographs (top left corner) and slide-scanned images (bottom left corner). Small folds in the physical section create small artifacts in the gray matter (arrowhead in (B)). (**C,D**) The same sections after deparaffinization, before staining. The sections lack brightfield contrast for slide-scanning so only photographs are shown (left). Retrieved fiber orientations are similar to those in (A,B), especially in the white matter. (**E,F**) The same sections after staining (hippocampus: hematoxylin and eosin (H&E), cortex: luxol fast blue (LFB)), where orientations follow the same orientation patterns. Overall, gray matter orientations are more pronounced in paraffinized compared to unstained sections. In the hippocampus, gray matter orientations were pronounced in the stained section as well, similar to the hippocampi in Fig. 2 and Supplementary Fig. 1, owing to the multiple tracts crossing the hippocampal gray matter[20].

Supplementary Fig. 3 | Multiple sclerosis (MS) and leukoencephalopathy (LE) brain samples from Fig. 3, additional histology stains (LFB+H&E and neurofilament). (**A**) MS brain section with Luxol-fast-blue (LFB) and hematoxylin & eosin (H&E) stain; LFB stains myelin blue, showing the demyelinating gradient, with minimal myelin left near the ventricle and the cortex (arrowheads). (**B**) Consecutive neurofilament (NF)-stained section, with orange boxes zoomed-in in C-E. (**C**) Normal-appearing white matter with NF stain, with high density of NF stain and healthy cell nuclei. (**D**) High myelin loss region, with oblique axons and tissue separation leading to low ComSLI signal in Fig. 3B. (**E**) Region of lower myelin loss, with preserved axons, lower stain and nuclei density, leading to a bit lower scattering signal only in Fig. 3B. (**F**) LE brain section LFB+H&E stained, with distinct fully demyelinated lesions (black arrowheads). (**G**) Consecutive NF-stained section, with orange boxes zoomed-in in H-J. Examples of axonal spheroids and thickened dystrophic axons are indicated by red arrowheads and ellipsoids, respectively. (**H**) Relatively normal-appearing white matter with NF stain, with high density of NF stain and healthy cell nuclei, with only occasional axonal spheroids and dystrophic axons. (**I, J**) More involved areas of partial spongiosis and disrupted axons, with region-wise sparser stain and lower nuclei density.

Supplementary Fig. 4 | Human hippocampal pathology measured with ComSLI (FFPE sections, H&E stained).
Left images show a sclerotic hippocampus from an epileptic patient with atrophy largely involving CA1 extending to the medial subiculum, right images a hippocampus from a healthy control, both coronal sections at the level of the
lateral geniculate nucleus. (**A**) Brightfield microscopy image of the sclerotic hippocampus, 0.5µm/pixel. (**B**) Corresponding average scattering signal from ComSLI, 8µm/pixel. DG: dentate gyrus, CA1: cornu ammonis 1,
CA3: cornu ammonis 3. SUB: subiculum. (**C**) Fiber orientations of the sclerotic hippocampus; fiber orientations are displayed as colored lines according to the color wheel. Dashed arrow indicates a very weak, heavily degenerated
perforant pathway (p). Colored lines representing orientations of 15x15 pixels are overlaid for visual clarity. (**D**) Zoom-in of white box in (C) showing detail of subiculum and cornu ammonis, where few, clear tracts are visible, in
contradistinction to the healthy hippocampus (E-H). (**E**) Brightfield image of the control hippocampus. (**F**) Corresponding average scattering signal. (**G**) Fiber orientations of the control hippocampus. White arrows indicate a
strong perforant pathway (p) crossing the subiculum towards the cornu ammonis. (**H**) Zoom-in of white box in (G) showing detail of the intricate perforant pathway crossings. Note that tissue tears are present in the subiculum in
both specimens.

Supplementary Fig. 5 | Nissl-ST fiber orientations computed for different kernel sizes. (**A**) Fiber orientations of the Cresyl-violet stained brain section no. 3301 from Fig. 4A computed from the brightfield microscopy image (1µm/pixel) with the provided code for Nissl-ST[25], using 15µm blur radius and a kernel size of 200x200µm, 100x100µm, and 50x50µm. (**B**) Zoom-in of the rectangular areas marked in (A), with anatomical regions labeled as in Fig. 4B.

Supplementary Fig. 6 | Comparison of ComSLI, Nissl-ST, and dMRI fiber orientations for a silver-stained human brain section. ComSLI and Nissl-ST were performed on the same brain section (second *BigBrain* dataset,
section no. 3452, FFPE, silver-stained, cf. Fig. 1) – the ComSLI measurement with 9µm pixel size, and Nissl-ST on the brightfield microscopy image with 1µm pixel size. Diffusion MRI was performed *in-vivo* on a healthy volunteer
with 0.76mm isotropic voxel size, and the orientation distribution functions were evaluated at a plane similar to the evaluated brain section (see Methods). (**A**) Color-coded fiber orientations depicted for each image pixel. The
orientations were computed with *SLIX*[32] for ComSLI and dMRI, and with the provided code for Nissl-ST[25], using 15µm blur radius and 100x100µm kernel. (**B**) Enlarged views for the rectangular area marked in (A), featuring
crossing fibers in the corona radiata. ComSLI fiber orientations were visualized as colored lines and overlaid on 200x200 pixels for better visualization and comparison; Nissl-ST orientations were overlaid on 5x5 pixels; dMRI
orientation distribution functions were visualized with *MRtrix3*'s *mrview*[33].

Supplementary Fig. 7 | 3D-PLI fiber orientation map of an FFPE brain section in comparison to those obtained by ComSLI and Nissl-ST. One hemisphere of a Cresyl-violet stained, human coronal brain section (no.
2520) was measured with 3D-PLI (1.85µm/pixel) and ComSLI (3µm/pixel), and the fiber orientations were computed for comparison. Due to brain preparation, the organization of myelin was destroyed and led to the loss of
birefringence, going along with an impairment of fiber contrast in 3D-PLI measurements. The Nissl-ST fiber orientations were computed from the brightfield microscopy image (1µm/pixel) with 15µm blur radius and
200x200µm kernel size.

**Supplementary Fig. 8 | Human tongue sections of 4µm, 10µm, and 15µm thickness measured with ComSLI at
different steps of sample preparation.** Each row corresponds to three different sample preparation steps of a single
884 section of the indicated thickness. The left and middle images of each panel show the average scattering signal and
the color-coded fiber orientations from ComSLI respectively (3µm/pixel), the right images show the corresponding
886 brightfield microscopy images (0.46µm/pixel for H&E-stained sections, 10µm/pixel for the other sections). Panels in
the left column (A,D,G) show the sections still in paraffin, unstained and without coverslip. Panels in the middle
888 column (B,E,H) show the same sections after deparaffinization. Panels in the right column (C,F,I) show these
sections after staining with hematoxylin and eosin (H&E) and cover-slipping.