Reprogramming of host energy metabolism mediated by the TNF-iNOS-HIF-1α axis plays a key role in host resistance to *Plasmodium* infection

KELY C. MATTEUCCI[1,2*], NATHALIA P. S. LEITE[1,2,7], PATRICIA A. ASSIS[3], ISABELLA C. HIRAKO[4], FRANCIELLE PIOTO[1], OGOOLUWA OJELABI[5], JULIANA E. TOLLER-KAWAHISA[6], LEONARDO G. VAZ[7], DIEGO L. COSTA[2], JOÃO S. DA SILVA[1], JOSÉ C. ALVES-FILHO[6], RICARDO T. GAZZINELLI [1,2,4,5,7*]

[1] Translational Medicine Platform Oswaldo Cruz Foundation, Ribeirão Preto Medical School, University of Sao Paulo, Brazil.
[2] Department of Biochemistry and Immunology, Ribeirão Preto Medical School, University of Sao Paulo, Brazil.
[3] Department of Pathology, University of Michigan Medical School, USA
[4] Laboratory of Immunopathology, Instituto René Rachou, Fundação Oswaldo Cruz - Minas, Brazil.
[5] Department of Medicine, Division of Infectious Diseases and Immunology, University of Massachusetts Chan Medical School, USA.
[6] Department of Pharmacology, Ribeirão Preto Medical School, University of Sao Paulo, Brazil.
[7] Technology Center for Vaccines, Federal University of Minas Gerais, Brazil

*Contact info

Gazzinelli, Ricardo T (Ricardo.gazzinelli@ummassmed.edu) and Kely Catarine Matteucci (kelycatarine@gmail.com).

Summary (128 words)

TNF has a dual effect in *Plasmodium* infection, bolstering the host's immune defense while also inducing sickness behavior. Here, we show that TNF signaling hampers physical activity, food intake, and energy expenditure while enhancing glucose uptake by the liver and spleen as well as controlling parasitemia in *P. chabaudi* (*Pc*)-infected mice. We also report that TNF is required for expression of inducible nitric oxide synthase (iNOS), stabilization of HIF-1α, expression of glucose transporter GLUT1 and enhanced glycolysis in monocytic cells from *Pc*-infected mice. Importantly, *Pc*-infected iNOS$^{-/-}$, TNFR$^{\Delta Lyz2}$ and *HIF-1α*$^{\Delta Lyz2}$ mice show impaired release of TNF and glycolysis in monocytes, along with increased parasitemia and disease tolerance. Altogether, our results indicate that TNF-iNOS-HIF-1α-induced glycolysis in monocytes plays a critical role in host defense and sickness behavior in *Pc*-infected mice.

Keywords: Glucose, TNF, iNOS, HIF-1α, monocytic cells, *P. chabaudi*, myeloid cells, malaria.

Tease

The role of host energy metabolism and glycolysis in monocytes as determinant of host resistance to *Plasmodium* infection and tolerance to disease.

Introduction

Malaria is the most common mosquito-borne infectious disease caused by parasites from the *Plasmodium* genus (*1*). In 2021, approximately 247 million cases were reported, which resulted in nearly 619 000 deaths worldwide (*2*). The clinical symptoms include periodic fever episodes that occur synchronously with the rupture of infected red blood cells and the release of erythrocyte and parasite debris (*3*). Besides the characteristic development of fever, which is caused by systemic pro-inflammatory cytokine release (*3*), metabolic disorders like hypoglycemia and hyperlactatemia also occur in patients with malaria. High lactate plasma levels are related to the development of acidosis, which is an important prognostic factor in patients with severe malaria (*4–9*).

Successful immunity to *Plasmodium* infection requires the participation of monocytic cells, which have a central role in sensing and phagocytosing parasitized red blood cells (*10*). In experimental models, monocytes are the major cells responsible for secreting pro-inflammatory pyrogenic cytokines, such as IFN-γ, IL-1β and TNF, during acute malaria episodes (*11–13*). Among these pro-inflammatory cytokines, TNF plays an important role in the pathophysiology of malaria (*14*). In other contexts, TNF has been identified as a major inducer of glucose uptake and metabolism in host immune cells (*14–17*). Glucose is metabolized inside cells through glycolysis to generate energy and biosynthetic intermediates for cell growth, activation and proliferation (*18, 19*). Moreover, the numerous intermediate metabolites that participate in the glycolysis-mediated conversion of glucose into pyruvate also play important roles in maintaining the activity and effector functions of innate immunity cells (*20*).

The hypoxia-inducible factor 1 alpha (HIF-1α) has emerged as one of the central regulators of inflammation and glucose metabolism in myeloid cells (*21*). Under normoxic conditions, HIF-1α is hydroxylated and degraded by the proteasome. Under hypoxic conditions, such as high levels of reactive nitrogen intermediates (RNI) and altered mitochondria function and release of reactive oxygen species (ROS), the activity of prolyl-hydroxylase domain (PHD) enzymes is

inhibited, leading to stabilization of HIF-1α, and subsequent translocation to the

nucleus (*22, 23*). In addition, the expression of HIF-1α is upregulated in response to

TNF. Consequently, HIF-1α activity in macrophages confers resistance to infection

in different experimental models (*21, 24–28*).

However, it is unclear how glucose metabolism, HIF-1α and their interplay

with TNF affects innate immune cells and host resistance to *Plasmodium* parasites.

Therefore, we investigated the involvement of TNF, RNI and HIF-1α in regulating

glucose metabolism in monocytic cells and if this mechanism is relevant for host

resistance and disease tolerance in experimental malaria. Our findings

demonstrate that both TNF and inducible nitric oxide synthase (iNOS) promotes

the expression and stabilization of HIF-1α enhance expression of glucose

transporter GLUT1 and glycolysis in monocytic cells from *P. chabaudi-* (*Pc-*)

infected mice. Altogether, our results reveal that this metabolic shift in monocytes

has an important role in regulating the host energy metabolism, resistance to

infection and disease outcome in an experimental malaria model.

Results

TNF signaling modulates signs of disease and energy expenditure in *P. chabaudi*-infected mice.

TNF is involved in the development of protective immunity and clinical signs of malaria both in humans and experimental models (*29–31*). We first analyzed the time course of parasitemia in *Pc*-infected C57BL/6 mice and the blood glucose levels measured daily at the end of the dark cycle. Parasites were detected in the circulation, starting at 3 days post-infection (dpi), and the peak of parasitemia was observed at 8 dpi (Figure 1A). Blood glucose levels significantly decreased at 8 dpi, coinciding with the peak of parasitemia. At later time points, parasitemia decreased, and glycaemia returned to homeostatic levels (Figure 1B). As previously demonstrated (*32*), the circulating levels of TNF and expression of TNF mRNA in the liver peaked at 6 am (end of dark cycle) at 8 dpi (Figure 1C and 1D), and has been reported to peak between days 6 and 10 post-infection, with a consistent pattern observed on days 6 and 8 (*32*). We also observed that the percentage of iRBCs was higher in TNFR$^{-/-}$ (TNFR p55p75$^{-/-}$) compared to C57BL/6 mice (Figure 1E), as previously reported (*31*). In addition, while infected C57BL/6 mice displayed lower temperature in response to infection, the rectal temperature of infected and uninfected TNFR$^{-/-}$ mice was similar (Figure 1F). Furthermore, the circulating levels of MCP1, TNF and IFN-γ were significantly lower in TNFR1$^{-/-}$ (TNFR p55$^{-/-}$) compared to C57BL/6 mice (Figures 1G-I), whereas no difference was observed in IL-10 levels (Figure 1J). We next evaluated glucose homeostasis during *Pc* infection. Infected C57BL/6 mice exhibited increased glucose uptake, which was associated with the development of hypoglycemia at the peak of infection. In contrast, this metabolic alteration was not observed in TNFR$^{-/-}$ mice, indicating that TNF signaling contributes to infection-induced changes in glucose metabolism and the subsequent development of hypoglycemia (Figure 1K). These results further indicate that TNF signaling promotes the systemic release of pro-inflammatory cytokine and plays a key role in controlling parasitemia, along with signs of disease in *Pc*-infected mice.

To further address the role of TNF signaling in promoting disease signs in
Pc-infected mice, we evaluated alteration on host energy metabolism, as indicated
in physical activity, food intake, overall energy expenditure and respiratory
exchange rate in naïve or *Pc*-infected C57BL/6 and TNFR[-/-] mice during 24h, at the
8 dpi. We observed that in naïve animals, all these parameters were similar in
TNFR[-/-] and C57BL/6 mice (Figures 2A-D, top panels and Figures 2E-H). In
contrast, all the evaluated parameters were decreased in infected C57BL/6 mice
compared to their naïve counterparts during the light and dark cycles. When we
analyzed only infected mice, the alterations in all parameters were milder in TNFR[-/-]
/- compared to C57BL/6 mice (Figures 2A-D bottom panels and 2E-H).
We then asked whether *Pc* infection might also affect body fat mass and
tissue glucose uptake. We observed a major depletion of fat mass in infected
C57BL/6 mice at 9 dpi, which was less pronounced in TNFR[-/-] animals (Figure 2I),
while the percentage of lean mass was not altered in either wild type or TNFR[-/-]
mice (Figure 2J). We also found that *Pc* infection resulted in a marked increase in
glucose uptake in the liver and spleen of C57BL/6 mice, unlike skeletal muscle or
white and brown adipose tissue (Figure 2K). The increase in these organs was not
seen in TNFR[-/-] animals (Figure 2L). Altogether, these results demonstrate the role
of TNF in regulating glucose and energy metabolism in *Pc*-infected mice and,
consequently, with important implications in the pathophysiology of experimental
malaria.
173

Hepatic non-parenchymal cells display enhanced GLUT1 receptor expression
and shift to glycolytic metabolism following *P. chabaudi* infection.

The liver has a key role in regulating host glucose metabolism (*33*).
Therefore, we asked whether *Pc* infection alters the expression of genes related to
carbohydrate metabolism in the liver. There was an overall enhancement in the
expression of genes related to glycolysis and a reduction in the expression of
genes associated with the Tricarboxylic (TCA) cycle and gluconeogenesis (Figure
3A). Enhanced glycolysis relies on the heightened glucose uptake by cells,
mediated by different glucose carriers (*34, 35*). However, the expression of GLUT2

(*Slc2a2),* GLUT5 *(Slc2a5)* and GLUT9 *(Slc2a9)* was decreased, whereas of

GLUT1 (*Slc2a1),* GLUT3 (*Slc2a3)* and GLUT6 *(Slc2a6)* expression was increased

in the liver of C57BL/6 infected, as compared to naïve mice (Figure 3A). Among

the transporters whose expression was increased, GLUT3 is expressed mainly in

neurons (*36*), while *Slc2a6* GLUT6 has been shown to be located in lysosomes and

does not mediate glucose uptake (*37*). GLUT1, on the other hand, is a ubiquitously

expressed and highly effective transporter of glucose. Also, GLUT1 is the most

well-characterized surface glucose transporter in immune cells (*18, 38, 39*).

Therefore, we next quantified GLUT1 protein, and, as expected, an enhancement

in GLUT1 levels was found in the liver of *Pc*-infected C57BL/6 mice, as compared

to the uninfected controls (Figure 3B).

Hepatocytes have an important role in glucose uptake from the circulation,

and they do this majorly through GLUT2 (*38*), whose mRNA expression was

downregulated (Figure 3A). Since GLUT1 expression can also be induced in

hepatocytes (*18*), we evaluated if GLUT1 expression was enhanced in hepatocytes

and non-parenchymal cells from the liver of infected mice. We found that at 8 dpi,

GLUT1 expression was not altered in hepatocytes (Figure 3C), but increased in

non-parenchymal cells (Figure 3D). We have also assessed the metabolic activity

of these same hepatic cell populations by quantifying their extracellular acidification

rate (ECAR) and oxygen consumption rate (OCR), which are indicatives of how

much glucose is being metabolized through glycolysis and the TCA cycle,

respectively. Consistent with GLUT1 expression, we observed no change in the

metabolic profile of hepatocytes (Figure 3E) but an increase in ECAR of non-

parenchymal cells (Figure 3F) from the liver of *Pc*-infected mice.

CD11b[+] cells from the spleen shift towards glycolytic metabolism in

response to *P. chabaudi* infection.

The spleen is the main tissue involved in immunity to *Plasmodium* infection,

where the response of immune cells to infection results in remarkable

splenomegaly (*40*). Moreover, the magnitude of glucose uptake in spleens was

higher than in the livers of infected mice (Figure 2K). Hence, we next asked

whether the increased tissue glucose uptake in experimental malaria was associated with the enhancement of GLUT1 expression and glycolysis in different subsets of immune cells. The levels of GLUT1 in spleens from *Pc*-infected C57BL/6 mice were also increased at 8 dpi (Figure 3G). We assessed GLUT1 expression in CD4$^+$ and CD8$^+$ T lymphocytes, B lymphocytes and CD11b$^+$F4/80$^+$/CD11c$^+$/Ly6g$^-$ myeloid cells by flow cytometry (gated as shown in Supplemental 1A and 1B). We found that the CD11b$^+$ subset and, to a lesser extent, CD4$^+$ T cells, but not CD8$^+$ T and B lymphocytes, displayed increased expression of GLUT1 in response to infection (Figures 3G, 3H and Supplemental 1C). Importantly, the magnitude of GLUT1 expression in monocytic cells was ~410%, whereas in CD4$^+$ T cells was ~29% (Figure 3H). In accordance with the expression of GLUT1 in those cell populations, the metabolic profiles of T and B lymphocytes were not altered (Figure 3I). In contrast, splenic CD11b$^+$ cells displayed enhanced glycolysis in infected mice as compared to naive mice (Figure 3J).

We have previously demonstrated that infection with *P. berghei ANKA* induces a marked increase in monocyte-derived dendritic cells (MO-DCs) in the spleens of mice (*41, 42*). We then analyzed splenic CD11b$^+$F4/80$^+$ cells (which are ≥90% CD11c$^+$Ly6G$^-$) (Supplemental 1), and selected monocytes (G1: DC-SIGN$^-$ MHCII$^-$), inflammatory monocytes (G2: DC-SIGN$^+$MHCII$^-$) and MO-DCs (G3: DC-SIGN$^+$MHCII$^+$). As shown in Figures 3K and 3L, *Pc* infection resulted in increased frequencies of MO-DCs, and reduced percentages of inflammatory monocytes and monocytes in spleens. At 8 dpi, expression of GLUT-1 was increased in inflammatory monocytes and MO-DCs but not in monocytes (Figure 3 M). Moreover, when we compared cells from infected mice, the levels of GLUT1 were similar between MO-DCs and inflammatory monocytes, but significantly higher in MO-DCs compared to monocytes (Figure 3M). Altogether, our results suggest that activated monocytes are the primary cells responsible for increased glucose uptake in the spleens in *Pc*-infected mice*.*

TNF signaling induces glycolysis in tissues from *P. chabaudi*-infected mice.

As shown above, TNF signaling promotes glucose uptake in livers and
spleens from *Pc*-infected mice (Figure 2K). We next asked whether TNF modulates
glucose metabolism in cells from these same organs. We observed that, except for
Hexokinase-3, the expression of mRNA from glycolytic enzymes (Hexokinase-1,
PFKP and PKM) was increased in C57BL/6 but not TNFR$^{-/-}$ 8-dpi (Figure 4B-E),
whereas the signature of gluconeogenesis enzymes (G6PC, G6PC3, FBP1, PCK1
and PCK2) mRNA remained unchanged (Figures 4F-J). The expression of GLUT2,
the main glucose transporter of hepatocytes, was similar in naïve and infected
mice, both in C57BL/6 and TNFR$^{-/-}$ mice (Figure 4K). The expression of GLUT1
protein mirrored the mRNA expression of glycolytic enzymes (Figure 4K and L),
and the increased expression of GLUT1 was also lower in splenic CD11b^{+} cells
from *Pc*-infected TNFR$^{-/-}$ mice (Figure 4L). To directly assess the role of TNF
during infection in this phenotype, splenocytes were stimulated with TNF *in vitro*,
which reproduced the effects observed during infection, supporting a TNF-
dependent regulation of GLUT1 expression (Figure 4M). In agreement with the
reduced levels of GLUT1, splenic CD11b^{+} cells from TNFR$^{-/-}$ infected mice
displayed reduced ECAR when compared with CD11b^{+} from infected wild-type
animals (Figure 4N), denoted as lower basal and compensatory glycolysis (Figure
4O).

To further confirm that the observed effects were due to TNF signaling in
myeloid cells we generated mice with conditional deletion of TNF receptor 1 in
lysozyme M- expressing cells (*TNFR1$^{\Delta Lyz2}$*). As expected, we observed a higher
parasitemia in *TNFR1$^{\Delta Lyz2}$* mice than in WT animals (Figure 4P). Moreover,
TNFR1$^{\Delta Lyz2}$ infected mice did not decreased the rectal temperature (Figure 4Q) and
blood glucose levels (Figure 4R). These data demonstrate that during experimental
malaria, TNF signaling plays a key role in inducing GLUT1 expression and glucose
metabolism in myeloid cells.

TNF and HIF-1α signaling crosstalk regulates glycolytic metabolism in
myeloid cells from *P. chabaudi*-infected mice.

The hypoxia-inducible factor 1 alpha (HIF-1α) is an oxygen-regulated
transcriptional activator that plays essential roles in mammalian development,
metabolism and pathogenesis of several diseases (*43*). Its functions are primarily
associated with the reprogramming of cellular energetic metabolism in response to
hypoxic conditions, but in innate immune cells, HIF-1α can also promote glycolysis
and quickly generate ATP and intermediate metabolites that fuel the activation and
release of pro-inflammatory cytokines (*43*). Because of that, we next asked if HIF-
1α is involved in the TNF-induced reprogramming of glucose metabolism in
immune cells during *Pc* infection. We observed a higher nuclear expression of HIF-
1α in the spleen and liver of infected mice; however, the levels of HIF-1α were
lower in livers of TNFR[-/-] compared to C57BL/6 infected mice (Figure 5A-B).

We next assessed the importance of the HIF-1α pathway for host resistance
to malaria by using mice with conditional deletion of HIF-1α in myeloid cells (*HIF-
1a[ΔLyz2]*) and their WT counterparts (*HIF-1a[fl/fl]*). It has been previously described that
HIF-1α activity can induce the expression of GLUT1 in different mammalian cell
populations (*20*). We found that glycolysis (ECAR) and GLUT1 expression were
impaired, though partially, in monocytic and splenic CD11b+ cells from infected
HIF-1aΔLyz2 mice (Figures 5C-5E) compared to infected WT mice. The level of
GLUT1 expression that is still maintained is likely due to other host or parasite
factors, such as IFN-γ (*44*).
Consistent with these findings, stimulation of splenic cells with TNF *in vitro* induced
a similar pattern, further supporting the link between TNF signaling and HIF-1α
activation (Figure 5D). Consistent with the impaired GLUT1 expression, CD11b□
splenocytes isolated from HIF-1αΔLyz2 infected mice exhibited a marked decrease
in ECAR compared to CD11b□ cells from infected control animals (Figures 5E and
F). Importantly, we observed a higher parasitemia in *HIF-1a[ΔLyz2]* mice than in WT
animals (Figure 5G and Supplemental 2). At 8dpi with *Pc*, the parasitemia was
higher and the circulating levels of TNF lower in *HIF-1a[ΔLyz2]* mice compared to WT
animals (Figures 5G and 5H). In addition, *Pc*-infected *HIF-1a[fl/fl]* mice splenocytes
displayed enhanced TNF secretion following LPS stimulation compared to cells
from *HIF-1a[ΔLyz2]* infected animals (Figure 5I). Altogether, these data demonstrate

that the induction of the HIF-1a in myeloid cells is important for inducing glycolysis,
TNF production and control of *Pc* infection.

iNOS expression promotes HIF-1α stability, glycolytic metabolism and
modulates signs of disease in *P. chabaudi*-infected mice

Our next step was to understand how TNF promotes HIF-1α expression. It is
known that RNI induces HIF-1α expression in particular by enhancing the stability
of this transcription factor through a mechanism that inhibits its degradation (*45*).
TNF is also widely described as a major inducer of iNOS expression and RNI
release by myeloid cells, such as monocytes (*46*). As expected, we observed that
splenocytes from *Pc*-infected TNFR[-/-] mice stimulated *ex vivo* with LPS released
lower levels of RNI than LPS-stimulated cells from C57BL/6-infected mice (Figure
6A). Importantly, iNOS-deficient mice presented higher parasitemia than C57BL/6
animals at 8dpi with *Pc* (Figure 6B), as previously described (*47*). Similar to what
was observed in TNFR[-/-] animals, infected iNOS[-/-] mice did show lower temperature
at 8dpi with *Pc* (Figure 6C). Additionally, at this same time point, we did not
observe hypoglycemia in the iNOS[-/-] mice infected with *Pc* (Figure 6D). As
expected, splenic CD11b[+] cells from iNOS[-/-] infected mice displayed higher OCR
when compared to C57BL/6 infected with *Pc* (Figure 6E), indicating increased
basal and maximal mitochondrial respiratory capacities in the absence of iNOS
(Figure 6F).

In accordance with the results obtained in *Pc*-infected TNFR[-/-] mice, iNOS
deficiency resulted in reduced hepatic expression of the glycolytic enzymes HK1,
HK3, PFKP and PKM (Figures 6 G-J) as well as reduced GLUT1 expression in
splenic monocytes (Figure 6K). TNF stimulation *in vitro* resulted in a similar profile
to that observed *in vivo*, further supporting a TNF–iNOS–dependent regulation of
glycolytic pathways (Figure 6L). Accordingly, glycolytic metabolism in splenic
monocytic cells of iNOS[-/-] infected animals was decreased compared to those of
C57BL/6 infected mice, as denoted by a lower ECAR (Figure 6M). Finally, to
directly link glycolytic metabolism and TNF production during *Pc* infection, we
inhibited glycolysis *in vivo* using 2-deoxy-D-glucose (2-DG). Treatment with 2-DG

resulted in a marked increase in parasitemia compared to untreated infected mice
(Figure 6O), resembling the impaired parasite control observed in HIF-1αΔLyz2,
TNFR[-/-] and iNOS[-/-] mice. Consistent with effective glycolytic blockade, 2-DG–
treated mice displayed reduced lactate levels during infection (Figure 6N and O).
Thus, the inhibition of glycolysis mirrored the metabolic and immunological
alterations observed upon disruption of the TNF–HIF-1α–iNOS axis, supporting the
conclusion that this pathway is critical for sustaining glycolytic metabolism, TNF
production and effective parasite control during *Pc* infection. Altogether, these data
demonstrated that RNI induces HIF-1α expression, glycolysis, and TNF release by
monocytic cells, leading to control of parasitemia but also promoting clinical signs
of disease, such as hypothermia and hypoglycemia, in *Pc*-infected mice.
348

Discussion

Malaria disease manifestations include severe metabolic changes in the
host organism. Different studies report that decreased glycaemia and increased
lactate plasma levels followed by acidosis are common manifestations in patients
with severe malaria (*5, 48–50*). However, the mechanisms that lead to
hypoglycemia and increased lactate levels during malaria are poorly understood. A
better understanding of the mechanisms that mediate host metabolic alterations
and the connections of such events with the immune response against the parasite
may provide new insights for therapeutic interventions in malaria patients. In this
study, we found that TNF signaling mediates changes in host energy metabolism,
accompanied by increased expression of GLUT1 and enhanced glycolysis in
monocytes. In addition, we found that TNF-induced RNI stabilizes the transcription
factor HIF-1α, which promotes both a metabolic shift towards glycolysis and the
expression of pro-inflammatory cytokines by monocytes. Furthermore, we report
that TNF, iNOS and HIF-1α have an important role in controlling *Pc* infection and
signs of systemic inflammation in this mouse malaria model.

Previous studies have shown that TNF induces the production of RNI
through the upregulation of iNOS via the NF-κB pathway (*51–54*). TNF-mediated
iNOS expression is critical for RNI production, which in turn stabilizes HIF-1α by

inhibiting prolyl hydroxylases (PHDs) even under normoxic conditions (*52*). HIF-1α

then upregulates the expression of glycolytic genes, including GLUT1 (*55–57*).

TNF has been described as a critical mediator in malaria, driving cytokine release

and parasitemia control (*58*). It also enhances glucose uptake in tissues, aligning

with our findings of increased glycolysis in monocytes (*59*). The role of iNOS in

malaria is well documented. IFN-γ and TNF induced RNI inhibits parasite growth,

but can cause tissue damage and organ dysfunction, especially in severe malaria

(*60*). Recent studies also highlight the complexity of glycemia regulation during

Plasmodium infection describing its role in modulating parasite virulence and

transmission (*61*). These studies demonstrate the critical function of TNF and iNOS

in immune responses against *Plasmodium*. Aligning with our findings that this axis

and metabolic rewiring are essential for monocyte activation and outcome of *Pc*

infection

Interestingly, we found that glucose uptake was significantly increased in

spleens and livers of *Pc*-infected mice and that this effect was partially dependent

on TNF signaling. Consistently, TNF promoted increased expression of GLUT1

and glycolytic energy metabolism in non-parenchymal monocytic cells, but not

hepatocytes or lymphocytes in livers and spleens, respectively. These results

indicate that the hypoglycemia of *Pc*-infected mice is, at least in part, caused by

TNF-driven GLUT1-mediated enhanced glucose uptake and glycolysis in

monocytes. Insulin is the major hormone responsible for the upregulation of

glucose uptake in several organs, including adipose and skeletal muscle tissues,

which happens mainly through the upregulation of GLUT4 (*62, 63*). In mice, GLUT1

expression is recognized as independent of insulin, in contrast to GLUT4 (*64*). In

our model, this regulation appears to be driven by pro-inflammatory cytokines,

particularly TNF. Supporting this, our results show that *in vitro* stimulation with

TNF, significantly increases GLUT1 expression in monocytes, accordingly to the *ex*

vivo phenotype observed in infected animals. Altogether, our findings suggest that

the increase in glucose uptake and glycolysis in monocytes is insulin-independent.

Although the frequency of MO-DCs increases during infection, other cell

populations may also contribute to glucose consumption. Further experiments,

including the assessment of GLUT1 function in these populations, are needed to
clarify their relative contribution to glucose consumption during infection

Following the rupture of parasitized red blood cells, there is a release of
pathogen and danger-associated molecular patterns that activate Toll-like
receptors (TLRs), Nod-like receptors (NLRs), and the cyclic GMP–AMP (cGAS)
synthase (*10, 65*). Activation of these innate immune receptors and sensors triggers
the production of high levels of pro-inflammatory cytokines, such as TNF, IL-6, IL1β
and IL-12, all of which participate in host immunity against *Plasmodium* infection
(*66*). The importance of the glycolytic pathway for innate immune cell activation has
been demonstrated in different studies (*20, 67*). For instance, monocyte
differentiation from a resting to an inflammatory state requires a shift in energy
metabolism to high glucose consumption and rapid energy generation by glycolysis
(*68*). Activated monocytes, along with dendritic cells, then produce IL-12, which is
critical for the differentiation of Th1 lymphocytes that produce IFN-γ, a cytokine that
activates splenic macrophages to eliminate *Plasmodium*-infected red blood cells
(*10, 66*).

Importantly, we found that similarly to TNFR[-/-] mice, iNOS-deficient mice
display impaired glycolytic metabolism in monocytic cells from *Pc*-infected mice.
RNI are produced in macrophages once they receive two signals. The first signal is
the priming with IFN-γ. The second signal is the activation of monocytes with TNF
induced by microbial products via TLRs. Like in TNFR[-/-] mice, iNOS deficiency
resulted in increased parasitemia following *Pc* infection (*69, 70*). HIF-1α is required
for the optimal expression of many glycolytic genes in macrophages, including
those encoding GLUT1, hexokinase, phosphofructokinase and pyruvate kinase
(*43*). Recently, it was suggested that RNI promotes *S*-nitrosylation at the Cys533
residue of HIF-1α molecules (*71*), promoting HIF-1α stabilization by direct
inactivation of PHDs (*72*). Indeed, the impairment of RNI release during *Pc*
infection, resulted in reduced translocation of HIF-1α to the nuclei of splenic
monocytes and non-parenchymal liver cells, which was markedly increased in
infected WT mice. Therefore, we propose that the mechanism by which TNF

signaling regulates HIF-1α stabilization/expression during *Pc* infection is mediated
by RNI.

Different studies have demonstrated that HIF-1α is critical for host
resistance to different pathogens (*73*), but both the participation of this transcription
factor in malaria as well as the role of glycolytic metabolism in resistance to
Plasmodium infection have not been explored. The *HIF-1α*$^{\Delta Lyz2}$ animals, which are
genetically deficient for HIF-1α and have impaired glycolysis only in myeloid cells,
were more susceptible to *Pc* infection, further demonstrating the importance of
myeloid cells in controlling parasitemia. This might be explained by the importance
of glycolysis for the release of TNF and other pro-inflammatory cytokines by
monocytes that are activated during *Plasmodium* infection (*3, 40*). Although we
have found that *Pc* infection-induced increase in nuclear translocation of HIF-1α
was impaired in the absence of TNF signaling, *HIF-1α*$^{\Delta Lyz2}$ animals also displayed
lower TNF secretion in response to *Pc* infection than WT mice. Therefore, these
results indicate that the in vivo relationship between TNF signaling and HIF-1α-
driven glycolysis is complex and that a positive feedback loop between TNF
production, HIF-1α expression and glycolysis must exist during *Pc* infection.

The link between TNF and HIF has been explored in different cell types.
Previously, studies demonstrated a crosstalk between NF-kB and HIF-1α. TNF
signaling recruits different intracellular adaptors that activate multiple signal
transduction pathways. One consists of the TRAF family of proteins, which can
lead to IKK-dependent activation of NF-kB, thereby promoting inflammatory
responses (*74*). Conversely, HIF-1α can also regulate NF-kB activation.
Koedderitzch and collaborators (*75*) have recently shown that TNF induces
glycolytic shift via GLUT-1 and HIF-1α in rheumatoid arthritis (RA). As with malaria,
TNF is a pivotal cytokine involved in the pathogenesis of RA. *In vitro* treatment of
fibroblast-like synoviocytes with TNF induces upregulation of GLUT-1 and HIF-1α,
which depends on TAK1-induced NF-kB activation downstream of TNF receptor 1
(*75*). In addition, NF-kB plays a central role in inflammatory responses,
orchestrating the expression of TNF, IL-6, adhesion molecules and enzymes.

Moreover, an NF-kB binding site is present in the proximal promoter site of the
HIF-1α gene, indicating that NF-kB activation can regulate its expression (*76, 77*).

Plasmodium parasites do not synthesize glucose, being solely dependent on
host glucose as a source of energy required for parasite replication. On the one
hand, the increased glucose uptake and subsequent glycolytic metabolism in
monocytes promote hypoglycemia, helping to starve the erythrocytic stage of the
parasite, controlling its replication. On the other hand, hypoglycemia may also be
detrimental to the host. For instance, lactate, the final product of glycolytic
metabolism of glucose, when secreted in high amounts, causes acidosis with
decreased blood pH and severe consequences to the host, such as extreme
fatigue, pain, overall feelings of physical discomfort and decreased appetite (*5*).

Indeed, we found that the absence of TNF signaling and iNOS[-/-] mice
reverted the development of hypothermia, which is a known sickness manifestation
in *Pc* infection-induced experimental malaria in mice (*78–80*). Moreover, the
attenuation of clinical symptoms in TNFR[-/-] infected mice extended beyond thermal
regulation. Despite increased parasitemia, these animals displayed restored
physical activity, food consumption, energy expenditure and respiratory exchange
rate compared to WT animals. Although restored physical activity, food
consumption and energy expenditure in knockout mice may contribute to the
observed systemic metabolic parameters by altering energy balance, these effects
are not mutually exclusive with the TNF-driven, cell-intrinsic metabolic mechanisms
described here. These findings demonstrate that TNFR[-/-] mice are more "tolerant"
to disease. Hence, our results allow us to conclude that although TNF promotes
host resistance during malaria, it plays a detrimental role during *Plasmodium*
infection.

In summary, our results indicate that TNF-induced RNI induces HIF-1α-
mediated glycolysis and pro-inflammatory response by monocytes, promoting host
resistance to infection with *Plasmodium*. However, this exacerbated activation of
monocytes may be detrimental to the host. Hence, our findings enlighten a
fundamental mechanistic question related to the pathophysiological role of

Plasmodium infection in the vertebrate host. These findings may provide insights
for developing novel therapeutic interventions to treat this devastating disease.
491
492

Materials and Methods

Ethics statement

495 All experiments were carried out in accordance with institutional guidelines for
496 animal ethics and approved by the Ethics Committee on Animal Use of Ribeirão
497 Preto Medical School University of São Paulo (CEUA 94 / 2020), Institutional
498 Ethics Committees from Oswaldo Cruz Foundation (Fiocruz-Minas,
499 CEUA/LW15/14, and LW16/18) and UMMS (IACUC A-1369-14-5), respectively.
500
501

Mice

C57BL/6 and TNFR$^{-/-}$ (TNFR p55/p75 chains double knockout mice) were obtained
from the Center for Breeding of Transgenic Mice from the Ribeirao Preto Medical
School. Conditional knockout mice (*TNFR1*$^{\Delta Lyz2}$) were generated by crossing
LysMCre$^{+/-}$ to *TNFR1*^{flfl} mice, which were maintained on a C57BL/6 genetic
background. Mice used in experiments were sex and age-matched: female mice
between 8–12 weeks of age. All mouse lineages were housed in micro-isolators in
a maximum number of six mice per cage, in a specific pathogen-free facility at the
Oswaldo Cruz Foundation, or Ribeirão Preto Medical School or University of
Massachusetts Medical School, under controlled temperature (22–25°C) and 12-h
light-dark cycle and provided with water and food ad libitum.
513

Experimental Infections

Plasmodium chabaudi chabaudi AS strain (*Pc*) was used for experimental
infections (*81, 82*). First, *Pc* was maintained in C57BL/6 mice by serial passages
once a week up to eight times. For experimental infection, mice were injected
intraperitoneally (i.p.) with 10^5 infected red blood cells (iRBCs) diluted in 100 µl
PBS-1X. The percentage of RBCs containing parasites (parasitemia) was

measured in the blood with Giemsa at different days post-infection (dpi). Infected
and non-infected mice were treated daily with a glycolytic inhibitor 2-deoxy-D-
glucose (2-DG) at a dose of 250 mg/kg, administered from day 4 to day 15 post-
infection (dpi). Control groups, including both infected and non-infected animals,
received PBS 1X following the same treatment schedule.

Glucose Measurement
The glucose uptake was measured using 2-[14C]-deoxyglucose. Experiments were
performed at the National Mouse Metabolic Phenotyping Center (MMPC) at
UMASS Medical School. At 6 am on day 8 dpi, basal glucose uptake in individual
organs was measured using an intravenous injection of 2-[14C]-deoxyglucose.
After 1h, mice were anesthetized, and tissue samples were taken for organ-specific
levels of 2-[14C]-deoxyglucose-6-phosphate. Blood glucose levels were measured
with an *Accu-chek* glucometer.

Mouse RNA-Seq
RNA-seq was performed in biological replicates (3 mice per group). Liver samples
were collected from C57BL/6 mice at different time points: 12 am; 6 am; 12 pm,
and 6 pm from mice at day 8 post-infection with *P. ch.* or uninfected control. RNA-
seq libraries were prepared using the TruSeq Stranded mRNA Kit (*Illumina*)
following the manufacturer's instructions. Briefly, poly-A-containing mRNA
molecules were purified using poly-T oligo-attached magnetic beads and
fragmented using divalent cations. The RNA fragments were transcribed into cDNA
using SuperScript II Reverse Transcriptase (*Invitrogen*), followed by second strand
cDNA synthesis using DNA Polymerase I and RNase H. Finally, cDNA fragments
then have the addition of a single 'A' base and subsequent ligation of the adapter.
The products were then purified and enriched by PCR using paired-end primers
(*Illumina*) for 15 cycles to create the final cDNA library. The library quality was
verified by fragmentation analysis (*Agilent Technologies 2100 Bioanalyzer*) and
submitted for sequencing on the Illumina NextSeq 500 (*Bauer Core Facility*

Harvard University). These samples were collected in according to the circadian
cycle published by our group (*32*).

RNA extraction and Real-time PCR

Liver samples were collected from C57BL/6 mice at 8 dpi with *P. ch.* or uninfected
controls. Tissue fragments were stabilized   in RNAlater (Qiagen) and   RNA
extraction was performed using an RNeasy kit (Qiagen) according to manufacturer
instructions. A total of 1mg of RNA was converted to cDNA using iScript cDNA
Synthesis Kit (Bio-Rad). Real-time PCR was performed with the SYBR Green
master mix (Bio-Rad). Primer sequences are: CCCTCACACTCAGATCATCTTCT
(Foward   primer)   and   GCTACGACGTGGGCTACAG   (Reverse   primer)
(NM_013693).

Isolation of splenocytes and purification of CD11b$^+$, B and T cells

Spleens from control and infected mice at 8 dpi were dissociated through a 100-µm
nylon cell strainer for obtaining single-cell suspensions. For *ex vivo* cultures, known
concentrations   of   splenocytes   were   resuspended   in   RPMI   1640   medium
supplemented with penicillin, streptomycin, and 10% fetal bovine serum (FBS)
(*Gibco, ThermoFisher*). In some experiments, splenocytes (2 × 10^6 cells) were
stimulated in vitro with 1 µg/ml of LPS for 24h, or TNF-α [20ng/ml] or not during 15
minutes. For CD11b+ cell-, B cell- or T cell- purification, immunomagnetic beads
for positive selection of CD11b$^+$ cells, B cells and T cells (*Miltenyi Biotec*) were
used, respectively, according to manufacturer's instructions.

Hepatocyte's isolation

576   First, the liver portal vein was cannulated with a 24G catheter. The inferior cava
577   vein was cutted to allow perfusion flow. Perfusion was done with 50 mL of Hanks A
578   and then 25 mL of Hanks B with 20 mg of collagenase. After that, tissue was
579   dissociated   and   samples   were   filtered   in   a   40   mM   membrane.   Cells   were

centrifuged: 60x g (5 min, 4oC), disposing of supernatants, 60x g (5 min, 4oC), and resuspended in cold RPMI 10% SFB.

Liver non-parenchymal-cells

Livers were first digested in a collagenase solution (5 mg/liver in 10 mL of RPMI 2% SFB) at 37oC for 35 min. Then, the solution was added PBS/BSA 0,5% + EDTA in order to inactivate collagenase and cells were differentially separated by centrifugation: 300 x g (10 min, 4oC) disposing supernatants, 60 x g (3 min, 4oC) twice preserving supernatants and finally, 300 x g (10 min, 4oC) disposing supernatants. The samples were filtered through 100 µm cell strainers and centrifuged for 300 x g (10 min, 4oC). Red blood cells were lysed using ACK (ammonium chloride potassium) buffer. Concentrations of single-cell suspensions were then adjusted following counting on the hematocytometer.

Western blotting

Mouse splenocytes, magnetically purified CD11b☐ splenic cells, isolated hepatocytes, isolated liver non-parenchymal cells, or total liver samples from infected or non-infected mice were lysed using RIPA buffer (Sigma) supplemented with protease and phosphatase inhibitor cocktail (Thermo Fisher Scientific).After 15 min on ice, lysates were centrifuged at 13,000 × g for 10 min at 4 °C. The proteins were separated in a 10%-acrylamide NUPAGE *Bis-tris* Protein gels (*Invitrogen*) and transferred onto nitrocellulose membranes. Membranes were blocked with 5% (wt/vol) nonfat milk (*Molico*) in Trisbuffered saline with 0.1% Tween-20 (TBST) for 1 h at room temperature and then incubated overnight at 4°C with primary antibodies. The membranes were incubated with GLUT1 (*Abcam*, 1:1000), GLUT2 (*Abcam*, 1:1000), Na,K-ATPase (*Cell Signaling, 1:2000)* and β-actin (Sigma, 1:5000) specific antibodies. Subsequently, membranes were repeatedly washed with TBST and then incubated for 2 h with the anti-Rabbit HRP-conjugated secondary antibody (1:30.000 dilution; *Sigma-Aldrich*). Immunoreactivity was detected with Clarity Max ECL Substrate (Biorad) and then the chemiluminescence

611 signal was recorded on the iBright FL1500 (*ThermoFisher*). Data were analyzed
612 with iBright FL1500.

Metabolic Profiling.

Metabolic profiling of CD11b$^+$ cells, B cells, T cells, hepatocytes, and non-
parenchymal cells was undertaken using a Seahorse XFe96 Extracellular Flux
Analyzer (*Agilent*) in microplates. Basal oxygen consumption rate (OCR) and
extracellular acidification production (ECAR) were measured by Phenotypic
Seahorse Kit (KIT 103325-100), or Seahorse XF Cell Mito Stress Tests (KIT
103015-100), or Glycolytic Rate Assay Kit (KIT 103344-100) according to the
manufacturer's instructions.

Flow cytometry

Splenocytes (2×10^6 cells) from mice at 0 (uninfected controls) and 8 days post-
infection were stained with fluorophore-labeled monoclonal antibodies (mAbs)
specific for cell surface markers. The following flow cytometry specific mAbs were
used: CD11c-PECy7 clone N418 (1:200), CD11b-APCCy7 clone M1/70 (1:200),
Ly6c PERCP clone H.K1.4 (1:200), F4/80- PE clone BM8 (1:200), Ly6g FITC clone
1A8 (1:200), CD45 Pacific Blue- clone30F-11 (1:200), CD3 FITC clone 145-2c11
(1:200), CD4 PE clone RM4-5 (1:200), CD8 PERCP clone 53-6.7 (1:200), CD19
PeCy7 clone 6D5(1:200) and Bv510 (Live/Dead kit, ThermoFisher). Intracellular
staining of GLUT1 Alexa Fluor 647 [EPR3915] (1:100) was performed following cell
fixation and permeabilization using the eBioscience Foxp3
Fixation/Permeabilization Kit. Data were acquired in a FACSCanto II machine (BD
Biosciences) and analyzed using FlowJo software (Tree Star).

Cytokine measurement.

638 Plasma was collected from whole blood and supernatants were collected from cell
639 cultures. Cytokine concentration in these samples were determined using CBA
640 mouse Inflammation Kit (BD^{TM}) or ELISA according to the manufacturers'
641 instructions (*Biolegend*).

Food Intake and Physical Activity

Mouse food intake and physical activity were assessed for 3 days prior to infection and at days 6, 7 and 8 dpi with *Pc* using metabolic cages (*TSE Systems*, Chesterfield, MO) located at the National Mouse Metabolic Phenotyping Center (MMPC) at UMASS Medical School. Mice were housed under controlled temperature and lighting, with food and water ad libitum. Food intake and physical activity monitoring was fully automated using the *TSE Systems LabMaster* platform. *LabMaster* cages allowed the use of bedding, thus, minimizing any animal anxiety during the experimental period. Physical activity was calculated based on quantitative measurement of horizontal and vertical movement (XYZ-axis). In addition, non-invasive measures of O_2 consumption and CO_2 production were used to calculate the respiratory exchange ratio to reflect energy expenditure.

RNA-Seq Analysis

Gene expression of each experimental group was shown at different time points 12 am; 6 am; 12 pm and 6 pm. The average expression of all time points was calculated for visualization in heatmaps. The heatmaps presented in Figures 1 and 2 were developed with a free version of the software Morpheus (https://software.broadinstitute.org/morpheus). For hierarchical clustering, we used linkage from the average expression of each gene in metric One Minus Pearson Correlation.

Statistical analysis

GraphPad Prism 7.0 software was used for statistical analysis. Multiple-group comparisons were performed with either one-way ANOVA (followed by Tukey's post hoc test) or two-way ANOVA (followed or not by Sidak's post hoc test). Unpaired two-tailed Student's t-test was used for the comparison of two conditions. Results are expressed as means ± SEM. P value ≤ 0.05 was considered significant.

References

1.   X. Z. Su, K. D. Lane, L. Xia, J. M. Sá, T. E. Wellems, Plasmodium genomics and genetics: New insights into malaria pathogenesis, drug resistance, epidemiology, and evolution. *Clin Microbiol Rev* 32, 1–29 (2019).

2.   *World Malaria Report 2022* (2022).

3.   C. A. Moxon, M. P. Gibbins, D. McGuinness, D. A. Milner, M. Marti, New Insights into Malaria Pathogenesis. *Annual Review of Pathology: Mechanisms of Disease* 15, 315–343 (2020).

4.   T. Planche, S. Krishna, Severe Malaria: Metabolic Complications. *Curr Mol Med* 6, 141–153 (2006).

5.   H. Possemiers, L. Vandermosten, P. E. Van Den Steen, Etiology of lactic acidosis in malaria. *PLoS Pathog* 17, 1–17 (2021).

6.   N. P. J. Day, N. H. Phu, N. T. H. Mai, T. T. H. Chau, P. P. Loc, L. Van Chuong, D. X. Sinh, P. Holloway, T. T. Hien, N. J. White, The pathophysiologic and prognostic significance of acidosis in severe adult malaria. *Crit Care Med* 28, 1833–1840 (2000).

7.   J. Hanson, S. J. Lee, S. Mohanty, M. A. Faiz, N. M. Anstey, P. Charunwatthana, E. Bin Yunus, S. K. Mishra, E. Tjitra, R. N. Price, R. Rahman, F. Nosten, Y. Htut, G. Hoque, T. T. H. Chau, N. H. Phu, T. T. Hien, N. J. White, N. P. J. Day, A. M. Dondorp, A simple score to predict the outcome of severe malaria in adults. *Clinical Infectious Diseases* 50, 679–685 (2010).

8.   J. A. Kraut, N. E. Madias, Metabolic acidosis: Pathophysiology, diagnosis and management. *Nat Rev Nephrol* 6, 274–285 (2010).

9.   E. M, S. R, W. C, M. M, O. J, L. B, E. Al., "Acidosis in severe childhood malaria" (1997).

10.   R. T. Gazzinelli, P. Kalantari, K. A. Fitzgerald, D. T. Golenbock, Innate sensing of malaria parasites. [Preprint] (2014). https://doi.org/10.1038/nri3742.

11.   C. A. Dinarello, Immunological and inflammatory functions of the interleukin-
1   family.   [Preprint]   (2009).
https://doi.org/10.1146/annurev.immunol.021908.132612.

12.   G. A. Awandare, P. Kempaiah, D. O. Ochiel, P. Piazza, C. C. Keller, D. J.
Perkins, Mechanisms of erythropoiesis inhibition by malarial pigment and malaria-
induced proinflammatory mediators in an in vitro model. *Am J Hematol* 86, 155–
162 (2011).

13.   K. C. Matteucci, A. S. Correa, D. L. Costa, Recent Advances in Host-
Directed Therapies for Tuberculosis and Malaria. 12, 1–23 (2022).

14.   G. L. Popa, M. I. Popa, Recent Advances in Understanding the Inflammatory
Response in Malaria: A Review of the Dual Role of Cytokines. *J Immunol Res*
2021, 10–12 (2021).

15.   Y. Sakurai, X. J. Zhang, A. R. Wolfe, TNF directly stimulates glucose uptake
and leucine oxidation and inhibits FFA flux in conscious dogs. *Am J Physiol*
Endocrinol Metab 270 (1996).

16.   Z. Spolarics, A. Schuler, G. J. Bagby, C. H. Lang, K. Meszaros, J. J. Spitzer,
Tumor necrosis factor increases in vivo glucose uptake in hepatic nonparenchymal
cells. *J Leukoc Biol* 49, 309–312 (1991).

17.   J. Ciesla, I. Moreno, J. Munger, TNFα-induced metabolic reprogramming
drives an intrinsic anti-viral state. *PLoS Pathog* 18, e1010722 (2022).

18.   A. L. McCall, Glucose transport. *Stress: Physiology, Biochemistry, and*
Pathology Handbook of Stress Series, Volume 3, 293–307 (2019).

19.   D. G. Ryan, L. A. J. O'Neill, Krebs cycle rewired for macrophage and
dendritic cell effector functions. *FEBS Lett* 591, 2992–3006 (2017).

20.   L. O'Neill, K. Rigel, J. Rathmell, A guide to immunometabolism for
immunologists. *Nat Rev Immunol.*, doi: 10.1038/nri.2016.70. (2016).

21.   T. Cramer, Y. Yamanishi, B. Clausen, I. Förster, R. Pawlinski, N. Mackman,
V. Haase, R. Jaenisch, M. Corr, V. Nizet, G. Firestein, H.-P. Gerber, N. Ferrara, R.
S. Johnson, HIF-1 Is Essential for Myeloid Cell Inflammation. *Cell* 112, 645–657
(2015).

22.   G. L. Semenza, Hypoxia-Inducible Factors in Physiology and Medicine. *Cell*
148, 399–408 (2012).

23.   A. J. Majmundar, W. J. Wong, M. C. Simon, Hypoxia-inducible factors and
the response to hypoxic stress. *Mol Cell* 40, 294–309 (2010).

24.   F. Li, J. Luo, H. Xu, Y. Wang, W. Jiang, K. Chang, S. Deng, M. Chen, Early
secreted antigenic target 6-kDa from Mycobacterium tuberculosis enhanced the
protective innate immunity of macrophages partially via HIF1α. *Biochem Biophys*
Res Commun 522, 26–32 (2020).

25.   Q. Li, Y. Xie, Z. Cui, H. Huang, C. Yang, B. Yuan, P. Shen, C. Shi,
Activation of hypoxia-inducible factor 1 (Hif-1) enhanced bactericidal effects of
macrophages to Mycobacterium tuberculosis. *Tuberculosis* 126 (2021).

26.   A. C. Codo, G. G. Davanzo, L. de B. Monteiro, G. F. de Souza, S. P.
Muraro, J. V. Virgilio-da-Silva, J. S. Prodonoff, V. C. Carregari, C. A. O. de Biagi
Junior, F. Crunfli, J. L. Jimenez Restrepo, P. H. Vendramini, G. Reis-de-Oliveira, K.
Bispo dos Santos, D. A. Toledo-Teixeira, P. L. Parise, M. C. Martini, R. E.
Marques, H. R. Carmo, A. Borin, L. D. Coimbra, V. O. Boldrini, N. S. Brunetti, A. S.
Vieira, E. Mansour, R. G. Ulaf, A. F. Bernardes, T. A. Nunes, L. C. Ribeiro, A. C.
Palma, M. V. Agrela, M. L. Moretti, A. C. Sposito, F. B. Pereira, L. A. Velloso, M. A.
R. Vinolo, A. Damasio, J. L. Proença-Módena, R. F. Carvalho, M. A. Mori, D.
Martins-de-Souza, H. I. Nakaya, A. S. Farias, P. M. Moraes-Vieira, Elevated
Glucose Levels Favor SARS-CoV-2 Infection and Monocyte Response through a
HIF-1α/Glycolysis-Dependent Axis. *Cell Metab* 32, 437-446.e5 (2020).

27.   D. Friedrich, D. Zapf, B. Lohse, R. A. Fecher, G. S. Deepe, J. Rupp, The
HIF-1/LC3-II axis impacts fungal immunity in human macrophages. *Infect Immun*
87 (2019).

28.   E. Riboldi, C. Porta, S. Morlacchi, A. Viola, A. Mantovani, A. Sica, Hypoxia-
mediated regulation of macrophage functions in pathophysiology. *Int Immunol* 25,
67–75 (2013).

29.   O. Agudelo, J. Bueno, A. Villa, A. Maestre, High IFN-gamma and TNF
production by peripheral NK cells of Colombian patients with different clinical
presentation of Plasmodium falciparum. *Malar J* 11, 1–7 (2012).

30.   L. N. Cruz, Y. Wu, H. Ulrich, A. G. Craig, C. R. S. Garcia, Tumor necrosis
factor reduces Plasmodium falciparum growth and activates calcium signaling in
human malaria parasites. *Biochim Biophys Acta* 1860, 1489 (2016).

31.   G. H. Long, B. H. K. Chan, J. E. Allen, A. F. Read, A. L. Graham, Blockade
of TNF receptor 1 reduces disease severity but increases parasite transmission
during Plasmodium chabaudi chabaudi infection. *Int J Parasitol* 38, 1073–1081
(2008).

32.   I. C. Hirako, P. A. Assis, N. S. Hojo-Souza, G. Reed, H. Nakaya, D. T.
Golenbock, R. S. Coimbra, R. T. Gazzinelli, Daily Rhythms of TNFα Expression
and Food Intake Regulate Synchrony of Plasmodium Stages with the Host
Circadian Cycle. *Cell Host Microbe* 23, 796-808.e6 (2018).

33.   E. Trefts, M. Gannon, D. H. Wasserman, The liver. *Current Biology* 27,
R1147–R1151 (2017).

34.   I. Vrhovac, D. Breljak, I. Sabólic, Glucose transporters in the mammalian
blood cells. *Period Biol* 116, 131–138 (2014).

35.   M. Mueckler, C. Caruso, S. A. Baldwin, M. Panico, I. Blench, H. R. Morris,
W. J. Allard, G. E. Lienhard, H. F. Lodish, Sequence and Structure of a Human
Glucose Transporter. *Science (1979)*, doi: 229(4717):941e945. (1985).

36.   I. A. Simpson, D. Dwyer, D. Malide, K. H. Moley, A. Travis, S. J. Vannucci,
The facilitative glucose transporter GLUT3: 20 Years of distinction. *Am J Physiol*
Endocrinol Metab 295, 242–253 (2008).

37.   S. Maedera, T. Mizuno, H. Ishiguro, T. Ito, T. Soga, H. Kusuhara, GLUT6 is
a lysosomal transporter that is regulated by inflammatory stimuli and modulates
glycolysis in macrophages. *FEBS Lett* 593, 195–208 (2019).

38.   H. Yaribeygi, F. R. Farrokhi, A. E. Butler, A. Sahebkar, Insulin resistance:
Review of the underlying molecular mechanisms. *J Cell Physiol* 234, 8152–8161
(2019).

39.   G. Litwack, Insulin and Sugars. *Human Biochemistry*, 131–160 (2018).

40.   D. Ghosh, J. S. Stumhofer, The spleen: "epicenter" in malaria infection and
immunity. *J Leukoc Biol*, 1–17 (2021).

41.   I. C. Hirako, P. A. Assis, B. Galvão-Filho, A. D. Luster, L. R. Antonelli, R. T.
Gazzinelli, Monocyte-derived dendritic cells in malaria. *Curr Opin Microbiol* 52,
139–150 (2019).

42.   I. C. Hirako, M. A. Ataide, L. Faustino, P. A. Assis, E. W. Sorensen, H. Ueta,
N. M. Araújo, G. B. Menezes, A. D. Luster, R. T. Gazzinelli, Splenic differentiation
and emergence of CCR5+CXCL9+CXCL10+ monocyte-derived dendritic cells in
the brain during cerebral malaria. *Nat Commun* 7, 13277 (2016).

43.   A. F. McGettrick, L. A. J. O'Neill, The Role of HIF in Immunity and
Inflammation. *Cell Metab* 32, 524–536 (2020).

44.   T. Ramalho, P. A. Assis, O. Ojelabi, L. Tan, B. Carvalho, L. Gardinassi, O.
Campos, P. L. Lorenzi, K. A. Fitzgerald, C. Haynes, D. T. Golenbock, R. T.
Gazzinelli, Itaconate impairs immune control of Plasmodium by enhancing mtDNA-
mediated PD-L1 expression in monocyte-derived dendritic cells. *Cell Metab* 36,
484-497.e6 (2024).

45.   N. Olson, A. Van Der Vliet, Interactions between nitric oxide and hypoxia-
inducible factor signaling pathways in inflammatory disease. *Nitric Oxide* 25, 125–
137 (2011).

46.   H. Blaser, C. Dostert, T. W. Mak, D. Brenner, TNF and ROS Crosstalk in
Inflammation. *Trends Cell Biol* 26, 249–261 (2016).

47.   M. M. Stevenson, M. F. Tam, S. F. Wolf, A. Sher, IL-12-induced protection
against blood-stage Plasmodium chabaudi AS requires IFN-gamma and TNF-
alpha and occurs via a nitric oxide-dependent mechanism. *The Journal of*
Immunology 155 (1995).

48.   T. E. Taylor, M. E. Molyneux, J. J. Wirima, K. A. Fletcher, K. Morris, Blood
Glucose Levels in Malawian Children before and during the Administration of
Intravenous Quinine for Severe falciparum Malaria. *New England Journal of*
Medicine 319, 1040–1047 (1988).

49.   N. J. White, D. A. Warrell, P. Chanthavanich, S. Looareesuwan, M. J.
Warrell, S. Krishna, D. H. Williamson, R. C. Turner, Severe Hypoglycemia and
Hyperinsulinemia in Falciparum Malaria. *New England Journal of Medicine* 309,
61–66 (1983).

50. H. V. Thien, P. A. Kager, H. P. Sauerwein, Hypoglycemia in falciparum
malaria: is fasting an unrecognized and insufficiently emphasized risk factor?
Trends Parasitol 22, 410–415 (2006).

51. Q. W. Xie, C. Nathan, "The high-output nitric oxide pathway: Role and
regulation" in *Journal of Leukocyte Biology* (Federation of American Societies for
Experimental Biology, 1994)vol. 56, pp. 576–582.

52. J. Zhou, T. Schmid, B. Brü, Tumor Necrosis Factor-Causes Accumulation of
a Ubiquitinated Form of Hypoxia Inducible Factor-1 through a Nuclear Factor-B-
Dependent Pathway. *Mol Biol Cell* 14, 2216–2225 (2003).

53. J. L. M. Madrigal, O. Hurtado, M. A. Moro, I. Lizasoain, P. Lorenzo, A.
Castrillo, L. Boscá, J. C. Leza, "The Increase in TNF-Levels Is Implicated in NF-B
Activation and Inducible Nitric Oxide Synthase Expression in Brain Cortex after
Immobilization Stress" (2002).

54. C. Bogdan, Nitric oxide synthase in innate and adaptive immunity: An
update. Elsevier Ltd [Preprint] (2015). https://doi.org/10.1016/j.it.2015.01.003.

55. M. Hayashi, M. Sakata, T. Takeda, T. Yamamoto, Y. Okamoto, K. Sawada,
A. Kimura, R. Minekawa, M. Tahara, K. Tasaka, Y. Murata, Induction of glucose
transporter 1 expression through hypoxia-inducible factor 1α under hypoxic
conditions in trophoblast-derived cells. *Journal of Endocrinology* 183, 145–154
(2004).

56. G. L. Semenza, P. H. Roth, H. M. Fang, G. L. Wang, Transcriptional
regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1.
Journal of Biological Chemistry 269, 23757–23763 (1994).

57. G. L. Semenza, Hypoxia-inducible factors in physiology and medicine. *Cell*
148, 399–408 (2012).

58. H. Sam, Z. Su, M. M. Stevenson, "Deficiency in Tumor Necrosis Factor
Alpha Activity Does Not Impair Early Protective Th1 Responses against Blood-
Stage Malaria" (1999).

59. D. A. Evans, D. 0 Jacobs, D. W. Wilmore, D. Evans, D. ~ Andrew, D. W. W.
Jacobs, "Tumor necrosis factor enhances glucose uptake by peripheral tissues"
(1989).

60. "Nitric Oxide and Infection."

61. S. Ramos, T. W. Ademolue, E. Jentho, Q. Wu, J. Guerra, R. Martins, G.
Pires, S. Weis, A. R. Carlos, I. Mahú, E. Seixas, D. Duarte, F. Rajas, S. Cardoso,
A. G. G. Sousa, J. Lilue, T. Paixão, G. Mithieux, F. Nogueira, M. P. Soares, A
hypometabolic defense strategy against malaria. *Cell Metab* 34, 1183-1200.e12
(2022).

62. K. E. Merz, D. C. Thurmond, "Role of Skeletal Muscle in Insulin Resistance
and Glucose Uptake" in *Comprehensive Physiology* (Wiley, 2020)vol. 176, pp.
785–809.

63. A. Chadt, H. Al-Hasani, Glucose transporters in adipose tissue, liver, and
skeletal muscle in metabolic health and disease. *Pflugers Arch* 472, 1273–1298
(2020).

64. P. Ebeling, H. A. Koistinen, V. A. Koivisto, Insulin-independent glucose
transport regulates insulin sensitivity. [Preprint] (1998).

65. P. Parroche, F. N. Lauw, N. Goutagny, E. Latz, B. G. Monks, A. Visintin, K.
A. Halmen, M. Lamphier, M. Olivier, D. C. Bartholomeu, R. T. Gazzinelli, D. T.
Golenbock, Malaria hemozoin is immunologically inert but radically enhances
innate responses by presenting malaria DNA to Toll-like receptor 9. *Proc Natl Acad
Sci U S A*, doi: 10.1073/pnas.0608745104 (2007).

66. L. R. Antonelli, C. Junqueira, J. M. Vinetz, D. T. Golenbock, M. U. Ferreira,
R. T. Gazzinelli, The immunology of Plasmodium vivax malaria. (2020).
https://doi.org/10.1111/imr.12816.

67. C. Dumitru, A. M. Kabat, K. J. Maloy, Metabolic adaptations of CD4+ T cells
in inflammatory disease. *Front Immunol* 9, 1–17 (2018).

68. E. L. Mills, B. Kelly, L. A. J. O'Neill, Mitochondria are the powerhouses of
immunity. *Nat Immunol* 18, 488–498 (2017).

69. P. Jacobs, D. Radzioch, M. M. Stevenson, Nitric oxide expression in the
spleen, but not in the liver, correlates with resistance to blood-stage malaria in
mice. *The Journal of Immunology* 155 (1995).

70. M. M. Stevenson, M. F. Tam, S. F. Wolf, A. Sher, IL-12-induced protection
against blood-stage Plasmodium chabaudi AS requires IFN-gamma and TNF-

alpha and occurs via a nitric oxide-dependent mechanism. *J Immunol* 155, 2545–
2556 (1995).

71.   F. Li, P. Sonveaux, Z. N. Rabbani, S. Liu, B. Yan, Q. Huang, Z. Vujaskovic,
M. W. W. Dewhirst, C. Y. Li, Regulation of HIF-1alpha stability through S-
nitrosylation. *Mol Cell* 26, 63–74 (2007).

72.   E. Metzen, J. Zhou, W. Jelkmann, J. Fandrey, B. Brüne, Nitric oxide impairs
normoxic degradation of HIF-1alpha by inhibition of prolyl hydroxylases. *Mol Biol*
Cell 14, 3470–3481 (2003).

73.   M. Knight, S. Stanley, HIF-1α as a central mediator of cellular resistance to
intracellular pathogens. *Curr Opin Immunol* 60, 111–116 (2019).

74.   K. J. Tracey, A. Cerami, Tumor necrosis factor: A pleiotropic cytokine and
therapeutic target. [Preprint] (1994). https://doi.org/10.1146/annurev.med.45.1.491.

75.   K. Koedderitzsch, E. Zezina, L. Li, M. Herrmann, N. Biesemann, TNF
induces glycolytic shift in fibroblast like synoviocytes via GLUT1 and HIF1A.
Scientific Reports 2021 11:1 11, 1–11 (2021).

76.   P. Van Uden, N. S. Kenneth, S. Rocha, Regulation of hypoxia-inducible
factor-1alpha by NF-kappaB. *Biochem J* 412, 477–484 (2008).

77.   J. Rius, M. Guma, C. Schachtrup, K. Akassoglou, A. S. Zinkernagel, V.
Nizet, R. S. Johnson, G. G. Haddad, M. Karin, NF-kappaB links innate immunity to
the hypoxic response through transcriptional regulation of HIF-1alpha. *Nature* 453,
807–811 (2008).

78.   M. Utsuyama, K. Hirokawa, Differential expression of various cytokine
receptors in the brain after stimulation with LPS in young and old mice. *Exp*
Gerontol 37, 411–420 (2002).

79.   L. R. Leon, Invited review: cytokine regulation of fever: studies using gene
knockout mice. *J Appl Physiol (1985)* 92, 2648–2655 (2002).

80.   C. Li, L. A. Sanni, F. Omer, E. Riley, J. Langhorne, Pathology of
Plasmodium chabaudi chabaudi infection and mortality in interleukin-10-deficient
mice are ameliorated by anti-tumor necrosis factor alpha and exacerbated by anti-
transforming growth factor beta antibodies. *Infect Immun* 71, 4850–4856 (2003).

81.   P. B. Falanga, M. Regina, D. I. Limaa, A. Coutinho, primary infection by
Plasmodium chabaudi and in. 599–603 (1987).
82.   M. A. Ataide, W. A. Andrade, D. S. Zamboni, D. Wang, M. do C. Souza, B.
S. Franklin, S. Elian, F. S. Martins, D. Pereira, G. Reed, K. A. Fitzgerald, D. T.
Golenbock, R. T. Gazzinelli, Malaria-Induced NLRP12/NLRP3-Dependent
Caspase-1 Activation Mediates Inflammation and Hypersensitivity to Bacterial
Superinfection. *PLoS Pathog* 10 (2014).

Acknowledgments

The authors thank members of the D.L.C., J.S.S., J.C.A.F. and R.T.G. groups for
scientific discussions and technical help. Denise Brufato Ferraz and Francielle
Pioto for the excellent technical assistance. Cristiane Gomes and Patricia Palhares
for excellent financial technical assistance.
Funding: This work was supported by the Fundação de Amparo de Pesquisa do
Estado de São Paulo (2016/23618-8) (2020/01043-9), the National Institute of
Infectious Diseases and Allergy (1R21 AI150546-01, R01NS098747 and
R01AI079293) and the Instituto Nacional de Ciência e Tecnologia de Vacinas
(CNPq/ FAPEMIG/ CAPES 465293/2014-0).

Author contributions: K.C.M., J.C.A.F. and R.T.G. designed, K.C.M., N.P.S.L. P.A.A., I.C.H., L.G.V., J.E.T.K., D.L.C. performed experiments. I.C.H, P.A.A., O.O., N.P.S.L., K.C.M. and R.T.G. analyzed data, D.L.C., J.S.S., J.C.A.F. and R.T.G. contributed with reagents, materials, and/or analysis tools, and K.C.M., N.P.S.L. and R.T.G. wrote the manuscript. All authors revised and approved the final version of the manuscript.

Competing interests: The authors declare no competing interests.

Data and materials availability:

GEO (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE109908) accession number is GEO: GSE109908.

Figures

Fig. 1. *P. chabaudi*-**infected mice display enhanced TNF production associated with increased disease parameters.** C57BL/6 mice were infected or not i.p. with *P. chabaudi* (10^5 iRBCs). **(A)** Parasitemia was determined at 0, 3, 5, 7, 8, 10 and 12 days-post infection (dpi). **(B)** Blood glucose levels were measured with a glucometer at 0, 3, 5, 7, 8, 10 and 12 dpi. Means of blood glucose levels were compared between infected (red) and uninfected (blue) mice at the same time points. **(C)** Blood TNF Levels were measured by ELISA at 8 dpi. Means of blood TNF levels were compared between infected (red) and uninfected (blue) mice at the same time points. **(D)** TNF m-RNA from mice liver was determined at 8 days post- infection (dpi). **(E)** Parasitemia was determined at 8 days-post-infection (dpi) in C57BL/6 and TNFR deficient mice. **(F)** Rectal temperature was determined at 8 dpi in C57BL/6 and TNFR deficient mice. **(G-J)** MCP-1, TNF, IFN and IL-10 was measured by CBA Kit at 8 dpi. **(K)** Blood glucose levels were measured with a glucometer at 8dpi. A, B and F: Graphs show mean ± SEM of 1 representative experiment of at least 3 independent ones performed with 4-5 mice per group; Statistical analysis: One-way ANOVA with Tukey post-hoc test. C and D: Graph shows mean± SEM of combined data from 3 independent experiments with 4 - 6 mice per group; Statistical analysis: One-way ANOVA with Bonferroni post-hoc test. E, G, H, I, J, K: Graphs show mean ± SEM of combined data from 3 independent experiments with 4-6 mice per group; Statistical analysis: One-way ANOVA with Tukey post-hoc test. Asterisks above the bars indicate comparisons between mice of the same strain before infection and at 8 dpi. Asterisks connected by horizontal lines indicate comparisons between infected groups. *P≤0.05, **P≤0.01, ***P≤0.001, **** P≤0.0001, ns = non-significant.

Fig. 2. The detrimental effects of *P. chabaudi* **infection on physical activity, food intake and energy metabolism are partially dependent on TNF receptor signaling.** Mouse energy balance from C57BL/6 and TNFR$^{-/-}$ was assessed using metabolic cages (TSE 959 Systems, Chesterfield, MO) at 3 days prior to infection

(uninfected) (C57BL/6: blue, TNFR-/-: gray) and at 8 days postinfection (dpi) with

P. chabaudi (10^5 infected RBCs) (C57BL/6: red, TNFR$^{-/-}$: orange). Scores for **(A**

and E) physical activity, **(B and F)** food intake, **(C and G)** energy expenditure, **(D**

and H) respiratory exchange, and **(I and J)** percentual of fat mass and lean mass

are depicted. Hourly measurements of such parameters are shown in top row

panels for mice 3 days prior to infection (uninfected) and in middle row panels for

infected mice. The average values for light and dark cycles in all groups are

depicted in bottom row panels. **(K and L)** Glucose uptake was measured 2-[14C]

deoxyglucose. Experiments were performed at the National Mouse Metabolic

Phenotyping Center (MMPC) at UMASS Medical School. Basal glucose uptake in

individual organs of infected (red) and uninfected (blue) mice was measured using

an intravenous injection of 2-[14C] deoxyglucose. After 1h, mice were

anesthetized, and tissue samples were taken for organ-specific levels of 2-[14C]

deoxyglucose-6-phosphate. A - L: Graph shows mean ± SEM of combined data

from 2 independent experiments; Statistical analysis: One-way ANOVA with Tukey

post-hoc test. Asterisks above the bars indicate comparisons between mice of the

same strain before infection and at 8 dpi. Asterisks connected by horizontal lines

indicate comparisons between infected groups. *$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$, ****

$P \leq 0.0001$, ns = non-significant.

Fig. 3. *P. chabaudi* infection stimulates Increased GLUT1 expression in

hepatic non-parenchymal and splenic CD11b+ cells associated with an

altered host metabolic profile. C57BL/6 mice were infected or not i.p. with P.

chabaudi (10^5 iRBCs). **(A)** RNA-seq was performed in livers from infected and

uninfected mice. **(B)** GLUT1 expression in livers of infected (8 dpi) or uninfected

mice was evaluated on by western blot. **(C and D)** GLUT1 expression in **(C)**

hepatocytes or **(D)** hepatic non-parenchymal cells isolated from livers from

C57BL/6 infected (8dpi) or uninfected mice, quantified by western blot (expression

of β-actin was used as control). **(E)** Hepatocytes or **(F)** Hepatic non-parenchymal

cells were purified from livers harvested from infected (red - 8 dpi) and uninfected

(blue) mice and cultured ex vivo for quantification of extracellular acidification rate

(ECAR) by Seahorse XFe96. **(G, I and J)** T cells, B cells and myeloid cells
(CD11b+) were purified from splenocytes of infected and uninfected mice using
magnetic beads and **(G)** GLUT1 expression was evaluated by western blot. **(I)** T
and B cells or **(J)** CD11b+ purified from spleens were cultured *ex vivo* for
quantification ECAR. **(H, K, L and M)** graphs showing GLUT1 expression
evaluated by flow cytometry in splenic cells from infected (red – 8 dpi) and
uninfected (blue) mice. A. Heatmap represents 3 biological replicates performed in
RNA-seq. B, C, D and G: Western blot images representative of at least 3
independent experiments. E, F, I and J: Graphs show mean ± SEM of 1
representative experiment of at least 3 independent ones performed with 4-5 mice
per group; Statistical analysis: Two-way ANOVA followed by Sidak's post-hoc test
employing mixed effect analysis provided by Agilent Seahorse XFe96 analyzer
software. H, K L and M: Representative dot plots and Graphs show mean ± SEM of
1 representative experiment of at least 3 independent ones performed with 4-5
mice per group; Statistical analysis: Student's t test. Asterisks above the bars
indicate comparisons between mice of the same strain before infection and at 8
dpi. Asterisks connected by horizontal lines indicate comparisons between infected
groups. *$P \leq 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$, **** $P \leq 0.0001$, ns = non-significant.
1028
Fig. 4. *P. chabaudi* infection triggers increased glycolysis in monocytic cells
in a TNF-receptor dependent way. C57BL/6 or TNFR-/- mice were infected or not
i.p. with *P. chabaudi* (10^5 iRBCs). All experimental groups survived *Pc* infection
throughout the observation period. (**A**) Glycolytic and gluconeogenesis pathways.
Relative expression of glycolytic **(B-E)** or gluconeogenesis **(F-J)** enzymes in livers
of control or infected (8 dpi) mice. **(K)** GLUT1, GLUT2 and NAK-ATPase
expression in livers of infected (8 dpi) or uninfected mice evaluated by western
blot. **(L)** Representative histogram and graph showing GLUT1 expression
evaluated by flow cytometry in CD11b+/F4/80+/CD11c+/Ly6G- cells from spleens
of infected (8 dpi) or uninfected C57BL/6 or TNFR$^{-/-}$ mice. **(M)** Representative
histogram and graph showing GLUT1 expression evaluated by flow cytometry in
CD11b+/F4/80+/CD11c+/MHCII+ cells from spleens of uninfected C57BL/6 or

TNFR$^{-/-}$ mice stimulated or not with TNF (100 ng/ml) for 18h. **(N and O)** CD11b+

cells were purified from spleens harvested from infected (8 dpi) C57BL/6 (red) and

TNFR-/- (orange) mice or uninfected C57BL/6 (blue) and TNFR-/- (gray) mice and

cultured *ex vivo* for evaluation of ECAR by Seahorse XFe96 Analyzers. **(P)**

Parasitemia was determined at 8 days-post-infection (dpi) from *TNF-R1aΔLyz2*

and Wild-type mice. **(Q)** Rectal temperature was determined at 8 dpi in WT and

TNF- R1aΔLyz2 at 8 dpi. **(R)** Blood glucose levels were measured with a

glucometer at 8dpi. B, C, D, E, F, G, H, I, J and M: Graphs show mean ± SEM of

combined data from 2 independent experiments with 3-5 mice per group; Statistical

analysis: One-way ANOVA followed by Tukey post-test. K: representative dot plot

of 1 representative experiment representative of 3 independent ones; L, P, Q, R:

Graphs show mean ± SEM of 1 representative experiment of at least 3

independent ones performed with 4-5 mice per group, Statistical analysis:

Student's t test. N: Graphs show mean ± SEM of 1 representative experiment of at

least 3 independent ones performed with 4-5 mice per group; O: Violin plots of

combined data from 3 independent experiments with 3-5 mice per group –

individual dots represent the average of triplicates in each experiment – the

difference refers to the number of cells isolated from naïve vs infected animals,

which varied. N and O: Statistical analysis: Two-way ANOVA followed by Sidak's

post-test employing mixed effect analysis provided by Agilent Seahorse XFe96

analyzer software. Asterisks above the bars indicate comparisons between mice of

the same strain before infection and at 8 dpi. Asterisks connected by horizontal

lines indicate comparisons between infected groups. *P≤0.05, **P≤0.01,

***P≤0.001, **** P≤0.0001, ns = non-significant.

1065

Fig. 5. HIF-1α contributes to glycolysis induction during *Pc* infection. HIF-1α

expression in the nuclear extract of cells from spleen **(A)** and liver **(B)** of infected

(8 dpi) or uninfected C57BL/6 and TNFR-/- mice evaluated by western blot.

(expression of nucleofosmin -NFM- was used as control). **(C)** Representative

histogram and and graph showing GLUT1 expression evaluated by flow cytometry

in CD11b+/F4/80+/CD11c+/MHCII+ cells from spleens of infected (8 dpi) or

uninfected HIF-1aΔLyz2 and Wild-type mice. **(D)** Representative histogram and
and graph showing GLUT1 expression evaluated by flow cytometry in
CD11b+/F4/80+/CD11c+/MHCII+ cells from spleens of uninfected HIF-1aΔLyz2
and wild-type mice stimulated or not with TNF (100 ng/ml) for 18h. **(E and F)**
CD11b+ cells were purified from spleens harvested from infected (8 dpi) WT (red)
and HIF-1aΔLyz2 (pink) mice or uninfected WT (blue) and HIF-1aΔLyz2 (purple)
mice and cultured ex vivo for evaluation of ECAR by Seahorse XFe96 Analyzers.
(G) Parasitemia was determined at 8 days-post- infection (dpi) from HIF-1aΔLyz2
and Wildtype mice. **(H)** Blood TNF Levels were measured by ELISA at 8 dpi. **(I)**
TNF levels in supernatants from splenic cells harvested from infected (8 dpi) or
uninfected HIF-1aΔLyz2 and Wild-type mice mice and stimulated ex vivo with LPS
[1ug/mL] or not during 24H. A and B: representative dot plot of 1 representative
experiment representative of 3 independent ones. C: Graphs show mean ± SEM of
1 representative experiment of 3 independent ones with 3-5 mice per group;
Statistical analysis: Student's t test. D: Graphs show mean ± SEM of 1
representative experiment of 2 independent ones with 3-5 mice per group;
Statistical analysis: Student's t test. E: Graphs show mean ± SEM of 1
representative experiment of at least 3 independent ones performed with 4-5 mice
per group. F: Violin plots of combined data from 3 independent experiments with 3-
5 mice per group – individual dots represent the average of triplicates in each
experiment – the difference refers to the number of cells isolated from naïve vs
infected animals, which varied. E and F: Statistical analysis: Two-way ANOVA
followed by Sidak's post-hoc test employing mixed effect analysis provided by
Agilent Seahorse XFe96 analyzer software. G and H: Graphs show mean ± SEM
of combined data from 3 independent experiments with 3-5 mice per group;
Statistical analysis: Student's t test. I: Graphs show mean ± SEM of 1
representative experiment of 3 independent ones with 3-5 mice per group;
Statistical analysis: One-way ANOVA followed by Tukey post-hoc test. Asterisks
above the bars indicate comparisons between mice of the same strain before
infection and at 8 dpi. Asterisks connected by horizontal lines indicate comparisons

between infected groups. *P≤0.05, **P≤0.01, ***P≤0.001, **** P≤0.0001, ns = non-

significant.

1104

Fig. 6. Malaria disease parameters in iNOS deficient mice mirror those

observed in TNF receptor deficient animals. (A) Nitrite levels in supernatants

from CD11b+ splenic cells harvested from infected (8 dpi) C57BL/6 or TNF-R1-/-

mice and stimulated ex vivo with LPS [1ug/mL] or not during 48H. **(B)** Parasitemia

was determined at 8 days-post- infection (dpi) from C57BL/6 and iNOS knockout

mice. **(C)** Rectal temperature was determined at 8 dpi. **(D)** Blood glucose levels

were measured with a glucometer at 8dpi. **(E and F)** CD11b+ cells were purified

from spleens harvested from infected (8 dpi) C57BL/6 (red) and iNOS knockout

(green) mice or uninfected C57BL/6 (blue) and iNOS knockout (beige) mice and

cultured ex vivo for evaluation of ECAR by Seahorse XFe96 Analyzers. **(G-J)**

Relative expression of glycolytic enzymes in livers of naïve or infected (8 dpi) mice.

(K) Representative histogram and graph showing GLUT1 expression evaluated by

flow cytometry in CD11b+/F4/80+/CD11c+/MHCII+ cells from spleens of infected (8

dpi) or uninfected C57BL/6 or iNOS$^{-/-}$ mice. **(L)** Representative histogram and

graph showing GLUT1 expression evaluated by flow cytometry in

CD11b+/F4/80+/CD11c+/MHCII+ cells from spleens of uninfected C57BL/6 or

iNOS$^{-/-}$ mice stimulated or not with TNF (100 ng/ml) for 18h. **(M)** CD11b+ splenic

cells were purified from infected C57BL/6 (red - 8 dpi), iNOS$^{-/-}$ (green - 8 dpi), and

uninfected mice and cultured ex vivo for quantification of proton efflux rate (PER)

using glycolytic rate assay by Seahorse XFe96. **(N)** Parasitemia was determined

during 20 days-post-infection (dpi) from wild-type infected mice treated with 2-

[14C]-deoxyglucose (250mg/kg daily from 4 to 15 dpi) or PBS. **(O)** Lactate was

measured in the plasma of mice. A: Graphs show mean ± SEM of 1 representative

experiment of 3 independent ones with 4-5 mice per group; Statistical analysis:

One-way ANOVA followed by Tukey post-hoc test. B, C and D: Graphs show mean

± SEM of 1 representative experiment of 3 independent ones with 3-5 mice per

group (B and C) or of combined data from 2 independent experiments with 3-5

mice per group (D); Statistical Analysis: Student's t test. E: Graphs show mean ±

SEM of 1 representative experiment of at least 3 independent ones performed with 4-5 mice per group. F: Violin plots of combined data from 3 independent experiments with 3-5 mice per group – individual dots represent the average of triplicates in each experiment – the difference refers to the number of cells isolated from naïve vs infected animals, which varied. E and F: Statistical analysis: Two-way ANOVA followed by Sidak's post-hoc test employing mixed effect analysis provided by Agilent Seahorse XFe96 analyzer software. G, H, I, J, N, O: Graphs show mean ± SEM of combined data from 2 independent experiments with 3-5 mice per group; Statistical Analysis: Student's t test: One-way ANOVA followed by Tukey post-hoc test. K: Graphs show mean ± SEM of 1 representative experiment of 3 independent ones with 3-5 mice per group; Statistical analysis: Student's t test. L: Graphs show mean ± SEM of 1 representative experiment of 2 independent ones with 3-5 mice per group; Statistical analysis: Student's t test. M: Graphs show mean ± SEM of 1 representative experiment of at least 3 independent ones performed with 4-5 mice per group; Statistical analysis: Two-way ANOVA followed by Sidak's post-hoc test employing mixed effect analysis provided by Agilent Seahorse XFe96 analyzer software. Asterisks above the bars indicate comparisons between mice of the same strain before infection and at 8 dpi. Asterisks connected by horizontal lines indicate comparisons between infected groups. *P≤0.05, **P≤0.01, ***P≤0.001, **** P≤0.0001, ns = non-significant.