1 **Microbiome turnover during offspring development varies with**

2 **maternal care, but not moult, in a hemimetabolous insect**

Marie-Charlotte Cheutin[1], Manon Boucicot[1], Joël Meunier[1]

[1] Institut de Recherche sur la Biologie de l'Insecte, UMR 7261, CNRS, University of

Tours, Tours, France

Corresponding author: MC Cheutin, mccheutin@gmail.com

J Meunier ORCID #0000-0001-6893-2064

MC Cheutin ORCID #0000-0001-7711-7512

Abstract

The ecological success of insects often depends on their association with beneficial microbes. However, insect development involves repeated moults, which can have dramatic effects on their microbial communities. Here, we investigated whether and how moulting affects the microbiome of a hemimetabolous insect, and whether maternal care can modulate these effects. We reared European earwig juveniles with or without mothers and used 16S rRNA metabarcoding to analyse the prokaryotic fraction of the core microbiome of eggs, recently and old moulted individuals at four developmental stages and the resulting adults. The 218 samples obtained showed that the microbiome diversity changed non-linearly during development and that these changes were associated with bacterial biomarkers. Surprisingly, these changes did not occur during moulting, but rather between the beginning and end of certain developmental stages. We also found that access to maternal care affected the microbiome of both juveniles and adults, even when the last contact with mothers was two months before adulthood. Overall, these results provide new insights into our understanding of the (in)stability of the prokaryotic microbiome in hemimetabolous insects and its independence from moult. More generally, they question the role of microbiome acquisition through maternal care in maintaining family life in species where this behaviour is facultative.

Keywords: Microbiome, Horizontal transmission, Vertical transmission, Metamorphosis, Sociality

Introduction

Insects are the most diverse and abundant animal taxon on Earth, comprising more than half of the animal kingdom (Mora *et al.*, 2011; Berenbaum, 2017; Samways, 2018). One reason for their evolutionary success is their frequent association with a large and complex diversity of beneficial microorganisms (Shapira, 2016; Sudakaran *et al.*, 2017). This is because such associations can help their host colonize new ecological niches and mediate insect speciation and adaptation to a variety of environments (Berlanga and Guerrero, 2016; Shapira, 2016). However, maintaining these associations is challenging for insects, as it requires hosts to retain their microbes despite the numerous and successive moults (or ecdysis) that occur between the egg and adult stages (Bright and Bulgheresi, 2010). From a microbial perspective, these moults are risky as they can lead to a partial purge from the host, inducing a microbial bottleneck in which almost all symbionts are lost (McFrederick *et al.*, 2014; Zhukova *et al.*, 2017). Moults can also lead to critical changes in the habitat used by microbes within the host, which in turn can rapidly induce major turnover in their communities (Engel and Moran, 2013). For the hosts, moulting also poses a significant challenge, as they need to retain or (re)acquire the beneficial symbionts (Arce *et al.*, 2012; Wang and Rozen, 2017) while coping with surrounding pathogenic microorganisms susceptible to take their place in the host (Salem *et al.*, 2015; Hammer and Moran, 2019). Therefore, it is necessary for both host insects and beneficial microorganisms to develop strategies to ensure the continuity of their association throughout host development.

To ensure the (re)inoculation and maintenance of beneficial microorganisms throughout development, insects can adopt two non-mutually exclusive strategies. On one hand, they can acquire these microorganisms from direct contact with their

environment and conspecifics, a process called horizontal transmission. This process can be particularly important when individuals live in groups, share a nesting environment and/or when moulting leads to the loss of these microorganisms at each development stage (Raina *et al.*, 2008; Nalepa, 2020; Rose *et al.*, 2023). On the other hand, hosts can acquire beneficial microorganisms from their parents, a process called vertical transmission (Bright and Bulgheresi, 2010; Sachs *et al.*, 2011; Hosokawa and Fukatsu, 2020; Michaud *et al.*, 2020). In insects, vertical transmission was long thought to occur mainly through the transfer of microorganisms directly into the eggs, which the resulting offspring then had to maintain throughout their development. However, many insect parents provide care to their eggs and juveniles after oviposition (Meunier *et al.*, 2022) and recent studies show that this care can mediate a vertical transmission of microorganisms. For instance, mothers can deposit microorganisms directly on the eggshell of their future juveniles or produce symbiont capsules that they place next to the eggs for future ingestion by newly hatched juveniles, as reported in stinkbugs of the families Pentomatidae and Scutelleridae (Fukatsu and Hosokawa, 2002; Kikuchi *et al.*, 2008; Hosokawa *et al.*, 2013). After hatching, other mechanisms of vertical and horizontal transmissions can also occur by social interactions between family members, such as trophallaxis through mouth-to-mouth or mouth-to-anus contacts (Powell *et al.*, 2014; Zhukova *et al.*, 2017) or allo-coprophagy through consumption of parental feces (vertical) and/or sibling feces (horizontal) (Lombardo, 2008; Onchuru *et al.*, 2018). Access to maternal care and family life can thus ensure the acquisition and reacquisition of beneficial microbes by moulting juveniles, thus possibly strengthening the stability and evolutionary trajectory of symbiotic associations.

While our current understanding of the consequences of moulting and

maternal care on the dynamics of the host microbiome during development is mainly

based on holometabolous insects, little is known about these consequences on

hemimetabolous species (Hammer and Moran, 2019; Girard *et al.*, 2023). The focus

on holometabolous species is explained by the fact that their immature stages have a

morphology and sometimes an ecology very different from those of adults, which

raises obvious questions about the fate of their microbiome during metamorphosis

(Johnston *et al.*, 2019). In contrast, hemimetabolous insects have immature stages

called nymphs that are very similar in morphology and ecology to the adult, and their

juveniles undergo only gradual morphological changes through successive moults

(Johnston *et al.*, 2019). The impact of these gradual changes on the dynamics of the

host microbiome has received comparatively much less attention, and a few studies

suggest that it may be stable throughout development such as in Blattodea,

Orthoptera, and Hemiptera (Sudakaran *et al.*, 2012; Manthey *et al.*, 2022). However,

it is not clear whether this stability is universal across species. Importantly, more

information is needed to determine whether this stability is due either to the non-

purging effect of moulting on the microbial community, to the host microbial niche not

changing during development and therefore selecting for the same microbial

community, and/or to maternal care ensuring maintenance of the microbial

community through vertical transmission.

In this study, we investigated whether and how the microbiome of the

hemimetabolous European earwig *Forficula auricularia* L*.* (Order Dermaptera:

Forficulidae) changes during juvenile development and tested whether these

potential changes were due to moulting events, stage-specific microbial niches

and/or offspring access to maternal care. In this species, females oviposit in

individual burrows in early winter (Meunier et al., 2012; J. Tourneur & Meunier, 2020)

after which they stop their foraging activity and provide extensive forms of care to

their eggs. For instance, mothers fiercely protect their eggs from predators (Trumbo,

2012; Wong & Kölliker, 2012), move their clutches when faced with extreme

temperature changes (Tourneur *et al.*, 2022), and frequently groom their eggs to

remove fungal spores and deposit cuticular hydrocarbons to protect them from

desiccation (Boos et al., 2014; Diehl & Meunier, 2018). About 50 days later, the eggs

hatch and the mothers stay with their new juveniles for about two more weeks.

During this time, they continue to provide care to their nymphs, such as allo-grooming

and food provisioning (Kölliker, 2007; Lamb, 1976). Interestingly, maternal presence

is not required after hatching, as nymphs can develop and survive without contact

with a mother (Kölliker, 2007; Kramer *et al.*, 2015; Thesing *et al.*, 2015). The family

naturally splits shortly after the nymphs have moulted for the second time (the first

moult occurs at the time of hatching), and the nymphs then moult three more times

before reaching adulthood two months later (Thesing et al., 2015; Tourneur et al.,

2020). Whether the offspring microbiome changes during development and whether

maternal care influences these changes are unknown in the European earwig.

However, the microbiome of the eggshell is known to change over 16 days and to be

partly influenced by maternal presence in the maritime earwig *Anisolabis maritima*

(Greer *et al.*, 2020).

We conducted two experiments in which we analysed the prokaryotic

microbiome of 20 families of the European earwig *F. auricularia*. The first experiment

tested whether and how their microbial community changes during development. We

reared fifteen earwig families to adulthood using a standard protocol in which

mothers remained with the clutch until 14 days after hatching, *i.e.* for the normal

duration of family life. During this time, we sampled offspring from each

developmental stage (from eggs to adult male and female offspring) at both the

beginning and end of each developmental stage to determine the impact of moulting

on microbial communities. We then used 16S rRNA metabarcoding to analyse the

prokaryotic fraction of their core microbiome, *i.e.,* the sequences non-randomly

distributed across all datasets (see details below). The second experiment tested

whether and how the presence of the mother affected the microbiome of her first

instar nymphs and resulting adult offspring. To do this, we reared five additional

earwig families following the same protocol, but where the mother was removed from

the clutch shortly after the eggs hatched. Overall, we found that the microbiome of

earwig offspring surprisingly changed during development. We also showed that

these changes are not due to a purging event during moulting, but rather likely reflect

stage-specific microbial niches in the nymphs. Finally, we found that access to

maternal care has both short- and long-term effects on the microbiome of offspring.

Material and methods

Earwig sampling and laboratory rearing

The eggs, nymphs and adults analysed in this study were the first-generation

progeny of 20 females of *F. auricularia* sp "A" (Wirth *et al.*, 1998; González-Miguéns

et al., 2020). These 20 females were part of a large field sampling of earwig males

and females conducted in an orchard near Valence, France (44°58′38″N, 4°55′43″E)

in the summer of 2021. Just after field sampling, these individuals were randomly

distributed into plastic containers with 100 females and 100 males and then

maintained under standard laboratory conditions to allow for uncontrolled mating

(Koch and Meunier, 2014; Sandrin *et al.*, 2015). In November 2021, each female was

isolated to mimic her natural dispersal from the groups and to stimulate oviposition

(Kölliker, 2007). These females were transferred to individual Petri dishes (55 mm x

12 mm) lined with moist sand (Körner *et al.*, 2018) and maintained in constant

darkness at 10°C until oviposition and egg hatching. From isolation until oviposition,

females were fed ad libitum with a laboratory-prepared food consisting mainly of

carrots, cat food, seeds and pollen (Kramer *et al.*, 2015). Food was renewed each

week, but removed from the day of oviposition until the day of egg hatching, as this is

when the mothers typically stop their foraging activity (Kölliker, 2007; Van Meyel and

Meunier, 2020). From this large pool of females, we haphazardly selected 20

clutches in which the mothers produced 50 eggs (clutch size typically varies from 30

to 60 eggs, Tourneur and Gingras, 1992) for our measurements. The remaining

females were used in other experiments not presented here.

On the day of egg hatching, the 20 selected clutches were transferred to larger

Petri dishes (145 mm x 13 mm) lined with moist sand (Körner *et al.*, 2018) to

manipulate maternal presence during offspring development. Of these 20 clutches,

five (randomly selected) had their mothers removed to subsequently prevent post-

hatching maternal care and any mother-offspring interactions. For the remaining 15

clutches, the nymphs were kept with their mother for 14 days after egg hatching

(which is the natural length of family life in this species) and then separated from their

mother for the rest of their development. All clutches were maintained under

laboratory conditions at 18-20°C under a 12:12 light:dark photoperiod, and received

the laboratory-prepared food twice a week (see above).

Experimental design and sample collection

Overall, we analysed the microbiome of 218 samples collected throughout earwig

development (Figure 1, Table S1). As ecdysis is known to induce a microbial shift in

many insect species, we sampled all developmental stages (except eggs and adults)

both immediately after the moult (freshly moulted) and several days after the moult,

before the next moult (old moulted). Freshly moulted nymphs have a white

(compared to dark) colour, which typically lasts for maximum 3h in early instar

nymphs and up to 6h in late instar (MC Cheutin, pers. obs.). For the 15 clutches with

maternal care, we obtained a total of 12 egg samples (Egg), 37 first instar nymphs of

which 12 were freshly moulted (later called L1-Y) and 15 were old moulted (L1-O), 30

second instar nymphs of which 15 were freshly moulted (L2-Y) and 15 were old

moulted (L2-O), 30 third instar nymphs of which 15 were freshly moulted (L3-Y) and

15 were old moulted (L3-O), 30 fourth instar nymphs of which 15 were freshly

moulted (L4-Y) and 15 were old moulted (L4-O), and finally 32 adults of which 11

field-sampled mothers (Mother), 10 adult female offsprings (Adult-F) and 11 adult

male offsprings (Adult-M). For the 5 clutches without maternal care, we only used a

total of 21 old moulted first instar nymphs, 16 adult females and 20 adult males as we

were only interested in the short and long-term effect of maternal presence/absence

on offsprings (Figure 1, see details in Table S1). Note that the first observable

developmental stage of the nymphs is called L1, as the moult that occurs at hatching

is called L0 in this species (Tourneur *et al.*, 2020). Moreover, the sex of an individual

can only be determined in adults, where the shape of the forceps is straight in

females and curved in males. Each sample was collected individually and

immediately transferred to Eppendorf tubes at -80°C until DNA extraction.

Figure 1: Overview of the experimental design. In Experiment 1, we sampled each developmental stage from egg to adult for microbiome analysis. We sampled L1 to L4 nymphs both just after they had moulted to the new stage (grey - Y for young nymphs) and just before they moulted to the next stage (coloured - O for old nymphs). We also sampled female (Adult-F) and male (Adult-M) adults. This gave us 150 samples across all developmental stages in families with mothers during early nymph development. In Experiment 2, we sampled old first instar nymphs and adult males and females. This gave us 58 samples in families without mothers.

Genomic extraction and 16S amplification

Total genomic DNA was extracted using the NucleoMag® Tissue extraction kit (Macherey-Nagel™, Düren, Germany) and the V3-V4 region of the 16S rDNA gene was amplified with the prokaryotic primers 343F (5′- ACGGRAGGCAGCAG – 3′) and 784R (5′- TACCAGGGTATCTAATC – 3′) (Muyzer *et al.*, 1993) coupled with platform-specific Illumina linkers. We performed PCR reactions using the Taq Polymerase Phusion® High-Fidelity PCR Master Mix with GC buffer and prepared them according to the manufacturer's instructions (Qiagen, Hilder, Germany). PCR amplification steps involved an initial denaturing step for 30 sec at 98°C, followed by 22 amplification cycles (denaturation for 10 sec at 98°C; annealing for 30 sec at 61.5°C;

extension for 30 sec at 72°C), and ended by a final extension step of 10 min at 72°C.

Electrophoresis migration was run to check the probe specificity and amplification

success. Extraction and amplification steps involved several blank controls to confirm

that samples were not contaminated by environmental microorganisms. Samples

were amplified in duplicates and equally pooled for a final product of 30 µL further

sequenced with 2x250bp Illumina MiSeq technology at the Bio-Environnement

platform (University of Perpignan, France). As the blanks were all negative, we did

not send them for sequencing.

Bioinformatic process

The obtained libraries were trimmed and filtered using the quality profiles from the

DADA2 algorithm v1.24.0 (Callahan *et al.*, 2016), cleaned for errors, dereplicated and

inferred towards Amplicon Sequence Variants (ASVs) (Glassman and Martiny, 2018).

Chimeras were removed and taxonomy assignment was performed on a count table

where forward and reverse ASVs were merged and pooled, using the SILVA

reference database (release 138) (Quast *et al.*, 2012). The multiple alignment of the

sequences was provided with the MAFFT program (Katoh, 2002) and we inferred the

phylogenetic tree with the FastTree 2 tool (Price *et al.*, 2010) and Phangorn package

v2.8.1 (Schliep, 2011). The table was then transformed into a phyloseq object using

the phyloseq package v1.40.0 (McMurdie and Holmes, 2013) of which we removed

the 529 sequences of mitochondrial origin and the 30 800 unknown sequences. As

the 2 700 Eukarya sequences had no affiliation at Phylum level and were sparsely

distributed, they were also removed. Finally, we obtained a final dataset constituted

by 4 163 989 sequences (3 780 ASVs) ranging from 3 570 to 32 946

sequences/sample.

Identification of the core microbiomes

We defined the core microbiome as the set of microbial taxa (*i.e.,* ASVs) that are characteristic of all samples. We obtained this core microbiome using the species abundance distribution (SAD) patterns of each ASVs (Magurran and Henderson, 2003). This approach is frequently used in the literature (Fillol *et al.*, 2016; Jeanbille *et al.*, 2016; Cheutin *et al.*, 2021; Neu *et al.*, 2021) and allows us to distinguish between core and satellite ASVs, while avoiding the use of subjective and arbitrary occurrence and abundance thresholds (Magurran and Henderson, 2003). We first calculated an index of dispersion (*i.e.,* the variance to mean ratio, VMR) for each ASV within each developmental stage. We then tested whether these indices followed a Poisson distribution, falling between the 2.5 and 97.5% confidence limits of the Chi^2 distribution (Krebs, 1999). ASVs with index values outside these confidence limits were considered as part of the core microbiome, while the others were considered as satellite ASVs. This process provided us with stage-dependent core microbiomes for the eggs, each instar nymph stages, adult offspring and mothers, and we finally merged all these cores to obtain a global core microbiome (Figure S1).

Functional predictions of the core microbiomes

We used the PICRUSt2 algorithm (Douglas *et al.*, 2020) to predict the potential functions associated with each of the core microbiomes generated above. In brief, the tool inserts each ASV sequence into a reference tree (EPA-ng, Barbera *et al.*, 2019) using a hidden-state prediction (HSP, castor, Louca and Doebeli, 2018) and infers a KEGG ortholog function (later named KOs) based on the functional profiles

obtained with the nearest-sequenced taxon (MinPath, Ye and Doak, 2009). The

functional table counts with KOs obtained for each sample are count-normalized for

the ASVs copy counts and multiplied by the gene content prediction resulting from

the HSP algorithm. Finally, a table with three levels of KEGG pathways is constructed

according to the MetaCyc database v27.1 (Caspi *et al.*, 2020).

Evolution of bacterial diversity during offspring development

To test how the microbiomes of offspring changed throughout host development, we

calculated the alpha and beta diversities of each sample using qualitative and

quantitative indices for both taxonomic and phylogenetic diversity. Regarding alpha

diversity, which was only calculated for ASVs (not for functions), we normalized all

samples at the minimum sampling size of sequences per individual and checked their

accuracy with rarefaction curves (Cameron *et al.*, 2021; Figure S2). For alpha

diversity proxies, we used direct observed richness for the qualitative taxonomic

richness and the Shannon entropy for its quantitative equivalent. We calculated their

phylogenetic equivalents with the Faith and the Allen's indices (Chao *et al.*, 2010).

For beta diversity, we calculated the relative abundance of each ASV and KOs by

sample, and we calculated the Jaccard distance between samples with ASVs (and

KOs) presence/absence and its quantitative equivalent Bray-Curtis. For ASVs, we

also used their phylogenetic equivalents with the Unifrac metrics for qualitative

(unweighted) and quantitative (weighted) beta diversity distances (Chao *et al.*, 2010;

Yang *et al.*, 2021).

To identify which microbes and/or potential functions, if any, are specific to

each developmental stage, we assessed the differential abundance of ASVs and KOs

with a negative binomial Wald test using the DESeq2 package (Love *et al.*, 2014). In

this model, we entered the developmental stage, which includes all stages with

young- and old-nymphs. Results are presented under heatmaps using the package

pheatmap v1.0.12 where ASVs are merged by Genera and KOs are merged by

KEGG pathway names (third level).

Impact of maternal presence on offspring bacterial diversity

Finally, we tested whether maternal presence influenced the microbiome of their

offspring at both the first developmental stage and the adult stage. To this end, we

repeated the analyses described above using the 5 families in which the mothers

were removed at egg hatching and the 15 families in which the mothers remained

with their nymphs for 15 days after hatching (*i.e.,* until the end of family life). We draw

volcano plots with the package *EnhancedVolcano* v1.14.0 to contrast discriminants

between the presence or absence of the mother.

Statistics

We first tested the effect of developmental stage and moult (fresh versus old moult)

on the microbiome diversity of offspring from the 15 families in which nymphs were

kept with a mother during family life. For alpha diversity, we conducted four linear

mixed models (LMM) in which we entered each of the four microbial alpha diversity

indices as a response variable, the sampling time (Egg, L1-Y, L1-O, L2-Y, L2-O, L3-Y,

L3-O, L4-Y, L4-O, Adult-F, Adult-M and Mother) as the explanatory variable and the

clutch used as a random effect to control for non-independence of the biological

samples. As observed richness can be considered as counts, we performed a

negative binomial mixed model instead of mixed linear models that were used for the

three other alpha diversity proxies. We did not use a classical 2-way ANOVA

approach because the two levels of moult were not available for each developmental

stage (*i.e.,* eggs and adults). We conducted pairwise comparisons between each

sampling time using the estimated marginal means of the models, with P values

corrected for multiple testing using Tukey methods with the *emmeans* package

(Lenth, 2022). For each pairwise comparison, we also calculated R^2 using the *MuMIn*

package (Bartoń, 2022). For beta diversity, the dissimilarity based on the ASVs and

KOs assemblies between each sample was first illustrated in a two-dimensional

Principal Coordinates Analyses (PCoA). We then performed Permutational Analyses

of Variances (PERMANOVAs) to test the effect of the clutch (as randomized block),

the sampling stage and the sex on each distance matrices based on bacterial

composition (*i.e.,* Jaccard, Bray-Curtis, weighted and unweighted Unifrac) and on

functional predictions (*i.e.*, Jaccard and Bray-Curtis). Post-hoc pairwise tests

between stages were performed for each dissimilarity matrix with the package

pairwiseAdonis v0.4.1 (Martinez Arbizu, 2017). These pairwise comparisons allowed

us to address five questions, namely whether (1) microbiome diversity changes

between each successive development stage (*i.e.,* Egg *vs* L1-Y, L1-Y *vs* L1-O, L1-O

vs L2-Y, etc), (2) moulting causes a shift in microbiome diversity between two

successive developmental stages (*i.e.,* Egg *vs* L1-Y, L1-O *vs* L2-Y, etc), (3) old

nymphs exhibit an instar-specific microbiome diversity (*i.e.,* Egg *vs* L1-O, L1-O *vs* L2-

O, L2-O *vs* L3-O, etc), (4) freshly moulted nymphs exhibit an instar-specific

microbiome diversity (*i.e.,* L1-Y *vs* L2-Y, L2-Y *vs* L3-Y, etc) and finally, whether (5)

adults offspring exhibit a sex-specific microbiome diversity (Adult-M *vs* Adult-F).

We tested whether maternal presence had short and/or long-term effects on

the microbiome diversity of their offspring using the 15 families with mothers and the

5 families without mothers. For alpha diversity, we conducted a negative binomial

mixed model for the observed richness and three LMMs. In these models, we entered

each of the index values as a response variable, as well as maternal presence (yes

or no), the developmental instar (L1-O or adult) and the interaction between these

two factors as explanatory variables. When the latter was non-significant, we

removed it after model simplification by AIC comparison. We also entered the clutch

as a random effect. For beta diversity, we repeated the approach detailed above by

testing the effect of the clutch and the presence/absence of a mother in interaction

with the stage (L1-O or adult), on both microbial and functional composition.

We performed all statistical analyses using R v4.2.0 (R Core Team, 2024). We

checked all model assumptions with the *DHARMa* package (Hartig, 2022). We

verified variance homoscedasticity between groups by comparing the distance

dispersion within group with the *betadisper* function (all P > 0.05).

Results

Experiment 1 - Microbiome changes during offspring development (with mothers)

Description of the core microbiome

In the pool of the 218 earwig samples, we detected a total of 915 ASVs core (24.21%

of the initial ASVs diversity), which encompasses 97.67% of the complete sequence

dataset (Figure S1, Table S2). This core microbiome consisted mainly of

Proteobacteria (52.5%), Firmicutes (26.8%), Bacteroidota (13.9%) and

Actinobacteriota (6.6%). They were distributed among 66 bacterial families, of which

Lactobacillaceae (Phylum Firmicutes - 23.5%) and Enterobacteriaceae (Phylum

Proteobacteria - 15.6%) were the most abundant. The abundance of the other

bacterial families depended on the developmental stage of the host (Figure 2).

Figure 2: Composition of the European earwig core microbiome. Data from
Experiments 1 and 2. Individual core microbiome at family scale, ordered by relative
importance, grouped, and coloured by bacterial family. Specimens are ordered
according to their developmental stage with freshly moulted young nymphs (grey) on
one side of the circle, and old nymphs (black) on the other side, with adult females
(Adult-F) and males (Adult-M) separated. The orphan specimens related with the first
instar and adult stages are indicated with dotted circles.

For instance, the families Nocardiaceae (Phylum Actinobacteriota) and

Comamonadaceae (Phylum Proteobacteria) dominated only the egg stage (27.6%

and 13.2% respectively) and the newly hatched nymphs L1-Y (39.2% and 16.9%

respectively). On a finer scale, 50 ASVs were present at all developmental stages, a

number that increased to 142 ASVs when eggs were excluded (Figure S3A). The 50

common ASVs covered 29 genera, represented mainly by *Chryseobacterium,*

Sphingobacterium (Phylum Bacteroidota), *Smaragdicoccus* (Phylum

Actinobacterioa), *Kosakonia, Stenotrophomonas*, *Paracoccus, Pseudomonas* and

Acinetobacter (Phylum Proteobacteria) (Figure S3B). Only a few ASVs were stage

specific. For instance, 2 ASVs (ASV66|*Pseudomonas* and

ASV482|*Sphingobacterium*) were specific to the eggs, 52 to the nymphs and 26 to

the adults (Figure S3A).

Changes in alpha and beta diversity during offspring development

The alpha diversity of the earwig gut microbiome changed during the development of

the offspring (Figure 3A; Figure S4; Shannon index, Likelihood Ratio χ^2_{11} = 59.70, P =

10^{-8} and P < 10^{-10} for all other alpha diversity indices). Markedly, the nature of these

changes did not follow a linear or systematic pattern. Alpha diversity did not change

between eggs and freshly moulted 1^{st} instar nymphs (pairwise comparisons using

log-rank test contrasts for Shannon index; Egg *vs* L1-Y, P = 0.921), nor did it change

during the entire 1^{st} instar (L1-Y *vs* L1-O, P = 0.621). However, there was a gradual

increase in alpha diversity during this time, as it was greater at the end of the 1^{st}

instar than in the eggs (Egg *vs* L1-O, P = 0.052). While this alpha diversity remained

at a high level just after the moult into 2^{nd} instar (L1-O *vs* L2-Y, P = 1.000), it then

dropped to the lowest level at the end of the 2^{nd} instar (L2-Y *vs* L2-O, *P* = 0.029). This

level remained low just after the moult into 3^{rd} instar (L2-O *vs* L3-Y, P = 0.998), but

then increased to its highest level at the end of the 3^{rd} instar (L3-Y *vs* L3-O, P =

0.010). Alpha diversity decreased slightly just after the moult into 4^{th} instar (L3-O *vs*

L4-Y, P = 0.086, but other indices are significant for this contrast) to finally remain

stable until the end of the 4^{th} instar (L4-Y *vs* L4-O, P = 0.702) and in the following

adult females (L4-O *vs* Adult-F, P = 0.496). By contrast, it decreased slightly between

the end of the 4th instar and the following adult males (L4-O *vs* Adult-M, P = 0.055,

but other indices are significant for this contrast).

Figure 3: Microbiome variation during offspring development. The diversity is

compared before (fill) and after moults (empty circles) and between each

developmental stage and between females (Adult-F) and males (Adult-M) (coloured).

(A) Shannon index comparisons for bacterial alpha diversity. Letters above the boxes

indicate significant grouping contrasts (P ≤ 0.05). Boxplots represent the median

(middle bar) and the interquartile range (box) with whiskers representing the 1.5-fold

the interquartile range. Dots represent the value for an individual (with red dot are

outliers). **(B-C)** Principal coordinates analyses (PCoA) plots in the two first axes

illustrating Bray-Curtis distances calculated between samples based on their ASVs

and their KOs compositions.

Overall, moult was not associated with changes in offspring alpha diversity,

except for the final moult from 4th instar to adult males. In old nymphs, alpha diversity

was lower in 2nd compared to 3rd and 4th instars, whereas values were intermediate

in 1st instars and adults (Figure 3A). In contrast, in newly moulted nymphs, alpha

diversity was lower in 3rd instar compared to 2nd instar, while values were

intermediate in 1st and 4th instar (Figure 3A). Finally, in adult offspring, alpha

diversity was similar in males and females (Figure 3A, Figures S3, Supplemental

File). According to what was reported for alpha diversity, the beta diversity of the

earwig gut microbiome changed as the offspring developed (PERMANOVA

performed on all beta diversity metrics, Stage effect $0.074 < R^2 < 0.103$, all P =

0.001) (Table 1).

Table 1: Effect of developmental stage (from eggs to adults) on microbiome alpha
and beta diversity at the bacterial (white) and functional (grey) levels. Significant p-
values (P <.005) are indicated in bold.

	Alpha diversity				Beta diversity			
	Index	R^2	Chi^2	**P**	**Distance**	R^2	**F**	**P**
	Obs. richness	.626	124.29	**<.001**	Jaccard	.074	1.94	**<.001**
Bacteria (ASVs)	Shannon	.335	59.00	**<.001**	Bray-Curtis	.082	2.50	**<.001**
	Faith	.398	71.13	**<.001**	Un. Unifrac	.065	2.73	**<.001**
	Allen	.335	78.57	**<.001**	W. Unifrac	.103	4.82	**<.001**
Functions (KOs)	-	-	-	-	Jaccard	.092	3.42	**<.001**
	-	-	-	-	Bray-Curtis	.089	3.54	**<.001**

Here, however, the beta diversity of each collection time was different from all

other collection times for at least one metric in 61 of the 65 pairwise comparisons (P

< 0.05). Two major microbial turnovers occurred after hatching and during the

passage from the 2^{nd} to the 3^{rd} instar nymphs. The PCoA performed on taxonomical

distance matrices (*i.e.,* Bray-Curtis and Jaccard) showed a marked separation on the

first axis with both microbial assemblies of old 2^{nd} instar nymphs (L2-O) and the

young 3^{rd} instar nymphs (L3-Y) on one side, and all the microbial assemblies

associated with the other stages on the other side (Figure 3B, Figure S5A). The

PCoA based on Unifrac distances shows an additional separation with eggs and

newly hatched nymphs (L1-Y) on one side and the other stages on the other side

(Figure S5B-C) showing a strong phylogenetic turnover. The only four comparisons

that did not present any beta diversity difference were before and after the three first

moults, i.e. between eggs and newly hatched nymphs (L1-Y), between old 1^{st} instar

nymphs (L1-O) and young 2^{nd} instar nymphs (L2-Y), and between old 2^{nd} instar

nymphs (L2-O) and young 3^{rd} instar nymphs (L3-Y), as well as the comparison

between adult males and females (Supplemental File).

Stage-specific microbes in the offspring microbiome

A total of 231 ASVs that belonged to 62 bacterial genera were stage-specific (Figure

4). The hierarchical clustering analysis based on the Z-score of these discriminants

showed a structure that does not strictly follow the chronology of the developmental

stages of the offspring, but rather clustered individuals at the end of one instar to

individuals at the beginning of the following instar. In particular, they grouped the

eggs with newly hatched nymphs (L1-Y), the old 1^{st} (L1-O) with the young 2^{nd} instar

nymphs (L2-Y), and the old 2^{nd} instar (L2-O) with the young 3^{rd} (L3-Y), and the old 4^{th}

instar nymphs (L4-O) with adults (Figure 4).

Figure 4: Heatmaps of hierarchical clustering based on the 62 genera indicators of developmental stages in the European earwig. Raw represent the Z-score associated to the bacterial genus in each sampling stage.

Experiment 2 - Short- and long-term effects of maternal care on offspring microbiome

The presence of the mother after hatching increased the alpha-diversity of their offspring microbiome at both 1st instar and adult stages, but only when the phylogeny was taken into account (Figure 5A; Faith: X^2 = 9.73, P = 0.002 and Hill1 : X^2 = 3.97, P = 0.046). By contrast, it affected neither richness nor entropy (Observed richness: X^2 = 1.86, P = 0.172 and Shannon: X^2 = 1.21, P = 0.271) (Figure S6). In terms of beta-diversity, maternal presence affected the structure of the bacterial communities (for all metrics, 0.009 ≥ P ≥ 0.001) (Figure 5B, C; Figure S7; Table 1), although it only explained a limited proportion of the total variance (for all metrics, 0.074 ≥ R^2 ≥ 0.023). Note that the PERMANOVA applied to Unweighted Unifrac distances calculated on bacterial assemblies might be affected by a non-homogeneous dispersion of the data as individuals that lived with their mother appeared to be more dispersed (*betadisper*, F = 7.7, P = 0.008). However, the ordination in Figure S7 supports a clustering, for both N1 and adult stages, between the samples that lived with their mother or not.

The presence of a mother and the developmental stage were discriminated by 19 ASVs belonging to 12 genera in the microbiomes of the offspring. Among these genera, *Rurimicrobium*, *Rhodoferax* and *Raoutella* were strongly discriminant in individuals that had a mother during early life (Log2Fold change > |20|) while no sequence belonging to these genera was present in the orphaned nymphs (Figure 5D).

Figure 5: Effect of maternal presence (orange) or absence (blue) on the microbiome of 1ˢᵗ instar nymphs (empty circles) and adults (fill circles). (A) Alpha diversity is represented with the phylogenetic diversity index of Faith. **(B, C)** Beta diversity is showed through a two-dimensional PCoA according to the Bray-Curtis distances between microbiome sample pairs for bacterial and functional assemblies. (**C, D**) Volcano plots representing the 19 ASVs and the 33 KEGG pathways discriminating the absence (blue) or presence (orange) of mothers during family life. Boxplots represent the median (middle bar) and the interquartile range (box) with whiskers representing the 1.5-fold the interquartile range. P 0.001 < ** ≤ 0.01.

Experiment 1 & 2 - Predictions of potential functions of the different microbiomes

In terms of function, the 915 ASVs of the global core microbiome were predicted to be associated with 4 952 KOs involved in 226 different KEGG pathways. Most of these pathways are related to metabolic routes such as lipid, carbohydrate or xenobiotics biodegradation and biosynthesis of secondary metabolites (Supplemental File). These functional predictions changed throughout the development of the host (Stage effect $0.089 < R^2 < 0.092$, all $P = 0.001$) (Table 1). They followed the same patterns of microbial assemblies, with a clear distinction between the eggs and newly hatched nymphs, the old 2^{nd} and young 3^{rd} instar nymphs and the rest of all stages (Figure 3C, Figure S8; see Pairwise results in Supplemental File). Among the 226 different KOs, 212 were stage-specific (Figure 6). As for bacteria, the hierarchical clustering of these discriminants showed a structure that does not strictly follow the chronological order of the developmental stages of the offspring (Figure 6).

520
521 **Figure 6: Heatmaps of hierarchical clustering based on the KEGG pathways**
522 **indicators of developmental stages in the European earwig.** Raw represent the
523 Z-score associated to the pathway name in each sampling stage (young-nymphs are
524 coloured in grey).

525 Maternal presence affected the predicted functional structure of the microbiome of

526 offspring (Mother presence $0.035 < R^2 < 0.040$, all $P < 0.009$) (Table 2). This

527 presence discriminated 12 of the 226 KEGG pathways, but only three had a

528 Log2Fold change $> |2.5|$. These pathways included the Fc gamma R-mediated

529 phagocytosis and GnRH signalling pathway in orphans and the isoflavonoid

530 biosynthesis in offspring that had lived with their mother (Figure 5E).

532 **Table 2:** Effect of access to (A) maternal care during family life and (B) offspring age

533 on microbiome alpha and beta diversity at bacterial (white) and functional (grey)

534 levels. Significant p-values (P <.005) are indicated in bold.

		Alpha diversity			**Beta diversity**			
	Index	R^2	Chi^2	P	Distance	R^2	F	P
(A) Maternal care								
Bacteria (ASVs)	Obs. richness	.243	2.72	.099	Jaccard	.023	2.50	**<.001**
	Shannon	.087	.98	.323	Bray-Curtis	.028	3.28	**<.001**
	Faith	.234	8.15	**.004**	Un. Unifrac	.074	9.79	**<.001**
	Allen	.095	2.80	.093	W. Unifrac	.037	5.13	**.003**
Functions (KOs)	-	-	-	-	Jaccard	.035	4.13	**.009**
	-	-	-	-	Bray-Curtis	.040	4.82	**<.001**
(B) Nymphs vs Adults								
Bacteria (ASVs)	Obs. richness	.277	6.97	**.008**	Jaccard	.052	5.68	**<.001**
	Shannon	.137	6.05	**.014**	Bray-Curtis	.077	9.07	**<.001**
	Faith	.209	4.15	**.042**	Un. Unifrac	.108	14.16	**<.001**
	Allen	.181	12.52	**<.001**	W. Unifrac	.178	24.82	**<.001**
Functions (KOs)	-	-	-	-	Jaccard	.083	9.72	**<.001**
	-	-	-	-	Bray-Curtis	.084	10.06	**<.001**

Discussion

The gradual and subtle changes that hemimetabolous juveniles undergo during moulting are often thought to limit profound changes in their microbiota during development (Carrasco and Pérez-Cobas, 2014; Manthey *et al.*, 2022). Our data show that this assumption is not always true, as these changes occur in the European earwig. Using 16S rRNA metabarcoding on 218 samples from egg to adult stages, we found that the bacterial microbiome of earwig offspring shows substantial variation throughout their development both in terms of beta diversity (thereafter called "structure") and alpha diversity (thereafter called "diversity"). Interestingly, these changes did not occur during moulting, but rather between the beginning and end of certain developmental stages. In addition, we found that maternal care partly shapes the bacterial microbiome of offspring, even if this behaviour is facultative in the European earwig. Access to maternal care during the first few weeks after hatching affected not only the bacterial microbiome of first instar nymphs during family life, but also that of the resulting adults two months after family life has ended.

Offspring bacterial microbiome varies during development

Our data first reveal that the bacterial microbiome diversity (alpha) and structure (beta) of earwig changed during offspring development. This is striking for two main reasons. First, these changes were not linear and only occurred at certain stages of offspring development, which contrasts with the general pattern where microbiome diversity gradually increases with natural growth in offspring body size (Sherrill-Mix *et al.*, 2018). Second, some of these changes occurred in the second and third nymphal instars, even though all instars were fed the same diet and developed in the same laboratory conditions throughout their development.

The first marked change in bacterial microbiome structure occurred shortly

after the nymphs hatched. We showed that both eggs and newly hatched nymphs

had a microbiome structure different from that of all other stages. From the egg stage

to the freshly moulted 2nd instar nymphs, we observed an increase in diversity due to

the gradual colonization of the 1st instar nymphs by new genera, including *Erwinia,*

Raoultella, Flavobacterium and *Lactobacillus*. These new genera are present in a

wide range of insects, where they are often known to perform beneficial functions in

juveniles (Malacrinò, 2022). For instance, *Erwinia* is known to reduce the maturation

time of the bark beetle *Ips typographus* (Peral-Aranega *et al.*, 2023).

The second marked change in bacterial microbiome diversity occurred during

the development of the 2nd instar nymphs. We found that the diversity drops to its

lowest level during the transition from newly moulted to old 2nd instar nymphs and

remains low in the newly moulted 3rd instar nymphs (Figure 3A). This fall is reflected

by the loss of several genera, including *Smaragdicoccus, Rhodococcus* or

Alkanindiges (Figure 4). The bacterial communities of the newly moulted 3rd instar

nymphs are very different from those of any other stage, because of its structural

monotony where *Lactobacillus* largely dominates the bacterial composition (Figures

1, 4). Such a decrease in microbiome diversity has been reported in the 3rd instar

larvae of the German cockroach *Blattella germanica* (Carrasco and Pérez-Cobas,

2014), where it has been suggested to result from the physiological state of the host

at that particular stage (Kirkland *et al.*, 2020). In lower termite workers, the gut

flagellates are totally lost prior the ecdysis, probably due to a combination of host

starvation and hormonal variation linked to changes in the host social status

(Cleveland, 1949; Raina *et al.*, 2008; Nalepa, 2017). This may also be the case with

earwigs, although our knowledge of the physiological and hormonal peculiarities of

each instar is still very limited (Meunier, 2024). This decrease may also result from

the end of mother-offspring interactions, as we removed mothers from their nymphs

14 days after hatching, which is often shortly after they have reached the 2nd instar

(Thesing *et al.*, 2015). This end of family life inherently means that any potential form

of coprophagy and trophallaxis between mother and offspring (and thus potential

bacterial vertical transmission) ceases, and the juveniles begin to process the food

source exclusively on their own. Whether physiological peculiarities of 2nd instar

nymphs, suppression of maternal bacterial transfer and/or changes in dietary habits

are the drivers of the reported changes in the microbiome diversity remains to be

further investigated.

The last marked change in bacterial microbiome diversity was an increase

during the development of the 3rd instar nymphs, mostly explained by the colonization

of new genera, such as *Nubsella, Massilia, Gemmobacter* or *Chitinophaga.* These

genera are common colonizers of insect guts (Da Silva Correia *et al.*, 2018; Guégan

et al., 2018; Paddock *et al.*, 2022) but can also be found in the environment

(Mayoral-Peña *et al.*, 2022). For example, bacteria of the genus *Nubsella* are

mutualistic with certain phytophagous insects and are likely to be important for the

fitness of walking sticks (Lü *et al.*, 2019; Li *et al.*, 2020). Interestingly, the genus

Chitinophaga is known to be involved in the degradation of chitin, a major component

of the insect skeleton (Glavina Del Rio *et al.*, 2010). Whether their presence is a

mutualistic association with earwigs aiding in cuticle digestion during cannibalism or

during the reingestion of their own exuviae is an open question and will be developed

below.

Moulting does not alter the bacterial microbiome diversity

Moulting did not induce any changes in the bacterial microbiome diversity and

structure of earwig offspring, except for the moults between the 3rd and 4th instar

nymphs. This overall lack of moulting effect is in line with studies carried out in other

hemimetabolous insects showing that the gut microbiome remains stable during

development (Manthey *et al.*, 2022). For earwigs, we propose several possible

explanations. First, nymphs could possess a microbial reservoir that is preserved

during moulting. Such a reservoir is present, for example, in the bean bug *Riptortus*

clavatus to maintain their *Burkholderia* symbionts during moult (Kikuchi and Yumoto,

2013). However, the presence of such a reservoir has never been documented in

earwigs. Another possibility is that earwig nymphs re-inoculate themselves with their

own bacteria by eating their shed cuticle after moulting, as reported in cockroaches

(Mira, 2000). Even if earwig nymphs are regularly observed eating their shed cuticle

during laboratory rearing (J Meunier, pers. obs), the role of this behaviour to inoculate

a lost microbiome has never been investigated in this species. Finally, earwig nymphs

may be able to re-inoculate themselves through their frequent (in)direct contact with

their siblings by means of coprophagy and trophallaxis (Falk *et al.*, 2014; Kramer and

Meunier, 2016). These social acquisitions allow the persistence and the

(re)acquisition of lost bacteria in numerous arthropods, such as termites (Raina *et al.*,

2008; Michaud *et al.*, 2020), cockroaches (Nalepa, 2020), and spiders (Rose *et al.*,

2023). Overall, these data call for future experiments to disentangle which parameter

explains the resistance of the earwig microbiome to moulting, and to understand why

they are no longer efficient for the moult between the 3rd and 4th instar nymphs.

Bacterial functions could play a role throughout juvenile development

The reported changes in the structure of the offspring bacterial microbiome during

development are associated with changes in the predicted potential functions of their

bacteria that could be beneficial to the host. For instance, many of the bacteria

overrepresented in eggs and newly hatched nymphs are known to produce and

accumulate lipids such as *Rhodococcus, Delftia* and *Pedobacter* (Alvarez *et al.*,

1997; Liu *et al.*, 2016; Franks *et al.*, 2021). As completing the transition from eggs to

nymphs is a highly energetic process, our data suggest that earwig embryos may not

only obtain this energy from egg lipid reserves, but also from these bacteria (Ziegler

and Vanantwerpen, 2006; Diether and Willing, 2019). Moreover, this would be

consistent with the predictions of PICRUSt2, which show a strong positive correlation

in lipogenesis processes such as adipocytokine signalling pathway and unsaturated

fatty acid biosynthesis (Figure 6). The bacterial loss observed in the third nymph

instar was also found in terms of predicted functions, as all pathways except cell

developmental pathways, amino acids and sugar metabolism were underrepresented

in this developmental stage compared to the subsequent ones. Finally, the

acquisition of new genera such as *Nubsella, Massilia, Gemmobacter* or *Chitinophaga*

during the development of the third nymphal instar came with new predicted functions

linked to energy uptake, amino acids and vitamins B biosynthesis, which act as a

coenzyme in numerous pathways involved in the fatty acid synthesis, glucogenesis or

amino acids synthesis. These vitamins are essential during insect development, but

they cannot be synthetized by animals themselves and are often acquired through

alimentation or provided by the microbiota (Douglas, 2017; Kinjo *et al.*, 2022).

Although these predicted functions may provide insights into our general

understanding of the driver of microbiome changes during offspring development,

they must be considered with caution. Indeed, metabarcoding, as a qualitative

approach might be biased by the compositional nature of the data and must be

validated by quantitative approach (Gloor *et al.*, 2017). In addition, the further

functional predictions should be confirmed by transcriptomic analyses as the

approach is debatable due to the short length of the amplicons and the lack of

genome reference concerning insect-associated microbial communities (Djemiel *et*

al., 2022).

Maternal care shapes the bacterial microbiome structure of offspring

In animals, maternal care is often considered to be an important mediator of microbial

transmission from parents to offspring (Bright and Bulgheresi, 2010; Sachs *et al.*,

2011; Hosokawa and Fukatsu, 2020). Our data suggests that the European earwig is

no exception. Access to maternal care not only shapes the bacterial microbiome

structure, phylogenetic diversity and functions of 1st instar nymphs, but also those of

the resulting adults, even though none of these adults had any contact with their

mothers in the previous two months. We found that the bacterial microbiome of

nymphs with maternal care contained a higher diversity of phylogenetically distant

bacteria compared to their orphaned counterparts. Not surprisingly, the bacteria

found in these nymphs were also found in the microbiome of their mothers (including

Raoultella, *Rurimicrobium* and *Sphingobacterium*). This is likely due to direct or

indirect maternal transmission during post-hatching family life. However, even if

maternal transmission contributes to the shape of the bacterial microbiome, the non-

significant effect of maternal care on alpha diversity in terms of taxonomic richness,

Shannon entropy or Allen index, combined with the small, albeit significant effect on

microbial structure, suggests that maternal presence is not the main route of bacterial

acquisition in this species. Indeed, we also found an overabundance of some ASVs

in the orphan offspring, such as sequences related to *Comamonas*, *Lactobacillus* or

Serratia. Since these bacteria are often generalists, associated with laboratory

rearing conditions (Malacrinò, 2022), and common in the mothers tested, they are

likely to come from the non-sterile rearing environment. However, their

overabundance in orphaned nymphs suggests that they were outcompeted by

maternally transmitted bacteria in non-orphaned nymphs. In addition to these

differences, we found one major discriminant (potential) function in the tended

nymphs, related to isoflavonoid biosynthesis, and two in the orphan nymphs, related

to the endocrine and immune systems. These predicted functions are consistent with

previous phenotypic studies showing that orphaned nymphs develop faster to adults,

produce larger adults with longer male appendages, but contradict with other studies

showing that orphaning has limited long-term effects on the basal immunity of the

nymphs and resulting adults (Meunier and Kölliker, 2012; Thesing *et al.*, 2015;

Vogelweith *et al.*, 2017; Körner *et al.*, 2020). Here again, these predicted functions

need to be taken with caution, especially as recent studies demonstrated that the

alteration of the microbiome does not affect mother earwigs (Van Meyel *et al.*, 2021)

and that its natural variation does not explain its aggregation behaviour (Cheutin *et*

al., 2024), calling into question the need to have a microbiome (Hammer *et al.*, 2017,

2019). Whether all or some of these offspring phenotypes are indeed due to

maternally derived bacteria remains to be further explored, *e.g.* with the use of

gnotobiotic lineages.

Conclusion

Overall, we showed that the European earwig bacterial microbiome changes multiple

times during offspring development. Interestingly, these changes were independent

of moulting. The fact that moulting did not induce any purge or shift in the bacterial

communities of the nymphs calls for future studies to test whether this is due to the

presence of a bacterial reservoir, moult consumption and/or social interactions with

siblings. Our data suggest that the predicted functions of some components of these

microbiomes are relevant to the developmental stage at which they occur, such as

lipogenesis or steroid synthesis in early stages, and nutrient and vitamin synthesis in

late stages. However, future studies are required to confirm the functional role of the

microbiome changes in this species. Finally, we showed that maternal care is an

important short- and long-term determinant of the offspring bacterial microbiome.

Given that earwig nymphs do not require maternal care to develop and survive and

that nymph can also develop in absence of any social interactions, our results call for

future studies to investigate the role of these socially-acquired bacteria (and other

potential members of their microbiota such as fungi, viruses and other

microorganisms) in the biology of the European earwig and, more generally, in the

early evolution and maintenance of facultative family life in insects.

Acknowledgements

We thank Franck Dedeine, Maximilian Körner and Jean-Claude Tourneur for their

comments on this manuscript. We also thank all members of the EARWIG group at

IRBI for their help with animal rearing and setup installation, Marina Querejeta for her

help on bioinformatical analyses, and the Bio-environment platform of the University

of Perpignan (France) for providing the MiSeq Illumina sequencing. Finally, we thank

Armand Guillermin and the INRAE unité expérimentale Recherche Intégrée in

Gotheron for giving us access to their orchards for earwig field sampling. Graphical

abstract was made with Biorender.com.

Author Contributions

All authors contributed to the study conception. The manuscript was written by M-C Cheutin and J. Meunier; Analyses were carried by M-C Cheutin and J. Meunier; Molecular analyses, sampling and animal rearing were performed by M-C Cheutin and M. Boucicot. All authors read and approved the final version of the manuscript.

Data Availability

The data set and scripts are available on Zenodo with the DOI 10.5281/zenodo.10776543. Libraries for each sample are deposited in the NCBI Sequence Read Archive (SRA) under BioProject accession no. PRJNA1066258.

Funding statement

This research has been supported by a research grant from the Agence Nationale de la Recherche (ANR-20-CE02-0002 to J.M.).

Conflict of interest

We declare that our work does not have any conflict of interest.

Ethics approval statements

Our investigation complies with the current European Directive 2010/63/EU that does not acquire ethical approval on invertebrates. All animals were handled with care until necessary sacrifices.

References

Alvarez, H.M., Kalscheuer, R., and Steinbüchel, A. (1997) Accumulation of storage lipids in species of *Rhodococcus* and *Nocardia* and effect of inhibitors and polyethylene glycol. *Fett/Lipid* **99**: 239–246.

Arce, A.N., Johnston, P.R., Smiseth, P.T., and Rozen, D.E. (2012) Mechanisms and fitness effects of antibacterial defences in a carrion beetle: Antibacterial defence in a carrion beetle. *Journal of Evolutionary Biology* **25**: 930–937.

Barbera, P., Kozlov, A.M., Czech, L., Morel, B., Darriba, D., Flouri, T., and Stamatakis, A. (2019) EPA-ng: Massively parallel evolutionary placement of genetic sequences. *Systematic Biology* **68**: 365–369.

Bartoń, K. (2022) MuMIn: Multi-Model Inference.

Berenbaum, M. (2017) Insect biodiversity - Millions and millions. In *Insect Biodiversity*. Foottit, R.G. and Adler, P.H. (eds). Chichester, UK: John Wiley & Sons, Ltd, pp. 783–792.

Berlanga, M. and Guerrero, R. (2016) The holobiont concept: the case of xylophagous termites and cockroaches. *Symbiosis* **68**: 49–60.

Boos, S., Meunier, J., Pichon, S., and Kölliker, M. (2014) Maternal care provides antifungal protection to eggs in the European earwig. *Behavioral Ecology* **25**: 754–761.

Bright, M. and Bulgheresi, S. (2010) A complex journey: transmission of microbial symbionts. *Nat Rev Microbiol* **8**: 218–230.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes, S.P. (2016) DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* **13**: 581–583.

Cameron, E.S., Schmidt, P.J., Tremblay, B.J.-M., Emelko, M.B., and Müller, K.M. (2021) Enhancing diversity analysis by repeatedly rarefying next generation sequencing data describing microbial communities. *Sci Rep* **11**: 22302.

Carrasco, P. and Pérez-Cobas, A.E. (2014) Succession of the gut microbiota in the cockroach *Blattella germanica*. *International Microbiology* 99–109.

Caspi, R., Billington, R., Keseler, I.M., Kothari, A., Krummenacker, M., Midford, P.E., et al. (2020) The MetaCyc database of metabolic pathways and enzymes - a 2019 update. *Nucleic Acids Research* **48**: D445–D453.

Chao, A., Chiu, C.-H., and Jost, L. (2010) Phylogenetic diversity measures based on Hill numbers. *Phil Trans R Soc B* **365**: 3599–3609.

Cheutin, M.-C., Leclerc, B., and Meunier, J. (2024) With or without you: gut microbiota does not predict aggregation behavior in European earwig females. *Behavioral Ecology* **35**: arae022.

Cheutin, M.-C., Villéger, S., Hicks, C.C., Robinson, J.P.W., Graham, N.A.J., Marconnet, C., et al. (2021) Microbial shift in the enteric bacteriome of coral reef fish following climate-driven regime shifts. *Microorganisms* **9**: 1711.

Cleveland, L.R. (1949) Hormone-induced sexual cycles of flagellates. I. Gametogenesis, fertilization, and meiosis in *Trichonympha*. *Journal of Morphology* **85**: 197–295.

Da Silva Correia, D., Passos, S.R., Proença, D.N., Morais, P.V., Xavier, G.R., and Correia, M.E.F. (2018) Microbial diversity associated to the intestinal tract of soil invertebrates. *Applied Soil Ecology* **131**: 38–46.

Diehl, J.M.C. and Meunier, J. (2018) Surrounding pathogens shape maternal egg care but not egg production in the European earwig. *Behavioral Ecology* **29**: 128–136.

Diether, N. and Willing, B. (2019) Microbial fermentation of dietary protein: An important factor in diet–microbe–host interaction. *Microorganisms* **7**: 19.

Djemiel, C., Maron, P.-A., Terrat, S., Dequiedt, S., Cottin, A., and Ranjard, L. (2022)
Inferring microbiota functions from taxonomic genes: a review. *GigaScience* **11**:
giab090.

Douglas (2017) The B vitamin nutrition of insects: the contributions of diet, microbiome
and horizontally acquired genes. *Current Opinion in Insect Science* **23**: 65–69.

Douglas, G.M., Maffei, V.J., Zaneveld, J.R., Yurgel, S.N., Brown, J.R., Taylor, C.M., et al.
(2020) PICRUSt2 for prediction of metagenome functions. *Nat Biotechnol* **38**: 685–
688.

Engel, P. and Moran, N.A. (2013) The gut microbiota of insects – diversity in structure
and function. *FEMS Microbiol Rev* **37**: 699–735.

Falk, J., Wong, J.W.Y., Kölliker, M., and Meunier, J. (2014) Sibling cooperation in earwig
families provides insights into the early evolution of social life. *The American*
Naturalist **183**: 547–557.

Fillol, M., Auguet, J.-C., Casamayor, E.O., and Borrego, C.M. (2016) Insights in the
ecology and evolutionary history of the *Miscellaneous Crenarchaeotic Group* lineage.
The ISME Journal **10**: 665–677.

Franks, K., Kooienga, E., Sanders, M., Pendarvis, K., Yang, F., Tomberlin, J.K., and
Jordan, H.R. (2021) The effect of *Rhodococcus rhodochrous* supplementation on
black soldier fly (Diptera: Stratiomyidae) development, nutrition, and waste
conversion. *JIFF* **7**: 397–408.

Fukatsu, T. and Hosokawa, T. (2002) Capsule-Transmitted Gut Symbiotic Bacterium of
the Japanese Common Plataspid Stinkbug, *Megacopta punctatissima*. *Appl Environ*
Microbiol **68**: 389–396.

Girard, M., Luis, P., Valiente Moro, C., and Minard, G. (2023) Crosstalk between the
microbiota and insect postembryonic development. *Trends in Microbiology* **31**: 181–
196.

Glassman, S.I. and Martiny, J.B.H. (2018) Broadscale ecological patterns are robust to
use of Exact Sequence Variants versus Operational Taxonomic Units. *mSphere* **3**:
e00148-18.

Glavina Del Rio, T., Abt, B., Spring, S., Lapidus, A., Nolan, M., Tice, H., et al. (2010)
Complete genome sequence of *Chitinophaga pinensis* type strain (UQM 2034T).
Stand Genomic Sci **2**: 87–95.

Gloor, G.B., Macklaim, J.M., Pawlowsky-Glahn, V., and Egozcue, J.J. (2017) Microbiome
datasets are compositional: and this is not optional. *Front Microbiol* **8**: 2224.

González-Miguéns, R., Muñoz-Nozal, E., Jiménez-Ruiz, Y., Mas-Peinado, P., Ghanavi,
H.R., and García-París, M. (2020) Speciation patterns in the *Forficula auricularia*
species complex: cryptic and not so cryptic taxa across the western Palaearctic
region. *Zoological Journal of the Linnean Society* **190**: 788–823.

Greer, J.A., Swei, A., Vredenburg, V.T., and Zink, A.G. (2020) Parental care alters the
egg microbiome of maritime earwigs. *Microb Ecol* **80**: 920–934.

Guégan, M., Zouache, K., Démichel, C., Minard, G., Tran Van, V., Potier, P., et al. (2018)
The mosquito holobiont: fresh insight into mosquito-microbiota interactions.
Microbiome **6**: 49.

Hammer, T.J., Janzen, D.H., Hallwachs, W., Jaffe, S.P., and Fierer, N. (2017)
Caterpillars lack a resident gut microbiome. *Proc Natl Acad Sci USA* **114**: 9641–
9646.

Hammer, T.J. and Moran, N.A. (2019) Links between metamorphosis and symbiosis in
holometabolous insects. *Phil Trans R Soc B* **374**: 20190068.

Hammer, T.J., Sanders, J.G., and Fierer, N. (2019) Not all animals need a microbiome.
FEMS Microbiology Letters **366**: fnz117.

Hartig, F. (2022) DHARMa: Residual diagnostics for hierarchical (multi-Level / mixed)
regression models. *R package version 046*.

Hosokawa, T. and Fukatsu, T. (2020) Relevance of microbial symbiosis to insect
behavior. *Current Opinion in Insect Science* **39**: 91–100.

Hosokawa, T., Hironaka, M., Inadomi, K., Mukai, H., Nikoh, N., and Fukatsu, T. (2013)
Diverse strategies for vertical symbiont transmission among subsocial stinkbugs.
PLoS ONE **8**: e65081.

Jeanbille, M., Gury, J., Duran, R., Tronczynski, J., Agogué, H., Ben Saïd, O., et al.
(2016) Response of core microbial consortia to chronic hydrocarbon contaminations
in coastal sediment habitats. *Front Microbiol* **7**:.

Johnston, P.R., Paris, V., and Rolff, J. (2019) Immune gene regulation in the gut during
metamorphosis in a holo- versus a hemimetabolous insect. *Phil Trans R Soc B* **374**:
20190073.

Katoh, K. (2002) MAFFT: a novel method for rapid multiple sequence alignment based
on fast Fourier transform. *Nucleic Acids Research* **30**: 3059–3066.

Kikuchi, Y., Hosokawa, T., and Fukatsu, T. (2008) Diversity of bacterial symbiosis in
stinkbugs. In *Microbial Ecology Research Trends*. pp. 39–63.

Kikuchi, Y. and Yumoto, I. (2013) Efficient colonization of the bean bug *Riptortus*
pedestris by an environmentally transmitted *Burkholderia* symbiont. *Appl Environ*
Microbiol **79**: 2088–2091.

Kinjo, Y., Bourguignon, T., Hongoh, Y., Lo, N., Tokuda, G., and Ohkuma, M. (2022)
Coevolution of metabolic pathways in Blattodea and their *Blattabacterium*
endosymbionts, and comparisons with other insect-bacteria symbioses. *Microbiol*
Spectr **10**: e02779-22.

Kirkland, L.S., Maino, J., Stuart, O., and Umina, P.A. (2020) Ontogeny in the European
earwig (*Forficula auricularia*) and grain crops interact to exacerbate feeding damage
risk. *J Applied Entomology* **144**: 605–615.

Koch, L.K. and Meunier, J. (2014) Mother and offspring fitness in an insect with maternal
care: phenotypic trade-offs between egg number, egg mass and egg care. *BMC Evol*
Biol **14**: 125.

Kölliker, M. (2007) Benefits and costs of earwig (*Forficula auricularia*) family life. *Behav*
Ecol Sociobiol **61**: 1489–1497.

Körner, M., Foitzik, S., and Meunier, J. (2018) Extended winters entail long-term costs for
insect offspring reared in an overwinter burrow. *Journal of Thermal Biology* **74**: 116–
122.

Körner, M., Vogelweith, F., Libbrecht, R., Foitzik, S., Feldmeyer, B., and Meunier, J.
(2020) Offspring reverse transcriptome responses to maternal deprivation when
reared with pathogens in an insect with facultative family life. *Proc R Soc B* **287**:
20200440.

Kramer, J. and Meunier, J. (2016) Kin and multilevel selection in social evolution: a
never-ending controversy? *F1000Res* **5**: 776.

Kramer, J., Thesing, J., and Meunier, J. (2015) Negative association between parental
care and sibling cooperation in earwigs: a new perspective on the early evolution of
family life? *J of Evolutionary Biology* **28**: 1299–1308.

Krebs, C. (1999) Ecological methodology, 2nd ed. New York: Harper and Row.

Lamb, R.J. (1976) Parental behavior in the Dermaptera with special reference to
Forficula auricularia (Dermaptera: Forficulidae). *Can Entomol* **108**: 609–619.

Lenth, R. (2022) emmeans: Estimated Marginal Means, aka Least-Squares Means.

Li, Y., Huang, Y., Chen, T., Wu, S., Tang, H., Hsiao, C., et al. (2020) Comparison of gut
microbiota of healthy and diseased walking sticks, *Phasmotaenia lanyuhensis*. *Arch*

Insect Biochem Physiol **105**: e21749.

Liu, W., Li, Y., Guo, S., Yin, H., Lei, C.-L., and Wang, X.-P. (2016) Association between
gut microbiota and diapause preparation in the cabbage beetle: a new perspective
for studying insect diapause. *Sci Rep* **6**: 38900.

Lombardo, M.P. (2008) Access to mutualistic endosymbiotic microbes: an
underappreciated benefit of group living. *Behav Ecol Sociobiol* **62**: 479–497.

Louca, S. and Doebeli, M. (2018) Efficient comparative phylogenetics on large trees.
Bioinformatics **34**: 1053–1055.

Love, M.I., Huber, W., and Anders, S. (2014) Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**: 550.

Lü, J., Guo, W., Chen, S., Guo, M., Qiu, B., Yang, C., et al. (2019) Host plants influence
the composition of the gut bacteria in *Henosepilachna vigintioctopunctata*. *PLoS*
ONE **14**: e0224213.

Magurran, A.E. and Henderson, P.A. (2003) Explaining the excess of rare species in
natural species abundance distributions. *Nature* **422**: 714–716.

Malacrinò, A. (2022) Host species identity shapes the diversity and structure of insect
microbiota. *Molecular Ecology* **31**: 723–735.

Manthey, C., Johnston, P.R., Nakagawa, S., and Rolff, J. (2022) Complete
metamorphosis and microbiota turnover in insects. *Molecular Ecology* mec.16673.

Martinez Arbizu, P. (2017) pairwiseAdonis: Pairwise Multilevel Comparison using Adonis.

Mayoral-Peña, Z., Lázaro-Vidal, V., Fornoni, J., Álvarez-Martínez, R., and Garrido, E.
(2022) Studying plant–insect interactions through the analyses of the diversity,
composition, and functional inference of their bacteriomes. *Microorganisms* **11**: 40.

McFrederick, Q.S., Wcislo, W.T., Hout, M.C., and Mueller, U.G. (2014) Host species and
developmental stage, but not host social structure, affects bacterial community
structure in socially polymorphic bees. *FEMS Microbiol Ecol* **88**: 398–406.

McMurdie, P.J. and Holmes, S. (2013) phyloseq: an R package for reproducible
interactive analysis and graphics of microbiome census data. *PLoS ONE* **8**: e61217.

Meunier, J. (2024) The biology and social life of Earwigs (Dermaptera). *Annu Rev*
Entomol **69**: annurev-ento-013023-015632.

Meunier, J. and Kölliker, M. (2012) When it is costly to have a caring mother: food
limitation erases the benefits of parental care in earwigs. *Biol Lett* **8**: 547–550.

Meunier, J., Körner, M., and Kramer, J. (2022) Parental care. In *Reproductive Strategies*
in Insects. Boca Raton: CRC Press.

Meunier, J., Wong, J.W.Y., Gómez, Y., Kuttler, S., Röllin, L., Stucki, D., and Kölliker, M.
(2012) One clutch or two clutches? Fitness correlates of coexisting alternative female
life-histories in the European earwig. *Evol Ecol* **26**: 669–682.

Michaud, C., Hervé, V., Dupont, S., Dubreuil, G., Bézier, A.M., Meunier, J., et al. (2020)
Efficient but occasionally imperfect vertical transmission of gut mutualistic protists in
a wood-feeding termite. *Molecular Ecology* **29**: 308–324.

Mira, A. (2000) Exuviae eating: a nitrogen meal? *Journal of Insect Physiology* **46**: 605–
610.

Mora, C., Tittensor, D.P., Adl, S., Simpson, A.G.B., and Worm, B. (2011) How many
species are there on Earth and in the Ocean? *PLoS Biol* **9**: e1001127.

Muyzer, G., de Waal, E.C., and Uitterlinden, A.G. (1993) Profiling of complex microbial
populations by denaturing gradient gel electrophoresis analysis of polymerase chain
reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* **59**: 695–700.

Nalepa (2020) Origin of mutualism between termites and flagellated gut protists:
transition from horizontal to vertical transmission. *Front Ecol Evol* **8**: 14.

Nalepa, C. (2017) What kills the hindgut flagellates of lower termites during the host

molting cycle? *Microorganisms* **5**: 82.

Neu, A.T., Allen, E.E., and Roy, K. (2021) Defining and quantifying the core microbiome:
Challenges and prospects. *Proc Natl Acad Sci USA* **118**: e2104429118.

Onchuru, T.O., Javier Martinez, A., Ingham, C.S., and Kaltenpoth, M. (2018)
Transmission of mutualistic bacteria in social and gregarious insects. *Current Opinion*
in Insect Science **28**: 50–58.

Paddock, K.J., Finke, D.L., Kim, K.S., Sappington, T.W., and Hibbard, B.E. (2022)
Patterns of microbiome composition vary across spatial scales in a specialist insect.
Front Microbiol **13**: 898744.

Peral-Aranega, E., Saati-Santamaría, Z., Ayuso-Calles, M., Kostovčík, M., Veselská, T.,
Švec, K., et al. (2023) New insight into the bark beetle *Ips typographus* bacteriome
reveals unexplored diversity potentially beneficial to the host. *Environmental*
Microbiome **18**: 53.

Powell, J.E., Martinson, V.G., Urban-Mead, K., and Moran, N.A. (2014) Routes of
acquisition of the gut microbiota of the honey bee *Apis mellifera*. *Appl Environ*
Microbiol **80**: 7378–7387.

Price, M.N., Dehal, P.S., and Arkin, A.P. (2010) FastTree 2 – Approximately maximum-
likelihood trees for large alignments. *PLoS ONE* **5**: e9490.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., et al. (2012) The
SILVA ribosomal RNA gene database project: improved data processing and web-
based tools. *Nucleic Acids Research* **41**: D590–D596.

R Core Team (2024) R: A language and environment for statistical computing.

Raina, A., Park, Y.I., and Gelman, D. (2008) Molting in workers of the formosan
subterranean termite *Coptotermes formosanus*. *Journal of Insect Physiology* **54**:
155–161.

Rose, C., Lund, M.B., Søgård, A.M., Busck, M.M., Bechsgaard, J.S., Schramm, A., and
Bilde, T. (2023) Social transmission of bacterial symbionts homogenizes the
microbiome within and across generations of group-living spiders. *ISME*
Communications **3**: 60.

Sachs, J.L., Skophammer, R.G., and Regus, J.U. (2011) Evolutionary transitions in
bacterial symbiosis. *Proc Natl Acad Sci USA* **108**: 10800–10807.

Salem, H., Florez, L., Gerardo, N., and Kaltenpoth, M. (2015) An out-of-body experience:
the extracellular dimension for the transmission of mutualistic bacteria in insects.
Proc R Soc B **282**: 20142957.

Samways, M.J. (2018) Insect conservation for the twenty-first century. In *Insect science-*
diversity, conservation and nutrition. IntechOpen.

Sandrin, L., Meunier, J., Raveh, S., Walser, J.-C., and Kölliker, M. (2015) Multiple
paternity and mating group size in the European earwig, *Forficula auricularia*. *Ecol*
Entomol **40**: 159–166.

Schliep, K.P. (2011) phangorn: phylogenetic analysis in R. *Bioinformatics* **27**: 592–593.

Shapira, M. (2016) Gut microbiotas and host evolution: Scaling up symbiosis. *Trends in*
Ecology & Evolution **31**: 539–549.

Sherrill-Mix, S., McCormick, K., Lauder, A., Bailey, A., Zimmerman, L., Li, Y., et al.
(2018) Allometry and ecology of the bilaterian gut microbiome. *mBio* **9**: e00319-18.

Sudakaran, S., Kost, C., and Kaltenpoth, M. (2017) Symbiont acquisition and
replacement as a source of ecological innovation. *Trends in Microbiology* **25**: 375–
390.

Sudakaran, S., Salem, H., Kost, C., and Kaltenpoth, M. (2012) Geographical and
ecological stability of the symbiotic mid-gut microbiota in European firebugs,
Pyrrhocoris apterus (Hemiptera, Pyrrhocoridae). *Molecular Ecology* **21**: 6134–6151.

Thesing, J., Kramer, J., Koch, L.K., and Meunier, J. (2015) Short-term benefits, but
transgenerational costs of maternal loss in an insect with facultative maternal care.
Proc R Soc B **282**: 20151617.
Tourneur, Cole, C., and Meunier, J. (2020) The first of five moults of *Forficula auricularia*
L. (Dermaptera: Forficulidae). *Can Entomol* **152**: 783–789.
Tourneur and Gingras, J. (1992) Egg laying in a northeastern north american (Montreal,
Quebec) population of *Forficula auricularia* L. (Dermaptera: Forficulidae). *Can*
Entomol **124**: 1055–1061.
Tourneur, J. and Meunier, J. (2020) Variations in seasonal (not mean) temperatures
drive rapid adaptations to novel environments at a continent scale. *Ecology* **101**:
e02973.
Tourneur, J.-C., Cole, C., Vickruck, J., Dupont, S., and Meunier, J. (2022) Pre- and post-
oviposition behavioural strategies to protect eggs against extreme winter cold in an
insect with maternal care. *Peer Community Journal* **2**: e21.
Trumbo, S. (2012) Patterns of parental care in invertebrates. In *The evolution of parental*
care. Oxford: Oxford University Press, pp. 81–93.
Van Meyel, S., Devers, S., Dupont, S., Dedeine, F., and Meunier, J. (2021) Alteration of
gut microbiota with a broad-spectrum antibiotic does not impair maternal care in the
European earwig. *J of Evolutionary Biology* **34**: 1034–1045.
Van Meyel, S. and Meunier, J. (2020) Filial egg cannibalism in the European earwig: its
determinants and implications in the evolution of maternal egg care. *Animal*
Behaviour **164**: 155–162.
Vogelweith, F., Körner, M., Foitzik, S., and Meunier, J. (2017) Age, pathogen exposure,
but not maternal care shape offspring immunity in an insect with facultative family life.
BMC Evol Biol **17**: 69.
Wang, Y. and Rozen, D.E. (2017) Gut microbiota colonization and transmission in the
burying beetle *Nicrophorus vespilloides* throughout development. *Appl Environ*
Microbiol **83**: e03250-16.
Wirth, T., Le Guellec, R., Vancassel, M., and Veuille, M. (1998) Molecular and
reproductive characterization of sibling species in the European earwig (*Forficula*
auricularia). *Evolution* **52**: 260–265.
Wong, J.W.Y. and Kölliker, M. (2012) The effect of female condition on maternal care in
the European earwig.
Yang, Z., Xu, F., Li, H., and He, Y. (2021) Beyond samples: a metric revealing more
connections of gut microbiota between individuals. *Computational and Structural*
Biotechnology Journal **19**: 3930–3937.
Ye, Y. and Doak, T.G. (2009) A parsimony approach to biological pathway
reconstruction/inference for genomes and metagenomes. *PLoS Comput Biol* **5**:
e1000465.
Zhukova, M., Sapountzis, P., Schiøtt, M., and Boomsma, J.J. (2017) Diversity and
transmission of gut bacteria in *Atta* and *Acromyrmex* leaf-cutting ants during
development. *Front Microbiol* **8**: 1942.
Ziegler, R. and Vanantwerpen, R. (2006) Lipid uptake by insect oocytes. *Insect*
Biochemistry and Molecular Biology **36**: 264–272.
1052