MftG is crucial for alcohol metabolism of mycobacteria by linking mycofactocin oxidation to respiration

Ana Patrícia Graça[1], Vadim Nikitushkin[1,2], Mark Ellerhorst[1,2], Cláudia Vilhena[3], Tilman E. Klassert[4,5], Andreas Starick[6,7], Malte Siemers[6,7], Walid K. Al-Jammal[8], Ivan Vilotijevic[8], Hortense Slevogt[4,5], Kai Papenfort[6,7], and Gerald Lackner[1,2,7]

[1]Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute, Junior Research Group Synthetic Microbiology, Beutenbergstr. 11a, 07745 Jena, Germany

[2]University of Bayreuth, Chair of Biochemistry of Microorganisms, Fritz-Hornschuch-Str. 13, 95326, Kulmbach, Germany

[3]Leibniz Institute for Natural Product Research and Infection Biology– Hans Knöll Institute, Department of Infection Biology, Beutenbergstr. 11a, 07745 Jena, Germany

[4]Respiratory Infection Dynamics, Helmholtz Centre for Infection Research - HZI Braunschweig, Germany

[5]Department of Respiratory Medicine and Infectious Diseases, Hannover Medical School, German Center for Lung Research (DZL), BREATH, Hannover, Germany

[6]Friedrich Schiller University Jena, Institute of Microbiology, Winzerlaer Str. 2, 07745, Jena, Germany

[7]Microverse Cluster, Friedrich Schiller University Jena, 07743 Jena, Germany.

[8]Friedrich Schiller University Jena, Institute of Organic Chemistry and Macromolecular Chemistry, Humboldtstraße 10, 07743 Jena, Germany

Abstract

Mycofactocin is a redox cofactor essential for the alcohol metabolism of Mycobacteria. While the biosynthesis of mycofactocin is well established, the gene *mftG*, which encodes an oxidoreductase of the glucose-methanol-choline superfamily, remained functionally uncharacterized. Here, we show that MftG enzymes strictly require *mft* biosynthetic genes and are found in 75% of organisms harboring these genes. Gene deletion experiments in *Mycolicibacterium smegmatis* demonstrated a growth defect of the Δ*mftG* mutant on ethanol as a carbon source, accompanied by an arrest of cell division reminiscent of mild starvation. Investigation of carbon and cofactor metabolism implied a defect in mycofactocin reoxidation. Cell-free enzyme assays and respirometry using isolated cell membranes indicated that MftG acts as a mycofactocin dehydrogenase shuttling electrons toward the respiratory chain. Transcriptomics studies also indicated remodeling of redox metabolism to compensate for a shortage of redox equivalents. In conclusion, this work closes an important knowledge gap concerning the mycofactocin system and adds a new pathway to the intricate web of redox reactions governing the metabolism of mycobacteria.

Keywords

Mycofactocin, mycobacteria, respiration, redox-cofactor, oxidoreductase

Introduction

Mycobacteria are a metabolically versatile group of microorganisms that are able to utilize various organic compounds for their carbon and energy metabolism. Especially environmental mycobacteria like *M. smegmatis* utilize for instance sugars, polyols, small organic acids, fatty acids, or sterols for their growth (1). While obligate pathogens like *M.*

tuberculosis, the causative agent of tuberculosis, are more limited in their menu, they still retain the ability to metabolize various nutrients available in the host organism (2, 3). Intriguingly, alcohols like ethanol are readily consumed by many mycobacterial species, but the metabolic pathway for ethanol consumption requires genes related to the biosynthesis of the unusual redox cofactor mycofactocin (MFT) (4, 5). Indeed, the inactivation of the mycofactocin biosynthetic gene cluster *mftA-F* (**Figure 1A**) showed a severe impact on the growth of *M. smegmatis* and *M. marinum* cultivated on ethanol as the sole carbon source. For *M. tuberculosis,* a similar growth deficit on media containing ethanol and cholesterol was observed (5). In *M. smegmatis*, it was shown that the gene MSMEG_6242, encoding a MFT-associated alcohol dehydrogenase (termed Mno or Mdo), is strictly required for alcohol utilization. Mno/Mdo was also demonstrated to catalyze methanol metabolism in the same organism (6) and involvement of mycofactocin in this process was suggested (7). Furthermore, a metabolomics study conducted in our lab revealed that MFT production in *M. smegmatis* was significantly increased on ethanol as carbon source in comparison to a control cultivated on glucose (8). Recently, a homolog of Mdo was shown to be essential for MFT-dependent ethylene glycol oxidation in *Rhodococcus jostii* (9). All of this evidence supported the hypothesis that mycofactocin enables alcohol metabolism by acting as an electron acceptor of alcohol dehydrogenases (**Figure 1B**) (4, 5).

Although the role of MFT in alcohol metabolism is well established, further biological roles of mycofactocin appear to exist. Intriguingly, the MFT gene locus seems to be de-repressed by long-chain fatty acid-CoA esters in *M. smegmatis,* which bind to the repressor *mftR* (10). Furthermore, a comparison of the proteome of active and dormant cells revealed that *mftD* (Rv0694) was upregulated in the dormant state, which is associated with the persistence of *M. tuberculosis* in macrophages (11). Lastly, a study

on the impact of mycofactocin inactivation (Δ*mftD*) on *M. tuberculosis* showed increased

growth on glucose containing media as well as a decreased number of mycobacterial

cells in a mouse model after the onset of hypoxia (12).

Mycofactocin is a ribosomally synthesized and post-translationally modified peptide

(RiPP) (13). After translation, the precursor peptide MftA is bound by its chaperone MftB

and undergoes several modifications until it matures into a redox-active molecule. More

precisely, its C-terminal core peptide (Val-Tyr) is decarboxylated and cyclized by the

radical SAM maturase MftC (14). The cyclized core peptide (AHDP) is liberated from the

precursor by the peptidase MftE (15). The flavoenzyme MftD further deaminates AHDP

to premycofactocin (PMFT) thus creating a redox-active ketoamide moiety (16). *In vivo*,

mycofactocins exist in oligoglycosylated forms (MFT-n), where n denotes the number of

β-1,4-linked glucose moieties. The glycosylation requires the presence of the

glycosyltransferase MftF (8). In some mycobacteria, the glycosyl chain of MFT-n is

further modified by methylation at the second sugar moiety (yielding MMFT-n) by the

mycofactocin-associated methyltransferase MftM (17).

The reduction of M/MFT-n to M/MFT-nH_2 during growth on ethanol is likely to be

catalyzed by the above-mentioned alcohol dehydrogenase Mno/Mdo (5, 7). However,

two important questions remained open concerning the mycofactocin system. First, it is

unclear how MFT-nH_2 is re-oxidized to MFT-n after reduction by Mno/Mdo or other

enzymes (**Figure 1B**). Such a step would be crucial to regenerate the cofactor and allow

for further electron transfer to terminal electron acceptors such as oxygen. Second, the

role of the *mftG* gene (**Figure 1A**), the terminal gene in the *mft* gene cluster of

mycobacteria, encoding a flavoenzyme of the glucose-methanol-choline (GMC)

oxidoreductase superfamily remained unknown. GMC oxidoreductases are present in

bacteria, fungi, plants, and insects (18, 19) and are known to catalyze redox reactions

using a variety of different types of substrates and electron acceptors. Prominent

examples are the fungal glucose oxidases, which are used in portable glucose

biosensors for diabetes patients (20). Other important representatives are the alcohol

oxidases from the methylotrophic yeast *Pichia pastoris,* which uses oxygen as an

electron acceptor (21), or the choline dehydrogenase of *E. coli*, which feeds electrons

into the electron transport chain (22). We therefore hypothesized that MftG may be an

oxidoreductase involved in mycofactocin-related metabolism (4). Here, we show that

mftG (MSMEG_1428), encoding a flavoenzyme of the GMC family, is crucial for ethanol

metabolism of the model organism *Mycolicibacterium smegmatis.* We further present

evidence that MftG acts as a mycofactocin dehydrogenase and promotes MFT

regeneration via electron transfer to the respiratory chain of mycobacteria.

Figure 1: The mycofactocin redox system. (A) Schematic representation of the *mft* gene cluster of *M. smegmatis. mftA-F:* MFT biosynthetic genes. *mftR*: TetR-like regulator. *mftG*: GMC oxidoreductase (subject of this study). (B) Chemical structures of MMFT-n (oxidized methylmycofactocin) and MMFT-nH$_2$ (reduced form) and hypothetical scheme of MFT reduction by the ethanol dehydrogenase Mdo/Mno. The proposed mycofactocin dehydrogenase is the subject of this study. X: Unknown electron acceptor.

Results

Phylogeny and co-occurrence of *mftG* and mycofactocin

Primary sequence alignments with known members of the GMC family and AlphaFold (23, 24) prediction of the tertiary structure of MftG (WP_014877070.1) from *M. smegmatis* MC^2 155 (**Figure 2A**) confirmed that MftG is a protein of the GMC superfamily with an intact FAD binding pocket (Rossmann fold) and a conserved active site histidine (25). While these properties are general characteristics of the GMC superfamily, a hallmark of the MftG subfamily of GMC oxidoreductases is their tight genetic linkage to the mycofactocin biosynthetic gene cluster *mftA-F* (4). These MFT-associated GMC homologs are defined by three protein families. The first, TIGR03970.1 (dehydrogenase, Rv0697 family) describes proteins from various actinobacteria and includes the *M. smegmatis* homolog. The other two families are TIGR04542 (GMC_mycofac_2) and NF038210.1 (GMC_mycofac_3). TIGR04542 is specific to the *Gordonia* genus, while NF038210.1 is exclusive to the *Dietzia* genus (4).

Figure 2 – Bioinformatics analysis of MftG. (A) Structural model of MftG from *M*. *smegmatis* retrieved

from the Alphafold database (26) with the FAD prosthetic group (yellow) modeled into the structure. Green:

Rossman fold motif (GxGxxG), red: active site histidine (His411). (B) Collapsed phylogenetic tree

(maximum likelihood) of GMC enzymes showing major MftG subfamilies. FastTree support values are

shown on branches. The full tree is provided as **Supplementary Figure S1** (C) Venn diagram representing

the frequency of co-occurrence of *mftC* (left-medium blue) and *mftG* (right-dark blue) genes in 312

organisms that encode the MFT gene locus or MftG-like proteins.

To further investigate the co-occurrence of *mftG* with the *mft* gene cluster we retrieved a

set of genomes encoding MftG or MftC homologs, the latter serving as a proxy for the

mft biosynthetic gene cluster (**Supplementary Table S1**). To refine the annotation of

MftGs, we first performed a phylogenetic analysis. To the MftG candidates, we added

further GMC-superfamily enzymes from the same genome set as well as sixteen

experimentally characterized GMC enzymes used in a previous study (25). Maximum

Likelihood analysis (**Figure 2B** and **Supplementary Figure S1**) clustered the MftG

candidates described by TIGR03970 into a well-supported clade, which comprised MftG

proteins from *Rhodococcus*, other *Nocardiaceae* and *Mycobacteriaceae*. The *Gordonia*

and *Dietzia* MftG candidates were also phylogenetically related to the main MftG clade.

We therefore defined all sequences belonging to this clade as MftG. Two sequences

previously annotated as MftG, however, were placed outside of the proposed MftG clade.

Their corresponding genomes (*Nocardia terpenica* and *Peterkaempfera bronchialis*) did

not contain any MftC candidate and the MftG candidates were therefore treated as

misannotations. We conclude that all MftG candidates are monophyletic. To further

improve the annotations, the gene neighborhood of the MftG candidates was

investigated, confirming that MftG candidates are frequently encoded within a 5 kb

distance from *mft* genes (**Supplementary Table S1**). After refinement of the MftG

annotation, we proceeded with co-occurrence analysis of MftC and MftG in our microbial

genome set. In a total of 312 genomes that contained either MftC or MftG, 311 encoded

an MftC homolog, and 235 encoded a putative MftG homolog. In 234 genomes the two

genes co-occurred (**Figure 2C**). This result reinforced the hypothesis that MftG enzymes

strictly require mycofactocin. On the other hand, about 25% of the genomes encoding

MftC, lack an MftG enzyme. It remains an open question for future investigations,

whether other enzymes can complement the function of MftG or whether the function of

MftG is dispensable in these organisms.

2 The role of MftG in growth and metabolism of mycobacteria

Figure 3 – Effect of *mftG* gene deletion on mycobacterial ethanol metabolism. (A) Growth curve of

M. smegmatis WT, Δ*mftG*, Δ*mftG-mftG,* and WT-*mftG* growing in HdB-Tyl with 10 g L^{-1} ethanol as the sole

carbon source. (B) Growth curve of WT and Δ*mftG* growing on HdB-Tyl with 10 g L^{-1} glucose as the sole

carbon source. (C, E) Ethanol and acetic acid quantification over time in *M. smegmatis* WT, Δ*mftG*, Δ*mftG-*

mftG, and WT-*mftG* cultures in HdB-tyloxapol with 10 g L^{-1} of ethanol and in uninoculated media as control.

(D) Growth curve of WT and Δ*mftG* on HdB-Tyl with 10 g L^{-1} glucose and 10 g L^{-1} ethanol combined. (F)

Acetaldehyde quantification in culture supernatants of the WT and Δ*mftG* strains grown on HdB-Tyl with

10 g L^{-1} glucose and/or 10 g L^{-1} ethanol. (●) *M. smegmatis* WT; (■) *M. smegmatis* Δ*mftG* mutant; (♦-

dashed) *M. smegmatis* Δ*mftG* mutant grown with starting OD_{600} 1; (▼) *M. smegmatis* Δ*mftG-mftG*

complementation mutant; (♦) *M. smegmatis* double presence of the *mftG* gene; (▲) Medium HdB-Tyl with

10 g L^{-1} of ethanol without bacterial inoculation. Measurements were performed in biological replicates,

(growth curves: n≥3, ethanol and acetate quantification: n=3). Error bars represent standard deviations.

Statistical analysis was performed with ordinary one-way ANOVA with Tukey's multiple comparison test,

p-values depicted in the figure.

To investigate the physiological role of MftG, a *mftG* deletion mutant (Δ*mftG*) was

generated in *M. smegmatis* MC^2 155. Since *mft* mutant strains typically display defects

in ethanol utilization (5), we compared the growth of *M. smegmatis* MC^2 155 Δ*mftG* with

the WT (wild-type) strain on media containing 10 g L^{-1} of ethanol as the sole carbon

source (**Figure 3A**). Indeed, almost no growth of Δ*mftG* was detected on ethanol (**Figure**

3A), whereas the growth curve of the Δ*mftG* mutant on glucose-containing media was

indistinguishable from the isogenic WT strain (**Figure 3B**).

We also investigated the growth of the WT and the Δ*mftG* strain on several related

carbon sources and recorded growth curves (**Supplementary Figure S2**). The growth

of WT and mutant strains was not supported by 10 g L^{-1} methanol, 5 g L^{-1} hexanol, 0.01

g L^{-1} acetaldehyde as sole carbon sources as previously reported (5). Notably, WT and

Δ*mftG* cells displayed significant growth with 5 g L^{-1} acetate or 10 g L^{-1} glycerol.

However, evaluation of the growth using other alcohols showed differential behavior. A

prolonged lag phase of the Δ*mftG* strain was detected on 1-propanol (29 h), 1-butanol

(18 h), and 1,3-propanediol (20 h) compared to the WT strain. Interestingly, a putative

1,3-propanediol dehydrogenase (MSMEG_6239) is present in the same operon as

MSMEG_6242, indicating that 1,3-propanediol degradation might be also MFT-

dependent (27). We further assessed the growth of ∆*mftG* using a panel of phenotype

microarrays. While no further defect regarding carbon source utilization was detected,

the ∆*mftG* mutant showed the ability to grow to a small extent on formic acid in contrast

to the WT. Besides the different usage of carbon sources, only minor differences in

bacterial growth were found in some of the sensitivity plates (**Supplementary Figure**

S3) demonstrating that the lack of the *mftG* gene did not induce relevant sensitivity

towards antibiotics and stressors compared to WT when grown on glucose alone.

The role of ∆*mftG* in ethanol metabolism of mycobacteria

Since the utilization of ethanol was the process that was most strikingly impacted by

mftG inactivation, it was investigated further. First, the complementation of the ∆*mftG*

deletion using an integrative vector carrying the *mftG* gene (∆*mftG attB*::pMCpAINT-

mftG, here named ∆*mftG-mftG*) was performed and resulted in the restoration of growth

on 10 g L^{-1} ethanol. Intriguingly, the growth curve of the complemented strain, which

could be dysregulated in *mftG* expression, displayed a shorter lag phase, a faster second

exponential growth phase, and a higher final biomass yield compared to the WT (**Figure**

3A). Altered growth kinetics of complement mutants on ethanol are a known

phenomenon, albeit not always well understood (28). However, duplication of the *mftG*

gene using the same vector in the WT strain (WT *attB*::pMCpAINT-*mftG*, here named

WT-*mftG*) also showed enhanced growth. These results indicate that MftG might

catalyze a rate-limiting step during ethanol utilization.

To further characterize the metabolic processes during ethanol consumption, the amount

of ethanol and acetic acid present in the growth media with and without mycobacterial

strains (WT, ∆*mftG*, ∆*mftG-mftG* and WT-*mftG*) was measured (**Figure 3C, E**). As

reported previously, ethanol consumption during exponential growth of WT mycobacteria

was accompanied by acetic acid production (**Figure 2E**), a process known as overflow

metabolism where the rate of ethanol oxidation to acetate exceeds the rate of carbon

assimilation for biomass formation during fast growth (8, 29). Although the Δ*mftG* strain

could achieve residual growth after 45 h, the consumption of ethanol and the production

of acetic acid remained at a very low level. (**Figure 2C**). While this finding is in line with

the occurrence of a metabolic roadblock in the Δ*mftG* mutant, it indicates, however, that

enzymatic activities supporting ethanol oxidation to acetate are not completely abolished

in Δ*mftG* mutants. This interpretation is also supported by growth curves recorded on a

combination of ethanol and glucose, where a simultaneous feeding enhanced the growth

yield of the Δ*mftG* mutant (**Figure 2D**) suggesting at least partially intact ethanol

utilization in Δ*mftG* cells.

Typically, ethanol is oxidized to acetaldehyde first, which is further oxidized to acetic

acid. To test whether the metabolic roadblock affects acetaldehyde formation or

consumption, we also quantified acetaldehyde from supernatants of WT and Δ*mftG* cells

grown on ethanol or ethanol combined with glucose **(Figure 3F)**. Interestingly, when

comparing WT and the Δ*mftG* mutant*,* we noticed that both strains produced a similar

amount of acetaldehyde regardless of whether ethanol was used as the sole carbon

source or supplemented (**Figure 3F**), indicating that acetaldehyde formation is not

altered in the mutant strain. At this point it may be helpful to revisit the fact that Δ*mftG*

mutants grew well on acetate (**Supplementary Figure S2**). This observation together

with the detection of acetaldehyde and acetate in the supernatants of Δ*mftG* cultures

excludes the hypothesis that *mftG* is required for acetaldehyde and acetate assimilation.

When investigating the carbon metabolism of the Δ*mftG-mftG* (complement) and WT-

mftG (overexpression) strains, an inverted phenotype became visible. Parallel to the

accelerated and enhanced growth described above (**Figure 3A**), the overexpression

strains displayed higher rates of ethanol consumption as well as an earlier onset of

acetate overflow metabolism and acetate consumption (**Figure 3D**). These results

indicate that MftG activity is, at least indirectly, related to ethanol oxidation. Notably, the

accelerated turnover of the volatile substrate ethanol, which is subject to substantial

evaporation during the cultivation process, to acetate, which is less volatile, could explain

the enhanced final growth yield of the complement and overexpression strains.

The impact of ethanol on survival and cell division of Δ*mftG* mutants

To investigate whether the reduced growth of Δ*mftG* mutants on ethanol is due to limited

carbon and energy supply or rather a consequence of insufficient ethanol detoxification,

we cultivated bacteria on combinations of glucose and ethanol and recorded growth

curves. Surprisingly, simultaneous feeding with 10 g L^{-1} glucose and 10 g L^{-1} ethanol

even promoted the growth of the Δ*mftG* mutant. In addition, we determined the

percentage of dead cells upon cultivation with different carbon sources using propidium

iodide staining followed by flow cytometry quantification (**Figure 4A**). Regardless of the

genotype, the bacterial cultures grown on glucose or ethanol as the sole carbon source

and under starvation displayed a similar percentage of dead cells after a cultivation

period of 72 h. We, therefore, conclude that the inability of Δ*mftG* cells to grow on ethanol

alone is not due to ethanol toxicity. This conclusion was further supported by

measurements of the transmembrane potential that did not reveal any disturbance of the

proton motif force (PMF) in the mutant (**Figure 4B**). Previous studies of the

transmembrane potential of *M. smegmatis* Δ*mftC* cells, another mutant unable to grow

on ethanol as a carbon source, showed the same pattern with no differences compared

to WT (5). However, unexpectedly, an elevated proportion of dead cells was detected

1 when ΔmftG was cultivated on combined carbon sources, while the WT tolerated this

2 condition well. This finding emphasizes the role of the mycofactocin system in enhancing

3 the metabolic adaptability of mycobacterial cells under increasingly complex

4 environmental conditions.

5 To reveal potential morphological changes induced by ethanol treatment, we further

6 examined the ΔmftG mutant incubated with different carbon sources by super-resolution

7 microscopy. The identification of elongation and replication sites was achieved by the

8 sequential incubation of the cells with the fluorescent D-amino acid labeling probes NADA

9 (green) and RADA (red) (**Figure 4C-E).** These probes are incorporated via extracellular

10 routes of mycobacterial peptidoglycan assembly, labeling both elongation poles as well

11 as the sidewall (30). The *M. smegmatis* WT and ΔmftG-mftG cultivated on glucose as

12 the sole carbon source appear elongated with mainly polar growth (accumulation of

13 RADA at the poles indicates the most recent active site of peptidoglycan incorporation).

14 NADA occurs mainly at a non-polar location (previous active replication site). The ΔmftG

15 mutant grown on glucose, however, showed few yellow cells (overlap of green and red)

16 indicating growth arrest on a small part of the culture. Some cells are not actively

17 synthesizing peptidoglycan and thus not growing, which leads to the overlap of both dyes

18 (**Figure 4C**).

19 Mycobacteria grown on glucose and ethanol showed a regular dispersion profile of both

20 dyes, with polar RADA, non-polar NADA, and occasional mid-cell septum formation with

21 no obvious differences between the three genotypes. In contrast, cultivation on 10 g L^{-1}

22 ethanol alone as well as under starvation condition (no carbon source) had a strong

23 influence on peptidoglycan synthesis as the dyes show decreased incorporation into the

24 cell wall. The simultaneous incorporation of both dyes in WT and ΔmftG-mftG cells grown

25 on ethanol as the sole carbon source reflected the reduced growth rate, as shown by the

growth curves (**Figure 3A, B**) compared to the glucose condition. Interestingly, Δ*mftG*

mutant cells struggled to divide on ethanol displaying multiple septa and extreme

elongation compared to WT on ethanol, leading to a higher ratio of foci per cell compared

to WT grown on the same condition (**Figure 4D, E**). Independent of the genotype, the

cultivation in the presence of ethanol significantly decreased cell size. A similar

phenotype was previously observed in *Mycobacterium vaccae* cells exposed to ethanol

(31). Starving cells were also significantly reduced in size in WT and Δ*mftG* strains

compared to growth on glucose alone (**Figure 4C, E)**, reflecting the limitation of

resources available for propagation and elongation. Especially under starvation

conditions, the cells appeared predominantly yellow, suggestive of growth arrest. A

previous study on the effect of starvation on mycobacteria also showed the formation of

large cells with multiple septa as a hallmark of starvation, from which after 14 days, mildly

starved cells (with traces of carbon source available) remodeled into small resting cells

(32). Our results indicated that the major phenotype of Δ*mftG* on ethanol as a sole source

of carbon is a growth defect comparable to a starvation effect. When incubated solely

with this carbon source, Δ*mftG* cells are almost unable to divide, but remain alive and in

a metabolically active state, most likely consuming ethanol in trace amounts.

Figure 4 – Phenotypic characterization of mycobacterial strains grown on HdB-Tyl with glucose

and/or ethanol or starvation. (A) Quantification of dead cells by flow cytometry using propidium iodide of

M. smegmatis strains grown throughout 72 h. Biological replicates, starvation cells n=2, other conditions

n=3. (B) Quantification of cells with normal transmembrane potential by flow cytometry of the *M. smegmatis*

cultures throughout 72 h. Biological replicates n=3. (C) Super-resolution microscopy images of *M.*

smegmatis strains at exponential phase or 48h of starvation, labeled with NADA (green), RADA (red),

superposition of NADA and RADA (yellow). Bar size: 3 µm. (D) Cell size distribution obtained from super-

resolution microscopy of the *M. smegmatis* strains at exponential phase or 48h of starvation. (E) Ratio of

the number of replication sites to the number of cells of the *M. smegmatis* strains cultures at exponential

phase or 48h of starvation, together with a microscopy image of a single Δ*mftG* cell at 48h grown on

ethanol, with arrows pointing to the several septa stained with NADA (green) and RADA (red). Bar size: 3

μm.

Color legend: (A,B): ● – sample at 24h; ■– sample at 48h; ▲ – 72h. (C) (D,E): orange – 10 g L^{-1} glucose;

green – 10 g L^{-1} glucose and 10 g L^{-1} ethanol; blue – 10 g L^{-1} ethanol; white – starvation for 48h. Statistical

analysis was performed for PI, cell size and ratio of replication sites per cell with ordinary one-way ANOVA,

for transmembrane potential with ordinary two-way ANOVA, all using Tukey's multiple comparison test.

The p-values are depicted on the figure, microscopy-based analysis performed with technical replicates

(n=3).

10 Impact of *mftG* on mycobacterial cofactor metabolism

Figure 5 – Cofactor metabolism of *M. smegmatis* strains. (A) NADH/NAD$^+$ ratio of *M. smegmatis* WT

and Δ*mftG* grown on HdB-Tyl with either 10 g L^{-1} glucose or 10 g L^{-1} ethanol at exponential phase. (B)

NADH and NAD$^+$ quantification of *M. smegmatis* WT and Δ*mftG* grown on HdB-Tyl with either 10 g L^{-1}

glucose or 10 g L^{-1} ethanol at exponential phase. (C) ADP/ATP ratio of *M. smegmatis* WT and Δ*mftG*

grown on HdB-Tyl with 10 g L^{-1} ethanol at 24 h, 48 h and 72 h. (D, E, F) Targeted comparative metabo-

lomics of *M. smegmatis* WT, Δ*mftG*, Δ*mftG-mftG,* and WT-*mftG* strains. The most representative MFT

species, methylmycofactocinone with 8 glucose moieties (MMFT-8H$_2$, sum formula: C$_{62}$H$_{99}$NO$_{43}$, RT: 6.82

min, *m/z* 1546.5665 [M+H]$^+$) and methylmycofactocinol with 8 glucose moieties (MMFT-8, sum formula:

C$_{62}$H$_{97}$NO$_{43}$, RT: 7.18 min, *m/z* 1544.5507 [M+H]$^+$), was used to reflect MFT obtained from *M. smegmatis*

strains. The bacteria were grown in HdB-Tyl with either (D) 10 g L^{-1} glucose, (E) 10 g L^{-1} ethanol, or (F) 10

g L^{-1} glucose combined with 20 g L^{-1} ethanol. Samples of the different growth phases are represented in

the chart. A sampling at 60h of Δ*mftG* was chosen to sample the residual growth of the strain on ethanol

as the sole carbon source. Statistical analysis was performed with one- or two-way ANOVA with Dunnett's

multiple comparison test for NADH/NAD$^+$ ratio and, Tukey's test for the rest, with most relevant p-values

depicted on the figure. Measurements were performed in biological replicates (n=3).

Based on the results above, we concluded that MftG is crucial for proper ethanol

metabolism in mycobacteria. However, deletion of its corresponding gene still allowed

for basal metabolic activity and ethanol oxidation. One way to explain these findings is

that MftG is involved in cofactor regeneration during growth on ethanol as a carbon

source. The genetic linkage of *mftG* with mycofactocin biosynthesis supported the

hypothesis that MftG might be responsible for mycofactocin regeneration. However,

before directly addressing mycofactocin metabolism, we decided to monitor the central

respiratory redox cofactor NAD. We therefore determined the NADH/NAD$^+$ ratios of WT

and KO during the exponential phase when bacteria were grown on glucose or ethanol

as the sole carbon sources. However, the presence or absence of *mftG* did not

significantly influence the NADH/NAD$^+$ ratio of bacteria grown on either carbon source

(**Figure 5A**). This finding indicated that NAD homeostasis remained intact in the Δ*mftG*

mutant. Notably, when looking at absolute NAD$^+$ levels, (**Figure 5B**) an increase of the

NAD pool in cells growing on ethanol was detected, however, independent of the genotype. Along with central redox (NAD$^+$) metabolism, we also monitored the central energy metabolism of the cells by determining ADP/ATP ratios. These clearly supported the hypothesis that $\Delta mftG$ cells suffered from starvation. The mutant strain exhibited a significant energy deficit at all three sampled time points, with the energy depletion progressively worsening between 24 and 48 hours of incubation (**Figure 5C)**. Previous studies of *M. smegmatis* under starvation also showed a reduction of ATP levels as a result of hampered energy metabolism (32). It should be mentioned that ATP synthesis strictly depends on respiration in *M. smegmatis*. Mycobacteria typically cannot sustain growth, even on fermentable substrates, via substrate-level phosphorylation alone (33).

After confirming energy shortage but intact NAD homeostasis in $\Delta mftG$ cells, we decided to test whether MftG is involved in mycofactocin regeneration. To this end, we directly analyzed the MFT pool using targeted liquid chromatography-high resolution mass spectrometry (LC-MS). We have previously shown that the total pool size of MFT species is highly expanded when *M. smegmatis* uses ethanol as a sole carbon source and that both reduced (M/MFT-nH_2) as well as oxidized species (M/MFT-n) are stable when extracted and can thus be detected by LC-MS (8). Here, we performed a comparative analysis of the metabolome of *M. smegmatis* WT and $\Delta mftG$ grown in HdB-Tyl supplemented with either 10 g L^{-1} glucose, 10 g L^{-1} ethanol, or both carbon sources combined (10 g L^{-1} glucose and 20 g L^{-1} ethanol). Metabolites were extracted during the exponential or stationary growth phase. Additionally, metabolome extracts from complemented $\Delta mftG$-*mftG* and WT-*mftG* strains grown on HdB-Tyl with 10 g L^{-1} ethanol sampled only at the stationary phase were analyzed. Strikingly, this analysis revealed significantly elevated levels of MMFT-$8H_2$ in $\Delta mftG$ compared to the other strains tested (**Figure 5D-F**). MMFT-$8H_2$ is the major representative of reduced mycofactocins

(mycofactocinols) in *M. smegmatis*. Interestingly, Δ*mftG* strains contained almost none of the corresponding oxidized mycofactocinone (MMFT-8) in any of the conditions tested. These results strongly supported the hypothesis that MFT regeneration was hampered in the Δ*mftG* mutant while MFT reduction was still taking place, thus resulting in a near-total conversion of mycofactocinones to mycofactocinoles. This phenotype was also observed when Δ*mftG* cells were cultivated on glucose and ethanol combined (**Figure 5E**) and even on glucose alone (**Figure 5D**). Complementation of the Δ*mftG* mutant (Δ*mftG-mftG*) reverted the MMFTH$_2$ to MMFT ratio back to WT levels. In contrast, the overexpression strain WT-*mftG* grown in ethanol contained significantly more MMFT-8 (oxidized) compared to other strains in the same condition (**Figure 5F**), thus providing further evidence that MftG is involved in reoxidation of mycofactocin and re-enforcing that this step might even represent a rate-limiting step during ethanol utilization.

MFT dehydrogenase assay with recombinant MftG

To further support the hypothesis that MftG is responsible for MFT reoxidation, purification of MftG as a recombinant fusion protein with hexahistidine tag followed by *in-vitro* enzyme assays was attempted. For homologous production of recombinant, hexahistidine-tagged MftG, *M. smegmatis* Δ*mftG* was chosen as an expression host (resulting genotype: Δ*mftG-mftG*His$_6$). The restoration of the growth on ethanol of the Δ*mftG-mftG*His$_6$ expression strain indicated that the recombinant protein was produced in an active form. Semi-purified cell-free extracts of Δ*mftG-mftG*His$_6$ incubated with metabolome extract from *M. smegmatis* Δ*mftG* as substrate showed that the *mftG-mftG*His$_6$ lysate significantly increased the amount of MMFT-8 and decreased the amount of the MMFT-8H$_2$ while the Δ*mftG* lysate did not affect the ratio of the two compounds (**Figure 6A**). Inspired by these results we conducted assays with semi-

purified cell-free extracts from *M. smegmatis* Δ*mftG-mftG*His$_6$ with higher amounts of purified cofactors as substrates. We added synthetic PMFTH$_2$ and purified MMFT-2H$_2$ as substrates as well as synthetic PMFT as a negative control. The negative control using PMFT contained trace amounts of PMFTH$_2$ from the start and no clear trend was observed over the time course of the reaction (**Figure 6B**). In contrast to this, we observed conversion of both reduced substrates PMFTH$_2$ (**Figure 6C**) and MMFTH-2H$_2$ (**Figure 6D**) to the corresponding oxidized species PMFT and MMFT-2, respectively. However, the reaction proceeded only with partial conversion of the substrate. Additionally, since some enzymes of the GMC family can utilize oxygen as electron acceptors, the role of oxygen in MftG activity was accessed. To this end, the assays were performed under a nitrogen atmosphere at < 0.1% oxygen. Under these conditions the MftG activity remained unchanged (**Figure 6C, D**), thus ruling out that MftG acts as a mycofactocin oxidase using O$_2$ as an electron acceptor.

Figure 6 – MftG assays with recombinant enzymes and MFTs as substrates. (A) Mycofactocinol oxi-

dation assay with semi-purified cell-free extract of *M. smegmatis* Δ*mftG* or Δ*mftG-mftG*His$_6$ using Δ*mftG*

metabolome extract as substrate (naturally enriched in reduced MFT as described in Figure 5F). Result

showing the oxidation of MMFT-8H$_2$ to MMFT-8 after overnight incubation (ON) when semi-purified cell-

free extract from Δ*mftG-mftG*His$_6$ is used. t_0 – start of the assay. (B) Assay with semi-purified cell-free

extract of Δ*mftG-mftG*His$_6$ using synthetic PMFT as a control substrate showed no relevant reaction. (C)

Successful oxidation of PMFTH$_2$ when synthetic PMFTH$_2$ was used as substrate (C$_{13}$H$_{17}$NO$_3$, RT: 7.84

min, *m/z* 236.1281 [M+H]$^+$) to PMFT (C$_{13}$H$_{15}$NO$_3$, RT: 8.40 min, *m/z* 234.1125 [M+H]$^+$). Black and grey

lines depict reactions performed in an anaerobic chamber. (D) Successful oxidation of MMFT-2H$_2$

(C$_{26}$H$_{39}$NO$_{13}$, RT: 7.10 min, *m/z*: 574.2494 [M+H]$^+$) to MMFT-2 (C$_{26}$H$_{37}$NO$_{13}$, RT: 7.47 min, *m/z* 572.2338

[M+H]$^+$). Black and grey lines depict assays performed in the anaerobic chamber (E) Mycofactocinol

(MMFT-2H$_2$) oxidation using MftG heterologously produced in *E. coli* and DCPIP, NAD$^+$, NDMA, and PMS

as potential electron acceptors. Control – no MftG added; Enzyme – MftG added. (F) Dose-dependent

effect of heterologously expressed MftG on the oxidation of MMFT-2H$_2$ to MMFT-2 was observed after a

24-hour incubation period. Sample size of all experiments n=3. Error bars represent standard deviations.

In order to confirm that the activity detected in cell-free lysates was not due to

background effects, the heterologous production of MftG in *Escherichia coli* followed by

activity assays was attempted. Despite poor production and solubility of MftG in *E. coli*,

we detected MftG activity in semi-purified fractions of *E. coli* producing MftG tagged with

maltose-binding-protein (pPG36). While mycofactocin dehydrogenase activity was

clearly linked to the overexpression of MftG, only partial turnover of substrate was still

observed suggesting the absence of an appropriate electron acceptor. A screening for

several potential electron acceptors was performed using MMFT-2H$_2$ as substrate.

However, none of the potential acceptors tested, DCPIP, NAD$^+$, NDMA and PMS, were

able to increase substrate turnover (**Figure 6E**). Assays with increasing concentrations

of protein suggested that the observed activity was due to single-turnover reactions,

most likely using the FAD cofactor in the active site as an electron acceptor (**Figure 6F**).

We, therefore, concluded that MftG can indeed interact with mycofactocins as electron

donors but might require complex electron acceptors, for instance, proteins present in

the respiratory chain.

Influence of MftG and MFT in mycobacterial respiration

The phenotypes observed for cell growth and division together with an increased ratio of ADP/ATP in the $\Delta mftG$ strain suggested a potential involvement of MftG in mycobacterial respiration. To further test this hypothesis, the consumption of O_2 from whole cells of *M. smegmatis* WT and $\Delta mftG$ grown on HdB-Tyl with 10 g L^{-1} of ethanol was measured in a respirometer equipped with a Clark-type electrode. This experiment revealed that the respiration rate of $\Delta mftG$ reached only about 45% of the WT level (**Figure 7**). Notably, the *M. smegmatis* carrying a duplicated *mftG* gene, WT-*mftG*, showed accelerated oxygen consumption, supporting the idea that MftG is a limiting factor in the respiration of mycobacteria grown on ethanol.

Figure 7 – Respiratory activity (oxygen consumption) of *M. smegmatis* WT and Δ*mftG* mutants. (A)

Respiration of intact cells. Average of n=3 (B) Respiration of WT isolated cell membranes and addition of

electron donors as indicated in the figure. (C) Respiration of Δ*mftG* isolated cell membranes and addition

of electron donors as indicated in the figure. NADH and succinate served as positive controls, and water

as a negative control. KCN treatment served as inhibitor control. MMFT-2H$_2$ and PMFTH$_2$ were added to

confirm MFT's role as an electron donor. (D) Oxidation of PMFTH$_2$ to PMFT in WT isolated membranes

(combined LC-MS profiles). (E) Oxidation of PMFTH$_2$ to PMFT of the Δ$mftG$ isolated membranes

(combined LC-MS profiles). Each inset depicts the profile after KCN treatment. Representative data was

selected from independent experiments n≥3.

To test whether MftG might interact with components of the respiratory chain, we

investigated mycobacterial membrane preparations for respiratory activities. The

influence of mycofactocin on mycobacterial respiration was tested using isolated

membranes from *M. smegmatis* WT supplemented with either NADH, succinate,

synthetic PMFTH$_2$, or purified MMFT-2H$_2$. In these respiration assays, the consumption

of oxygen after supplementation with NADH or succinate was highly increased showing

the expected respiratory activity of the isolated membranes via NADH or succinate

dehydrogenases (**Figure 7**). The addition of PMFTH$_2$ or MMFT-2H$_2$ stimulated oxygen

consumption to a rate that was comparable to succinate-induced respiration of the WT

strain. In *M. smegmatis*, cyanide is a known inhibitor of the cytochrome *bc*/*aa3* but not

of cytochrome *bd* (34), therefore, the decrease of oxygen consumption when MFTs were

added to the membrane fractions in combination with KCN (**Figure 7**), revealed that

MFT-induced oxygen consumption is indeed linked to mycobacterial respiration.

The membrane fraction of the Δ$mftG$ strain showed a decreased consumption of oxygen

compared to WT. Although the auto-respiration, i.e., the consumption of oxygen in the

absence of a stimulant, was lower in the Δ$mftG$ strain, the supplementation of NADH and

succinate stimulated oxygen consumption demonstrating that the membranes contained

a functional electron transport chain. The addition of reduced mycofactocins to the

reaction revealed strongly reduced respiration when compared to the WT membrane. To

test whether increased O$_2$ consumption was linked to the oxidation of mycofactocinols,

we also we carried out LC-MS analysis of the samples used in the respiration assays.

Indeed, mycofactocinols were oxidized to mycofactocinols (e.g. PMFTH$_2$ to PMFT) under

active assay conditions, while only low amounts of PMFTH$_2$ were oxidized in membranes of the $\Delta mftG$ strain or by membranes treated with KCN. Two observations suggest a low level of MftG-independent mycofactocin oxidation: (I) The residual induction of oxygen consumption in the $\Delta mftG$ strain by mycofactocinols, (II) the formation of low amounts of oxidized mycofactocins in the $\Delta mftG$ strain (at a similar level to WT inhibited by KCN). From the respirometry experiments, we conclude that MftG catalyzes the electron transfer from mycofactocinol to membrane-bound components of the respiratory chain. These could be the quinone pool as known from succinate dehydrogenase complex, or cytochromes, for instance. For example, MftG in mycobacteria might play a similar role as the subunit II of the pyrroquinoline quinone-dependent alcohol dehydrogenase (PQQ-ADH) complex from acetic acid bacteria. The subunit II transfers electrons from the alcohol dehydrogenase (subunit I/III) to the ubiquinone pool, from which further electron transfer to oxygen occurs (35).

Transcriptomic analysis of *M. smegmatis* $\Delta mftG$ cells

To monitor genome-wide regulatory adaptation caused by *mftG* deletion, we employed high-throughput sequencing to record transcriptomic changes in WT and $\Delta mftG$ cells when grown to exponential phase (60 h of incubation for $\Delta mftG$ on ethanol) on glucose or ethanol. In the glucose condition, only nine out of 6.506 annotated genes classified significantly upregulated in $\Delta mftG$ when compared to the WT (adjusted p-value <0.05 and log2FoldChange > 2). Since these genes showed neither functional nor spatial clustering, we conclude that no meaningful differences in the mycobacterial transcriptome between WT and mutant grown on glucose were observed (**Table S2).**

On the contrary, transcriptomics analysis of $\Delta mftG$ and WT grown on ethanol detected 578 genes as significantly up-regulated and 697 as down-regulated (adjusted p-

value<0.05 and log2FoldChange > 2 or <-2) in the mutant when compared to the WT (**Table S2**). The deletion of the *mftG* gene induced mainly processes related to transmembrane transporter activity, iron binding, monooxygenase activity, quinone activity and NADH dehydrogenase (**Figure 8A**). Downregulated processes were related to ATP binding and ATPase transmembrane movement, respiration, and DNA replication (**Figure 8B**). Significant changes, presumably due to energy limitation, took place in the Δ*mftG* mutant compared to the WT. For instance, the *nuo* genes (MSMEG_2050 - 2063) encoding NADH dehydrogenase I (NADH-I), were amongst the most highly expressed genes (**Figure 8, Table S2**). In mycobacteria, two NADH dehydrogenases (NDH-I and II) are active, while in *M. smegmatis* the NADH-I dehydrogenase typically contributes only a low proportion of total NADH activity (36). Notably, enhanced expression of the high-efficiency, proton-pumping NADH-I (*nou*) system was previously observed under energy-limited starvation conditions (37). In addition, alternative dehydrogenases fostering electron transfer from other electron donors to the respiratory chain were strongly upregulated (**Figure 8C**). These comprise enzymes like proline dehydrogenase (MSMEG_5117), pyruvate dehydrogenase (MSMEG_4710 – 4712), carbon monoxide oxidase (MSMEG_0744 – 0749), the hydrogenases 1 (MSMEG_2262 – 2276) and 2 (MSEMG_2720-2719). Furthermore, quinone-dependent succinate dehydrogenase SDH1 (MSMEG_0416-0420) was highly upregulated when compared to the WT. Again, similar patterns were previously described in cultures growing under carbon limitation and can be seen as an attempt of the cell to compensate for poor loading of the electron transport chain (37). This finding is in line with our hypothesis that electron transfer to the respiratory chain is hampered and mutant cells suffer from mild starvation.

Figure 8 – Representation of the main metabolic activities affected by *mftG* deletion in *M.*

***smegmatis* grown on 10 g L⁻¹ ethanol as the sole carbon source compared to WT.** (A,B) Functional

annotation chart (Gene ontology enrichment analysis) of the (A) up- and (B) down-regulated processes.

Gene ratio denotes the ratio of the involved genes (count) to the quantity of the genes making up the

enriched terms. (C) Impact on the respiration of the mutant strain $\Delta mftG$ grown on 10 g L⁻¹ ethanol

compared to WT strain. Blue represents genes downregulated $p<0.05$ and log2FC< -2. Red represents

genes upregulated $p<0.05$ and log2FC> 2. Protein figures retrieved from public databases NADH-II:

A0QYD6, NADH-I: 8e9g, Cyt bd: 7d5i, Cyt bcc-aa3: 7rh5, SDH-II: 6LUM, Sdh-I: 7d6x, ATP synthase:

7NJK.

Reflecting the division arrest phenotype observed by super-resolution microscopy,

peptidoglycan biosynthesis, represented by the genes *murABCDEFG* (e.g.

MSMEG_1661, MSMEG_6276, MSMEG_2396, MSMEG_4194) was found to be

downregulated. DNA replication was also affected in $\Delta mftG$ grown on ethanol by the

downregulation of RNaseH (MSMEG_4306), DNA polymerase (MSMEG_3839),

helicase (MSMEG_6892), and all the modules of the DNA polymerase III holoenzyme

(MSMEG_4259, MSMEG_3178, MSMEG_6285, MSMEG_6153, MSMEG_4572 and

dnaN).

In general, the transcriptomic analysis of the Δ*mftG* strain on ethanol compared to WT

in the same condition showed that the observed phenotype resembled energy-limiting

conditions, accompanied by remodeling of the respiration to compensate for a lack of

redox equivalents as well as inhibition of peptidoglycan synthesis and DNA replication.

Discussion

Alcohols are widespread ingredients of various foods and beverages and are used as

active principles of many disinfectants. Therefore, the impact of alcohols on commensal

and clinically important bacterial strains have high relevance. However, ethanol

metabolism in mycobacteria is still poorly characterized. While it is established that the

cofactor mycofactocin plays a major role in this process, the fate of electrons during

alcohol oxidation remained unknown (5, 12).

Based on bioinformatics analysis we hypothesized that MftG homologs, flavoenzymes

of the GMC superfamily, might be involved in redox metabolism connected to MFT-

dependent alcohol metabolism. Using *M. smegmatis* as a model system we showed that

MftG is indeed essential for ethanol metabolism, a process known to depend on both an

intact mycofactocin biosynthetic pathway as well as the putative MFT-dependent

alcohol-dehydrogenase Mdo/Mno. Mutants lacking the *mftG* gene incubated on ethanol

as the sole carbon source displayed strongly impaired growth. Super-resolution

microscopy combined with peptidoglycan labeling, transcriptomics as well as direct

measurements of ADP/ATP ratios demonstrated that mutants grown on ethanol suffer

from energy limitation and growth arrest similar to a mild starvation condition. However,

the mutant maintained a level of functional metabolism, conserving its transmembrane

potential. The ethanol oxidation pathway was proven to be affected in the KO strain but not fully interrupted as suggested by partially intact co-metabolism with glucose and production of intermediate products.

Investigation of cofactor metabolism revealed that deletion of *mftG* resulted in a drastic increase in the ratio of reduced/oxidized mycofactocins in the mycobacterial cell while the $NADH/NAD^+$ ratio remained unchanged. This finding prompted the hypothesis that MftG could be involved in MFT cofactor regeneration. Assays with homologously and heterologously produced, semi-purified enzymes showed that MftG indeed oxidized $PMFTH_2$ and $MMFT-2H_2$, but only single-turnover reactions could be observed. Respirometry studies further demonstrated that reduced mycofactocins feed redox equivalents into the intact mycobacterial respiratory chain. Strikingly, this process was severely hampered in $\Delta mftG$ mutants, suggesting that MftG, while directly accepting electrons from mycofactocinol, delivers electrons either directly or indirectly into the respiratory chain. Transcriptomic analysis revealed a profound re-organization of the respiration machinery when MftG was absent suggestive of starvation and low availability of redox equivalents to the respiratory chain.

We, therefore, conclude that MftG is a mycofactocin dehydrogenase requiring electron acceptors associated with the membrane-bound respiratory chain. This assumption is plausible considering that other members of the GMC superfamily, e.g., choline dehydrogenase from *E. coli*, were reported to couple substrate oxidation with the electron transport chain (22). While the exact electron acceptor remains a subject for future studies, it is plausible that cytochromes might fulfill this function. Interestingly, cellobiose dehydrogenase, a fungal GMC enzyme, is a multidomain enzyme where the dehydrogenase domain (GMC) is linked to a cytochrome domain. (38). This situation is reminiscent of pyrroloquinoline quinone (PQQ)-dependent alcohol dehydrogenases (39).

Those enzymes, that mainly occur in the periplasm of Gram-negative bacteria, are typically coupled to a dedicated cytochrome c subunit, which is genetically clustered along with the other subunits (39). It is also likely that the membrane-associated quinone pool serves as the electron acceptor. For instance, the GMC-family oxidoreductase enzymes PhcC and PhcD from *Sphingobium* sp. involved in lignin oxidation were suggested to transfer electrons to the ubiquinone pool (40). Reduction of menaquinone ($E_0' = -74$ mV), the dominant respiratory quinone of *Mycobacteria*, by mycofactocinol would be thermodynamically favorable considering the midpoint potential of premycofactocins (PMFT/PMFTH$_2$: $E_0' = -255$ mV) (16). However, other redox coenzymes, redox proteins or protein complexes of the respiratory chain might act as direct reaction partners of MftG.

Our study sets the ground for future investigations into the connection of the mycofactocin system to further (known or unknown redox) systems. For instance, background MFTH$_2$ oxidation by membrane preparations of the Δ*mftG* mutant points towards the existence of alternative MFT-dependent dehydrogenases or oxidases. Our results provide insights that are important for mechanistic studies of previous effects shown to be related to MftG. For instance, previous studies reported growth defects connected to *mftG* inactivation by transposon mutagenesis in *M. tuberculosis* (41). More recently, the transcriptome analysis of a dose-dependent response of *M. tuberculosis* to the succinate dehydrogenase inhibitor BB2-50F showed that this new antibiotic induced upregulation of *mftG* (42). Furthermore, vulnerability studies of *M. smegmatis*, *M tuberculosis* H37Rv and the hypervirulent strain HN878 using a CRISPRi library revealed that, although disruption of *mftG* did not induce any phenotype in *M. smegmatis*, still the disturbance of this gene induced growth defects in both *M. tuberculosis* strains (43). Moreover, disruption of the *mftG* gene by this technique showed increased fitness of *M.*

tuberculosis H37Rv when exposed to various drugs, e.g., bedaquiline and ethambutol,

while reducing fitness when exposed to isoniazid and vancomycin (43). Having these

results in mind, further investigations of the role of MftG in *M. tuberculosis* survival and

infection might disclose important implications of the MFT system for TB treatment.

Taken together, we close an important knowledge gap by showing that the flavoprotein

MftG plays a role in MFT regeneration and acts as a redox shuttle between alcohol

substrates and the respiratory chain. With the clarification of the MftG function, the entire

set of *mft* genes are now elucidated in mycobacteria. Thereby, this study adds another

important element to the amazingly flexible network of mycobacterial redox metabolism

and opens up an avenue for future investigations of the mycofactocin system.

Material and methods

Bacterial strains and culture media

The *Mycolicibacterium smegmatis* MC^2 155 strain was used in this study. All generated

mycobacterial strains were maintained at 37 °C in lysogeny broth (LB) supplemented

with 0.5 g L^{-1} Tween 80 (LBT). *Escherichia coli* TOP10 (Thermo Fischer Scientific) was

used to propagate plasmids on LB medium supplemented with either 100 µg mL^{-1}

hygromycin B (Carl Roth) or 50 µg mL^{-1} kanamycin (Carl Roth). *E. coli* NiCo21(DE3)

(NEB BioLabs) was used to express recombinant proteins. Genomic manipulation of *M.*

smegmatis was performed in LBT supplemented with 60 µg mL^{-1} hygromycin B and 10

g L^{-1} sucrose for double crossover event or 50 µg mL^{-1} kanamycin for *gfp-hyg* cassette

removal and complementation/overexpression strain generation. Studies for comparison

between WT and mutants were performed with adapted Hartmans de Bont medium (17)

(HdB-Tyl) containing (NH$_4$)$_2$SO$_4$ 2 g L^{-1}, MgCl$_2$·6H$_2$O 0.1 g L^{-1}, Na$_2$HPO$_4$ anhydrous 3 g

L^{-1}, and KH_2PO_4 anhydrous 1.07 g L^{-1} supplemented with 0.10% [v/v] 1000 × trace element solution (ethylenediaminetetraacetic acid 10.0 g L^{-1}, $CaCl_2 \cdot 2H_2O$ 1.0 g L^{-1}, $Na_2MoO_4 \cdot 2H_2O$ 0.2 g L^{-1}, $CoCl_2 \cdot 6H_2O$ 0.4 g L^{-1}, $MnCl_2 \cdot 2H_2O$ 1.0 g L^{-1}, $ZnSO_4 \cdot 7H_2O$ 2.0 g L^{-1}, $FeSO_4 \cdot 7H_2O$ 5.0 g L^{-1}, and $CuSO_4 \cdot 5H_2O$ 0.2 g L^{-1}) and 0.5 g L^{-1} tyloxapol (Sigma Aldrich) and supplemented with the appropriate carbon source: 10 g L^{-1} absolute ethanol (TH.Geyer), 10 g L^{-1} glucose (Carl Roth), 1% [v/v] glycerol (Carl Roth), 5 g L^{-1} acetate (Carl Roth), 10 g L^{-1} methanol (Carl Roth), 10 g L^{-1} 1-propanol (Sigma-Aldrich), 5 g L^{-1} 1-butanol (Sigma-Aldrich), 5 g L^{-1} hexanol (Sigma-Aldrich), 0.010 g L^{-1} acetaldehyde, 10 g L^{-1} propane-1,2-diol (Sigma-Aldrich) and 10 g L^{-1} propane-1,3-diol (Sigma-Aldrich).

Bioinformatics analysis of MftG, co-occurrence and phylogenetic analysis

The AlphaFold (24) structure prediction of MftG from *M. smegmatis* MC^2 155 (model AF-I7F8I2-F1) was retrieved from the AlphaFold Protein Structure Database (23). In order to investigate the FAD binding pocket, the closest homolog of MftG with a solved crystal structure, HMFO oxidase (PDB: 4udp) (44) was retrieved from the Protein Data Bank (PDB). The structure of HMFO oxidase was superimposed with the AlphaFold model of MftG using the matchmaker feature in ChimeraX version 1.2.5 (45). The FAD molecule present in the HMFO oxidase crystal structure was used to complement the MftG model. The characteristic Rossman fold (GxGxxG) and histidine active site were identified through the alignment feature (MUSCLE) in Geneious Prime version 2022.2.2. comparing the aforementioned *M. smegmatis* protein sequence with examples of different activity GMCs previously described (25). The regions were highlighted in the predicted structure.

The co-occurrence of *mftG* and *mftC* was analyzed similarly as described before (17). Since *mftC* is essential for the biosynthesis of mycofactocin, the presence of an *mftC*

gene served as a proxy for the presence of the *mft* gene cluster. To obtain genomes

encoding either an *mftC* or *mftG* homolog, the tables for the Hidden Markov Model

(HMM) hits in the NCBI RefSeq protein database were retrieved from the respective

NCBI protein family models' entries. The NCBI HMM accession numbers were

TIGR03962.1 for *mftC* and TIGR03970.1, TIGR04542.1. and NF038210.1 for *mftG*. The

MftG family was described by three distinct HMMs at the time of writing, with

TIGR03970.1 as the main family comprising a variety of genera while the other two MftG

models TIGR04542.1 and NF038210.1 describe homologs restricted to the genera

Gordonia and *Dietzia*, respectively. The entries were filtered for origin in complete

genomes and each entry was checked manually for the presence of potentially missed

mftC or *mftG* homologs.

The search for other GMC proteins that could be present in the genomes of strains that

contain the mycofactocin cluster (but no MftG) was performed by collecting a list of pu-

tative homologs of *M*. smegmatis MC^2 155 Uniprot accession number A0QSC2 through

BLASTP search of the non-redundant protein database from NCBI with the NCBI tool

search with the following settings: max target sequences 5000, expect threshold 0.0002,

word size 6, BLOSUM62, gap cost exist 11, gap cost extend 1, conditional compositional

score matrix adjustment and filter low complexity regions. Partial sequences were not

included in the analysis. The protein sequences of all GMC proteins obtained, as well as

16 previously published reference GMC enzymes, were used for phylogenetic analysis

and are listed in **Supplementary Table S1**. The sequences were aligned using MAFFT

(46) and the phylogenetic tree was built using the maximum likelihood methods imple-

mented in FastTree 2.1.11 (47) with WAG substitution model (both programs imple-

mented as Geneious plugins). The tree was further edited using TreeViewer Version 2.1

(48) to condense phylogenetic clades and full tree was colored using FigTree v1.4.4.

Plasmids and *M. smegmatis* genetic manipulation

The *M. smegmatis* MC^2 155 mutants were produced using the pML2424/pML2714 (49) system combined with the pMCpAINT complementation vector (5). The plasmid for *mftG* removal (pPG17) resulted from the combination of the vector pML2424 with SpeI/SwaI and PacI/NsiI ligation with the amplified upstream and downstream regions of *mftG* using primers described in Table 1, flanking the *gfp-hyg* cassette. Preparation of mycobacterial competent cells and transformation was conducted as previously described (17). The confirmation of the mutant (Δ*mftG*) was achieved with PCR amplification of external and internal primers (Table 1) after the removal of the selection cassette using Cre recombinase from pML2714. The Δ*mftG* was made competent for complementation (Δ*mftG-mftG*) and overexpression (WT-*mftG*) of *mftG* by genome integration using plasmid pPG29, which originated from pMCpAINT as a backbone by the addition of a NdeI restriction site using the primers listed in Table 1. The linearized plasmid was cut with NdeI and BamHI for the introduction of the native *M. smegmatis* MC^2 155 *mftG* gene (synthesized by BioCat). To obtain the overexpression strain (WT-*mftG*), the WT strain was transformed with pPG29. The plasmid pPG32 was obtained by PCR using pPG29 as a template and appropriate primers to add the hexa-histidine tag at the C-terminus (Table 1). It was subsequently used to transform the Δ*mftG* strain and used for *mftG*his$_6$ expression (Δ*mftG-mftG*His$_6$). The plasmid pPG23 was constructed using pPG20 as a backbone, introducing at the NcoI/HindIII cutting sites the PCR-amplified mycofactocin operon *mftA-F* (primers listed in Table 1) and used to generate the strain WT-*mftABCDEF* with the integration of the plasmid in the WT strain. The PCR amplifications were performed using Q5 High-Fidelity DNA Polymerase (New England Biolabs) with supplementation with High GC-enhancer to the reaction following the manufacturer's instructions and primers synthesized by Eurofins Genomics. The plasmids were

constructed using T4 DNA ligase (New England Biolabs) with a ratio of 3:1 of insert to backbone. All the molecular Biology work was planned using Geneious Prime version 2022.2.2. (https://www.geneious.com/). All plasmids and strains are listed in Table 1, plasmid sequences are available in the **Supplementary Information** Text 1.

Growth curves of *M. smegmatis*

Growth curves of *M. smegmatis* WT, $\Delta mftG$, $\Delta mftG$-*mftG,* and WT-*mftG* using HdB-Tyl supplemented with 10 g L^{-1} ethanol or 10 g L^{-1} glucose as the sole carbon source at 37 °C and 210 rpm were performed three times in biological triplicates. Evaluation of growth on 1% [v/v] glycerol, 5 g L^{-1} acetate, 10 g L^{-1} methanol, 10 g L^{-1} 1-propanol, 5 g L^{-1} 1-butanol, 0.5% hexanol [v/v], 0.010 g L^{-1} acetaldehyde, 10 g L^{-1} propane-1,2-diol, 10 g L^{-1} propane-1,3-diol were performed in at least duplicates. Culture of the strains grown on HdB-Tyl supplemented with 10 g L^{-1} glucose at 37 °C and 210 rpm for 24h was used as pre-inoculum. Cultures were centrifuged at 4,000 $\times$ *g* for 5 min and the supernatant was removed. The pellet was resuspended with base HdB-Tyl and used as inoculum for new cultures in a total volume of 40 mL in 250 mL Erlenmeyer flasks (50 mm Ø) with starting OD_{600} of 0.1 and closed with breathable rayon film (VWR). The cultures were monitored using the Cell Growth Quantifier (Aquila Biolabs) with readings every 60 s and incubated at 37 °C and 210 rpm until the late stationary phase was reached. Recorded data were further processed using CGQquant (Aquila Biolabs) for merging the replicates. The resulting average and standard deviation data were plotted in GraphPad Prism 9.

Ethanol, acetaldehyde, and acetate quantification

Cultures of the WT, $\Delta mftG$, $\Delta mftG$-*mftG,* and WT-*mftG* were grown (starting OD_{600} of 0.1, 37 °C, 210 rpm) in triplicates in HdB-Tyl supplemented with 10 g L^{-1} ethanol for 24

h, 32 h, 48 h, and 72 h and quantified for ethanol, acetaldehyde, and acetate concentration. The culture of $\Delta mftG$ starting at OD_{600} = 1 and medium without inoculum were also analyzed. The samples were centrifuged at 17,000 × g for 20 min and supernatants were sterilized using 0.2 µm cellulose acetate membrane filters (VWR) and kept at -20 °C until further analysis. The samples were diluted 1:10 with 0.005 mol L^{-1} H_2SO_4 and 50 µL injected in an HPLC X-LC (JASCO International Co) using a pre-column Kromasil 100 C 18, 40 mm x 4 mm, 5 µm (Dr. Maisch GmbH, Ammerbuch-Entringen) combined with the column Aminex HPX-87H Ion Exclusion Column, 300 mm x 7.8 mm, 9 µm (Bio-Rad) with isocratic mobile phase 0.005 mol L^{-1} H_2SO_4, flow 0.5 mL min^{-1}, heated to 50 °C. Detection was performed via refractive index (RI) and UV (210 nm) and compared with standards of ethanol (Uvasol for spectroscopy, Merck), acetaldehyde (Reagent plus, Sigma-Aldrich), and acetic acid (100%, water-free, p.a., Merck). Acetaldehyde content was further quantified using the supernatant of WT and $\Delta mftG$ grown in HdB-Tyl supplemented with 10 g L^{-1} ethanol alone and 10 g L^{-1} glucose combined with 10 g L^{-1} ethanol for 48 h as described in the acetaldehyde Assay Kit (Sigma-Aldrich) by the manufacturers.

Flow cytometry measurements

The cultures of WT, $\Delta mftG$, $\Delta mftG$-$mftG$, and WT-$mftG$ grown in HdB-Tyl supplemented with 10 g L^{-1} glucose, 10 g L^{-1} ethanol, 10 g L^{-1} glucose and 10 g L^{-1} ethanol combined or no carbon source (starvation) incubated at 37 °C and 210 rpm were sampled at 24 h, 48 h, and 72 h, centrifuged at 4,000 × g for 5 min and the supernatant removed. Samples were re-suspended in 500 µL HdB-Tyl with no carbon source with OD_{600} adjusted to 0.2. To samples of each condition were added either 2.5 µL of 750 µM of propidium iodide; 3 µL of 3 mM 3,3'-diethyloxacarbocyanine iodide [$DIOC_2(3)$]; 20 µL of 500 mM of the

protonophore (uncoupler) carbonyl cyanide 3-chlorophenylhydrazone (CCCP, Sigma-Aldrich) followed by 3 µL of 3 mM DIOC$_2$ (3) combined. Samples were incubated for 15 min and briefly vortexed. All the samples were injected on a FACS AriaFusion (BD Biosciences) and for each sample, FSC and SSC were detected using the blue laser (488 nm) and a threshold set to 400. The propidium iodide is excited by 488 nm and has an emission of 630 nm and emission was detected with the help of a 600 nm long-pass filter and a 610/20 nm bandpass filter. DiOC2(3) has the excitation maximum at 488 nm and the cells with a low level of transmembrane potential (or the cells treated with uncoupler CCCP) have a maximum emission at 530 nm (green color), the cells with a higher level of electric potential energy accumulate more concentration of this lipophilic dye, which results in its accumulation within the cells and these aggregates of dye have a maximum of emission at 600 nm (red fluorescence) (50, 51). The emission of DIOC$_2$(3) was detected with the help of a 502 nm long-pass filter and a 530/30 nm bandpass filter and 600 nm long-pass filter and a 610/20 nm bandpass. PMT voltages were adjusted to values of 600 V for the red channel and 350 V for the blue channel. The propidium iodide data were analyzed in a batch using flowCore and ggcyto R packages in R (version 4.1.0). Single cells were gated using FSC.A vs. FSC.H (intensities below 0.5e3 and above 1e5 have been discarded) and in the following step the highly fluorescent single cells were gated using YG610.A vs. FSC.A. The transmembrane potential data (DIOC$_2$(3) and CCCP) were directly gated using FlowJo™ v10.8 Software (BD Life Sciences) in coordinates B530.A ~ Y610.A (51)

Phenotype microarrays

The full panel of phenotype microarrays (PMs) on BioLog plates was used to detect additional phenotypes. The initial cultures of *M. smegmatis* WT and Δ*mftG* were grown

in LB overnight at 37 °C and 210 rpm, after 24 h the cultures were upscaled to 50 mL in HdB-Tyl supplemented with 10 g L^{-1} glucose. Preparation of *M. smegmatis* cells for PMs was performed with minor changes compared to the one previously described (52). After 24 h the cultures were centrifuged at 4,000 $\times$ *g* for 5 min and the supernatant was removed. The cells were resuspended in base HdB-Tyl and stored for 22 h at room temperature for starvation. The cultures were once again centrifuged at 4,000 $\times$ *g* for 5 min, the supernatant discarded and the cells resuspended in GN/IF-0 to an OD$_{600}$ of 0.68. The culture was then supplemented with a PM mixture. The culture mixture of the WT and $\Delta mftG$ was inoculated with 100 µL in each well of the 20 plates and incubated at 37 °C for 48 hours. The OD$_{595}$ was measured in a CLARIOStar microplate reader (BMG Labtech) after 48 h and 72 h. The sensitivity plates data was analyzed towards differences of growth compared to control growth of each strain and between $\Delta mftG$ and WT. Only differences in growth percentages comparing WT and $\Delta mftG$ [(%$\Delta mftG$ *100)/ % WT] above 150% or under 50% were considered for discussion and results were plotted using GraphPad Prism 9.

3.6 Fluorescent D-amino acid (FDAA) labeling

To target peptidoglycan biosynthesis, fluorescently labeled D-amino acids were incorporated into the nascent peptidoglycan as previously reported (53). Briefly, WT, $\Delta mftG,$ and $\Delta mftG$-*mftG* cells were grown in HdB-Tyl supplemented with either 10 g L^{-1} glucose, 10 g L^{-1} ethanol, or 10 g L^{-1} glucose combined with 10 g L^{-1} ethanol. The starvation condition was accomplished with the incubation of bacteria in a plain HdB-Tyl medium with no carbon source added. The cultures were incubated at 37 °C and 210 rpm until the mid-logarithmic growth phase. For FDAA incorporation, the green NADA was added to a final concentration of 250 µM, and cells were left to grow for 2,5 h. To

stop incorporation, cells were placed on ice for 2 min followed by one washing step with

ice-cold PBS. The red RADA was added to the same final concentration for 2,5 h. After

the final washing step with PBS, bacterial cells were fixed with 4% (v/v)

paraformaldehyde (PFA) solution for 20 min at 4 ºC.

Super-resolved structured illumination microscopy (SR-SIM)

For the SR-SIM imaging, 10 µL of the sample was spotted on 10 g L^{-1} agarose pads.

The agarose pads were covered with 1.5H coverslips (Roth) and stored at 4 °C for further

imaging. The SR-SIM data were acquired on an Elyra 7 system (Zeiss) equipped with a

63×/1.4 NA Plan-Apochromat oil-immersion DIC M27 objective lens (Zeiss), a Piezo

stage, and a PCO edge sCMOS camera with 82% QE and a liquid cooling system with

16-bit dynamic range. Using Lattice SIM mode, images were acquired with 13 phases.

NADA was detected with a 488 nm laser and a BP 495-590 emission filter; RADA was

detected with a 561 laser and an LP 570 emission filter. Super-resolution images were

computationally reconstructed from the raw data sets using default settings on ZenBlack

software (Zeiss). Images were analyzed using the Fiji ImageJ software (54).

NADH/NAD$^+$ and ADP/ATP ratio measurement

The *M. smegmatis* WT and Δ*mftG* strains were grown in triplicates on HdB-Tyl

supplemented with 10 g L^{-1} ethanol as the sole carbon source and incubated at 37 °C

and 210 rpm with starting OD_{600} = 0.1 for WT and OD_{600} = 1 for Δ*mftG* for a period of 48

h. Samples of culture were adjusted to OD_{600} 0.5 mL^{-1} and processed according to the

manufacturer's instruction ADP/ATP Ratio Assay Kit (Sigma-Aldrich). For NADH/NAD$^+$

quantification samples from WT and Δ*mftG* grown in triplicates on HdB-Tyl

supplemented with 10 g L^{-1} glucose for 24h were also quantified. Different cell

concentrations were tested with the best result OD_{600} of 0.1. Bacterial suspensions were

diluted in ice-cold PBS (8 g L^{-1} NaCl, 0.2 g L^{-1} KH$_2$PO$_4$, 1.15 g L^{-1} Na$_2$HPO$_4$, 0.2 g L^{-1}

KCl, at pH 7.4), samples were centrifuged at 10,000 x *g*, 4 °C, for 5 min, and the resulting

pellets were extracted with 100 µL of extraction buffer either for NAD+ or NADH

extraction. The resulting extracts were treated and measured according to the

NAD+/NADH Assay Kit manufactures' protocol (MAK460, Sigma-Aldrich). The readings

of each plate were acquired using a CLARIOStar microplate reader (BMG Labtech) and

data were further plotted and statistical analysis performed in GraphPad Prism9.

MFT profiling

The workflow for metabolome extraction was based on our previously described protocol

with small changes (17). The *M. smegmatis* WT and Δ*mftG*, cultures were incubated in

HdB-Tyl with either 10 g L^{-1} glucose or 10 g L^{-1} ethanol at 37 °C and 210 rpm in triplicates.

Cultures were quenched at the exponential phase, for WT and Δ*mftG* strain at 30 h in

glucose and WT strain at 35 h in ethanol. The Δ*mftG* strain grown in ethanol was

quenched at 60 h when it presented a small growth. Cultures were also sampled at

stationary phases, for WT and Δ*mftG* strains at 45 h in glucose and WT strain at 60 h in

ethanol. The *M. smegmatis* Δ*mftG-mftG* and WT-*mftG* strains were grown in HdB-Tyl

with 10 g L^{-1} ethanol at 37 °C and 210 rpm in triplicates and quenched only at the

stationary phase (60 h). The samples were normalized upon sampling to 10 mL of 1 unit

of OD_{600}, quenched in 20 mL of cold extraction mix (acetonitrile:methanol:water: FA;

60:20:19:1, v/v), and extracted as described previously (8). The lyophilized samples

were dissolved with 450 µL LC-MS grade water (VWR) twice, combined, and the extracts

were centrifuged at 17,000 $\times$ *g* for 20 min twice to remove debris and kept at -20 °C until

analysis. The metabolomes were analyzed using 10 µL injection in LC-MS/MS, Dionex

UltiMate 3000 UHPLC connected to a Q Exactive Plus mass spectrometer (Thermo Fisher Scientific), using the previously described method (8).

LC-MS/MS raw files were loaded into MZmine version 2.53 (55) and processed via the following pipeline for targeted peak analysis. Mass lists were created on MS^1 and MS^2 levels using the mass detector module set to centroid with a noise level of 0. Mass detection was followed by targeted peak integration using the targeted feature detection module set to MS level: 1, intensity tolerance: 50%, noise level: 0, *m/z* tolerance: 0 *m/z* or 5 ppm, retention time tolerance: 0.2 min, and a target list comprising expected protonated ions of the most abundant mycofactocin congener found *in vivo*: Methylmycofactocinol-8 (MMFT-8H$_2$, sum formula: $C_{62}H_{99}NO_{43}$, expected retention time: 6.81 min, expected *m/z* 1546.5663 $[M+H]^+$) and the corresponding oxidized form Methylmycofactocinone-8 (MMFT-8, sum formula: $C_{62}H_{97}NO_{43}$, expected retention time: 7.18 min, expected *m/z* 1544.5507 $[M+H]^+$). Theoretical mass-to-charge ratios were calculated using the respective sum formulae and the enviPat web interface (56). Resulting feature lists were filtered for the presence of at least one entry to remove empty rows and aligned using the RANSAC aligner module with the following settings: *m/z* tolerance: 0 *m/z* or 5 ppm, retention time tolerance: 0.2 min before and after correction, RANSAC iterations: 10^4, the minimum number of points: 20%, threshold value: 1, linear model: false, require same charge state: false. The resulting aligned feature list was manually inspected for misidentifications and exported as a comma-separated value table. Statistical analysis and plotting were performed with GraphPad Prism 9. Ratios between oxidized and reduced forms were calculated using MMFT-8H$_2$/ MMFT-8.

Purification of methylmycofactocinol-2 (MMFT-2H$_2$) from WT-*mftABCDEF*

To isolate MMFT-2H$_2$ from *M. smegmatis,* the mycofactocin gene cluster duplication strain WT-*mftABCDEF* was built by integration of the plasmid pPG23 in the WT strain. A first preculture was prepared through inoculation of 3 mL LBT with the strain and incubation for 24 h. The first preculture was used to inoculate a second preculture of 50 mL HdB-Tyl supplemented with 10 g L^{-1} glucose in a 250 mL shake flask and cultivated for 48 h. From the second preculture 1:100 [v/v] was used to inoculate the 5 L main culture of 2x HdB, i.e. all components except tyloxapol were concentrated twice compared to standard HdB-Tyl, supplemented with 3% [v/v] ethanol. The main culture was cultivated as 10x 500 mL cultures in 2 L shake flasks sealed with breathable rayon film (VWR) for reproducible aeration for 72 h. All cultivation steps were performed at 37 °C and 210 rpm in a shaker with a 25 mm throw. The culture was harvested through centrifugation at 8,000 × *g* and 18 °C for 3 min. The cell pellet of the harvested culture was washed with 500 mL ultrapure water once followed by centrifugation at 8,000 × *g* and 18 °C for 5 min and resuspension in 500 mL of 50 mM citrate phosphate buffer (Na$_2$HPO$_4$ anhydrous 7.1 g L^{-1}, citric acid 9.6 g L^{-1}, pH 5.2). The cell suspension was autoclaved at 134 °C for 30 min to disrupt the cells followed by treatment with 20 µg mL^{-1} of cellulase mix from *Trichoderma reesei* (Sigma Aldrich) at 37 °C for 18 h, which was subsequently heat-inactivated at 95 °C for 15 min to condense the pool of different long-chain mycofactocinols present *in vivo* to MMFT-2H$_2$ and PMFTH$_2$. The treated crude extract was cleared through centrifugation at 15,000 × *g* and 18 °C for 20 min. To avoid column overloading, the extract was split into two 250 mL parts before fractionation on 70 mL/10 g octadecyl modified silica solid phase extraction columns (Chromabond SPE, Machery-Nagel). Before fractionation, the SPE column was washed with three column volumes of methanol (CV, 1 CV = 40 mL) followed by equilibration with three CVs of 50 mM citrate phosphate buffer. After application of the crude extract, bound molecules

were washed with 1 CV water and step-wise eluted with 4 CV 80:20 [v/v] water:methanol, 3 CV 70:30 [v/v] water:methanol, and 1 CV 50:50 [v/v] water:methanol. Samples of each fraction were tested for the presence of MMFT-$2H_2$ via LC-MS as described above. The 70:30 fraction showed the highest concentration of MMFT-$2H_2$ and thus was subjected to further purification. The fraction was lyophilized at 1 mbar and -90 °C until all solvent was removed. The lyophilizate was resuspended in 500 µL water and subjected to a final purification via size-exclusion chromatography (SEC) using a Superdex 30 Increase 10/300 GL (Cytiva) equilibrated with 50 mL water operated at 500 µL min^{-1} on an NGC medium-pressure chromatography system (Bio-Rad). Per chromatography, 250 µL resuspended enriched fractions were separated. Each of the SEC fractions was tested for the presence of MMFT-$2H_2$ via LC-MS/MS as described above and the fractions with the highest concentrations of MMFT-$2H_2$ and lowest amounts of LC-MS/MS detectable impurities were combined and lyophilized at 0.01 mbar and -90 °C until complete dryness.

Synthesis of PMFT and PMFTH$_2$

Premycofactocinone (PMFT) and premycofactocinol (PMFTH$_2$) were synthesized as reported previously (17).

Homologous MftG expression and assay

The strain Δ*mftG-mftG*His$_6$ was designed for the homologous expression of a hexahistidine-tagged MftG protein. In order to confirm the correct expression and activity of the induced protein the Δ*mftG- mftG*His$_6$ was grown in HdB-Tyl supplemented with 10 g L^{-1} ethanol to confirm recovery from the phenotype. The strains Δ*mftG* and Δ*mftG- mftG*His$_6$ were inoculated in 200 ml HdB-Tyl supplemented with 5 g L^{-1} glucose and 20 g L^{-1} ethanol in 1L flasks covered with breathable membrane and starting OD_{600} of 0.1.

The cultures were incubated for 48 h at 37 °C and 210 rpm and the cell pellet were retrieved after centrifugation at 8,000 rpm for 20 min at 4 °C and stored at -20 °C. The cell pellet of both strains was resuspended in 8 mL PBS buffer with 5 mM imidazole, 300 mM NaCl, 50 µL protease inhibitor (1/100 solution), and 10 µM flavin adenine dinucleotide (FAD). The lysates were retrieved by sonicating (Bandelin Sonoplus UW 2070 homogenizer) the cells for 7.5 min (30 s on/off) at 70% cycle time and 70% amplitude in cold conditions and collection of supernatants after centrifugation at 15,000 rpm for 30 min at 4 °C. Lysates were incubated for 1 h 30 min with 2 mL Ni-NTA beads on ice under constant shaking. The beads and lysate were transferred to an empty column and flow-through discarded, the column was washed with 5 mM (2CV) and 20 mM (2CV) imidazole by gravity flow. The partially purified fraction was eluted using 100 mM imidazole and the buffer was changed to 0.1 M sodium phosphate at pH 7 using a PD-10 column. The proteins were concentrated using a Vivaspin6 10 kDa filter unit (Sartorius). The protein concentration was measured via absorption at 280 nm using a Nanodrop spectrophotometer (ThermoFisher) and adjusted to 0.500 mg mL^{-1}.

From each extract, 20 µL were assayed with 20 µL of Δ*mftG* metabolite extract and incubated at 37 °C, an initial sample was quenched with acetonitrile (1:1) directly after the mixture and final samples were quenched after 18 h of incubation. Pure substrates at 2 µM of MMFT-2H$_2$, synthetic PMFTH$_2$, and PMFT (17) were also used in the enzymatic assays, and samples every 20 min for a total of 2 hours were retrieved and quenched as before. Heat-inactivated cell-free extract was used to confirm enzyme-dependent activity. All variables were performed in triplicates. The extracts were centrifuged at 17,000 $\times$ *g* for 20 min and the supernatant was retrieved and centrifuged a second time in the same conditions to remove debris. From each assay, 10 µL were injected in LC-MS/MS and analyzed as described above. Differences of MMFT-8H$_2$ and

MMFT-8, MMFT-2H$_2$ and MMFT-2, PMFTH$_2$ and PMFT amounts between initial and different incubation points were retrieved with targeted analysis to the later compounds using MZmine version 2.53 (55).

6.3 Heterologous MftG expression and activity

Escherichia coli NiCo21(DE3) was transformed with pPG36_biocat as described. The pMAL-c4x-based plasmid harbored a gene for the expression of a maltose-binding protein fused N-terminally to MftG from *M. smegmatis* (WP_014877070.1) codon-optimized for the expression in *E. coli* obtained from BioCat. A single colony of *E. coli* NiCo21(DE3)/pPG36_biocat was used to inoculate a preculture on 5 mL LB which was incubated at 37 °C and 210 rpm for 18 h. Three mL of the preculture were used to inoculate 300 mL of LB in a 1 L shake flask. The expression culture was cultivated at 37 °C and 210 rpm until an OD$_{600}$ of 0.6, when gene expression was induced through the addition of isopropyl β-D-1-thiogalactopyranoside (IPTG) at a final concentration of 1 mM. The induced expression culture was incubated at 18 °C and 210 rpm for 20 h. Media were supplemented with 100 µg mL^{-1} of ampicillin and cultivation was performed using an incubator with a throw of 25 mm. Cells were harvested through centrifugation at 8,000 x *g* and 4 °C for 3 min. The cell pellet was resuspended in 12 mL lysis buffer (20 mM Tris, 200 mM NaCl, 1 mM EDTA, pH 7.4) and disrupted through sonication using a Bandelin Sonoplus UW 2070 homogenizer equipped with an MS 73 microtip operated at 70% cycle time and 70% amplitude for a total on-time of 7 min with cycles of 1 min sonication followed by 30 s pause to cool the lysate. Sonication was performed on ice. The lysate was cleared through centrifugation at 20,000 x *g* and 4 °C for 30 min followed by filtration through a 0.22 µm cellulose-acetate filter. The cleared lysate was subjected to affinity chromatography using a 1 mL MBPTrap HP maltose-binding protein affinity

chromatography column (Cytiva) equilibrated with 10 mL lysis buffer. Bound proteins were washed with 20 mL of lysis buffer and eluted using 10 mL of elution buffer (20 mM Tris, 200 mM NaCl, 1 mM EDTA, 10 mM Maltose, pH 7.4). Chromatography was performed with cooled buffers and at a flow-rate of 1 mL min^{-1}. Target protein-containing fractions were combined and concentrated using an Amicon Ultra-4 50 kDa filter (Merck) operated at 7,500 x g and 4 °C. Protein concentrations were determined using the A$_{280}$ nm method of a NanoDrop (Thermo Fisher Scientific) with the calculation setting of 1 ABS = 1 mg mL^{-1}. Mycofactocinol oxidation assays were performed in a total reaction volume of 30 µL in lysis buffer including 2 µM MMFT-2H$_2$, 100 µM FAD, and either one of the tested potential electron acceptors: 2,6-dichlorophenolindophenol (DCPIP), nicotinamide adenine dinucleotide (NAD), N,N-dimethyl-4-nitrosoaniline (NDMA), phenazine methosulfate (PMS), at a concentration of 100 µM and 87 µg protein, or no artificial electron acceptor and at different protein concentrations as indicated (Figure 6E). The reactions were incubated at 37 °C for 24 h and quenched through heat-inactivation of the enzyme at 100 °C for 10 min. The quenched reactions were centrifuged at 17,000 x g and 18 °C for 20 min and 10 µL of the supernatant were subjected to LC-MS/MS analysis with subsequent targeted data analysis as described. The target features for data analysis were the mycofactocin substrate and expected product of the assay, namely methylmycofactocinol-2 (MMFT-2H$_2$, sum formula: C$_{26}$H$_{39}$NO$_{13}$, expected retention time: 7.10 min, expected m/z 574.2494 [M+H]$^+$) and methylmycofactocinone-2 (MMFT-2, sum formula: C$_{26}$H$_{37}$NO$_{13}$, expected retention time: 7.47 min, expected m/z 572.2338 [M+H]$^+$). All experiments were performed in triplicates and controls were set up with lysis buffer replacing the enzyme. All tested potential electron acceptors were purchased from Sigma-Aldrich.

Whole-cell respiration assay

The WT and $\Delta mftG$ cells were incubated on HdB-Tyl supplemented with 10 g L^{-1} ethanol for 48h at starting OD$_{600}$ of 0.1 for WT and 0.7 for $\Delta mftG$ at 37 °C and 210 rpm. For the starvation experiment, WT was cultivated for 24 h in the same conditions as the latter in HdB-Tyl without carbon sources and starting OD$_{600}$ of 0.7. After this period the cells were collected by centrifugation at 5,000 × g for 10 min and pellets were resuspended in fresh HdB supplemented with 0.25 g L^{-1} Tyloxapol without carbon sources and normalized to OD$_{600}$ of 0.7.

Aliquots of 200 µL of bacterial samples were placed in the electrochemical chamber supplied with a Clark-type polarographic sensor (Oxytherm^{+}, Hansatech Instruments). Before experiments, the chamber was calibrated according to the manufacturer's manual ('liquid phase calibration') and set to 37 °C and 50 rpm stirring. The oxygen consumption was monitored over 10 min. The retrieved data was further plotted using R version 2023.03.0+.

Isolation of mycobacterial membranes

M. smegmatis WT and $\Delta mftG$ strains were grown in 320 mL HdB-Tyl medium supplemented with 1 g L^{-1} ethanol as the sole carbon source in 2 L flasks covered by breathable rayon film (VWR) in triplicates and incubated at 37 °C and 210 rpm with a starting OD$_{600}$ of 0.1 for WT and 2.0 for $\Delta mftG$ for a period of 48 h.

The cells were concentrated by centrifugation at 7,000 × g for 10 min and the resulting pellets were resuspended in buffer (38 mL) of the following composition: 50 mM Tris-HCl, 300 mM sucrose, 5 mM EDTA, 50 mM MgCl$_2$ at pH 6.7. Pellets were disrupted with 0.1 mm ZrO$_2$ beads (CarlRoth, GmbH) in 15 mL plastic tubes using a bead beater (MP Biomedicals, FastPrep-24 5G). A total of 6 cycles of beating were applied (3 times of

application of the pre-installed program for *M. tuberculosis* cells from the manufacturer).

After each step of beating, tubes were cooled down on ice for at least one minute. To

get rid of the beads and undisrupted cells the resulting material was centrifuged twice at

4 °C and 6,000 $\times$ g for 15 min. The obtained lysate was further ultra-centrifuged (Thermo

Scientific™ Sorvall LYNX 6000 Superspeed Centrifuge) at 4 °C, 100,000 $\times$ g for 2 h to

obtain the sedimentation of the mycobacterial membrane fraction and clarified soluble

fraction (cytoplasm and disrupted periplasm). The isolated membranes were stored on

ice for no longer than 12 hours.

Analysis of MMFT-2H$_2$ and PMFTH$_2$ oxidation in respiration assay

Oxygen consumption was measured using Oxytherm$^+$ (Hansatech Instruments)

advanced oxygen electrode system at 37 °C and stirred at 50 rpm. For homogenization

of mycobacterial membranes, the assay buffer of the following composition was used:

50 mM sucrose, 5 mM MgCl$_2$, 5 mM KF, 2 mM AMP, 4 mM K$_2$HPO$_4$, 50 mM Tris at pH 6.7.

All experiments were performed with 250 µL of membrane homogenates. The respiration

stimulation started after circa 100 s from the beginning of the oxygen measurement

experiments by the addition of 1 mM of the substrates: MMFT-2H$_2$, PMFTH$_2$, NADH

(CarlRoth), or sodium succinate (Sigma-Aldrich). Inhibition of the reaction was tested by

the addition of a final concentration of 2.5 mM KCN (Sigma-Aldrich) to the solution. To

exclude non-enzymatic oxidation, controls such as reaction mixtures without substrates,

as well as microwave-inactivated membrane fractions were included. Total protein

concentration was quantified using Roti-Nanoquant 5x Concentrate (Carl Roth) and

normalization on the total protein concentration of the fractions was used to make the

results comparable. The respirometry curves were plotted using R version 2023.03.0+.

For the analysis of PMFTH$_2$ conversion into PMFT, 200 µL of the sample from the respirometry analysis was quenched by 200 µL of acetonitrile followed by two centrifugations at 17,000 $\times$ g for 15 min to remove debris. To assess if PMFTH$_2$ conversion was membrane-bound, an assay was carried out with soluble fractions and further quenched as previously described. The LC-MS/MS profiling and the obtained chromatograms were analyzed as described above in this study. Generated csv tables were further plotted and analyzed in R version 2023.03.0+ .

RNA extraction, library preparation and analysis of *M. smegmatis* MC^2 155 WT and Δ*mftG*

M. smegmatis MC^2 155 WT and Δ*mftG* were grown on HdB-Tyl with 10 g L^{-1} for 24h. Cultures were centrifuged, supernatant removed and resuspended on HdB-Tyl plain medium and used as inoculum for triplicate cultures on HdB-Tyl with 10 g L^{-1} of ethanol at a starting OD$_{600}$ concentration of 0.1. Samples of 4mL of WT at exponential phase and 8ml of Δ*mftG* after 60h of incubation were retrieved and further processed for RNA extraction following the manufacturer's instructions (InnuPREP RNA Mini kit 2.0 – Analytik Jena). An additional DNAse step was included to remove traces of gDNA. RNA samples were then subjected to RNA-Seq library preparation following the stranded total RNA prep Ligation with Ribo-Zero plus kit (Illumina). After depletion of the rRNA by hybridization with directed probes, the RNA samples were normalized to 500 ng/sample. The RNA samples were then subjected to fragmentation and denaturation, followed by reverse transcription through first-strand (and subsequently second-strand) cDNA synthesis. After 3′end adenylation, anchors and indexes were ligated. Barcoding was accomplished using the IDT® for Illumina® RNA UD Indexes Set A, Ligation (Ref# 20040553, Ilumina, Berlin, Germany). Standard AMPURE XP Beads protocols (Ref#

A63881; Beckman, Krefeld, Germany) were employed to purify the final library fragments, and their size and molarity were verified using a Tape Station 2200 (Agilent, Waldbronn, Germany). The equimolarly pooled libraries were subsequently subjected to a 100-cycle run on a NovaSeq 6000 apparatus (Ilumina, Berlin, Germany) using an SP flowcell v. 1.5 (Ref# 20028401; Agilent, Waldbronn, Germany).

Raw reads were checked for quality using fastP (50), and polyX tails (-x 10), low quality ends (-3, -5, -M 25) were removed, and resulting sequences shorter than 25 were discarded (-l 25). The remaining reads were mapped to the reference genome (RefSeq accession GCF_000015005.1) using bwa-mem2 (55) with default parameters for the affine gap scoring model. For each sample and every gene annotation, all overlapping reads were counted and a counts table was generated for every experiment condition using custom scripts. The counts' tables were used as an input to the R package DESeq2 (56) to compute fold changes and false discovery rates of differentially expressed genes between the control condition and each experiment condition.

Statistical analysis

The data retrieved in this study were further plotted and analyzed statistically using GraphPad Prism 9. Significance was calculated using either two-way ANOVA with Tukey's multiple comparisons test once interaction index was not significant, or one-way ANOVA for multiple comparisons. Significance values were plotted directly on the graphs and only more relevant comparisons are shown.

Data Availability

The transcriptomic datasets produced in this study are available via the Gene Expression Omnibus (GEO) database (GEO accession numbers: GSE250373, GSM7978029 – GSM7978040).

Author contributions

Ana Patrícia Graça: conceptualization, data curation, formal analysis, investigation, methodology, validation, visualization, writing- original draft, writing- review and editing,

Vadim Nikitushkin: conceptualization, data curation, formal analysis, investigation, methodology, validation, visualization, writing- original draft, writing- review and editing,

Mark Ellerhorst: data curation, formal analysis, investigation, methodology, validation, writing- original draft, writing- review and editing

Cláudia Vilhena: conceptualization, data curation, formal analysis, investigation, methodology, validation, visualization, writing- original draft, writing- review and editing

Tilman E. Klassert: conceptualization, formal analysis, methodology, resources, writing- review and editing

Andreas Starick: data curation, methodology, validation, writing- review and editing,

Malte Siemers: data curation, formal analysis, methodology, validation, writing- review and editing

Walid K. Al-Jammal: methodology, validation

Ivan Vilotijevic: conceptualization, supervision, validation, methodology, funding acquisition

Hortense Slevogt: conceptualization, validation, funding acquisition

Kai Papenfort: validation, supervision, funding acquisition, writing-review and editing

Gerald Lackner: conceptualization, validation, visualization, funding acquisition, project administration, resources, supervision, writing- original draft, writing- review and editing

Acknowledgments

We thank Bettina Bardl (Bio Pilot Plant, Leibniz-HKI) for HPLC analysis of metabolites. We thank the Microverse Imaging Center for providing microscope facility support for data acquisition. The ELYRA 7 was funded by the Free State of Thuringia with grant number 2019 FGI 0003. The Microverse Imaging Center is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy - EXC 2051 - Project-ID 390713860. DAAD, German Academic Exchange Service funding is greatly acknowledged (graduate fellowship to W.K.A-J, fellowship for university academics and scientists to V.N). A.P.G and G.L. thank the Carl Zeiss Foundation for financial support.

Conflict of interest

The authors declare to not have competing interests.

References

1. Tsukamura, M. (1966) Adansonian classification of mycobacteria J Gen Microbiol **45**, 253-273

2. Bloch, H., and Segal, W. (1956) Biochemical differentiation of *Mycobacterium tuberculosis* grown *in vivo* and *in vitro* J Bacteriol **72**, 132-141

3. Zimmermann, M., Kogadeeva, M., Gengenbacher, M., McEwen, G., Mollenkopf, H. J., Zamboni, N. *et al.* (2017) Integration of metabolomics and transcriptomics reveals a complex diet of *Mycobacterium tuberculosis* during early macrophage infection mSystems **2**,

4. Haft, D. H. (2011) Bioinformatic evidence for a widely distributed, ribosomally produced electron carrier precursor, its maturation proteins, and its nicotinoprotein redox partners BMC Genomics **12**, 21

5. Krishnamoorthy, G., Kaiser, P., Lozza, L., Hahnke, K., Mollenkopf, H. J., and Kaufmann, S. H. E. (2019) Mycofactocin is associated with ethanol metabolism in mycobacteria mBio **10**, e00190-00119

6. Dubey, A. A., Wani, S. R., and Jain, V. (2018) Methylotrophy in mycobacteria: Dissection of the methanol metabolism pathway in *Mycobacterium smegmatis* J Bacteriol **200**,

7. Dubey, A. A., and Jain, V. (2019) Mycofactocin is essential for the establishment of methylotrophy in *Mycobacterium smegmatis* Biochem Biophys Res Commun **516**, 1073-1077

8. Peña-Ortiz, L., Graça, A. P., Guo, H. J., Braga, D., Köllner, T. G., Regestein, L. *et al.* (2020) Structure elucidation of the redox cofactor mycofactocin reveals oligo-glycosylation by MftF Chem Sci **11**, 5182-5190

9. Shimizu, T., Suzuki, K., and Inui, M. (2024) A mycofactocin-associated dehydrogenase is essential for ethylene glycol metabolism by *Rhodococcus jostii* RHA1 Appl Microbiol Biotechnol **108**, 58

10. Mendauletova, A., and Latham, J. A. (2022) Biosynthesis of the redox cofactor mycofactocin is controlled by the transcriptional regulator MftR and induced by long-chain acyl-CoA species J Biol Chem **298**, 101474

11. Nikitushkin, V., Shleeva, M., Loginov, D., Dycka, F. F., Sterba, J., and Kaprelyants, A. (2022) Shotgun proteomic profiling of dormant, 'non-culturable' *Mycobacterium tuberculosis* PloS One **17**, e0269847

12. Krishnamoorthy, G., Kaiser, P., Constant, P., Abu Abed, U., Schmid, M., Frese, C. K. *et al.* (2021) Role of premycofactocin synthase in growth, microaerophilic adaptation, and metabolism of *Mycobacterium tuberculosis* mBio **12**, e0166521

13. Ayikpoe, R., Govindarajan, V., and Latham, J. A. (2019) Occurrence, function, and biosynthesis of mycofactocin Appl Microbiol Biotechnol **103**, 2903-2912

14. Khaliullin, B., Ayikpoe, R., Tuttle, M., and Latham, J. A. (2017) Mechanistic elucidation of the mycofactocin-biosynthetic radical S-adenosylmethionine protein, MftC J Biol Chem **292**, 13022-13033

15. Ayikpoe, R., Salazar, J., Majestic, B., and Latham, J. A. (2018) Mycofactocin biosynthesis proceeds through 3-amino-5-[(p-hydroxyphenyl)methyl]-4,4-dimethyl-2-pyrrolidinone (AHDP); direct observation of MftE Specificity toward MftA Biochemistry **57**, 5379-5383

16. Ayikpoe, R. S., and Latham, J. A. (2019) MftD catalyzes the formation of a biologically active redox center in the biosynthesis of the ribosomally synthesized and post-translationally modified redox cofactor, mycofactocin J Am Chem Soc **141**, 13582-13591

17. Ellerhorst, M., Barth, S. A., Graça, A. P., Al-Jammal, W. K., Peña-Ortiz, L., Vilotijevic, I. *et al.* (2022) *S*-Adenosylmethionine (SAM)-dependent methyltransferase MftM is responsible for methylation of the redox cofactor mycofactocin ACS Chem Biol **17**, 3207–3217

18. Sützl, L., Foley, G., Gillam, E. M. J., Boden, M., and Haltrich, D. (2019) The GMC superfamily of oxidoreductases revisited: analysis and evolution of fungal GMC oxidoreductases Biotechnol Biofuels **12**, 118

19. Cavener, D. R. (1992) GMC oxidoreductases. A newly defined family of homologous proteins with diverse catalytic activities J Mol Biol **223**, 811-814

20. Ferri, S., Kojima, K., and Sode, K. (2011) Review of glucose oxidases and glucose dehydrogenases: a bird's eye view of glucose sensing enzymes J Diabetes Sci Technol **5**, 1068-1076

21. Koch, C., Neumann, P., Valerius, O., Feussner, I., and Ficner, R. (2016) Crystal structure of alcohol oxidase from *Pichia pastoris* PloS One **11**, e0149846

22. Landfald, B., and Strom, A. R. (1986) Choline-glycine betaine pathway confers a high level of osmotic tolerance in *Escherichia coli* J Bacteriol **165**, 849-855

23. Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G. *et al.* (2022) AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models Nucleic Acids Res **50**, D439-D444

24. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O. *et al.* (2021) Highly accurate protein structure prediction with AlphaFold Nature **596**, 583-589

25. Aleksenko, V. A., Anand, D., Remeeva, A., Nazarenko, V. V., Gordeliy, V., Jaeger, K. E. *et al.* (2020) Phylogeny and structure of fatty acid photodecarboxylases and glucose-methanol-choline oxidoreductases Catalysts **10**,

26. Gao, C. H., Yu, G. C., and Cai, P. (2021) ggVennDiagram: An intuitive, easy-to-use, and highly customizable R package to generate Venn diagram Front Genet **12**,

27. Dubey, A. A., and Jain, V. (2019) MnoSR is a *bona fide* two-component system Involved in methylotrophic metabolism in *Mycobacterium smegmatis* Appl Environ Microbiol **85**,

28. Kpebe, A., Guendon, C., Payne, N., Ros, J., Khelil Berbar, M., Lebrun, R. *et al.* (2023) An essential role of the reversible electron-bifurcating hydrogenase Hnd for ethanol oxidation in *Solidesulfovibrio fructosivorans* Front Microbiol **14**, 1139276

29. Peña-Ortiz, L., Schlembach, I., Lackner, G., and Regestein, L. (2020) Impact of oxygen supply and scale up on *Mycobacterium smegmatis* cultivation and mycofactocin formation Front Bioeng Biotechnol **8**,

30. Garcia-Heredia, A., Pohane, A. A., Melzer, E. S., Carr, C. R., Fiolek, T. J., Rundell, S. R. *et al.* (2018) Peptidoglycan precursor synthesis along the sidewall of pole-growing mycobacteria Elife **7**,

31. Pacifico, C., Fernandes, P., and de Carvalho, C. (2018) Mycobacterial response to organic solvents and possible implications on cross-resistance with antimicrobial agents Front Microbiol **9**, 961

32. Wu, M. L., Gengenbacher, M., and Dick, T. (2016) Mild nutrient starvation triggers the development of a small-cell survival morphotype in mycobacteria Front Microbiol **7**, 947

33. Tran, S. L., and Cook, G. M. (2005) The F_1F_0-ATP synthase of *Mycobacterium smegmatis* is essential for growth J Bacteriol **187**, 5023-5028

34. Kana, B. D., Weinstein, E. A., Avarbock, D., Dawes, S. S., Rubin, H., and Mizrahi, V. (2001) Characterization of the cydAB-encoded cytochrome bd oxidase from *Mycobacterium smegmatis* J Bacteriol **183**, 7076-7086

35. Yakushi, T., and Matsushita, K. (2010) Alcohol dehydrogenase of acetic acid bacteria: structure, mode of action, and applications in biotechnology Appl Microbiol Biotechnol **86**, 1257-1265

36. Cook, G. M., Hards, K., Vilcheze, C., Hartman, T., and Berney, M. (2014) Energetics of respiration and oxidative phosphorylation in Mycobacteria Microbiol Spectr **2**,

37. Berney, M., and Cook, G. M. (2010) Unique flexibility in energy metabolism allows mycobacteria to combat starvation and hypoxia PLoS One **5**, e8614

38. Tan, T. C., Kracher, D., Gandini, R., Sygmund, C., Kittl, R., Haltrich, D. *et al.* (2015) Structural basis for cellobiose dehydrogenase action during oxidative cellulose degradation Nat Commun **6**, 7542

39. Anthony, C. (2001) Pyrroloquinoline quinone (PQQ) and quinoprotein enzymes Antioxid Redox Signal **3**, 757-774

40. Takahashi, K., Hirose, Y., Kamimura, N., Hishiyama, S., Hara, H., Araki, T. *et al.* (2015) Membrane-associated glucose-methanol-choline oxidoreductase family enzymes PhcC and PhcD are essential for enantioselective catabolism of dehydrodiconiferyl alcohol Appl Environ Microbiol **81**, 8022-8036

41. DeJesus, M. A., Gerrick, E. R., Xu, W., Park, S. W., Long, J. E., Boutte, C. C. *et al.* (2017) Comprehensive essentiality analysis of the *Mycobacterium tuberculosis* genome via saturating transposon mutagenesis mBio **8**,

42. Adolph, C. R. (2021) Essentiality of succinate metabolism in *Mycobacterium tuberculosis* Thesis, University of Otago

43. Bosch, B., DeJesus, M. A., Poulton, N. C., Zhang, W., Engelhart, C. A., Zaveri, A. *et al.* (2021) Genome-wide gene expression tuning reveals diverse vulnerabilities of *M. tuberculosis* Cell **184**, 4579-4592 e4524

44. Dijkman, W. P., Binda, C., Fraaije, M. W., and Mattevi, A. (2015) Structure-based enzyme tailoring of 5-hydroxymethylfurfural oxidase ACS Catal **5**, 1833-1839

45. Goddard, T. D., Huang, C. C., Meng, E. C., Pettersen, E. F., Couch, G. S., Morris, J. H. *et al.* (2018) UCSF ChimeraX: Meeting modern challenges in visualization and analysis Protein Sci **27**, 14-25

46. Katoh, K., Misawa, K., Kuma, K., and Miyata, T. (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform Nucleic Acids Res **30**, 3059-3066

47. Price, M. N., Dehal, P. S., and Arkin, A. P. (2010) FastTree 2--approximately maximum-likelihood trees for large alignments PLoS One **5**, e9490

48. Bianchini, G., and Sánchez-Baracaldo, P. (2023) TreeViewer - Cross-platform software to draw phylogenetic trees (Version 2.1.0) [Computer software] Zenodo

49. Ofer, N., Wishkautzan, M., Meijler, M., Wang, Y., Speer, A., Niederweis, M. *et al.* (2012) Ectoine biosynthesis in *Mycobacterium smegmatis* Appl Environ Microbiol **78**, 7483-7486

50. Novo, D., Perlmutter, N. G., Hunt, R. H., and Shapiro, H. M. (1999) Accurate flow cytometric membrane potential measurement in bacteria using diethyloxacarbocyanine and a ratiometric technique Cytometry **35**, 55-63

51. Nikitushkin, V. D., Trenkamp, S., Demina, G. R., Shleeva, M. O., and Kaprelyants, A. S. (2020)
Metabolic profiling of dormant *Mycolicibacterium smegmatis* cells' reactivation reveals a gradual
assembly of metabolic processes Metabolomics **16**, 24
52. Karlikowska, M., Singh, A., Bhatt, A., Ott, S., Bottrill, A. R., Besra, G. S. *et al.* (2021) Biochemical
and phenotypic characterisation of the *Mycobacterium smegmatis* transporter UspABC Cell Surf
7, 100052
53. Kuru, E., Radkov, A., Meng, X., Egan, A., Alvarez, L., Dowson, A. *et al.* (2019) Mechanisms of
incorporation for D-amino acid probes that target peptidoglycan biosynthesis ACS Chem Biol **14**,
2745-2756
54. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T. *et al.* (2012) Fiji:
an open-source platform for biological-image analysis Nat Methods **9**, 676-682
55. Pluskal, T., Castillo, S., Villar-Briones, A., and Oresic, M. (2010) MZmine 2: modular framework
for processing, visualizing, and analyzing mass spectrometry-based molecular profile data BMC
Bioinformatics **11**, 395
56. Loos, M., Gerber, C., Corona, F., Hollender, J., and Singer, H. (2015) Accelerated isotope fine
structure calculation using pruned transition trees Anal Chem **87**, 5738-5744

Tables

Table 1: List of *M. smegmatis* and *E. coli* strains, vectors, plasmids and primers

used and generated on the course of this study.

Strain	Description	Reference
WT	*Mycolicibacterium smegmatis* MC^2 155	(5)
Δ*mftG*	derivate of WT without *mftG* replaced with *loxP* site	this study
Δ*mftG*-*mftG*	derivate of Δ*mftG* integrated with pPG29 at the attB site	this study
WT-*mftG*	derivate of WT integrated with pPG29 at the attB site	this study
Δ*mftG*- *mftG*His$_6$	derivate of Δ*mftG* integrated with pPG32 at the attB site	this study
WT-*mftABCDEF*	derivate of WT integrated with pPG23 at the attB site	this study
NiCo21(DE3)Competent *E. coli*	Derived from *E. coli* BL21 (DE3)	NEB BioLabs
Escherichia coli TOP10	F⁻*mcr*A Δ(*mrr-hsd*RMS-*mcr*BC) φ80*lac*ZΔM15 Δ*lac*X74 *rec*A1 *ara*D139 Δ(*ara-leu*)7697 *gal*U *gal*K λ⁻*rps*L(Str^R) *end*A1 *nup*G	Thermo Fischer
Plasmid name	**Backbone**	**Reference**

pML2424	vector for double crossover event with tdTomato, gfp-hyg cassette, and PAL5000ts	(49)
pML2714	vector with kanamycin resistance for Cre recombinase expression and *gfp-hyg* cassette removal	(49)
pPG20	pMCpAINT derivate with kanamycin resistance, potential mycofactocin promotor, and *mftF*	(8)
pPG17	pML2424 with up and downstream regions of *mftG*	this study
pPG23	pMCpAINT derivate with kanamycin resistance, potential mycofactocin promotor, and *mftABCDEF*	this study
pPG29	pPG20 with *mftF* replaced with *mftG*	this study
pPG32	pPG29 with mftG replaced with *mftG*His$_6$	this study
pPG36	pMAL-C4X with *malE* fused with *mftG* codon optimized for *E. coli* expression	this study
Primer name	**Primer sequence 5'-3'**	**Amplicon**
GMC_up_F1	GCTACACTAGTCGGTGTCGTATGTGCCGAG	upstream region of *mftG*
GMC_up_R1	GCTACATTTAAATTCAAAGTCGGCGGCTAACTC	
GMC_dn_F1	GCTACTTAATTAATCGACGGCTCGATCATGC	Downstream region of *mftG*
GMC_dn_R1	GCTACATGCATGTTGTCGAGGCTCCGGTG	
INT_GMC_F1	CACTATGGGTCGACGCTGAC	Internal region of *mftG*
INT_GMC_R1	GCGTGACTTACCAATTCGCG	
EXT_GMC_F1	AACATCGTGGCCCGGTAC	External region of *mftG*
EXT_GMC_R1	CTCCTCACGCGACGACTC	
pMCpAINT_FC_F	GCTACAAGCTTATCGATGTCGACGTAGTTAAC	Backbone pMCpAINT introducing NdeI
pPG20_NdeI_R	GCTACCCATATGCGTATGGTCTCGACAGTTGT	
GMC_COMP_F1	GCTACCCATATGGAGTTAGCCGCCGACTTT	Insertion of 6 histidines C-terminally
GMC_Hist_R3	GCTACAAGCTTACTATTAG-TGGTGGTGGTGGTGGTGGGTCGCGATG AACTCGGC	
pMCpAINT_conf_2_F	CTGATACCGCTCGCCGCA	Sequencing confirmation
pMCpAINT_conf_2_R	CTTTCGACTGAGCCTTTCGT	

MFTKIMS_FC_KI_CLUSTER_F2	GCTACCCATGGTCGGACATCTCTCACACCCC	Region from hypothetical mycofactocin precursor until end of *mftF*
MS_FC_KI_CLUSTER_R1	GTTAACTACGTCGACATCGATAA-GCTTTCAAAGTCGGCGGCTAACTC	

| MFTKIMS_FC_KI_CLUSTER_F2 | GCTACCCATGGTCGGACATCTCTCACACCCC | Region from hypothetical mycofactocin precursor until end of *mftF* |

Supplementary Information Files

Supplementary Figure S1 (PDF): **Phylogenetic tree (maximum likelihood) of the MftG and GMC sequences described in Supplementary Table S1.**

Supplementary Information (PDF): Single file containing Supplementary Figures S2 – S3 and Supplementary Text 1.

Supplementary Table S1 (.xlsx): Set of organisms encoding MftG or MftC homologs (co-occurrence table). The co-occurrence table (sheet 1) contains accession numbers of MftG and MftC candidate proteins, as well as further GMC proteins found in each organism. The phylogenetic analysis was based on all GMC enzymes including MftG listed in the co-occurrence table as well as the GMC proteins listed, under 'Reference GMC enzymes' (sheet 2).

Supplementary Table S2 (.xlsx): Gene expression analysis of *M. smegmatis* MC^2 155 Δ*mftG* compared to WT grown on either ethanol or glucose as a carbon source. Sheet 1: Gene expression in Δ*mftG* mutants compared to WT of *M. smegmatis* MC2 155 grown on 10 g L^{-1} ethanol as the sole carbon source. Sheet 2: Gene expression in Δ*mftG* mutants compared to WT of *M. smegmatis* MC2 155 grown on 10 g L^{-1} glcuose as the sole carbon source. Black: all the genes with adjusted p-value ≤0.05. Grey: all the genes that with adjusted p-value > 0.05.