A high-resolution haplotype pangenome uncovers somatic hybridization,
recombination and intercontinental migration in oat crown rust

Eva C. Henningsen[1,2], David Lewis[1], Eric Nazareno[3], Yung-Fen Huang[4], Brian J.
Steffenson[3], Brendan Boesen[1], Shahryar F. Kianian[3,5], Eric Stone[2], Peter N. Dodds[1],
Jana Sperschneider[1], Melania Figueroa[1*]

[1]Commonwealth Scientific and Industrial Research Organisation, ACT Australia
[2]Australian National University, ACT Australia
[3]University of Minnesota, Saint Paul, MN U.S.A.
[4]National Taiwan University, Taipei, Taiwan
[5]Cereal Disease Laboratory, USDA-ARS, St. Paul, MN U.S.A.
*Corresponding author: M. Figueroa; E-mail: Melania.Figueroa@csiro.au

Keywords: haplotype, mating type, pangenome, *Puccinia coronata*, somatic
hybridization, virulence

Funding
This project was supported by the Bioplatforms Australia Plant Pathogen 'Omics
Initiative, GRDC project CSP2204 007RTX, USDA-NIFA BBSRC award 2022-67013-
36505, the CSIRO Research Office, and grant 109-2313-B-002-028-MY3 from the
National Science and Technology Council of Taiwan. ECH was supported by the ANU
University Research Scholarship and ANU/CSIRO Digital Agriculture PhD
Supplementary Scholarship.

Eva Henningsen

Abstract

Basidiomycetes like rust fungi have complex reproductive cycles and dikaryotic life stages which influence their population structure and evolution. *Puccinia coronata* f. sp. *avenae* (*Pca*), the causal agent of oat crown rust, is a pathogen of global economic importance. To investigate the genetic diversity of the species, as well as the role of mating type system and nuclear exchange (somatic hybridization) in host adaptation of *Pca* we acquired whole genome sequencing data of Taiwanese and Australian isolates, adding to existing data for USA and South African populations. An atlas of 30 chromosome-level, fully-phased nuclear haplotypes from six USA isolates and nine Australian isolates was generated to capture the genomic composition of key oat crown rust lineages. This study provides evidence of nuclear exchange and recombination of haplotypes in both the USA and Australian *Pca* populations as mechanisms for the introduction of genetic diversity. The limitations of assuming clonal evolutionary history from virulence phenotyping is demonstrated by the detection of either sexual or cryptic genetic recombination in the *Pca* Australian population. Overall, the characterization of intercontinental migration of *Pca* at the haplotype level provides further impetus for molecular monitoring of rust pathogen populations on a global scale.

Eva Henningsen

Introduction

Rust fungi are a diverse group of biotrophic plant pathogens belonging to the

Basidiomycete order Pucciniales (Bauer *et al*., 2006; Aime & McTaggart, 2020). With

over 7,000 species and diverse hosts, these organisms cause many significant

agricultural diseases (Kirk *et al*., 2008). Notably, many rust fungal species such as

Puccinia coronata f. sp. *avenae* (*Pca* - oat crown rust), *P. graminis* f. sp. *tritici* (*Pgt* –

wheat stem rust), *P. striiformis* f. sp. *tritici* (*Pst* – wheat stripe rust), and *P. triticina* (*Pt*

– wheat leaf rust) propagate as dikaryons (two haploid nuclei) while infecting their

cereal hosts, which has complicated the identification of virulence factors and

genomics informed surveillance (Petersen, 1974; Figueroa *et al*., 2020).

The flax – flax rust (*Melampsora lini*) pathosystem was the basis for Flor's pioneering

work in plant pathology establishing the gene-for-gene model (Dodds, 2023), in which

recognition of pathogen avirulence (Avr) effectors by plant resistance (*R*) proteins

triggers signalling cascades and ultimately leads to cell death, an effective defence

strategy against biotrophs (Jones & Dangl, 2006; Dodds & Rathjen, 2010; Chen *et al*.,

2022; Dodds, 2023). However, understanding virulence evolution and host adaptation

in rust populations has been hampered by their complex dikaryotic genome structures.

The development of key molecular and bioinformatics technologies has resolved

challenges posed by the ploidy changes in rust fungi through their complex lifecycles

(Figueroa *et al*., 2022). Long-read sequencing and haplotype-aware assembly

software enabled the first partial dissection of haplotypes to uncover high

heterozygosity between nuclei of *Pca* and *Pst,* which highlighted the importance of

nuclear phasing to create accurate rust genome references (Miller *et al*., 2018;

Schwessinger *et al*., 2018). Further improvements in chromatin contact sequencing

and phasing pipelines for rusts delivered the first chromosome-level and nuclear

phased genomes for *Pgt*, *Pt*, and *Pca* (Li *et al*., 2019; Duan *et al*., 2022; Henningsen

et al., 2022). In *Pgt*, this revealed that whole nuclear exchange without recombination

precipitated the emergence of the devastating wheat stem rust Ug99 lineage (Li *et al*.,

2019), leading to a paradigm shift that rust epidemiology must consider the movement

of entire haplotypes. This has been made possible by the advent of high-fidelity long-

read sequencing technology allowing the assembly of more and larger nuclear-phased

rust genome references (Duan *et al*., 2022). This has been explored recently in *Pt*,
where extensive somatic hybridization and intercontinental migration was uncovered
through comparative genomics (Sperschneider *et al*., 2023).

Although molecular resources for studying rust fungi continue to grow, well-
established areas of research for other plant pathogens remain poorly characterized
in these pathogens. For instance, about 90% of Basidiomycota require different mating
types to mate (heterothallic) (Kües *et al*., 2011), with compatibility controlled by two
mating type loci, *a* and *b*. These loci encode a pheromone and receptor at the *a* (*PR*)
locus and homeodomain transcription factor subunits at the *b* (*HD*) locus (Bölker *et*
al., 1992; Burkhard *et al*., 1990). Heterothallic fungi are further divided into two main
types; bipolar and tetrapolar, in which the *a* and *b* loci are linked and unlinked,
respectively. The presence and location of conserved mating type genes in the rust
species *Pgt*, *Pt*, and *Pst* suggests heterothallic and tetrapolar systems (Cuomo *et al*.,
2017; Holden *et al*., 2023), although the role of these presumed mating type loci has
not been functionally validated in rust fungi (Cuomo *et al*., 2017). It has been proposed
that this system also plays a role in somatic combability for nuclear exchanges to occur
between strains (Flor, 1964).

Oat crown rust disease caused by *Pca* results in significant yield losses worldwide and
is difficult to control due to rapid host adaptation by the pathogen (Nazareno *et al*.,
2018). Molecular evidence shows that sexuality is a clear driver of this rapid virulence
evolution in USA populations and influences population structure in Sweden (Berlin *et*
al., 2018; Miller *et al*., 2020; Hewitt *et al*., 2023). However, asexual *Pca* populations
have also been observed to gain virulence following *R* gene deployment (Nazareno *et*
al., 2018; Figueroa *et al*., 2023). Like for other rusts, key aspects of *Pca* biology and
epidemiology remain unclear.

The role of somatic hybridization in *Pca* virulence evolution cannot be assessed
without haplotype-phased genomes (Li *et al*., 2019; Sperschneider *et al*., 2023). There
is one fully phased, chromosome level assembly (Pca203) and two partially haplotype-
separated reference genomes from USA isolates (12SD80 and 12NC29) (Miller *et al*.,
2018; Henningsen *et al*., 2022). Given the strong genetic differentiation between USA

and South African isolates detected by Hewitt *et al*. (2023), the existing references are

unlikely to capture the genetic variation of *Pca* globally. Thus, there exists ripe

opportunity to address key questions for rust biology in *Pca* by further expanding

global resources for the pathosystem that allow genomic comparisons at the

haplotype-level. This study combines a haplotype-aware pangenome with the

extensive collection of publicly available short-read data for *Pca* to further our

understanding of the biology and epidemiology of *Pca* through the application of

contemporary genomic-based comparative approaches.

Materials and Methods

Pca isolates, amplification, and phenotyping

Australian isolates were collected from 2020 to 2023 across six states and territories

(ACT – Australian Capital Territory; NSW - New South Wales; QLD – Queensland; SA

– South Australia; VIC – Victoria; WA – Western Australia). Isolates from Taiwan (TW)

were collected from the College of Bioresources and Agriculture Experimental Farm

at the National Taiwan University in 2020 and 2021 (Taipei, Taiwan). Oat lines for

phenotyping were sourced from existing CSIRO seed stocks and the Australian Grains

Genebank (AGG) (Henningsen *et al*., 2024).

Differences between infection procedures for reviving field samples, single pustule

purification, and phenotyping are detailed in full in Methods S1. In all infections,

susceptible oat seedlings were grown at 23C° for 16 hours light and 18C° for 8 hours

dark. Plants were treated with 15 mL maleic hydrazide per pot at 9 days of growth

immediately prior to inoculation. Inoculations were conducted using urediniospores in

an oil or talc suspension and infected plants were kept in humidity chambers (90-99%

relative humidity) for two days before removal to growth chambers under the same

conditions as before. For phenotyping, infection types were recorded at 10-11 days

post inoculation (dpi), with the final score chosen to reflect the most prevalent infection

type across the biological replicates of the same differential line.

DNA extraction and sequencing

DNA extractions for Illumina sequencing were performed with the G-Biosciences

OmniPrep Genomic DNA isolation kit using 20-40 mg of rust spores as input. Libraries

were generated using either the Illumina DNA PCR-Free Prep or the IDT Prism library

preparation protocol depending on sample needs. Libraries were sequenced to 15-

30X depth, 150 bp paired end reads with Illumina Novaseq by Azenta or the Australian

Genome Research Facility (AGRF) (Table S1).

Extractions for high molecular weight (HMW) DNA from rust spores were completed

as described previously by Li *et al*. (2019). HMW DNA was used for PacBio HiFi library

preparation and sequencing to 30-50X depth at Azenta Life Sciences (formerly

Genewiz) through Phase Genomics (US isolates) or AGRF (Australian isolates) (Table

S2). Isolates 21ACT116, 20WA94, and 21WA134 were sequenced to 100-200X

coverage due to high yield from a single lane of PacBio Revio.

Spores for Hi-C were prepared as described by Sperschneider *et al*. (2023) and

detailed in Methods S2. Hi-C libraries were prepared at Phase Genomics, Seattle WA

USA and sequenced with Illumina Novaseq by Azenta Life Sciences (formerly

Genewiz) or were prepared and sequenced with Illumina Novaseq at the Ramaciotti

Centre for Genomics, NSW Australia (Methods S2).

Genome assembly, phasing, scaffolding, and annotation

Some contamination was apparent in the high-coverage HiFi read data for 20WA89,

21ACT116, and 21WA134 based on large and highly fragmented initial assemblies,

so these reads were filtered with mash v2.0 (Ondov *et al*., 2019) as described in

Methods S3. Raw reads for 20WA94 and 21WA139 and cleaned reads for 20WA89,

21ACT116, and 21WA134 were filtered on length with a 10 kb cutoff and were then

randomly down-sampled with seqkit (v2.7.0) sample to 30-50X coverage (Shen *et al*.,

2016). PacBio HiFi reads were assembled using Hifiasm with Hi-C integration and

resulting contigs were cleaned, phased, scaffolded, and annotated according to

Sperschneider *et al*. (2023) and described in detail in Methods S4. BUSCO values for

each haplotype were determined using compleasm v0.2.1 (Huang & Li, 2023) with the

odb10 Basidiomycota lineage dataset (Manni *et al*., 2021).

Identification of putative mating type genes and population screening

Protein sequences for *STE3.2.1*, *STE3.2.2*, *STE3.2.3*, *bW-HD1* and *bE-HD2* from *Pgt*

were BLASTed (tblastn) against the Pca203 genome reference with BLAST+ v2.13.0

(Camacho *et al*., 2009; Cuomo *et al*., 2017). Proteins overlapping the best hits were

compared to the *Pgt* alleles and examined for conserved domains known to be present

and intron/exon count before use in searching the other 30 haplotypes (Kües *et al.*,

2011; Cuomo *et al.*, 2017; Methods S4). Protein sequences for *STE3.2*, *bW-HD1*, and

bE-HD2 alleles were aligned with CLUSTALW, and phylogenetic trees were

constructed with the built-in 'raxml-bootstrap' option (Larkin *et al.*, 2007). Protein trees

were rooted on alleles from *Pgt* CRL 75-36-700-3 (*Pgt STE3.2.1* and *Pgt bW-HD1*)

with R package 'ape' and visualized with iTOL (Cuomo *et al.*, 2017; Paradis & Schliep,

2019; Letunic & Bork, 2021).

To screen the *Pca* population for the presence of the *STE3.2* and *HD* alleles, the 3' to

3' DNA sequence of the *HD* locus and 5' to 3' sequences of *STE3.2.2* and *STE3.2.3*

were extracted from each haplotype. These loci were sketched with mash v2.0 (-s

1000; Ondov *et al.*, 2019) and then screened with the Illumina reads for 352 isolates

(Miller *et al.*, 2018, 2020; Henningsen *et al.*, 2022; Hewitt *et al.*, 2023; Ho *et al.*, 2024;

Table S1). Alleles were considered contained at or above 99% *k*-mer identity and 94%

shared *k*-mers.

To compare the repetitive regions on chromosome 9, haplotypes were aligned with

minimap2 v2.22 (-k19 -w19 -m200 -DP -r1000) and visualized with ggplot (Li, 2018;

Wickham, 2016).

Population screening for haplotype containment

The 352 *Pca* isolates with Illumina data (Table S1) were screened against the 32

available *Pca* haplotypes using mash v2.0 (Ondov *et al.*, 2019). These haplotypes

were processed with mash sketch (-s 100000) and mash screen was run comparing

the Illumina data against the haplotype sketch file. Isolates with *k*-mer identity of ≥

99.98 % and shared *k*-mers ≥ 99.65% were considered as likely containing the

screened haplotype. Candidate hybrids were identified for having high identity to only

one haplotype, or as having high identity to two haplotypes which were not contained

within the same reference isolate.

Phylogenetic analysis

Illumina reads for 352 isolates (Table S1) were mapped to the diploid assemblies for

isolates Pca203, 90TX52, 20NSW19, 20QLD86, 20WA94, and 21ACT116 using bwa-

mem2 v2.2.1 (Vasimuddin *et al.*, 2019). Variants were called against the diploid

assembly of Pca203 and individual haplotypes (3, 4, 5, 6, 13, 14, 23, 24, 25, 26), using

freebayes v1.3.6 (--use-best-n-alleles 6) (Garrison & Marth, 2012). Variants were

filtered with vcflib v1.0.1 (-f "QUAL > 20 & QUAL / AO > 10 & SAF > 0 & SAR > 0 &

RPR > 1 & RPL > 1 & AC > 0") and vcftools v0.1.16 (--min-alleles 2 --max-alleles 2 --

max-missing 0.9 --maf 0.05) (Danecek *et al.*, 2011; Garrison *et al.*, 2022). Variants

were converted to PHYLIP format using vcf2phylip (Ortiz, 2019). RAxML v8.2.12 was

used to construct the Maximum Likelihood (ML) tree with 500 bootstraps (-f a -m

GTRCAT -# 500 --no-bfgs) (Stamatakis, 2014). The resulting ML tree was visualized

with iTOL (Letunic & Bork, 2021).

Variants called against the Pca203 complete reference for the 137 Australian isolates

were filtered to include only biallelic single-nucleotide polymorphisms (SNPs). Variants

were converted to PHYLIP format with vcf2phylip (Ortiz, 2019), and heterozygous calls

were converted to missing (N) resulting in 391,118 SNPs. Splitstree CE v.6.2.1-beta

was used to calculate Hamming distances and generate a neighbor net with 301 splits

(Bryant & Huson, 2023; https://github.com/husonlab/splitstree6). The network was

evaluated with the phi test for recombination in Splitstree CE v6.2.1-beta (Bruen *et al.*,

2005).

Comparison of pathotype clustering and phylogenetic tree structures

Infection types were converted to a binary system of 0 (avirulent, infection types 0 to

2) and 1 (virulent, infection types 3 to 4) and clustered with R v4.3.2 hclust() before

conversion to a dendrogram (R Core Team, 2023). The Pca203 ML tree was pruned

to contain only phenotyped Australian isolates with iTOL (Letunic & Bork, 2021) and

was modified to be binary and ultrametric, with R package 'ape' before conversion to

a dendrogram (Paradis & Schliep, 2019). The two dendrograms were compared with

untangle_step_rotate_1side() from 'dendextend', with the phenotype clustering

dendrogram being rotated to find the best structural match to the RAxML-derived

dendrogram (Galili, 2015). The comparison was visualized with

tanglegram(rank_branches = TRUE) from 'dendextend' to visualize topology without

considering branch lengths (Galili, 2015). Finally, the two dendrograms were

compared with cor_bakers_gamma() from 'dendextend' to calculate the correlation

between the tree structures (Galili, 2015).

Haplotype comparisons

Nucmer and DNAdiff from mummer v4.0.0 were run to determine the identity between

all nuclear haplotype pairs (Kurtz *et al.*, 2004; Marçais *et al.*, 2018). Haplotypes were

aligned with D-Genies v1.5.0 and visualized with ggplot2 (Wickham, 2016; Cabanettes

& Klopp, 2018). The cactus-pangenome pipeline from cactus v2.6.6 was run on all

haplotypes and variants were called for each haplotype as the reference (Hickey *et*

al., 2023). The full procedure for determining shared haplotype blocks is detailed in

Methods S5. Briefly, variant counts were binned and a cutoff of <= 50 non-ref SNPs

per 100kb was applied. Adjacent shared regions for each sample-reference pair were

merged, subjected to an ordered bedtools (v2.31.1; Quinlan & Hall, 2010) subtraction,

and regions <= 50 Kb were removed before visualization.

Results

Contextualization of Australian and Taiwanese Pca isolates within the global

population

From 2020 to 2023, 137 Australian *Pca* isolates were collected through community

submissions across six Australian states and territories (ACT = 9, NSW = 36, QLD =

9, WA = 55, SA = 10, VIC = 18), and isolates were recovered from both wild (n=111)

and cultivated oats (n=26) (Table S1). Furthermore, four Taiwanese isolates were

collected from cultivated oat in 2020 and 2021. Whole genome sequence (WGS) data

(Illumina) was generated from genomic DNA from all isolates (42X average genome

coverage; Table S1). We generated a ML phylogenetic tree using the WGS data of all

isolates and including published data of 211 isolates from the USA, South Africa, and

Taiwan (total n=352) (Miller *et al.*, 2018, 2020; Henningsen *et al.*, 2022; Hewitt *et al.*,

2023; Ho *et al.*, 2024). Reads were mapped to both haplotypes of the Pca203 genome

reference (Henningsen *et al.*, 2022) and filtered variants (n=376,646) were used to

construct the ML phylogenetic tree.

A total of 18 lineages were detected in the Australian *Pca* collection, which is

unexpected as the population is proposed to consist of only four asexually-reproducing

(clonal) genetic groups (Fig. **2a**; Park *et al.*, 2022). Although clonality is influential in

Australia as shown by the size and temporal persistence of the largest lineages (L18

– 48 individuals, L1 – 35 individuals), nine lineages consist of only one isolate. Further,

only seven lineages were sampled across multiple years (4 years – L1, L9, L18; 3

years – L2, L11; 2 years – L5, L16), suggesting there are more lineages to be sampled

(Fig. **2a**). Lineage diversity was greatest in WA (n=15), followed by SA (n=5),

NSW/ACT (n=4), VIC (n=3), and QLD (n=2; Fig. **2b**). The five most abundant lineages

(L1, L3, L9, L11, L18) were sampled on both wild and cultivated oat, suggesting that

the populations on these different host types are not separate (Table S1). Only one

lineage was detected from the four Taiwanese isolates, which are clones of isolate

NTU-01.

To assess the robustness of virulence phenotypes as the basis for lineage assignment

within the Australian *Pca* population, 95 of the 137 sequenced *Pca* isolates were

tested on 27 genotypes from the Australian oat crown rust differential set (Henningsen

et al., 2024; Fig. S2; Table S3). Most isolates in lineage L1 (20ACT25, 22NSW08)

displayed broad virulence across the differential lines, while others had virulence to

fewer lines (22QLD110, 22NSW107) (Fig. S2). Lineage L2 had the least virulence,

with 21ACT116 and 23NSW13 only defeating resistance for two lines, Swan (carrying

Pc1) and Pc70 (also known as H547). The remaining lineages (L3-L18) share many

virulence traits. For example, isolates 22WA31 and 22WA54 from lineages L11 and

L5 had nearly identical reactions. Clustering isolates by these phenotypic results is

inconsistent with the phylogenetic relationships (Baker's Gamma = 0.64, Fig. S3).

Thus, virulence phenotypes (pathotypes) are poor predictors of lineage relationships

in the Australian population.

Construction of a haplotype-aware pangenome of Pca

From the phylogenetic analysis, we selected representative *Pca* isolates to generate

haplotype-phased chromosome-level genome references that capture genetic

diversity in Australia (n=9) and USA (n=6) (Fig. **S4a,b**). References were generated

from eight Australian clonal groups (20QLD86 – L1, 21ACT116 – L2, 20WA72 and

20WA95 – L9, 20WA94 – L11, 21WA139 – L13, 20WA89 – L14, 21WA134 – L16,

20NSW19 – L18), with lineage L9 sampled twice (20WA72, 20WA95) as a baseline

for comparing near-identical haplotypes (Fig. **S4a**). We also chose three contemporary

buckthorn-derived isolates (18MNBT34, 18MNBT36, 18MNBT50) from the USA

sexual population (Hewitt *et al*., 2023; Fig. **S4b**) and three isolates from USA clonal

lineages (90TX52, 90AR100, 90MN4B; Miller *et al*., 2018; Fig. **S4a,b**).

Eva Henningsen

Figure 1. Midpoint rooted ML phylogenetic tree constructed by mapping reads from 352 *Puccinia coronata* f. sp. *avenae* (*Pca*) isolates and calling variants against the full Pca203 reference (hap1, hap2, and unplaced contigs). A total of 376,646 biallelic SNPs and 500 bootstraps were used for this analysis. Tree branches are colored by country of origin: AUS = Australia; SA = South Africa; TW = Taiwan; USA = United States of America. Bootstrap values above 80% shown as white circles. Yellow stars indicate isolates chosen for the haplotype atlas and red stars indicate isolates with existing references. Tree scale is mean substitutions per site.

Eva Henningsen

Figure 2. **a)** View of Australian lineages with clones collapsed in the midpoint rooted
ML phylogenetic tree constructed for 352 *Puccinia coronata* f. sp. *avenae* (*Pca*)
isolates based on 376,646 biallelic SNPs called against the full Pca203 reference.
Bootstrap values (500 cycles) above 80% shown as white circles. Tree scale is mean
substitutions per site. Stacked barplots show the number of isolates of each lineage

collected in 2020 (black), 2021 (dark grey), 2022 (light grey), and 2023 (white) across
Australian lineages. Tree branches for Australian lineage L1-L18 are colored by
lineage to match points and pie charts in panel **b)** Distribution of *Pca* isolates and
lineages across Australia. Pie charts are scaled on a logarithmic scale according to
the total number of *Pca* isolates collected in their respective states. Pie chart for
NSW/ACT includes isolates from both regions due to the small area of the ACT relative
to NSW. Colors reflect lineage assignments.

We assembled nuclear phased genome references of 15 isolates following established
computational workflows using hifiasm with incorporated Hi-C chromatin contact data
(Duan *et al*., 2022; Henningsen *et al*., 2022; Sperschneider *et al*., 2023), resulting in
30 complete nuclear haplotypes (hap3-hap32; Fig. **1**, see yellow stars; Fig. **3a**; Table
S2). Nuclearphaser detected only five phase swaps (in three isolates) that required
correction. Four within-haplotype mis-assemblies were also corrected in hap5, hap9,
hap10, and hap19. Scaffolding resulted in 18 chromosomes in each haplotype (chr1
to chr18; Fig. **3b**). An average of 84.44% of *trans* and 96.75% of *cis* and *trans* Hi-C
contacts occurred on chromosomes within the same haplotype which reflects accurate
haplotype separation and nuclear assignment. *Pca* haplotypes were on average 98.4
Mb long with 42.7% repeat coverage and 19,700 gene annotations (Table S4).
Haplotype BUSCO completeness averaged 94.85% with less than 2% duplication,
which is comparable to haplotype-phased assemblies for *Pgt* and *Pt* (Table S4; Li *et*
al., 2019; Duan *et al*., 2022; Sperschneider *et al*., 2023). Haplotypes were consistent
for most genome characteristics (Table S4). Chromosome sizes were likewise
consistent except for chr9, which harbours a large repetitive region of approximately
0.1 to 1.4 Mb (Fig. **3b,c**). Most assemblies break within this locus, so the exact size of
the region is not certain.
Pairwise haplotype comparisons validated the clonal relationship between 20WA72
and 20WA95 from L9, as their matching haplotypes (hap7 ≈ hap22; hap8 ≈ hap21)
had very high sequence identity (99.98%). In contrast, comparison of the two
haplotype sequences within any isolate showed high levels of divergence and
heterozygosity (98.73 - 99.45% identity), with most inter-isolate haplotype
comparisons similarly diverse (98.65 - 99.98% identity). The only exception to this was
the haplotype pair hap5 (20QLD86 from L1) and hap24 (21ACT116 from L2) which

showed high 99.94% sequence identity (99.78-99.86% alignment coverage, 9,228

SNPs) comparable to haplotypes in the clonal 20WA72 and 20WA95 isolates (99.44-

99.97% alignment coverage, 99.98% identity, 1,051-1,701 SNPs; Fig. **3d**). As the

other haplotypes from 20QLD86 (hap6) and 21ACT116 (hap23) were extremely

dissimilar (98.39% identity, 96.17-97.85% alignment coverage, 481,662 SNPs), it is

likely that lineages L1 and L2 are related through somatic hybridization and nuclear

exchange (Fig. **3d**).

Characterization of mating loci in the Pca pangenome and overall population

To better understand the role of putative *a* and *b* mating type loci in *Pca*, we

characterized the *bW-HD1/bE-HD2* and *STE3.2* loci using sequences identified in *Pgt*

(Cuomo *et al*., 2017). Pca203 has one copy of *STE3.2.3* on chr9A and one copy of

STE3.2.2 on chr9B. Similarly, all isolates in the *Pca* pangenome carry one near-

identical copy each of *STE3.2.2* and *STE3.2.3* on opposite haplotypes of chromosome

9 (Fig. S**5a,b**). The *mfa* pheromone precursors were also identified close to these

STE3.2 alleles (Fig. **S5a**). The related gene *STE3.2.1* was invariant and present in a

single copy on chr1 in all haplotypes except in hap11 (90MN4B) which had two copies,

supporting previous findings that it is unlikely to be involved in mating compatibility

(Cuomo *et al*., 2017). The *STE3.2.2* and *STE3.2.3* alleles are adjacent to the large

repetitive region on chr9 which is allele-specific, explaining the unusual chr9 length

distribution (Fig. **3b**, Fig. **S5c,d**). We then characterized the *bW-HD1 bE-HD2* (*HD*)

allele pairs across haplotypes in the pangenome. All isolates contained different *HD*

alleles in each haplotype on chr4 and a total of 13 alleles were recorded (Fig**. S5e,f**).

Using these characterized loci, we screened Illumina short reads from the entire *Pca*

collection (*n*=352) and found that all isolates contain both *STE3.2.2* and *STE3.2.3*

alleles and two different alleles at the *HD* locus (Fig. **S6a**, Table S5, Table S6). In *Pca*

isolates where one or both *HD* alleles were not characterized in the pangenome, short

read mapping still indicates heterozygosity at the *HD* locus (Fig. S7). The observed

genotype frequency for both the *PR* and *HD* loci in the USA sexual population (*n*=88

isolates derived as aecia from the sexual host buckthorn in the UMN nursery) are 0%

homozygous and 100% heterozygous (*PR* locus alleles *n*=2, *HD* locus alleles *n*=11).

This is a significant departure from the expected 50:50 (homozygous:heterozygous)

Eva Henningsen

ratio for the biallelic *PR* locus in Hardy-Weinberg equilibrium (p = 6.55 x 10^{-21}).

Likewise, the *HD* locus genotype ratio is significantly different than expected (12%

homozygous, 88% heterozygous; p = 5.70 x 10^{-4}). This suggests that both loci

contribute to mating type compatibility resulting in exclusive heterozygosity, meaning

that *Pca* likely has a tetrapolar mating system given that the *PR* and *HD* loci are on

separate chromosomes.

Figure 3. **a**) Haplotype numbers assigned to nuclear genotypes of 15 *Puccinia coronata* f. sp. *avenae* (*Pca*) isolates from the USA (purple) and Australia (yellow), in addition to previously published haplotypes from Pca203 (1 = "A"; 2 = "B"). **b**) Boxplots of chromosome lengths for the 32 *Pca* haplotypes with black dots indicating

Eva Henningsen

chromosome size of each haplotype. Box top and bottom boundaries are the upper

and lower quartiles, respectively. Lines within boxes represent the mean. Lines

extending below and above boxes delimit minimum and maximum values. **c**) In

descending order: alignments of chromosome 9 regions containing STE3.2 genes

between hap13 and hap25 (*STE3.2.2*), hap26 and hap27 (*STE3.2.3*), and hap13 and

hap26 containing different *STE3.2* alleles. **d**) Dotplot of genome alignment between

Puccinia coronata f. sp. *avenae* (*Pca*) haplotypes hap5 and hap24 (left), hap8 and

hap21 (middle) and hap6 and hap23 (right). Chromosomes are numbered along the

x- and y-axis. Greyscale indicates percent identity along alignment.

Pca isolates from the USA, Australia, and Taiwan are related through somatic
hybridization and migration

To assess the role of somatic hybridization in the evolution of *Pca* we employed *k*-mer
containment, phylogenetics, and comparative genomics approaches used to identify
somatic hybridization events in other rust species (Li *et al.*, 2019; Sperschneider *et al.*,
2023). In our analysis, *k*-mer identity of ≥ 99.98 % and shared *k*-mers ≥ 99.65%
accurately identified clones of the *Pca* reference isolates. For example, L18 isolate
20NSW20 has high *k*-mer containment for hap3 and hap4 from the L18 reference
20NSW19 (100% *k*-mer identity and ≥ 99.93% shared *k*-mers). As established in *Pgt*
and *Pt*, phylogenetic trees constructed from an isolate's entire genome show clonal
lineages as discrete clades, while single-haplotype trees merge lineages that share
the reference haplotype into a single clade (Li *et al.*, 2019; Sperschneider *et al.*, 2023).
Multiple putative nuclear exchange events were identified across the entire *Pca*
dataset using these criteria (Table S1; Table S7).
The full haplotype comparisons we described earlier are the strongest evidence for
somatic hybridization between L1 and L2 and *k*-mer containment and phylogenetic
tree results also support this hypothesis. L1 isolates had high containment for hap5,
hap6, and hap24 (> 99.98% identity, > 99.75% shared *k*-mers), but not hap23 (<
99.86% identity, < 96.99% shared *k*-mers) (Fig. **4a**; Table S7). Conversely, L2 isolates
had high containment for hap23, hap24, and hap5 (> 99.98% identity, > 99.82%
shared *k*-mers) and low containment for hap6 (< 99.64% identity, < 92.50% shared *k*-
mers) (Fig. **4a**; Table S7). Further, trees constructed from these haplotypes
individually support hybridization between L1 and L2. Phylogenies generated from
hap5 and hap24 both show L1 and L2 isolates as members of a single clade, while
those for hap6 and hap23 show L1 and L2 as discrete clades (Fig. **4b-e**). These
analyses also suggest a somatic hybridization event linking Australia and Taiwan,
implicating direct or indirect migration of *Pca* (Fig. **4f**). The five Taiwanese isolates (L-
TW) have high *k*-mer containment for hap6 (99.99% identity, > 99.84% shared *k*-mers)
but not hap5 (< 99.70% identity, < 93.65% shared *k*-mers) (Fig. **4a**; Table S7). This is
also supported by the phylogenetic trees, as L1 and L-TW isolates comprise separate
clades in the hap5 tree but cluster together in the hap6 tree (Fig. **4b,d**). Since we did

not sample the other haplotype for L-TW in the current pangenome, we cannot infer

the history of haplotype exchange.

The *k*-mer containment and phylogenetic analyses also support the occurrence of

somatic hybridization and nuclear exchange within the USA *Pca* population. Members

of the 90TX52 lineage (L-1990) have high containment for both 90TX52 haplotypes

(100% identity, > 99.92% shared *k*-mers) (Fig. **S8a**; Table S7). Seven clonal isolates

collected in 2017 (L-2017) from the southern USA (LA, TX, FL) contain hap14 (100%

identity, > 99.89% shared *k*-mers), but not hap13 (< 99.85% identity, < 96.65% shared

k-mers) (Fig. **S8a**; Table S7). Phylogenetic trees constructed separately from hap13

and hap14 SNPs further support somatic hybridization between L-1990 and L-2017,

as they appear as one clade in the hap14 phylogeny and separate clades in the hap13

phylogeny (Fig. **S8b,c**). As the second haplotype of the 2017 isolates was not sampled

in the current pangenome, we cannot establish the history of hap14 inheritance (e.g.,

donor in the hybridization event) (Fig. **S8d**).

Mating allele composition of putative hybrid isolates agrees with the postulated nuclear

exchange events, as somatic hybrid isolates have the same *HD* locus allele as the

haplotype they are proposed to contain (Fig. **4b-e**; Fig. **S8b,c**; Table S5). The

assemblies for hap5 and hap24 likewise have the same alleles at both mating loci

(*STE3.2.2* and *bW3bE3*; Fig. S5). The other pangenome haplotypes do not appear to

be related through somatic hybridization events to *Pca* isolates in the existing

collection. However, 90MN14B and 90MN7B were identified as likely clones of

90MN4B (100% identity, > 99.92% shared *k*-mers) for which short-read data is not

available (Table S7). Altogether, our results support the occurrence of somatic

hybridization in the broad *Pca* population.

Figure 4. **a)** Plot of % shared *k*-mers (x-axis) versus % *k*-mer identity (y-axis) for 352

Pca isolates on hap5, hap6, hap23, and hap24. Colors indicate relevant lineages (L1-

L3, TW =Taiwan); light grey points are all other isolates. **b-e)** Maximum likelihood trees

for **b)** hap5 (225,727 SNPs) **c)** hap24 (239,896 SNPs) **d)** hap6 (198,467 SNPs) and

e) hap23 (191,316 SNPs). Tree branches are colored by country of origin: AUS =

Australia; SA = South Africa; TW = Taiwan; USA = United States of America. 500

bootstraps were generated for each tree and bootstraps 80% and higher are shown

as white circles on branches. Collapsed branches are indicated with circles at leaf tips.

Mating type alleles for hybrid lineages are shown as rectangles to the right of

branches. Tree scales are mean substitutions per site. **f)** Proposed relationships

between *Pca* lineages and haplotypes overlaid on a map of Oceania and Asia.

The Australian and USA Pca populations are shaped by genetic recombination

The *k*-mer containment analysis also yielded some results that were incompatible with

the hypothesis of nuclear exchanges. For instance, L18 isolates had high containment

for hap3 and hap4 (L18) and hap25 (L11) (100% identity, > 99.91% shared *k*-mers)

but not hap26 (L11) (< 99.94% identity, < 98.56% shared *k*-mers; Table S7). However,

L11 isolates only had high *k*-mer containment for hap25 and hap26 (100% identity, >

99.95% shared *k*-mers) but neither hap3 nor hap4 (< 99.95% identity, < 98.70%

shared *k*-mers), as would be the case if L11 and L18 were somatic hybrids (Table S7).

Haplotype-specific phylogenetic trees reflected the same relationships, with L18 and

L11 isolates forming a single clade in the hap25 phylogeny and discrete clades in the

hap3, hap4, and hap26 phylogenetic trees (Fig. **S9a-d**). Whole-haplotype alignments

between hap3, hap4, and hap25 clarified this non-reciprocal relationship. Hap25 has

high-identity alignments from hap3 and hap4 in large blocks covering >95% of this

nuclear genome, with 0 to 3 breakpoints between hap3 and hap4 sequences per

chromosome. This is consistent with the frequency of recombination breakpoints in a

single meiotic event reported in other rust fungi (Fig. **S9e**; Anderson *et al*., 2016). As

further confirmation, we obtained variants from a pangenome graph of all haplotypes

and recovered the same recombination breakpoints as before from hap3 and hap4

SNP-sparse regions on hap25 (Fig. **S9f**). This is consistent with hap25 being derived

as a haploid product of meiosis from lineage L18 as part of a sexual cross.

We next explored the contradictory finding that L3 isolates had high containment for

haplotype hap5 (= hap24) as well as hap26 from 20WA94 (> 99.98% identity, >

99.84% shared *k*-mers), despite these all having the *bW3bE3 HD* allele pair (Fig. **4b**;

Fig. S5; Table S5; Table S7). In support of the haplotype containment results, we found

that L3 isolates cluster with L1 and L2 isolates in the hap5 and hap24 phylogenetic

trees but with L11 in the hap26 tree, while these lineages all form discrete clades in

the hap6, hap23, and hap25 phylogenetic trees (Fig. **4c-f**; Fig. **S9a,b**). The hap5/24

and hap26 haplotypes all contain the *bW3bE3 HD* allele, while L3 is heterozygous for

b3/b12 alleles suggesting some recombination involved in the relationships between

these isolates. Alignment of hap5 and hap26 again revealed numerous high-identity

segments covering about 45% of haplotype genome, in a pattern that suggests

recombination (Fig. **S10a**; Table S8). A model to explain these results is that L3 shares

a nucleus (hap5) with lineage L1 and L2 through a nuclear exchange event, while

hap26 in L11 is a meiotic recombinant between the two nuclei (hap5 and an unknown

haplotype) in L3 (Fig. **S10b**). In this scenario, L11 results from a sexual cross between

L3 and L18. The unique sequence amounting to approximately half of the unknown L3

haplotype should also contain *bW12bE12*.

Given the evidence for recombination in the hap25 and hap26 genomes, we evaluated

the entire Australian *Pca* population for recombination by generating a splitstree

network which showed reticulation between most Australian lineages. The *phi* test was

significant ($p < 10^{-10}$), suggesting that recombination has occurred to shape the extant

Australian population. We next characterized haplotype recombination across the

entire *Pca* pangenome using the pangenome graph-derived variants approach

validated above in our parent-F1 comparison. A subset of six USA haplotypes were

assessed first, which showed a pattern of small recombination blocks that reflects

frequent sexual recombination in the USA facilitated by the prevalence of *R. cathartica*

(Fig. **5a**). Further, each haplotype added only 4 to 9% coverage, with hap18 having

approximately 30% coverage by the preceding five USA haplotypes. This suggests

the full diversity of the USA sexual population has not been sampled in the haplotype

pangenome.

Among the Australian haplotypes, hap6 and hap23 are clearly divergent from the

others, while all other haplotypes appear related through recombination (Fig. **5b**).

Hap3 and hap4 have contributed to all recombinant haplotypes, while hap5 and hap26

have contributed to all except hap25. The large size of recombination blocks between

Australian haplotypes relative to those shared between USA haplotypes suggests that

recombination is relatively infrequent. However, Australian haplotypes in L3-L18 are

approximately 75% covered by hap3, hap4, hap5, and hap26, consistent with founding

by four haplotypes. Considering our earlier results with hap5 and hap26, we

hypothesize that the remaining 25% is likely present in L3, which has not been

represented in the pangenome yet. Therefore, Australian lineages L4-L17 may have

been founded by limited recombination between the L3 and L18 lineages (Fig. **6a**).

The divergent hap6 and hap23 haplotypes in L1 and L2 have not contributed to the

L3-L18 recombining population.

Eva Henningsen

Recombination analysis performed on all haplotypes demonstrated that the USA and

Australian haplotypes overall have not had recent recombination, as the total amount

of shared sequence between USA and Australian haplotypes was generally less than

5% (Table S8). The only exceptions were hap6 and hap23, which share approximately

50% and 15% of their sequence with the USA haplotypes, respectively (Fig. **S12b**).

Hap1 and hap2 from Pca203 contribute most to hap6 (31.4%), while the other USA

haplotypes share small regions (<5%) with the remaining hap6 sequence (Fig. **S12b**).

Hap9 and hap14 are divergent from other USA haplotypes, but hap14 and hap6 have

approximately 11% of their sequence in common (Table S8). The USA population is

shaped by frequent recombination between diverse haplotypes, which is illustrated by

hap20 having only 45% of its sequence represented by 11 other USA haplotypes (Fig.

S12b). Historic isolate Pca203 clearly had an influential founding effect on the USA

population, with hap1 and hap2 contributing from 21.3 to 57.6% of the sequence of

other USA haplotypes (Fig. **S12b**; Table S8).

Figure 5. **a)** Shared recombination blocks across six *Puccinia coronata* f. sp. *avenae* (*Pca*) haplotypes identified by identifying 100 kb sequence bins with less than 50 non-reference variants hierarchically across hap11, hap12, hap15, hap16, hap17, and hap18. Shared sequences with hap11 were assigned across the other five haplotypes first, after which the unassigned areas of hap12 were evaluated on the four haplotypes below in the hierarchy, and so on. **b)** Shared haplotype blocks between 15 Australian

Pca haplotypes from founder representatives hap3, hap4, hap5, hap26 and divergent
haplotypes hap6 and hap23 generated in a similar manner to panel **a)** with hap3 being
first in the hierarchy. The haplotype with the shorter name was kept as the
representative in cases where haplotypes are nearly identical, as they show redundant
information in this analysis (hap8, hap7, and hap5 shown; hap21, hap22, and hap24
not shown). Chromosome fill color represents unassigned regions or regions shared
with haplotypes earlier in the hierarchy. Unassigned regions from hap7, hap8, hap27,
hap28, hap29, hap30, hap31, and hap32 are filled white. Table on the right side shows
percent coverage of each haplotype by those listed in each column. *HD* locus alleles
are indicated with fill color of rectangles at the chromosome 4 midpoint. Asterisk (*)
indicates that the haplotype blocks on hap4 from hap3 are not shown for visual clarity.

Figure 6. **a)** Diagram showing the proposed relationship of haplotypes in the Australian *Puccinia coronata* f. sp. *avenae* (*Pca*) population to the two postulated founder lineages L18 and L3. Four colors are assigned to founding haplotypes (brown – hap3, orange – hap4, blue – hap5, green – hap'unknown') and derivative haplotypes are color combinations of their ancestors. **b)** Known distribution of haplotypes in the Australian *Pca* collection in either Western Australia or eastern Australia (Australian Capital Territory, New South Wales, Queensland, South Australia, Victoria). **c)**

586   Recombination between closely related USA *Pca* haplotypes (shades of pink) and

587   hap6 with simplified possible pathways for hap6 migration.

Discussion

In this study we investigated the genetic relationships between newly sequenced Australian and Taiwanese *Pca* isolates and previously evaluated populations from the USA and South Africa (Miller *et al*., 2018, 2020; Hewitt *et al*., 2023). The rich genotypic diversity of this pathogen necessitated the construction of a haplotype-aware pangenome for the species to capture genotypic variability across different geographic areas. We focused on haplotypes from fifteen isolates representing Australian and USA *Pca* lineages as an initial step. This haplotype atlas allowed us to examine mechanisms that contribute to virulence evolution such as migration, somatic hybridization, and sexual recombination in *Pca* (Li *et al*., 2019; Figueroa *et al*., 2020; Sperschneider *et al*., 2023).

The role of somatic hybridization (nuclear exchange) in the evolution of dikaryotic rust fungi was proposed in the 1950-1960's (Flor, 1964; Dodds, 2023) and was investigated in *Pca* under controlled inoculations and conditions (Bartos *et al*., 1969). Unfortunately, most early studies assessing somatic hybridization in cereal rust fungi failed to ensure that isolate cross-contamination did not explain detection of new races/virulence profiles. Indisputable proof of nuclear exchange in rust fungi was found by comparing the first chromosome level assembly of wheat stem rust *Pgt*21-0 and sequences from epidemic isolate *Pgt* Ug99 (Li *et al*., 2019), which showed that one nuclear haplotype of *Pgt*21-0 is nearly identical to one of the haplotypes of *Pgt* Ug99. The power of haplotype resolution to track the migration and nuclear exchanges shaping global rust populations was subsequently demonstrated for *Pt* (Sperschneider *et al*., 2023)*.* This study provides evidence that somatic hybridization is also occurring in *Pca* populations in Australia, USA and possibly Taiwan, supporting that nuclear exchanges are common to rust fungi.

Newly acquired and existing whole-genome sequence short-read data and haplotype atlas also enabled population-level *in silico* characterization of mating type control in *Pca*. In other Basidiomycota such as smuts and mushrooms, mating type has been shown to maintain dikaryotic state, regulate life cycle, and enforce self/non-self recognition during mating (Nieuwenhuis *et al*., 2013; Coelho *et al*., 2017). Although

the role of mating loci has not been functionally characterized in the rusts, our finding

that all 353 *Pca* individuals in this study are heterozygous at both the *PR* and *HD* loci

suggests that mating type controls critical biological functions in *Pca*. Consistent with

the findings of Luo et al. (2024), pheromone precursor *mfa2* is linked closely to

STE3.2.2 in *Pca*; however, we identified a 55 amino acid *mfa3* pheromone precursor

linked to *STE3.2.3* which was not reported in their study. Analysis of the *HD* alleles in

the haplotype pangenome released by our study support most sequences imputed for

Pca HD locus alleles published by Luo et al. (2024). There is one discrepancy between

the *bW5* allele between the studies in which the first 12 N-terminal amino acids of *bW5*

defined by Luo et al. (2024) are not encoded in the haplotypes used to define *bW5* in

this study. In addition, the '*bW8bE8*' allele pair reported by Luo et al. (2024) was only

found in a single isolate (90MN5B) that was not included in our pangenome.

The *HD* locus configuration has been used in *Pt* and *Pst* to support conclusions

regarding migration and population structure derived from genomic and phylogenetic

approaches (Holden *et al.*, 2023; Sperschneider *et al.*, 2023). However, several

divergent haplotypes (~500-600K SNPs) included in the *Pca* pangenome have

identical alleles at both mating type loci (i.e., hap6 and hap23, hap3 and hap13, hap5

and hap17). Thus, *HD* locus allele containment solely should not be used to infer

lineage membership or haplotype composition.

We investigated the contribution of genetic recombination to the evolution of *Pca* in

the USA and Australia. Small and frequent recombination blocks among USA

haplotypes were detected and are consistent with widespread sexual reproduction

enable by the prevalence of *R. cathartica* in North America (Miller *et al.*, 2020; Hewitt

et al., 2023). Prior to this study, molecular characterization of the Australian population

was limited to 12 *Pca* isolates using DNA amplification fingerprinting and virulence

phenotypes (Brake *et al.*, 2001). Based on this, it was proposed the Australian *Pca*

population was evolving by mutations in clonal lineages, despite the high phenotypic

diversity of the pathogen.

We uncovered 18 lineages in the Australian *Pca* population, in contrast to the *Pca*

collections from Taiwan and South Africa, which contain only one or two clonal

lineages. This is a contrasting result to cereal rusts in Australia, like *Pgt* and *Pt*, which

appear to be exclusively clonal, with the presence of only two and five lineages,

respectively (Li *et al*., 2019; Sperschneider *et al*., 2023). The discovery of many

lineages in the Australian *Pca* population was unexpected given its presumed

asexuality (Park *et al*., 2022) and is evidence against mutations and incursions as the

sole drivers of *Pca* evolution in the country. The best-described sexual host (*R.*

cathartica) is not present in Australia or Oceania; however, other *Rhamnus* species

(e.g., *R. alaternus, R. lycioides*, *R. palaestina*) have been reported as aecial hosts for

Pca (Dinoor, 1962; Hemmami *et al*., 2006; Nazareno *et al*., 2018). Notably, *R.*

alaternus was introduced to Australia as an ornamental and is currently managed as

an invasive weed (Stajsic & VicFlora, 2023)*.*

Both network analysis and whole-genome comparisons to define recombination

blocks provided evidence of recombination among most Australian lineages, although

infrequent when compared to the USA *Pca* population. It is yet unclear where or when

these events occurred, and their rarity could be explained by sporadic access to a

sexual host or uncommon parasexual events. Parasexuality akin to somatic

hybridization that involves karyogamy and haploidization to generate recombinants

without the aecial host is also proposed to occur in rust fungi (Harder, 1984), although

this phenomenon has not been experimentally validated using molecular and genomic

tools. The use of field collections hampers validation of parasexuality as isolates

resulting from either somatic recombination or a sexual cross would be genotypically

indistinguishable. The recombination among Australian lineages likely explains the

high phenotypic diversity of *Pca* in Australia (Park *et al*., 2022), which was poorly

correlated to lineage structure as has been reported in other *Pca* populations (Hewitt

et al., 2023).

The splitstree network showed clear reticulation between lineages L3 through L18.

Interestingly L3 and L18, which are positioned at opposite ends of the network, are

also implicated in recombination events connected to L11. Hap25 is likely the meiotic

product of hap3 and hap4, while hap26 may be a recombinant derived from L3

haplotypes (hap5 and unknown). Therefore, it seems that L3 and L18 could have

undertaken a sexual cross to produce L11. While intercontinental migrations could

explain these results, the likelihood of F1 haplotypes migrating together or in separate
events is much lower than the hypothesis that rare sexual/parasexual cycles may
occur. Together with the evidence of somatic hybridization in L1 and L2, we propose
that the extant Australian *Pca* population was derived from three ancestral isolates.
Two of these have undergone sexual and/or parasexual processes and the third
donated nuclei via somatic nuclear exchange.
693
Our results suggest that it would be prudent to continue molecular monitoring of *Pca*
and investigate the relationship between *R. alaternus* or other *Rhamnus* species and
Pca in the Australian context, as the presence of these invasive species may present
significant risk for virulence evolution to the detriment of the oat industry. Ongoing
efforts to expand this haplotype-aware pangenome to include members from other
continents will further our understanding of the evolution and global movement of *Pca*.
700

Eva Henningsen

Acknowledgements

We would like to thank Jakob Riddle (USDA-ARS) for his technical support. We would

like to acknowledge the contribution of the Plant Pathogen 'Omics Initiative consortium

in the generation of data used in this publication. The Initiative is supported by funding

from Bioplatforms Australia, enabled by the Commonwealth Government National

Collaborative Research Infrastructure Strategy (NCRIS).

We also thank rust sample donors, with special thanks to Allan Rattey.

Competing Interests

The authors declare no competing interests.

Author Contributions

MF, PD, SFK, JS, BJS, and ES planned and supported the project. DL, ECH, Y-FH,

and ESN amplified and prepared pathogen materials for sequencing and phenotyping.

ECH and JS analysed the data. ECH, JS, MF, and PD prepared the manuscript and

figures. All authors reviewed and edited the manuscript.

Data Availability

Reference genomes are available in the CSIRO Data Access Portal

(https://data.csiro.au/collection/csiro:61932). PacBio, HiC and genomic short reads

are available on NCBI under BioProject PRJNA1063754. Scripts used to perform

analyses and generate figures are available on GitHub

(https://github.com/henni164/Pca_pangenome).

References

Aime MC, McTaggart AR. 2020. A higher-rank classification for rust fungi, with notes on genera. *Fungal Sys. Evol.* 7: 21-47.

Anderson C, Khan MA, Catanzariti AM, Jack CA, Nemri A, Lawrence GJ, Upadhyaya NM, Hardham AR, Ellis JG, Dodds PN, et al. 2016. Genome analysis and avirulence gene cloning using a high-density RADseq linkage map of the flax rust fungus, *Melampsora lini. BMC Genomics*. 17: 667.

Bartos P, Fleischmann G, Samborski DJ, Shipton WA. 1969. Studies on asexual variation in the virulence of oat crown rust, *Puccinia coronata* f. sp. *avenae*, and wheat leaf rust, *Puccinia recondita. Can. J. Botany.* 47: 1383-1387.

Bauer R, Begerow D, Sampaio JP, Weiß M, Oberwinkler F. 2006. The simple-septate basidiomycetes: a synopsis. *Mycol. Prog.* 5: 41-66.

Berlin A, Wallenhammar A-C, Andersson B. 2018. Population differentiation of *Puccinia coronata* between hosts – implications for the epidemiology of oat crown rust. *Eur. J. Plant Pathol*. 152: 901–907.

Bölker M, Urban M, Kahmann R. 1992. The *a* mating type locus of *U. maydis* specifies cell signaling components. *Cell*. 68: 441-450.

Brake VM, Irwin JAG, Park RF. 2001. Genetic variability in Australian isolates of *Puccinia coronata* f. sp. *avenae* assessed with molecular and pathogenicity markers*. Australas. Plant Path*. 30: 259-266.

Bruen TC, Philippe H, Bryant D. 2005. A simple and robust statistical test for detecting the presence of recombination. *Genetics.* 17: 2665-81.

Bryant D, Huson DH. 2023. NeighborNet: improved algorithms and implementation. *Front. Bioinform*. 3: 1178600.

Burkhard S, Banuet F, Dahl M, Schlesinger R, Schäfer W, Martin T, Herskowitz I, Kahmann R. 1990. The *b* alleles of *U. maydis*, whose combinations program pathogenic development, code for polypeptides containing a homeodomain-related motif. *Cell*. 60: 295-306.

Cabanettes F, Klopp C. 2018. D-GENIES: dot plot large genomes in an interactive, efficient and simple way T. Tullius, ed. *PeerJ*. 6: e4958.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden T. L. 2009. BLAST+: architecture and applications. *BMC Bioinformatics*. 10: 421.

Chen J, Zhang X, Rathjen JP, Dodds PN. 2022. Direct recognition of pathogen
effectors by plant NLR immune receptors and downstream signalling. *Essays*
Biochem. 66: 471–483.

Coelho MA, Bakkeren G, Sun S, Hood ME, Giraud T. 2017. Fungal sex: the
Basidiomycota. *Microbiol. Spectr.* 5. 10.1128/microbiolspec.funk-0046-2016.

Cuomo CA, Bakkeren G, Khalil HB, Panwar V, Joly D, Linning R, Sakthikumar S,
Song X, Adiconis X, Fan L, et al. 2017. Comparative analysis highlights variable
genome content of wheat rusts and divergence of the mating loci. *G3*. 7: 361–376.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE,
Lunter G, Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools.
Bioinformatics. 27: 2156–2158.

Dinoor A. 1962. Note on germination of crown rust teliospores and infection of
buckthorn. *Can. J. Plant Sci*. 42: 537-538.

Dodds PN. 2023. From gene-for-gene to resistosomes: Flor's enduring legacy. *MPMI*.
36: 461-467.

Dodds PN, Rathjen JP. 2010. Plant immunity: towards an integrated view of plant–
pathogen interactions. *Nat Rev Genet*. 11: 539–548.

Duan H, Jones AW, Hewitt T, Mackenzie A, Hu Y, Sharp A, Lewis D, Mago R,
Upadhyaya NM, Rathjen JP, et al. 2022. Physical separation of haplotypes in
dikaryons allows benchmarking of phasing accuracy in Nanopore and HiFi assemblies
with Hi-C data. *BMC Genome Biol*. 23: 84.

Figueroa M, Dodds PN, Henningsen EC. 2020. Evolution of virulence in rust fungi –
multiple solutions to one problem. *Curr. Opin. Plant Biol*. 56: 20-27.

Figueroa M, Dodds PN, Henningsen EC, Sperschneider J. 2023. Global landscape of
rust epidemics by *Puccinia* species: current and future perspectives. In: Scott B,
Mesarich C, eds. *Plant Relationships: Fungal-Plant Interactions*. Springer, 183-203.

Flor HH. 1964. Genetics of somatic variation for pathogenicity in *Melampsora lini*.
Phytopathol. 54: 823-826.

Galili T. 2015. dendextend: an R package for visualizing, adjusting, and comparing
trees of hierarchical clustering. *Bioinformatics*. 31: 3718-3720.

Garrison E, Kronenberg ZN, Dawson ET, Pedersen BS, Prins P. 2022. A spectrum of
free software tools for processing the VCF variant call format: vcflib, bio-vcf, cyvcf2,
hts-nim and slivar. *PLoS Comput. Biol*. 18: e1009123.

Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read
sequencing. *arXiv*. doi: 10.48550/arXiv.1207.3907.

Harder DE. 1984. Developmental Ultrastructure of Hyphae and Spores. In: Bushnell
WR, Roelfs AP, eds. *The Cereal Rusts Volume I: Origins, specificity, structure, and*
physiology. Orlando, Florida: Academic Press.

Hemmami I, Allagui MB, Chakroun M, Gazzah ML. 2006. *Rhamnus lycoides* in Tunisia
is a new aecial host of oat crown rust. *Eur. J. Plant Pathol*. 115: 357-361.

Henningsen EC, Hewitt T, Dugyala S, Nazareno ES, Gilbert E, Li F, Kianian SF,
Steffenson BJ, Dodds PN, Sperschneider J, et al. 2022. A chromosome-level, fully
phased genome assembly of the oat crown rust fungus *Puccinia coronata* f. sp.
avenae: a resource to enable comparative genomics in the cereal rusts. *G3*. 12:
jkac149.

Henningsen EC, Lewis D, Nguyen DT, Sperschneider J, Kianian SF, Stone E, Dodds
PN, Figueroa M. 2024. Virulence patterns of oat crown rust in Australia – season 2022.
bioRxiv. doi: 10.1101/2023.09.24.558528.

Hewitt T, Henningsen EC, Pereira D, McElroy K, Nazareno ES, Dugyala S, Nguyen-
Phuc H, Li F, Miller ME, et al. 2023. Genome-enabled analysis of population
dynamics and virulence associated loci in the oat crown rust fungus *Puccinia*
coronata f. sp. *avenae*. *MPMI*. doi: 10.1094/MPMI-09-23-0126-FI.

Hickey G, Monlong J, Ebler J, Novac AM, Eizenga JM, Gao Y, Human Pangenome
Reference Consortium, Marschall T, Li H, Paten B. 2023. Pangenome graph
construction from genome alignments with Minigraph-Cactus. *Nat. Biotechnol*. doi:
10.1038/s41587-023-01793-w.

Holden S, Bakkeren G, Hubensky J, Bamrah R, Abbasi M, Qutob D, de Graaf ML,
Kim SH, Kutcher HR, McCallum BD, et al. 2023. Uncovering the history of
recombination and population structure in western Canadian stripe rust populations
through mating-type alleles. *BMC Biology*. 21: 233.

Ho C-Y, Henningsen EC, Chen S-T, Ariyawansa HA, Nazareno ES, Sperschneider
J, Dodds PN, Riddle J, Kianian SF, Figueroa M, et al. 2024. Detection of oat crown
rust disease in Taiwan (2019-2021). *bioRxiv*. doi: 10.1101/2024.03.12.584714.

Huang N, Li H. 2023. compleasm: a faster and more accurate reimplementation of
BUSCO. *Bioinformatics*. 39: btad595.

Jones JDG, Dangl JL. 2006. The plant immune system. *Nature*. 444: 323–329.

Kirk P, Cannon P, Minter D, Stalpers J. 2008. *Dictionary of the Fungi.* 10th ed.

Wallingford, UK: CABI, eISBN:978-1-84593-728-7.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL.

2004. Versatile and open software for comparing large genomes. *Genome Biol*. 5:

R12.

Kües U, James TY, Heitman J. 2011. Mating type in Basidiomycetes: Unipolar, bipolar,

and tetrapolar patterns of sexuality. In: Pöggeler S, Wöstemeyer J, eds. *Evolution of*

Fungi and Fungal-Like Organisms. The Mycota, vol. 14. Berlin, Heidelberg: Springer.

https://doi.org/10.1007/978-3-642-19974-5_6.

Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H,

Valentin F, Wallace IM, Wilm A, Lopez R, et al. 2007. Clustal W and Clustal X version

2.0. *Bioinformatics*. 23: 2947-2948.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for

phylogenetic tree display and annotation. *Nucleic Acids Res*. 49: W293–W296.

Li F, Upadhyaya NM, Sperschneider J, Matny O, Nguyen-Phuc H, Mago R, Raley C,

Miller ME, Silverstein KAT, Henningsen EC, et al. 2019. Emergence of the Ug99

lineage of the wheat stem rust pathogen through somatic hybridisation*. Nat. Commun*.

10: 5068.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*.

34: 3094–3100.

Luo Z, McTaggart A, Schwessinger B. 2024. Genome biology and evolution of mating-

type loci in four cereal rust fungi. *PLoS Genet.* 20: e1011207.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO update:

novel and streamlined workflows along with broader and deeper phylogenetic

coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Mol. Biol. Evol*. 38:

4647-4654.

Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018.

MUMmer4: A fast and versatile genome alignment system. *PLoS Comp. Biol*. 14:

e1005944.

Miller ME, Zhang Y, Omidvar V, Sperschneider J, Schwessinger B, Raley C, Palmer

JM, Garnica D, Upadhyaya N, Rathjen J, et al. 2018. *De novo* assembly and phasing

of dikaryotic genomes from two isolates of *Puccinia coronata* f. sp. *avenae*, the causal

agent of oat crown rust. *mBio*. 9: 10.1128/mbio.01650-17.

Miller ME, Nazareno ES, Rottschaefer SM, Riddle J, Dos Santos Pereira D, Li F,

Nguyen-Phuc H, Henningsen EC, Persoons A, Saunders DGO, et al. 2020. Increased

virulence of *Puccinia coronata* f. sp. *avenae* populations through allele frequency

changes at multiple putative *Avr* loci. *PLoS Genet*. 16: e1009291.

Nazareno ES, Li F, Smith M, Park RF, Kianian SF, Figueroa M. 2018. *Puccinia*

coronata f. sp. *avenae*: a threat to global oat production. *Mol. Plant Pathol*. 19: 1047–

1060.

Nieuwenhuis BPS, Billiard S, Vuilleumier S, Petit E, Hood ME, Giraud T. 2013.

Evolution of uni- and bifactorial sexual compatibility systems in fungi. *Heredity.* 111:

445-455.

Ondov BD, Starrett GJ, Sappington A, Kostic A, Koren S, Buck CB, Phillippy AM. 2019.

Mash Screen: high-throughput sequence containment estimation for genome

discovery. *Genome Biol*. 20: 232.

Ortiz EM. 2019. vcf2phylip v2.0: convert a VCF matrix into several matrix formats for

phylogenetic analysis. doi: 10.5281/zenodo.2540861.

Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics*. 35: 526-528.

Park RF, Chhetri M, Singh D, Ding Y. 2022. *Cereal rust situation, August 2022*.

Cereal Rust Report 19. Plant Breeding Institute, University of Sydney. URL

https://www.sydney.edu.au/content/dam/corporate/documents/faculty-of-

science/research/life-and-environmental-sciences/rust-reports/cereal-rust-report-

2022-vol-19-2-revised-updated.pdf.

Petersen RH. 1974. The rust fungus life cycle. *Bot. Rev*. 40: 453-513.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics*. 26: 841–842.

R Core Team. 2023. *R: A language and environment for statistical computing v4.3.2*.

R Foundation for Statistical Computing, Vienna, Austria.

Schwessinger B, Sperschneider J, Cuddy WS, Garnica DP, Miller ME, Taylor JM,

Dodds PN, Figueroa M, Park RF, Rathjen JP. 2018. A near-complete haplotype-

phased genome of the dikaryotic wheat stripe rust fungus *Puccnia striiformis* f. sp.

tritici reveals high interhaplotype diversity. *mBio*. 9: 10.1128/mbio.02275-17.

Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A cross-platform and ultrafast toolkit for

FASTA/Q file manipulation. *PLoS One*. 11: e0163962.

Sperschneider J, Hewitt T, Lewis DC, Periyannan S, Milgate AW, Hickey LT, Mago R,

Dodds PN, Figueroa M. 2023. Nuclear exchange generates population diversity in the

wheat leaf rust pathogen *Puccinia triticina*. *Nat. Microbiol.* 8: 2130-2141.

Stajsic T, VicFlora. 2023. *VicFlora: Rhamnus alaternus*. Royal Botanic Gardens

Victoria. [WWW document] URL: https://vicflora.rbg.vic.gov.au/flora/taxon/1e3789d3-

489c-4071-ae6e-05f722c10a18. [accessed 6 March 2024].

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. *Bioinformatics*. 30: 1312–1313.

Vasimuddin M, Misra S, Li H, Aluru S. 2019. Efficient architecture-aware acceleration

of BWA-MEM for multicore systems. In: *2019 IEEE International Parallel and*

Distributed Processing Symposium (IPDPS). Rio de Janeiro, Brazil: IEEE, 314-324.

Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New

York. https://ggplot2-book.org/.

Supplementary Figures

Figure S1. Examples of three main unplaced contig types: **a**) extra copies of chromosome 9 repeat-rich region **b**) extra copies of chromosome 17 ribosomal repeats **c**) extra copies of other genome sequences which are already represented in chromosomes.

Eva Henningsen

Figure S2. Phylogenetic tree constructed by mapping reads from 352 *Puccinia*
coronata f. sp. *avenae* isolates to Pca203 hap1, and hap2, and unplaced contigs.
376,646 biallelic SNPs and 500 bootstraps were used to produce the maximum
likelihood phylogenetic tree. Tree branches are colored by country of origin: AUS =
Australia; SA = South Africa; TW = Taiwan; USA = United States of America. Bootstrap
values are percentages (100 = 100%). Tree scale is mean substitutions per site.
Heatmap represents isolate virulence on the differential lines (virulent = red, avirulent
= yellow). Isolates that were not phenotyped were pruned from the tree, resulting in
the removal of L7 and L17.

Eva Henningsen

Pathotype clustering   **Genetic phylogeny**

Figure S3. Comparison between tree topologies of 16 Australian *Puccinia coronata* f. sp. *avenae* lineages from clustering by pathotype (left) versus phylogenetic relationships (right). The ML tree (right) was generated with 376,646 SNPs from 352 *Pca* isolates against the complete Pca203 genome (hap1, and hap2, and unplaced contigs), which was pruned to contain only phenotyped Australian isolates and midpoint rooted. The R package 'tanglegram' was used to rotate the pathotype clustering tree branches until the best match to the phylogenetic tree was found. Branch lengths are arbitrary in the visualization and assessment of similarity. Solid black lines in the tree structure indicate edges found in both trees. Lines in the center

Eva Henningsen

929   connect the same isolate across trees, with colored lines showing clusters containing

930   more than one isolate which are structurally identical.

Figure S4. Midpoint rooted maximum likelihood phylogenetic tree of 352 *Puccinia coronata* f. sp. *avenae* (*Pca*) isolates constructed by mapping reads and calling variants against the full Pca203 reference (hap1, hap2, and unplaced contigs). 376,646 biallelic SNPs and 500 bootstraps were used. **a)** Collapsed tree view of primarily Australian lineages (L1 to L18). Except for Pca203, the first two digits of the name of the *Pca* isolate reflect year of collection, followed by state and sample identifier. **b)** Collapsed tree view showing USA lineages. USA *Pca* lineages were not numbered as the population is highly diverse. Tree branches are colored by country of origin: AUS = Australia; SA = South Africa; TW = Taiwan; USA = United States of America. Bootstrap values above 80% shown as white circles. Yellow labels indicate isolates chosen for the haplotype atlas and red labels indicate isolates with published references. Tree scales are mean substitutions per site.

Figure S5. **a**) Orientation and arrangement of *STE3.2* and *mfa* (*PR*) alleles on chromosome 9 (chr9) and *HD* alleles on chromosome 4 (chr4) of hap25 and hap26 from *Puccinia coronata* f. sp. *avenae* (*Pca*) isolate 20WA94. **b)** Phylogenetic tree of *STE3.2* alleles in 32 *Pca* haplotypes, rooted at *Puccinia graminis* f. sp. *tritici* (*Pgt*)

STE3.2.1. **c**) Diagram showing chromatin contact maps, and gene and annotations

around the *STE3.2* locus (approximate position shown by labelled red vertical lines)

on chromosome 9 in hap25 and hap26. Repetitive sequences adjacent to the *STE3.2*

alleles correspond to multimapping reads resulting in loss of signal in the contact

maps. **d**) Boxplots showing chromosome 9 length distribution separated by which

STE3.2 allele is present. **e**) Phylogenetic tree of *bW-HD1* alleles rooted at *Pgt bW1-*

HD1 and **f)** *bE-HD2* alleles rooted at *Pgt bE1-HD2* in 32 *Pca* haplotypes. Colors

indicate identical *HD* alleles within each tree and allele pairs across trees (i.e. *bW3*

and *bE3* are always found in the same haplotype), as recombination between *bW* and

bE was not detected in our haplotype atlas**.** Branches with bootstraps over 80% are

shown with white circles at the midpoint. Tree scales are mean substitutions per site.

Figure S6. **A)** Midpoint rooted phylogenetic tree constructed by mapping short reads from 352 *Puccinia coronata* f. sp. *avenae* isolates and calling variants against the complete Pca203 genome (hap1, hap2, unplaced contigs). 376,646 biallelic SNPs were used over 500 bootstraps to produce the maximum likelihood tree. Bootstrap

Eva Henningsen

965 values are percentages (100 = 100%). **b)** Collapsed view showing Australian lineages

966 L1-L18. *HD* alleles are shown as rectangles next to branches. Tree branches are

967 colored by country of origin: AUS = Australia; SA = South Africa; TW = Taiwan; USA

968 = United States of America. Tree scales are mean substitutions per site.

Figure S7. Variants called for 49 *Puccinia coronata* f. sp. *avenae* isolates with one or two unknown *HD* locus alleles in the *HD* locus regions of Pca203 hap1 (*bW1bE1*) and hap2 (*bW2bE2*). Line colors indicate missing (white), reference (black) and alternative

Eva Henningsen

(pink, green, blue, orange, yellow) genotypes. Half-length lines indicate heterozygous

sites and full-length lines indicate homozygous sites.

Figure S8. **a)** Plot of % *k*-mer identity (y-axis) versus % shared *k*-mers (x-axis) for 352

Puccinia coronata f. sp. *avenae* isolates on hap13 and hap14. Colors indicate relevant

lineages; light grey points are all other isolates. **b-c)** Maximum likelihood phylogenetic

trees constructed from variants from individual haplotypes **b**) hap13 (194,540 SNPs),

c) hap14 (209,542 SNPs) with percentages for 500 bootstraps shown on branches.

Tree branches are colored by country of origin: AUS = Australia; SA = South Africa;

TW = Taiwan; USA = United States of America. Collapsed clades are indicated by

circles at tips. *HD* alleles are shown as colored rectangles next to relevant clonal

lineages. Tree scales are mean substitutions per site. **d)** Diagram of the proposed

relationship between USA lineages and haplotypes involved in somatic hybridization.

Figure S9. Maximum likelihood phylogenetic trees for 352 *Puccinia coronata* f. sp. *avenae* isolates constructed from variants from individual haplotypes **a**) hap25 (205,993 SNPs) **b**) hap26 (188,803 SNPs) **c**) hap3 (228,300 SNPs) **d**) hap4 (226,617 SNPs). Bootstrap supports 80% or higher are shown as circles at branch midpoints. Tree branches are colored by country of origin: AUS = Australia; SA = South Africa; TW = Taiwan; USA = United States of America. Collapsed branches are indicated by

Eva Henningsen

circles at tips. *HD* alleles are shown as colored rectangles next to relevant clonal

lineages. Tree scales are mean substitutions per site. **e)** Genome alignment between

20NSW19 haplotypes (hap3, hap4) and hap25 from 20WA94. Chromosomes for the

three haplotypes are shown with high-identity alignments shown with colored fill (hap3

= pink, hap4 = orange). **f)** Non-reference variant counts within 100 kb bins from hap3

(pink) and hap4 (orange) on hap25 are shown in the histogram above hap25

chromosomes. Hap25 chromosome fill color is determined by identifying bins with a

low density (<50 SNPs/100 kb) of non-reference variants from hap3 (pink) and hap4

(orange).

a

b

Figure S10. **a)** Regions of 20QLD86 hap5 against 20WA94 hap26 with over 95% identity when aligned are shown in dark blue. Unique sequences are shown in light blue (hap5) and green (hap26). **b)** Considering the mating type composition and shared sequences between hap5 and hap26, it is possible that hap26 from L11 is the progeny of hap5 and an unknown haplotype from L3. L3 may have been founded by somatic hybridization with L1 or L2.

Eva Henningsen

1009

1010 Figure S11. Splitstree network of 137 Australian *Puccinia coronata* f. sp. *avenae*
1011 isolates from 18 lineages generated from 391,118 SNPs from the entire Pca203
1012 genome. Heterozygous sites were converted to missing. Scale indicates nucleotide
1013 substitutions per site.

Eva Henningsen

Figure S12. **a)** Shared haplotype blocks across six *Puccinia coronata* f. sp. *avenae*

(*Pca*) haplotypes from 100 kb bins with less than 50 non-reference variants which were

assigned hierarchically across hap3, hap4, hap5, hap26, hap9, and hap14. Shared

sequences with hap3 were assigned across the other five haplotypes first, after which

the unassigned areas of hap4 were evaluated on the four haplotypes below in the

hierarchy, and so on. Regions in hap14 were not assigned to other haplotypes, so the

unassigned hap14 sequence has a white fill. **b)** Generated the same way as panel **a)**,

Eva Henningsen

except shared haplotype blocks are assigned across 14 *Pca* haplotypes (hap1, hap2,

hap10, hap11, hap12, hap13, hap15, hap16, hap17, hap18, hap19, hap20, hap6,

hap23) with hap1 being the first in the hierarchy. Regions from hap6 were not assigned

across hap23, so both haplotypes have unassigned sequences filled in white.

Chromosome fill color represents unassigned regions or regions shared with

haplotypes earlier in the hierarchy. Percent coverage by preceding haplotypes is

shown on the right side. *HD* locus alleles indicated by fill color of rectangles at the

chromosome 4 midpoint.