1 **Title:** Deciphering the Timing and Impact of Life-extending Drugs: A Novel Analytic Approach
2 that Differentiates Early, Midlife, and Senescence Phase Efficacies
3

4 **Authors:** Nisi Jiang[1,2]†, Catherine J. Cheng[1,2]†, Qianqian Liu[3]†, Randy Strong[1,4,5], Jonathan
5 Gelfond[1,3]*, James F. Nelson[1,2]*
6
Affiliations:
[1]The Sam and Ann Barshop Institute for Longevity and Aging Studies, UT Health San Antonio;
San Antonio, TX, U.S.A.
[2]Department of Cellular and Integrative Physiology, UT Health San Antonio; San Antonio, TX,
U.S.A.
[3]Department of Population Health Sciences, UT Health San Antonio; San Antonio, TX, U.S.A.
[4]Department of Pharmacology, UT Health San Antonio; San Antonio, TX, U.S.A.
[5]South Texas Veterans Health Care System, San Antonio, TX, U.S.A.
*Corresponding authors. Emails: nelsonj@uthscsa.edu; gelfondjal@uthscsa.edu.
† These authors contributed equally to this work.

17
18 **Abstract:**
19

Evidence that life-extending interventions are not uniformly effective across the lifespan calls for
an analytic tool that can estimate age-specific treatment effects on mortality hazards. Here we
report such a tool, applying it to mouse data from 42 agents tested in the NIA Interventions
Testing Program. This tool identified agents that either reduced (22) or increased (16) mortality
hazards or did both (6), all with marked variation in the duration of efficacy and magnitude of
effect size. Only 7 reduced mortality hazards after the 90% mortality, when the burden of
senescence is greatest. Sex differences were apparent in all parameters. This new analytic tool
complements the commonly used log-rank test. It detects more potential life-extending
candidates (22 versus 10) and indicates when during the life course they are effective. It also
uncovers adverse effects. Most importantly, it identifies agents that specifically reduce mortality
hazards during the senescent phase of life.

31
32 **Keywords:** mortality hazard, life-extending interventions, statistical analysis, mice, longevity
33

Introduction:

The search for pharmacological interventions that extend the healthy lifespan has increased markedly in recent years, spurred by the discovery of a wide range of compounds, such as rapamycin and acarbose, that lengthen life of model organisms[1-3]. Whether these life-extending agents act broadly by extending survival throughout the lifespan or only affect survival during part of the life course remains unclear, in part due to the inadequacy of statistical tests used in interventional research. The log-rank test[4], widely employed in clinical trials, is the statistical tool most commonly used in aging research to determine whether an intervention, be it pharmacologic, genetic, or nutritional, is life-extending. However, its use as the primary and often only tool for this purpose is questionable for several reasons. First is its requirement for proportional hazards between compared groups, implying that treatment effects on mortality remain constant over time[5]. This assumption does not align with the evidence. Many aging interventions exert varying impacts at different life stages. For example, in an earlier analysis of data from the Interventions Testing Program (ITP), we found that many tested interventions do not adhere to the PH assumption, challenging the applicability of the log-rank test in these contexts[6]. For these interventions, we utilized the Gehan test, more robust to the PH consistency requirement and more sensitive to the effects during early adulthood. Notably, it identified five new life-extending candidates[6]. Despite its strengths, the Gehan test has its own drawback: a diminished sensitivity to effects manifesting at later life stages[7].

To assess the effects of interventions on the final phase of the aging process, methods like the Wang-Allison test have been developed to determine if treatments extend the "maximum" lifespan[8]. However, these approaches predominantly assess cumulative survival rates[9]. They do not evaluate whether an intervention specifically reduces age-specific mortality in the last phase of life when frailty, cognitive impairment, chronic disease, and other burdens of senescence peak. Although the Gompertz model has been used for evaluating age-specific or time-varying effects, it is limited by its strict parametric assumptions about the shape of the hazard function[10]. The limitations of these approaches underscore the need for a flexible tool for evaluating longevity interventions, one that accommodates the variable impacts of treatments across an organism's lifespan. Such methods should pinpoint when, for how long, and to what extent an intervention significantly alters the mortality risk. This capability is particularly crucial for identifying interventions that mitigate mortality toward the end of life when the exponential increase in the burden of senescence is greatest.

Here we introduce a novel analytic tool, a time-varying hazard ratio analysis, that detects age-specific drug effects on the mortality hazard with notable precision, thus addressing the major limitation of the log-rank test. For this investigation, we utilized publicly available data from the ITP up to 2022, comprising 42 drugs evaluated in over 27,000 genetically heterogeneous mice at 3 geographically distinct sites[11]. These agents were tested alone or in combination in 132 trials, examining the effects of sex, dosage, and age of treatment initiation. Ten of these agents have been identified by log-rank testing to significantly extend lifespan in at least one sex[12]. This is the largest compendium of mouse survival data from tests of compounds with lifespan-extension potential, an exemplary resource for testing the efficacy of our analytic tool.

Results

<u>A new analytic tool to determine the timing and impact of life-extending candidates</u>.

Figure 1 illustrates how the application of the time-varying hazard analysis identifies age-specific effects of an intervention on the mortality hazard, using the ITP test of green tea extract

85   (GTE) in females as an example. Details of the analysis are described in "Online Methods". It
86   should be noted that GTE had no effect on survival by log-rank testing[13]. Figure 1A shows the
87   Kaplan-Meier survival plots for treatment and control groups. These plots indicate that the
88   proportional hazard assumption is likely violated due to the crossing survival curves, which was
89   confirmed by the z-test (Table S1).
90
91   Figure 1B is a graphical representation of the mortality hazards of the control and GTE-treated
92   groups throughout the period of testing, using a method described previously[14,15]. The mortality
93   hazard of the GTE-treated group is reduced relative to that of the control group before the
94   median lifespan, but shortly thereafter crosses over, exceeding that of the control group.
95
Figure 1C shows the application of the time-varying hazard ratio analysis to the GTE data. The
log ratio of the mortality hazards of GTE treated and control groups shown in Figure 1B is
calculated along with its 95% confidence intervals. Negative values (i.e., log hazard ratios < 0)
indicate that GTE has a beneficial effect (i.e., lower mortality hazard than the control group),
while positive values (i.e., log hazard ratio > 0) reveal adverse effects of GTE treatment. The
95% confidence intervals of the mortality hazard ratio were estimated using 1,000 bootstrapped
replications[15], shown as dashed lines in the figure. Ages when the hazard ratio is < 0 and the
upper 95% confidence limit is also < 0 indicate when the treatment is significant for reducing
mortality. The duration (age range) of significance is bounded by the ages when the upper 95%
confidence interval crosses 0, as illustrated. Conversely, ages when the hazard ratio is > 0 and
the lower 95% confidence limit is > 0, correspond to ages when the treatment is significantly
increasing mortality, and the duration of significance is bounded in the same way. This analysis
reveals that GTE both reduced mortality hazards during midlife and increased mortality hazards
toward the end of life.
110
Figure 1D is the graphical representation of the results of the new analysis. It combines the
features of Figure 1C into an annotated horizontal heatmap to facilitate cross-drug comparisons.
Colors encode the drug effects on a single horizontal band from birth to the death of the last
mouse in either the control group or treated group, whichever is first. The heatmap is blank until
treatment begins. During the period of treatment, gray designates ages with no significant
treatment effect. Green and red designate ages when the treatment is significantly reducing or
increasing the mortality rate, respectively. Color intensity is directly proportional to effect size
(log HR). This graphic representation facilitates comparisons across various treatments as
shown in Figures 2 and 3. Unlike the log-rank test, this tool can identify at what ages, for how
long, and to what extent an intervention significantly alters the age-specific mortality hazard. It
also can show whether the intervention decreases or increases the mortality rate**.**
122
123   Greater sensitivity and precision in identifying mortality-modifying interventions
124
Figure 2 shows all the interventions identified by the new analytic tool that significantly reduced
(or increased) the age-specific mortality hazard during treatment, using the annotated horizontal
heatmap graphical representation. The hazard ratio plots used to generate these heatmaps,
calculated by the time-varying hazard ratio analysis, are shown in Figures S2 and S3, for males
and females, respectively. In this Figure, the interventions are ranked from the earliest to the
oldest age of cessation of beneficial effect in males. Figure S1 shows the results ranking by the
age of cessation of beneficial effect in females. Thirty-two compounds, consisting either of a
single agent or a combination of two agents, at one or more doses, initiated at varying ages,
significantly modified the mortality hazard in one or both sexes at one or more periods during

the treatment period. This analysis identified 12 new compounds that significantly reduced
mortality in at least one sex during treatment but were overlooked by the log-rank test: namely,
candesartan cilexetil (CC), caffeic acid phenethyl ester (CAPE), 17-dimethylaminoethylamino-
17-demethoxygeldanamycin hydrochloride (DMAG), enalapril, GTE, L-leucine, metformin,
oxaloacetic acid (OAA), PB125, simvastatin, syringaresinol (Syr), and ursodeoxycholic acid
(UDCA). The new analysis also identified 16 compounds that were detrimental (i.e., increased
mortality) in one or both sexes at one or more periods of treatment. The duration of significant
benefit or detriment varied markedly, from a few days (e.g., simvastatin, minocycline) to the
entire treatment period (e.g., rapamycin + acarbose). Most drugs only reduced mortality or only
increased mortality. Two exceptions were UDCA in males and GTE in females. Both were
beneficial for several months before the median lifespan and detrimental for several months in
very old mice. Effect sizes, indicated by the color intensity, varied markedly during the periods of
benefit and detriment. Acarbose had its greatest benefit at the initiation of treatment, waning
progressively thereafter. Effect sizes of other compounds, such as butanediol and captopril in
males and many of the different rapamycin trials in females peaked during the middle of
treatment. A few interventions showed a steady increase in effect with continued treatment (e.g.,
glycine in males and leucine in females).
151
152 <u>Only a fraction of interventions reduced mortality at later ages</u>
153
One of the benefits of the new analytic tool is its ability to estimate when during the life course
and for how long an agent exerts its effect on survival. In males, 17 drugs reduced mortality
hazards at some point during the life course (Figure 2). Of these, 9 compounds only reduced
mortality risk in early and mid-adulthood (i.e., before reaching the median lifespan): Syr, (R/S)-
1,3-butanediol (BD), CC, captopril, enalapril, UDCA, metformin, and DMAG, and
nordihydroguaiaretic acid (NDGA) at 800 ppm. The two higher doses of NDGA had a slightly
longer period of benefit, but only a few days beyond the median lifespan. By contrast, in
females, of the 11 agents that reduced mortality risk at some stage of life, a much smaller
fraction only reduced mortality during early- to mid-adulthood: GTE and OAA. In males, five
compounds tested in 11 trials demonstrated reduced mortality after attainment of median
lifespan, although these effects vanished before mice attained ages reaching the 90% mortality
benchmark: 17α-estradiol, aspirin at 21 ppm, Protandim, high doses of NDGA, and 3 of 4 late-
onset (20 mo) rapamycin treatments. Notably, only 6 of the 17 compounds that reduced
mortality in males did so at ages beyond the 90% mortality threshold: canagliflozin, acarbose,
17α-estradiol, glycine, simvastatin, rapamycin, and cocktails of either acarbose or metformin
with rapamycin. In females, in contrast to males, most of the drugs with beneficial effects (i.e., 9
of 11) reduced mortality mainly at ages after attainment of median lifespan. 8 trials involved
drugs that reduced mortality at ages after the median lifespan, but half of these lost efficacy
before reaching 90% mortality, including CAPE, glycine, CC, PB125, acarbose, and one trial of
late-onset rapamycin treatments; 14 trials had mortality-reducing compounds with efficacy
beyond the 90% mortality milestone: predominantly rapamycin-related (10 out of 14), and BD, L-
leucine, and captopril (Figure S1).
176
177 <u>Some compounds have adverse effects on mortality hazards</u>
178
Unexpectedly, the new analysis identified 16 compounds that adversely affected mortality at
specific stages in the life course in at least one sex (Figure 3). Since the inception of the ITP,
concerns have lingered about the potential lifespan-shortening and mortality hazard-increasing
effects of some compounds. However, no substances were identified with significant adverse
effects using the log-rank test. The analytic tool used here revealed 20 trials with 16 compounds
that increased the mortality hazard at one or more periods of treatment: 5 in males and 15 in

females. In males, UDCA, 3-(3-hydroxybenzyl)-5-methylbenzo[d]oxazol-2(3H)-one (MIF098), 2-(2-Hydroxyphenyl)benzoxazole (HBX), resveratrol, and INT-767 had significant negative impacts on the mortality hazard ratio. In females, CC, metformin, DMAG, canagliflozin, metformin combined with rapamycin, 17α-estradiol, GTE, minocycline, beta-guanidinopropionic acid (bGPA), geranylgeranyl acetone (GGA), fish oil, nicotinamide riboside (NR), UDCA, and MIF098 exhibited detrimental effects. UDCA and MIF098 affected both sexes adversely. Notably, some compounds displayed dual effects. In males, UDCA treatment showed early protective effects during pre-median lifespan stages but at later ages manifested significant negative impacts. In females, three experiments demonstrated mixed outcomes. GTE, akin to UDCA's pattern in males, had early beneficial effects but turned detrimental nearing the 90% survival mark. CC exhibited significant early adverse effects but became protective post-median lifespan. Most intriguing was the metformin and rapamycin combination (MetRapa), which presented pronounced benefits beyond the 90% mortality threshold but briefly exhibited significant detrimental effects at advanced ages in females. Metformin given alone also had detrimental effects briefly at the end of life in females.

Sex differences in the effect of pharmacological interventions

Marked sex differences in the responses to life-extending drugs are one of the key outcomes of the ITP[12]. In addition to those already noted, this new analytic tool unveiled even more sex differences in the response to the pharmacological interventions of the ITP. It identified 6 additional compounds that only benefited males: Syr, enalapril, simvastatin, metformin, DMAG, and UDCA, and 5 drugs that only reduced mortality in females: OAA, CAPE, PB125, Leu, and GTE. Notably, 7 interventions, including UDCA, CC, metformin, DMAG, canagliflozin, MetRapa, and 17α-estradiol, exhibited beneficial effects in males but detrimental effects in females (Figure 3). More compounds adversely affected survival in females (13) than in males (4). Moreover, most drugs with negative effects exerted their effect on females almost from the beginning of treatment. The detrimental effects waned during the 2^{nd} year of life but sometimes reappeared in the final stage of life. Six agents only had deleterious effects late in life: fish oil, GTE, metformin, and MetRapa in females, and MIF098, INT-767, and resveratrol in males. These results underscore the necessity of including both sexes when testing longevity interventions.

Discussion

The analytic method presented here promises to be broadly useful and impactful for interventional research on aging. Revealing the age-specific effects of interventions on the mortality hazard opens the door to asking more nuanced and targeted questions about the actions of an intervention. Answers to such questions can ultimately lead to greater life-extending efficacy of interventions and a better understanding of the underlying mechanisms that the interventions target. The analysis does this by providing estimates of when and for how long during the life course an intervention reduces (or in the case of detrimental effects, increases) age-specific mortality. It also provides an estimate of the effect size of an intervention and how the strength of its effect changes over the course of treatment. None of this information is attainable by traditional methods such as the log-rank test, the current standard for evaluating longevity interventions.

This new method can distinguish interventions that specifically reduce mortality during senescence from those that only affect survival during midlife or earlier. This is an important distinction in the search for therapeutic interventions that benefit individuals of advanced age when the burdens of senescence are greatest. This analytic tool is also sensitive to adverse effects—critically important for pre-clinical models that aim to be translatable. Furthermore, the

method is sensitive to sex differences in timing, duration, and efficacy of interventions, as well
as adverse outcomes—providing further impetus to probe the mechanisms underlying the
growing number of sexually dimorphic traits in aging. Here we discuss some of the ways the
new information provided by this analytic tool can assist drug discovery, the search for the
underlying mechanisms that drive aging, and other areas of Geroscience. These are only a few
examples of how this analytical tool can be utilized. Additional applications will likely emerge as
its adoption spreads within the geroscience community.
243
A major discovery using this tool is that the effect of virtually every intervention analyzed was
non-uniform across the life course. This observation is not readily apparent by visual inspection
of most Kaplan-Meier plots and is not obtainable from log-rank tests. Very few interventions
significantly reduced (or increased) mortality through the entire course of treatment. Most were
only effective for less than half of the treatment duration. This calls for explanation, and the
answers are likely to lead to better interventions and greater insight into the mechanisms of
aging. One possibility is that the aging process may impact drug efficacy. The decrease,
increase, or loss of efficacy of an intervention may reflect age-related changes in
pharmacokinetics or pharmacodynamics, leading to suboptimal (over or under) dosage. The ITP
database provides some insights into this question, because some interventions were tested at
several doses. Acarbose, for example, was tested at three doses. Acarbose efficacy in males,
measured as the reduced mortality hazard ratio, increased with increasing doses during the
initial period of treatment, but paradoxically, its beneficial effect ceased at progressively earlier
ages with increasing doses. This finding opens the door to developing age-specific doses to
sustain efficacy for longer periods and raises awareness of the importance of understanding the
role of aging in pharmacokinetics. An alternative explanation for the complex response to
varying doses of acarbose is an age-related change in pharmacodynamics. It is plausible that
the aging processes or causes of mortality change with age and the intervention loses efficacy
because it no longer targets the underlying pathways. Whatever the reason, this tool has
uncovered a critical variable that needs to be considered in interventional geroscience.
264
Another important outcome of the application of this analytic tool to longevity data is the finding
that only a subset of the interventions in the ITP database affected age-specific mortality rates in
the last half of the lifespan, and even fewer affected mortality rates at ages after the age when
90% of the control cohort has died (maximum lifespan)[8]. Diet restriction has long been
considered an example of an intervention that retards aging processes broadly, because it
extends the age of 90% mortality, distinguishing it from many interventions that do not—the
latter often only extending the median lifespan[16,17]. Several studies, including the ITP, use the
Wang-Allison test as a discriminator for interventions that do or do not extend the maximum
lifespan based on the 90% mortality measure. However, this test does not distinguish whether
an increase in age of 90% mortality reflects the effects of reduced mortality accumulated during
earlier ages from the effects of the age-specific mortality reduction at or near the age of 90%
mortality. Also, it provides no information on the effect of the intervention on the last 10% of the
population. This distinction is of particular importance to a major goal of Geroscience: namely, to
identify compounds and discover the underlying mechanisms that unequivocally extend the
maximum lifespan of a species by reducing age-specific mortality during the later stages of life
when the burden of senescence is greatest. The analytic tool described here provides such a
measure by indicating whether the intervention specifically reduces mortality rates in the final
stage of life, whether it be after the age at which 90% of the control population has died or some
other late age. Only a subset of the interventions that have been reported by the ITP as
"lifespan extending" using log-rank analysis reduced mortality hazard after the median lifespan,
and even fewer did so during what is considered the maximum lifespan.
286

Nevertheless, compounds that only reduce mortality during the first half of adult life should not
be discounted. Reducing mortality at any stage of life can be impactful, especially when
considering its potential translatability to humans. For example, the male mortality
disadvantage, compared to females, is greatest in the first half of adult life in both humans and
UM-HET3 mice[15]. It is noteworthy that most of the compounds that are only effective in males
are only effective during the first half of the lifespan. Castration of UM-HET3 males before
puberty eliminates this mortality disadvantage[14]. If any of the drugs that only eliminate the male
mortality disadvantage during this period can do so without interfering with male reproductive
function, the societal impact if clinically translatable would be great[12,18].

This method is not only more sensitive to agents that reduce age-specific mortality, it also is
more sensitive to those that increase mortality. The ITP never identified adverse effects using
the log-rank test. This new tool revealed 20 trials involving 16 compounds that increased
mortality hazards at certain life stages in at least one gender. There was a marked sex
difference. Only 5 trials showed detrimental effects in males compared to 15 trials in females.
MIF098 was the only drug that adversely affected both sexes. Some compounds, including
canagliflozin and high doses of 17α-estradiol markedly reduced mortality in males but were
harmful in females. These findings underscore the need for sex-specific testing of life-extending
candidates.

This new analytic tool can detect reversals of the benefit of compounds across the life course.
UDCA in males and GTE in females reduced mortality before the median lifespan but increased
mortality at later ages—another discriminator not possible using the log-rank test. There is
precedence for this reversal. In humans, individuals reporting the lowest intake of dietary protein
had reduced mortality from cardiovascular disease and cancer before 65 years of age, but this
relationship reversed after 65[19]. Mice with reduced branch chain amino acid intake had
extended life when the diet began in early adulthood, but their lifespan was unaffected when the
diet was initiated at a later age[20]. Age-related changes in pharmacokinetics and
pharmacodynamics may play a role here. For example, blood levels of canagliflozin, whose
beneficial effects in males diminish with age and are absent in females are 2-3-fold higher in
older males and similarly elevated at all ages in females[21].

Another strength of this tool is its heightened sensitivity to potential life-extending candidates. It
identified over twice as many as the log-rank test. This is due in part to its ability to identify age-
specific effects on the mortality hazard unimpeded by the requirement of the log-rank test for
consistent proportional hazard across the duration of treatment. The newly identified
compounds generally have smaller effect sizes and shorter durations of positive effect
compared to those identified by the log-rank test. Given their geroprotective potential, they
deserve further study. It is important to emphasize that neither this nor any other statistical tool
should be used as a final arbiter of any candidate for mortality reduction and lifespan extension
(or adverse effect), but rather should be considered a screening tool for identifying potential
candidates that deserve follow-up—for example with different doses. We would argue that Type
1 errors (i.e., false positives) during initial screens are more acceptable and preferable to false
negatives.

It is important to note the limitations of this method and consider ways to increase its utility.
While the flexible estimation of the hazard ratio makes few assumptions about the proportional
hazard, the precision of the hazard ratio varies throughout the lifespan. The hazard ratio
precision tends to be low during early life due to the lower rate of mortality. The method requires
a larger sample size than the log-rank test, and the log-rank test has higher power when the PH
assumption is met. The bootstrap confidence intervals mitigate the false positive findings for a

wide range of sample sizes, but the Type I error rate is controlled at each specific time point
which does not ensure tight Type I error rates for the full lifespan that a permutation test might
achieve. The method currently does not explicitly consider uncertainty in the Time axis so the
ages at which the treatment effect becomes nonzero are presented as point-estimates without
confidence intervals. However, this limitation did not prevent the consistent findings between
similar treatments such as the early effects of ACE inhibitors (Enalapril and Captopril) or early
effects of different doses of NDGA. Statistically testing whether two different treatments have
the same effect relative to control is more complex (testing whether the ratio of hazard ratios is
1) and may require comparisons across cohorts. While this method allows the estimation of
time-varying treatment effects relative to control, future extensions of the method could explicitly
test and estimate differences in active treatments in terms of time and magnitude of reduction or
increase in the mortality hazard ratio.
350
351
352

Author contributions
Conceptualization: CJC, NJ, JG, QL, JN, RS
Methodology: CJC, NJ, JG, QL, JN
Funding acquisition: RS, JN
Data analysis: QL, NJ, CJC, JG, JN
Writing: NJ, JN, JG

Acknowledgments

Dr. Strong has been honored with the Senior Research Career Scientist award (# IK6 BX006289) from the Department of Veterans Affairs. The funding for this research was provided by the Center for Testing Potential Anti-Aging Interventions (5U01AG022307), the Nathan Shock Center of Excellence in Basic Biology of Aging (5P30AG013319), as well as NIH grant 5T32AG021890-15, alongside a fellowship from the Glenn Foundation awarded to CJC.

Conflict of interest
The authors declare no conflict of interest.

Funding

National Institute on Aging grant 5U01AG022307 (RS, JN)
National Institute on Aging grant 5P30AG013319 (RS, JN)

Online Methods:

<u>Data availability, mouse model, and husbandry</u>

The datasets employed in this study are sourced from the Mouse Phenome Database (MPD; phenome.jax.org), encompassing all data from the Interventions Testing Program (ITP) spanning from 2004 to 2022. This dataset incorporates 13 distinct cohorts, integrating data across three research facilities to ensure the robustness and reproducibility of the findings. The ITP employed the UM-HET3 mouse line, a genetically heterogeneous model, chosen for its relevance to the genetically diverse human population. UM-HET3 mice are bred according to a specific crossbreeding protocol: BALB/cByJ females are mated with C57BL/6J males to produce F1 hybrid females, which are then bred with F1 hybrid males derived from mating C3H/HeJ females with DBA/2J males. This breeding strategy is designed to maximize genetic diversity within the model, thereby approximating the genetic variability inherent in human populations and increasing the translational value of the research findings. The mice designated for longevity assays were maintained under controlled environmental conditions, with a constant ambient temperature of 25°C and a regulated photoperiod of 12 hours light/12 hours darkness. Nutritional needs were met with ad libitum access to the Purina 5LG6 diet, alongside specific drugged food formulations as per experimental requirements. Housing protocols were optimized for social enrichment and welfare, accommodating up to three males or five females per standard laboratory enclosure, in accordance with established ethical guidelines. Rigorous daily health assessments were conducted by trained staff to monitor the well-being of the subjects, promptly identify morbidity signs, and implement early intervention strategies as necessary. This proactive health management approach minimized unnecessary suffering and ensured the reliability of longevity data. The specifics of drug administration, including dosage, frequency, and duration, as well as the rationale behind the selection of intervention agents, are detailed in the original published reports, providing a comprehensive overview of the therapeutic strategies explored in this body of research.

<u>Description of the time-varying hazard ratio test</u>

Our investigation into the impact of various treatments on age-specific mortality utilized a piecewise polynomial B-spline hazard model. This model, which assumes a Poisson distribution, was applied using the bshazard package in R, offering a robust tool for analyzing the complex interplay between treatment effects and mortality over time. By integrating this model, we were able to capture the nuanced variations in mortality risk associated with different treatment regimens across the lifespan of the subjects. A key aspect of our analytical strategy was the generation of a nonparametric smoothed estimate of the baseline hazard rate. This was achieved by stratifying the survival data by both treatment and sex within each cohort, thereby allowing for a precise adjustment for the site-specific effects that might otherwise confound the treatment impact assessment. Importantly, this approach facilitated a refined understanding of how baseline mortality risks shift in response to treatment interventions, while accounting for potential biological differences in treatment efficacy between males and females. In our analysis, mortality events occurring prior to the initiation of treatment were excluded to ensure that the hazard ratio estimates accurately reflect the treatment's effect on survival. This exclusion criterion is crucial for eliminating bias arising from pre-treatment mortality, thus enhancing the validity of our findings. The confidence intervals for the treatment hazard ratio were estimated using 1,000 bootstrapped replications. This age-specific analysis is similar to that reported in the estimated age-specific effects of sex[15]. The conventional evaluation of the time-varying hazard (age-specific mortality) was conducted using the test of proportional

hazards (PH) assumption (z-test)[22]. This test assesses whether the hazard ratio (treatment
effect) varies across the lifespan. This test was performed for each sex and treatment
combination. Test results are shown in Table S1. The distribution of PH violation p-values was
5% with p<.01 (5 times expected), 12% with p<.05 (2.4 times expected), 19% with p<.1 (1.9
times expected). The p-values were combined using Fisher's Method indicating that the PH
assumption does not hold (p<.0001) for a subset of interventions.
432
433   <u>Data Visualization</u>
434
The visualization method uses a color-coded band to depict treatment effects on hazard ratios,
with the pre-treatment phase shown as a blank band. Upon treatment initiation, a gray color
indicates no detectable effect, while significant effects are represented by changes in color
intensity: beneficial effects cause the band to turn green, with the intensity reflecting the
magnitude of negative log hazard ratios, and detrimental effects are shown in red, with intensity
corresponding to positive log hazard ratios. The transition points where significant effects begin,
or end are marked by dashed lines. Additionally, key lifespan metrics for the control group, such
as median and maximum lifespan (when 90% have died), are highlighted to facilitate
interpretation. All computational analyses were conducted in R (version 4.3, Vienna, Austria).
444
445
446

References:

Harrison, D. E. *et al.* Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature* **460**, 392-395, doi:10.1038/nature08221 (2009).

Harrison, D. E. *et al.* Acarbose, 17-alpha-estradiol, and nordihydroguaiaretic acid extend mouse lifespan preferentially in males. *Aging Cell* **13**, 273-282, doi:10.1111/acel.12170 (2014).

Knufinke, M., MacArthur, M. R., Ewald, C. Y. & Mitchell, S. J. Sex differences in pharmacological interventions and their effects on lifespan and healthspan outcomes: a systematic review. *Front Aging* **4**, 1172789, doi:10.3389/fragi.2023.1172789 (2023).

Mantel, N. Evaluation of survival data and two new rank order statistics arising in its consideration. *Cancer Chemother Rep* **50**, 163-170 (1966).

Bouliotis, G. & Billingham, L. Crossing survival curves: alternatives to the log-rank test. *Trials* **12**, A137, doi:10.1186/1745-6215-12-S1-A137 (2011).

Jiang, N., Gelfond, J., Liu, Q., Strong, R. & Nelson, J. F. The Gehan test identifies life-extending compounds overlooked by the log-rank test in the NIA Interventions Testing Program: Metformin, Enalapril, caffeic acid phenethyl ester, green tea extract, and 17-dimethylaminoethylamino-17-demethoxygeldanamycin hydrochloride. *bioRxiv*, 2024.2002.2017.579363, doi:10.1101/2024.02.17.579363 (2024).

Harrington, D. P. & Fleming, T. R. A class of rank test procedures for censored survival data. *Biometrika* **69**, 553-566 (1982).

Wang, C., Li, Q., Redden, D. T., Weindruch, R. & Allison, D. B. Statistical methods for testing effects on "maximum lifespan". *Mech Ageing Dev* **125**, 629-632, doi:10.1016/j.mad.2004.07.003 (2004).

Gao, G., Wan, W., Zhang, S., Redden, D. T. & Allison, D. B. Testing for differences in distribution tails to test for differences in 'maximum' lifespan. *BMC Med Res Methodol* **8**, 49, doi:10.1186/1471-2288-8-49 (2008).

Pletcher, S. D., Khazaeli, A. A. & Curtsinger, J. W. Why do life spans differ? Partitioning mean longevity differences in terms of age-specific mortality parameters. *J Gerontol A Biol Sci Med Sci* **55**, B381-389, doi:10.1093/gerona/55.8.b381 (2000).

Miller, R. A. *et al.* An Aging Interventions Testing Program: study design and interim report. *Aging Cell* **6**, 565-575, doi:10.1111/j.1474-9726.2007.00311.x (2007).

Jiang, N. & Nelson, J. F. Sex Differences in Mouse Longevity and Responses to Geroprotective Drugs: Implications for Human Intervention. *Public Policy Aging Rep* **33**, 120-124, doi:10.1093/ppar/prad026 (2023).

Strong, R. *et al.* Evaluation of resveratrol, green tea extract, curcumin, oxaloacetic acid, and medium-chain triglyceride oil on life span of genetically heterogeneous mice. *J Gerontol A Biol Sci Med Sci* **68**, 6-16, doi:10.1093/gerona/gls070 (2013).

Jiang, N. *et al.* Prepubertal castration eliminates sex differences in lifespan and growth trajectories in genetically heterogeneous mice. *Aging Cell* **22**, e13891, doi:10.1111/acel.13891 (2023).

Cheng, C. J., Gelfond, J. A. L., Strong, R. & Nelson, J. F. Genetically heterogeneous mice exhibit a female survival advantage that is age- and site-specific: Results from a large multi-site study. *Aging Cell* **18**, e12905, doi:10.1111/acel.12905 (2019).

Yu, B. P., Masoro, E. J. & McMahan, C. A. Nutritional influences on aging of Fischer 344 rats: I. Physical, metabolic, and longevity characteristics. *J Gerontol* **40**, 657-670, doi:10.1093/geronj/40.6.657 (1985).

Masoro, E. J. Biology of Aging: Current State of Knowledge. *Archives of Internal Medicine* **147**, 166-169, doi:10.1001/archinte.1987.00370010164033 (1987).

496   18   Jiang, N., Cheng, C. J., Strong, R. & Nelson, J. F. Castration reduces mortality and
497   increases resilience in male mice: what is next? *Geroscience* **46**, 2787-2790,
498   doi:10.1007/s11357-023-00973-5 (2024).
499   19   Levine, M. E. *et al.* Low protein intake is associated with a major reduction in IGF-1,
500   cancer, and overall mortality in the 65 and younger but not older population. *Cell Metab*
501   **19**, 407-417, doi:10.1016/j.cmet.2014.02.006 (2014).
502   20   Richardson, N. E. *et al.* Lifelong restriction of dietary branched-chain amino acids has
503   sex-specific benefits for frailty and lifespan in mice. *Nat Aging* **1**, 73-86,
504   doi:10.1038/s43587-020-00006-2 (2021).
505   21   Miller, R. A. *et al.* Canagliflozin extends life span in genetically heterogeneous male but
506   not female mice. *JCI Insight* **5**, doi:10.1172/jci.insight.140019 (2020).
507   22   Grambsch, P. M. & Therneau, T. M. Proportional hazards tests and diagnostics based on
508   weighted residuals. *Biometrika* **81**, 515-526 (1994).
509   23   Strong, R. *et al.* Lifespan benefits for the combination of rapamycin plus acarbose and
510   for captopril in genetically heterogeneous mice. *Aging Cell* **21**, e13724,
511   doi:10.1111/acel.13724 (2022).
512
513

Figures:

 Figure 1. Graphical representation of the analytic tool for determining the timing and impact of
 life-extending candidates. Survival data are from the test of Green Tea Extract in females[13]. **A)**
 Kaplan-Meier survival curves of the GTE-treated female mice (Red) and control female mice
 (Black); **B)** Age-specific mortality hazards of GTE-treated and control mice groups with 95%
 confidence intervals; **C)** Mortality hazard ratio between GTE-treated and control mice groups
 and 95% confidence intervals shown as dashed lines; **D)** Life course heat map visualization of
 the age-specific effects of GTE on the mortality hazard ratio. Vertical dashed lines mark the
 boundaries of significant effects on the mortality hazard ratio based on the ages when the 95%
 confidence intervals in Figure C cross 0.

Figure 2. Interventions that significantly modified mortality hazard using the time-varying hazard
analytic tool. Each row represents an individual trial of one intervention in a single cohort. Each
intervention involved one compound or a combination of two, with dosage and starting age of
treatment listed. Trials are ranked by the cessation age of beneficial effects in males, from
earliest to latest age. The color-coded bands denote the temporal significance of drug effects:
white indicates the period before treatment onset, gray marks periods with no significant effects,
green indicates periods of significant beneficial effects, and red denotes intervals of significant
detrimental effects. The solid black triangle indicates the median lifespan of the control group for
each trial, and the open triangle marks the age of 90% mortality of the control group.
539
540 Footnotes:
541 # New Compounds that significantly affect mortality (i.e., not identified by the log-rank test) are
542 also noted in **bold** font.
543 $ PB125 is a mixture of luteolin, withaferin A, and carnosol, dosages refer to publication[23].

Compounds	Dose (ppm)	Age of Onset (months)	Age-specific Drug Effects on Mortality Hazard	
			Female	Male
Ursodeoxycholic Acid (UDCA)	5000	5		
MIF098	240	8		
Canagliflozin	180	7		
Nicotinamide Riboside (NR)	1000	8		
Fish Oil	15000	9		
17-a-estradiol	14	10		
Candesartan Cilexetil (CC)	30	8		
DMAG	30	6		
Fish Oil	50000	9		
Green Tea Extract	2000	4		
Geranylgeranyl Acetone (GGA)	600	9		
b-Guanidinopropionic Acid (bGPA)	3300	6		
Minocycline	300	6		
Rapa_Met	1000, 14	9		
Metformin	1000	9		
INT-767	180	10		
2-(2-Hydroxyphenyl) benzoxazole (HBX)	1	15		
Resveratrol	300	4		

Figure 3. Trials with Drug-Induced Detrimental Effects on Mortality Hazard. These trials are shown ranked by the length of time (longest to shortest) during which significant detrimental effects were observed in females. Each color-coded band across the timeline represents the drug effect phase in relation to the treatment period: white for the pre-treatment phase, gray for phases without significant impact, green for phases with significant beneficial effects, and red for phases with significant adverse effects. Key lifespan indicators are marked by triangles; a solid black triangle denotes the median lifespan in the control group, and an open triangle indicates the point at which 10% of the control group remains alive.

Age-specific Drug Effects on Mortality Hazard

Compounds	Dose (ppm)	Age of Onset (months)	Male	Female
Green Tea Extract#	2000	4		
Oxaloacetic acid (OAA)#	2200	4		
Caffeic Acid Phenethyl Ester (CAPE)#	300	4		
Glycine	80000	9		
Candesartan Cilexetil (CC)#	30	8		
Rapa_start_stop	42	20 (until 23)		
PB125#	See Pub$	5		
Acarbose	400	8		
Acarbose	1000	8		
Acarbose	2500	8		
(R/S)-1,3-butanediol (BD)	100000	6		
Rapa_cycle	42	20 (on off)		
Acarbose	1000	4		
Rapamycin	42	20		
Rapa_Met	1000, 14	9		
Rapamycin	14	20		
Rapamycin	14	9		
Rapa_Aca	14.7, 1000	9		
Rapamycin	42	9		
Rapamycin	14	9		
Rapamycin	4.7	9		
Rapa_Aca	14.7, 1000	16		
Captopril	180	5		
L-leucine#	40000	5		
Minocycline#	300	6		
b-Guanidinopropionic Acid (bGPA)#	3300	6		
Geranylgeranyl Acetone (GGA)#	600	9		
Fish Oil#	15000	9		
Nicotinamide Riboside (NR)#	1000	8		
Canagliflozin				
MIF098#				
Ursodeoxycholic A (UDCA)#				
DMAG				
17-a-estradiol				
Fish Oil#	C2010_FO_50000_9 Male			
Metformin#				
Syringaresinol#				
Enalapril#				
Nordihydroguaiareti acid (NDGA)				
Nordihydroguaiareti acid (NDGA)				
Nordihydroguaiareti acid (NDGA)				
17-a-estradiol	14.4	20		
17-a-estradiol	4.8	10		
Aspirin	21	4		
Nordihydroguaiaretic acid (NDGA)	2500	9		
Protandim	600	10		
17-a-estradiol	14.4	16		
Acarbose	1000	16		
Simvastatin#	120	10		
2-(2-Hydroxyphenyl) benzoxazole (HBX)#	1	15		
Resveratrol#	300	4		
INT-767#	180	10		

Age (days): 0 400 800 1200 1600   0 400 800 1200 1600

time

Mean Log Hazard Ratio: 1 / 0 / -1

Figure S1. Interventions that significantly modified mortality hazard using the time-varying
hazard analytic tool, which is ranked by the cessation age of beneficial effects in females. Each
row represents an individual trial of one intervention in a single cohort. Each intervention
involved one compound or a cocktail of two, with dosage and starting age of treatment listed.
The color-coded bands denote the temporal significance of drug effects: white indicates the
period before treatment onset, gray marks periods with no significant effects, green indicates
periods of significant beneficial effects, and red denotes intervals of significant detrimental
effects. The solid black triangle indicates the median lifespan of the control group for each trial,
and the open triangle marks the age of 90% mortality of the control group.
565
566 Footnotes:
567 # New Compounds that significantly affect mortality (i.e., not identified by the log-rank test) are
568 also noted in **bold** font.
569 $ PB125 is a mixture of luteolin, withaferin A, and carnosol, dosages refer to publication[23].