Title Local nutrient addition drives plant diversity losses but not biotic homogenization in global
grasslands

Author list
Qingqing Chen[1,2*], Shane A. Blowes[1,3], W. Stanley Harpole[1,4,5], Emma Ladouceur[1,3,6–8], Elizabeth
T. Borer[9], Andrew MacDougall[10], Jason P. Martina[11], Jonathan D. Bakker[12], Pedro M. Tognetti[13],
Eric W. Seabloom[9], Pedro Daleo[14], Sally Power[15], Christiane Roscher[1,4], Peter B. Adler[16], Ian
Donohue[17], George Wheeler[18], Carly Stevens[19], G.F. (Ciska) Veen[20], Anita C. Risch[21], Glenda M.
Wardle[22], Yann Hautier[23], Catalina Estrada[24], Erika Hersch-Green[25], Yujie Niu[26], Pablo L. Peri[27],
Anu Eskelinen[1,4,28],Daniel S. Gruner[29], Harry Olde Venterink[30], Carla D'Antonio[31],Marc W.
Cadotte[32], Sylvia Haider[1,33], Nico Eisenhauer[1,34], Jane Catford[35,36], Risto Virtanen[28], John W.
Morgan[37], Michelle Tedder[38], Sumanta Bagchi[39], Maria C. Caldeira[40], Miguel N. Bugalho[41],
Johannes M. H. Knops[42], Chris R. Dickman[43], Nicole Hagenah[44], Anke Jentsch[26], Petr Macek[45],
Brooke B. Osborne[46], Lauri Laanisto[47], Jonathan M. Chase[1,3]
14

Affiliations
1. German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig.
2. Senckenberg Museum for Natural History Görlitz, D-02826, Görlitz, Germany.
3. Department of Computer Science, Martin Luther University Halle-Wittenberg, Halle (Saale),
Germany.
4. Helmholtz Center for Environmental Research – UFZ, Department of Physiological Diversity,
Permoserstrasse 15, 04318 Leipzig, Germany.
5. Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108 Halle (Saale), Germany.
6. Department of Biology, University of Prince Edward Island, PE, Canada.
7. Canadian Centre for Climate Change and Adaptation, University of Prince Edward Island, St.
Peter's Bay, PE, C0A 2A0, Canada.
8. School of Climate Change and Adaptation, University of Prince Edward Island, Charlottetown,
PE, C1A 4P3, Canada.
9. Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN 55108
USA.
10. Department of Integrative Biology, University of Guelph, Guelph, Ontario, Canada N1G2W1.
11. Department of Biology, Texas State University, San Marcos, TX, 78666 USA.
12. School of Environmental and Forest Sciences, University of Washington, Seattle, WA USA.
13. Instituto de Investigaciones Fisiológicas y Ecológicas Vinculadas a La Agricultura (IFEVA),
Facultad de Agronomía, Universidad de Buenos Aires and CONICET, Buenos Aires, Argentina.
14. Instituto de Investigaciones Marinas y Costeras (IIMyC), UNMdP-CONICET, CC 1260 Correo
Central, Mar Del Plata B7600WAG, Argentina.
15. Hawkesbury Institute for the Environment, Locked Bag 1797, Penrith, New South Wales, 2751,
Australia.
16. Department of Wildland Resources and the Ecology Center, Utah State University, Logan, UT,
84322 USA.
17. Zoology, School of Natural Sciences, Trinity College Dublin, Dublin, Ireland.
18. University of Nebraska-Lincoln, Lincoln, NE, USA.
19. Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK.
20. Department of Terrestrial Ecology, Netherlands Institute of Ecology, Droevedaalsesteeg 10,
6708 PB, Wageningen, The Netherlands.
21. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zuercherstrasse 111,
8903 Birmensdorf, Switzerland.
22. Desert Ecology Research Group, School of Life and Environmental Sciences, ARC Training
Centre in Data Analytics for Resources and Environments (DARE), The University of Sydney,
NSW 2006, Australia.

23. Ecology and Biodiversity Group, Department of Biology, Utrecht University, Padualaan 8, 52 3584 CH Utrecht, The Netherlands.
24. Department of Life Sciences, Imperial College London, Silwood Park Campus, Buckhurst 54 Road, Ascot SL5 7PY, UK.
25. Department of Biological Sciences, Michigan Technological University, Houghton, MI 49930, 56 USA.
26. Disturbance Ecology and Vegetation Dynamics, Bayreuth Center of Ecology and 58 Environmental Research (BayCEER), University of Bayreuth, Germany.
27. Instituto Nacional de Tecnología Agropecuaria (INTA), Universidad Nacional de La Patagonia 60 Austral (UNPA).CONICET. Río Gallegos, Santa Cruz, Argentina.
28. Ecology & Genetics Unit, University of Oulu, Finland.
29. Department of Entomology, University of Maryland, College Park, MD USA.
30. Vrije Universiteit Brussel, Department Biology, WILD, Pleinlaan 2, 1050 Brussels, BELGIUM.
31. University of California, Santa Barbara.
32. Dept. of Biological Sciences, University of Toronto Scarborough, Toronto, ON, Canada.
33. Institute of Ecology, Leuphana University of Lüneburg, Universitätsallee 1, 21335 Lüneburg, 67 Germany.
34. Institute of Biology, Leipzig University; Puschstrasse 4, 04103 Leipzig, Germany.
35. Department of Geography, King's College London, 30 Aldwych, London, WC2B 4BG, UK.
36. School of Ecosystem & Forest Sciences, University of Melbourne, Parkville, VIC 3010, 71 Australia.
37. Department of Environment & Genetics, La Trobe University, Bundoora 3083, Victoria, 73 Australia.
38. Centre for Functional Biodiversity, School of Life Sciences, University of KwaZulu-Natal, 75 Pietermaritzburg, South Africa.
39. Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India.
40. Forest Research Centre, Associate Laboratory TERRA, School of Agriculture, University of 78 Lisbon, Portugal.
41. Centre for Applied Ecology 'Prof. Baeta Neves' (CEABN-InBIO), School of Agriculture, 80 University of Lisbon, Portugal.
42. Health & Environmental Sciences Department, Xi'an Jiaotong-Liverpool University, Suzhou, 82 Jiangsu, China.
43. Desert Ecology Research Group, School of Life and Environmental Sciences, The University of 84 Sydney, Sydney, NSW 2006, Australia.
44. Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria, 86 Pretoria, South Africa.
45. Institute of Hydrobiology, Biology Centre of the Czech Academy of Sciences, Na Sadkach 7, 88 Ceske Budejovice 370 05, Czech Republic.
46. Department of Environment and Society, Utah State University, Moab, UT, USA.
47. Chair of Biodiversity and Nature Tourism, Estonian University of Life Sciences, Kreutzwaldi 5, 91 51006, Tartu, Estonia.
*corresponding author: chqq365@gmail.com

Abstract:

Nutrient enrichment typically causes local plant diversity declines. A common but untested
expectation is that nutrient enrichment also reduces variation in nutrient conditions among localities
and selects for a smaller pool of species, causing greater diversity declines at larger than local scales
and thus biotic homogenization. Here we apply a framework that links changes in species richness
across scales to changes in the numbers of spatially restricted and widespread species for a
standardized nutrient addition experiment across 72 grasslands on six continents. Overall, we find
proportionally similar species loss at local and larger scales, suggesting similar declines of spatially
restricted and widespread species, and no biotic homogenization after 4 years and up to 14 years of
treatment. These patterns of diversity changes are generally consistent across species groups. Thus,
nutrient enrichment poses threats to plant diversity, including for widespread species that are often
critical for ecosystem functions.

105
106

Introduction

Agricultural fertilization practices and atmospheric nutrient deposition have led to increased availability and redistribution of soil nutrients globally[1–3]. At local scales (i.e., α diversity), nutrient enrichment tends to reduce plant diversity[4,5]. This diversity decline is typically ascribed to disproportionate losses of rare species (i.e., species with relatively low cover) because small populations are more susceptible to extinction[6–8]. In addition, nutrient enrichment often leads to the removal of species with traits ill-suited for effectively competing in high nutrient conditions[6]. While species can vary widely in their nutrient requirements and tolerances, groups of species with similar (shared) characteristics can be lost from a flora. For example, native species are more likely to be lost than non-native species when nutrients are enriched because non-natives are often better-adapted to nutrient-rich conditions[6,9,10]. Similarly, nitrogen-fixing legumes may be more vulnerable than other species in high nutrient environments due to their decreased competitive advantage[6,11].

Despite clear evidence that nutrient enrichment causes losses of α diversity in grasslands[4,5], how these losses are reflected at larger spatial scales (i.e., γ diversity; calculated by aggregating local communities) is less straightforward[12–14]. Yet, it is diversity loss and change at larger spatial scales that is most often relevant for biodiversity conservation and management as well as for the provision of ecosystem functions and services[15,16]. It is often assumed that nutrient enrichment, like many other global changes, results in biotic homogenization (i.e., increasing similarity in composition among local communities, quantified as a decrease in β diversity)[17–21]. This is because local nutrient enrichment is expected to create homogeneous nutrient conditions among localities and consistently select for a smaller pool of species that are nutrient-demanding, fast-growing, and highly competitive for light[22,23]. Previous investigations of scale-dependent diversity change under nutrient enrichment have tended to be short term or limited in spatial extent[24–28]. These short-term or spatially-restricted studies have found mixed results, indicating that nutrient enrichment leads to biotic homogenization[29–31], no changes in β diversity[24,26,27,32] or even differentiation (i.e., increase in β diversity)[25,28,33–37].

Fig 1. A framework linking diversity changes at the α-, γ-, and β-scales to changes in the number of spatially restricted and widespread species. $\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$ are log response ratios (LRR) of average α-, γ- and β- diversity under nutrient enrichment relative to that under ambient conditions (control). $\Delta\beta$ is equal to zero along the 1:1 diagonal line. Below the diagonal line, $\Delta\gamma < \overline{\Delta\alpha}$, $\Delta\beta < 0$, nutrient enrichment causes biotic homogenization. Above the diagonal line, $\Delta\gamma > \overline{\Delta\alpha}$, $\Delta\beta > 0$, nutrient enrichment causes biotic differentiation. Moreover, homogenization can be attributed to three scenarios including I: gain of widespread species ($\Delta\gamma < \overline{\Delta\alpha}$, and $\Delta\gamma > 0$, $\overline{\Delta\alpha} > 0$); II: spatially restricted species replaced by widespread species ($\Delta\gamma < \overline{\Delta\alpha}$, $\Delta\gamma < 0$ and $\overline{\Delta\alpha} > 0$); III: Loss of spatially restricted species ($\Delta\gamma < \overline{\Delta\alpha}$, and $\Delta\gamma < 0$, $\overline{\Delta\alpha} < 0$). Conversely, differentiation can be attributed to three scenarios that include IV: Loss of widespread species ($\Delta\gamma > \overline{\Delta\alpha}$, and $\Delta\gamma < 0$, $\overline{\Delta\alpha} < 0$); V: Widespread species replaced by spatially restricted species ($\Delta\gamma > \overline{\Delta\alpha}$, $\Delta\gamma > 0$ and $\overline{\Delta\alpha} < 0$); VI: gain of spatially restricted species ($\Delta\gamma > \overline{\Delta\alpha}$, and $\Delta\gamma > 0$, $\overline{\Delta\alpha} > 0$). Adapted from[12].

A recent extension to Whittaker's multiplicative β diversity partition enables linking changes in average α diversity ($\overline{\Delta\alpha}$), γ, and β diversity (in log scale) to changes in the numbers of spatially restricted and widespread species (Fig. 1)[12,38]. This framework illustrates how nutrient enrichment

could cause biotic homogenization if local communities gain widespread species (Fig. 1: scenario I), if spatially restricted species are replaced by widespread species (Fig. 1: scenario II), or if the number of spatially restricted species decreases (Fig. 1: scenario III). Conversely, nutrient enrichment could cause biotic differentiation if local communities lose widespread species (Fig. 1: scenario IV), if widespread species are replaced by spatially restricted species (Fig. 1: scenario V), or if the number of spatially restricted species increases (Fig. 1: scenario VI). Finally, if gains or losses of species at the α and γ scale are similar (i.e., approximately equal or proportional), then we would observe no change in β diversity (1:1 diagonal line in Fig. 1). However, clear links between changes in diversity across spatial scales and changes in the number of spatially restricted and widespread species under nutrient enrichment are yet to be made.

Here, we use this framework to synthesize scale-dependent plant diversity change (for the entire community and groups of species) under nutrient enrichment using a long-term standardized experiment in 72 grasslands distributed across six continents (i.e., NutNet[39]; Fig. S1; Table S1). We use two treatments: Ambient (Control) and fertilization by nitrogen, phosphorus, and potassium together (i.e., NPK). Nutrients were added at a rate of 10 g m^{-2} annually. Treatments were randomly assigned to 5 m × 5 m plots and were replicated in three or more blocks. Species cover was recorded in one 1 m × 1 m permanent subplot using a standardized protocol. At each site, α diversity is determined as the number of species in each permanent subplot (i.e., species richness), and γ diversity as the total number of species occurring in three permanent subplots (for each treatment separately). We exclude additional blocks from sites that have more than three because γ and β diversity depend on the number of local communities used. We calculate Δα as the richness difference in local communities (subplots) and Δγ as the difference in the sum of the subplots under nutrient addition relative to that of control on the log scale. That is, $\Delta\alpha = \log(\alpha_{NPK}/\alpha_{Control})$ and $\Delta\gamma = \log(\gamma_{NPK}/\gamma_{Control})$. We then calculate Δβ as Δγ minus $\overline{\Delta\alpha}$, where $\overline{\Delta\alpha}$ is the average of Δα over three blocks. Overall, we find proportionally similar species loss at local and larger scales, suggesting similar magnitudes of declines of spatially restricted and widespread species. Thus, we find no clear biotic homogenization or differentiation four years, and even up to 14 years, after nutrient additions began. Moreover, these overall patterns of little change in β diversity hold consistent across species groups.

Results and discussion

Changes in α-, γ-, and β diversity for the entire communities
Overall, adding nutrients decreased α and γ diversity, but it had no significant effects on β diversity (Δβ = 0.03; 95% credible interval: -0.02 to 0.08) (Fig. 2; Table S3), see also[27]. While we observed substantial variation in $\overline{\Delta\alpha}$, Δγ, and to a lesser extent, Δβ among sites (Table S4), we found no strong relationships between site-level $\overline{\Delta\alpha}$, Δγ, and Δβ and distance among blocks, drought intensity, grazing intensity, productivity, and/or the size of the species pool[24,25,34,40] (Fig. S3). On a site level, we found biotic homogenization at 24 sites, differentiation at 47 sites, and no change in β diversity at one site. However, the site-level 95% credible intervals (see Methods) overlapped 0 for all sites, suggesting no significant change in β diversity under nutrient addition (Table S4). Importantly, the overall effects of nutrient addition on α, γ, and β diversity were similar when we used effective numbers of species based on either Shannon diversity or Simpson diversity that account for species relative abundances[40] (Fig. S4; Table S3). Because species richness is more strongly influenced by rare species, while Shannon and Simpson diversity increasingly weigh abundant species, this result suggests that relatively rare and abundant species responded similarly to nutrient addition.

Gain of widespread species · Loss of widespread species

Spatially restricted replaced by widespread species · Widespread replaced by spatially restricted species

Loss of spatially restricted species · Gain of spatially restricted species

Fig. 2. Changes in average α, γ, and β diversity ($\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$) with nutrient addition. LRR: log response ratio. The white 1:1 diagonal line indicates no effects of nutrient addition on β diversity. Numbers in the parentheses are the number of sites. When a site has $\overline{\Delta\alpha}$ = 0, $\Delta\gamma$ = 0, or $\Delta\beta$ = 0, it was not counted into any of the six scenarios as shown in the framework. The small points represent site-level $\overline{\Delta\alpha}$ and $\Delta\gamma$ at 72 sites. The large open point and error bars are the estimated mean and 95% credible intervals for $\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$ across all sites. See Table S3 for model fit and estimated overall means and 95% credible intervals for $\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$. See Table S4 for site-level estimates and 95% credible intervals. Source data are provided as a Source Data file.

Changes in α-, γ-, and β diversity for species groups

The overall proportional species loss within the community at local and larger scales on average may result if different species groups have contrasting patterns of response to nutrient addition. For instance, this result could be the case if native species loss is greater at the larger spatial scale than at the local scale, while non-native species loss is lower at the larger than the local scale. To test this

possibility, we investigate changes in α, γ, and β diversity for native and non-native species
separately. Extending previous studies[6,41], we found that nutrient addition decreased native species
more than non-native species. Compared to non-native species, nutrient addition resulted in a 10%
greater reduction in α diversity and a 16% greater reduction in γ diversity of native plant species,
respectively (Table S5). The overall pattern of diversity change across spatial scales for native
species largely followed that of the entire communities with similar magnitudes of decline in α and
γ diversity and little change in β diversity (Δβ = 0; 95% credible interval: -0.05 to 0.06; Fig. 3A).
For non-native species, overall, nutrient addition decreased α diversity 6% more than γ diversity
(Table S5). But nutrient addition had no significant effects on β diversity of non-native species (Δβ
= 0.04; 95% credible interval: -0.05 to 0.14; Fig. 3B).
227

228
Fig. 3. Changes in average α, γ, and β diversity ($\overline{\Delta\alpha}$, Δγ, and Δβ) with nutrient addition for
native and non-native species groups. (A) native and (B) non-native species. LRR: log response
ratio. The white 1:1 diagonal line indicates no effects of nutrient addition on β diversity. Numbers
in the parentheses are the number of sites. When a site has $\overline{\Delta\alpha}$= 0, Δγ = 0, or Δβ = 0, it was not
counted into any of the six scenarios as shown in the framework. The small points represent site-
level $\overline{\Delta\alpha}$ and Δγ. The large open point and error bars are the estimated mean and 95% credible
intervals for $\overline{\Delta\alpha}$, Δγ, and Δβ across all sites. See Table S5 for model fit and estimated overall means
and 95% credible intervals for $\overline{\Delta\alpha}$, Δγ, and Δβ. Source data are provided as a Source Data file.
237

Fig. 4. Changes in average α, γ, and β diversity ($\overline{\Delta\alpha}$, Δγ, and Δβ) with nutrient addition for different functional species groups. (A) forb, (B) graminoid, (C) legume, and (D) woody species. LRR: log response ratio. The white 1:1 diagonal line indicates no effects of nutrient addition on β diversity. Numbers in the parentheses are the number of sites. When a site has $\overline{\Delta\alpha}$= 0, Δγ = 0, or Δβ = 0, it was not counted into any of the six scenarios as shown in the framework. The small points represent site-level $\overline{\Delta\alpha}$ and Δγ. The large open point and error bars are the estimated mean and 95% credible intervals for $\overline{\Delta\alpha}$, Δγ, and Δβ across all sites. See Table S6 for model fit and estimated overall means and 95% credible intervals for $\overline{\Delta\alpha}$, Δγ, and Δβ. Source data are provided as a Source Data file.

We also separated species into graminoid, forb, legume, and woody species to investigate scale-dependent diversity change within species groups. Nutrient addition led to the greatest reduction of α diversity in forb species and of γ diversity in woody species (Table S6). Similar to that of entire

communities, nutrient addition decreased α and γ diversity by similar magnitudes and it had no effects on β diversity for graminoid species (Δβ = 0.01; 95% credible interval: -0.04 to 0.05; Fig. 4B; Table S6) and legume species (Δβ = 0.00; 95% credible interval: -0.17 to 0.18; Fig. 4C; Table S6). Overall, nutrient addition decreased α diversity 8% more than γ diversity for forb species, while it decreased γ diversity 11% more than α diversity for woody species (Table S6). However, nutrient addition also did not have significant effects on β diversity for forb species (Δβ = 0.09; 95% credible interval: -0.02 to 0.19; Fig. 4C; Table S6). Nutrient addition caused a weak biotic homogenization for woody species (Δβ = -0.14; 95% credible interval: -0.30 to 0.003; Table S6), this was primarily linked to loss of spatially restricted species (Fig. 4D).

Robustness and limitations

We tested the robustness of our results by performing multiple sensitivity tests. We redid the analyses for the effects of nutrient addition on α, γ, and β diversity for the entire communities as well as for species groups using a subset of 16 sites that had data 14 years after nutrient additions began (Fig. S5 ~ Fig. S7). In all, we found that the overall effects were largely similar in these longer-term sites to that of 72 sites that had data four years after nutrient additions began. Because three spatial blocks may be somewhat limited in spatial extent for estimating effects on β diversity, we tested whether including more blocks to create the γ scale altered our results. We redid the analyses for the effects of nutrient addition on α, γ, and β diversity for the entire communities as well as for species groups using using 11 sites that had five spatial blocks (Fig. S8~Fig. S10). Again, the overall effects were largely similar to that we found for the full analysis of the 72 sites with three blocks.

Despite our evidence for little change in β diversity under nutrient addition across the entire experiment, we recognize limitations of extrapolating these results to the landscape scale (e.g., >1 km × 1 km). This is because many ecological processes that directly and indirectly influence plant diversity can be very different at the landscape scales[14,42]. The scale at which we inferred changes in the number of spatially restricted and widespread species, by examining how many local communities they were lost from, is a relatively small spatial scale. Linking estimates of species' geographic range size and other key traits with changes in plant diversity across larger spatial scales[19,42] will deepen our understanding of the mechanisms of diversity change.

To summarize, we apply a framework to a globally distributed long-term experiment to provide a comprehensive synthesis of the impact of nutrient addition on scale-dependent plant diversity change in grasslands. The framework links changes in species richness across scales to that changes in the numbers of spatially restricted and widespread species. Overall, we found similar proportional plant diversity declines at local and larger spatial scales under nutrient addition, and little evidence for either biotic homogenization or differentiation within sites. These overall patterns were largely consistent for diversity metrics that incorporate relative species covers, across species groups, and over long time periods. This demonstrates that nutrient enrichment poses a potential threat to all plant species groups, including widespread and native species that often drive ecosystem functions and services.

Methods

Experimental setup

The experimental sites used in this study are part of the Nutrient Network (NutNet, Fig. S1 and Table S1). The experimental design includes a factorial manipulation of nutrients (N, P, and K) plus two fences to exclude herbivores, see ref[39] for more details. For the analyses here, we used plots under two treatments: Ambient (Control) and fertilization by nitrogen, phosphate, and potassium together (i.e., NPK). Treatments were randomly assigned to 5 m × 5 m plots and were replicated in

three or more blocks. A micronutrient mix consists of Fe (15%), S (14%), Mg (1.5%), Mn (2.5%),
Cu (1%), Zn (1%), B (0.2%), and Mo (0.05%) was added once only at the start of the experiment
(i.e., year 1) for the nutrient addition plots, but not in subsequent years to avoid toxicity. Nitrogen,
phosphate, potassium were added annually before the growing season of each treatment year at
most sites. Nitrogen was added as $10 \, \text{g m}^{-2} \, \text{yr}^{-1}$ time-release urea [$(NH_2)_2CO$], phosphate was added
as $10 \, \text{g m}^{-2} \, \text{yr}^{-1}$ triple-super phosphate [$Ca(H_2PO_4)_2$], while potassium was added as $10 \, \text{g m}^{-2} \, \text{yr}^{-1}$
sulfate [K_2SO_4].
310
311 Data were retrieved from the NutNet database in November 2023. We analyzed data from 72 sites
312 where 1) nutrients were applied for at least four years; and 2) each site had at least three blocks.
313 These sites are distributed across six continents and include a wide range of grassland types. See
314 Fig. S1 and Table S1 for details of geolocation, grassland types, and experimental years used.
315
Sampling protocol
Scientists at NutNet sites followed standard sampling protocols[39]. Specifically, a 1 m × 1 m subplot
within each plot was permanently marked for annual recording of plant species composition.
Species cover (%) was estimated visually for individual species in the subplots; thus the total cover
of living plants may sometimes exceed 100% for multilayer canopies. At most sites, cover was
recorded once per year at peak biomass. At some sites with strong seasonality, cover was recorded
twice per year to include a complete list of species. For those sites, the maximum cover for each
species and total biomass were used in the analyses. When taxa could not be identified to the
species level, they were aggregated at the genus level but referred to as "species" for simplicity.
325
Quantifying changes in α, γ, and β diversity
We measured α and γ diversity using species richness (i.e., number of species) because it is the
most commonly examined diversity metric[43]. At each site, α diversity was estimated as the number
of species in each permanent subplot (1 m × 1 m), and γ diversity as the total number of species
occurring in three permanent subplots (for each treatment separately). To standardize sampling
effort, for sites with more than three blocks, we selected the first three blocks according to the block
number recorded by site PIs. The framework relies on Whittaker's multiplicative β diversity
partition, and it quantifies β diversity using the effective number of communities[12]. As such, if all
subplots share the same species, then β diversity would equal to one. In contrast, if each subplot has
unique species, then β diversity would equal to three. We calculated Δα as the richness difference in
local communities (subplots) and Δγ as the difference in the sum of the subplots under nutrient
addition relative to that of control treatment on the log scale. That is, $\Delta\alpha = \log(\alpha_{NPK}/\alpha_{Control})$ and $\Delta\gamma$
$= \log(\gamma_{NPK}/\gamma_{Control})$. We calculated Δβ as Δγ minus $\overline{\Delta\alpha}$, where $\overline{\Delta\alpha}$ is the average of Δα over three
blocks. A decrease in Δβ indicates nutrient addition causes species composition to be more similar
among three subplots than that among control subplots. Because sites are not evenly distributed
around the world, many sites are aggregated in North America, we checked spatial autocorrelation
of diversity change under nutrient addition using Moran's I[44]. We found that $\overline{\Delta\alpha}$, Δγ, and Δβ did not
appear to be more similar for sites that are closer to each other (Table S2).
344
We fitted multilevel (also referred as mixed effects or hierarchical) models for Δα, Δγ, and Δβ (as
the response variable; all on the log scale) separately. We included random intercept for each site,
model was coded as: richness change ~ 1 + (1 |sites) to estimate site-level variation. We used
Bayesian analysis because it yields full posterior distributions of parameters rather than point
estimates and p-values, which provides a deeper understanding of the uncertainty and variability in
the results[45]. Models described above were fitted using the Hamiltonian Monte Carlo (HMC)
sampler in Stan and coded using the package 'brms' (version 2.21.0) in R (version 4.4.1)[46,47].
Models were fitted without explicitly specifying priors, allowing brms to assign its default priors.
Models were fitted with 6 chains and 3000 iterations (1000 iterations for warm up). Visual
inspection of the HMC chains and Rhat summaries showed model convergence (all Rhats < 1.03;

Table S3; Table S5; Table S6). We visually checked posterior predictive plots to determine how
well models can reproduce the data (Fig. S2).
357
To examine whether diversity changes were sensitive to species relative covers, we redid the above
analyses (i.e., based on species richness) using Shannon diversity and Simpson diversity (both
converted to effective numbers)[48](Fig. S4). Species richness is most sensitive to rare species,
followed by Shannon diversity, and Simpson diversity is more sensitive to the numbers of
relatively abundant species. We calculated the exponential of Shannon diversity and the inverse
form of Simpson diversity using the R package vegan (version 2.6-6.1)[49]. These three diversity
metrics equal to diversity with order q = {0, 1, 2}, where increasing q decreases the influence of
rare species, and $D_q = \left(\sum_{i=1}^{s} p_i^q\right)^{1/(1-q)}$, where p is the relative cover of species i, s is the total
number of species. These diversity metrics are also referred to as Hill numbers[48,50].
367
Site covariates. We investigated whether the effects of nutrient addition on $\overline{\Delta\alpha}$, γ, and β diversity
based on species richness were mediated by site characteristics. We included site characteristics that
have been shown in previous literature to influence Δα, Δγ, and Δβ in grasslands: site species pool,
site productivity, drought intensity, and grazing intensity[24,25,34,40]. We quantified drought intensity
as the sum of annual evapotranspiration/precipitation, and averaged it from year 0 to 4 at each site.
Precipitation and potential evapotranspiration used to calculate SPEI were downloaded from
https://crudata.uea.ac.uk/cru/data/hrg/cru_ts_4.07/. We quantified the site species pool as the total
number of species and site productivity as the average aboveground biomass from year 0 to 4 under
the control treatment in the three blocks at each site. Aboveground biomass was harvested within
two 1 × 0.1 m strips (in total 0.2 m^2), strips were moved from year to year to avoid resampling the
same location. For subshrubs and shrubs occurring within strips, we collected all leaves and current
year's woody growth. All biomass was dried at 60 °C (to constant mass) before weighing to the
nearest 0.01 g. We used published methods to quantify an integrated grazing intensity metric from
vertebrate herbivores at each site. Specifically, herbivore species (> 2 kg) that consume grassland
biomass were documented at each site by site PIs, and each species was assigned an importance
value from 1 (present, but low impact and frequency) to 5 (high impact and frequency). An index
value was calculated for each site as the sum of herbivore importance values for all herbivores
following ref[51,52]. We also investigated relationships between change in diversity and distance
among blocks, because species composition may become less similar as the distance between
sampled communities increases. The average pairwise distance among the three blocks within sites
ranged from 23.04 to 12538.09 m, with a mean of 513.01 m and a median of 118.7 m across 54
sites that have geolocation data for each block. We first calculated three Euclidean distances
between pairs of blocks, we then used the mean of these pairwise distances as the average distance
among blocks. We used the average distance among blocks instead of area, because blocks are
arranged in parallel at some sites. We fitted linear regression models with $\overline{\Delta\alpha}$, Δγ, and Δβ as the
response variable separately, and each of the site characteristics was used as a predictor variable.
394
Species groups. We then investigated the effects of nutrient addition on α, γ, and β diversity within
groups of species with similar characteristics following the method for changes in α, γ, and β
diversity in the entire communities. We eliminated sites where no species occurred in control,
nutrient addition, or both plots for a particular group because the value of the log (0) is undefined.
We ran the analyses separately for native and non-native species. Native and non-native species
were classified by site PIs. Then, we investigated effects of nutrient addition on species richness for
different life forms including forb, graminoid, legume, and woody species because previous studies
have shown that different life forms may show distinct responses to nutrient addition[6,11,53].
403
Sensitivity test. We tested whether effects of nutrient addition on species richness across spatial
scales depend on experimental duration because a few single-site experiments have shown that the

effects of nutrient additions on changes in diversity, especially β diversity, may take several years
to emerge[29,31]. To that end, we used a subset of 16 sites that had data 14 years after nutrient
additions began. Also, because three blocks may be limited in spatial extent, we tested whether
combining more blocks to create the γ scale would alter our results. We redid the analyses using
data from 11 sites that had five spatial blocks.
411
412 **Data availability**: The species cover and species richness data, site abiotic and biotic
413 environmental data used and generated in this study have been deposited in the Figshare database
414 and are publicly available (https://doi.org/10.6084/m9.figshare.26412295.v3). The NutNet data are
415 publicly available on the Environmental Data Initiative (EDI)
416 (https://portal.edirepository.org/nis/advancedSearch.jsp). Source data are provided with this paper.
417
Code Availability
419 The R codes used to produce results in this study have been deposited in the GitHub
420 (https://github.com/chqq365/plant-diversity-and-biotic-homogenization.git) and archived through
421 Zenodo (https://doi.org/10.5281/zenodo.14902812).
422

423 **References:**

1. Sardans, J. & Peñuelas, J. Potassium: a neglected nutrient in global change. *Global Ecology and Biogeography* **24**, 261–275 (2015).
2. Galloway, J. N., Bleeker, A. & Erisman, J. W. The Human Creation and Use of Reactive Nitrogen: A Global and Regional Perspective. *Annual Review of Environment and Resources* **46**, 255–288 (2021).
3. Yuan, Z. *et al.* Human Perturbation of the Global Phosphorus Cycle: Changes and Consequences. *Environ. Sci. Technol.* **52**, 2438–2450 (2018).
4. Bobbink, R. *et al.* Global assessment of nitrogen deposition effects on terrestrial plant diversity: A synthesis. *Ecological Applications* **20**, 30–59 (2010).
5. Seabloom, E. W. *et al.* Increasing effects of chronic nutrient enrichment on plant diversity loss and ecosystem productivity over time. *Ecology* **102**, 1–11 (2021).
6. Suding, K. N. *et al.* Functional- and abundance-based mechanisms explain diversity loss due to N fertilization. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 4387–4392 (2005).
7. Yang, Z., Hautier, Y., Borer, E. T., Zhang, C. & Du, G. Abundance- and functional-based mechanisms of plant diversity loss with fertilization in the presence and absence of herbivores. *Oecologia* **179**, 261–270 (2015).
8. Zhang, P. *et al.* Space resource utilization of dominant species integrates abundance- and functional-based processes for better predictions of plant diversity dynamics. *Oikos* **2023**, e09519 (2023).
9. MacDougall, A. S. *et al.* The Neolithic Plant Invasion Hypothesis: the role of preadaptation and disturbance in grassland invasion. *New Phytologist* **220**, 94–103 (2018).
10. Seabloom, E. W. *et al.* Plant species' origin predicts dominance and response to nutrient enrichment and herbivores in global grasslands. *Nat Commun* **6**, 7710 (2015).
11. Tognetti, P. M. *et al.* Negative effects of nitrogen override positive effects of phosphorus on grassland legumes worldwide. *Proceedings of the National Academy of Sciences* **118**, (2021).
12. Blowes, S. A. *et al.* Synthesis reveals approximately balanced biotic differentiation and homogenization. *Science Advances* **10**, eadj9395 (2024).
13. Keil, P. *et al.* Spatial scaling of extinction rates: Theory and data reveal nonlinearity and a major upscaling and downscaling challenge. *Global Ecology and Biogeography* **27**, 2–13 (2018).
14. Chase, J. M. *et al.* Species richness change across spatial scales. *Oikos* **128**, 1079–1091 (2019).
15. Daleo, P. *et al.* Environmental heterogeneity modulates the effect of plant diversity on the

spatial variability of grassland biomass. *Nat Commun* **14**, 1809 (2023).

16. Le Provost, G. *et al.* The supply of multiple ecosystem services requires biodiversity across spatial scales. *Nat Ecol Evol* **7**, 236–249 (2022).

17. Daru, B. H. *et al.* Widespread homogenization of plant communities in the Anthropocene. *Nat Commun* **12**, 6983 (2021).

18. Gossner, M. M. *et al.* Land-use intensification causes multitrophic homogenization of grassland communities. *Nature* **540**, 266–269 (2016).

19. Staude, I. R. et al. Replacements of small- by large-ranged species scale up to diversity loss in Europe's temperate forest biome. Nat Ecol Evol 4, 802–808 (2020).

20. McKinney, M. L. & Lockwood, J. L. Biotic homogenization: a few winners replacing many losers in the next mass extinction. *Trends in Ecology & Evolution* **14**, 450–453 (1999).

21. Chen, Q. *et al.* Long-term cross-scale comparison of grazing and mowing on plant diversity and community composition in a salt-marsh system. *Journal of Ecology* **109**, 3737–3747 (2021).

22. Huston, M. A General Hypothesis of Species Diversity. *The American Naturalist* **113**, 81–101 (1979).

23. Tilman, D. Plant strategies and the dynamics and structure of plant communities. Princeton University Press. (1988)

24. Seabloom, E. W. *et al.* Species loss due to nutrient addition increases with spatial scale in global grasslands. *Ecology Letters* **24**, 2100–2112 (2021).

25. Chalcraft, D. R. *et al.* Scale-dependent responses of plant biodiversity to nitrogen enrichment. *Ecology* **89**, 2165–2171 (2008).

26. Lu, W. *et al.* Similar effects of seasonal nitrogen enrichment on plant species–area relationship in a temperate grassland. *Ecological Indicators* **158**, 111430 (2024).

27. Chen, Q. *et al.* Nutrients and herbivores impact grassland stability across spatial scales through different pathways. *Global Change Biology* **28**, 2678–2688 (2022).

28. Zhou, X., Liu, X., Zhang, P., Guo, Z. & Du, G. Increased community compositional dissimilarity alleviates species loss following nutrient enrichment at large spatial scales. *Journal of Plant Ecology* **12**, 376–386 (2019).

29. DeSiervo, M. H., Sullivan, L. L., Kahan, L. M., Seabloom, E. W. & Shoemaker, L. G. Disturbance alters transience but nutrients determine equilibria during grassland succession with multiple global change drivers. *Ecology Letters* **26**, 1132–1144 (2023).

30. Eskelinen, A. & Harrison, S. Erosion of beta diversity under interacting global change impacts in a semi-arid grassland. *Journal of Ecology* **103**, 397–407 (2015).

31. Inouye, R. S. & Tilman, D. Convergence and Divergence of Old-Field Vegetation After 11 yr of Nitrogen Addition. *Ecology* **76**, 1872–1887 (1995).

32. Zhang, Y. *et al.* Nitrogen addition does not reduce the role of spatial asynchrony in stabilising grassland communities. *Ecology Letters* **22**, 563–571 (2019).

33. Foster, B. L. *et al.* Seed availability constrains plant species sorting along a soil fertility gradient. *Journal of Ecology* **99**, 473–481 (2011).

34. Koerner, S. E. *et al.* Nutrient additions cause divergence of tallgrass prairie plant communities resulting in loss of ecosystem stability. *Journal of Ecology* **104**, 1478–1487 (2016).

35. Lan, Z. *et al.* Testing the scaling effects and mechanisms of N-induced biodiversity loss: evidence from a decade-long grassland experiment. *Journal of Ecology* **103**, 750–760 (2015).

36. Xu, Z. *et al.* Antithetical effects of nitrogen and water availability on community similarity of semiarid grasslands: evidence from a nine-year manipulation experiment. *Plant Soil* **397**, 357–369 (2015).

37. Yang, X. *et al.* Resource addition drives taxonomic divergence and phylogenetic convergence of plant communities. *Journal of Ecology* **107**, 2121–2132 (2019).

38. Whittaker, R. H. Vegetation of the Siskiyou Mountains, Oregon and California. **30**, 279–338 (1960).

39. Borer, E. T. *et al.* Finding generality in ecology: A model for globally distributed experiments. *Methods in Ecology and Evolution* **5**, 65–73 (2014).

40. Chase, J. M. & Leibold, M. A. Spatial scale dictates the productivity-biodiversity relationship. *Nature* **416**, 427–430 (2002).

41. Seabloom, E. W. *et al.* Predicting invasion in grassland ecosystems: Is exotic dominance the real embarrassment of richness? *Global Change Biology* **19**, 3677–3687 (2013).

42. Staude, I. R. *et al.* Directional turnover towards larger-ranged plants over time and across habitats. *Ecology Letters* **25**, 466–482 (2022).

43. Chase, J. M. *et al.* Embracing scale-dependence to achieve a deeper understanding of biodiversity and its change across communities. *Ecology Letters* **21**, 1737–1751 (2018).

44. Getis, A. A History of the Concept of Spatial Autocorrelation: A Geographer's Perspective. *Geographical Analysis* **40**, 297–309 (2008).

45. Kruschke, J. K. Bayesian estimation supersedes the t test. *Journal of Experimental Psychology: General* **142**, 573–603 (2013).

46. Bürkner, P.-C. Advanced Bayesian Multilevel Modeling with the R Package brms. *The R Journal* **10**, 395 (2018).

47. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. (2022).

48. Jost, L. Entropy and diversity. *Oikos* **113**, 363–375 (2006).

49. Oksanen, A. J. *et al.* Package ' vegan '. (2019).

50. Hill, M. O. Diversity and Evenness: A Unifying Notation and Its Consequences. *Ecology* **54**, 427–432 (1973).

51. Borer, E. T. *et al.* Nutrients cause grassland biomass to outpace herbivory. *Nature Communications* **11**, 1–8 (2020).

52. Anderson, T. M. *et al.* Herbivory and eutrophication mediate grassland plant nutrient responses across a global climatic gradient. *Ecology* **99**, 822–831 (2018).

53. Van Sundert, K. *et al.* Towards comparable assessment of the soil nutrient status across scales—Review and development of nutrient metrics. *Global Change Biology* **26**, 392–409 (2020).

Acknowledgements

This work was generated using data from the Nutrient Network (http://www.nutnet.org) experiment, funded at the site-scale by individual researchers. We thank the Minnesota Supercomputer Institute for hosting project data and the Institute on the Environment for hosting Network meetings. We thank researchers from the NutNet who contributed data to our analysis, table S7 lists these researchers. We thank Shuli Niu for constructive suggestions to improve our manuscript. Nitrogen fertilizer was donated to NutNet by Crop Production Services, Loveland, CO.

Funding: German Center for Integrative diversity Research (iDiv) Halle-Jena-Leipzig (DFG FZT 118-202548816) for the contribution of J.M.C., S.A.B., E.L., Q.C., W.S.H., and other iDiv co-authors. ERC Advanced Grant (MetaChange) to J.M.C. for the contribution of J.M.C., S.A.B.. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

National Science Foundation grant NSF-DEB-1042132 (E.T.B., E.W.S.; for NutNet coordination and data management) National Science Foundation grant NSF-DEB-1234162 (E.T.B., E.W.S.; for Long-Term Ecological Research at Cedar Creek). National Science Foundation grant NSFDEB-1831944 (E.T.B., E.W.S.; for Long-Term Ecological Research at Cedar Creek), and the Institute on the Environment (DG-0001-13; E.T.B., E.W.S.).

Author Contributions Statement

Q.C., J.M.C., S.A.B., E.L., W.S.H. conceived the study; Q.C., J.M.C., S.A.B. developed the methodology; Q.C. analyzed data with contribution from S.A.B.; W.S.H., E.T.B., A.M., J.P.M., J.D.B., P.M.T., E.W.S., P.D., S.A.P., C.R., P.BA., I.D., G.W., C.S., G.F.V., A.C.R., G.M.W., Y.H., C.E., Y.N., P.L.P., A.E., D.S.G., H.O., C.D., M.W.C., S.H., N.E., J.C., R.V., J.W.M., M.T., S.B.,

M.C.M., M.B., J.M.H.K., C.R.D., N.H., A.J., P.M., B.B.O., L.L. contributed data; Q.C. visualized the results; Q.C. wrote the original draft of the manuscript; Q.C., S.A.B., W.S.H., E.L., E.T.B., A.M., J.P.M., J.D.B., P.M.T., E.W.S., P.D., S.A.P., C.R., P.B.A., I.D., G.W., C.S., G.F.V., A.C.R., G.M.W., Y.H., C.E., E.H., Y.N., P.L.P., A.E., D.S.G., H.O., C.D., M.W.C., S.H., N.E., J.C., R.V., J.W.M., M.T., S.B., M.C.M., M.B., J.M.H.K., C.R.D., N.H., A.J., P.M., B.B.O., L.L. J.M.C. reviewed and edited the manuscript. See table S8 for more details for the contribution of each author.

Competing Interests Statement
Authors declare that they have no competing interests.