Title: Local nutrient addition drives plant diversity losses but not biotic homogenization in global grasslands

Authors:

Qingqing Chen[1,2*], Shane A. Blowes[1,3], W. Stanley Harpole[1,4,5], Emma Ladouceur[1,3], Elizabeth T. Borer[6], Andrew MacDougall[7], Jason P. Martina[8], Jonathan D. Bakker[9], Pedro M. Tognetti[10,11], Eric W. Seabloom[6], Pedro Daleo[12], Sally A. Power[13], Christiane Roscher[1,4], Peter Adler[14], Ian Donohue[15], George Wheeler[16], Carly Stevens[17], Ciska Veen[18], Anita C. Risch[11], Glenda M. Wardle[19], Yann Hautier[20], Catalina Estrada[21], Erika Hersch-Green[22], Yujie Niu[23], Pablo L. Peri[24], Anu Eskelinen[1,4,25], Daniel S Gruner[26], Harry Olde Venterink[27], Carla D'Antonio[28], Marc W. Cadotte[29], Sylvia Haider[30], Nico Eisenhauer[1,31], Jane Catford[32,33], Risto Virtanen[25], John W. Morgan[34], Michelle Tedder[35], Sumanta Bagchi[36], Maria C. Caldeira[37], Miguel Bugalho[38], Johannes M. H. Knops[39], Chris R. Dickman[40], Nicole Hagenah[41], Anke Jentsch[23], Petr Macek[42,44], Brooke B. Osborne[43], Lauri Laanisto[44], Jonathan M. Chase[1,3]

Affiliations:

1. German Centre for Integrative diversity Research (iDiv) Halle-Jena-Leipzig.
2. Senckenberg Museum for Natural History Görlitz, D-02826, Görlitz, Germany.
3. Department of Computer Science, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany.
4. Helmholtz Center for Environmental Research – UFZ, Department of Physiological Diversity, Permoserstrasse 15, 04318 Leipzig, Germany.
5. Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108 Halle (Saale), Germany.
6. Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN 55108 USA.
7. Department of Integrative Biology, University of Guelph, Guelph, Ontario, Canada N1G2W1.
8. Department of Biology, Texas State University, San Marcos, TX, 78666 USA.
9. School of Environmental and Forest Sciences, University of Washington, Seattle, WA USA.
10. Instituto de Investigaciones Fisiológicas y Ecológicas Vinculadas a La Agricultura (IFEVA), Facultad de Agronomía, Universidad de Buenos Aires and CONICET, Buenos Aires, Argentina.
11. Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland.
12. Instituto de Investigaciones Marinas y Costeras (IIMyC), UNMdP-CONICET, CC 1260 Correo Central, Mar Del Plata B7600WAG, Argentina.
13. Hawkesbury Institute for the Environment, Locked Bag 1797, Penrith, New South Wales, 2751, Australia.
14. Department of Wildland Resources and the Ecology Center, Utah State University, Logan, UT, 84322 USA.
15. Zoology, School of Natural Sciences, Trinity College Dublin, Dublin, Ireland.
16. University of Nebraska-Lincoln, Lincoln, NE, USA.
17. Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK.
18. Department of Terrestrial Ecology, Netherlands Institute of Ecology, Droevedaalsesteeg 10, 6708 PB, Wageningen, The Netherlands.
19. Desert Ecology Research Group, School of Life and Environmental Sciences, ARC Training Centre in Data Analytics for Resources and Environments (DARE), The University of Sydney, NSW 2006, Australia.
20. Ecology and diversity Group, Department of Biology, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands.
21. Department of Life Sciences, Imperial College London, Silwood Park Campus, Buckhurst Road, Ascot SL5 7PY, UK.
22. Department of Biological Sciences, Michigan Technological University, Houghton, MI 49930,

USA.

23. Disturbance Ecology and Vegetation Dynamics, Bayreuth Center of Ecology and
Environmental Research (BayCEER), University of Bayreuth, Germany.

24. Instituto Nacional de Tecnología Agropecuaria (INTA), Universidad Nacional de La Patagonia
Austral (UNPA).CONICET. Río Gallegos, Santa Cruz, Argentina.

25. Ecology & Genetics Unit, University of Oulu, Finland.

26. Department of Entomology, University of Maryland, College Park, MD USA.

27. Vrije Universiteit Brussel, Department Biology, WILD, Pleinlaan 2, 1050 Brussels, BELGIUM.

28. Dept. of Biological Sciences, University of Toronto Scarborough, Toronto, ON, Canada.

29. University of California, Santa Barbara.

30. Institute of Ecology, Leuphana University of Lüneburg, Universitätsallee 1, 21335 Lüneburg,
Germany.

31. Institute of Biology, Leipzig University; Puschstrasse 4, 04103 Leipzig, Germany.

32. Department of Geography, King's College London, 30 Aldwych, London, WC2B 4BG, UK.

33. School of Ecosystem & Forest Sciences, University of Melbourne, Parkville, VIC 3010,
Australia.

34. Department of Environment & Genetics, La Trobe University, Bundoora 3083, Victoria,
Australia.

35. Centre for Functional diversity, School of Life Sciences, University of KwaZulu-Natal,
Pietermaritzburg, South Africa.

36. Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India.

37. Forest Research Centre, Associate Laboratory TERRA, School of Agriculture, University of
Lisbon, Portugal.

38. Centre for Applied Ecology 'Prof. Baeta Neves' (CEABN-InBIO), School of Agriculture,
University of Lisbon, Portugal.

39. Health & Environmnetal Sciences Department, Xi'an Jiaoting-Liverpool University, Suzhou,
Jiangsu, China.

40. Desert Ecology Research Group, School of Life and Environmental Sciences, The University of
Sydney, Sydney, NSW 2006, Australia.

41. Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria,
Pretoria, South Africa.

35. Centre for Functional diversity, School of Life Sciences, University of KwaZulu-Natal,
Pietermaritzburg, South Africa.

36. Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India.

37. Forest Research Centre, Associate Laboratory TERRA, School of Agriculture, University of
Lisbon, Portugal.

38. Centre for Applied Ecology 'Prof. Baeta Neves' (CEABN-InBIO), School of Agriculture,
University of Lisbon, Portugal.

39. Health & Environmnetal Sciences Department, Xi'an Jiaoting-Liverpool University, Suzhou,
Jiangsu, China.

40. Desert Ecology Research Group, School of Life and Environmental Sciences, The University of
Sydney, Sydney, NSW 2006, Australia.

41. Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria,
Pretoria, South Africa.

42.   Institute of Hydrobiology, Biology Centre of the Czech Academy of Sciences, Na Sadkach 7,
Ceske Budejovice 370 05, Czech Republic

43.   Department of Environment and Society, Utah State University, Moab, UT, USA

44.   Chair of Diversity and Nature Tourism, Estonian University of Life Sciences, Kreutzwaldi 5,
51006, Tartu, Estonia

*corresponding author: chqq365@gmail.com

Abstract:

Nutrient enrichment typically causes local plant diversity decline. A common but untested expectation is that it reduces variation in nutrient conditions among localities and selects for a smaller pool of species, causing greater diversity declines at larger scales and biotic homogenization. We apply a topology linking changes in species richness across scales that characterizes changes in the numbers of spatially restricted and widespread species to a standardized experiment adding nutrients across 72 grasslands on six continents. Overall, we found similar declines of spatially restricted and widespread species, and proportional species loss at local and larger scales, and no biotic homogenization after 4 years and up to 14 years of treatment. These patterns of diversity changes were generally consistent across species groups (e.g., native species, graminoids and legume species). Thus, nutrient enrichment poses threats to plant diversity that includes widespread species that are often critical for ecosystem functions.

Main Text:

Agricultural fertilization practices and atmospheric nutrient deposition have led to increased availability and redistribution of soil nutrients globally[1–3]. At local scales (i.e., α diversity), nutrient enrichment tends to reduce plant diversity[4,5]. This diversity decline is typically ascribed to disproportionate losses of rare species (i.e., species with low cover) because small populations are more susceptible to extinction[6–8]. In addition, nutrient enrichment often leads to the removal of species with traits ill-suited for effectively competing in high nutrient conditions[6]. While species can vary widely in their nutrient requirements and tolerances, groups of species with similar (shared) characteristics can be lost from a flora. For example, native species are more likely to be lost than non-native species when nutrients are enriched because non-natives are often better-adapted to nutrient-rich conditions[6,9,10]. Similarly, nitrogen-fixing legumes may be more vulnerable than other species in high nutrient environments due to their decreased competitive advantage[6,11].

Despite clear evidence that nutrient enrichment causes losses of α diversity in grasslands[4,5], how these losses are reflected at larger spatial scales (i.e., γ diversity; calculated by aggregating local communities) is less straightforward[12–14]. Yet, it is diversity loss and change at larger spatial scales that is most often relevant for biodiversity conservation and management as well as for the provision of ecosystem functions and services[15,16]. It is often assumed that nutrient enrichment, like many other global changes, results in biotic homogenization (i.e., increasing similarity in composition among local communities, quantified as a decrease in β diversity)[17–21]. This is because local nutrient enrichment is expected to create homogeneous nutrient conditions among localities and consistently select for a smaller pool of species that are nutrient-demanding, fast-growing, and highly competitive for light[22,23]. Previous studies investigating scale-dependent diversity change under nutrient enrichment have tended to be short term and limited in scope[24–28]. While some experimental studies support the notion that nutrient enrichment leads to biotic homogenization[29–31], other studies have found no changes in β diversity[24,26,27,32] or even differentiation (i.e., increase in β diversity)[25,28,33–37].

A recent extension to Whittaker's multiplicative β diversity partition enables linking changes in average α diversity ($\overline{\Delta\alpha}$), γ, and β diversity (in log scale) to changes in the numbers of spatially restricted and widespread species (Fig. 1)[12,38]. This topology illustrates how nutrient enrichment could cause biotic homogenization if local communities gain widespread species (Fig. 1: scenario I), if spatially restricted species are replaced by widespread species (Fig. 1: scenario II), or if the number of spatially restricted species decreases (Fig. 1: scenario III). Conversely, nutrient enrichment could cause biotic differentiation if local communities lose widespread species (Fig. 1: scenario IV), if widespread species are replaced by spatially restricted species (Fig. 1: scenario V), or if the number of spatially restricted species increases (Fig. 1: scenario VI). Finally, if gains or losses of species at the α and γ scale are similar (i.e., approximately equal or proportional), then we would observe no change in β diversity (1:1 diagonal line in Fig. 1). However, clear links between changes in diversity across spatial scales and changes in the number of spatially restricted and widespread species under nutrient enrichment are yet to be made.

Here, we use this topology to synthesize scale-dependent plant diversity change (for the entire community and groups of species) under nutrient enrichment using a long-term standardized experiment in 72 grasslands distributed across six continents (i.e., NutNet[39]; Fig. S1; Table S1). We used two treatments: Ambient (Control) and fertilization by nitrogen, phosphorus, and potassium together (i.e., NPK). Nutrients were added at a rate of 10 g m^{-2} annually. Treatments were randomly assigned to 5 m × 5 m plots and were replicated in three blocks at most sites. Species cover was reecorded in one 1 m × 1 m permanent subplot using a standardized protocol. At each site, α diversity was determined as the number of species in each permanent subplot (i.e., species richness), and γ diversity as the total number of species occurring in three permanent subplots (for

each treatment separately). We excluded additional blocks from sites that had more than three
because γ and β diversity depend on the number of local communities used. We calculated Δα as the
richness difference in local communities (subplots) and Δγ as the difference in the sum of the
subplots under nutrient addition relative to that of control on the log scale. That is, Δα =
log(α$_{NPK}$/α$_{Control}$) and Δγ = log(γ$_{NPK}$/γ$_{Control}$). We then calculated Δβ as Δγ minus $\overline{\Delta\alpha}$, where $\overline{\Delta\alpha}$ is
the average of Δα over three blocks. In all, we found similar magnitudes of loss for spatially
restricted and widespread species, and proportional losses of species at local and larger scales, and
no clear biotic homogenization or differentiation four years, and even up to 14 years, after nutrient
additions began. Moreover, these overall patterns of little change in β diversity held consistent
across species groups.

Fig 1. A topology linking diversity changes at the α-, γ-, and β-scales to changes in the number
of spatially restricted and widespread species. $\overline{\Delta\alpha}$, Δγ, and Δβ are log response ratios (LRR) of
average α-, γ- and β- diversity under nutrient enrichment relative to that under ambient conditions
(control). Δβ is equal to zero along the 1:1 diagonal line. Below the diagonal line, Δγ < $\overline{\Delta\alpha}$, Δβ <
0, nutrient enrichment causes biotic homogenization. Above the diagonal line, Δγ > $\overline{\Delta\alpha}$, Δβ > 0,
nutrient enrichment causes biotic differentiation. Moreover, homogenization can be attributed to
three scenarios including I: gain of widespread species (Δγ < $\overline{\Delta\alpha}$, and Δγ >0, $\overline{\Delta\alpha}$ > 0); II:
spatially restricted species replaced by widespread species (Δγ < $\overline{\Delta\alpha}$, Δγ < 0 and $\overline{\Delta\alpha}$ > 0); III:
Loss of spatially restricted species (Δγ < $\overline{\Delta\alpha}$, and Δγ < 0, $\overline{\Delta\alpha}$ < 0). Conversely, differentiation

184 can be attributed to three scenarios that include IV: Loss of widespread species ($\Delta\gamma > \overline{\Delta\alpha}$, and $\Delta\gamma$
185 < 0, $\overline{\Delta\alpha}$ < 0); V: Widespread species replaced by spatially restricted species ($\Delta\gamma > \overline{\Delta\alpha}$, $\Delta\gamma > 0$
186 and $\overline{\Delta\alpha}$ < 0); VI: gain of spatially restricted species ($\Delta\gamma > \overline{\Delta\alpha}$, and $\Delta\gamma > 0$, $\overline{\Delta\alpha}$ > 0). Adapted
187 from[12].
188

Results and discussion
Overall, adding nutrients decreased α and γ diversity, but it had no significant effects on β diversity
($\Delta\beta = 0.03$; 95% credible interval: -0.02 to 0.07) (Fig. 2; Table S3), see also[27]. While we observed
substantial variation in $\overline{\Delta\alpha}$, $\Delta\gamma$, and to a lesser extent, $\Delta\beta$ among sites (Table S4), we found no
strong relationships between site-level $\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$ and distance among blocks, drought
intensity, grazing intensity, productivity, and/or the size of the species pool[24,25,34,40] (Fig. S3). On a
site level, we found biotic homogenization at 24 sites, differentiation at 47 sites, and no change in β
diversity at 1 site. However, the site-level 95% credible intervals (see Methods) overlapped 0 for all
sites, suggesting no significant change in β diversity under nutrient addition (Table S4).
Importantly, the overall effects of nutrient addition on α, γ, and β diversity was similar when we
used effective numbers of species based on either Shannon diversity or Simpson diversity that
account for species relative abundances[40] (Fig. S4; Table S3). Because species richness is more
strongly influenced by rare species, while Shannon and Simpson diversity increasingly weigh
common species, this suggests that relatively rare and common species responded similarly to
nutrient addition.

Legend:
- Gain of widespread species
- Loss of widespread species
- Spatially restricted replaced by widespread species
- Widespread replaced by spatially restricted species
- Loss of spatially restricted species
- Gain of spatially restricted species

Fig. 2. Effects of nutrient addition on α, γ, and β diversity. LRR: log response ratio. The white 1:1 diagonal line indicates no effects of nutrient addition on β diversity. Numbers in the parentheses are the number of sites. When a site has $\overline{\Delta\alpha} = 0$, $\Delta\gamma = 0$, or $\overline{\Delta\alpha} = 0$, $\Delta\gamma = 0$, and $\Delta\beta = 0$, it was not counted into any of the six scenarios. The small points represent site-level $\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$ at 72 sites. The large open point and error bars are the estimated mean and 95% credible intervals for $\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$ across all sites. See Table S3 for model fit and estimated overall means and 95% credible intervals for $\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$. See Table S4 for site-level estimates and 95% credible intervals.

The overall proportional species loss within the community at local and larger scales may result if different species groups have contrasting patterns. For instance, this could be the case if native species loss is greater at the larger spatial scale than at the local scale, while non-native species loss is lower at the larger than the local scale. To test this, we investigate changes in α, γ, and β diversity for native and non-native species separately. Extending previous studies[6,41], we found that nutrient addition decreased native species more than non-native species. Compared to non-native species,

nutrient addition resulted in a 9% greater reduction in α diversity and a 16% greater reduction in γ diversity of native plant species, respectively (Table S5). The overall pattern of diversity change across spatial scales for native species largely followed that of the entire communities with similar magnitudes of decrease in α and γ diversity and little change in β diversity (Δβ = 0; 95% credible interval: -0.06 to 0.06; Fig. 3A). For non-native species, overall, nutrient addition decreased α diversity 7% more than γ diversity (the effects on γ diversity was not significant; Table S5), but it had no significant effects on β diversity (Δβ = 0.04; 95% credible interval: -0.04 to 0.13; Fig. 3B).

Fig. 3. Effects of nutrient addition on α, γ, and β diversity for (A) native and (B) non-native species. LRR: log response ratio. The white 1:1 diagonal line indicates no effects of nutrient addition on β diversity. Numbers in the parentheses are the number of sites. When a site has $\overline{\Delta\alpha}$ = 0, Δγ = 0, or $\overline{\Delta\alpha}$ = 0, Δγ = 0, and Δβ = 0, it was not counted into any of the six scenarios. The small points represent site-level $\overline{\Delta\alpha}$, Δγ, and Δβ. See Table S5 for model fit and estimated overall means and 95% credible intervals for $\overline{\Delta\alpha}$, Δγ, and Δβ.

We also separated species into graminoid, forb, legume, and woody species to investigate scale-dependent diversity change. We found that nutrient addition led to the greatest reduction of α diversity in forb and legume species (Table S6). These results were consistent with expectations that legumes are more likely to be lost than other groups under nitrogen addition[6]. However, nutrient addition led to the greatest reduction of γ diversity in woody species (Table S6). Similar to that of entire communities, nutrient addition decreased α and γ diversity to similar magnitudes and it had no effects on β diversity for graminoid species (Δβ = 0.01; 95% credible interval: -0.04 to 0.05; Fig. 4B; Table S6) and legume species (Δβ = 0.01; 95% credible interval: -0.16 to 0.19; Fig. 4C; Table S6). Overall, nutrient addition decreased α diversity 7% more than γ diversity for forb species, while it decreased γ diversity 11% more than α diversity for woody species (Table S6). However, nutrient addition also did not have significant effects on β diversity for forb species (Δβ = 0.08; 95% credible interval: -0.02 to 0.18; Fig. 4C; Table S6) and woody species (Δβ = -0.14; 95% credible interval: -0.29 to 0.01; Fig. 4D; Table S6).

Fig. 4. Effects of nutrient addition on α, γ, and β diversity for (A) forb, (B) graminoid, (C) legume, and (D) woody species. LRR: log response ratio. The white 1:1 diagonal line indicates no effects of nutrient addition on β diversity. Numbers in the parentheses are the number of sites. When a site has $\overline{\Delta\alpha}$ = 0, $\Delta\gamma$ = 0, or $\overline{\Delta\alpha}$ = 0, $\Delta\gamma$ = 0, and $\Delta\beta$ = 0, it was not counted into any of the six scenarios. The small points represent site-level $\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$. See Table S6 for model fit and estimated overall means and 95% credible intervals for $\overline{\Delta\alpha}$, $\Delta\gamma$, and $\Delta\beta$.

We tested the robustness of our results by performing multiple sensitivity tests. We redid the analyses for the effects of nutrient addition on $\Delta\alpha$, $\Delta\gamma$, and $\Delta\beta$ for the entire communities as well as species groups using a subset of 29 sites that had data 10 years after nutrient additions began (Fig. S5 ~ Fig. S7), and another subset of 16 sites that had data 14 years after nutrient additions began (Fig. S8 ~ Fig. S10). In all, we found that the overall effects were largely similar in these longer-term sites to that of 72 sites that had data four years after nutrient additions began. Because three spatial blocks may be somewhat limited in spatial extent for estimating effects on β diversity, we

tested whether including more blocks to create the γ scale altered our results. We redid the analyses
for the effects of nutrient addition on Δα, Δγ, and Δβ for the entire communities as well as species
groups using 16 sites that had four spatial blocks (Fig. S11~Fig. S13), and again using 11 sites that
had five spatial blocks (Fig. S14 ~ Fig. S16). Again, the overall effects were largely similar to that
we found for the full analysis of the 72 sites with three blocks.

Despite our evidence for little change in β diversity under nutrient addition across the entire
experiment, we recognize limitations of extrapolating these results to the landscape scale (e.g., >1
km × 1 km). This is because many ecological processes that directly and indirectly influence plant
diversity can be very different at the landscape scales[14,42]. Also, the scale in which we inferred
changes in the number of spatially restricted and widespread species, by examining how many local
communities they were lost from, is on a small scale. Linking estimates of species' geographic
range size and their traits with changes in plant diversity across spatial scales[19,42] will deepen our
understanding of the mechanisms of diversity change under nutrient enrichment.

To summarize, we apply a topology that characterizes changes in the number of spatially restricted
and widespread species to a globally distributed long-term experiment to provide a comprehensive
synthesis of the impact of nutrient addition on scale-dependent plant diversity change in grasslands.
Overall, we found proportional plant diversity declines at local and larger spatial scales under
nutrient addition, and little evidence for either biotic homogenization or differentiation within sites.
These overall patterns were largely consistent for diversity metrics less sensitive to rare species than
species richness, across species groups, and over long time periods. This demonstrates that nutrient
enrichment poses a potential threat to all plant species groups, including widespread and native
species that often drive ecosystem functions and services.

Materials and Methods

Experimental setup
The experimental sites used in this study are part of the Nutrient Network (NutNet, Fig. S1 and
Table S1). The experimental design includes a factorial manipulation of nutrients (N, P, and K) plus
two fences to exclude herbivores, see ref[39] for more details. For the analyses here, we used plots
under two treatments: Ambient (Control) and fertilization by nitrogen, phosphate, and potassium
together (i.e., NPK). A micronutrient mix consists of Fe (15%), S (14%), Mg (1.5%), Mn (2.5%),
Cu (1%), Zn (1%), B (0.2%), and Mo (0.05%) was added once only at the start of the experiment
(i.e., year 1) for the nutrient addition plots, but not in subsequent years to avoid toxicity. Nitrogen,
phosphate, potassium were added annually before the beginning of the growing season at most
sites. Nitrogen was added as $10\,\mathrm{g\,m^{-2}\,yr^{-1}}$ time-release urea $[(NH_2)_2CO]$, phosphate was added as
$10\,\mathrm{g\,m^{-2}\,yr^{-1}}$ triple-super phosphate $[Ca(H_2PO_4)_2]$, while potassium was added as $10\,\mathrm{g\,m^{-2}\,yr^{-1}}$
sulfate $[K_2SO_4]$.

Data were retrieved from the NutNet in November 2023. We analyzed data from 72 sites where 1)
nutrients were applied for at least four years; 2) each site had at least three blocks. These sites are
distributed across six continents and include a wide range of grassland types. See Fig. S1 and Table
S1 for details of site codes, grassland types, geolocation, and experimental years used.

Sampling protocol
Scientists at NutNet sites followed standard sampling protocols[39]. Specifically, a 1 m × 1 m subplot
within each plot was permanently marked for annual recording of plant species composition.
Species cover (%) was estimated visually for individual species in the subplots; thus the total cover
of living plants may sometimes exceed 100% for multilayer canopies. At most sites, cover was
recorded once per year at peak biomass before nutrient addition. At some sites with strong

seasonality, cover was recorded twice per year to include a complete list of species. For those sites,
the maximum cover for each species and total biomass were used in the analyses. The taxonomy
was adjusted within sites to ensure consistent naming over time. That is, when taxa could not be
identified to the species level, they were aggregated at the genus level but referred to as "species"
for simplicity. Aboveground biomass was measured within the treatment plot adjacent to the
permanent subplot. All aboveground biomass within two 1×0.1 m strips (in total 0.2 m^2) were
clipped, strips were moved from year to year to avoid resampling the same location. For subshrubs
and shrubs occurring within strips, we collected all leaves and current year's woody growth. All
biomass was dried at 60 °C (to constant mass) before weighing to the nearest 0.01 g.

Quantifying changes in α, γ, and β diversity

We measured α and γ diversity using species richness (i.e., number of species) because it is the
most commonly examined diversity metric[43]. At each site, α diversity was estimated as the number
of species in each permanent subplot (1 m × 1 m), and γ diversity as the total number of species
occurring in three permanent subplots (for each treatment separately). To standardize sampling
effort, for sites with more than three blocks, we selected the first three blocks according to the block
number recorded by site PIs. The typology relies on Whittaker's multiplicative β diversity partition,
it measures changes in the effective number of communities[12]. As such, if all subplots share the
same species, then β diversity would equal to one. In contrast, if each subplot has unique species,
then β diversity would equal to three. We calculated Δα as the richness difference in local
communities (subplots) and Δγ as the difference in the sum of the subplots under nutrient addition
relative to that of control treatment on the log scale. That is, $\Delta\alpha = \log(\alpha_{NPK}/\alpha_{Control})$ and $\Delta\gamma =$
$\log(\gamma_{NPK}/\gamma_{Control})$. We calculated Δβ as Δγ minus $\overline{\Delta\alpha}$, where $\overline{\Delta\alpha}$ is the average of Δα over three
blocks. A decrease in Δβ indicates nutrient addition causes species composition to be more similar
among three subplots than that among control subplots. Because sites are not evenly distributed
around the world, many sites are aggregated in Norther America, we checked spatial autocorrelation
of diversity change under nutrient addition using Moran's I. We found that $\overline{\Delta\alpha}$, Δγ, and Δβ did not
appear to be more similar for sites that are closer to each other (Table S2).

We fitted multilevel (also referred as mixed effects or hierarchical) models for Δα, Δγ, and Δβ (as
the response variable; all on the log scale) separately. We included random intercept for each site,
model was coded as: richness change ~ 1 + (1 |sites) to estimate site-level variation. We used
Bayesian analysis because it yields full posterior distributions of parameters rather than point
estimates and p-values, which provides a deeper understanding of the uncertainty and variability in
the results[44]. Models described above were fitted using the Hamiltonian Monte Carlo (HMC)
sampler in Stan and coded using the R package 'brms' in R[45,46]. Models were fitted with 6 chains
and 3000 iterations (1000 iterations for warm up). Visual inspection of the HMC chains and Rhat
summaries showed model convergence (all Rhats < 1.02; Table S3; Table S5; Table S6). We
visually checked posterior predictive plots to determine how well models can reproduce the data
(Fig. S2).

To examine whether diversity change were sensitive to species relative covers, we redid the above
analyses (i.e., based on species richness) using Shannon diversity and Simpson diversity (both
converted to effective numbers)[47](Fig. S4). Species richness is most sensitive to rare species,
followed by Shannon diversity, and Simpson diversity is more sensitive to the numbers of common
(or relatively abundant) species. We calculated the exponential of Shannon diversity and the inverse
form of Simpson diversity using the R package vegan[48]. These three diversity metrics equal to
diversity with order q = {0, 1, 2}, where increasing q decreases the influence of rare species, and D_q
$= \left(\sum_{i=1}^{s} p_i^q\right)^{1/(1-q)}$, where p is the relative cover of species i, s is the total number of species. These
diversity metrics are also referred to as Hill numbers[47,49].

Site covariates. We investigated whether the effects of nutrient addition on $\overline{\Delta\alpha}$, γ, and β diversity based on species richness were mediated by site characteristics. We included site characteristics that have been shown in previous literature to influence Δα, Δγ, and Δβ in grasslands: site species pool, site productivity, drought intensity, and grazing intensity[24,25,34,40]. We quantified drought intensity as the sum of annual evapotranspiration/precipitation, and averaged it from year 0 to 4 at each site. We quantified the site species pool as the total number of species occurring from year 0 to 4 at each site, and site productivity as the average biomass from year 0 to 4 under the control treatment. We used published methods to quantify an integrated grazing intensity metric from vertebrate herbivores at each site. Specifically, herbivore species (> 2 kg) that consume grassland biomass were documented at each site by site PIs, and each species was assigned an importance value from 1 (present, but low impact and frequency) to 5 (high impact and frequency). An index value was calculated for each site as the sum of herbivore importance values for all herbivores following ref[50,51]. We also investigated relationships between change in diversity and distance among blocks, because species composition may become less similar as the distance between sampled communities increases. The average pairwise distance among the three blocks within sites ranged from 23.04 to 12538.09 m, with a mean of 513.01 m and a median of 118.7 m across 54 sites that have geolocation data for each block. We first calculated three euclidean distances between pairs of blocks, we then used the mean of these pairwise distances as the average distance. We used the average distance instead of area, because blocks are arranged in parallel at some sites. We fitted linear regression models for $\overline{\Delta\alpha}$, Δγ, and Δβ as the response variable separately, and each of the site characteristics was used as a predictor variable.

Species groups. We then investigated the effects of nutrient addition on Δα, Δγ, and Δβ within groups of species with similar characteristics following the method for changes in α, γ, and β diversity in the entire communities. We eliminated sites where no species occurred in control, nutrient addition, or both plots because the value of the log (0) is undefined. We ran the analyses separately for native and non-native species. Native and non-native species were classified by site PIs. Then, we investigated effects of nutrient addition on species richness for different life forms including forb, graminoid, legume, and woody species because previous studies have shown that different life forms may show distinct responses to nutrient addition[6,11,52].

Sensitivity test. We tested whether effects of nutrient addition on species richness across spatial scales depend on experimental duration because a few single-site experiments have shown that the effects of nutrient additions on changes in diversity, especially β diversity, may take several years to emerge[29,31]. To that end, we used a subset of 29 sites that had data 10 years after nutrient additions began, and another subset of 16 sites that had data 14 years after nutrient additions began. Also, because three blocks may be limited in spatial extent, we tested whether combining more blocks to create the γ scale would alter our results. We redid the analyses using data from 16 sites that had four spatial blocks, and again using data from 11 sites that had five spatial blocks.

References:

1. Sardans, J. & Peñuelas, J. Potassium: a neglected nutrient in global change. *Global Ecology and Biogeography* **24**, 261–275 (2015).
2. Galloway, J. N., Bleeker, A. & Erisman, J. W. The Human Creation and Use of Reactive Nitrogen: A Global and Regional Perspective. *Annual Review of Environment and Resources* **46**, 255–288 (2021).
3. Yuan, Z. *et al.* Human Perturbation of the Global Phosphorus Cycle: Changes and Consequences. *Environ. Sci. Technol.* **52**, 2438–2450 (2018).
4. Bobbink, R. *et al.* Global assessment of nitrogen deposition effects on terrestrial plant diversity: A synthesis. *Ecological Applications* **20**, 30–59 (2010).

5. Seabloom, E. W. *et al.* Increasing effects of chronic nutrient enrichment on plant diversity loss and ecosystem productivity over time. *Ecology* **102**, 1–11 (2021).
6. Suding, K. N. *et al.* Functional- and abundance-based mechanisms explain diversity loss due to N fertilization. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 4387–4392 (2005).
7. Yang, Z., Hautier, Y., Borer, E. T., Zhang, C. & Du, G. Abundance- and functional-based mechanisms of plant diversity loss with fertilization in the presence and absence of herbivores. *Oecologia* **179**, 261–270 (2015).
8. Zhang, P. *et al.* Space resource utilization of dominant species integrates abundance- and functional-based processes for better predictions of plant diversity dynamics. *Oikos* **2023**, e09519 (2023).
9. MacDougall, A. S. *et al.* The Neolithic Plant Invasion Hypothesis: the role of preadaptation and disturbance in grassland invasion. *New Phytologist* **220**, 94–103 (2018).
10. Seabloom, E. W. *et al.* Plant species' origin predicts dominance and response to nutrient enrichment and herbivores in global grasslands. *Nat Commun* **6**, 7710 (2015).
11. Tognetti, P. M. *et al.* Negative effects of nitrogen override positive effects of phosphorus on grassland legumes worldwide. *Proceedings of the National Academy of Sciences* **118**, (2021).
12. Blowes, S. A. *et al.* Synthesis reveals approximately balanced biotic differentiation and homogenization. *Science Advances* **10**, eadj9395 (2024).
13. Keil, P. *et al.* Spatial scaling of extinction rates: Theory and data reveal nonlinearity and a major upscaling and downscaling challenge. *Global Ecology and Biogeography* **27**, 2–13 (2018).
14. Chase, J. M. *et al.* Species richness change across spatial scales. *Oikos* **128**, 1079–1091 (2019).
15. Daleo, P. *et al.* Environmental heterogeneity modulates the effect of plant diversity on the spatial variability of grassland biomass. *Nat Commun* **14**, 1809 (2023).
16. Le Provost, G. *et al.* The supply of multiple ecosystem services requires biodiversity across spatial scales. *Nat Ecol Evol* **7**, 236–249 (2022).
17. Daru, B. H. *et al.* Widespread homogenization of plant communities in the Anthropocene. *Nat Commun* **12**, 6983 (2021).
18. Gossner, M. M. *et al.* Land-use intensification causes multitrophic homogenization of grassland communities. *Nature* **540**, 266–269 (2016).
19. Staude, I. R. *et al.* Disentangling non-random structure from random placement when estimating β-diversity 1 through space or time. *Nat Ecol Evol* **4**, 802–808 (2020).
20. McKinney, M. L. & Lockwood, J. L. Biotic homogenization: a few winners replacing many losers in the next mass extinction. *Trends in Ecology & Evolution* **14**, 450–453 (1999).
21. Chen, Q. *et al.* Long-term cross-scale comparison of grazing and mowing on plant diversity and community composition in a salt-marsh system. *Journal of Ecology* **109**, 3737–3747 (2021).
22. Huston, M. A General Hypothesis of Species Diversity. *The American Naturalist* **113**, 81–101 (1979).
23. *Tilman, D. 1988. Plant Strategies and the Dynamics and Structure of Plant Communities. Princeton University Press, Princeton, New Jersey, USA.*
24. Seabloom, E. W. *et al.* Species loss due to nutrient addition increases with spatial scale in global grasslands. *Ecology Letters* **24**, 2100–2112 (2021).
25. Chalcraft, D. R. *et al.* Scale-dependent responses of plant biodiversity to nitrogen enrichment. *Ecology* **89**, 2165–2171 (2008).
26. Lu, W. *et al.* Similar effects of seasonal nitrogen enrichment on plant species–area relationship in a temperate grassland. *Ecological Indicators* **158**, 111430 (2024).
27. Chen, Q. *et al.* Nutrients and herbivores impact grassland stability across spatial scales through different pathways. *Global Change Biology* **28**, 2678–2688 (2022).
28. Zhou, X., Liu, X., Zhang, P., Guo, Z. & Du, G. Increased community compositional dissimilarity alleviates species loss following nutrient enrichment at large spatial scales. *Journal of Plant Ecology* **12**, 376–386 (2019).

29. DeSiervo, M. H., Sullivan, L. L., Kahan, L. M., Seabloom, E. W. & Shoemaker, L. G. Disturbance alters transience but nutrients determine equilibria during grassland succession with multiple global change drivers. *Ecology Letters* **26**, 1132–1144 (2023).

30. Eskelinen, A. & Harrison, S. Erosion of beta diversity under interacting global change impacts in a semi-arid grassland. *Journal of Ecology* **103**, 397–407 (2015).

31. Inouye, R. S. & Tilman, D. Convergence and Divergence of Old-Field Vegetation After 11 yr of Nitrogen Addition. *Ecology* **76**, 1872–1887 (1995).

32. Zhang, Y. *et al.* Nitrogen addition does not reduce the role of spatial asynchrony in stabilising grassland communities. *Ecology Letters* **22**, 563–571 (2019).

33. Foster, B. L. *et al.* Seed availability constrains plant species sorting along a soil fertility gradient. *Journal of Ecology* **99**, 473–481 (2011).

34. Koerner, S. E. *et al.* Nutrient additions cause divergence of tallgrass prairie plant communities resulting in loss of ecosystem stability. *Journal of Ecology* **104**, 1478–1487 (2016).

35. Lan, Z. *et al.* Testing the scaling effects and mechanisms of N-induced biodiversity loss: evidence from a decade-long grassland experiment. *Journal of Ecology* **103**, 750–760 (2015).

36. Xu, Z. *et al.* Antithetical effects of nitrogen and water availability on community similarity of semiarid grasslands: evidence from a nine-year manipulation experiment. *Plant Soil* **397**, 357–369 (2015).

37. Yang, X. *et al.* Resource addition drives taxonomic divergence and phylogenetic convergence of plant communities. *Journal of Ecology* **107**, 2121–2132 (2019).

38. Whittaker, R. H. Vegetation of the Siskiyou Mountains, Oregon and California. **30**, 279–338 (1960).

39. Borer, E. T. *et al.* Finding generality in ecology: A model for globally distributed experiments. *Methods in Ecology and Evolution* **5**, 65–73 (2014).

40. Chase, J. M. & Leibold, M. A. Spatial scale dictates the productivity-biodiversity relationship. *Nature* **416**, 427–430 (2002).

41. Seabloom, E. W. *et al.* Predicting invasion in grassland ecosystems: Is exotic dominance the real embarrassment of richness? *Global Change Biology* **19**, 3677–3687 (2013).

42. Staude, I. R. *et al.* Directional turnover towards larger-ranged plants over time and across habitats. *Ecology Letters* **25**, 466–482 (2022).

43. Chase, J. M. *et al.* Embracing scale-dependence to achieve a deeper understanding of biodiversity and its change across communities. *Ecology Letters* **21**, 1737–1751 (2018).

44. Kruschke, J. K. Bayesian estimation supersedes the t test. *Journal of Experimental Psychology: General* **142**, 573–603 (2013).

45. Bürkner, P.-C. Advanced Bayesian Multilevel Modeling with the R Package brms. *The R Journal* **10**, 395 (2018).

46. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. (2022).

47. Jost, L. Entropy and diversity. *Oikos* **113**, 363–375 (2006).

48. Oksanen, A. J. *et al.* Package ' vegan '. (2019).

49. Hill, M. O. Diversity and Evenness: A Unifying Notation and Its Consequences. *Ecology* **54**, 427–432 (1973).

50. Borer, E. T. *et al.* Nutrients cause grassland biomass to outpace herbivory. *Nature Communications* **11**, 1–8 (2020).

51. Anderson, T. M. *et al.* Herbivory and eutrophication mediate grassland plant nutrient responses across a global climatic gradient. *Ecology* **99**, 822–831 (2018).

52. Van Sundert, K. *et al.* Towards comparable assessment of the soil nutrient status across scales —Review and development of nutrient metrics. *Global Change Biology* **26**, 392–409 (2020).

410 **Acknowledgments:**
411 This work was generated using data from the Nutrient Network (http://www.nutnet.org)
412 experiment, funded at the site-scale by individual researchers. We thank the Minnesota

Supercomputer Institute for hosting project data and the Institute on the Environment for hosting Network meetings. We thank researchers from the NutNet who contributed data to our analysis, table S7 lists these researchers. We thank the Minnesota Supercomputing Institute for hosting project data and the Institute on the Environment for hosting Network meetings. Nitrogen fertilizer was donated to some NutNet sites by Crop Production Services, Loveland, CO. This experiment is funded by individual researchers at the site scale.

Funding: German Center for Integrative diversity Research (iDiv) Halle-Jena-Leipzig (DFG FZT 118-202548816) for the contribution of JMC, SAB, EL, QC, WSH, and other iDiv co-authors. ERC Advanced Grant (MetaChange) to JMC for the contribution of JMC, SAB. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.
National Science Foundation grant NSF-DEB-1042132 (ETB, EWS; for NutNet coordination and data management) National Science Foundation grant NSF-DEB-1234162 (ETB, EWS; for Long-Term Ecological Research at Cedar Creek). National Science Foundation grant NSFDEB-1831944 (ETB, EWS; for Long-Term Ecological Research at Cedar Creek), and the Institute on the Environment (DG-0001-13).

Author contributions:
Conceptualization: QC, JMC, SAB, EL, WSH; **Methodology**: QC, JMC, SAB; **Investigation and data collection**: all authors except QC, JMC, SAB, EL, EHG; **Visualization**: QC; **Writing—original draft**: QC; **Writing—review & editing**: all authors. See table S8 for more details for the contribution of each author.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: The NutNet data are publicly available on the Environmental Data Initiative (EDI) (https://doi.org/10.6073/pasta/583874460a0af70f93d 3eee2f22f9a13). The data used in this study are publicly available on https://figshare.com/s/2ef4e1edf817721d3f6f. The R codes used to produce results in this study will be made publicly available once the manuscript is accepted.

Supplementary Materials

Materials and Methods

Figs. S1 to S16

Tables S1 to S8