Loss of LasR function leads to decreased repression of *Pseudomonas aeruginosa* PhoB activity at physiological phosphate concentrations

Amy Conaway[1], Igor Todorovic[1], Dallas L. Mould[1], and Deborah A. Hogan[1#]

[1]Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth, Hanover, NH USA

Running Title: Increased PhoB activity in LasR- strains

Key Words: PhoB, LasR, PhoR, quorum sensing, phosphate scavenging, phenazines, CbrB, carbon catabolite repression, *Pseudomonas aeruginosa*

#To whom correspondence should be addressed:

Department of Microbiology and Immunology

Geisel School of Medicine at Dartmouth

Rm 208 Vail Building, Hanover, NH 03755

E-mail: dhogan@dartmouth.edu

Tel: (603) 650-1252

Abstract

While *Pseudomonas aeruginosa* LasR plays a role in quorum sensing (QS) across all phylogenetically-distinct lineages, isolates with loss-of-function mutations in *lasR* (LasR– strains) are commonly found in diverse settings including infections where they are associated with worse clinical outcomes. In LasR– strains, another QS transcription factor RhlR can be reactivated in low inorganic phosphate (Pi) concentrations via the two-component system PhoR-PhoB, restoring expression of some of the genes in the QS network. Here, we demonstrate a new link between LasR and PhoB in which the absence of LasR increases PhoB activity at physiological Pi concentrations and raises the Pi concentration necessary for PhoB inhibition. PhoB activity was also less repressed by Pi in mutants lacking different QS regulators (RhlR, and PqsR) and in mutants lacking genes required for the production of QS-regulated phenazines suggesting that decreased phenazine production was one reason for decreased PhoB repression by Pi in LasR–. In addition, the CbrA-CbrB two-component system, which is elevated in LasR– strains, was necessary for reduced PhoB repression by Pi and a Δcrc mutant, which lacks the CbrA-CbrB-controlled translational repressor, activated PhoB at higher Pi concentrations than the in the wild type. The $\Delta lasR$ mutant had a PhoB-dependent growth advantage in a medium with no added Pi and increased virulence-determinant gene expression in a medium with 0.7 mM Pi. Reanalysis of published RNA-seq data found evidence for PhoB activity in *P. aeruginosa* in cystic fibrosis sputum indicating that potential strain differences in PhoB activation may impact disease state.

Introduction

Pseudomonas aeruginosa is a pernicious pathogen that infects burn wounds, nonhealing wounds, eyes, and the airways of people with cystic fibrosis (pwCF) or chronic obstructive pulmonary disease (COPD). One key P. aeruginosa attribute that contributes to pathogenicity is its ability to acquire nutrients, such as phosphorus, that are restricted in the host environment. Phosphate is required for cell membranes, nucleic acids, metabolic intermediates, and is utilized in signal transduction. The most accessible form of phosphorus is free inorganic phosphate (Pi), which is found in human serum at concentrations near 1.3 mM Pi in healthy adults [1, 2]. However, during infection, Pi is highly restricted by the host as part of nutritional immunity through mechanisms including the secretion of phosphate-binding proteins [3-5]. Additionally, the majority of phosphorus in the body is present in organic forms which require degradation prior to uptake and utilization by microbes [6].

In response to low Pi levels, P. aeruginosa induces the production of enzymes such as phosphatases, phospholipases, and DNases to access phosphate from organic sources and high-affinity phosphate transporters via the transcription factor PhoB and its sensor kinase PhoR [7-10]. PhoB activity can also induce the production of specific phenazine small molecules [11-15] which solubilize phosphate from minerals [16]. PhoR activity is regulated by interactions with the high-affinity Pi transporter PstABC via PhoU [9, 17-23] such that PhoR is inhibited by Pi transport into the cell [23, 24]. PhoB positively regulates its own expression [25]. However, published evidence suggests a negative feedback loop also exists in P. aeruginosa [26] as has been described in Escherichia coli [27], presumably to control intracellular Pi. Homologous PhoR-PhoB two-component

systems play significant roles in Pi acquisition across gram-negative bacteria including *E.*

coli [23, 28] *Salmonella enterica* [29]*, and Vibrio cholerae* [30].

In addition to nutrient acquisition, PhoB also regulates virulence in multiple

pathogens [9]. In *E. coli*, type III secretion system (T3SS) components and effectors are

upregulated during Pi-limitation in a PhoB-dependent manner [31]. A different relationship

has been found in *V. cholerae* where PhoB directly binds and inhibits transcription of

genes encoding virulence regulators TcpPH [30]. In *P. aeruginosa,* cross-regulation

between PhoR-PhoB and quorum sensing (QS) has been described [12-15, 32, 33]. *P.*

aeruginosa QS is largely controlled by three transcription factors, LasR, RhlR, and PqsR,

that are active when QS inducers are at sufficient concentrations i.e. high cell densities

or environments with decreased diffusion. While LasR positively regulates both RhlR and

PqsR, many groups have shown that RhlR and PqsR activity can occur in the absence of

LasR [34-37]. It has also been established that activation of PhoB by Pi limitation induces

QS [13-15, 38, 39] and this can occur in a LasR-independent manner [12, 32, 33]. QS regulation

contributes to virulence and biofilm formation in many contexts [40-42]. One group of

virulence factors induced by both PhoB and QS are phenazines which inhibit immune cell

function [43, 44] and kill other microbes [45-49] through oxidative stress. Though PhoB-

binding sites have been identified upstream of some phenazine biosynthesis genes [15],

PhoB induction of phenazine production is largely attributed to increased expression of

rhlR [12, 15, 38]. Furthermore, *phoR, phoB, rhlR,* and *pqsR* are all required for *P. aeruginosa*

phenazine production in co-culture with *Candida albicans* [26], a fungus that frequently co-

infects the airways of pwCF [50]. PhoB regulation of QS suggests low Pi could be a signal

for QS remodeling to circumvent reliance on LasR.

LasR– *P. aeruginosa* are present in the environment [51-53] and also often evolve

during chronic infections like those observed in pwCF [34, 54] where they make up

approximately one-third of clinical isolates [55]. As QS positively correlates with virulence,

it is of significant interest that these LasR– isolates are associated with worse disease

progression in pwCF [55] and worse lesions during acute ocular infections [52]. LasR– *P.*

aeruginosa competitive fitness in laboratory evolution experiments depends on increased

CbrA-CbrB activity [56, 57], which provides growth advantages in complex media [35, 56, 58].

CbrA-CbrB is a two-component system that relieves <u>c</u>arbon <u>c</u>atabolite <u>r</u>epression (CCR)

in *P. aeruginosa* by inducing the production of the small RNA *crcZ* which sequesters Crc

[59-62]. Crc interacts with Hfq to bind to the ribosome binding site of multiple mRNA targets,

inhibiting their translation. Crc-Hfq target mRNAs encode transporters, enzymes, pili, and

more [63]. CbrA is required for full *P. aeruginosa* virulence [64] and both CbrA and CbrB

are rarely mutated in clinical isolates [65].

Investigations of *P. aeruginosa* virulence regulation are often conducted in

laboratory media with millimolar Pi concentrations that repress PhoR activation of PhoB

(e.g. synthetic CF medium (SCFM), 5.1 mM [66]; RPMI medium, 5.6 mM [67, 68]; Luria Broth,

6 mM [69]; or M9 medium, 64 mM [69]). Further, PhoR-PhoB activity is most often studied

in media with less than 0.5 mM Pi [12, 19, 23, 33]. These two extremes either strongly repress

or strongly induce PhoB activity, making it difficult to investigate the contributions of PhoB

to *P. aeruginosa* virulence inside the host at physiological Pi concentrations (1.3 mM Pi

in serum [1, 4]) or when Pi is transiently available. Thus, we aimed to elucidate the

relationship between LasR and PhoB at physiologically relevant Pi concentrations. In

these studies, we show that LasR– isolates and a Δ*lasR* mutant had elevated PhoR-

dependent PhoB activity relative to comparator strains with functional LasR at Pi

concentrations in the 0.7 – 1.1 mM range. In contrast, LasR+ and LasR– strains had

similar PhoB activity in both low (0.2 mM) and high (10 mM) Pi conditions. Our data

demonstrate that a lack of phenazines or reduced Crc activity via CbrA-CbrB led to higher

PhoB activity and decreased repression of PhoR-PhoB by Pi. We show that PhoB is

required for the Δ*lasR* mutant growth advantages in a medium with no added Pi and the

increased expression of multiple virulence factors including phenazine biosynthetic

enzymes and phospholipases. This work establishes a novel connection between QS and

PhoB wherein LasR represses PhoB activity and PhoB is required for increased virulence

gene expression in the Δ*lasR* mutant in part through reactivation of QS. This model for

virulence regulation may aid in understanding why *P. aeruginosa* LasR– strains are

associated with poor clinical outcomes.

Results

PhoB activity is repressed at lower Pi concentrations in *P. aeruginosa* LasR+

strains than in their *P. aerugiosa* LasR– counterparts.

Though PhoB regulates virulence factor production across species [31, 70], there are

only a limited number of *in vitro* studies on PhoB activity at Pi concentrations similar to

those found *in vivo*. Thus, we sought to determine the Pi concentrations necessary to

repress PhoB activity in *P. aeruginosa* strain PA14. PhoB activity was monitored by

assessing the activity of alkaline phosphatase (AP), which is encoded by the PhoB-

regulated gene *phoA* [7, 71, 72], on agar containing the colorimetric AP substrate bromo-4-

chloro-3-indolyl phosphate (BCIP) [8, 73] (Fig. 1A). Using gradient plates with range of Pi

concentrations (0.1 mM - 1 mM) [26] in a MOPS-glucose medium base [74], we found 0.7

mM Pi repressed PhoB activity in the wild-type *P. aeruginosa*. Interestingly, a Δ*lasR*

mutant showed AP activity across the entire gradient though AP activity showed some

reduction at higher Pi concentrations (Fig. 1A). Consistent with PhoB regulation of AP

activity, the Δ*phoB* mutant had no detectable AP activity at any concentration, and the

Δ*pstB* mutant, which has constitutive PhoB activity due to de-repression of PhoR [26, 75],

had strong AP production at all Pi concentrations tested (Fig. 1A). At 0.7 mM Pi, which

repressed PhoB activity in the wild type but not the Δ*lasR* mutant, the Δ*lasR* mutant

complemented with *lasR* at the native locus phenocopied the wild type (Fig. 1B).

Consistent with PhoR activation of PhoB, we found neither the Δ*lasR*Δ*phoB* or

Δ*lasR*Δ*phoR* mutants had AP activity (Fig. 1B). Furthermore, a clinical isolate from a

corneal infection, DH2590 (262K in Hammond, *et al.* [52]), which has a loss-of-function

mutation in *lasR,* also expressed AP at 0.7 mM Pi and this high level was reduced by

complementing a functional copy of *lasR* at the native locus (Fig. 1B). In addition, we

compared AP activity in two clinical isolate pairs (Fig. 1C) from CF sputum samples

wherein one isolate has a LOF mutation in *lasR* and the other does not [76]. In both cases,

the LasR– isolates had more AP activity than their LasR+ counterparts at 0.7 mM Pi.

Both the Δ*lasR* mutant and the wild type showed PhoB activity at lower Pi

concentrations (< 0.7 mM). We found that at low Pi (0.2 mM), the concentration frequently

used to study PhoB activity [12, 13, 19, 23, 77], the wild type and Δ*lasR* strains had similar AP

activity (Fig. 1D). Additionally, neither strain showed AP production at 10 mM Pi.

Additionally, no AP activity was observed in either the wild-type or the Δ*lasR* strain on LB

agar which is reported to have 6 mM Pi [69] (Fig. S1A). The Δ*phoB* mutant did not have

AP activity even at 0.2 mM Pi and the Δ*pstB* mutant had AP production at both 0.2 mM

Pi, 10 mM Pi, and on LB. AP activity was quantified using the soluble colorimetric

substrate p-nitrophenyl phosphate (PNPP). The Δ*lasR* mutant had significantly more AP

activity than the wild type at 0.7 mM Pi (Fig. 1E) and this difference was not attributable

to differences in growth (Fig. S1B). Similarly, qRT-PCR analysis of *phoA*, which encodes

AP, normalized to the reference transcript *ppiD* found significantly lower levels in the wild

type when compared to the Δ*lasR* mutant (Fig. 1F).

To directly assess the levels of transcripts from other genes within the PhoB

regulon, we utilized NanoString multiplex technology as previously published [26]. At 0.7

mM Pi, the Δ*lasR* mutant had significantly higher levels of PhoB-regulated transcripts

including *phoB* itself and genes encoding phosphatases (*phoA* and *lapA*), a phosphate

transporter (*oprO*), a type two secretion system protein required for secretion of LapA

(*hxcT*) [78], a putative TonB transporter (*exbD2*) [79], a phosphodiesterase (*glpQ*), a low-

phosphate ornithine lipid biosynthetic protein (*olsA*) [80], and a Pi chemotaxis protein (*ctpL*)

[81]. None of these transcripts were significantly different when the Δ*phoB* and

Δ*lasR*Δ*phoB* mutants were compared (Fig. 1G and Supplemental File 1).

Other quorum-sensing mutants have active PhoB at higher Pi concentrations than

the wild type which correlates with decreased phenazine production.

LasR positively regulates other QS transcription factors including RhlR and PqsR

and some Δ*lasR* mutant phenotypes are due to their decreased activity. At the same time,

RhlR and PqsR are capable of inducing QS-controlled genes in the absence of LasR in

specific strains and conditions [34, 35, 57, 82]. Thus, we assessed the role of RhlR and PqsR

in the repression of PhoB. As shown in Fig. 2A, AP activity was similarly significantly

elevated in the $\Delta lasR$, $\Delta rhlR$, and $\Delta pqsR$ mutants from the wild type at 0.7 mM Pi. Across

a gradient plate of 0.5 – 1.5 mM Pi, PhoB activity in all QS mutants was inhibited at a

significantly higher Pi concentration than the wild type (Fig. 2B). However, the absence

of LasR ($\Delta lasR$, $\Delta lasR\Delta rhlR$, and $\Delta lasR\Delta pqsR$) led to decreased repression by Pi at

concentrations >1 mM, while the $\Delta rhlR$ and $\Delta pqsR$ were inhibited by 0.9 mM Pi.

As each of the QS transcription factor mutants, $\Delta lasR$, $\Delta rhlR$, and $\Delta pqsR$, produce

fewer phenazines than the wild type in late exponential and early stationary phase

cultures [83, 84], we tested whether the absence of phenazines was sufficient to increase

PhoB activity at moderate Pi concentrations. *P. aeruginosa* produces multiple phenazines

including phenazine-1-carboxylic acid (PCA) as well as the PCA derivatives 5-methyl-

PCA (5MPCA), pyocyanin (PYO), and phenazine-1-carboxamide. PCA is synthesized by

proteins encoded by two highly similar operons, *phzA1-G1* and *phzA2-G2.* The mutant

lacking both *phz* operons ($\Delta phzA1-G1\Delta phzA2-G2$), referred to as Δphz, had significantly

elevated AP activity compared to the wild type at 0.7 mM Pi (Fig. 3A). The $\Delta phz1$ mutant

($\Delta phzA1-G1$) was not different from the wild type, and deletion of either of the adjacent

phenazine-modifying genes required for 5MPCA and PYO biosynthesis, *phzM* or *phzS*,

did not lead to changes in AP production at 0.7 mM (Fig. S2A). In contrast, the $\Delta phz2$

mutant ($\Delta phzA2-G2$) phenocopied Δphz and had significantly more AP production than

the $\Delta phz1$ mutant or the wild type (Fig. 3A). We and others have previously published

data identifying *phzA2-G2* as the predominant contributor of PCA [26, 84-86].

We found no difference in AP activity at 0.7 mM Pi between the $\Delta lasR$ and Δphz

mutants (Fig. 3B). However, a $\Delta lasR\Delta phz$ mutant had significantly elevated AP activity

compared to both single mutants (Fig. 3B). Both the Δ*lasR* and Δ*lasRΔphz* mutants had

detectable AP activity at a significantly higher Pi concentration than Δ*phz* (Fig. 3C)

suggesting that PhoB activity was derepressed by an additional phenazine-independent

mechanism in the Δ*lasR* mutant. Consistent with this model, the Δ*phz* mutant was

repressed by a similar Pi concentration as the Δ*rhlR* and Δ*pqsR* mutants, but not their

LasR– counterparts (Fig. 2B).

CbrA-CbrB-Crc impacts PhoB activity in the Δ*lasR* mutant.

Several studies suggest that *P. aeruginosa* strains lacking LasR activity have

elevated CbrA-CbrB activity in many conditions [56, 58]. CbrA-CbrB activity leads to relief

of Crc-mediated translation repression of its targets including some that encode catabolic

enzymes and transporters [62, 63, 87] (Fig. 4A). In both BCIP and PNPP assays, we found

that deletion of *cbrB* eliminated AP activity at 0.7 mM Pi in a Δ*lasR* mutant and that the

Δ*lasR* phenotype was restored by complementation with the restoration of *cbrB* at the

native locus (Fig 4B). PhoB activity in both the Δ*cbrB* and Δ*lasRΔcbrB* mutants was

similarly repressed below 0.5 mM Pi (Fig. S2B). Additionally, AP activity at 0.7 mM Pi on

BCIP is restored to a Δ*lasRΔcbrB* mutant by deletion of *crc* (Fig. 4B) suggesting that the

increase in Pi repression of PhoB in the Δ*lasRΔcbrB* is due to elevated Crc activity (Fig.

4A for pathway). A Δ*crc* single mutant also had significantly elevated AP activity

compared to the wild type and *crc* complemented strain (Fig. 4C). There was no difference

in AP activity at 0.7 mM Pi between the Δ*lasR* and Δ*lasRΔcrc* mutants, though both were

significantly elevated compared to the Δ*crc* mutant (Fig. 4D). Additionally, the Δ*lasR* and

Δ*lasRΔcrc* mutants both showed significantly decreased PhoB sensitivity to Pi than the

Δ*crc* mutant (Fig. 4E).

206 PhoB activity in the *P. aeruginosa* Δ*lasR* strain provides fitness advantages when

207 Pi is limited.

To determine if the increased PhoB activity in *P. aeruginosa* Δ*lasR* cells increased

fitness upon depletion of Pi, we grew the wild-type, the Δ*lasR* mutant, and their respective

phoB mutants overnight in LB, then sub-cultured into MOPS-glucose medium with no

added Pi (Fig. 5A). The wild type and the Δ*lasR* mutant showed similar initial growth rates.

However, in the post-exponential phase, the Δ*lasR* mutant reached significantly higher

densities than the wild type. Both the Δ*phoB* and Δ*lasR*Δ*phoB* mutants showed minimal

growth in the no added Pi media and were not significantly different from each other.

215 PhoB activity mediates the expression of virulence determinants in LasR– *P.*

216 *aeruginosa,* in part through QS*.*

Multiple studies have shown that there are alternative mechanisms for the

activation of RhlR in LasR– strains [12, 32-36] and that phosphate limitation promotes RhlR

activity [12-15, 32, 33]. We have previously published that both PhoB and RhlR are required

for antagonistic 5MPCA production by *P. aeruginosa* in co-culture with *Candida albicans*

[26]. Here we show that the Δ*lasR* mutant produces more of the antifungal phenazine

5MPCA in co-culture with *Candida albicans* and that 5MPCA production by both LasR+

and LasR– *P. aeruginosa* required PhoB and CbrB (Fig. 5B). Using NanoString multiplex

technology and a previously published code set [26], we compared virulence determinant

transcripts from the wild type and Δ*phoB, ΔlasR, ΔlasRΔphoB* mutants grown as colony

biofilms on MOPS agar with 0.7 mM Pi. Transcripts encoding multiple virulence

associated proteins, including phenazine biosynthetic enzymes (*phzM* and *phzS*),

phospholipases (*plcH* and *plcN*), and an exopolysaccharide (EPS) matrix regulator

(*phdA*) [88], were significantly higher in the Δ*lasR* mutant compared to the wild type (Fig.

5C and Supplemental File 1). These differences were not detected when the Δ*phoB* and

Δ*lasR*Δ*phoB* mutants were compared, though all transcripts were significantly lower in

the Δ*lasR*Δ*phoB* mutant compared to the Δ*lasR* strain. As expected, expression of *plcN*

[11, 89] and *phdA* [79, 90], which are impacted by PhoB activity, were significantly lower in the

Δ*phoB* mutant compared to the wild type. However, it is interesting that *phzM, phzS,* and

plcH expression were only PhoB-dependent in the Δ*lasR* mutant under these conditions.

As the production of phenazines by LasR– *P. aeruginosa* requires RhlR [12, 32, 33, 35]

and Pi limitation promotes RhlR activity [12-14, 32, 33], we sought to determine if PhoB

contributes to QS at 0.7 mM Pi. We found significantly higher abundance of transcripts

encoding QS proteins (*lasI, rhlR, pqsR,* and *pqsH)* and those regulated by QS

transcription factors (*lasB* and *rhlA*) in the Δ*lasR* mutant compared to the Δ*lasR*Δ*phoB*

mutant at 0.7 mM Pi (Fig. 5D and Supplemental File 1). The transcripts *lasB, rhlR, pqsR,*

and *pqsH* were significantly lower in the Δ*lasR* mutant compared to the wild type, though

there were no differences in *lasI* or *rhlA* abundance. We found no significant differences

in QS transcript abundance between the Δ*phoB* mutant and the wild type at 0.7 mM Pi.

Cystic fibrosis sputum isolates have active PhoB.

CF sputum is frequently colonized by *P. aeruginosa* but is difficult to assess free

Pi levels in sputum as there are many organic forms also present. Thus, we sought to

determine if conditions in sputum, including Pi levels, suppressed PhoB activity by

analyzing high-quality RNA-seq data of *P. aeruginosa* from sputum published by Lewin,

et al. [91]. In this study, RNA-seq analysis of RNA from *in vivo* CF sputum samples was

compared to that of the laboratory strain PAO1 grown in synthetic cystic fibrosis medium

(SCFM2) and other *in vitro* conditions. SCFM2 contains 5.1 mM Pi [92]. We used the *in*

vivo and SCFM2 samples from this published dataset to compare expression of 85 genes

in the PhoB regulon, as defined by a gene ADAGE signature [93] (Supplemental File 2).

Overall, we found higher transcripts of PhoB-regulated genes including those encoding

phosphatases, phospholipases, and transporters *in vivo* than *in vitro* (Fig. 6). In fact, 42%

of the Adage-defined PhoB regulon was significantly higher *in vivo* (Supplemental File 2).

Across the sputum samples, there was heterogeneity suggesting that sputum conditions

and/or infecting strains led to varied levels of PhoB activity.

Discussion

In this manuscript, we showed that LasR– *P. aeruginosa* laboratory strains and

clinical isolates had elevated PhoB-dependent AP activity and increased expression of

the PhoB regulon at 0.7 mM Pi. We found that the range of PhoB-permissive Pi

concentrations was broader for LasR– *P. aeruginosa*, though PhoB was still repressed at

high Pi concentrations. Importantly, these intermediate Pi concentrations where LasR–

P. aeruginosa showed more PhoB activity are similar to those found in the serum of

healthy adults [1, 2]. We also found that elevated PhoB activity was common across QS

mutants and that this was possibly due to reduced phenazine production. However,

LasR– *P. aeruginosa* had active PhoB at even higher Pi concentrations than the other

QS mutants, suggesting differences in PhoB regulation.

LasR– *P. aeruginosa* are more fit than their LasR+ counterparts in many contexts.

LasR– *P. aeruginosa* are known to have elevated CbrA-CbrB activity, conferring growth

advantages in complex media through decreased Crc activity [56, 58]. Other studies have

shown that LasR– *P. aeruginosa* are more resistant to cell lysis at high pH [94] and more

fit in microoxic conditions and normoxically grown colony biofilms [95]. Future studies are

necessary to determine if these phenotypes are interrelated. We showed that CbrB was

required for PhoB activation in LasR– *P. aeruginosa* and a *crc* mutation was sufficient to

induce PhoB activity, even in a LasR+ strain. Importantly, the Δ*crc* mutant had less AP

activity and was inhibited by lower Pi concentrations than the Δ*lasR* mutant. These data

suggest Crc- *P. aeruginosa* do not phenocopy LasR– strains.

The elevated PhoB activity in the Δ*lasR* mutant contributed to fitness upon a shift

from LB to a medium without added Pi. As phosphate accessibility can be transient inside

the host, increased PhoB activity may be beneficial in building up stores of polyphosphate

for utilization when cells experience low Pi stress. In addition to regulating phosphate

acquisition, PhoB activity has also been shown to promote RhlR expression and

subsequent PYO production, swarming motility, and cytotoxicity [12, 14, 32, 33, 89]. Here we

showed that the Δ*lasR* mutant had PhoB-dependent increased expression of virulence

genes including phenazine biosynthetic enzymes and phospholipases like PlcH. PlcH is

known to degrade airway surfactant and we have previously shown that its expression

leads to a decline in lung function [96]. We also showed PhoB and CbrB are both required

for increased production of the antifungal phenazine 5MPCA by the Δ*lasR* mutant in co-

culture with *C. albicans*.

The roles for phenazines in PhoB regulation are complex. As all QS mutants

produce fewer phenazines in late exponential phase and early stationary phase [83, 84], we

proposed phenazines may repress PhoB activity. We demonstrated that phenazine-

deficient *P. aeruginosa* had similarly elevated PhoB activity to the single QS mutants.

Unlike the Δ*rhlR* and Δ*pqsR* mutants, LasR– *P. aeruginosa* are capable of producing

phenazines in certain contexts [12, 34-36]. We observed that the Δ*lasR* mutant appeared

greener on BCIP agar at 0.7 mM Pi which we speculate is due to the production of the

blue-green phenazine PYO as this was not observed in the phenazine-deficient

Δ*lasR*Δ*phz* mutant (Fig. 3B). Thus, the higher AP activity in the Δ*lasR*Δ*phz* compared to

the Δ*lasR* may be due to the absence of phenazines that reduce PhoB activation at

moderate Pi concentrations. It is also of note that PhoB activity promotes phenazine

production, potentially through direct binding upstream of the *phz* operons [15], but more

likely through RhlR activity [12, 32, 33]. This increased phenazine production could then

inhibit further PhoB activity, providing negative feedback. However, our data suggests

PCA, not PYO or 5MPCA, inhibits PhoB as loss of the biosynthetic enzymes PhzM or

PhzS did not induce PhoB activity. It cannot be assumed that cells producing visible PYO

also have high PCA and reduced PhoB activity. An example of this is the Δ*crc* mutant

which we showed had elevated AP activity and is known to produce more PYO through

de-repression of PhzM [97]. In some instances, increased PCA conversion to PYO may

even result in PCA depletion and thus aid in the activation of PhoB. Further work is

needed to fully understand the interplay between PhoB activity and phenazines.

It is well established that in permissive Pi conditions, PhoB activity can directly

promote the expression of QS-regulated genes through RhlR [14, 15, 32, 89], allowing cells to

circumvent reliance on LasR as a QS regulator [12, 33]. Our data support this existing model

by showing QS gene expression is PhoB-dependent in a Δ*lasR* mutant but not the wild

type at 0.7 mM Pi. Importantly, we have shown increased PhoB activity in LasR– *P.*

aeruginosa compared to LasR+ strains and this has not been identified in previous

publications. The relatively low (0.4 – 0.5 mM) and high (4 – 4.5 mM) Pi concentrations

used in the past likely hindered the observation of increased PhoB activity in LasR– *P.*

aeruginosa and demonstrates the utility of gradient agar plates. As we found PhoB is

active in the ∆*lasR* mutant at higher Pi concentrations than the wild type, the previously

reported PhoB-driven reorganization of QS may occur in LasR– *P. aeruginosa* under

conditions that otherwise repress PhoB activity in LasR+ strains.

It is challenging to determine extracellular Pi concentrations in the infection

environment due to the abundance of organic phosphate in membrane lipids, DNA, and

metabolites. To address this, we reanalyzed an existing RNAseq data set of CF sputum

samples and PAO1 grown in SCFM2 [91]. We found the majority of the PhoB regulon was

elevated in the sputum samples, suggesting they may contain less Pi than what is

provided in SCFM2 (5.1 mM) [92]. Our data indicate that PhoB activity in LasR– *P.*

aeruginosa differs from the wild type at intermediate Pi concentrations (0.7 – 1.1 mM).

increased levels of the PhoB regulon in sputum was variable across samples from

different donors. This variability could be due to differences in the genotype of infecting

P. aeruginosa, differences in inflammation, or the presence of other microbes that may

compete for Pi.

Further work is needed to elucidate the mechanisms by which LasR activity

represses PhoB activity at Pi concentrations in the intermediate Pi range. There is

evidence that PhoB can be spontaneously phosphorylated by acetyl phosphate *in vitro*

[98] but the concentrations required for this reaction suggest it is unlikely to occur inside

cells [99]. Therefore, we expect changes in PhoB activity are mediated by its sensor kinase

PhoR. PhoR activity is suppressed by interactions with the Pi transporter PstABC and

PhoU [17, 20, 22]. Thus, changes in expression of these proteins could alter PhoR-PhoB

activity. We observed no differences in transcript abundance of *phoU* or *pstA* between

the wild type and Δ*lasR* mutant (Supplemental File 1) and so believe this model is unlikely.

However, other changes in Pi transport into the cell could still contribute to differences in

PhoB activity. While the mechanisms of PhoR-PhoU-PstABC interactions are not well

understood, PhoR has been shown to interact with PhoU through its Per-Arnt-Sim (PAS)

domain [22]. PAS domains in other organisms bind a broad range of ligands [100] and can

modify kinase activity [101, 102]. Thus, we propose that changes to the intracellular

environment in phenazine-deficient cells or cells with low Crc activity may stimulate PhoR

activity through its PAS domain. Future work into the mechanisms of the PhoR PAS

domain would also be beneficial for understanding similar kinases in *P. aeruginosa* and

other organisms as PAS domains are abundant but remain poorly understood [103].

Understanding LasR– *P. aeruginosa* virulence in physiologically relevant conditions is

critical as LasR– isolates are frequently found in both acute and chronic infections and

these findings may be relevant to *P. aeruginosa* when in settings that do not induce QS

regulation. We propose that increased PhoB-mediated QS and virulence gene expression

in LasR– *P. aeruginosa* may contribute to their association with worse outcomes [52, 55].

Our data support a novel model where LasR represses PhoB activity and thus

LasR– *P. aeruginosa* have elevated PhoB activity at physiological Pi concentrations.

Others have shown that PhoB activity can induce RhlR activity in the absence of LasR,

leading to increased production of virulence factors like PYO [12, 33]. Here we have shown

QS and virulence-related gene expression in the Δ*lasR* mutant is highly PhoB dependent

at a physiologically relevant Pi concentration.

Materials and Methods

Strains and growth conditions

Bacterial strains and plasmids used in this study are listed in **Table S1**. Bacteria were

streaked from frozen stocks onto LB (lysogeny broth) with 1.5% agar [104]. Planktonic

cultures were grown in 5 mL LB medium in 18 mm Borosilicate glass tubes on a roller

drum at 37 °C in the indicated medium.

Construction of in-frame deletion, complementation, and expression plasmids

Construction of in-frame deletion and complementation constructs was performed using

yeast cloning techniques in *Saccharomyces cerevisiae* as previously described [104] or

Gibson assembly [105, 106]. In-frame deletion and chromosomal complementation

constructs were made using the allelic replacement vector pMQ30 [107]. Plasmids were

purified from yeast using Zymoprep Yeast Plasmid Miniprep II according to the

manufacturer's protocol and transformed into *E. coli* strain S17/λpir by electroporation.

Plasmids were introduced into *P. aeruginosa* by conjugation and recombinants were

obtained using sucrose counter-selection. Genotypes were screened by PCR and

plasmid constructs were confirmed by sequencing.

Alkaline phosphatase activity assessment

Agar plates were made as previously described [26] with MOPS minimal medium [74] with

0.7 mM K_2HPO_4/KH_2PO_4, 60 µg/mL BCIP (Sigma Aldrich # 11383221001), 20 mM

glucose and 15 g/L agar (referred to here as MOPS agar). Overnight cultures of *P.*

aeruginosa were grown in LB, normalized to an OD_{600} of 1, then 5 µl were spotted on

MOPS agar and incubated at 37 °C for 24 h. Plates with a gradient of phosphate were

made based on the methodology described for pH gradient plates [108]. First 35 mL of

molten MOPS agar at one end of the gradient was pipetted into a 10 cm square petri dish

(Corning, BP124-05) that rested in a custom 3D-printed prop that held the plate slanted

at a 30° angle. Once the bottom layer had solidified, the plate was laid flat and 35 mL of

molten medium agar containing the second desired concentration on the gradient was

poured atop. BCIP was added as described above.

398   Calculating alkaline phosphatase enzymatic activity

A colorimetric assay using the substrate p-Nitrophenyl phosphate (PNPP) (New England

Biolabs) was used to quantify alkaline phosphatase activity in colonies grown on MOPS

agar plates with no added BCIP using the inoculation regime described above. After 12-

24 h incubation, the colonies were scraped from the agar with a pipette tip and

resuspended in 1 mL 10 mM Tris-HCL buffer at pH 8. Fifty microliters of the cell

suspension was mixed with 50 µl of room temperature 1-step PNPP solution or 50 µl of

Tris-HCL buffer at pH 8 and incubated for 1 h. A_{405} and A_{600} were recorded at t = 0 min

and t = 60 min. AP activity was calculated using the equation 1000 x (ΔA_{405}/(time (min) x

A_{600})) where ΔA_{405} = (A_{405}PNPP − A_{405}NoPNPP)$_{t60}$ − (A_{405}PNPP − A_{405}NoPNPP)$_{t0}$.

NanoString analysis

Total RNA was collected from *P. aeruginosa* colony biofilms grown on MOPS agar with

0.7 mM K_2HPO_4/KH_2PO_4. Bacteria were grown overnight in LB then sub-cultured in LB

until OD_{600} = 0.5 before 5 µl were spotted on MOPS agar and incubated at 37 °C for 12

h. Cells were then scraped from the agar and resuspended in 200 µl Tris-EDTA buffer.

RNA was extracted using the Qiagen RNeasy kit (Qiagen 74004) using the protocol

provided by the manufacturer. Samples were not treated with DNAse. NanoString

analysis of 80 ng of isolated RNA using codeset PaV5 [26] was performed as previously

reported [109]. Counts were normalized to the geometric mean of spiked-in technical

controls and five housekeeping genes (*ppiD*, *rpoD*, *soj*, *dnaN*, *pepP*, *dapF*). Normalized

counts were used for heatmap construction.

Growth curves

P. aeruginosa was grown in LB medium for 16 h and then sub-cultured to a starting OD_{600}

= 0.05 in 5 mL of MOPS minimal medium + 16.7 mM glucose in glass culture tubes. OD_{600}

was recorded using a Spectronic 20D+.

Transcriptomic analysis

Publicly available transcriptomics data was accessed from Lewin *et al.* [91]. Transcripts

were Log_2 transformed, z-scored, and represented in a heatmap generated in R [110]. Full

details of this analysis and the code used to produce the heatmap are accessible through

the Github repository associated with this publication (https://github.com/hoganlab-

dartmouth/PhoB-regulon-in-sputum).

Statistical analysis

Statistical analyses were performed using GraphPad Prism version 10.1.1 for macOS.

Acknowledgments

We thank Jay Goddard for their assistance in the reanalysis of existing RNA-seq data. The research reported in this publication was supported by grants from the Cystic Fibrosis Foundation, GREEN19G0) and STANTO19R0, and the National Institutes of Health (NIH), NHLBI T32HL134598 (A.C.) and NIDDK P30-DK117469 (Dartmouth Cystic Fibrosis Research Center or DartCF). Additional support came from bioMT through (NIGMS P20GM113132) and the Dartmouth Molecular Biology Shared Resource (NCI 5P30CA023108).

Citations

1. Bansal VK. 1990. Clinical Methods: The history, physical, and laboratory examinations. Serum inorganic phosphorous. *In* HK W, HD, Hall, JW, Hurst (ed), Clinical Methods: The History, Physical, and Laboratory Examinations, 3 ed. Butterworths, Boston.

2. Oh TK, Jo J, Oh AY. 2018. Perioperative serum calcium and phosphorus levels are associated with hospital costs and length of stay after major abdominal surgery. J Clin Med 7.

3. Salem RR, Tray K. 2005. Hepatic resection-related hypophosphatemia is of renal origin as manifested by isolated hyperphosphaturia. Ann Surg 241:343-8.

4. Long J, Zaborina O, Holbrook C, Zaborin A, Alverdy J. 2008. Depletion of intestinal phosphate after operative injury activates the virulence of *P. aeruginosa* causing lethal gut-derived sepsis. Surgery 144:189-97.

5. Riedler GF, Scheitlin WA. 1969. Hypophosphataemia in septicaemia: higher incidence in gram-negative than in gram-positive infections. Br Med J 1:753-6.

6. Berner YN, Shike M. 1988. Consequences of phosphate imbalance. Annual review of nutrition 8:121-148.

7. Filloux A, Bally M, Soscia C, Murgier M, Lazdunski A. 1988. Phosphate regulation in *Pseudomonas aeruginosa*: Cloning of the alkaline phosphatase gene and identification of *phoB*- and *phoR*-like genes. Molecular and General Genetics 212:510-513.

8. Monds RD, Newell PD, Schwartzman JA, O'Toole GA. 2006. Conservation of the Pho regulon in *Pseudomonas fluorescens* Pf0-1. Applied and Environmental Microbiology 72:1910-1924.

9. Lamarche MG, Wanner BL, Crépin S, Harel J. 2008. The phosphate regulon and bacterial virulence: a regulatory network connecting phosphate homeostasis and pathogenesis. FEMS Microbiology Reviews 32:461-473.

10. Gallarato LA, Sanchez DG, Olvera L, Primo ED, Garrido MN, Beassoni PR, Morett E, Lisa AT. 2014. Exopolyphosphatase of *Pseudomonas aeruginosa* is essential for the production of virulence factors, and its expression is controlled by NtrC and PhoB acting at two interspaced promoters. Microbiology 160:406-417.

11. Matilla MA, Udaondo Z, Maaß S, Becher D, Krell T. 2022. Virulence induction in *Pseudomonas aeruginosa* under inorganic phosphate limitation: a proteomics oerspective. Microbiology Spectrum 10:e0259022.

12. Soto-Aceves MP, Cocotl-Yañez M, Servín-González L, Soberón-Chávez G. 2021. The Rhl quorum-sensing system is at the top of the regulatory hierarchy under phosphate-limiting conditions in *Pseudomonas aeruginosa* PAO1. Journal of Bacteriology 203:e00475-20.

13. Bains M, Fernandez L, Hancock RE. 2012. Phosphate starvation promotes swarming motility and cytotoxicity of *Pseudomonas aeruginosa*. Appl Environ Microbiol 78:6762-8.

14. Blus-Kadosh I, Zilka A, Yerushalmi G, Banin E. 2013. The effect of *pstS* and *phoB* on quorum sensing and swarming motility in *Pseudomonas aeruginosa*. PloS one 8:e74444.

488   15.   Jensen V, Löns D, Zaoui C, Bredenbruch F, Meissner A, Dieterich G, Münch R,
489   Häussler S. 2006. RhlR expression in *Pseudomonas aeruginosa* is modulated by
490   the *Pseudomonas* quinolone signal via PhoB-dependent and-independent
491   pathways. Journal of Bacteriology 188:8601-8606.
492   16.   McRose DL, Newman DK. 2021. Redox-active antibiotics enhance phosphorus
493   bioavailability. Science 371:1033-1037.
494   17.   Muda M, Rao N, Torriani A. 1992. Role of PhoU in phosphate transport and
495   alkaline phosphatase regulation. Journal of Bacteriology 174:8057-8064.
496   18.   Rice CD, Pollard JE, Lewis ZT, McCleary WR. 2009. Employment of a promoter-
497   swapping technique shows that PhoU modulates the activity of the PstSCAB2 ABC
498   transporter in *Escherichia coli*. Applied and environmental microbiology 75:573-
499   582.
500   19.   Lee SJ, Park YS, Kim SJ, Lee BJ, Suh SW. 2014. Crystal structure of PhoU from
501   *Pseudomonas aeruginosa*, a negative regulator of the Pho regulon. Journal of
502   Structural Biology 188:22-9.
503   20.   de Almeida LG, Ortiz JH, Schneider RP, Spira B. 2015. *phoU* inactivation in
504   *Pseudomonas aeruginosa* enhances accumulation of ppGpp and polyphosphate.
505   Applied and environmental microbiology 81:3006-3015.
506   21.   Munévar NFV, de Almeida LG, Spira B. 2017. Differential regulation of
507   polyphosphate genes in *Pseudomonas aeruginosa*. Molecular Genetics and
508   Genomics 292:105-116.
509   22.   Gardner SG, Johns KD, Tanner R, McCleary WR. 2014. The PhoU protein from
510   *Escherichia coli* interacts with PhoR, PstB, and metals to form a phosphate-
511   signaling complex at the membrane. Journal of Bacteriology 196:1741-1752.
512   23.   Wanner BL. 1996. Phosphorus assimilation and control of the phosphate regulon.
513   *Escherichia coli* and *Salmonella*: cellular and molecular biology 1:1357-1381.
514   24.   Nikata T, Sakai Y, Shibata K, Kato J, Kuroda A, Ohtake H. 1996. Molecular
515   analysis of the phosphate-specific transport (*pst*) operon of *Pseudomonas
516   aeruginosa*. Molecular and General Genetics MGG 250:692-698.
517   25.   Anba J, Bidaud M, Vasil ML, Lazdunski A. 1990. Nucleotide sequence of the
518   *Pseudomonas aeruginosa phoB* gene, the regulatory gene for the phosphate
519   regulon. J Bacteriol 172:4685-9.
520   26.   Doing G, Koeppen K, Occipinti P, Harty CE, Hogan DA. 2020. Conditional
521   antagonism in co-cultures of *Pseudomonas aeruginosa* and *Candida albicans*: an
522   intersection of ethanol and phosphate signaling distilled from dual-seq
523   transcriptomics. PLoS genetics 16:e1008783.
524   27.   Gao R, Stock AM. 2018. Overcoming the cost of positive autoregulation by
525   accelerating the response with a coupled negative feedback. Cell Reports
526   24:3061-3071.e6.
527   28.   Makino K, Shinagawa H, Amemura M, Kawamoto T, Yamada M, Nakata A. 1989.
528   Signal transduction in the phosphate regulon of *Escherichia coli* involves
529   phosphotransfer between PhoR and PhoB proteins. Journal of Molecular Biology
530   210:551-559.
531   29.   Bruna RE, Kendra CG, Pontes MH. 2023. Phosphorus starvation response and
532   PhoB-independent utilization of organic phosphate sources by *Salmonella
533   enterica*. Microbiology Spectrum 11:e02260-23.

30.   Pratt JT, Ismail AM, Camilli A. 2010. PhoB regulates both environmental and
virulence gene expression in *Vibrio cholerae*. Molecular microbiology 77:1595-
1605.
31.   Chekabab SM, Jubelin G, Dozois CM, Harel J. 2014. PhoB activates *Escherichia*
coli O157:H7 virulence factors in response to inorganic phosphate limitation. PLoS
One 9:e94285.
32.   Mellbye B, Schuster M. 2014. Physiological framework for the regulation of quorum
sensing-dependent public goods in *Pseudomonas aeruginosa*. Journal of
Bacteriology 196:1155-1164.
33.   Welsh MA, Blackwell HE. 2016. Chemical genetics reveals environment-specific
roles for quorum sensing circuits in *Pseudomonas aeruginosa*. Cell Chem Biol
23:361-9.
34.   Feltner JB, Wolter DJ, Pope CE, Groleau MC, Smalley NE, Greenberg EP, Mayer-
Hamblett N, Burns J, Déziel E, Hoffman LR, Dandekar AA. 2016. LasR variant
cystic fibrosis isolates reveal an adaptable quorum-sensing hierarchy in
Pseudomonas aeruginosa. mBio 7:e01513-16.
35.   Mould DL, Botelho NJ, Hogan DA, Goldberg JB. 2020. Intraspecies signaling
between common variants of *Pseudomonas aeruginosa* increases production of
quorum-sensing-controlled virulence factors. mBio 11:e01865-20.
36.   Kostylev M, Kim DY, Smalley NE, Salukhe I, Greenberg EP, Dandekar AA. 2019.
Evolution of the *Pseudomonas aeruginosa* quorum-sensing hierarchy.
Proceedings of the National Academy of Sciences 116:7027-7032.
37.   Cugini C, Morales DK, Hogan DA. 2010. *Candida* albicans-produced farnesol
stimulates *Pseudomonas* quinolone signal production in LasR-defective
Pseudomonas aeruginosa strains. Microbiology 156:3096.
38.   Meng X, Ahator SD, Zhang L-H. 2020. Molecular mechanisms of phosphate stress
activation of *Pseudomonas aeruginosa* quorum sensing systems. MSphere
5:e00119-20.
39.   Zaborin A, Gerdes S, Holbrook C, Liu DC, Zaborina OY, Alverdy JC. 2012.
Pseudomonas aeruginosa overrides the virulence inducing effect of opioids when
it senses an abundance of phosphate. PLoS One 7:e34883.
40.   Antunes LCM, Ferreira RBR, Buckner MMC, Finlay BB. 2010. Quorum sensing in
bacterial virulence. Microbiology 156:2271-2282.
41.   Chen G, Lim ED, Winkelman BT, Winkelman JT, Mukherjee S. 2022.
Combinatorial control of biofilm development by quorum-sensing and nutrient-
sensing regulators in *Pseudomonas aeruginosa*. bioRxiv
doi:10.1101/2022.09.27.509822:2022.09.27.509822.
42.   Hentzer M, Wu H, Andersen JB, Riedel K, Rasmussen TB, Bagge N, Kumar N,
Schembri MA, Song Z, Kristoffersen P, Manefield M, Costerton JW, Molin S, Eberl
L, Steinberg P, Kjelleberg S, Høiby N, Givskov M. 2003. Attenuation of
Pseudomonas aeruginosa virulence by quorum sensing inhibitors. The EMBO
Journal 22:3803-3815.
43.   Nutman J, Berger M, Chase PA, Dearborn DG, Miller K, Waller R, Sorensen R.
1987. Studies on the mechanism of T cell inhibition by the *Pseudomonas*
aeruginosa phenazine pigment pyocyanine. Journal of immunology (Baltimore,
Md: 1950) 138:3481-3487.

44. Rada B, Leto TL. 2013. Pyocyanin effects on respiratory epithelium: relevance in *Pseudomonas aeruginosa* airway infections. Trends in microbiology 21:73-81.

45. Gibson J, Sood A, Hogan DA. 2009. *Pseudomonas aeruginosa-Candida albicans* interactions: localization and fungal toxicity of a phenazine derivative. Applied and environmental microbiology 75:504-513.

46. Morales DK, Jacobs NJ, Rajamani S, Krishnamurthy M, Cubillos-Ruiz JR, Hogan DA. 2010. Antifungal mechanisms by which a novel *Pseudomonas aeruginosa* phenazine toxin kills *Candida albicans* in biofilms. Mol Microbiol 78:1379-92.

47. Morales DK, Grahl N, Okegbe C, Dietrich LE, Jacobs NJ, Hogan DA. 2013. Control of *Candida albicans* metabolism and biofilm formation by *Pseudomonas aeruginosa* phenazines. MBio 4:e00526-12.

48. Zhang L, Tian X, Kuang S, Liu G, Zhang C, Sun C. 2017. Antagonistic activity and mode of action of phenazine-1-carboxylic acid, produced by marine bacterium *Pseudomonas aeruginosa* PA31x, against *Vibrio anguillarum in vitro* and in a zebrafish *in vivo* model. Frontiers in microbiology 8:289.

49. Truong-Bolduc QC, Yonker LM, Wang Y, Lawton BG, Hooper DC. 2024. NorA efflux pump mediates S*taphylococcus aureus* response to *Pseudomonas aeruginosa* pyocyanin toxicity. Antimicrobial Agents and Chemotherapy 68:e01001-23.

50. Kahl LJ, Stremmel N, Esparza-Mora MA, Wheatley RM, MacLean RC, Ralser M. 2023. Interkingdom interactions between *Pseudomonas aeruginosa* and *Candida albicans* affect clinical outcomes and antimicrobial responses. Current Opinion in Microbiology 75:102368.

51. O'Connor K, Zhao CY, Mei M, Diggle SP. 2022. Frequency of quorum-sensing mutations in *Pseudomonas aeruginosa* strains isolated from different environments. Microbiology (Reading) 168.

52. Hammond JH, Hebert WP, Naimie A, Ray K, Van Gelder RD, DiGiandomenico A, Lalitha P, Srinivasan M, Acharya NR, Lietman T, Hogan DA, Zegans ME. 2016. Environmentally endemic *Pseudomonas aeruginosa* strains with mutations in *lasR* are associated with increased disease severity in corneal ulcers. mSphere 1:00140-16.

53. Trottier MC, Pereira TdO, Groleau M-C, Hoffman LR, Dandekar AA, Déziel E. 2024. The end of the reign of a "master regulator"? A defect in function of the LasR quorum sensing regulator is a common feature of *Pseudomonas aeruginosa* isolates. mBio 15:e02376-23.

54. Wilder CN, Allada G, Schuster M. 2009. Instantaneous within-patient diversity of *Pseudomonas aeruginosa* quorum-sensing populations from cystic fibrosis lung infections. Infect Immun 77:5631-9.

55. Hoffman LR, Kulasekara HD, Emerson J, Houston LS, Burns JL, Ramsey BW, Miller SI. 2009. *Pseudomonas aeruginosa lasR* mutants are associated with cystic fibrosis lung disease progression. Journal of Cystic Fibrosis 8:66-70.

56. Mould DL, Stevanovic M, Ashare A, Schultz D, Hogan DA. 2022. Metabolic basis for the evolution of a common pathogenic *Pseudomonas aeruginosa* variant. eLife 11:e76555.

57. Mould Dallas L, Finger Carson E, Conaway A, Botelho N, Stuut Stacie E, Hogan Deborah A. 2024. Citrate cross-feeding between *Pseudomonas aeruginosa* genotypes supports *lasR* mutant fitness. mbio doi:01278-23:e01278-23.

58. D'Argenio DA, Wu M, Hoffman LR, Kulasekara HD, Déziel E, Smith EE, Nguyen H, Ernst RK, Larson Freeman TJ, Spencer DH. 2007. Growth phenotypes of *Pseudomonas aeruginosa lasR* mutants adapted to the airways of cystic fibrosis patients. Molecular microbiology 64:512-533.

59. Yeung ATY, Bains M, Hancock REW. 2011. The Sensor Kinase CbrA Is a Global Regulator That Modulates Metabolism, Virulence, and Antibiotic Resistance in *Pseudomonas aeruginosa*. Journal of Bacteriology 193:918-931.

60. García-Mauriño SM, Pérez-Martínez I, Amador CI, Canosa I, Santero E. 2013. Transcriptional activation of the CrcZ and CrcY regulatory RNAs by the CbrB response regulator in *Pseudomonas putida*. Molecular microbiology 89:189-205.

61. Valentini M, García-Mauriño SM, Pérez-Martínez I, Santero E, Canosa I, Lapouge K. 2014. Hierarchical management of carbon sources is regulated similarly by the CbrA/B systems in *Pseudomonas aeruginosa* and *Pseudomonas putida*. Microbiology 160:2243-2252.

62. Sonnleitner E, Abdou L, Haas D. 2009. Small RNA as global regulator of carbon catabolite repression in *Pseudomonas aeruginosa*. Proceedings of the National Academy of Sciences 106:21866-21871.

63. Sonnleitner E, Valentini M, Wenner N, Haichar FeZ, Haas D, Lapouge K. 2012. Novel targets of the CbrAB/Crc carbon catabolite control system revealed by transcript abundance in *Pseudomonas aeruginosa*. PloS one 7:e44637.

64. Yeung ATY, Janot L, Pena OM, Neidig A, Kukavica-Ibrulj I, Hilchie A, Levesque RC, Overhage J, Hancock REW. 2014. Requirement of the *Pseudomonas aeruginosa* CbrA sensor kinase for full virulence in a murine acute lung infection model. Infection and Immunity 82:1256-1267.

65. Tajmir Riahy M, Honarmand Jahromi S, Khaledi M, Afkhami H, Shafaati M, Lava Khamseh H, Zare Karizi S. 2021. Frequency of *cbrA*, *cbrB, ndvB,* and *phoBR* Genes in Relation to Biofilm Formation in *Pseudomonas aeruginosa* Clinical Isolates. Journal of Medical Microbiology and Infectious Diseases 9:97-102.

66. Palmer KL, Aye LM, Whiteley M. 2007. Nutritional cues control *Pseudomonas aeruginosa* multicellular behavior in cystic fibrosis sputum. Journal of Bacteriology 189:8079-8087.

67. Moore GE, Gerner RE, Franklin HA. 1967. Culture of normal human leukocytes. JAMA 199:519-524.

68. Mössinger J. 1991. In vitro cultivation of adult *Litomosoides carinii*: evaluation of basic culture media, gas phases and supplements. Parasitology 103:85-95.

69. Schurig-Briccio LA, Rintoul MR, Volentini SI, Farías RN, Baldomà L, Badía J, Rodríguez-Montelongo L, Rapisarda VA. 2008. A critical phosphate concentration in the stationary phase maintains ndh gene expression and aerobic respiratory chain activity in *Escherichia coli*. FEMS Microbiology Letters 284:76-83.

70. Chekabab SM, Harel J, Dozois CM. 2014. Interplay between genetic regulation of phosphate homeostasis and bacterial virulence. Virulence 5:786-793.

71. Hou CI, Gronlund AF, Campbell JJR. 1966. Influence of phosphate starvation on cultures of *Pseudomonas aeruginosa*. Journal of Bacteriology 92:851-855.

670 72. Xin L, Deliang L, Heng Y, Dingpeng Y, Shuang Z, Shuting Z, Mengqiu L, Miao H,
671   Min X, Xingguo W. 2016. Phosphatidylcholine affects the secretion of the alkaline
672   phosphatase PhoA in *Pseudomonas* strains. Microbiological Research 192:21-29.
673 73. Horwitz JP, Chua J, Noel M, Donatti JT, Freisler J. 1966. Substrates for
674   cytochemical demonstration of enzyme activity. II. Some Dihalo-3-indolyl
675   Phosphates and Sulfates1. Journal of Medicinal Chemistry 9:447-447.
676 74. Neidhardt FC, Bloch PL, Smith DF. 1974. Culture medium for *enterobacteria*.
677   Journal of Bacteriology 119:736-747.
678 75. Wanner B. 1993. Gene regulation by phosphate in enteric bacteria. Journal of
679   Cellular Biochemistry 51:47-54.
680 76. Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, D'Argenio DA,
681   Miller SI, Ramsey BW, Speert DP, Moskowitz SM, Burns JL, Kaul R, Olson MV.
682   2006. Genetic adaptation by *Pseudomonas aeruginosa* to the airways of cystic
683   fibrosis patients. Proceedings of the National Academy of Sciences 103:8487-
684   8492.
685 77. Gellatly SL, Bains M, Breidenstein EBM, Strehmel J, Reffuveille F, Taylor PK,
686   Yeung ATY, Overhage J, Hancock REW. 2018. Novel roles for two-component
687   regulatory systems in cytotoxicity and virulence-related properties in
688   *Pseudomonas aeruginosa*. AIMS Microbiology 4:173-191.
689 78. Ball G, Durand E, Lazdunski A, Filloux A. 2002. A novel type II secretion system
690   in *Pseudomonas aeruginosa*. Mol Microbiol 43:475-85.
691 79. Otero-Asman JR, Quesada JM, Jim KK, Ocampo-Sosa A, Civantos C, Bitter W,
692   Llamas MA. 2020. The extracytoplasmic function sigma factor σVreI is active
693   during infection and contributes to phosphate starvation-induced virulence of
694   *Pseudomonas aeruginosa*. Scientific Reports 10:3139.
695 80. Lewenza S, Falsafi R, Bains M, Rohs P, Stupak J, Sprott GD, Hancock RE. 2011.
696   The olsA gene mediates the synthesis of an ornithine lipid in *Pseudomonas*
697   *aeruginosa* during growth under phosphate-limiting conditions, but is not involved
698   in antimicrobial peptide susceptibility. FEMS Microbiology Letters 320:95-102.
699 81. Wu H, Kato J, Kuroda A, Ikeda T, Takiguchi N, Ohtake H. 2000. Identification and
700   characterization of two chemotactic transducers for inorganic phosphate in
701   *Pseudomonas aeruginosa*. Journal of Bacteriology 182:3400-3404.
702 82. Dekimpe V, Déziel E. 2009. Revisiting the quorum-sensing hierarchy in
703   *Pseudomonas aeruginosa*: the transcriptional regulator RhlR regulates LasR-
704   specific factors. Microbiology 155:712-723.
705 83. Higgins S, Heeb S, Rampioni G, Fletcher MP, Williams P, Cámara M. 2018.
706   Differential regulation of the phenazine biosynthetic operons by quorum sensing in
707   *Pseudomonas aeruginosa* PAO1-N. Frontiers in cellular and infection microbiology
708   8:252.
709 84. Recinos DA, Sekedat MD, Hernandez A, Cohen TS, Sakhtah H, Prince AS, Price-
710   Whelan A, Dietrich LE. 2012. Redundant phenazine operons in *Pseudomonas*
711   *aeruginosa* exhibit environment-dependent expression and differential roles in
712   pathogenicity. Proceedings of the National Academy of Sciences 109:19420-
713   19425.

85. Cui Q, Lv H, Qi Z, Jiang B, Xiao B, Liu L, Ge Y, Hu X. 2016. Cross-Regulation between the *phz1* and *phz2* operons maintain a balanced level of phenazine biosynthesis in *Pseudomonas aeruginosa* PAO1. PLOS ONE 11:e0144447.

86. Schmitz S, Rosenbaum MA. 2020. Controlling the production of *Pseudomonas* phenazines by modulating the genetic repertoire. ACS Chemical Biology 15:3244-3252.

87. Smyth P, Clarke PH. 1975. Catabolite repression of *Pseudomonas aeruginosa* amidase: the effect of carbon source on amidase synthesis. Microbiology 90:81-90.

88. Petrova OE, Schurr JR, Schurr MJ, Sauer K. 2011. The novel *Pseudomonas aeruginosa* two-component regulator BfmR controls bacteriophage-mediated lysis and DNA release during biofilm development through PhdA. Molecular microbiology 81:767-783.

89. Bains M, Fernández L, Hancock RE. 2012. Phosphate starvation promotes swarming motility and cytotoxicity of *Pseudomonas aeruginosa*. Applied and environmental microbiology 78:6762-6768.

90. Faure LM, Llamas MA, Bastiaansen KC, De Bentzmann S, Bigot S. 2013. Phosphate starvation relayed by PhoB activates the expression of the *Pseudomonas aeruginosa* σvreI ECF factor and its target genes. Microbiology 159:1315-1327.

91. Lewin GR, Kapur A, Cornforth DM, Duncan RP, Diggle FL, Moustafa DA, Harrison SA, Skaar EP, Chazin WJ, Goldberg JB. 2023. Application of a quantitative framework to improve the accuracy of a bacterial infection model. Proceedings of the National Academy of Sciences 120:e2221542120.

92. Turner KH, Wessel AK, Palmer GC, Murray JL, Whiteley M. 2015. Essential genome of *Pseudomonas aeruginosa* in cystic fibrosis sputum. Proceedings of the National Academy of Sciences 112:4110-4115.

93. Tan J, Huyck M, Hu D, Zelaya RA, Hogan DA, Greene CS. 2017. ADAGE signature analysis: differential expression analysis with data-defined gene sets. BMC Bioinformatics 18:512.

94. Heurlier K, Dénervaud V, Haenni M, Guy L, Krishnapillai V, Haas D. 2005. Quorum-sensing-negative *lasR* mutants of *Pseudomonas aeruginosa* avoid cell lysis and death. Journal of Bacteriology 187:4875-4883.

95. Clay ME, Hammond JH, Zhong F, Chen X, Kowalski CH, Lee AJ, Porter MS, Hampton TH, Greene CS, Pletneva EV. 2020. *Pseudomonas aeruginosa lasR* mutant fitness in microoxia is supported by an Anr-regulated oxygen-binding hemerythrin. Proceedings of the National Academy of Sciences 117:3167-3173.

96. Wargo MJ, Gross MJ, Rajamani S, Allard JL, Lundblad LKA, Allen GB, Vasil ML, Leclair LW, Hogan DA. 2011. Hemolytic phospholipase C inhibition protects lung function during *Pseudomonas aeruginosa* infection. American Journal of Respiratory and Critical Care Medicine 184:345-354.

97. Huang J, Sonnleitner E, Ren B, Xu Y, Haas D. 2012. Catabolite repression control of pyocyanin biosynthesis at an intersection of primary and secondary metabolism in *Pseudomonas aeruginosa*. Applied and environmental microbiology 78:5016-5020.

98. Hiratsu K, Nakata A, Shinagawa H, Makino K. 1995. Autophosphorylation and
activation of transcriptional activator PhoB of *Escherichia coli* by acetyl phosphate
in vitro. Gene 161:7-10.
99. McCleary W. 1996. The activation of PhoB by acetylphosphate. Molecular
Microbiology 20:1155-1163.
100. Henry JT, Crosson S. 2011. Ligand-Binding PAS domains in a genomic, cellular,
and structural context. Annual Review of Microbiology 65:261-286.
101. Qi Y, Rao F, Luo Z, Liang ZX. 2009. A flavin cofactor-binding PAS domain
regulates c-di-GMP synthesis in AxDGC2 from *Acetobacter xylinum*. Biochemistry
48:10275-85.
102. Dupré E, Clantin B, Yuan Y, Lecher S, Lesne E, Antoine R, Villeret V, Jacob-
Dubuisson F. 2021. Structural insight into the role of the PAS domain for signal
transduction in sensor kinase BvgS. Journal of Bacteriology 203:e00614-20.
103. Xing J, Gumerov VM, Zhulin IB. 2023. Origin and functional diversification of PAS
domain, a ubiquitous intracellular sensor. Science Advances 9:eadi4517.
104. Bertani G. 1951. Studies on lysogenesis. I. The mode of phage liberation by
lysogenic *Escherichia coli*. J Bacteriol 62:293-300.
105. Gibson DG, Glass JI, Lartigue C, Noskov VN, Chuang RY, Algire MA, Benders
GA, Montague MG, Ma L, Moodie MM, Merryman C, Vashee S, Krishnakumar R,
Assad-Garcia N, Andrews-Pfannkoch C, Denisova EA, Young L, Qi ZQ, Segall-
Shapiro TH, Calvey CH, Parmar PP, Hutchison CA, 3rd, Smith HO, Venter JC.
2010. Creation of a bacterial cell controlled by a chemically synthesized genome.
Science 329:52-6.
106. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. 2009.
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nature
Methods 6:343-5.
107. Shanks RM, Caiazza NC, Hinsa SM, Toutain CM, O'Toole GA. 2006.
Saccharomyces cerevisiae-based molecular tool kit for manipulation of genes from
gram-negative bacteria. Applied Environmental Microbiology 72:5027-36.
108. Sacks L. 1956. A pH gradient agar plate. Nature 178:269-270.
109. Grahl N, Dolben EL, Filkins LM, Crocker AW, Willger SD, Morrison HG, Sogin ML,
Ashare A, Gifford AH, Jacobs NJ, Schwartzman JD, Hogan DA. 2018. Profiling of
bacterial and fungal microbial communities in cystic fibrosis sputum using RNA.
mSphere 3:00292-18.
110. Team RC. 2021. R: A language and environment for statistical computing., R
Foundation for Statistical Computing, Vienna, Austria.
795

Fig. 1 *P. aeruginosa* LasR- strains require lower Pi to repress PhoB activity. A) *P. aeruginosa* WT, Δ*lasR*, Δ*phoB* and Δ*pstB* were spotted on gradient plates of MOPS agar with a range of Pi concentrations (0.1 - 1 mM) and 60 µg/mL BCIP to indicate alkaline phosphatase (AP) activity. **B)** *P. aeruginosa* strains, including DH2590 (a LasR- clinical isolate (CI) and a derivative in which the *lasR* gene was replaced with the gene from strain PA14) on MOPS agar with BCIP and 0.7 mM Pi C. **C)** *P. aeruginosa* CI LasR+/LasR- pairs from the sputum of two pwCF. **D)** Colony biofilms on MOPS agar with 0.2% glucose and either 0.2 or 10 mM Pi. For **A-D**, similar results were obtained in three replicate experiments; a representative experiment is shown. For panels **E-G**, *P. aeruginosa* was grown as colony biofilms on MOPS agar with 0.7 mM Pi. **E)** AP activity in WT and Δ*lasR* after 12 h at 37 °C. Data from replicates collected on the same day have the same shape. Data analyzed using a paired, two-tailed t-test (n = 12). **F)** *phoA* transcripts in WT and Δ*lasR* were measured by qRT-PCR on different days and normalized to the housekeeping gene transcript *ppiD*. Data analyzed using a paired, two-tailed t-test (n = 8). **G)** Levels of PhoB-controlled transcripts and *phoB* itself were assessed using NanoString multiplex technology. Nine PhoB-regulated transcripts are shown as normalized counts. Data were analyzed using a two-way ANOVA; significant differences between wild type-Δ*lasR,* wild type-Δ*phoB*, and Δ*lasR*-Δ*lasR*Δ*phoB* (p < 0.0001, n = 2-3). There were no significant differences between the Δ*phoB* and Δ*lasR*Δ*phoB* mutants (p = 0.46, n = 3). Statistical differences for each transcript are available in Supplemental File 1. For all panels, asterisks denote significance (p ≤ 0.01 = **, p ≤ 0.001 = ***, p ≤ 0.0001 = ****).

Fig. 2 *P. aeruginosa* quorum qensing (QS) mutants have active PhoB at higher Pi concentrations than the wild type. A) Colony biofilms of WT and the indicated QS mutants (Δ*lasR*, Δ*rhlR*, Δ*pqsR*) were grown on MOPS agar with 0.7 mM Pi and BCIP (top) or on medium without BCIP for analysis of AP activity using the colorimetric PNPP substrate. Data from replicates collected on the same day have the same shape. Data analyzed by an ordinary one-way ANOVA with Tukey's multiple comparisons tests (n = 4). **B)** *P. aeruginosa* was grown on a plate with a gradient of Pi concentrations with 0. 5 to 1.2 mM shown. Quantitative data analyzed using a one-way ANOVA (n = 3) (asterisks denote significance from the wild type). Asterisks denote significance (p ≤ 0.05 = *, p ≤ 0.01 = **, p ≤ 0.001 = ***, p ≤ 0.0001 = ****).

Fig. 3 The loss of phenazines promotes PhoB activity. A) AP activity in WT and mutants Δ*phz1*, Δ*phz2*, and Δ*phz* (lacking *phz1* and *phz2* operons) grown on MOPS agar with 0.7 mM Pi and BCIP (top) or on medium without BCIP for analysis of AP activity using the colorimetric PNPP substrate. Data from replicates collected on the same day have the same shape. Data were analyzed using a one-way ANOVA and Tukey's multiple comparisons test (n = 7). **B)** Analysis of wild type, and the Δ*lasR*, Δ*phz*, and Δ*lasR*Δ*phz* mutants. The Δ*lasR*Δ*phz* mutant had significantly more AP activity than the Δ*lasR* and Δ*phz* strains. There are no significant differences between Δ*lasR* and Δ*phz* (p=0.99) (n = 12). **C)** *P. aeruginosa* was grown on plates with a gradient of Pi; the average concentration of Pi that inhibits AP activity is graphed to the right (n =3). Data analyzed using a one-way ANOVA; asterisks denote significance from the wild type. Asterisks denote significance (p ≤ 0.05 = *, p ≤ 0.01 = **, p ≤ 0.001 = ***, p ≤ 0.0001 = ****).

Fig. 4. The CbrA-CbrB-Crc pathway promotes PhoR-PhoB activity in LasR- strains. A) A proposed model of the relationship between the CbrA-CbrB and PhoR-PhoB two component systems; *crcZ*, a small RNA; Crc, which acts as a translational repressor in complex with Hfq. For figures **B** and **C**, AP activity in indicated strains grown on MOPS agar with 0.7 mM Pi and BCIP (top) or on medium without BCIP for analysis of AP activity using the colorimetric PNPP substrate. Data from replicates collected on the same day have the same shape. Data were analyzed using a one-way ANOVA and Tukey's multiple comparisons test. **B)** AP activity in Δ*lasR*, Δ*lasR*ΔcbrB, Δ*lasR*Δ*cbrB*+cbrB, and Δ*lasR*Δ*cbrB* +*crc* mutants (n = 3). **C)** AP activity in the wild type (WT), Δ*crc* mutant and its complemented derivative is shown (n = 5). **D)** AP activity in Δ*lasR*, Δ*crc*, and Δ*lasR*Δ*crc* (n = 8). **E)** *P. aeruginosa* was grown on plates with a gradient of Pi (0.5 – 1.5 mM Pi) and BCIP. The average concentration of Pi which inhibited AP activity is graphed to the right (n = 3). Asterisks denote significance (p ≤ 0.05 = *, p ≤ 0.01 = **, p ≤ 0.001 = ***).

Fig. 5 PhoB activity mediates fitness advantages and antagonism in Δ*lasR P. aeruginosa* **A)** *P. aeruginosa* wild type (WT), Δ*lasR*, Δ*phoB*, and Δ*lasR*Δ*phoB* was grown in MOPS liquid medium with no added Pi at 37 °C. Data analyzed using a two-way ANOVA with Tukey's multiple comparisons test (n = 3). Asterisks denote significance (p ≤ 0.0001 = ****). **B)** *P. aeruginosa* colonies were grown on a lawn of *Candida albicans*. Wild-type (LasR+) *P. aeruginosa* produces 5MPCA which is converted to red pigment inside *C. albicans*. The Δ*lasR* mutant produced more 5MPCA than the wild type. 5MPCA production required PhoB and CbrB in both strains (n = 3). **C)** NanoString analysis of PhoB-regulated transcripts shown as normalized counts (n = 2 - 3). All transcripts shown were significantly higher in the Δ*lasR* mutant compared to both the wild type (p ≤ 0.0001) and the Δ*lasR*Δ*phoB* mutant (p ≤ 0.0016). **D)** PhoB contribution to regulation of QS genes by NanoString analysis of QS-regulated transcripts between the wild type (WT) or Δ*phoB* mutant (p>0.30) and the Δ*lasR* and Δ*lasR*Δ*phoB*. All transcripts were significantly lower in the Δ*lasR*Δ*phoB* mutant than the Δ*lasR* mutant (p ≤ 0.01). *lasB, rhlR, pqsR,* and *pqsH* were significantly lower in the Δ*lasR* mutant than the wild type (p < 0.0001). Complete NanoString dataset available in Supplemental File 1.

Fig. 6 PhoB-regulated transcripts are higher *in vivo*. RNA-seq data from Lewin, *et al.* (2023) were arranged by sample type (Human Sputum or PAO1 grown in SCFM) and PhoB-regulated transcripts were Z-scored and sorted by unsupervised hierarchical clustering via the R pheatmap function. RNA-seq data is available in Supplemental File 2.