Loss of LasR function leads to decreased repression of *Pseudomonas aeruginosa* PhoB activity at physiological phosphate concentrations

Amy Conaway[1], Igor Todorovic[1], Dallas L. Mould[1], and Deborah A. Hogan[1#]

[1]Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth, Hanover, NH USA

Running Title: Increased PhoB activity in LasR- strains

Key Words: PhoB, LasR, PhoR, quorum sensing, phosphate scavenging, phenazines, CbrB, RhlR, Crc, *Pseudomonas aeruginosa*

#To whom correspondence should be addressed:

Department of Microbiology and Immunology

Geisel School of Medicine at Dartmouth

Rm 208 Vail Building, Hanover, NH 03755

E-mail: dhogan@dartmouth.edu

Tel: (603) 650-1252

Abstract

While the *Pseudomonas aeruginosa* LasR transcription factor plays a role in quorum sensing (QS) across phylogenetically-distinct lineages, isolates with loss-of-function mutations in *lasR* (LasR– strains) are commonly found in diverse settings including infections where they are associated with worse clinical outcomes. In LasR– strains, the transcription factor RhlR, which is controlled by LasR, can be alternately activated in low inorganic phosphate (Pi) concentrations via the two-component system PhoR-PhoB. Here, we demonstrate a new link between LasR and PhoB in which the absence of LasR increases PhoB activity at physiological Pi concentrations and raises the Pi concentration necessary for PhoB inhibition. PhoB activity was also less repressed by Pi in mutants lacking different QS regulators (RhlR and PqsR) and in mutants lacking genes required for the production of QS-regulated phenazines suggesting that decreased phenazine production was one reason for decreased PhoB repression by Pi in LasR– strains. In addition, the CbrA-CbrB two-component system, which is elevated in LasR– strains, was necessary for reduced PhoB repression by Pi and a Δ*crc* mutant, which lacks the CbrA-CbrB-controlled translational repressor, activated PhoB at higher Pi concentrations than the wild type. The Δ*lasR* mutant had a PhoB-dependent growth advantage in a medium with no added Pi and increased virulence-determinant gene expression in a medium with physiological Pi, in part through reactivation of QS. This work suggests PhoB activity may contribute to the virulence of LasR– *P. aeruginosa* and subsequent clinical outcomes.

Importance

Loss-of-function mutations in the gene encoding the *Pseudomonas aeruginosa* quorum sensing (QS) regulator LasR occur frequently and are associated with worse clinical outcomes. We have found that LasR– *P. aeruginosa* have elevated PhoB activity at physiological concentrations of inorganic phosphate (Pi). PhoB activity promotes Pi acquisition as well as the expression of QS and virulence-associated genes. Previous work has shown that PhoB induce RhlR, another QS regulator, in a LasR- mutant in low Pi conditions. Here, we demonstrate a novel relationship wherein LasR represses PhoB activity, in part through the production of phenazines and Crc-mediated translational repression. This work suggests PhoB activity may contribute to the increased virulence of LasR– *P. aeruginosa.*

Introduction

Pseudomonas aeruginosa is a pernicious pathogen that infects burns, nonhealing wounds, eyes, and the airways of people with cystic fibrosis (pwCF) or chronic obstructive pulmonary disease (COPD). P. aeruginosa fitness in vivo is due to its ability to establish persistent biofilms and acquire critical nutrients, such as phosphorus, in the host environment. Phosphate is required for cell membranes, nucleic acids, metabolic intermediates, and is used in some signal transduction pathways. The most accessible form of phosphorus is free inorganic phosphate (Pi) which is at ~1.3 mM in the serum of healthy adults.[1,2] During infection, however, Pi is restricted by the host as part of nutritional immunity through different mechanisms including the secretion of phosphate-binding proteins.[3-5]

Most host phosphorus is in organic forms that require degradation prior to uptake and utilization by microbes.[6,7] P. aeruginosa induces the production of phosphatases, phospholipases, and DNases to access organic phosphate along with high-affinity phosphate transporters in response to low Pi levels via the transcription factor PhoB and its sensor kinase PhoR.[8-11] In P. aeruginosa, PhoB also induces the production of specific phenazine small molecules [12-16] which solubilize phosphate from minerals.[17] PhoR, which activates PhoB through phosphorylation, is regulated by interactions with the high-affinity Pi transporter PstABC via PhoU such that Pi transport inhibits PhoR activation.[10,18-25] While PhoB is known to positively regulate its own expression [26], data suggest that PhoB also participates in a negative feedback loop [27], presumably to limit intracellular Pi concentrations, as has been described in Escherichia coli.[28]

In addition to the role of the PhoR-PhoB two-component system in Pi acquisition

in *P. aeruginosa* and other species such as *E. coli* [24, 29], *Salmonella enterica* [7], and

Vibrio cholerae [30], these regulators also play diverse roles in the regulation of virulence

in multiple pathogens.[10] In *E. coli*, type III secretion system components and effectors

are upregulated during Pi limitation in a PhoB-dependent manner.[31] In *V. cholerae*, PhoB

inhibits transcription of the genes encoding TcpPH, which positively regulate expression

of virulence factors including toxin co-regulated pilus.[30, 32] In *P. aeruginosa,* cross-

regulation between PhoR-PhoB and quorum sensing (QS), which contributes to virulence

and biofilm formation [33-35], has been described. *P. aeruginosa* QS is largely controlled

by three transcription factors, LasR, RhlR, and PqsR, that are active when QS inducers

are at sufficient concentrations, i.e. high cell densities or environments with decreased

diffusion. While LasR positively regulates both RhlR and PqsR, many groups have shown

that RhlR and PqsR can be activated in the absence of LasR.[36-39] Furthermore, in low Pi

conditions, PhoB induces QS [14-16, 40, 41] and this can occur in a LasR-independent

manner.[13, 42, 43] One group of virulence factors induced by both PhoB and QS are

phenazines which inhibit immune cell function [44, 45] and kill other microbes [46-50] through

oxidative stress. Though PhoB binding sites have been identified upstream of some

phenazine biosynthesis genes [16], PhoB induction of phenazine production is largely

attributed to increased expression of *rhlR*.[13, 16, 40] Furthermore, *phoR, phoB, rhlR,* and

pqsR are all required for *P. aeruginosa* phenazine production in co-culture with *Candida*

albicans [27], a fungus that frequently co-infects the airways of pwCF.[51] PhoB regulation

of QS suggests low Pi could be a signal for QS remodeling to circumvent reliance on

LasR.

P. aeruginosa LasR– strains are found in the environment and isolates from both

acute and chronic infections like those in the lungs of pwCF where they make up

approximately one-third of clinical isolates.[36, 52-57] LasR– isolates are associated with

worse disease progression in pwCF [57] and worse lesions during acute ocular

infections.[54] In laboratory evolution experiments, the increased fitness of *P. aeruginosa*

LasR– lineages depends on increased CbrA-CbrB activity [58, 59] which leads to growth

advantages in complex media.[37, 58, 60] *P. aeruginosa* CbrA-CbrB is a two-component

system that induces expression of the small RNA *crcZ* which sequesters Crc.[61-64] Crc,

with Hfq, binds to multiple mRNA targets, many of which encode transporters and

catabolic enzymes [65], and inhibits their translation. CbrA is required for full *P. aeruginosa*

virulence [66].

P. aeruginosa virulence regulation is often studied in laboratory media with Pi

concentrations that repress PhoR activation of PhoB (e.g. synthetic CF medium (SCFM),

5.1 mM [67]; RPMI medium, 5.6 mM [68, 69]; Luria Broth, 6 mM [70]; or M9 medium, 64 mM

[70]). PhoR-PhoB activity is generally studied in media with less than 0.5 mM Pi.[13, 20, 24,

42] These two extremes in Pi concentrations either strongly repress or strongly induce

PhoB activity and thus may not be relevant to our understanding of any contributions of

PhoB to *P. aeruginosa* virulence at physiological Pi concentrations (1.3 mM Pi in serum

[1, 4]). Thus, we aimed to elucidate the relationship between LasR and PhoB at

physiologically relevant Pi concentrations. In these studies, we show that LasR– isolates

and a Δ*lasR* mutant had elevated PhoR-dependent PhoB activity relative to comparator

strains with functional LasR at Pi concentrations in the 0.7 – 1.1 mM range. In contrast,

LasR+ and LasR– strains had similar PhoB activity in both low (0.2 mM) and high (10

mM) Pi conditions. Our data demonstrate that a lack of phenazines or reduced Crc activity

via CbrA-CbrB led to higher PhoB activity and decreased repression of PhoR-PhoB by

Pi. We show that PhoB is required for the Δ*lasR* mutant growth advantages in a medium

with no added Pi and the increased expression of multiple virulence factors, including

phenazine biosynthetic enzymes and phospholipases, at physiological Pi. This work

establishes a novel connection between QS and PhoB wherein LasR represses PhoB

activity and PhoB is required for increased virulence gene expression in the Δ*lasR* mutant,

in part through reactivation of QS. This model for virulence regulation may aid in

understanding why *P. aeruginosa* LasR– strains are associated with poor clinical

outcomes.

Results

PhoB activity is repressed at lower Pi concentrations in *P. aeruginosa* LasR+

strains than in their *P. aeruginosa* LasR– counterparts.

Though PhoB regulates virulence factor production across species [10, 30, 31], there

are only a limited number of in vitro studies on PhoB activity at Pi concentrations similar

to those found in human hosts. Thus, we sought to determine the Pi concentrations

necessary to repress PhoB activity. In *P. aeruginosa* strain PA14, PhoB activity was

monitored by assessing the activity of alkaline phosphatase (AP), which is encoded by

the PhoB-regulated gene *phoA* [8, 71, 72] in colonies on agar containing the colorimetric AP

substrate bromo-4-chloro-3-indolyl phosphate (BCIP) [9, 73] (Fig. 1A). Using gradient

plates with range of Pi concentrations (0.1 mM - 1 mM) [27] in MOPS-glucose medium [74],

we found that 0.7 mM Pi repressed AP activity in the wild type. Interestingly, a Δ*lasR*

mutant had AP activity across the entire gradient with a reduction in activity at higher Pi

concentrations (Fig. 1A). Consistent with PhoB regulation of AP activity, the ΔphoB

mutant had no detectable AP activity at any concentration, and the ΔpstB mutant, which

has constitutive PhoB activity due to de-repression of PhoR [27, 75], had strong AP

production at all Pi concentrations tested (Fig. 1A). At 0.7 mM Pi, these was AP activity

in colonies of the ΔlasR mutant, but not the wild type colonies or those formed by the

ΔlasR mutant complemented with lasR at the native locus (Fig. 1B). As expected, neither

the ΔlasRΔphoB or ΔlasRΔphoR mutants had AP activity (Fig. 1B). Furthermore, a clinical

isolate from a corneal infection, DH2590 (262K in Hammond, et al. [54]), which has a loss-

of-function (LOF) mutation in lasR, encoding the non-functional LasR^{I215S} variant, also

expressed AP at 0.7 mM Pi and AP activity was reduced by replacing the endogenous

lasR allele with a functional PA14 lasR allele at the native locus (Fig. 1B). In addition, we

compared AP activity in two clinical isolate pairs (Fig. 1C) from CF sputum samples

wherein one isolate has a LOF mutation in lasR and the other does not.[76] In both cases,

the LasR– isolates had more AP activity than their LasR+ counterparts at 0.7 mM Pi.

Both the ΔlasR mutant and the wild type showed PhoB activity at lower Pi

concentrations (< 0.7 mM). We found that at 0.2 mM Pi, the concentration frequently used

to study PhoB activity [13, 14, 20, 24, 77], the wild type and ΔlasR strains had similar AP activity

(Fig. 1D). Additionally, neither strain showed AP production at 10 mM Pi (Fig. 1D) and no

AP activity was observed in either the wild-type or the ΔlasR strain on LB agar, which is

reported to have 6 mM Pi [70] (Fig. S1A). The ΔphoB mutant did not have AP activity even

at 0.2 mM Pi and the ΔpstB mutant had AP production at 0.2 mM Pi, 10 mM Pi, and on

LB. To quantify differences AP activity, we used the soluble colorimetric substrate p-

nitrophenyl phosphate (PNPP). The ΔlasR mutant had significantly more AP activity than

the wild type at 0.7 mM Pi (Fig. 1E) while growth was similar (Fig. S1B). Similarly, qRT-

PCR analysis of *phoA*, which encodes AP, found significantly higher transcript levels in

the ΔlasR mutant compared to the wild type (Fig. 1F).

To directly assess the levels of transcripts of other genes within the PhoB regulon,

we utilized NanoString multiplex technology as previously published [27]. At 0.7 mM Pi,

the ΔlasR mutant had significantly higher levels of PhoB-regulated transcripts including

phoB itself and genes encoding phosphatases (*phoA* and *lapA*), a phosphate transporter

(*oprO*), a type two secretion system protein required for secretion of LapA (*hxcT*) [78], a

putative TonB transporter (*exbD2*) [79], a phosphodiesterase (*glpQ*), a low-phosphate

ornithine lipid biosynthetic protein (*olsA*) [80], and a Pi chemotaxis protein (*ctpL*).[81] None

of these transcripts were significantly different when the ΔphoB and ΔlasRΔphoB mutants

were compared (Fig. 1G and Supplemental File 1).

Other QS mutants and a mutant lacking QS-controlled phenazines have active

PhoB at higher Pi concentrations than the wild type

LasR positively regulates other QS transcription factors including RhlR and PqsR

and some ΔlasR mutant phenotypes are due to their decreased activity. However, RhlR

and PqsR are capable of inducing QS-controlled genes in the absence of LasR in specific

strains and conditions.[36, 37, 59, 82] Thus, we assessed the role of RhlR and PqsR in the

repression of PhoB. As shown in Fig. 2A, AP activity was significantly elevated in the

ΔrhlR and ΔpqsR mutants, as in the ΔlasR mutant, relative to the wild type at 0.7 mM Pi.

Across a gradient plate of 0.5 – 1.5 mM Pi, AP activity in all QS mutants was inhibited at

a significantly higher Pi concentration than the wild type (Fig. 2B). While AP activity in the

Δ*rhlR* and Δ*pqsR* strains was inhibited by Pi > 0.9 mM, activity in the LasR— strains

(Δ*lasR,* Δ*lasR*Δ*rhlR,* and Δ*lasR*Δ*pqsR*) was inhibited by Pi > 1 mM (Fig. 2B).

As each of the QS transcription factor mutants, Δ*lasR,* Δ*rhlR,* and Δ*pqsR,* produce

fewer phenazines than the wild type in late exponential and early stationary phase

cultures [83, 84], we tested whether the absence of phenazines was sufficient to increase

PhoB activity at moderate Pi concentrations. *P. aeruginosa* produces multiple phenazines

including phenazine-1-carboxylic acid (PCA and the PCA derivatives 5-methyl-PCA

(5MPCA), pyocyanin (PYO), and phenazine-1-carboxamide. PCA is synthesized by

proteins encoded by two highly similar operons, *phzA1-G1* and *phzA2-G2.* The mutant

lacking both *phz* operons (Δ*phzA1-G1*Δ*phzA2-G2*), referred to as Δ*phz,* had significantly

elevated AP activity compared to the wild type at 0.7 mM Pi (Fig. 3A). The Δ*phz1* mutant

(Δ*phzA1-G1*) was not different from the wild type and deletion of either of the adjacent

phenazine-modifying genes required for 5MPCA and PYO biosynthesis, *phzM* or *phzS,*

did not lead to changes in AP production at 0.7 mM (Fig. S2A). In contrast, the Δ*phz2*

mutant (Δ*phzA2-G2*) phenocopied Δ*phz* and had significantly more AP production than

the Δ*phz1* mutant or the wild type (Fig. 3A). We and others have previously published

data identifying *phzA2-G2* as the predominant contributor of PCA.[27, 84-86]

We found no difference in AP activity at 0.7 mM Pi between the Δ*lasR* and Δ*phz*

mutants (Fig. 3B). However, a Δ*lasR*Δ*phz* mutant had significantly elevated AP activity

compared to both single mutants (Fig. 3B). Both the Δ*lasR* and Δ*lasR*Δ*phz* mutants had

detectable AP activity at significantly higher Pi concentrations than Δ*phz* (Fig. 3C)

suggesting that PhoB activity was derepressed by an additional phenazine-independent

mechanism in the Δ*lasR* mutant. Consistent with this model, the Δ*phz* mutant was

repressed by a similar Pi concentration as the Δ*rhlR* and Δ*pqsR* mutants, but not their

LasR– counterparts (Fig. 2B).

CbrA-CbrB-Crc impacts PhoB activity in the Δ*lasR* mutant.

Several studies suggest that *P. aeruginosa* strains lacking LasR activity have

elevated CbrA-CbrB activity in many conditions.[58, 60] CbrA-CbrB activity leads to relief of

Crc-mediated translational repression of diverse mRNA targets including many that

encode catabolic enzymes and transporters [64, 65, 87] (Fig. 4A). In both BCIP and PNPP

assays, we found that in a Δ*lasR*Δ*cbrB* mutant did not have AP activity at 0.7 mM Pi and

that AP production was restored by complementation of *cbrB* at the native locus (Fig 4B).

PhoB activity in both the Δ*cbrB* and Δ*lasR*Δ*cbrB* mutants was similarly repressed when

Pi > 3 mM (Fig. S2B). Additionally, AP activity in colonies grown on medium with 0.7 mM

Pi was restored to a Δ*lasR*Δ*cbrB* mutant by deletion of *crc* (Fig. 4B) suggesting that the

increase in Pi repression of PhoB in the Δ*lasR*Δ*cbrB* is due to elevated Crc activity (Fig.

4A for pathway). A Δ*crc* single mutant also had significantly elevated AP activity

compared to the wild type and *crc*-complemented strain (Fig. 4C). There was no

difference in AP activity between the Δ*lasR* and Δ*lasR*Δ*crc* mutants at 0.7 mM Pi, though

both were significantly elevated compared to the Δ*crc* mutant (Fig. 4D). Additionally, the

Δ*lasR* and Δ*lasR*Δ*crc* mutants both showed significantly decreased PhoB sensitivity to Pi

than the Δ*crc* mutant (Fig. 4E).

PhoB activity in the *P. aeruginosa* Δ*lasR* strain provides fitness advantages when

Pi is limited.

To determine if the increased PhoB activity in Δ*lasR* cells increased fitness upon

depletion of Pi, we grew the wild type, the Δ*lasR* mutant, and their respective Δ*phoB*

derivatives overnight in LB followed by sub-culture into MOPS-glucose medium with no

added Pi (Fig. 5A). The wild type and the Δ*lasR* mutant showed similar initial growth rates.

However, in the post-exponential phase, the Δ*lasR* mutant reached significantly higher

densities than the wild type. Both the Δ*phoB* and Δ*lasR*Δ*phoB* mutants showed minimal

growth in the no added Pi media and were not significantly different from each other.

PhoB activity mediates the expression of virulence determinants in LasR– *P.

aeruginosa, in part through QS*.*

Multiple studies have shown that there are alternative mechanisms for the

activation of RhlR in LasR– strains [13, 36-38, 42, 43] and that phosphate limitation promotes

RhlR activity.[13-16, 42, 43] We have previously published that both PhoB and RhlR are

required for 5MPCA production by *P. aeruginosa* resulting in antagonism of *C. albicans* in

co-culture.[27] A derivative of 5MPCA accumulates inside *C. albicans* cells as a red

pigment*.* Here we show evidence of increased 5MCPA production by the Δ*lasR* mutant in

co-culture with *C. albicans.* 5MPCA production by both LasR+ and LasR– *P. aeruginosa*

required PhoB and CbrB (Fig. 5B).

The NanoString codeset used in Fig. 1G included genes that encode proteins

associated with phosphate acquisition, virulence, and QS.[27] We analyzed the abundance

of multiple transcripts encoding virulence-associated proteins, including phenazine

biosynthetic enzymes (*phzM* and *phzS*), phospholipases (*plcH* and *plcN*), and an

exopolysaccharide matrix regulator (*phdA*) [88] in the wild type and Δ*lasR, ΔphoB,* and

Δ*lasR*Δ*phoB* mutants. These virulence-associated transcripts were significantly higher in

the Δ*lasR* mutant compared to the wild type (Fig. 5C and Supplemental File 1). Three of

these transcripts have also been identified as part of the PhoB regulon (*plcH* [89], *plcN* [12,

90, 91] and *phdA* [79, 92]) while others have not (*phzM* and *phzS*). However, all five transcripts

were significantly lower in the Δ*lasR*Δ*phoB* mutant compared to the Δ*lasR* strain and no

differences were detected when the Δ*phoB* and Δ*lasR*Δ*phoB* mutants were compared.

While *plcN* and *phdA* were significantly lower in Δ*phoB* compared to the wild type, *phzM,*

phzS, and *plcH*, were only PhoB-dependent in the Δ*lasR* mutant. We then sought to

determine if PhoB contributes to QS. We found significantly higher abundance of

transcripts encoding proteins involved in QS regulation (*rhlR* and *pqsR*) and autoinducer

synthesis (*lasI* and *pqsH*) as well as transcripts that can be induced by RhlR (*rhlA* and

lasB) in the Δ*lasR* mutant compared to the Δ*lasR*Δ*phoB* mutant (Fig. 5D and

Supplemental File 1). As expected, the *lasB, rhlR, pqsR*, and *pqsH* transcripts were also

significantly lower in the Δ*lasR* mutant compared to the wild type. Consistent with the

model that PhoB is not active in the wild type at 0.7 mM Pi, there were no significant

differences in QS-related transcript abundance between the Δ*phoB* mutant and the wild

type.

Discussion

In this manuscript, we showed that LasR– *P. aeruginosa* laboratory strains and

clinical isolates had elevated PhoB-dependent AP activity and increased expression of

the PhoB regulon at 0.7 mM Pi. We found that the range of PhoB-permissive Pi

concentrations was broader for LasR– *P. aeruginosa*, though PhoB was still repressed at

high Pi concentrations. Importantly, these intermediate Pi concentrations where LasR–

P. aeruginosa showed more PhoB activity are similar to those found in the serum of

healthy adults.[1, 2] We also found that elevated PhoB activity was common across QS

mutants and this was possibly due to reduced phenazine production. However, LasR– *P.*

aeruginosa had active PhoB at even higher Pi concentrations than the other QS mutants,

suggesting differences in PhoB regulation.

LasR– *P. aeruginosa* are more fit than their LasR+ counterparts in many contexts.

LasR– *P. aeruginosa* are known to have elevated CbrA-CbrB activity, conferring growth

advantages in complex media through decreased Crc activity.[58, 60] Other studies have

shown that LasR– *P. aeruginosa* are more resistant to cell lysis at high pH [93] and more

fit in microoxic conditions and normoxically grown colony biofilms.[94] Future studies are

necessary to determine if these phenotypes are interrelated. We showed that CbrB was

required for increased PhoB activation in LasR– *P. aeruginosa* and a *crc* mutation was

sufficient to induce PhoB activity, even in a LasR+ strain. Importantly, the Δ*crc* mutant

had less AP activity and was inhibited by lower Pi concentrations than the Δ*lasR* mutant.

These data suggest Crc– *P. aeruginosa* do not phenocopy LasR– strains.

The elevated PhoB activity in the Δ*lasR* mutant contributed to fitness upon a shift

from LB to a medium without added Pi. As phosphate accessibility can be transient inside

the host, increased PhoB activity may be beneficial in building up stores of polyphosphate

for utilization when cells experience low Pi stress. In addition to regulating phosphate

acquisition, PhoB activity has also been shown to promote RhlR expression and

subsequent PYO production, swarming motility, and cytotoxicity.[13, 15, 42, 43, 90] Here we

showed that the Δ*lasR* mutant had PhoB-dependent increased expression of virulence-

associated genes including phenazine biosynthetic enzymes and phospholipases like

PlcH. PlcH is known to degrade airway surfactant and we have previously shown that its

expression leads to a decline in lung function.[95] We also showed PhoB and CbrB are

both required for increased production of the antifungal phenazine 5MPCA by the Δ*lasR*

mutant in co-culture with *C. albicans*.

The roles for phenazines in PhoB regulation are complex. As all QS mutants

produce fewer phenazines in late exponential phase and early stationary phase [83, 84], we

proposed phenazines may repress PhoB activity. We demonstrated that phenazine-

deficient *P. aeruginosa* had similarly elevated PhoB activity to the single QS mutants.

Unlike the Δ*rhlR* and Δ*pqsR* mutants, LasR– *P. aeruginosa* can still produce phenazines

in certain contexts.[13, 36-38] We observed that the Δ*lasR* mutant appeared greener on BCIP

agar at 0.7 mM Pi which we speculate is due to the production of the blue-green

phenazine PYO as this was not observed in the phenazine-deficient Δ*lasR*Δ*phz* mutant

(Fig. 3B). Thus, the higher AP activity in the Δ*lasR*Δ*phz* compared to the Δ*lasR* may be

due to the absence of phenazines that reduce PhoB activation at moderate Pi

concentrations. It is also of note that PhoB activity promotes phenazine production,

potentially through direct binding upstream of the *phz* operons [16], but more likely through

RhlR activity.[13, 42, 43] This increased phenazine production could then inhibit further PhoB

activity, providing negative feedback. However, our data suggests PCA, not PYO or

5MPCA, inhibits PhoB as loss of the biosynthetic enzymes PhzM or PhzS did not induce

PhoB activity. It cannot be assumed that cells producing visible PYO also have high PCA

and reduced PhoB activity. An example of this is the Δ*crc* mutant which we showed had

elevated AP activity and is known to produce more PYO through de-repression of

PhzM.[96] In some instances, increased PCA conversion to PYO could result in PCA

depletion and thus aid in the activation of PhoB. Further work is needed to fully

understand the interplay between PhoB activity and phenazines.

It is well established that in permissive Pi conditions, PhoB can promote the

expression QS genes directly [40] or through RhlR [15, 16, 43, 90], allowing cells to circumvent

reliance on LasR as a QS regulator.[13, 42] Our data support this existing model by showing

QS gene expression is PhoB-dependent in a Δ*lasR* mutant but not the wild type at 0.7

mM Pi. Importantly, we have shown increased PhoB activity in LasR– *P. aeruginosa*

compared to LasR+ strains and this has not been identified in previous publications. The

relatively low (0.4 – 0.5 mM) and high (4 – 4.5 mM) Pi concentrations used in the past

likely hindered the observation of increased PhoB activity in LasR– *P. aeruginosa* and

demonstrates the utility of gradient agar plates. As we found PhoB is active in the Δ*lasR*

mutant at higher Pi concentrations than the wild type, the previously reported PhoB-driven

reorganization of QS may occur in LasR– *P. aeruginosa* under conditions that otherwise

repress PhoB activity in LasR+ strains.

Further work is needed to elucidate the mechanisms by which LasR activity

represses PhoB activity at intermediate Pi concentrations. There is evidence that PhoB

can be spontaneously phosphorylated by acetyl phosphate in vitro [97] but the

concentrations required for this reaction suggest it is unlikely to occur inside cells.[98]

Therefore, we expect changes in PhoB activity are mediated by its sensor kinase PhoR.

PhoR activity is suppressed by interactions with the Pi transporter PstABC and PhoU.[18,

21, 23] Thus, changes in expression of these proteins could alter PhoR-PhoB activity. We

observed no differences in transcript abundance of *phoU* or *pstA* between the wild type

and Δ*lasR* mutant (Supplemental File 1) and so believe this model is unlikely. However,

other changes in Pi transport into the cell could still contribute to differences in PhoB

activity. While the mechanisms of PhoR-PhoU-PstABC interactions are not well

understood, PhoR has been shown to interact with PhoU through its Per-Arnt-Sim (PAS)

domain.[23] PAS domains in other organisms bind a broad range of ligands [99] and can

modify kinase activity.[100, 101] Thus, we propose that changes to the intracellular

environment in phenazine-deficient cells or cells with low Crc activity may stimulate PhoR

activity through its PAS domain. Future work into the mechanisms of the PhoR PAS

domain would also be beneficial for understanding similar kinases in *P. aeruginosa* and

other organisms as PAS domains are abundant but remain poorly understood.[102]

Understanding LasR– *P. aeruginosa* virulence in physiologically relevant conditions is

critical as LasR– isolates are frequently found in both acute and chronic infections and

these findings may be relevant to *P. aeruginosa* when in settings that do not induce QS

regulation. We propose that increased PhoB-mediated QS and virulence gene expression

in LasR– *P. aeruginosa* may contribute to their association with worse outcomes.[54, 57]

In conclusion, our data support a novel model where LasR represses PhoB activity

and thus LasR– *P. aeruginosa* have elevated PhoB activity at physiological Pi

concentrations. Others have shown that PhoB activity can induce RhlR activity in the

absence of LasR, leading to increased production of virulence factors like PYO.[13, 42] Here

we have shown QS and virulence-related gene expression in the Δ*lasR* mutant is highly

PhoB dependent at a physiologically relevant Pi concentrations.

Materials and Methods

Strains and growth conditions

Bacterial strains and plasmids used in this study are listed in **Table S1**. Bacteria were

streaked from frozen stocks onto LB (lysogeny broth) with 1.5% agar.[103] Planktonic

cultures were grown in 5 mL LB medium in 18 mm Borosilicate glass tubes on a roller

drum at 37 °C in the indicated medium.

Construction of in-frame deletion, complementation, and expression plasmids

Construction of in-frame deletion and complementation constructs was performed using

yeast cloning techniques in *Saccharomyces cerevisiae* as previously described [103] or

Gibson assembly.[104, 105] In-frame deletion and chromosomal complementation

constructs were made using the allelic replacement vector pMQ30.[106] Plasmids were

purified from yeast using Zymoprep Yeast Plasmid Miniprep II according to the

manufacturer's protocol and transformed into *E. coli* strain S17/λpir by electroporation.

Plasmids were introduced into *P. aeruginosa* by conjugation and recombinants were

obtained using sucrose counter-selection. Genotypes were screened by PCR and

plasmid constructs were confirmed by sequencing.

Alkaline phosphatase activity assessment

Agar plates were made as previously described [27] with MOPS minimal medium [74] with

0.7 mM K_2HPO_4/KH_2PO_4, 60 μg/mL BCIP (Sigma Aldrich # 11383221001), 20 mM

glucose and 15 g/L agar (referred to here as MOPS agar). Overnight cultures of *P.*

aeruginosa were grown in LB, normalized to an OD_{600} of 1, then 5 μL were spotted on

MOPS agar and incubated at 37 °C for 24 h. Plates with a gradient of phosphate were

made based on the methodology described for pH gradient plates [107]. First 35 mL of

molten MOPS agar at one end of the gradient was pipetted into a 10 cm square petri dish

(Corning, BP124-05) that rested in a custom 3D-printed prop that held the plate slanted

at a 30° angle. Once the bottom layer had solidified, the plate was laid flat and 35 mL of

molten medium agar containing the second desired concentration on the gradient was

poured atop. BCIP was added as described above.

Calculating alkaline phosphatase enzymatic activity

A colorimetric assay using the substrate p-Nitrophenyl phosphate (PNPP) (New England

Biolabs) was used to quantify alkaline phosphatase activity in colonies grown on MOPS

agar plates with no added BCIP using the inoculation regime described above. After 12-

24 h incubation, the colonies were scraped from the agar with a pipette tip and

resuspended in 1 mL 10 mM Tris-HCL buffer at pH 8. Fifty µL of the cell suspension was

mixed with 50 µL of room temperature 1-step PNPP solution or 50 µL of Tris-HCL buffer

at pH 8 and incubated for 1 h. A_{405} and A_{600} were recorded at t = 0 min and t = 60 min.

AP activity was calculated using the equation 1000 x (ΔA_{405}/(time (min) x A_{600})) where

$\Delta A_{405} = (A_{405}PNPP - A_{405}NoPNPP)_{t60} - (A_{405}PNPP - A_{405}NoPNPP)_{t0}$.

NanoString analysis

Total RNA was collected from *P. aeruginosa* colony biofilms grown on MOPS agar with

0.7 mM K_2HPO_4/KH_2PO_4. Bacteria were grown overnight in LB then sub-cultured in LB

until OD_{600} = 0.5 before 5 µL were spotted on MOPS agar and incubated at 37 °C for 12

h. Cells were then scraped from the agar and resuspended in 200 µL Tris-EDTA buffer.

RNA was extracted using the Qiagen RNeasy kit (Qiagen 74004) using the protocol

provided by the manufacturer. Samples were not treated with DNAse. NanoString

analysis of 80 ng of isolated RNA using codeset PaV5 [27] was performed as previously

reported [108]. Counts were normalized to the geometric mean of five housekeeping genes

(*ppiD*, *rpoD*, *soj*, *dnaN*, *pepP*, *dapF*). Normalized counts were used for heatmap

construction.

Growth curves

P. aeruginosa was grown in LB medium for 16 h and then sub-cultured to a starting OD_{600}

= 0.05 in 5 mL of MOPS minimal medium + 16.7 mM glucose in glass culture tubes. OD_{600}

was measured using a Spectronic 20D+.

Statistical analysis

Statistical analyses were performed using GraphPad Prism version 10.1.1 for macOS.

Acknowledgments

The research reported in this publication was supported by grants from the Cystic Fibrosis

Foundation, GREEN19G0 and STANTO19R0, and the National Institutes of Health (NIH),

NHLBI T32HL134598 (A.C.) and NIDDK P30-DK117469 (Dartmouth Cystic Fibrosis

Research Center or DartCF). Additional support came from bioMT (NIGMS

P20GM113132) and the Dartmouth Molecular Biology Shared Resource (NCI

5P30CA023108).

Citations

1. Bansal VK. 1990. Clinical Methods: The history, physical, and laboratory examinations. Serum inorganic phosphorous. *In* HK W, HD, Hall, JW, Hurst (ed), Clinical Methods: The History, Physical, and Laboratory Examinations, 3 ed. Butterworths, Boston.
2. Oh TK, Jo J, Oh AY. 2018. Perioperative serum calcium and phosphorus levels are associated with hospital costs and length of stay after major abdominal surgery. J Clin Med 7.
3. Salem RR, Tray K. 2005. Hepatic resection-related hypophosphatemia is of renal origin as manifested by isolated hyperphosphaturia. Ann Surg 241:343-8.
4. Long J, Zaborina O, Holbrook C, Zaborin A, Alverdy J. 2008. Depletion of intestinal phosphate after operative injury activates the virulence of *P. aeruginosa* causing lethal gut-derived sepsis. Surgery 144:189-97.
5. Riedler GF, Scheitlin WA. 1969. Hypophosphataemia in septicaemia: higher incidence in gram-negative than in gram-positive infections. Br Med J 1:753-6.
6. Berner YN, Shike M. 1988. Consequences of phosphate imbalance. Annual review of nutrition 8:121-148.
7. Bruna RE, Kendra CG, Pontes MH. 2023. Phosphorus starvation response and PhoB-independent utilization of organic phosphate sources by *Salmonella enterica*. Microbiology Spectrum 11:e02260-23.
8. Filloux A, Bally M, Soscia C, Murgier M, Lazdunski A. 1988. Phosphate regulation in *Pseudomonas aeruginosa*: Cloning of the alkaline phosphatase gene and identification of *phoB*- and *phoR*-like genes. Molecular and General Genetics 212:510-513.
9. Monds RD, Newell PD, Schwartzman JA, O'Toole GA. 2006. Conservation of the Pho regulon in *Pseudomonas fluorescens* Pf0-1. Applied and Environmental Microbiology 72:1910-1924.
10. Lamarche MG, Wanner BL, Crépin S, Harel J. 2008. The phosphate regulon and bacterial virulence: a regulatory network connecting phosphate homeostasis and pathogenesis. FEMS Microbiology Reviews 32:461-473.
11. Gallarato LA, Sanchez DG, Olvera L, Primo ED, Garrido MN, Beassoni PR, Morett E, Lisa AT. 2014. Exopolyphosphatase of *Pseudomonas aeruginosa* is essential for the production of virulence factors, and its expression is controlled by NtrC and PhoB acting at two interspaced promoters. Microbiology 160:406-417.
12. Matilla MA, Udaondo Z, Maaß S, Becher D, Krell T. 2022. Virulence induction in *Pseudomonas aeruginosa* under inorganic phosphate limitation: a proteomics oerspective. Microbiology Spectrum 10:e0259022.
13. Soto-Aceves MP, Cocotl-Yañez M, Servín-González L, Soberón-Chávez G. 2021. The Rhl quorum-sensing system is at the top of the regulatory hierarchy under phosphate-limiting conditions in *Pseudomonas aeruginosa* PAO1. Journal of Bacteriology 203:e00475-20.
14. Bains M, Fernandez L, Hancock RE. 2012. Phosphate starvation promotes swarming motility and cytotoxicity of *Pseudomonas aeruginosa*. Appl Environ Microbiol 78:6762-8.

15. Blus-Kadosh I, Zilka A, Yerushalmi G, Banin E. 2013. The effect of *pstS* and *phoB* on quorum sensing and swarming motility in *Pseudomonas aeruginosa*. PloS one 8:e74444.

16. Jensen V, Löns D, Zaoui C, Bredenbruch F, Meissner A, Dieterich G, Münch R, Häussler S. 2006. RhlR expression in *Pseudomonas aeruginosa* is modulated by the *Pseudomonas* quinolone signal via PhoB-dependent and-independent pathways. Journal of Bacteriology 188:8601-8606.

17. McRose DL, Newman DK. 2021. Redox-active antibiotics enhance phosphorus bioavailability. Science 371:1033-1037.

18. Muda M, Rao N, Torriani A. 1992. Role of PhoU in phosphate transport and alkaline phosphatase regulation. Journal of Bacteriology 174:8057-8064.

19. Rice CD, Pollard JE, Lewis ZT, McCleary WR. 2009. Employment of a promoter-swapping technique shows that PhoU modulates the activity of the PstSCAB2 ABC transporter in *Escherichia coli*. Applied and environmental microbiology 75:573-582.

20. Lee SJ, Park YS, Kim SJ, Lee BJ, Suh SW. 2014. Crystal structure of PhoU from *Pseudomonas aeruginosa*, a negative regulator of the Pho regulon. Journal of Structural Biology 188:22-9.

21. de Almeida LG, Ortiz JH, Schneider RP, Spira B. 2015. *phoU* inactivation in *Pseudomonas aeruginosa* enhances accumulation of ppGpp and polyphosphate. Applied and environmental microbiology 81:3006-3015.

22. Munévar NFV, de Almeida LG, Spira B. 2017. Differential regulation of polyphosphate genes in *Pseudomonas aeruginosa*. Molecular Genetics and Genomics 292:105-116.

23. Gardner SG, Johns KD, Tanner R, McCleary WR. 2014. The PhoU protein from *Escherichia coli* interacts with PhoR, PstB, and metals to form a phosphate-signaling complex at the membrane. Journal of Bacteriology 196:1741-1752.

24. Wanner BL. 1996. Phosphorus assimilation and control of the phosphate regulon. *Escherichia coli* and *Salmonella*: cellular and molecular biology 1:1357-1381.

25. Nikata T, Sakai Y, Shibata K, Kato J, Kuroda A, Ohtake H. 1996. Molecular analysis of the phosphate-specific transport (*pst*) operon of *Pseudomonas aeruginosa*. Molecular and General Genetics MGG 250:692-698.

26. Anba J, Bidaud M, Vasil ML, Lazdunski A. 1990. Nucleotide sequence of the *Pseudomonas aeruginosa phoB* gene, the regulatory gene for the phosphate regulon. J Bacteriol 172:4685-9.

27. Doing G, Koeppen K, Occipinti P, Harty CE, Hogan DA. 2020. Conditional antagonism in co-cultures of *Pseudomonas aeruginosa* and *Candida albicans*: an intersection of ethanol and phosphate signaling distilled from dual-seq transcriptomics. PLoS genetics 16:e1008783.

28. Gao R, Stock AM. 2018. Overcoming the cost of positive autoregulation by accelerating the response with a coupled negative feedback. Cell Reports 24:3061-3071.e6.

29. Makino K, Shinagawa H, Amemura M, Kawamoto T, Yamada M, Nakata A. 1989. Signal transduction in the phosphate regulon of *Escherichia coli* involves phosphotransfer between PhoR and PhoB proteins. Journal of Molecular Biology 210:551-559.

30.   Pratt JT, Ismail AM, Camilli A. 2010. PhoB regulates both environmental and
virulence gene expression in *Vibrio cholerae*. Molecular microbiology 77:1595-
1605.
31.   Chekabab SM, Jubelin G, Dozois CM, Harel J. 2014. PhoB activates *Escherichia*
coli O157:H7 virulence factors in response to inorganic phosphate limitation. PLoS
One 9:e94285.
32.   Higgins DE, DiRita VJ. 1994. Transcriptional control of toxT, a regulatory gene in
the ToxR regulon of *Vibrio cholerae*. Mol Microbiol 14:17-29.
33.   Antunes LCM, Ferreira RBR, Buckner MMC, Finlay BB. 2010. Quorum sensing in
bacterial virulence. Microbiology 156:2271-2282.
34.   Chen G, Lim ED, Winkelman BT, Winkelman JT, Mukherjee S. 2022.
Combinatorial control of biofilm development by quorum-sensing and nutrient-
sensing regulators in *Pseudomonas aeruginosa*. bioRxiv
doi:10.1101/2022.09.27.509822:2022.09.27.509822.
35.   Hentzer M, Wu H, Andersen JB, Riedel K, Rasmussen TB, Bagge N, Kumar N,
Schembri MA, Song Z, Kristoffersen P, Manefield M, Costerton JW, Molin S, Eberl
L, Steinberg P, Kjelleberg S, Høiby N, Givskov M. 2003. Attenuation of
Pseudomonas aeruginosa virulence by quorum sensing inhibitors. The EMBO
Journal 22:3803-3815.
36.   Feltner JB, Wolter DJ, Pope CE, Groleau MC, Smalley NE, Greenberg EP, Mayer-
Hamblett N, Burns J, Déziel E, Hoffman LR, Dandekar AA. 2016. LasR variant
cystic fibrosis isolates reveal an adaptable quorum-sensing hierarchy in
Pseudomonas aeruginosa. mBio 7:e01513-16.
37.   Mould DL, Botelho NJ, Hogan DA, Goldberg JB. 2020. Intraspecies signaling
between common variants of *Pseudomonas aeruginosa* increases production of
quorum-sensing-controlled virulence factors. mBio 11:e01865-20.
38.   Kostylev M, Kim DY, Smalley NE, Salukhe I, Greenberg EP, Dandekar AA. 2019.
Evolution of the *Pseudomonas aeruginosa* quorum-sensing hierarchy.
Proceedings of the National Academy of Sciences 116:7027-7032.
39.   Cugini C, Morales DK, Hogan DA. 2010. *Candida* albicans-produced farnesol
stimulates *Pseudomonas* quinolone signal production in LasR-defective
Pseudomonas aeruginosa strains. Microbiology 156:3096.
40.   Meng X, Ahator SD, Zhang L-H. 2020. Molecular mechanisms of phosphate stress
activation of *Pseudomonas aeruginosa* quorum sensing systems. MSphere
5:e00119-20.
41.   Zaborin A, Gerdes S, Holbrook C, Liu DC, Zaborina OY, Alverdy JC. 2012.
Pseudomonas aeruginosa overrides the virulence inducing effect of opioids when
it senses an abundance of phosphate. PLoS One 7:e34883.
42.   Welsh MA, Blackwell HE. 2016. Chemical genetics reveals environment-specific
roles for quorum sensing circuits in *Pseudomonas aeruginosa*. Cell Chem Biol
23:361-9.
43.   Mellbye B, Schuster M. 2014. Physiological framework for the regulation of quorum
sensing-dependent public goods in *Pseudomonas aeruginosa*. Journal of
Bacteriology 196:1155-1164.
44.   Nutman J, Berger M, Chase PA, Dearborn DG, Miller K, Waller R, Sorensen R.
1987. Studies on the mechanism of T cell inhibition by the *Pseudomonas*

aeruginosa phenazine pigment pyocyanine. Journal of immunology (Baltimore, Md: 1950) 138:3481-3487.

45. Rada B, Leto TL. 2013. Pyocyanin effects on respiratory epithelium: relevance in *Pseudomonas aeruginosa* airway infections. Trends in microbiology 21:73-81.

46. Gibson J, Sood A, Hogan DA. 2009. *Pseudomonas aeruginosa-Candida albicans* interactions: localization and fungal toxicity of a phenazine derivative. Applied and environmental microbiology 75:504-513.

47. Morales DK, Jacobs NJ, Rajamani S, Krishnamurthy M, Cubillos-Ruiz JR, Hogan DA. 2010. Antifungal mechanisms by which a novel *Pseudomonas aeruginosa* phenazine toxin kills *Candida albicans* in biofilms. Mol Microbiol 78:1379-92.

48. Morales DK, Grahl N, Okegbe C, Dietrich LE, Jacobs NJ, Hogan DA. 2013. Control of *Candida albicans* metabolism and biofilm formation by *Pseudomonas aeruginosa* phenazines. MBio 4:e00526-12.

49. Zhang L, Tian X, Kuang S, Liu G, Zhang C, Sun C. 2017. Antagonistic activity and mode of action of phenazine-1-carboxylic acid, produced by marine bacterium *Pseudomonas aeruginosa* PA31x, against *Vibrio anguillarum in vitro* and in a zebrafish *in vivo* model. Frontiers in microbiology 8:289.

50. Truong-Bolduc QC, Yonker LM, Wang Y, Lawton BG, Hooper DC. 2024. NorA efflux pump mediates S*taphylococcus aureus* response to *Pseudomonas aeruginosa* pyocyanin toxicity. Antimicrobial Agents and Chemotherapy 68:e01001-23.

51. Kahl LJ, Stremmel N, Esparza-Mora MA, Wheatley RM, MacLean RC, Ralser M. 2023. Interkingdom interactions between *Pseudomonas aeruginosa* and *Candida albicans* affect clinical outcomes and antimicrobial responses. Current Opinion in Microbiology 75:102368.

52. Groleau M-C, Taillefer H, Vincent AT, Constant P, Déziel E. 2022. *Pseudomonas aeruginosa* isolates defective in function of the LasR quorum sensing regulator are frequent in diverse environmental niches. Environmental Microbiology 24:1062-1075.

53. O'Connor K, Zhao CY, Mei M, Diggle SP. 2022. Frequency of quorum-sensing mutations in *Pseudomonas aeruginosa* strains isolated from different environments. Microbiology (Reading) 168.

54. Hammond JH, Hebert WP, Naimie A, Ray K, Van Gelder RD, DiGiandomenico A, Lalitha P, Srinivasan M, Acharya NR, Lietman T, Hogan DA, Zegans ME. 2016. Environmentally endemic *Pseudomonas aeruginosa* strains with mutations in *lasR* are associated with increased disease severity in corneal ulcers. mSphere 1:00140-16.

55. Trottier MC, Pereira TdO, Groleau M-C, Hoffman LR, Dandekar AA, Déziel E. 2024. The end of the reign of a "master regulator"? A defect in function of the LasR quorum sensing regulator is a common feature of *Pseudomonas aeruginosa* isolates. mBio 15:e02376-23.

56. Wilder CN, Allada G, Schuster M. 2009. Instantaneous within-patient diversity of *Pseudomonas aeruginosa* quorum-sensing populations from cystic fibrosis lung infections. Infect Immun 77:5631-9.

57. Hoffman LR, Kulasekara HD, Emerson J, Houston LS, Burns JL, Ramsey BW,
Miller SI. 2009. *Pseudomonas aeruginosa lasR* mutants are associated with cystic
fibrosis lung disease progression. Journal of Cystic Fibrosis 8:66-70.
58. Mould DL, Stevanovic M, Ashare A, Schultz D, Hogan DA. 2022. Metabolic basis
for the evolution of a common pathogenic *Pseudomonas aeruginosa* variant. eLife
11:e76555.
59. Mould DL, Finger CE, Conaway A, Botelho N, Stuut Stacie E, Hogan Deborah A.
2024. Citrate cross-feeding between *Pseudomonas aeruginosa* genotypes
supports *lasR* mutant fitness. mbio doi:01278-23:e01278-23.
60. D'Argenio DA, Wu M, Hoffman LR, Kulasekara HD, Déziel E, Smith EE, Nguyen
H, Ernst RK, Larson Freeman TJ, Spencer DH. 2007. Growth phenotypes of
Pseudomonas aeruginosa lasR mutants adapted to the airways of cystic fibrosis
patients. Molecular microbiology 64:512-533.
61. Yeung ATY, Bains M, Hancock REW. 2011. The Sensor Kinase CbrA Is a Global
Regulator That Modulates Metabolism, Virulence, and Antibiotic Resistance in
Pseudomonas aeruginosa. Journal of Bacteriology 193:918-931.
62. García-Mauriño SM, Pérez-Martínez I, Amador CI, Canosa I, Santero E. 2013.
Transcriptional activation of the CrcZ and CrcY regulatory RNAs by the CbrB
response regulator in *Pseudomonas putida*. Molecular microbiology 89:189-205.
63. Valentini M, García-Mauriño SM, Pérez-Martínez I, Santero E, Canosa I, Lapouge
K. 2014. Hierarchical management of carbon sources is regulated similarly by the
CbrA/B systems in *Pseudomonas aeruginosa* and *Pseudomonas putida*.
Microbiology 160:2243-2252.
64. Sonnleitner E, Abdou L, Haas D. 2009. Small RNA as global regulator of carbon
catabolite repression in *Pseudomonas aeruginosa*. Proceedings of the National
Academy of Sciences 106:21866-21871.
65. Sonnleitner E, Valentini M, Wenner N, Haichar FeZ, Haas D, Lapouge K. 2012.
Novel targets of the CbrAB/Crc carbon catabolite control system revealed by
transcript abundance in *Pseudomonas aeruginosa*. PloS one 7:e44637.
66. Yeung ATY, Janot L, Pena OM, Neidig A, Kukavica-Ibrulj I, Hilchie A, Levesque
RC, Overhage J, Hancock REW. 2014. Requirement of the *Pseudomonas
aeruginosa* CbrA sensor kinase for full virulence in a murine acute lung infection
model. Infection and Immunity 82:1256-1267.
67. Palmer KL, Aye LM, Whiteley M. 2007. Nutritional cues control *Pseudomonas
aeruginosa* multicellular behavior in cystic fibrosis sputum. Journal of Bacteriology
189:8079-8087.
68. Moore GE, Gerner RE, Franklin HA. 1967. Culture of normal human leukocytes.
JAMA 199:519-524.
69. Mössinger J. 1991. In vitro cultivation of adult *Litomosoides carinii*: evaluation of
basic culture media, gas phases and supplements. Parasitology 103:85-95.
70. Schurig-Briccio LA, Rintoul MR, Volentini SI, Farías RN, Baldomà L, Badía J,
Rodríguez-Montelongo L, Rapisarda VA. 2008. A critical phosphate concentration
in the stationary phase maintains ndh gene expression and aerobic respiratory
chain activity in *Escherichia coli*. FEMS Microbiology Letters 284:76-83.
71. Hou CI, Gronlund AF, Campbell JJR. 1966. Influence of phosphate starvation on
cultures of *Pseudomonas aeruginosa*. Journal of Bacteriology 92:851-855.

72. Xin L, Deliang L, Heng Y, Dingpeng Y, Shuang Z, Shuting Z, Mengqiu L, Miao H,
Min X, Xingguo W. 2016. Phosphatidylcholine affects the secretion of the alkaline
phosphatase PhoA in *Pseudomonas* strains. Microbiological Research 192:21-29.
73. Horwitz JP, Chua J, Noel M, Donatti JT, Freisler J. 1966. Substrates for
cytochemical demonstration of enzyme activity. II. Some Dihalo-3-indolyl
Phosphates and Sulfates1. Journal of Medicinal Chemistry 9:447-447.
74. Neidhardt FC, Bloch PL, Smith DF. 1974. Culture medium for *enterobacteria*.
Journal of Bacteriology 119:736-747.
75. Wanner B. 1993. Gene regulation by phosphate in enteric bacteria. Journal of
Cellular Biochemistry 51:47-54.
76. Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, D'Argenio DA,
Miller SI, Ramsey BW, Speert DP, Moskowitz SM, Burns JL, Kaul R, Olson MV.
2006. Genetic adaptation by *Pseudomonas aeruginosa* to the airways of cystic
fibrosis patients. Proceedings of the National Academy of Sciences 103:8487-
8492.
77. Gellatly SL, Bains M, Breidenstein EBM, Strehmel J, Reffuveille F, Taylor PK,
Yeung ATY, Overhage J, Hancock REW. 2018. Novel roles for two-component
regulatory systems in cytotoxicity and virulence-related properties in
Pseudomonas aeruginosa. AIMS Microbiology 4:173-191.
78. Ball G, Durand E, Lazdunski A, Filloux A. 2002. A novel type II secretion system
in *Pseudomonas aeruginosa*. Mol Microbiol 43:475-85.
79. Otero-Asman JR, Quesada JM, Jim KK, Ocampo-Sosa A, Civantos C, Bitter W,
Llamas MA. 2020. The extracytoplasmic function sigma factor σVreI is active
during infection and contributes to phosphate starvation-induced virulence of
Pseudomonas aeruginosa. Scientific Reports 10:3139.
80. Lewenza S, Falsafi R, Bains M, Rohs P, Stupak J, Sprott GD, Hancock RE. 2011.
The olsA gene mediates the synthesis of an ornithine lipid in *Pseudomonas*
aeruginosa during growth under phosphate-limiting conditions, but is not involved
in antimicrobial peptide susceptibility. FEMS Microbiology Letters 320:95-102.
81. Wu H, Kato J, Kuroda A, Ikeda T, Takiguchi N, Ohtake H. 2000. Identification and
characterization of two chemotactic transducers for inorganic phosphate in
Pseudomonas aeruginosa. Journal of Bacteriology 182:3400-3404.
82. Dekimpe V, Déziel E. 2009. Revisiting the quorum-sensing hierarchy in
Pseudomonas aeruginosa: the transcriptional regulator RhlR regulates LasR-
specific factors. Microbiology 155:712-723.
83. Higgins S, Heeb S, Rampioni G, Fletcher MP, Williams P, Cámara M. 2018.
Differential regulation of the phenazine biosynthetic operons by quorum sensing in
Pseudomonas aeruginosa PAO1-N. Frontiers in cellular and infection microbiology
8:252.
84. Recinos DA, Sekedat MD, Hernandez A, Cohen TS, Sakhtah H, Prince AS, Price-
Whelan A, Dietrich LE. 2012. Redundant phenazine operons in *Pseudomonas*
aeruginosa exhibit environment-dependent expression and differential roles in
pathogenicity. Proceedings of the National Academy of Sciences 109:19420-
19425.

85. Cui Q, Lv H, Qi Z, Jiang B, Xiao B, Liu L, Ge Y, Hu X. 2016. Cross-Regulation
between the *phz1* and *phz2* operons maintain a balanced level of phenazine
biosynthesis in *Pseudomonas aeruginosa* PAO1. PLOS ONE 11:e0144447.

86. Schmitz S, Rosenbaum MA. 2020. Controlling the production of *Pseudomonas*
phenazines by modulating the genetic repertoire. ACS Chemical Biology 15:3244-
3252.

87. Smyth P, Clarke PH. 1975. Catabolite repression of *Pseudomonas aeruginosa*
amidase: the effect of carbon source on amidase synthesis. Microbiology 90:81-
90.

88. Petrova OE, Schurr JR, Schurr MJ, Sauer K. 2011. The novel *Pseudomonas*
aeruginosa two-component regulator BfmR controls bacteriophage-mediated lysis
and DNA release during biofilm development through PhdA. Molecular
microbiology 81:767-783.

89. Shortridge VD, Lazdunski A, Vasil ML. 1992. Osmoprotectants and phosphate
regulate expression of phospholipase C in *Pseudomonas aeruginosa*. Mol
Microbiol 6:863-71.

90. Bains M, Fernández L, Hancock RE. 2012. Phosphate starvation promotes
swarming motility and cytotoxicity of *Pseudomonas aeruginosa*. Applied and
environmental microbiology 78:6762-6768.

91. Bielecki P, Jensen V, Schulze W, Gödeke J, Strehmel J, Eckweiler D, Nicolai T,
Bielecka A, Wille T, Gerlach RG. 2015. Cross talk between the response regulators
PhoB and TctD allows for the integration of diverse environmental signals in
Pseudomonas aeruginosa. Nucleic acids research 43:6413-6425.

92. Faure LM, Llamas MA, Bastiaansen KC, De Bentzmann S, Bigot S. 2013.
Phosphate starvation relayed by PhoB activates the expression of the
Pseudomonas aeruginosa σvreI ECF factor and its target genes. Microbiology
159:1315-1327.

93. Heurlier K, Dénervaud V, Haenni M, Guy L, Krishnapillai V, Haas D. 2005.
Quorum-sensing-negative *lasR* mutants of *Pseudomonas aeruginosa* avoid cell
lysis and death. Journal of Bacteriology 187:4875-4883.

94. Clay ME, Hammond JH, Zhong F, Chen X, Kowalski CH, Lee AJ, Porter MS,
Hampton TH, Greene CS, Pletneva EV. 2020. *Pseudomonas aeruginosa lasR*
mutant fitness in microoxia is supported by an Anr-regulated oxygen-binding
hemerythrin. Proceedings of the National Academy of Sciences 117:3167-3173.

95. Wargo MJ, Gross MJ, Rajamani S, Allard JL, Lundblad LKA, Allen GB, Vasil ML,
Leclair LW, Hogan DA. 2011. Hemolytic phospholipase C inhibition protects lung
function during *Pseudomonas aeruginosa* infection. American Journal of
Respiratory and Critical Care Medicine 184:345-354.

96. Huang J, Sonnleitner E, Ren B, Xu Y, Haas D. 2012. Catabolite repression control
of pyocyanin biosynthesis at an intersection of primary and secondary metabolism
in *Pseudomonas aeruginosa*. Applied and environmental microbiology 78:5016-
5020.

97. Hiratsu K, Nakata A, Shinagawa H, Makino K. 1995. Autophosphorylation and
activation of transcriptional activator PhoB of *Escherichia coli* by acetyl phosphate
in vitro. Gene 161:7-10.

98. McCleary W. 1996. The activation of PhoB by acetylphosphate. Molecular
Microbiology 20:1155-1163.
99. Henry JT, Crosson S. 2011. Ligand-Binding PAS domains in a genomic, cellular,
and structural context. Annual Review of Microbiology 65:261-286.
100. Qi Y, Rao F, Luo Z, Liang ZX. 2009. A flavin cofactor-binding PAS domain
regulates c-di-GMP synthesis in AxDGC2 from *Acetobacter xylinum*. Biochemistry
48:10275-85.
101. Dupré E, Clantin B, Yuan Y, Lecher S, Lesne E, Antoine R, Villeret V, Jacob-
Dubuisson F. 2021. Structural insight into the role of the PAS domain for signal
transduction in sensor kinase BvgS. Journal of Bacteriology 203:e00614-20.
102. Xing J, Gumerov VM, Zhulin IB. 2023. Origin and functional diversification of PAS
domain, a ubiquitous intracellular sensor. Science Advances 9:eadi4517.
103. Bertani G. 1951. Studies on lysogenesis. I. The mode of phage liberation by
lysogenic *Escherichia coli*. J Bacteriol 62:293-300.
104. Gibson DG, Glass JI, Lartigue C, Noskov VN, Chuang RY, Algire MA, Benders
GA, Montague MG, Ma L, Moodie MM, Merryman C, Vashee S, Krishnakumar R,
Assad-Garcia N, Andrews-Pfannkoch C, Denisova EA, Young L, Qi ZQ, Segall-
Shapiro TH, Calvey CH, Parmar PP, Hutchison CA, 3rd, Smith HO, Venter JC.
2010. Creation of a bacterial cell controlled by a chemically synthesized genome.
Science 329:52-6.
105. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. 2009.
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nature
Methods 6:343-5.
106. Shanks RM, Caiazza NC, Hinsa SM, Toutain CM, O'Toole GA. 2006.
Saccharomyces cerevisiae-based molecular tool kit for manipulation of genes from
gram-negative bacteria. Applied Environmental Microbiology 72:5027-36.
107. Sacks L. 1956. A pH gradient agar plate. Nature 178:269-270.
108. Grahl N, Dolben EL, Filkins LM, Crocker AW, Willger SD, Morrison HG, Sogin ML,
Ashare A, Gifford AH, Jacobs NJ, Schwartzman JD, Hogan DA. 2018. Profiling of
bacterial and fungal microbial communities in cystic fibrosis sputum using RNA.
mSphere 3:00292-18.
765

Fig. 1 *P. aeruginosa* **LasR– strains require higher Pi to repress PhoB activity.** **A)** *P. aeruginosa* wild type (WT), Δ*lasR*, Δ*phoB* and Δ*pstB* were spotted on gradient plates of MOPS agar with a range of Pi concentrations (0.1 - 1 mM) and 60 µg/mL BCIP to indicate alkaline phosphatase (AP) activity. **B)** *P. aeruginosa* strains, including DH2590 (a LasR– clinical isolate (CI) and a derivative in which the *lasR* allele was replaced with PA14 *lasR*) on MOPS agar with BCIP and 0.7 mM Pi. **C)** *P. aeruginosa* CI LasR+/LasR– pairs from the sputum of two pwCF on MOPS agar with BCIP and 0.7 mM Pi. **D)** Colony biofilms on MOPS agar with 0.2% glucose and either 0.2 or 10 mM Pi and BCIP. For **A-D**, similar results were obtained in three replicate experiments; a representative experiment is shown. For panels **E-G**, *P. aeruginosa* was grown as colony biofilms on MOPS agar with 0.7 mM Pi. **E)** AP activity in WT and Δ*lasR* after 12 h at 37 °C. Data from replicates collected on the same day have the same shape. Data analyzed using a paired, two-tailed t-test (n = 12). **F)** *phoA* transcripts in WT and Δ*lasR* were measured by qRT-PCR on different days and normalized to the housekeeping gene transcript *ppiD*. Data analyzed using a paired, two-tailed t-test (n = 8). **G)** Levels of PhoB-controlled transcripts and *phoB* itself were assessed using NanoString multiplex technology. Nine PhoB-regulated transcripts are shown as normalized counts. Data were analyzed using a two-way ANOVA; significant differences between WT-Δ*lasR,* WT-Δ*phoB*, and Δ*lasR*-Δ*lasR*Δ*phoB* (p < 0.0001, n = 2-3). There were no significant differences between the Δ*phoB*-Δ*lasR*Δ*phoB* mutants (p = 0.46, n = 3). Statistical differences for each transcript are available in Supplemental File 1. For all panels, asterisks denote significance (p ≤ 0.01 = **, p ≤ 0.001 = ***, p ≤ 0.0001 = ****).

Fig. 2 *P. aeruginosa* **quorum sensing (QS) mutants have active PhoB at higher Pi than the wild type. A)** Colony biofilms of wild type (WT) and the indicated QS mutants (Δ*lasR*, Δ*rhlR*, Δ*pqsR*) were grown on MOPS agar with 0.7 mM Pi and BCIP (top) or on medium without BCIP for analysis of AP activity using the colorimetric PNPP substrate. Data from replicates collected on the same day have the same shape. Data analyzed by an ordinary one-way ANOVA with Tukey's multiple comparisons tests (n = 4). **B)** *P. aeruginosa* was grown on a plate with a gradient of Pi concentrations, with 0.5 to 1.2 mM shown. Quantitative data analyzed using a one-way ANOVA (n = 3) (asterisks denote significance from the wild type). For all panels, asterisks denote significance (p ≤ 0.05 = *, p ≤ 0.01 = **, p ≤ 0.001 = ***, p ≤ 0.0001 = ****).

Fig. 3 The loss of phenazines promotes PhoB activity. A) AP activity in wild type (WT) and mutants Δ*phz1*, Δ*phz2*, and Δ*phz* (lacking *phz1* and *phz2* operons) grown on MOPS agar with 0.7 mM Pi and BCIP (top) or on medium without BCIP for analysis of AP activity using the colorimetric PNPP substrate. Data from replicates collected on the same day have the same shape. Data were analyzed using a one-way ANOVA and Tukey's multiple comparisons test (n = 7). **B)** Analysis of WT and the Δ*lasR*, Δ*phz*, and Δ*lasR*Δ*phz* mutants. The Δ*lasR*Δ*phz* mutant had significantly more AP activity than the Δ*lasR* and Δ*phz* strains. There are no significant differences between Δ*lasR* and Δ*phz* (p=0.99) (n = 12). **C)** *P. aeruginosa* was grown on plates with a gradient of Pi; the average concentration of Pi that inhibits AP activity is graphed to the right (n = 3). Data analyzed using a one-way ANOVA; asterisks denote significance from the wild type. Asterisks denote significance (p ≤ 0.05 = *, p ≤ 0.01 = **, p ≤ 0.001 = ***, p ≤ 0.0001 = ****).

Fig. 4. The CbrA-CbrB-Crc pathway promotes PhoR-PhoB activity in LasR– strains. A) A proposed model of the relationship between the CbrA-CbrB and PhoR-PhoB two component systems; *crcZ*, a small RNA; Crc, which acts as a translational repressor in complex with Hfq. For figures **B** and **C**, AP activity in indicated strains grown on MOPS agar with 0.7 mM Pi and BCIP (top) or on medium without BCIP for analysis of AP activity using the colorimetric PNPP substrate. Data from replicates collected on the same day have the same shape. Data were analyzed using a one-way ANOVA and Tukey's multiple comparisons test. **B)** AP activity in ΔlasR, ΔlasRΔcbrB, ΔlasRΔcbrB+cbrB, and ΔlasRΔcbrB+crc mutants (n = 3). **C)** AP activity in the wild type (WT), Δcrc mutant and its complemented derivative (n = 5). **D)** AP activity in ΔlasR, Δcrc, and ΔlasRΔcrc (n = 8). **E)** *P. aeruginosa* was grown on plates with a gradient of Pi (0.5 – 1.5 mM Pi) and BCIP. The average concentration of Pi which inhibited AP activity is graphed to the right (n = 3). Asterisks denote significance ($p \leq 0.05$ = *, $p \leq 0.01$ = **, $p \leq 0.001$ = ***).

Fig. 5 PhoB activity mediates fitness advantages and antagonism in Δ*lasR P. aeruginosa*.
A) *P. aeruginosa* wild type (WT), Δ*lasR*, Δ*phoB*, and Δ*lasR*Δ*phoB* were grown in MOPS liquid medium with no added Pi at 37 °C. Data analyzed using a two-way ANOVA with Tukey's multiple comparisons test (n = 3). Asterisks denote significance (p ≤ 0.0001 = ****). **B)** *P. aeruginosa* colonies were grown on a lawn of *Candida albicans*. Wild-type (LasR+) *P. aeruginosa* produces 5MPCA which is converted to red pigment inside *C. albicans*. The Δ*lasR* mutant produced more 5MPCA than the wild type. 5MPCA production required PhoB and CbrB in both strains (n = 3). **C)** NanoString analysis of PhoB-regulated transcripts shown as Log$_{10}$ normalized counts (n = 2 - 3). All transcripts shown were significantly higher in the Δ*lasR* mutant compared to both the wild type (p ≤ 0.0001) and the Δ*lasR*Δ*phoB* mutant (p ≤ 0.0016). **D)** PhoB contribution to regulation of QS genes by NanoString analysis of QS-regulated transcripts between the WT or Δ*phoB* mutant (p > 0.30) and the Δ*lasR* and Δ*lasR*Δ*phoB*. All transcripts were significantly lower in the Δ*lasR*Δ*phoB* mutant than the Δ*lasR* mutant (p ≤ 0.01). *lasB, rhlR, pqsR,* and *pqsH* were significantly lower in the Δ*lasR* mutant than the wild type (p < 0.0001). Complete NanoString dataset available in Supplemental File 1.