Reliable estimates of Japanese encephalitis virus evolutionary rate supported

by a formal test of temporal signal

Polina Ukraintceva [a], Artem Bondaryuk [b,*], Rodion Sakhabeev [a]

[a] Saint-Petersburg State Institute of Technology, Moskovski ave. 26, 190013, St.

Petersburg, Russia;

[b] Irkutsk Antiplague Research Institute of Siberia and the Far East, Trilisser 78,

664047, Irkutsk, Russia.

[*] Correspondence at: Artem Bondaryuk. E-mail: ui.artem.ui@gmail.com

Polina Ukraintceva, polina.ukraintseva@bk.ru; ORCID: https://orcid.org/0000-0002-3242-1526;

Artem Bondaryuk, ui.artem.ui@gmail.com; ORCID: https://orcid.org/0000-0003-4422-0497;

Rodion Sakhabeev, helm505@mail.ru; ORCID: https://orcid.org/0000-0002-4367-4924.

Abstract

The single-stranded RNA viruses exhibit stupendously high evolutionary

rate spanned several orders of magnitude. The accurate and reliable

estimates of the virus evolutionary rate are of importance for the robust

interpretation of outbreak investigations and are critical for understanding

short-term virus transmission patterns. In the previous studies, the

substitution rate of the Japanese encephalitis virus (JEV) and time to the

most recent common ancestor (MRCA) were reported, however, the

estimates are substantially different between some of the studies.

Importantly, the temporal signal in data was not evaluated by a formal test.

The purpose of our study is to estimate the substitution rate and the age of

the JEV common clade (GI-GV) and two JEV genotypes (GI and GIII)

supported by the formal Bayesian analysis of temporal signal in the open

reading frame (ORF) nucleotide data. Additionally, we assessed temporal

signal in the data from the previous work to explain the observed

discrepancy between the reported JEV evolutionary rate estimates. Results

showed that the substitution rate of the JEV GI-GV was 2.41×10^{-4}

nucleotide substitution per site per year (s/s/y) with 95% highest posterior density (HPD) interval of 1.67×10^{-4}–3.14×10^{-4} or approximately 2.5 nucleotide substitutions per ORF (10,296 nucleotides) per year. That is one of the lowest substitution rates among the species closely related to JEV and other well-studied members of mosquito-borne flaviviruses. The substitution rate of GI and GIII was evaluated to be 4.13×10^{-4} s/s/y (95% HPD interval, 3.45×10^{-4}– 4.82×10^{-4}) and 6.17×10^{-5} s/s/y (95% HPD interval 3.84×10^{-5}– 8.62×10^{-5}), accordingly, with the ages of clades of 153 (95% HPD, 87–237) and 216 (95% HPD, 139–317), respectively. The mean root height of JEV is 1234 years (95% HPD, 407–2333).

Keywords: substitution rate, molecular clock, Orthoflavivirus, phylogeny, RNA virus, evolution

1. Introduction

Japanese encephalitis virus (JEV) belongs to the species *Orthoflavivirus japonicum,* family *Flaviviridae* (Postler et al. 2023). JEV is the member of the mosquito-borne flavivirus (MBFV) group. The JEV genome consists of single-stranded positive-sense RNA with a length of about 10,965 nucleotides (nt). The genome encompasses a 10,296 nt single open reading frame (ORF), encoding a polyprotein, which includes three structural proteins: capsid (C), precursor membrane (prM) and envelope (E) proteins, and seven non-structural ones: NS1, NS2a, NS2b, NS3, N4a, NS4b, NS5 (Sumiyoshi et al. 1987; Xu et al. 2023).

Mosquitoes are the vectors for JEV, in particular, *Culex tritaeniorhynchus* and *Culex annulirostris*, which were recognized as principal ones (Van den Eynde et al. 2022). The virus is zoonotic, and it is transmitted in most cases between pigs or birds by mosquitoes, however, Ricklin et al. (2016) described the direct way of virus transmission between two pigs via oronasal secretions. Humans, cattle and horses are dead-end hosts, due to insufficient viremia to infect mosquitoes. When these hosts are infected, encephalitis can develop with fever, convulsions, coma and even death (Van den Eynde et al. 2022), (Ladreyt et al. 2019).

Approximately 3 billion people live in JEV epidemic areas according to the
World Health Organization data (https://www.who.int/news-room/fact-
sheets/detail/japanese-encephalitis, Retrieved 28 March 2024). JEV is the major
causative agent of human virus encephalitis disease in Asia. Now the virus is
endemic in the Torres Strait region of Australia and Asia, including 24 countries,
namely China, Korea, Indonesia, Malaysia, Japan and even Pakistan (Van den
Eynde et al. 2022), (Mackenzie 2005). Additionally, the virus was observed
outside the endemic areas in Italy (however, the data of the study are limited with
one short sequence, wherein JEV has not been registered in Europe since then)
(Ravanini et al. 2012) and Angola (Simon-Loriere et al. 2017).

Today, between 30,000 (Van den Eynde et al. 2022) and 69,000 (Auerswald
et al., 2021) cases registered annually according to different sources, but it is
considered as underestimated values due to insufficient monitoring and diagnosis.
Despite the fact that effective and safe vaccines have been developed (Vannice et
al. 2021), JEV is still one of the significant agents causing viral encephalitis in the
territory of Asia. Thus, to monitor and control this disease, it is important to
scrutinize evolutionary driving forces, which determine virus spread, genetic
variability and pathogenicity.

While recombination has previously been shown to contribute to JEV
evolution (Sistrom et al. 2024), most research focuses on the substitution rate. The
single-stranded RNA viruses exhibit stupendously high evolutionary rate spanned
several orders of magnitude (Duffy et al. 2008; Duchene et al. 2014). The members
of MBFVs evolve at a rate in the order of 10^{-4} nucleotide substitutions per site per
year (s/s/y) (Delatorre et al. 2019; Liu et al. 2019; Srihi et al. 2021; Zecchin et al.
2021; Stica et al. 2022; Xu et al. 2023). The evolutionary rate of JEV has been
assessed several times but the estimates vary widely in the different sources
spanning two orders of magnitude (10^{-5} to 10^{-4} s/s/y) (Mohammed et al. 2011; Pan
et al. 2011; Gao et al. 2015; Li et al. 2023; Xu et al. 2023). It is important to note
that none of the previous works has performed formal or any evaluation of
temporal signal in data sets prior to molecular clock application, and, therefore, the

reliability of the reported estimates is questionable. The accurate estimates of the

evolutionary rate are of importance for the robust interpretation of outbreak

investigations and are critical for understanding short-term virus transmission

patterns (Holmes et al. 2016; Theze et al. 2018; O'Toole et al. 2023; Crits-

Christoph et al. 2024). Besides, inferring evolutionary timescales helps to describe

and quantify the population dynamic processes that gave rise to the virus genetic

diversity observed.

The aim of our study is the reliable assessment of substitution rates and ages

of JEV with its genotypes and the (re-)evaluation of temporal signal in the data sets

from the previous studies using the Bayesian phylogenetic approach. To assess

JEV substitution rate, we used 325 complete ORF nucleotide sequences of all five

JEV genotypes with known collection dates available at the start of a study

(January 2023). Evolutionary rate and divergence times obtained were compared

with the previous estimates of JEV evolutionary rate and the rate of virus species,

closely related to JEV, and other well-studied MBFVs. Bayesian evaluation of

temporal signal makes reported results the first formally tested assessment of JEV

substitution rate.

2. Materials and methods

2.1 Nucleotide data set

As of January 2023, 440 nucleotide sequences of complete or near complete

genomes   of   JEV   were   found   in   GenBank   using   BLAST

(http://blast.ncbi.nlm.nih.gov/Blast.cgi). Clones, vaccine and chimeric strains were

removed from the data set and then collection dates were identified for 367 strains.

Recombinant sequences detected by RDP5 (Martin et al. 2021) were removed from

the alignment. Thus, our research was conducted using 325 nucleotide sequences

of a polyprotein gene.

Next, sequence alignment was done with MAFFT v.7.490 (Katoh and

Standley 2013). After that, the alignment was visualized in AliView v.1.24

(Larsson 2014) to exclude ORFs with stop-codons. Non-coding regions were cut

off using the NCBI reference sequences (NC_001437.1).

JEV genotypes were determined in preliminary phylogenetic analysis in IQ-
TREE v.1.6.12 using the tree topology reported by (Takhampunya et al. 2011). It
was revealed that 134 samples belong to GI, 7 to GII, 171 to GIII, 6 to GIV and 7
to GV. Thus, only GI and GIII have a sufficient number of sequences to obtain
reliable statistics in the further independent Bayesian phylogenetic analyses.
Therefore, we subsequently analyzed three data sets—the common clade of JEV
(JEV GI-GV), GI and GIII.

NCBI accession numbers, as well as fasta and BEAST xml files can be
found in the supplementary materials.

2.2 Temporal signal assessment

Temporal signal (or structure) assessment is the key step in substitution rate
evaluation. In the case of absence of temporal signal, the program (BEAST in our
case) inferred values based on the priors (e.g., population size prior) but not the
data that potentially bias the entire analysis.

Temporal structure in each of the data sets (JEV GI-GV, GI and GIII) was
assessed with the Bayesian evaluation of temporal signal (BETS) (Duchene et al.
2020) in BEAST v.2.6.7.

BETS involves the comparison of time-stamped (heterochronous) and
isochronous models (M_{het} and M_{iso}) based on the difference between log marginal
likelihoods   (for   a   tutorial   of   using   BETS   see:
https://beast.community/bets_tutorial). The log marginal likelihoods for each
model were calculated using the path sampling estimator implemented in the
MODEL_SELECTION package from the BEAST2 Package Manager. The number
of steps was set as 200 and the lengths of MCMC for each step were determined so
that the product of the number of steps and MCMC length equaled the length of
the MCMC during the main phylogenetic analysis in BEAST with an effective
sample size (ESS) at least of 200. Priors on model parameters (molecular clock,
substitution model, population model) were chosen similar to those used in the
main analysis (Table S1) with the exception of the tree prior—in this case, we

replaced the Bayesian skyline with the coalescent constant population with a

proper prior (i.e., integrated to 1) on effective population size (Table 1).

Table 1. Priors used in the Bayesian evaluation of temporal signal (BETS) in all
three data sets (JEV GI-GV, GI, GIII)

Model	Parameter		Prior
Strict clock	Clock rate		$\Gamma(\alpha = 0.1, \beta = 100)$
Optimized relaxed clock	Relative branch rates		LogNormal(M = 1.0, S = 0.2)
	Sigma		$\Gamma(\alpha = 0.5396, \beta = 0.3819)$
	Mean clock rate		$\Gamma(\alpha = 0.1, \beta = 100)$
Rate heterogeneity	Shape of Γ distribution		Exp(M = 1.0)
	Proportion of invariant sites		Uniform(0.0, 1.0)
Substitution model	GTR rates	Transitions	$\Gamma(\alpha = 0.05, \beta = 20)$
		Transversions	$\Gamma(\alpha = 0.05, \beta = 10)$
Tree prior (coalescent constant size)	Effective population size		Exp(M = 100)

The comparison of different molecular clock can also be involved in BETS

for calculation of marginal likelihoods for each model. Therefore, we included a

strict clock (SC) and an optimized relaxed clock (ORC) (Douglas et al. 2021)

models into the comparison.

The difference between log marginal likelihoods of competitive models (i.e.,

log Bayes factor) of at least 5.0 indicates very strong support, a value of 3 means

strong support and a value of 1 is considered to be positive evidence (Kass and

Raftery 1995).

For comparison purposes, we conducted widely used root-to-tip (RTT)

regression, which is an informal test with well-known limitations: data points are

not independent and strict clocklike behavior is assumed. RTT statistics were

computed with Clockor2 (Featherstone et al. 2024). The trees for Clockor2 were

reconstructed in IQ-TREE v.1.6.12 with the best-fit model determined for each

data set using ModelFinder (Kalyaanamoorthy et al. 2017). The JEV GI-GV tree

was rooted in Clockor2 by maximizing the coefficient of determination (R^2)

values. To calculate R^2 for JEV GI and GIII genotypes, the trees were rooted using

information from the JEV GI-GV phylogeny.

Additionally, temporal signal was evaluated using BETS (path sampling

method) in the data sets from the previous works of Gao et al. (2015), Li et al.

(2023), Mohammed et al. (2011), Xu et al. (2023), and Pan et al. (2011) available

as of November 2023. In total, 9 data sets were gathered (Table 2):

Table 2. Data sets of JEV open reading frame sequences from the previous studies

Study	JEV genotype	Number of samples	Sampling window (years)
Mohammed et al. (2011)	GI-GV	34	74.4
	GI	10	15.4
	GIII	21	73.0
Pan et al. (2011)	GI-GV	95	74.0
	GI	43	32.0
	GIII	50	70.6
Gao et al. (2015)	GI-GV	98	74.0
Li et al. (2023)	GI	131	42.0
Xu et al. (2023)	GI-GV	115	87.1

174
One sample from the study of Mohammed et al. (2011) and three samples from the

study of Pan et al. (2011) previously genotyped as GIII are absent in our data sets

since these sequences were removed from GenBank at the request of the authors.

As a molecular clock model, we used an uncorrelated log-normal relaxed

clock (UCLD) as in the original works.

2.3 Dated phylogeny reconstruction and substitution rate evaluation

Bayesian phylogenetic analysis using JEV GI-GV, GI and GIII data sets was

performed in BEAST v.2.6.7 (Bouckaert et al. 2019). The selection of substitution

model was based on the BIC previously assessed by ModelFinder in IQ-TREE

(GTR+Γ_4+I for all data sets). Molecular clock models were compared using the log

marginal likelihoods calculated during the BETS step. As a tree prior, we specified

a flexible non-parametric Bayesian Skyline model. For details on all priors, see the

supplementary.

The lengths of MCMC in each run were chosen so that the ESS associated

with estimated parameters reached at least 200 samples. The inspection of model

parameters and MCMC convergence was carried out in Tracer v.1.7.1 (Rambaut et

al. 2018). The burn-in thresholds of MCMC chains were defined individually

depending on the chain behavior. Maximum clade credible (MCC) trees were
annotated in TreeAnnotator (Bouckaert et al. 2019). Due to high computational
cost, BEAST was run using a BEAGLE library (Ayres et al. 2012) on a GPU
(NVIDIA GeForce RTX 3070).

2.4 Comparing evolutionary rates among JEV subtypes estimated using
a relaxed molecular clock

Relaxed molecular clocks allow evolutionary rates vary among tree
branches. Therefore, the among-branch rate variation should be taken into account
when comparing the substitution rates of different JEV subtypes. To address this
problem, we calculated the posterior probability for a hypothesis that a random rate
value from a lognormal rate distribution of one genotype is larger than a random
rate value from a lognormal rate distribution of another genotype. To approximate
this probability, we performed Monte Carlo simulations in R (Figure 2):

205
Figure 1. Flow chart of the comparative analysis of substitution rates between a pair of
virus genotypes evaluated using relaxed molecular clock. Random samples from lognormal
distributions were drawn with Monte Carlo simulations in R. Asterix—ucldMean parameter
values in real space must be transformed to log space: $\log(ORCucldMean) - \frac{ORCsigma^2}{2}$.

For each genotype pair, we randomly sampled mean ($\mu\square$, $\mu\square$) and standard
deviation ($\sigma\square$, $\sigma\square$) values from their posterior distributions and generated two sets
of 100,000 draws each: $X_1 \sim lognormal(\mu_1, \sigma_1)$ and $X_2 \sim lognormal(\mu_2, \sigma_2)$.

For each simulated pair (x_1, x_2), where $x_1 \in X_1$ and $x_2 \in X_2$, we computed the

frequency of $x_1 > x_2$, yielding $P(X_1 > X_2 \mid data)$. This procedure was repeated

1000 times to estimate 95% credible intervals (CIs).

3. Results

3.1 Degree of clocklike behavior of the common JEV data set (GI-GV)

and GI and GIII genotypes

In the first step, using three JEV data sets (GI-GV, GI, GIII) we conducted

RTT regression to assess to which degree data behave in a clock-like fashion

(Figure 2).

223
Figure 2 – Root-to-tips (RTT) regression analysis of three JEV data sets; X-axes represent
calendar years, Y-axes—distances from a root to tips.

The JEV GI-GV and GIII data sets showed R^2 values close to zero (0.04 and

0.12 with a negative slope, respectively) indicating the weak clocklike behavior of

the data which can be explained by the presence of a large amount of rate variation

between lineages of the trees (95% HPD of the coefficient of rate variation (CV),

1.04–1.62 for GI-GV and 1.29–2.4 for GIII). GI, on the contrary, demonstrated a

stronger temporal structure with R^2 of 0.79 and the lowest CV values (95% HPD,

0.43–0.68) (Table 3).

Table 3. Comparison of the determination coefficients inferred in the root-to-tip
regression analysis and the coefficient of variation values.

Data set	Number of samples	Sampling window, years	Determination coefficient (R^2) inferred in RTT analysis [1]	95% HPD of the coefficient of rate variation [2]	95% HPD of the substitution rate (s/s/y) [3]
GI-GV	325	87	0.04	1.04–1.62	1.67×10^{-4}–3.14×10^{-4}
GI	134	45	0.79	0.43–0.68	3.45×10^{-4}–4.82×10^{-4}
GIII	171	87	0.12	1.29–2.4	3.84×10^{-5}–8.63×10^{-5}

[1] Root-to-tip (RTT) regression analysis was performed in Clockor2;
[2] 95% HPDs were inferred in BEAST analysis;
[3] S/s/y—substitutions per site per year.

Importantly, the low R^2 values or a negative slope should not be

unequivocally considered as the absence of temporal signal (Rambaut et al. 2016).

RTT regression analysis appears to be uninformative when the data were sampled

over a relatively narrow sampling window (GI-GV data set) and/or there is large

rate variation among lineages (GIII data set) (Duchene et al. 2020). These factors

result in decreasing R^2. Therefore, we applied BETS—a formal test of temporal

signal—to assess the statistical support for including sampling times in a Bayesian

analysis using both the strict clock and optimized relaxed clock to take into

account among-branch rate variation.

3.2 Presence of temporal signal in all JEV data sets

BETS displayed very strong support (log Bayes factor > 5) for M_{het} over M_{iso}

in each pair of competitive models (Table 4) with the lowest log Bayes factor of

47.8 for the GIII data set (ORC_{het} relative to ORC_{iso}) which possessed the largest

amount of rate variation among branches (Table 3). Despite the last fact, in the log

scale, such log Bayes factor values can be interpreted as overwhelming support for

including sequence sampling times into the model (Table 4, Figure 3).

Table 4. Results of the Bayesian test of temporal signal in Japanese encephalitis
virus data sets.

Data set	Molecular clock [1]	Log marginal likelihood	Log Bayes factor [2]
GI-GV	ORC_{iso}	−113559.93	140.38
	ORC_{het}	−113419.55	0
	SC_{iso}	−115098.41	1678.86
	SC_{het}	−114995.46	1575.91
GI	ORC_{iso}	−54633.51	129.02
	ORC_{het}	−54504.49	0
	SC_{iso}	−55182.96	678.48
	SC_{het}	−54728.94	224.45
GIII	ORC_{iso}	−46598.66	47.80
	ORC_{het}	−46550.86	0
	SC_{iso}	−47491.94	941.08
	SC_{het}	−47529.78	978.92

[1] ORC—optimized relaxed clock, SC—strict clock; iso—isochronous model, het—heterochronous model;

[2] Best-performing model has a log Bayes factor value of 0.

Figure 3. Relative log marginal likelihoods of each JEV data set (GI-GV, GI, GIII) analyzed in our study. Log marginal likelihoods were calculated with four different configurations of sampling times (heterochronous and isochronous trees) and molecular clocks. In the corners, there are models with combinations of molecular clocks and the inclusion or exclusion of sampling times. The outermost dashed borders of the squares correspond to the highest log marginal likelihood calculated. The extent to which the polygons are stretched depicts *relative* model support such that the perfect square indicates equal support for each model. The corner of a polygon that falls close to the center would correspond to a model with the lowest support. SC is a strict clock and ORC is an optimized relaxed clock. *Het* (heterochronous) includes sampling times, while *iso* (isochronous) exclude sampling times. An orange corner points at the model combination with the highest log marginal likelihood. The numbers near the polygon edges are the Bayes factors for the best model (orange corner) over the model with less support. The values above the charts are sampling windows (sw) in years and sample sizes (n).

The comparison of the SC_{het} and ORC_{het} models also showed very strong evidence in favor of ORC in each data set (Table 4). The lowest log Bayes factor of 224.45 was detected for, as expected, GI which had the highest R^2 in the RTT regression analysis.

Thus, all three data sets passed the Bayesian test of temporal signal and were

analyzed with ORC in the next step of phylogenetic analysis.

3.3 Analysis of temporal signal in the previous studies

We analyzed temporal structure in the nine data sets from the previous
studies (Figure 4):

Figure 4. Relative log marginal likelihoods of nine JEV data sets from the previous works
(note – data set of a JEV common clade from the study of Pan et al. (2011) exclude the genotype
V) calculated with four different configurations of sampling times (heterochronous and
isochronous trees) and molecular clocks. The rows of the charts are JEV genotypes, the columns
depict the studies from which the data sets were taken. In the corners of each chart, there are
models with combinations of molecular clocks and the inclusion or exclusion of sampling times.
The extent to which the polygons are stretched depicts *relative* model support such that the
perfect square indicates equal support for each model. The outermost dashed borders of the
squares correspond to the highest log marginal likelihood. The corner of a polygon that falls
close to the center would correspond to a model with the lowest support. SC is a strict clock and
UCLD is an uncorrelated log-normal relaxed clock. *Het* (heterochronous) includes sampling
times, while *iso* (isochronous) exclude sampling times. An orange corner points at the model
combination with the highest log marginal likelihood. The numbers near the polygon edges are
the Bayes factors for the best model (orange corner) over the model with less support. The values
above the charts are sampling windows (sw) in years and sample sizes (n). The polygons with
dashed edges display the data sets with no temporal signal, while solid polygons are the data sets
where temporal signal was detected (i.e., heterochronous models including sampling times are
favored).

BETS revealed that temporal signal is present in the five of nine JEV data

sets (Pan et al. (2011), GI-GIV, GIII; Gao et al. (2015), GI-GV; Li et al. (2023),

GI; Xu et al. (2023), GI-GV) with very strong support (the lowest log Bayes factor

is 11.02 for JEV GIII from Pan et al. (2011) that contains the lowest number of

complete ORF sequences equal to 50). Interestingly, the Bayes factors increased

along with sample size (Table S2, Figure S1) which was most likely due to the

increase in the number of alignment informative sites. In all data sets from the

study of Mohammed et al. (2011) temporal signal was not detected even in the GI-

GV data set with 34 complete ORF sequences (log Bayes factor of 5.9 for $UCLD_{iso}$

over $UCLD_{het}$). However, in the case of the GI (n=10) and GIII (n=21) data sets

from the same work, BETS displayed only positive support for $UCLD_{iso}$ over

$UCLD_{het}$ with the log Bayes factors of 1.8 and 2.1, respectively. The last data set

with no temporal signal was JEV GI from the study of Pan et al. (2011) despite the

sample size of 43 complete ORF sequences: path sampling showed very strong

support for $UCLD_{iso}$ over $UCLD_{het}$ (log Bayes factor = 6.6).

3.4 Substitution rates

The mean substitution rate of the common JEV GI-GV clade was 2.41×10^{-4}

s/s/y (95% HPD, $1.67 \times 10^{-4} - 3.14 \times 10^{-4}$ s/s/y) or approximately 2.5 nucleotide

substitutions per ORF (10,296 nucleotides) per year. GI has the highest mean rate

of evolution: 4.13×10^{-4} s/s/y (95% HPD, $3.45 \times 10^{-4} - 4.82 \times 10^{-4}$ s/s/y) or

approximately 4.2 nucleotide substitutions per ORF per year. The lowest mean

substitution rate was obtained for GIII: 6.17×10^{-5} s/s/y (95% HPD, $3.84 \times 10^{-5} -$

8.63×10^{-5} s/s/y) or approximately 0.6 nucleotide substitutions per ORF per year

(Figure 5).

Figure 5. Substitution rates of common JEV clade (GI-GV), genotypes I (GI) and III
(GIII). Upper and lower ends of the boxes are 95 % highest posterior density interval borders.
The Y-axis represents substitution rate and is expressed in the number of substitutions per site

per year (s/s/y). The values above the boxplots are sampling windows (sw) in years and sample
sizes (n)

Despite the largest CV (Table 3), the 95% HPD interval for JEV GIII was
the narrowest one (Figure 5). The issue of a high-degree of among-branch rate
variation in the JEV GIII data set is likely due to passaging some virus strains. In
particular, 34 of 171 JEV GIII were isolated from before 1980. In addition, there
are clusters with closely related sequences isolated contemporaneously (phylo-
temporal structure, Figure 6). Inspecting a rate variation pattern on the MCC tree
(Figure S2) revealed the most outstanding substitution rate value of 0.0014 s/s/y on
the external branch of BJ-1_BP28 strain (KU871371.1) collected in 2015 (the
study has not been published). However, the reanalysis without this strain inferred
the nearly identical estimate of mean substitution rate (95% HPD, 3.71×10^{-5}–
8.22×10^{-5} s/s/y) (Supplementary Material).

The mean substitution rate values of the JEV GI-GV clade lies between
estimates for JEV GI and GIII and represents the average of rates across branches
of the entire tree.

While the 95% HPD intervals of the mean rates do not overlap, among-
branch rate variation should be taken into account—particularly for the two data
sets (JEV GI-GV and GIII), which exhibit coefficients of rate variation above 1.0
and even 2.0 (Table 3). In other words, some branches of the JEV GIII phylogeny
can *probably* exhibit higher rates than those of JEV GI, even though the mean of
GIII substantially lower. To resolve this issue, we used 1,000 posterior-sampled
mean and sigma values to perform Monte Carlo simulations in R, calculating the
posterior probability that a rate drawn from a lognormal distribution of one
genotype would exceed that of another genotype. We tested the following three
pairs of JEV data sets (Table 5):

Table 5. Posterior probability that a lineage from Genotype 1 evolves faster than
that of Genotype 2 approximated by Monte Carlo simulations in R

Genotype 1 (X_1)	Genotype 2 (X_2)	95% CI for $P(X_1 > X_2 \mid data)$*
JEV GI	JEV GIII	0.64–0.79

JEV GI	JEV GI-GV	0.87–0.97
JEV GI-GV	JEV GIII	0.69–0.84

The results of the comparison indicate that while JEV GI, on average,
evolves faster than JEV GIII and the common JEV GI-GV clade, the reverse
scenario can also occur with some probability. In the pair of JEV GI and JEV GIII,
the 95% CI (0.64–0.79) indicates a 21–36% probability that rates from the
lognormal distribution of JEV GIII exceed those from JEV GI although the mean
evolutionary rate of JEV GI is substantially higher than that of JEV GIII (Figure
5). This is probably due to the largest coefficient of variation of JEV GIII (1.29–
2.4). The similar observation was made for the comparison of evolutionary rates of
JEV GI-GV > JEV GIII (0.69–0.84).

For the JEV GI versus JEV GI-GV comparison, the 95% credible interval
(0.87–0.97) indicates a high probability that a rate sampled from the JEV GI
distribution would exceed that of the JEV GI-GV common clade distribution.

3.5 Divergence times

The evaluation of JEV divergence times was performed using 325 complete
ORF nucleotide sequences (Figure 6). The results of the analysis in BEAST
suggest that the tMRCA of JEV is 1234 years before 2022 (95% HPD, 407–2333
years).

Figure 6. Maximum clade credibility tree of JEV summarized from a posterior tree distribution produced by an MCMC run in BEAST. The numbers nearby the main nodes indicate divergence times before the most recent sample collected on February 20, 2022, with 95% highest posterior density (HPD) intervals within parentheses. The horizontal color bars depict 95% HPD of the divergence times of JEV genotypes.

Concerning the emergence of JEV genotypes, 95% HPD intervals of time to MRCA of all JEV genotypes are overlapped (Figure 6). Therefore, we are unable to reliably establish the chronology of the occurrence of genotypes. Besides, only the GI and GIII are representative in terms of sample size, and, as a consequence, the estimates of the other genotype ages (GII, GIV, GV) are unreliable. GI and GIII ages, in their turn, are not statistically indistinguishable: GI occurred 153 years ago (95% HPD, 87–237) and GIII—216 years ago (95% HPD, 139–317). Generally, considering the bounds of 95 HPDs obtained, JEV genotypes occurred 77–540 years ago.

4. Discussion

Our study demonstrates the first assessment of JEV evolutionary rate supported by the formal analysis of temporal signal in the nucleotide data. In the literature, there is substantial discrepancy between JEV substitution rate values estimated by different authors. Therefore, we also evaluated temporal structure in the data sets from the previously reported studies. Our findings indicate which of those estimates are reliable and which were based on uninformative data and/or on

the priors set in a program by the researchers on their own. The results of our
analysis show that JEV GI is the fastest-evolving genotype and, conversely, GIII is
the slowest JEV genotype. Regarding the width of 95% HPD interval, our JEV rate
estimates are among the most precise.

4.1 Comparing the mean substitution rates of JEV yielded in our
analysis with estimates reported previously

We compared the JEV substitution rate values obtained in our study with the
previous estimates based on the analysis of complete ORF gene sequences (Figure
7).

407
Figure 7. Comparison of the Japanese encephalitis virus (JEV) substitution rate evaluated
in our study based on complete open reading frame nucleotide sequences and the JEV
substitution rates according to the literature. The vertical bars represent 95% highest posterior
density intervals, circles—median or mean values. The X-axis labels are references to
appropriate studies, the methods on temporal signal evaluation applied are indicated above the
X-axis: BETS—Bayesian evaluation of temporal signal, RTT—root-to-tips regression, *—
temporal signal was not assessed.

First of all, we found only one study where at least the degree of clock-like
behavior was assessed with informal RTT (Li et al. 2023) when analyzing GI.
Therefore, the substitution rate values reported in the literature should be treated

with caution, especially in the cases where the number of samples was relatively

low. Hence, to our knowledge, we provide here, for the first time, the JEV

evolutionary rate estimates confirmed by a formal statistical test. Besides, we

additionally assessed temporal signal with BETS in the data sets from the previous

studies to improve the rigor of the manuscript and attempt to explain the difference

between rate estimates observed. Four of nine data sets analyzed displayed the

absence of temporal signal (Figure 4, Table S2). Our results emphasize the

importance of temporal signal evaluation in the data sets represented by dozens of

sequences, even those that are complete or near complete genomes.

4.1.1 JEV GI-GV clade

The comparison showed that the evolutionary rate of the JEV common clade

including all five genotypes GI-GV obtained in our study (95% HPD, 1.67×10^{-4}–

3.14×10^{-4} s/s/y) is substantially lower than values reported by Mohammed et al.

(2011) (95% HPD, 3.49×10^{-4}–5.30×10^{-4}) who analyzed the data set with the

smaller sample size (n=35). Our evaluation of temporal signal in the data set of

JEV GI-GV from the study of Mohammed et al. (2011) revealed the absence of

temporal signal with very strong support for the isochronous models over

heterochronous ones (log Bayes factor = 6.6) (Figure 4) which explained the

discrepancy between estimates inferred in our work and in the study of

Mohammed et al. (2011).The fact is that, in the case of low-informative and

uninformative data, the Bayesian analysis turns out to be very sensitive to a prior

choice. In particular, under the coalescent process an inferred substitution rate is

largely rested on, for example, a continuous-time Markov chain (CTMC) rate

reference prior (default prior on the clock rate in BEAST1). The CTMC prior is a

gamma probability distribution with a shape parameter $\alpha=0.5$ and rate parameter

$\beta=T$, where T is tree length, meaning that the CTMC is ultimately conditioned by

an effective population size prior (under the coalescent process the expected time

of divergence between lineages is inversely proportional to the population size)

(Ferreira and Suchard 2010). In brief, the more population size, the longer tree and,

as a consequence, the slower evolutionary rate estimated. Thus, interactions

between the population size and CTMC priors may drastically affect the inference
of substitution rates (we demonstrated this effect using the data set of JEV GI from
the study of Mohammed et al. (2011), Figure S3).

In contrast, our estimates are substantially higher than the JEV substitution
rates provided by Pan et al. (2011) (95% HPD, 6.04×10^{-5}–1.69×10^{-4}) and Gao et
al. (2015) (95% HPD, 4.37×10^{-5}–1.56×10^{-4}) who analyzed 98 (GI-GIV) and 100
(GI-GV) sequences, respectively. Our analysis of temporal signal in the data sets
from those works revealed that BETS supported the UCLD model with sampling
times (log Bayes factors of 22.7 and 25.0 for $UCLD_{het}$ over $UCLD_{iso}$, respectively)
(Figure 4). The discrepancy in the substitution rate evaluation between our and the
previous works can be explained by the difference in data sets and sample sizes.
This is particularly important in the event of high degree of among-lineage rate
variation since excluding or including certain tips can drastically affect inferred
CV values and, as a consequence, virus evolutionary rate. In the case of the study
of Pan et al. (2011), the authors did not include JEV GV in the analysis of the
common clade of JEV which also explain the discrepancy between the estimates of
a JEV substitution rate.

In one of the last studies, Xu et al. (2023) reported a JEV GI-GV substitution
rate (95% HPD, 1.62×10^{-4}–3.97×10^{-4}) comparable with our estimates. BETS also
demonstrated strong support for temporal signal in this data set (Figure 4) with the
log Bayes factor of 32.4 for $UCLD_{het}$ over $UCLD_{iso}$. Notably, the sample size
(115), clock model (UCLD) and tree prior (Bayesian Skyline) applied in the work
of Xu et al. (2023) were similar to two previously discussed studies with
substantially lower JEV substitution rates (Pan et al. 2011; Gao et al. 2015). The
possible explanation is that Xu et al. (2023) provide new JEV nucleotide
sequences, which were not used in our study and in the previous works of the other
authors.

4.1.2 JEV GI clade

Our estimates of JEV GI evolutionary rate (95% HPD, 3.45×10^{-4}–
4.82×10^{-4}) are substantially higher than those we calculated for the GI-GV clade

(95% HPD, 1.67×10^{-4}–3.14×10^{-4}) (Figure 5). JEV GI rate values inferred in our

study are also substantially different from the literature data, except for the

estimates with the widest 95% HPD interval (2.09×10^{-8} 8.7×10^{-4}) provided by

Mohammed et al. (2011) using only 10 complete ORF sequences. BETS revealed

the absence of temporal structure in this data (Figure 4) with log Bayes factor for

$UCLD_{iso}$ over $UCLD_{het}$ of 2.1 (positive evidence).

The study of Pan et al. (2011) (n=43) reported the substitution rate estimates

with an extremely low border of 95% HPD (3.42^{-9}–2.47×10^{-4}), which indicates

weak or absent temporal signal. In our evaluation of temporal structure in this data

set, the UCLD model in which sampling times are contemporaneous ($UCLD_{iso}$)

was favored with very strong support (log Bayes factor = 6.6) over the UCLD

model in which the data was accompanied by the actual sampling times ($UCLD_{het}$)

(Figure 4).

The only study (Li et al. 2023) in which the extent of time-dependency in

data was assessed (RTT regression analysis) reported statistically higher JEV GI

evolutionary rates (95% HPD, 5.03×10^{-4}–6.94×10^{-4}) than our estimates. Our

assessment of temporal signal in this data set showed very strong support for the

presence of temporal signal (log Bayes factor for $UCLD_{het}$ over $UCLD_{iso}$ of 145.8).

We speculated on three possible explanations of the discrepancy—the first one is

the authors removed RTT regression outliers without explaining the reason (e.g.,

sequencing errors, long passage history, lab contamination, etc.) that is forcing the

data to fit a strict clock model, which does not assume among-lineage rate

variation, instead of using an alternative model (e.g., relaxed clock) to describe the

data; the second one is the new JEV nucleotide sequences generated in the study of

Li et al. (2023) and not included in our work; the third reason is the smaller sample

size (n=131) compared to our work.

4.1.3 JEV GIII clade

Among all three data sets analyzed (Table 3), JEV GIII was characterized by

the lowest mean evolutionary rate (95% HPD, 3.84×10^{-5}–8.63×10^{-5}) (Figure 4).

Thus, JEV GIII evolves generally on the other order of magnitude than JEV GI and
JEV GI-GV.

The very similar estimates (4.28×10^{-5}–1.18×10^{-4}) were reported by Pan et
al. (2011) with a sample size of 53 and a model combination comparable to the one
we applied in our work with the exception of a relaxed molecular clock (UCLD).
Our analysis of temporal structure with BETS very strongly supported
heterochronous model over an isochronous one (log Bayes factor of 11.02). In the
study of Mohammed et al. (2011), the 95% HPD interval was the widest one
(7.05×10^{-8}–6.03×10^{-4}). As in the case of JEV GI, BETS indicated the absence of
temporal signal in the JEV GIII data set from the work of Mohammed et al. (2011)
and the main reason is the small sample size (n=22) and, therefore, the small
number of variable sites in the alignment.

However, it is worth noting that regardless of precise estimates, the JEV GIII
data set includes a large number of strains isolated before the 1980s, which could
be serially passaged over a period of decades before being sequenced. This leads to
artificially accelerated accumulation nucleotide substitutions and, as a result, to the
overestimation of a substitution rate. Also, the long passage history is the most
likely cause of the high CV values (Table 3) and the RTT negative slope (Figure 2)
inferred. Thus, the true JEV GIII evolutionary rate is likely *lower* than the values
reported in our study.

4.2 Comparing the substitution rates of JEV (GI-GV clade) and other
mosquito-borne flaviviruses

To provide a better understanding of the JEV evolutionary rate, we
compared the rate of JEV GI-GV clade obtained with rates of JEV's most closely
related species and the other well-studied MBFV (Figure 8).

To compare the JEV substitution rate, we considered the data only if 95%
HPD intervals for mean/median were provided. Also, the estimates reported with
mistakes or obvious typos were excluded from the comparison. We mainly focused
on the substitution rates derived from the analysis of as many virus genotypes as

possible (i.e., the genetic diversity of viruses was meant to be represented to its
greatest extent).

538
Figure 8. Comparison of the Japanese encephalitis virus (JEV) substitution rate inferred
in our study using complete open reading frame nucleotide sequences of GI–GV, GI, and GIII
genotypes, and the substitution rates of the other mosquito-borne flaviviruses according to the
literature: USUV—Usutu virus (Engel et al. 2016; Zecchin et al. 2021), MVEV—Murray Valley
encephalitis virus (Williams et al. 2015), WNV—West Nile virus (May et al. 2011; McMullen et
al. 2013; Srihi et al. 2021), YFV—yellow fever virus (Delatorre et al. 2019; Moreira Salles et al.
2022), ZIKV—Zika virus (Sun et al. 2017; Liu et al. 2019), DENV—dengue virus (Stica et al.
2022). The person icons to the right of the abbreviations point to the viruses for which humans
are not dead-end hosts. The vertical bars represent 95% highest posterior density intervals,
circles—median or mean values if available. The X-axis labels are references to appropriate
studies. The methods on temporal signal evaluation applied are indicated above the X-axis:
BETS—Bayesian evaluation of temporal signal, RTT—root-to-tips regression, PT—permutation
test, *—temporal signal was not assessed.

Unlike the literature data on JEV, the most of MBFV rate estimates were
supported by the RTT regressions, and, furthermore, a permutation or
randomization test was applied in one study (Srihi et al. 2021). This makes the
comparison much more reliable, but, for some virus species, reported rates are
statistically different between studies.

First, we compared our JEV substitution rate estimates with the rates of JEV

most closely related species (Moureau et al. 2015)—Usutu virus (USUV), Murray

Valley encephalitis virus (MVEV), West Nile virus (WNV)—for which the sample

size allows the evolutionary rate of the virus to be reliably evaluated. Second, we

included in the comparison the most widely understood members of MBFVs—

yellow fever virus (YFV), Zika virus (ZIKV), and dengue virus (DENV).

As shown in Figure 8, JEV is one of the slowest changing MBFVs

represented in our analysis (95% HPD, 1.67×10^{-4}–3.14×10^{-4}). However, when

accounting for among-branch rate variation (Tables 3 and 5), JEV can produce

branches with evolutionary rates that exceed the mean rate of JEV GI, bringing it

closer to the other MBFVs.

The overlap of 95% HPD intervals is observed in the case of two

assessments of WNV (2.24×10^{-4}–5.34×10^{-4} according to McMullen et al. (2013)

and 2.50×10^{-4}–4.47×10^{-4} for WNV lineage 2 reported by Srihi et al. (2021)). There

is a slight overlap with the rate values of the USUV European lineage (95% HPD,

3.04×10^{-4}–3.77×10^{-4}). More distant MBFV members evolve faster, especially

ZIKV (Sun et al. 2017; Liu et al. 2019) and DENV (Stica et al. 2022), the

substitution rates of which are near the threshold of 10^{-3} and 10^{-2} orders of

magnitude or even above it.

The low substitution rate of virus can play a positive role in the virus strain

selection for the vaccine production. Vaccines for slowly evolving viruses remain

effective longer. To date, there are four types of effective JEV vaccines based on

the GIII strain (Srivastava et al. 2023). Regarding the more rapidly changing

viruses, as a rule, there is either no effective vaccine (e.g., ZIKV), or in the case of

DENV the vaccinated person had to be previously infected with the virus

(https://www.who.int/publications/i/item/who-wer-9918-203-224) (although there

is an effective vaccine against poliovirus despite the virus exhibiting a high

evolutionary rate (Guo et al. 2023)). Likewise, the vaccine strain should be

constantly updated as in the case of SARS-CoV-2 or influenza viruses.

Rates of evolution are determined by many factors including mutation rate
(influenced, for instance, by RNA replication errors), natural selection, generation
time (time between successive infections in transmission chains) and population
size (Duchene et al. 2016). There is also a negative correlation between virus
genome size and evolutionary rate (Ho 2020). Considering RNA viruses which
exhibit evolutionary rate between 10^{-4} and 10^{-3} orders of magnitude (Duchene et
al. 2014), the mean substitution rate of JEV (95% HPD, 1.67×10^{-4}–3.14×10^{-4} s/s/y)
is relatively slow. The reason why the JEV evolution rate was slower than its
closest relatives is unclear and the answer likely lies in the complex interaction of
the described factors. We assume that humans which are not dead-end hosts
(Figure 8) in the case of fastest evolving viruses (YFV, ZIK, DENV) may shorten
the time between successive infections. However, USUV and MVEV viruses
demonstrate relatively high evolutionary rates, (although, calculated without a
formal test of temporal signal). Therefore, humans as amplifying hosts can play a
rather indirect role in virus viability.

4.3 Divergence time
Despite the relatively large number of complete ORF sequences used in our
study, the uncertainty in the assessment of the evolutionary timescale of the entire
tree spans a large time interval (95% HPD of root height, 407–2333 years). A little
more precise estimates of time to the most recent common ancestor (MRCA) of
JEV GI-GV (95% HPD, 463–2100 years) were reported by Xu et al. (2023) with a
sample size of 115 complete genome sequences. In the other studies with smaller
sample sizes, 95% HPD intervals for tree roots were much wider: Gao et al. (2015)
conclude that the MRCA of JEV GI-GV is estimated to have occurred 3255 years
ago (95% HPD, 978–6125 years) encompassing much of the range of our estimate;
Pan et al. (2011) studied JEV GI-GIV defined the MRCA age of the entire clade as
1695 years ago (95% HPD, 548–3153 years). More recent JEV emergence was
described by Mohammed et al. (2011) (95% HPD, 374–500 years ago), but the
sample size was the smallest one (n=35) which implies lack of temporal structure
in the data and is also reflected in the substitution rate estimates (Figure 7).

The yielded ages of JEV subtypes are consistent with the previous
assessments (Xu et al. 2023). However, the "minor" JEV genotypes, such as GII,
GIV, GV, were mainly scrutinized in the studies of E gene data sets. The estimates
of time to MRCA based on E gene sequences are more precise (Schuh et al. 2013;
Sikazwe et al. 2022), whereas the substitution rate of E gene is substantially higher
(95% HPD, 3.92×10^{-4}–6.52×10^{-4}) and MRCA ages are younger (e.g., the MRCA
of JEV GI-GV, 95% HPD 216–921 years) (Schuh et al. 2013); JEV GI, 44–55
years (Li et al. 2025)). Rejuvenation of the tree root along with (or possibly due to)
an increase in the evolutionary rate indicates the impact of the priors on the
inference of the evolutionary timescale. In other words, even if the E gene of JEV
evolved faster than the entire ORF gene, then the dates of the emergence of JEV
MRCA should not substantially differ between the analyses of two loci in the case
of the obligatory presence of temporal signal in both data sets. However, the
sample sizes also can play a significant role. In the cited works dedicated to the E
gene analysis, only Sikazwe et al. (2022) performed the RTT analysis, however,
JEV GV samples were removed from the analysis as outliers without a justified
reason. Moreover, the R^2 and the regression visualization were not reported as well
as the substitution rate of the common data set (JEV GI-GIV). Thus, the problem
of the fitness of the E gene for JEV date-stamped phylogeny reconstruction should
be addressed in future studies.

Conclusions

1.   All complete ORF data sets analyzed (JEV GI-GV, GI, GIII) yielded
strong temporal signal in BETS. It means that a sufficient amount of molecular
evolution occurred over the sampling time window to permit collection dates of
sequences for clock calibration. Thus, JEV population can be treated as measurably
evolving and the reconstructions of JEV dated phylogenies is warranted and
reliable;

2.   JEV represented by the clade comprising all five genotypes (GI-GV)
has the lowest mean substitution rate in comparison to its most closely related
species and other well-studied members of MBFV: 2.41×10^{-4} s/s/y (95% HPD,

1.67×10^{-4}–3.14×10^{-4} s/s/y) or approximately 2.5 nucleotide substitutions per ORF

(10,296 nucleotides) per year. To our knowledge, it is the first assessment of the

JEV substitution rate confirmed by the formal Bayesian test of temporal signal;

3.   JEV GIII is characterized by the lowest mean substitution rate

(6.17×10^{-5} s/s/y (95% HPD, 3.84×10^{-5}–8.63×10^{-5} s/s/y) or approximately 0.6

nucleotide substitutions per ORF per year) in comparison with GI and the common

JEV clade. However, the passage history of a large number of strains was not taken

into account, hence the true evolutionary rate of JEV GIII. Furthermore, among-

lineage rate variation also plays a significant role. The Monte Carlo simulations

showed a 21–36% posterior probability that GIII substitution rates would exceed

those of GI despite GI having the highest mean;

4.   Time to JEV MRCA (GI-GV clade) has the mean value of 1234 years

(95% HPD, 407–2333 years);

5.   The mean values and appropriate 95% HPD intervals of divergence of

each JEV genotype are comparable and close. Based on the lower and upper

bounds of HPDs, JEV genotypes occurred in the period from 77 to 540 years ago.

The main limitations of our study are that JEV GII, GIV, and GV genotypes

represent an insufficient number of complete or near complete genomes. Therefore,

future studies including more JEV genomes will obtain more reliable and precise

substitution rate estimates given that temporal signal will access in proper way.

Declaration of competing interest

The authors declared no potential conflicts of interest with respect to the

research, author-ship, and/or publication of this article.

References

Ayres DL, Darling A, Zwickl DJ, Beerli P, Holder MT, Lewis PO,

Huelsenbeck JP, Ronquist F, Swofford DL, Cummings MP, Rambaut A, Suchard

MA (2012) BEAGLE: an application programming interface and high-performance

computing library for statistical phylogenetics. Syst Biol 61:170-173.

https://doi.org/10.1093/sysbio/syr100

Bouckaert R, Vaughan TG, Barido-Sottani J, Duchene S, Fourment M,

Gavryushkina A, Heled J, Jones G, Kuhnert D, De Maio N, Matschiner M, Mendes

FK, Muller NF, Ogilvie HA, du Plessis L, Popinga A, Rambaut A, Rasmussen D,

Siveroni I, Suchard MA, Wu CH, Xie D, Zhang C, Stadler T, Drummond AJ

(2019) BEAST 2.5: An advanced software platform for Bayesian evolutionary

analysis.   PLoS   Comput.   Biol   15:e1006650.

https://doi.org/10.1371/journal.pcbi.1006650

Crits-Christoph A, Levy JI, Pekar JE, Goldstein SA, Singh R, Hensel Z,

Gangavarapu K, Rogers MB, Moshiri N, Garry RF, Holmes EC, Koopmans MPG,

Lemey P, Peacock TP, Popescu S, Rambaut A, Robertson DL, Suchard MA,

Wertheim JO, Rasmussen AL, Andersen KG, Worobey M, Debarre F (2024)

Genetic tracing of market wildlife and viruses at the epicenter of the COVID-19

pandemic. Cell 187:5468-5482 e5411. https://doi.org/10.1016/j.cell.2024.08.010

Delatorre E, de Abreu FVS, Ribeiro IP, Gomez MM, Dos Santos AAC,

Ferreira-de-Brito A, Neves M, Bonelly I, de Miranda RM, Furtado ND, Raphael

LMS, da Silva LFF, de Castro MG, Ramos DG, Romano APM, Kallas EG, Vicente

ACP, Bello G, Lourenco-de-Oliveira R, Bonaldo MC (2019) Distinct YFV

Lineages Co-circulated in the Central-Western and Southeastern Brazilian Regions

From   2015   to   2018.   Front   Microbiol   10:1079.

https://doi.org/10.3389/fmicb.2019.01079

Douglas J, Zhang R, Bouckaert R (2021) Adaptive dating and fast proposals:

Revisiting the phylogenetic relaxed clock model. PLoS Comput Biol 17:e1008322.

https://doi.org/10.1371/journal.pcbi.1008322

Duchene S, Holmes EC, Ho SY (2014) Analyses of evolutionary dynamics

in viruses are hindered by a time-dependent bias in rate estimates. Proc Biol Sci

281. https://doi.org/10.1098/rspb.2014.0732

Duchene S, Holt KE, Weill FX, Le Hello S, Hawkey J, Edwards DJ,

Fourment M, Holmes EC (2016) Genome-scale rates of evolutionary change in

bacteria. Microb Genom 2:e000094. https://doi.org/10.1099/mgen.0.000094

Duchene S, Lemey P, Stadler T, Ho SYW, Duchene DA, Dhanasekaran V,
Baele G (2020) Bayesian Evaluation of Temporal Signal in Measurably Evolving
Populations.   Molecular   Biology   and   Evolution.
https://doi.org/10.1093/molbev/msaa163
Duffy S, Shackelton LA, Holmes EC (2008) Rates of evolutionary change in
viruses: patterns   and   determinants.   Nat   Rev   Genet   9:267-276.
https://doi.org/10.1038/nrg2323
Engel D, Jost H, Wink M, Borstler J, Bosch S, Garigliany MM, Jost A,
Czajka C, Luhken R, Ziegler U, Groschup MH, Pfeffer M, Becker N, Cadar D,
Schmidt-Chanasit J (2016) Reconstruction of the Evolutionary History and
Dispersal of Usutu Virus, a Neglected Emerging Arbovirus in Europe and Africa.
mBio 7:e01938-01915. https://doi.org/10.1128/mBio.01938-15
Featherstone LA, Rambaut A, Duchene S, Wirth W (2024) Clockor2:
Inferring global and local strict molecular clocks using root-to-tip regression. Syst
Biol. https://doi.org/10.1093/sysbio/syae003
Ferreira MAR, Suchard MA (2010) Bayesian analysis of elapsed times in
continuous‐time Markov chains. Canadian Journal of Statistics 36:355-368.
https://doi.org/10.1002/cjs.5550360302
Gao X, Liu H, Li M, Fu S, Liang G (2015) Insights into the evolutionary
history of Japanese encephalitis virus (JEV) based on whole-genome sequences
comprising the five genotypes. Virol J 12:43. https://doi.org/10.1186/s12985-015-
0270-z
Guo Q, Zhu S, Wang D, Li X, Zhu H, Song Y, Liu X, Xiao F, Zhao H, Lu H,
Xiao J, Yu L, Wang W, He Y, Liu Y, Li J, Zhang Y, Xu W, Yan D (2023) Genetic
characterization and molecular evolution of type 3 vaccine-derived polioviruses
from   an   immunodeficient   patient   in   China.   Virus   Res   334:199177.
https://doi.org/10.1016/j.virusres.2023.199177
Ho SYW (2020) The Molecular Clock and Evolutionary Rates Across the
Tree of Life. In: Ho SYW (ed) The Molecular Evolutionary Clock. Springer Cham

Holmes EC, Dudas G, Rambaut A, Andersen KG (2016) The evolution of
Ebola virus: Insights from the 2013-2016 epidemic. Nature 538:193-200.
https://doi.org/10.1038/nature19790

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS
(2017) ModelFinder: fast model selection for accurate phylogenetic estimates. Nat.
Methods 14:587-589. https://doi.org/10.1038/nmeth.4285

Kass RE, Raftery AE (1995) Bayes Factors. Journal of the American
Statistical Association 90:773-795.
https://doi.org/10.1080/01621459.1995.10476572

Katoh K, Standley DM (2013) MAFFT multiple sequence alignment
software version 7: improvements in performance and usability. Mol Biol Evol
30:772-780. https://doi.org/10.1093/molbev/mst010

Ladreyt H, Durand B, Dussart P, Chevalier V (2019) How Central Is the
Domestic Pig in the Epidemiological Cycle of Japanese Encephalitis Virus? A
Review of Scientific Evidence and Implications for Disease Control. Viruses 11.
https://doi.org/10.3390/v11100949

Larsson A (2014) AliView: a fast and lightweight alignment viewer and
editor for large data sets. Bioinformatics 30:3276-3278.
https://doi.org/10.1093/bioinformatics/btu531

Li F, Feng Y, Wang G, Zhang W, Fu S, Wang Z, Yin Q, Nie K, Yan J, Deng
X, He Y, Liang L, Xu S, Wang Z, Liang G, Wang H (2023) Tracing the
spatiotemporal phylodynamics of Japanese encephalitis virus genotype I
throughout Asia and the western Pacific. PLoS Negl Trop Dis 17:e0011192.
https://doi.org/10.1371/journal.pntd.0011192

Li G, Li X, Chen J, Lemey P, Vrancken B, Su S, Dellicour S, Gámbaro F
(2025) Tracing more than two decades of Japanese encephalitis virus circulation in
mainland China. Journal of Virology 99:e01575-01524.
https://doi.org/doi:10.1128/jvi.01575-24

Liu RF, He ZJ, Mei P, Xi JC, Cao XD, Yuan LH, Lu JH (2019) Risk
assessment and genomic characterization of Zika virus in China and its

surrounding   areas.   Chin   Med   J   (Engl)   132:1645-1653.

https://doi.org/10.1097/CM9.0000000000000317

Mackenzie JS (2005) Emerging zoonotic encephalitis viruses: lessons from

Southeast   Asia   and   Oceania.   J   Neurovirol   11:434-440.

https://doi.org/10.1080/13550280591002487

Martin DP, Varsani A, Roumagnac P, Botha G, Maslamoney S, Schwab T,

Kelz Z, Kumar V, Murrell B (2021) RDP5: a computer program for analyzing

recombination in, and removing signals of recombination from, nucleotide

sequence datasets. Virus Evol 7:veaa087. https://doi.org/10.1093/ve/veaa087

May FJ, Davis CT, Tesh RB, Barrett ADT (2011) Phylogeography of West

Nile Virus: from the Cradle of Evolution in Africa to Eurasia, Australia, and the

Americas. Journal of Virology 85:2964-2974. https://doi.org/10.1128/jvi.01963-10

McMullen AR, Albayrak H, May FJ, Davis CT, Beasley DWC, Barrett ADT

(2013) Molecular evolution of lineage 2 West Nile virus. J Gen Virol 94:318-325.

https://doi.org/10.1099/vir.0.046888-0

Mohammed MA, Galbraith SE, Radford AD, Dove W, Takasaki T, Kurane I,

Solomon T (2011) Molecular phylogenetic and evolutionary analyses of Muar

strain of Japanese encephalitis virus reveal it is the missing fifth genotype. Infect

Genet Evol 11:855-862. https://doi.org/10.1016/j.meegid.2011.01.020

Moreira Salles AP, de Seixas Santos Nastri AC, Ho YL, Vilas Boas Casadio

L, Emanuel Amgarten D, Justo Arevalo S, Soares Gomes-Gouvea M, Jose Carrilho

F, de Mello Malta F, Rebello Pinho JR (2022) Updating the Phylodynamics of

Yellow Fever Virus 2016-2019 Brazilian Outbreak With New 2018 and 2019 Sao

Paulo   Genomes.   Front   Microbiol   13:811318.

https://doi.org/10.3389/fmicb.2022.811318

Moureau G, Cook S, Lemey P, Nougairede A, Forrester NL, Khasnatinov M,

Charrel RN, Firth AE, Gould EA, de Lamballerie X (2015) New insights into

flavivirus evolution, taxonomy and biogeographic history, extended by analysis of

canonical   and   alternative   coding   sequences.   PLoS   One   10:e0117849.

https://doi.org/10.1371/journal.pone.0117849

O'Toole A, Neher RA, Ndodo N, Borges V, Gannon B, Gomes JP, Groves N,
King DJ, Maloney D, Lemey P, Lewandowski K, Loman N, Myers R, Omah IF,
Suchard MA, Worobey M, Chand M, Ihekweazu C, Ulaeto D, Adetifa I, Rambaut
A (2023) APOBEC3 deaminase editing in mpox virus as evidence for sustained
human transmission since at least 2016. Science 382:595-600.
https://doi.org/10.1126/science.adg8116

Pan XL, Liu H, Wang HY, Fu SH, Liu HZ, Zhang HL, Li MH, Gao XY,
Wang JL, Sun XH, Lu XJ, Zhai YG, Meng WS, He Y, Wang HQ, Han N, Wei B,
Wu YG, Feng Y, Yang DJ, Wang LH, Tang Q, Xia G, Kurane I, Rayner S, Liang
GD (2011) Emergence of genotype I of Japanese encephalitis virus as the dominant
genotype in Asia. J Virol 85:9847-9853. https://doi.org/10.1128/JVI.00825-11

Postler TS, Beer M, Blitvich BJ, Bukh J, de Lamballerie X, Drexler JF, Imrie
A, Kapoor A, Karganova GG, Lemey P, Lohmann V, Simmonds P, Smith DB,
Stapleton JT, Kuhn JH (2023) Renaming of the genus Flavivirus to Orthoflavivirus
and extension of binomial species names within the family Flaviviridae. Arch Virol
168:224. https://doi.org/10.1007/s00705-023-05835-1

Rambaut A, Drummond AJ, Dong X, Baele G, Suchard MA (2018) Posterior
summarization in Bayesian phylogenetics using tracer 1.7. Syst. Biol. 67:901-904.
https://doi.org/10.1093/sysbio/syy032

Rambaut A, Lam TT, Max Carvalho L, Pybus OG (2016) Exploring the
temporal structure of heterochronous sequences using TempEst (formerly Path-O-
Gen). Virus Evol. 2:vew007. https://doi.org/10.1093/ve/vew007

Ravanini P, Huhtamo E, Ilaria V, Crobu MG, Nicosia AM, Servino L, Rivasi
F, Allegrini S, Miglio U, Magri A, Minisini R, Vapalahti O, Boldorini R (2012)
Japanese encephalitis virus RNA detected in Culex pipiens mosquitoes in Italy.
Euro Surveill 17. https://doi.org/10.2807/ese.17.28.20221-en

Ricklin ME, Garcia-Nicolas O, Brechbuhl D, Python S, Zumkehr B,
Nougairede A, Charrel RN, Posthaus H, Oevermann A, Summerfield A (2016)
Vector-free transmission and persistence of Japanese encephalitis virus in pigs. Nat
Commun 7:10832. https://doi.org/10.1038/ncomms10832

Schuh AJ, Ward MJ, Brown AJ, Barrett AD (2013) Phylogeography of

Japanese encephalitis virus: genotype is associated with climate. PLoS Negl Trop

Dis 7:e2411. https://doi.org/10.1371/journal.pntd.0002411

Sikazwe C, Neave MJ, Michie A, Mileto P, Wang J, Cooper N, Levy A,

Imrie A, Baird RW, Currie BJ, Speers D, Mackenzie JS, Smith DW, Williams DT

(2022) Molecular detection and characterisation of the first Japanese encephalitis

virus belonging to genotype IV acquired in Australia. PLoS Negl Trop Dis

16:e0010754. https://doi.org/10.1371/journal.pntd.0010754

Simon-Loriere E, Faye O, Prot M, Casademont I, Fall G, Fernandez-Garcia

MD, Diagne MM, Kipela JM, Fall IS, Holmes EC, Sakuntabhai A, Sall AA (2017)

Autochthonous Japanese Encephalitis with Yellow Fever Coinfection in Africa. N

Engl J Med 376:1483-1485. https://doi.org/10.1056/NEJMc1701600

Sistrom M, Andrews H, Edwards DL (2024) Comparative genomics of

Japanese encephalitis virus shows low rates of recombination and a small subset of

codon positions under episodic diversifying selection. PLoS Negl Trop Dis

18:e0011459. https://doi.org/10.1371/journal.pntd.0011459

Srihi H, Chatti N, Ben Mhadheb M, Gharbi J, Abid N (2021) Phylodynamic

and phylogeographic analysis of the complete genome of the West Nile virus

lineage 2 (WNV-2) in the Mediterranean basin. BMC Ecol Evol 21:183.

https://doi.org/10.1186/s12862-021-01902-w

Srivastava KS, Jeswani V, Pal N, Bohra B, Vishwakarma V, Bapat AA,

Patnaik YP, Khanna N, Shukla R (2023) Japanese Encephalitis Virus: An Update

on the Potential Antivirals and Vaccines. Vaccines (Basel) 11.

https://doi.org/10.3390/vaccines11040742

Stica CJ, Barrero RA, Murray RZ, Devine GJ, Phillips MJ, Frentiu FD

(2022) Global Evolutionary History and Dynamics of Dengue Viruses Inferred

from Whole Genome Sequences. Viruses 14. https://doi.org/10.3390/v14040703

Sumiyoshi H, Mori C, Fuke I, Morita K, Kuhara S, Kondou J, Kikuchi Y,

Nagamatu H, Igarashi A (1987) Complete nucleotide sequence of the Japanese

encephalitis   virus   genome   RNA.   Virology   161:497-510.

https://doi.org/10.1016/0042-6822(87)90144-9

Sun JW, D., Zhong H, Guan D, Zhang H, Tan Q, Zhou H, Zhang M, Ning D,

Zhang B, Ke C, Song T, Lin J, Zhang Y, Koopmans M, Gao GF (2017) Returning

ex-patriot Chinese to Guangdong, China, increase the risk for local transmission of

Zika virus. J Infect 75:356-367. https://doi.org/10.1016/j.jinf.2017.07.001

Takhampunya R, Kim HC, Tippayachai B, Kengluecha A, Klein TA, Lee

WJ, Grieco J, Evans BP (2011) Emergence of Japanese encephalitis virus genotype

V in the Republic of Korea. Virol J 8:449. https://doi.org/10.1186/1743-422X-8-

449

Theze J, Li T, du Plessis L, Bouquet J, Kraemer MUG, Somasekar S, Yu G,

de Cesare M, Balmaseda A, Kuan G, Harris E, Wu CH, Ansari MA, Bowden R,

Faria NR, Yagi S, Messenger S, Brooks T, Stone M, Bloch EM, Busch M, Munoz-

Medina JE, Gonzalez-Bonilla CR, Wolinsky S, Lopez S, Arias CF, Bonsall D, Chiu

CY, Pybus OG (2018) Genomic Epidemiology Reconstructs the Introduction and

Spread of Zika Virus in Central America and Mexico. Cell Host Microbe 23:855-

864 e857. https://doi.org/10.1016/j.chom.2018.04.017

Van den Eynde C, Sohier C, Matthijs S, De Regge N (2022) Japanese

Encephalitis Virus Interaction with Mosquitoes: A Review of Vector Competence,

Vector   Capacity   and   Mosquito   Immunity.   Pathogens   11.

https://doi.org/10.3390/pathogens11030317

Vannice KS, Hills SL, Schwartz LM, Barrett AD, Heffelfinger J, Hombach J,

Letson GW, Solomon T, Marfin AA, Japanese encephalitis vaccination experts p

(2021) The future of Japanese encephalitis vaccination: expert recommendations

for achieving and maintaining optimal JE control. NPJ Vaccines 6:82.

https://doi.org/10.1038/s41541-021-00338-z

Williams DT, Diviney SM, Niazi AU, Durr PA, Chua BH, Herring B, Pyke

A, Doggett SL, Johansen CA, Mackenzie JS (2015) The Molecular Epidemiology

and Evolution of Murray Valley Encephalitis Virus: Recent Emergence of Distinct

Sub-lineages of the Dominant Genotype 1. PLoS Negl Trop Dis 9:e0004240.

https://doi.org/10.1371/journal.pntd.0004240

Xu G, Gao T, Wang Z, Zhang J, Cui B, Shen X, Zhou A, Zhang Y, Zhao J,

Liu H, Liang G (2023) Re-Emerged Genotype IV of Japanese Encephalitis Virus Is

the Youngest Virus in Evolution. Viruses 15. https://doi.org/10.3390/v15030626

Zecchin B, Fusaro A, Milani A, Schivo A, Ravagnan S, Ormelli S, Mavian

C, Michelutti A, Toniolo F, Barzon L, Monne I, Capelli G (2021) The central role

of Italy in the spatial spread of USUTU virus in Europe. Virus Evol 7:veab048.

https://doi.org/10.1093/ve/veab048