Dynamic Gamma Modulation of Hippocampal Place Cells

Predominates Development of Theta Sequences

Ning Wang[1], Yimeng Wang[1], Mingkun Guo[1], Ling Wang[1,2,3], Xueling Wang[1], Nan

Zhu[1], Jiajia Yang[1,2,3], Lei Wang[4*], Chenguang Zheng[1,2,3,5*], Dong Ming[1,2,3*]

1. Academy of Medical Engineering and Translational Medicine, Medical College,

Tianjin University, Tianjin, China

2. Tianjin Key Laboratory of Brain Science and Neuroengineering, Tianjin, China

3. Haihe Laboratory of Brain-Computer Interaction and Human-Machine Integration,

Tianjin, China

4. School of Statistics and Data Science, Nankai University, Tianjin, China

5. Lead contact

* Correspondence: L.W. (lwangstat@nankai.edu.cn) or C.Z. (cgzheng@tju.edu.cn) or

D.M. (richardming@tju.edu.cn)

ABSTRACT

The experience-dependent spatial cognitive process requires sequential

organization of hippocampal neural activities by theta rhythm, which develops to

represent highly compressed information for rapid learning. However, how the theta

sequences were developed in a finer time scale within theta cycles remains unclear.

In this study, we found that sweep-ahead structure of theta sequences developing

with exploration was predominantly dependent on a relatively large proportion of FG-

cells, i.e. a subset of place cells dominantly phase-locked to fast gamma rhythms.

These ensembles integrated compressed spatial information by cells consistently

firing at precessing slow gamma phases within the theta cycle. Accordingly, the

sweep-ahead structure of FG-cell sequences was positively correlated with the

intensity of slow gamma phase precession, in particular during early development of

theta sequences. These findings highlight the dynamic network-modulation by fast

and slow gamma in the development of theta sequences which may further facilitate

memory encoding and retrieval.

Keywords: Hippocampus, Place cells, Theta sequences, Gamma rhythms,

Sequence development

INTRODUCTION

The hippocampus is known as its central role in encoding and storing experience and thus is essential for forming episodic memories (Buzsaki, McKenzie, & Davachi, 2022; Cohen & Eichenbaum, 1993). In the context of active navigation in an environment, hippocampal place cells are organized to fire sequentially within individual cycles of theta oscillations (4-12Hz) and form time-compressed representations of behavioral experiences, referred to as theta sequences (Colgin, 2020; Comrie, Frank, & Kay, 2022; Dragoi & Buzsaki, 2006; W. E. Skaggs, McNaughton, Wilson, & Barnes, 1996; Wallis, 2018). By sweeping ahead from the past to the upcoming locations, theta sequences represent information about potential future paths that could guide goal-directed behaviors (Gupta, van der Meer, Touretzky, & Redish, 2012; Joshi et al., 2023; Zheng, Hwaun, Loza, & Colgin, 2021). These predictive theta sequences exhibited experience-dependent development when rats were exploring a novel environment or learning a novel goal location (Feng, Silva, & Foster, 2015; Gobbo et al., 2022; Wikenheiser & Redish, 2015; Zheng et al., 2021). However, how the hippocampal network modulated the development of theta sequences' temporospatial-compressed organization remains incompletely understood.

A common assumption was established by previous studies that theta sequences were manifestations of theta phase precession, in which an individual place cell fired at progressively earlier theta phase as the animal traversed the place field (Aoki et al., 2023; Foster & Wilson, 2007; O'Keefe & Recce, 1993; Qasim, Fried, & Jacobs, 2021; W. E. Skaggs et al., 1996). However, when examined the theta sequences with experience at single trial level, their sequential structures were not developed on the first experience of a novel environment even if the phase

precession of individual place cells has been observed (Colgin, 2020; Feng et al.,

2015). This suggests that the dynamic process of theta sequences development

could be modulated by another network-level mechanism with neuronal firing

organized at a finer timescale within theta cycles. Gamma (25-100Hz) could be a

candidate that is thought to interact with theta to temporally organize theta

sequences (Dragoi & Buzsaki, 2006; Fernandez-Ruiz, Sirota, Lopes-Dos-Santos, &

Dupret, 2023; Lisman & Jensen, 2013). As two subtypes of gamma, fast and slow

gamma rhythms (Colgin et al., 2009; Schomburg et al., 2014; Zhang, Lee, Rozell, &

Singer, 2019) occur at distinct theta phases and coordinate theta sequences in

different manners (Zheng, Bieri, Hsiao, & Colgin, 2016). This indicates that fast and

slow gamma may act as different tuners but collaborate together in modulating theta

sequence development.

Fast gamma (~65-100Hz), associated with the input from the medial entorhinal

cortex, is thought to rapidly encode ongoing novel information in the context

(Fernandez-Ruiz et al., 2021; Kemere, Carr, Karlsson, & Frank, 2013; Zheng et al.,

2016). During fast gamma rhythms, place cell spikes occurred across all positions

within a place field and displayed theta phase precession (Bieri, Bobbitt, & Colgin,

2014). This type of cells was identified recently as "phase-precessing" cells, in which

the phase-precession pattern emerges with strong fast gamma (Guardamagna,

Stella, & Battaglia, 2023). Their discrete distribution of theta phases is likely to

contribute to the sequential structure formation when the sequences have been

developed (M. Wang, Foster, & Pfeiffer, 2020). However, a question is raised about

how these fast gamma-modulated cells be coordinated for the developmental

process of theta sequences.

Slow gamma (~25-45Hz) in the hippocampal CA1 reflects information received

from CA3, which is responsible for the integration processing of learned information

and experience (Colgin et al., 2009; Zhu et al., 2023). In contrast to fast gamma,

spiking during slow gamma exhibited dominant theta phase locking and attenuated

theta phase precession (Guardamagna et al., 2023; Zhang et al., 2019). This

concentrated distribution of theta phase could be interpreted as they are coordinated

within slow gamma cycles, as a finer time-scale than a theta cycle. Place cells

spiking exhibited slow gamma phase precession, to generate mini-sequences in an

individual theta sequence, for highly time-compressed representation of ongoing and

future information (Zheng et al., 2016; Zheng et al., 2021). Thus, how fast and slow

gamma rhythms coordinate place cells spiking to modulate the dynamic process of

theta sequence development remains unclear.

Here, we proposed a scheme that fast gamma rhythms coordinate a subgroup

of place cells for rapidly encoding and processing sensory information since from the

early phase of theta sequence development (Figure 1). The developing process

would be then dependent on the slow gamma modulation, that the spikes are fired at

slow gamma phase-precessed pattern for spatial information compression within

theta cycles. To test this hypothesis, we investigated how fast and slow gamma

separately modulated place cells activities during the whole process of theta

sequence development. We found that the fast gamma phase-locked place cells

(FG-cells) were crucial for the development of theta sequences, without which the

temporospatially compressed structure of theta sequences could not be developed.

Furthermore, these cells exhibited slow gamma phase precession within theta cycles

associated with the sweep-ahead structure of theta sequences, which occurred

earlier than those cells not phase locked to fast gamma (NFG-cells). Our findings

highlight the dynamic network-modulation by neural oscillations fast and slow

gamma, at finer timescales within theta, in development of theta sequences which

may further facilitate memory encoding and retrieval.

RESULTS

A subgroup of hippocampal place cells phase locked to fast gamma rhythms

during active behaviors

To investigate the temporal organization of place cells' firing associated with the

gamma rhythms during the development of theta sequences, we trained 4 Long-

Evans rats traversed unidirectionally on a circular track, with 5 trials (laps) in each

recording session (day). Meanwhile, neuronal spiking of place cells and local field

potentials were recorded simultaneously in hippocampal CA1 (Figure 2-figure

supplement 1) during active running behaviors (Figure 2).

The slow and gamma episodes were detected in time periods when the power

of each gamma type (Figure 2A-B, fast gamma: 65-100 Hz, slow gamma: 25-45 Hz)

was above a preset threshold (see Materials and Methods). We could observe both

slow gamma and fast episodes occurred during active exploration. As shown in a

previous study, 32% of hippocampal pyramidal cells were phase-locked to gamma

rhythms during waking theta states (Senior, Huxter, Allen, O'Neill, & Csicsvari, 2008).

We further tested how the hippocampal place cells were modulated by slow and fast

gamma rhythms respectively, by quantifying the slow or fast gamma phase

distributions of all spikes across all trials (Figure 2B). In this study, 488 place cells

were included with place field distributed on the circular track but not located at the

resting box. We found that 120 (24.6%) out of these cells were significantly phase-

locked to gamma rhythms, a majority of which (113 cells, 23.2%) exhibited significant

phase-locking to fast gamma rhythms (Figure 2C). Their preferred fast gamma

phases (202.38±2.01 deg) were concentrated around the peak of fast gamma cycle,

which was consistent with the findings in Schomburg et al. 2014 (Schomburg et al.,

2014). However, a very small number (3.7%) of cells were found to phase-locked to

slow gamma rhythms. Thus, these place cells were subdivided according to whether

they were significantly phase-locked to fast gamma (FG-cells) or not (NFG-cells)

during active behaviors in the subsequent analyses. Furthermore, the fast gamma

phase-locking of FG-cells was stable across trials, with a significantly higher mean

vector length of fast gamma phase-locking than those of NFG-cells (Figure 2D,

repeated measure ANOVA, main effect of cell type, $F(1,22)=33.9$, $p=7.5 \times 10^{-6}$, partial

$\eta^2 =0.6$; main effect of trials, $F(4,88)=1.3$, $p=0.3$, partial $\eta^2 =0.06$, no cell type x trials

interaction, $F(4,88)=0.5$, $p=0.8$, partial $\eta^2 =0.02$). Meanwhile, we also found that a

larger proportion of neurons in FG-cells had significant theta phase precession than

NFG-cell (Figure 2-figure supplement 2, chi-squared test, ratio of theta phase

precession cell among the FG-cell vs. NFG-cell, $n=457$cell , $\chi^2=13.7$, $p=2.1 \times 10^{-4}$,

Cramer V=0.2). These results suggested that the FG-cells were stably modulated by

fast gamma rhythms during active behaviors.

The FG-cells were crucial for theta sequence development

As theta sequences were developed dependent on experience, we next

investigated whether FG-cells and NFG-cells played different roles in the

development of theta sequences. We then used a Bayesian decoding approach

(see Methods) to identify trajectories represented as rats running on a circular track,

and detected theta sequences in each single lap (Figure 3A). Another two decoders

were built by excluding either FG-cells spikes or down-sampled NFG-cells spikes

respectively, in order to compare the structures of theta sequences without each type

of cells. It can be observed that posterior probability distribution of theta sequences

was changed by excluding spikes from FG-cells (Figure 3B).   The weighted

correlation of each sequence (Feng et al., 2015) was then measured to quantify the

temporospatial correlation of a theta sequence structure.   We found that the

weighted correlation of theta sequences decoded by excluding FG-cells spikes

(exFG-sequences) was much lower than that of theta sequences decoded by

excluding down-sampled NFG-cells spikes (exNFG-sequences) in the same theta

cycles (Figure 3C, generalized linear mixed model, main effect of cell type,

$F(2,7255)=53.0$, p=0; post-hoc test, raw sequences vs. exFG-sequences, p=0, raw

sequences vs. exNFG-sequences, $p=4.6\times10^{-8}$, exFG-sequences vs. exNFG-

sequences, $p=1.1\times10^{-6}$).   This suggests that the sequential structure could be

dominantly interrupted by excluding FG-cells rather than NFG-cells.

Consistent with the previous studies (Feng et al., 2015; M. Wang et al., 2020),

we also observed the development of theta sequences along with exploration

experience (Figure 3D-E, repeated measure ANOVA, main effect of trials,

$F(4,44)=7.0$, $p=2.0\times10^{-4}$, partial $\eta^2=0.4$).   This could not be accounted for by

behavioral parameters, as the rats were running at stable running speed and head

direction across trials (Figure 3F, repeated measure ANOVA, main effect of trials:

$F(4,44)=0.9$, p=0.5 partial η^2 =0.07; Figure 3-figure supplement 1, Watson-Williams

multi-sample test, $F(4,55)=0.2$, p=0.9, partial $\eta^2=$ 0.01).   We further investigated

whether the FG-cells or NFG-cells were required for the development of theta

sequence structure.   Interestingly, the temporospatial structure of the theta

sequences cannot be developed with experience when the FG-cell spikes were

excluded, with significantly lower weighted correlation than theta sequences

constructed by excluding NFG-cell spikes (Figure 3D, G, repeated measure ANOVA,

trials x cell type interaction, $F(4,44)=3.0$, $p=0.03$, partial $\eta^2=0.2$, main effect of cell

type, $F(1,11)=7.8$, $p=0.02$, partial $\eta^2=0.4$; main effect of trials, $F(4,44)=4.0$, $p=0.008$).

Since the theta sequences have been developed after Lap2, we defined the early

development phase of sequences as the first 2 laps and late development phase of

sequences as the last 2 laps of exploration experience, respectively. We found that

the weighted correlation of both exFG-sequences and exNFG-sequences was low

during early development (Figure 3H, paired t-test, $t(24)=0.2$, $p=0.9$, Cohen's

d=0.04). During late development, however, the weighted correlation of exNFG-

sequences was higher than that of exFG-sequences (Figure 3H, paired t-test, $t(24)=$-

3.7, $p=0.001$, Cohen's d=0.8). These results suggest that fast gamma modulation of

place cells may contribute to the development of theta sequence with exploration

experience.

In order to estimate to what extent, the FG cells contributed to theta sequence

development, we investigated the relationship between the percentage of FG-cells

and the formation of sequential structure in different developments. In our dataset,

the percentage of FG-cells out of all place cells varied across days, thus we detected

theta sequences from trials with relatively low percentage (~10%) and high

percentage (~35%) of FG-cells. The number of excluded NFG-cells for decoding

was matched up with the number of excluded FG-cells by a pseudo-random method

in each day. During early development, the temporospatial structure of theta

sequences has not been developed by excluding either low or high percentage of

each type of cells (Figure 4A). During late development, however, the sweeping-

ahead structure of the theta sequence was significantly disrupted when excluding a

high percentage (~35%) of FG-cells, which was not observed when excluding a low

percentage (~10%) of FG-cells (Figure 4B). The sequential structure could be

maintained by excluding NFG-cells at either low or high percentage. To quantify the

relationship between sequence structure and the number of excluded neurons during

early and late development, we then measured the weighted correlation difference

between exFG-sequences and exNFG-sequences as a function of the fraction of

excluded cells. The results showed that the weighted correlation difference was

close to 0 regardless of the fraction of excluded cells during early development

(Figure 4C, Pearson's correlation, Corr=0.1, p=0.7). During late development,

however, we found that the weighted correlation difference was positively correlated

with the fraction of excluded cells (Figure 4D, Pearson's correlation, Corr=0.6,

p=0.03). These results suggest different roles of FG-cells and NFG-cells in the

development of the sweeping-ahead structure of theta sequence with experience,

that the more temporospatially compressed structure of theta sequences may

require a larger percentage of FG-cells.

In addition, we tested whether these results could be accounted for by the

different firing characteristics between FG-cells and NFG-cells. We found that

although these two types of cells took different proportion of all recorded place cells,

the place fields of both types of cells could cover the entire track uniformly (Figure

2C, Figure 5A). We measured the distribution of center of mess (COM) of main

place field of FG-cells and number-matched NFG-cells (Figure 5B), and found similar

distribution of place field COM between cell types (Figure 5C, two-sample

Kolmogorov-Smirnov test, Z=0.1, p=0.8). Furthermore, Although the mean firing rate

of FG-cell was significantly higher than NFG-cell (Figure 5D, Student's t test, t(22) =

2.1, p = 0.04, Cohen's D = 0.9), the theta sequences in the above analyses were

decoded by excluding similar numbers of spikes from FG-cells and NFG-cells

(Figure 5E, excluded spike counts, t(22) = 1.2, p = 0.3, Cohen's D = 0.5;), indicating

that the different development patterns between exFG-sequences and exNFG-

sequences were not due to quality difference between two decoders. The two type

of cells exhibited similar place fields size (Figure 5F, t(22)=0.7, p=0.5, Cohen's D=0.3)

and spatial information (Figure 5G, t(22)=1.0, p=0.3, Cohen's D=0.4).   These

findings support the hypothesis that FG-cells, but not NFG-cells, may contribute to

the development of theta sequences with experience.

241　FG-cells exhibited slow gamma phase precession for information compression

242　required in theta sequence development

In our previous study, we found that the place cell spikes occurred at

progressively earlier slow gamma phases across slow gamma cycles in theta

sequences, so-called slow gamma phase precession (Zheng et al., 2016).   As

sequences of locations were represented at different slow gamma phases, theta

sequences exhibited temporospatially compressed structure representing long paths.

We next investigated if the development of theta sequence structure was related to

the slow gamma phase modulation of place cells. By detecting the slow gamma

phases of spikes within all theta cycles, we observed that the slow gamma phases

significantly shifted across successive slow gamma cycles (Figure 6A, multi-sample

Mardia-Watson-Wheeler test, W(4)=369.4, $p<2.2\times10^{-16}$; post-hoc test, Cycle -1 vs.

Cycle 0, W(2)=97.5, $p<2.2\times10^{-16}$, Cycle 0 vs. Cycle 1, W(2)=95.4, $p<2.2\times10^{-16}$),

which was consistent with the previous finding (Zheng et al., 2016).   Then we

separated out the spike activity from the early and late development, and tested if

the slow gamma phase precession still existed during different developments.

Interestingly, we observed slow gamma phase shifting, but not phase precession,

across successive gamma cycles during early development. The distribution of slow

gamma phases in Cycle0 was similar as that in Cycle-1, then was dispersed in

Cycle1 (Figure 6B, multi-sample Mardia-Watson-Wheeler test, $W(4)=18.5$, $p=9.8\times10^{-4}$;

post-hoc test, Cycle -1 vs. Cycle 0, $W(2)=0.8$, $p=0.7$; Cycle 0 vs. Cycle 1,

$W(2)=15.5$, $p=4.4\times10^{-4}$). However, the slow gamma phase precession existed during

late development (Figure 6C, multi-sample Mardia-Watson-Wheeler test, $W(4)=21.2$,

$p=2.8\times10^{-4}$; post-hoc test, Cycle -1 vs Cycle 0, $W(2) = 8.8$, $p = 0.01$; Cycle 0 vs.

Cycle 1, $W(2)=8.2$, $p=0.02$). This finding suggests that slow gamma phase

precession of place cells may occur with exploring experience.

Furthermore, we investigated how the slow gamma phase precession occurred

in theta sequences developed with experience. We detected theta sequences

during slow gamma episodes and measured the distribution of phase shifts across

gamma cycles within single unites. In the theta sequences during slow gamma

episodes, we observed the slow gamma phase precession from single unit, that the

place cell spikes occurred at earlier slow gamma phases in later slow gamma cycles

(Figure 7A). To quantify the slow gamma phase precession at single unit level, we

calculated the phase shift as the phase difference within spike pairs from adjacent

active slow gamma cycles (See Methods). We found that slow gamma phase

difference was significantly lower than 0 between 2 active slow gamma cycles

(Figure 7B, t-test, $t(11)=-8.8$, $p=2.6\times10^{-6}$, Cohen's D=5.3), as well as among 3 or ≥4

active slow gamma cycles for strict limitation of theta sequences modulated by slow

gamma (3 active cycles, $t(11)=-7.3$, $p=1.5\times10^{-5}$, Cohen's D=4.4; ≥4 active cycles,

$t(8)=-3.6$, $p=0.007$, Cohen's D=2.5). Furthermore, we averaged the slow gamma

phase differences across different cells within a theta sequence and compare their

distribution along with learning trials. In the first lap, we found that the slow gamma

phase differences were not significantly different from their surrogate data (Figure 7C,

median of phase difference from lap1 was in 95% confidence intervals of those from

shuffled data, p=0.3). However, the distribution of slow gamma phase difference

was negatively biased compared with their surrogate data (Figure 7C, median of

phase difference from lap2-5 was beyond the 95% confidence intervals of those from

shuffled data; lap2 p=0.001, lap3 p=0.001, lap4 p=0.002, lap5 p=0.002). These

results can also be replicated by averaging the slow gamma phase differences from

a single unit across successive theta sequences (Figure 7D, lap1 p=0.1, lap2

p=0.001, lap3 p=0.001, lap4 p=0.002, lap5 p=0.01).

These findings raised a question of whether the development of slow gamma

phase presession existed for both FG-cells and NFG-cells. To address this question,

we quantified the distribution of slow gamma phase difference in FG-cells and NFG-

cells respectively. We found that the slow gamma phase difference of FG-cells

consistently exhibited a stable phase precession across trials, independent with

learning experience (Figure 7E, lap1 p=0.007, lap2 p=0.006, lap3 p=0.007, lap4

p=0.002, lap5 p=0.02). In contrast, the slow gamma phase precession of NFG-cells

was experience-dependent, with negative biased distribution of phase difference

occurring from Lap2 (Figure 7E, lap1 p=0.9, lap2 p=0.01, lap3 p=0.001, lap4

p=0.009, lap5 p=0.1). These results suggest that FG-cells and NFG-cells have

different dynamic modulated by slow gamma rhythms during exploration experience.

The slow gamma modulation of FG-cells may occur earlier than that of NFG-cells,

thereby driving the location information to be coded within theta cycles in a highly

temporospatially compressed way.

FG-cells and NFG-cells exhibited different dynamics of slow gamma modulation along with theta sequence development

Finally, we wondered if there was correlation between slow gamma phase precession and the development of theta sequence structure, and how the FG-cells and NFG-cells coordinated during this process. Within each theta sequence, we measured its weighted correlation of the sequence structure, and plot it as a function of slow gamma phase difference across slow gamma cycles. Interestingly, we found most data were distributed in the Quadrant1 (Q1) compared to the shuffled data distributed at the center (Figure 8A-B). The theta sequences with negative slow gamma phase difference and positive weighted correlation were significantly dominant (Figure 8C, the probability of observed data from lap1 exceeded 95% confidence intervals of those from shuffled data, p=0.001). This finding supports our hypothesis that the theta sequence development could be associated with the slow gamma phase precession.

We further investigated the contribution of FG-cells and NFG-cells to the slow gamma modulated theta sequence development during different exploring phases. FG-cell-dominant theta sequence (FG-cell sequence) was defined as the theta sequence with at least one FG-cell fired. And NFG-cell-dominant theta sequence (NFG-cell sequence) was defined as the theta sequence with no FG-cell fired. Importantly, the distribution of FG-cell sequences was significantly biased to Q1 than that of the shuffled data, not only in late development when the theta sequences have been developed, but also in early development (Figure 8D-F, early development, p=0.001; late development, p=0.004). However, for NFG-cell sequences, their distribution was significantly biased to Q1 than that of the shuffled data only in late development (Figure 8G-I bottom, p=0.046), but not in early

development (Figure 8G-I top, p=0.08). Taken together, these results support the

hypothesis that FG-cells and NFG-cells may exhibit different dynamical

characteristics in gamma modulation, thus their coordination may contribute to the

development of theta sequences. FG-cells were consistently modulated by slow

gamma phase precession, which would be precondition for theta sequence

development.

DISCUSSION

In current study, we employed a circular track exploration task to investigate the

different roles of fast and slow gamma rhythms in the coordination of neurons during

the experience-dependent development of theta sequences. We first found that a

subgroup of place cells (FG-cells) that were phase-locked to fast gamma played a

crucial role in the development of theta sequences. By excluding a sufficiently high

proportion of FG-cells, the sweep-ahead structure of theta sequences was

significantly disturbed compared to that of excluding equivalent number of NFG-cells.

These FG-cells exhibited slow gamma phase precession within single theta cycles

throughout the developing process of theta sequences. Finally, we showed positive

correlation between the intensity of slow gamma phase precession and development

of the sweep-ahead structure of theta sequences.

Fast gamma episodes were detected with dominantly high probability as initial

exploration of each day and decreased probability as being familiar to the

environment (Bieri et al., 2014). This could be related to the feedforward

communication from upstream to downstream regions by using fast gamma

frequencies (Vinck et al., 2023). Although the information propagation of visual

stimulation was unlikely synchronized between visual cortex and hippocampal CA1

(Schneider, Tzanou, Uran, & Vinck, 2023), the MEC inputs could also be entrained

by fast gamma to process ongoing sensory information to the hippocampus (Buzsaki

& Schomburg, 2015; Colgin et al., 2009). Thus, the fast gamma phase-locked

ensemble may act as a receiver of novel information that organized its intrinsic

individual neurons in a time-compressed and spatially-ordered sequence to integrate

the information under the mesoscopic cellular substrate modulated by fast gamma

rhythms (Fernandez-Ruiz et al., 2023). These neurons exhibited similar firing

characteristics as those of fast gamma non-phase-locked neurons, however, they

showed higher probability presenting theta phase precession. Therefore, this fast

gamma phase-locked ensemble likely has overlapping with "bimodal cells" involved

in the formation of theta sequence structure including reverse and forward windows

(Chu et al., 2023; M. Wang et al., 2020). In this case, regardless of these neurons,

the sweep-ahead structure of theta sequences could not be developed with learning,

with the extent of structure impairment positively correlated with the proportion of

excluded fast gamma phase-locked neurons.

Theta phase precession has long been recognized as the only cause of forming

the dominant theta sequence structure. The current findings offered some possible

additional insights to this point. Our finding that FG-cells displayed stronger theta

phase precession than NFG-cells was consistent with previous finding

(Guardamagna et al., 2023), that the theta phase precession pattern emerged with

strong fast gamma. Since slow gamma phase precession occurred within theta

cycles, it is hard to consider the contribution of these factors to theta sequences

development, without taking theta phase precession into account. But one should be

noted that the theta sequences could not be developed even if theta phase

precession existed from the very beginning of the exploration (Feng et al., 2015).

These findings suggest that theta phase precession, together with other factors,

impact theta sequence development. But the weight of these factors and their

interactions during memory and learning remain important aspects of further

investigation.

The important role of fast gamma in modulating theta sequence development

was also supported by a specific perturbation experiment. Liu and colleagues

optogenetically entrained the MEC GABAergic cells at an artificial gamma frequency

(53Hz) and observed the loss of phase precession patterns in CA1 place cells, and

thereby the disrupted development of theta sequences (Liu, Todorova, Tang, Oliva, &

Fernandez-Ruiz, 2023). The failure of sequential firing organization during behaviors

impaired the formation of a predictive map related to the highly time-compressed

awake replay sequences executing information storage. Fast gamma was also

found diminished between MEC and CA1 in transgenic model mice of Alzheimer's

Disease, linked with spatial memory impairment in AD as a circuit mechanism(Jun et

al., 2020). On the other hand, our previous study showed that fast gamma-

dominated theta sequences represented shorter trajectories with relative less spatial

information within a theta cycle (Zheng et al., 2016). This is indeed not contradictory

with the current results, as we found that slow gamma, related to information

compression, was also required to modulate fast gamma phase-locked cells during

sequence development. We replicated the results of slow gamma phase precession

at ensemble level (Zheng et al., 2016), and furthermore observed it at late

development, but not early development, of theta sequences. Thus, a scenario

could be drawn out that the network modulator was transitioned from fast gamma to

slow gamma throughout the sequence development with learning.

At single neuron level, we also observed that spikes of an individual place cell

occurred at precessed slow gamma phases across consecutive slow gamma cycles

under the time-scale of a theta window. At a finer scale within a theta cycle, slow

gamma cycles coordinated place cells in a "mini-sequence" way (Zheng et al., 2016),

in order to integrate information of long spatial trajectory related to the memory

retrieval and predictive behaviors.   This negatively-biased phase difference

distribution appeared from lap2, when the sweep-ahead structures of theta

sequences haven't been fully developed, suggesting that slow gamma phase

precession of place cells contributed to the information compression required to the

theta sequence development. More interestingly, we found that slow gamma phase

precession of a subset of neurons was absent at initial exploration when the theta

phase precession has already displayed (Feng et al., 2015). This finding was in line

with a previous study showing that silencing of CA3 output didn't affect theta phase

precession of a single cell, but impaired theta sequence structures (Middleton &

McHugh, 2016). A possibility was that recruitment of PV-interneurons during the

CA3-driven descending phase of the theta cycle, associated with the induction of

slow gamma oscillations in CA1, was found reduced in the absence of CA3 input

(Fernandez-Ruiz et al., 2023; Topolnik & Tamboli, 2022).   We also provided

additional evidence about the relationship between slow gamma phase precession

and theta sequence formation, that the more negative slow gamma phase difference

across cycles, the more sweep-ahead structure represented at late development of

theta sequences.

Our results supported such a possible model of gamma modulation to mediate

theta sequence development (Figure 1). The theta sequences were linked to a

balance between fast and slow gamma modulations, especially in FG-cells, which

displayed both fast gamma phase-locking and slow gamma phase precession. As

shown in previous work (Zheng et al., 2016), slow gamma phase precession

supported spatial information retrieval in a temporally compressed format. Thus, we

expected slow gamma precession in all cells during later laps when spatial

information retrieval occurs. However, during early laps, when novel information was

being encoded, only FG-cells exhibited slow gamma precession. This may suggest

that FG-cells help manage the balance between encoding and retrieval through

different phase-coding mechanisms: fast gamma for real-time encoding and slow

gamma for retrieval. This dual-modulation in FG-cells may be crucial for the

development of theta sequences, allowing them to include both FG and NFG cells in

a well-sweep-ahead pattern during later laps.

It is noteworthy that the subset of neurons (FG-cells) was modulated by both

fast gamma and slow gamma rhythms during the theta sequence development

(Figure 2), which was consistent with previous studies (Fernandez-Ruiz et al., 2017;

Lasztóczi & Klausberger, 2016; Sharif, Tayebi, Buzsaki, Royer, & Fernandez-Ruiz,

2021). These cells are seemingly challenging the above framework. However,

these place cells, associated with both memory encoding and retrieval, seem to

function like memory engram-like representations (Goode, Tanaka, Sahay, &

McHugh, 2020; Josselyn & Tonegawa, 2020; Lamothe-Molina et al., 2022; Pettit, Yap,

Greenberg, & Harvey, 2022). Indeed, the proportion of c-Fos positive place cells

(putative engram cells) had higher proportion being phase-locked to fast gamma

than the c-Fos negative place cells (Tanaka et al., 2018). Another possibility was

that these FG-cells were largely overlapped with the previously defined "fast-firing

neurons", which remained stable during learning process (Grosmark & Buzsáki,

2016). In contrast, the NFG-cells exhibited experience-dependent phase coding,

that they likely contributed to the path representation at the late learning process.

Thus, further studies would be needed to investigate how memory engram cells

contribute to theta sequence development and facilitate spatial memory and

navigation.

MATERALS AND METHODS

Subjects

Four male Long-Evans rats weighing 500 to 700g (~6-12 months old) were used

in this study. Rats were housed in custom-made acrylic cages (40cm×40cm×40cm)

on a reverse light cycle (Light: 8:00pm-8:00am). All waking behavior recordings

were performed during the dark phase of the cycle (i.e., from 8:00am-8:00pm). Rats

were handled prior to recording drive implantation surgery. Rats recovered from

surgery for at least one week before behavioral training resumed. Behavioral

training was initiated after at least one week of recovery from implantation surgery.

Rats were physically deprived during experimental data collection so that their body

weight was maintained at ~90% of their free-feeding body weight. All experiments

were conducted according to the guidelines of the Animal Care and Use Committee

of Tianjin University.

Behavioral paradigm

Following the postsurgery recovery period, rats were trained to run on a circular

track from our previous study (Y. Wang et al., 2024; Zheng et al., 2021). In each

recording day (session), rats ran 5 trials (or laps) on the track (inner diameter of 90

cm, height of 50 cm, and width of 10 cm, surrounded by a blue curtain) without

receiving any food reward. Each trial was starting and ending at a wooden rest box

attached to the track (starting point: 33 cm, ending point: 281 cm away from the box).

Surgery and tetrode implanting

Multi-tetrode drives containing 21 independently moveable tetrodes (18 recording tetrodes and 3 reference tetrodes) were implanted into the hippocampal dorsal CA1 region (anterior-posterior [AP] -3.5 mm, medial-lateral [ML] 3.0mm, dorsal-ventral [DV] 1mm on day of implantation). Bone screws were placed in the skull, and the screws along with the base of the drive were covered with dental cement to affix the drive to the skull. Two screws above the cerebellum in the skull were connected to the recording drive ground. Before surgery, tetrodes were built from 17µm polyimide-coated platinum-iridium (10/90%) wire (California Fine Wire, Grover Beach, California). The tips of tetrodes designated for single-unit recording were plated with platinum to reduce single-channel impedances under 300 kOhms. Tetrodes were gradually lowered to their target locations. Recording tetrodes were targeted to the dCA1 stratum pyramidale. One reference tetrode from each region was designated as the reference for differential recording and remained in a quiet area of the cortex throughout the experiments. This reference tetrode was moved up and down until a quiet location was found and was continuously recorded against ground to ensure that it remained quiet throughout data collection. The other reference tetrode in the dHPC was placed in the dCA1 apical dendritic layer to monitor and record local field potentials (LFPs) in the hippocampus during placement of the other tetrodes and to later obtain simultaneous recordings from a dendritic layer.

Data acquisition

Data were acquired using a Digital Lynx system and Cheetah recording software (Neuralynx, Bozeman, Montana). LFPs from one channel within each tetrode were continuously recorded in the 0.1-500 Hz band at a 2,000 Hz sampling rate. Input

amplitude ranges were adjusted before each recording session to optimize resolution

and prevent signal saturation. Input ranges for LFPs were typically within the range

of ±2,000 µV to 3000µV. To detect unit activity, signals from each channel in a

tetrode were bandpass filtered from 600 to 6,000 Hz. The spike was detected when

the signal on one or more of the channels exceeded a threshold set at 55 µV.

Detected events were acquired with a 32,000 Hz sampling rate. Signals were

recorded differentially against a dedicated reference channel (see "Surgery and

recording drive implantation" section above). The video was recorded through the

Neuralynx system with a resolution of 720 × 480 pixels and a frame rate of 25 frames

per second. The animal position was tracked via a red light-emitting-diode (LED) on

one of the three HS-27-Mini-LED headstages (Neuralynx, Bozeman, Montana)

attached to the hyperdrive.

Estimation of running speed and head direction

The running speed (v_t) at a time point (t) was estimated by calculating the

distance between the preceding position (x_{t-1}, y_{t-1}) and the following position (x_{t+1},

y_{t+1}), and dividing by the elapsed time (2×1/position sampling frequency). The

sampling frequency of the position data was 25 Hz, yielding a temporal resolution of

2/25 s (Figure 3F).

The head direction (d_t) at a given time point (t) was calculated as the arctangent

of the vector formed by the preceding (x_{t-1}, y_{t-1}) and the following (x_{t+1}, y_{t+1}) position:

$$d_t = \tan^{-1} \frac{(x_{t+1} - x_{t-1})}{(y_{t+1} - y_{t-1})}$$

Angles were adjusted to the range of 0–360° (Figure 3-figure supplement 1). If

the animal did not move at the time point (t), d_t was set to d_{t-1}.

Spike sorting and turning curve analysis

Detected spikes that occurred during trials were manually sorted using graphical

cluster-cutting software (MClust; A.D. Redish, University of Minnesota, Minneapolis).

Clusters of spikes were sorted using two-dimensional projections of three different

features of spike waveforms (i.e., energies, peaks, and peak-to-valley differences)

from four channels. Individual cells were identified as putative excitatory neurons

that they had pyramidal cell-like waveform shape, at least 1ms refractory period, and

their clusters were perfectly separated with other clusters or noise in at least three

projections. Units with mean firing rates that were higher than 15 Hz were identified

as putative fast-spiking interneurons and were excluded.

The turning curve of each unit was built at 90 angle bins (approximately 3.5cm

per bin) on the circular track by using spikes during locomotion (running speed >5

cm/s, Figure 5). The field size of the main place field was defined as the size of

contiguous position bins with a firing rate > 10% of the peak firing rate. Only units

with peak firing rates over 1 Hz and main field size over 3 angle bins (approximately

10.5cm) during trials were defined as putative neurons with spatial representation. A

putative excitatory neuron, which at least had one place field, was defined as a place

cell. Finally, we accepted 488 place cells for further analysis.

Spatial information score

Spatial information was calculated as previously described (William E Skaggs,

McNaughton, Gothard, & Markus, 1992). Briefly, for each cell, the spatial information

score in bits per spike was calculated from the recording during test trials (Figure 5),

as:

$$\text{Spatial information} = \sum_{i} P_i \frac{\lambda_i}{\lambda} \log_2 \frac{\lambda_i}{\lambda}$$

551   Where λ_i is the mean firing rate of a unit in the ith bin, λ is the overall mean firing

552   rate, and p_i is the probability of the animal being in the ith bin (occupancy in the ith

553   bin/total occupancies bin).

Detection of gamma episodes

LFP signals from each tetrode were band-pass filtered for slow gamma (25-45Hz)

and fast gamma (65-100Hz) rhythms. Time varying power for slow gamma and fast

gamma was computed, using a Morlet's wavelet transform method described

previously (Tallon-Baudry, Bertrand, Delpuech, & Pernier, 1997; Zheng et al., 2016),

and then averaged within each frequency range. To detect slow and fast gamma

episodes, we z-scored the time varying power for slow and fast gamma across time

within each trial. The timestamp of z-scored time varying power ≥ 3 was set as the

center of a slow or fast gamma episode, and a 160ms window around this center

was defined as a gamma episode. For the windows with overlapping time bins, the

one with the lower gamma amplitude was discarded. Also, the slow (or fast) gamma

episode with z-scored fast (or slow) gamma power ≥ 3 was removed, ensuring that

fast and slow gamma episodes are mutually exclusive (Figure 2A-B).

Detection of FG-cells and NFG-cells

The spikes of each place cell recorded during freely running on the track were

included. The time-varying slow or fast phases of local LFPs from the same tetrode

with recorded cells were calculated by Hilbert transform of the bandpass filtered

signals. Each spike time was matched to the closest LFP time stamp to determine

its corresponding slow and fast gamma phase, and the slow and fast gamma phase

distribution was established for each cell (Figure 2B). Mean vector length was

quantified based on the slow and fast gamma phase distributions. Significant phase-

locking to either slow or fast gamma was detected by Rayleigh test on phase

distributions with at least 10 spikes in each lap. Because of a relatively high

proportion of significant fast gamma phase-locked cells, we focused on these cells in

this study and defined them as FG-cells (Figure 2C). Meanwhile, NFG-cells were

defined as place cells which were not significantly phase-locked to fast gamma with

at least 10 spikes in each lap. In addition, regarding the contribution of spike

leakage to the local LFPs, we also selected an individual tetrode which located at

stratum pyramidale and at the center of the drive bundle for each rat. We detected a

similar proportion of FG-cells by using LFPs on this tetrode, compared with that

using local LFPs (Chi-squared test, χ2= 0.9, p=0.4, Cramer V=0.03).

Theta phase precession

LFPs from each tetrode were filtered through a 4-12 Hz bandpass filter. Theta

phases were estimated using angles of their Hilbert-transformed signals for each

channel. Theta phase precession was examined in linearized spatial turning curves

(see Spike sorting and turning curve analysis). For spikes within each place field,

phase precession was computed using a circular-linear fit (Kempter, Leibold, Buzsaki,

Diba, & Schmidt, 2012). The p-value from the circular-linear regression was reported

(Figure 2-figure supplement 2).

Single-trial Bayesian decoding analysis

In the decoding analysis, we included the recording sessions with 44 ± 1 HPC

place cells. To translate ensemble spiking activity into radian positions on the circular

track, the Bayesian decoding algorithm was implemented. For each trial, a

memoryless Bayesian decoder was built at 90 angle bins (ranging from 0 to 2pi) by

using spikes during locomotion (running speed >5 cm/s) in this trial. Decoding was

performed using a 20ms sliding time window with 5ms stepping. For each time

window, the number of spikes occurring is denoted as n_i, so that the vector of all

spiking activity simultaneously recorded for the cell is $\boldsymbol{n} = \{n_1, n_2, \cdots, n_i\}$. The

probability of a rat being at position x, given the number of spikes $\boldsymbol{n}$ from each unit

recorded in a time window, was estimated by Bayes' rule:

$$P(x|\boldsymbol{n}) = \frac{P(\boldsymbol{n}|x) \times P(x)}{P(n)}$$

where $P(\boldsymbol{n}|x)$ was approximated from the single directional firing turning curve of

each unit. The approximation assumed that the turning curve of individual units was

statistically independent and the number of spikes from each unit followed a Poisson

distribution. Prior knowledge of position $P(x)$ was set to 1 to avoid decoding bias to

any particular location on the track. The normalizing constant $P(n)$ was set to

ensure $P(x|\boldsymbol{n})$, or posterior probability, summed to 1 (Figure 3). The decoding

position was is the set of x that maximizes $P(x|\boldsymbol{n})$. For excluded cell decoding,

position reconstruction was performed by removing the FG-cells or NFG-cells from $\boldsymbol{n}$,

respectively. Bayesian decoding analyses were performed using custom routines in

MATLAB 2020b.

Theta sequences detection

Theta rhythms were detected from a central teteode of all recording tetrodes

which located in the stratum pyramidale of hippocampus. We cut sequences at each

theta phases (0°, 10°, 20°, …, 350°) and quantified averaged weighted correlation

corresponding to these 36 possible phases (see "Weighted correlation of theta

sequences" below). Then, the optimal cutting phase of the theta cycle was

determined when the weighted correlation of theta sequences was maximal,

whereby the sequences were separated within each theta cycle (Figure 3A).

Raw theta sequences were detected by the following criteria: (1) during the

sequence, the animal was running between the starting point and ending point of the

track, with running speed higher than 5 cm/s; (2) the duration of the sequence

ranged from 100 to 200 ms; (3) more than 3 place cells which emitted at least 5

spikes within each sequence. In order to investigate the role of FG and NFG cells in

theta sequence development, we decoded sequences without either FG or NFG

cells and cut out the sequences using the same theta cycles as above, defined as

"exFG-sequences" and "exNFG sequences" (Figure 3-4).

To investigate the relationship between sequence development and slow

gamma phase precession, we defined FG-cell sequences and NFG-cell sequences

(Fig8). FG-cell sequences are theta sequences with 1) at least one FG-cell and 2) at

least firing spikes across 3 slow gamma cycles for an individual cell. And the "NFG-

cell-dominant sequences" are theta sequences with 1) only NFG-cells and 2) at least

firing spikes across 3 slow gamma cycles for an individual cell.

Weighted correlation of theta sequences

637   Weighted correlation was calculated to identify sequential structure within a

638 theta cycle. Decoding probabilities (*prob*) were assigned as the weights of the

639 position estimates to calculate the correlation coefficient between the time (*t*) and the

640 decoding position (*p*) as follows:

$$corr_w(t, p; prob) = \frac{cov(t, p; prob)}{\sqrt{cov(t, t; prob)cov(p, p; prob)}}$$

641 and the weighted covariance between time and decoded position $cov(t, p; prob)$ is

642 as follows:

$$cov(t, p; prob) = \frac{\sum_i prob_i (t_i - m(t; prob))(p_i - m(p; prob))}{\sum_i prob_i}$$

643 Weighted means of time and decoded position are as follows:

$$m(t; prob) = \frac{\sum_i prob_i \, t_i}{\sum_i prob_i}$$

644   and

$$m(p; prob) = \frac{\sum_i prob_i\, p_i}{\sum_i prob_i}$$

Weighted correlation of each theta sequence was calculated from the decoded

probabilities of positions 35 cm behind and ahead of the rat's current location, in a

time window of ±30ms (approximate 1⁄4 theta cycle) before and after the mid-time

point of each theta sequence (Figure 3-4). A significant sweep-ahead structure in

the current running direction would produce a large positive correlation, whereas a

lack of sequential structure would produce a correlation close to zero.

Slow gamma phase shift analysis of cell ensemble

Slow gamma phase shift of cell ensemble was estimated as previously

described (Zheng et al., 2016). The gamma cycle with maximal spiking across all

simultaneously recorded cells was defined as cycle 0. Gamma cycles occurring

before cycle 0 are represented by decreasing integer values, after by increasing

integer values. Incomplete cycles at the beginning or end of the sequence or during

immobile (speed<5cm/s) were excluded from analyses. We estimated the gamma

phases and theta phases of spike times for place cells that exhibited spikes within at

least two slow gamma cycles or at least three fast gamma cycles within a single

theta cycle. The number of cycles analyzed within each theta sequence was limited

to three (cycles -1 to 1) for slow gamma. Two-dimensional histograms of gamma

phases and theta phases for spikes from each cycle number were binned into 40

bins and smoothed across nine bins (Figure 6).

Slow gamma phase shift analysis of individual cell

Time-varying slow gamma power (see Detection of gamma episodes) was

averaged within each theta cycle. To identify the theta cycle with high slow gamma

power, we z-scored time-varying slow gamma power across the theta cycle within

each trial. The theta cycle with z-scored time varying power ≥ 1.5 was defined as

slow gamma events.   The global 0 phase of slow gamma, set at the trough,

determined the phase shift. Slow gamma cycles with at least one spike was defined

as active slow gamma cycles (Figure 7A). The slow gamma phase difference of a

theta cycle with k (k ≥ 2) adjacent active slow gamma cycles (active cycle) is as

follows:

$$\text{slow gamma phase difference} = \frac{\sum_{i=2}^{k}(\varphi_i - \varphi_{i-1})}{k-1}$$

where φ_i and φ_{i-1} are respectively slow gamma phase of a spike in i^{th} and $(i-1)^{\text{th}}$

active cycle.   A negative slow gamma phase difference value indicates phase

precession, while a positive value indicates phase procession.

In order to test if the negatively biased distribution was robust for slow gamma

phase precession, we generated surrogate data and compared their distribution with

the real data. A random slow gamma phase within the same cycle was assigned to

each spike to keep its firing location constancy, and the distribution of slow gamma

phase difference was obtained for this mock data. This procedure was repeated

1000 times to get 1000 distributions and the 95% confident interval of their median

(black lines in Figure 7C-E).   In the analysis, we calculated one-sided, corrected

empirical p-values [(r+1)/(n+1)] by comparing real data to its corresponding shuffle

data. Where r is the number values from the shuffled distribution that are either

smaller or larger (depending on the hypothesis) than the observed value and n=1000

is the number of shuffling (Figure 7C-E).

Correlation between slow gamma phase precession and weighted correlation

FG-cell sequences and NFG-cell sequences were included in this analysis (see

"Theta sequences detection" section). For each type of sequences, a scatter plot

was made to visualize the relationship between slow gamma phase precession and

theta sequence structure, with slow gamma phase shift of each sequence on x-axis

and its weighted correlation on y-axis (Figure 8A,D,G). Meanwhile, a 2-dimensional

histogram was measured to show the distribution of sequences in four quadrants

(Figure 8B,E,H). We summed up the probability in Quadrant 1 (Q1) and compared it

with that of the mock data generated in Figure 7. The statistical significance was

determined by calculating the one-sided, corrected empirical p-value, with lower p-

value indicating significantly higher probability in Q1 of real data than that of the

mock data (Figure 8C,F,I).

Histology

At the end of the experiment, histological sections were prepared as follows to

verify the tetrodes position. Rats were intraperitoneally injected with a lethal dose of

urethane solution. The heart was subsequently perfused with phosphate buffer and

formalin to fix the brain tissue. Rat brains were then cut into 30-μm coronal sections

and Nissl Staining was performed to determine the location of tetrode in CA1.

Quantification and statistical analysis

Data analyses were performed using SPSS (IBM), custom Matlab (the Math

Work) and R scripts. Paired t-test was performed to compare weighted correlation

for exFG-sequences and exNFG-sequences during early and late development

(Figure 3H). Student's t test was performed to compare the mean firing rate, spike

counts, place fields size and spatial information for the FG-cell and NFG-cell (Figure

5D-G). One-sample t test was used to test whether the slow gamma phase shift was

significantly lower than 0 (Figure 7B). Repeated measures ANOVA was performed

to test the MVL for FG- and NFG-cell (Figure 2D), weighted correlation of theta

sequences (, Figure 3E,G), and running speed (Figure 3F) across laps. Generalized

linear mixed model was performed to compare weighted correlation among the Raw-,

exFG-, and exNFG-sequence. Chi-squared test was performed to compare the

proportion of neurons had significant theta phase precession in FG-cells and NFG-

cells (Figure 2-figure supplement 2). Kolmogorov-Smirnov test was performed to

test the distribution of place field COM (Figure 5C) on the track using the Matlab

function 'kstest2'. Watson-Williams multi-sample test (circular one-way ANOVA test)

was performed to test the for differences in head directions across laps (Figure 3-

figure supplement 1) using the 'circ_wwtest' function from the CircStat Toolbox. Multi-

sample Mardia-Watson-Wheeler test was performed to test the slow gamma phase

shift of place cells across successive slow gamma cycles (Figure 6) using the R

'circular-package'.

ACKNOWLEDGMENTS

We thank Shuang Meng for help with rat behavioral pre-training; and all Zheng

Lab members for helpful discussions. This work was supported by the grants from

National Science and Technology Innovation 2030 Major Project of China

(2022ZD0205000), National Natural Science Foundation of China (T2322021,

82271218 to C.Z., 12271272 to L.W., 81925020 to D.M., 82371886 to J.Y.,

82202797 to L.W.), The Space Brain Project from Lingang Laboratory (LG-TKN-

202204-01 to J.Y.), and the China Post-Doctoral Science Foundation (2022M712365

to L.W.).

AUTHOR CONTRIBUTIONS

Conceptualization, C.Z., D.M. and L.W.; Methodology and Software, N.W., L.W. and C.Z.; Investigation, N.W., Y.W., M.G., L.W. N.Z., X.W. and C.Z.; Formal Analysis, N.W. and L.W.; Writing – Original Draft, N.W. and C.Z.; Revision, N.W., L.W., D.M. and C.Z.; Funding Acquisition, C.Z., M.D., L.W., J.Y., and L.W.; Supervision, C.Z. and D.M.

COMPETING INTERSETS

The authors declare that they have no competing interests.

REFERENCES

Aoki, Y., Yokoi, T., Morikawa, S., Kuga, N., Ikegaya, Y., & Sasaki, T. (2023). Effects of theta phase precessing optogenetic intervention on hippocampal neuronal reactivation and spatial maps. *iScience, 26*(7), 107233. doi:10.1016/j.isci.2023.107233

Bieri, K. W., Bobbitt, K. N., & Colgin, L. L. (2014). Slow and fast gamma rhythms coordinate different spatial coding modes in hippocampal place cells. *Neuron, 82*(3), 670-681. doi:10.1016/j.neuron.2014.03.013

Buzsaki, G., McKenzie, S., & Davachi, L. (2022). Neurophysiology of Remembering. *Annual Review of Psychology, 73*, 187-215. doi:10.1146/annurev-psych-021721-110002

Buzsaki, G., & Schomburg, E. W. (2015). What does gamma coherence tell us about inter-regional neural communication? *Nature Neuroscience, 18*(4), 484-489. doi:10.1038/nn.3952

Chu, T., Ji, Z., Zuo, J., Mi, Y., Zhang, W., Huang, T., . . . Wu, S. (2023). Firing rate adaptation affords place cell theta sweeps, phase precession and procession. *Elife, 12*, RP87055.

Cohen, N. J., & Eichenbaum, H. (1993). *Memory, amnesia, and the hippocampal system.* Cambridge, MA, US: The MIT Press.

Colgin, L. L. (2020). Five Decades of Hippocampal Place Cells and EEG Rhythms in Behaving Rats. *J Neurosci, 40*(1), 54-60. doi:10.1523/JNEUROSCI.0741-19.2019

Colgin, L. L., Denninger, T., Fyhn, M., Hafting, T., Bonnevie, T., Jensen, O., . . . Moser, E. I. (2009). Frequency of gamma oscillations routes flow of information in the hippocampus. *Nature, 462*(7271), 353-357. doi:10.1038/nature08573

Comrie, A. E., Frank, L. M., & Kay, K. (2022). Imagination as a fundamental function of the hippocampus. *Philos Trans R Soc Lond B Biol Sci, 377*(1866), 20210336. doi:10.1098/rstb.2021.0336

Dragoi, G., & Buzsaki, G. (2006). Temporal encoding of place sequences by hippocampal cell assemblies. *Neuron, 50*(1), 145-157. doi:10.1016/j.neuron.2006.02.023

Feng, T., Silva, D., & Foster, D. J. (2015). Dissociation between the experience-dependent development of hippocampal theta sequences and single-trial phase precession. *Journal of Neuroscience, 35*(12), 4890-

4902. doi:10.1523/JNEUROSCI.2614-14.2015

Fernandez-Ruiz, A., Oliva, A., Nagy, G. A., Maurer, A. P., Berenyi, A., & Buzsaki, G. (2017). Entorhinal-CA3
Dual-Input Control of Spike Timing in the Hippocampus by Theta-Gamma Coupling. *Neuron, 93*(5),
1213-1226 e1215. doi:10.1016/j.neuron.2017.02.017

Fernandez-Ruiz, A., Oliva, A., Soula, M., Rocha-Almeida, F., Nagy, G. A., Martin-Vazquez, G., & Buzsaki, G.
(2021). Gamma rhythm communication between entorhinal cortex and dentate gyrus neuronal
assemblies. *Science, 372*(6537). doi:10.1126/science.abf3119

Fernandez-Ruiz, A., Sirota, A., Lopes-Dos-Santos, V., & Dupret, D. (2023). Over and above frequency: Gamma
oscillations as units of neural circuit operations. *Neuron, 111*(7), 936-953.
doi:10.1016/j.neuron.2023.02.026

Foster, D. J., & Wilson, M. A. (2007). Hippocampal theta sequences. *Hippocampus, 17*(11), 1093-1099.
doi:10.1002/hipo.20345

Gobbo, F., Mitchell-Heggs, R., Tse, D., Al Omrani, M., Spooner, P. A., Schultz, S. R., & Morris, R. G. M.
(2022). Neuronal signature of spatial decision-making during navigation by freely moving rats by using
calcium imaging. *Proc Natl Acad Sci U S A, 119*(44), e2212152119. doi:10.1073/pnas.2212152119

Goode, T. D., Tanaka, K. Z., Sahay, A., & McHugh, T. J. (2020). An Integrated Index: Engrams, Place Cells, and
Hippocampal Memory. *Neuron, 107*(5), 805-820. doi:10.1016/j.neuron.2020.07.011

Grosmark, A. D., & Buzsáki, G. (2016). Diversity in neural firing dynamics supports both rigid and learned
hippocampal sequences. *Science, 351*(6280), 1440-1443. doi:10.1126/science.aad1935

Guardamagna, M., Stella, F., & Battaglia, F. P. (2023). Heterogeneity of network and coding states in mouse
CA1 place cells. *Cell Reports, 42*(2), 112022. doi:10.1016/j.celrep.2023.112022

Gupta, A. S., van der Meer, M. A., Touretzky, D. S., & Redish, A. D. (2012). Segmentation of spatial experience
by hippocampal theta sequences. *Nature Neuroscience, 15*(7), 1032-1039. doi:10.1038/nn.3138

Joshi, A., Denovellis, E. L., Mankili, A., Meneksedag, Y., Davidson, T. J., Gillespie, A. K., . . . Frank, L. M.
(2023). Dynamic synchronization between hippocampal representations and stepping. *Nature,*
617(7959), 125-131. doi:10.1038/s41586-023-05928-6

Josselyn, S. A., & Tonegawa, S. (2020). Memory engrams: Recalling the past and imagining the future. *Science,*
367(6473), eaaw4325. doi:10.1126/science.aaw4325

Jun, H., Bramian, A., Soma, S., Saito, T., Saido, T. C., & Igarashi, K. M. (2020). Disrupted Place Cell
Remapping and Impaired Grid Cells in a Knockin Model of Alzheimer's Disease. *Neuron, 107*(6),
1095-1112 e1096. doi:10.1016/j.neuron.2020.06.023

Kemere, C., Carr, M. F., Karlsson, M. P., & Frank, L. M. (2013). Rapid and continuous modulation of
hippocampal network state during exploration of new places. *PLoS One, 8*(9), e73114.
doi:10.1371/journal.pone.0073114

Kempter, R., Leibold, C., Buzsaki, G., Diba, K., & Schmidt, R. (2012). Quantifying circular-linear associations:
hippocampal phase precession. *J Neurosci Methods, 207*(1), 113-124.
doi:10.1016/j.jneumeth.2012.03.007

Lamothe-Molina, P. J., Franzelin, A., Beck, L., Li, D., Auksutat, L., Fieblinger, T., . . . Oertner, T. G. (2022).
DeltaFosB accumulation in hippocampal granule cells drives cFos pattern separation during spatial
learning. *Nat Commun, 13*(1), 6376. doi:10.1038/s41467-022-33947-w

Lasztóczi, B., & Klausberger, T. (2016). Hippocampal Place Cells Couple to Three Different Gamma
Oscillations during Place Field Traversal. *Neuron, 91*(1), 34-40. doi:10.1016/j.neuron.2016.05.036

Lisman, J. E., & Jensen, O. (2013). The theta-gamma neural code. *Neuron, 77*(6), 1002-1016.
doi:10.1016/j.neuron.2013.03.007

Liu, C., Todorova, R., Tang, W., Oliva, A., & Fernandez-Ruiz, A. (2023). Associative and predictive
hippocampal codes support memory-guided behaviors. *Science, 382*(6668), eadi8237.
doi:10.1126/science.adi8237

Middleton, S. J., & McHugh, T. J. (2016). Silencing CA3 disrupts temporal coding in the CA1 ensemble. *Nature
Neuroscience, 19*(7), 945-951. doi:10.1038/nn.4311

O'Keefe, J., & Recce, M. L. (1993). Phase relationship between hippocampal place units and the EEG theta
rhythm. *Hippocampus, 3*(3), 317-330. doi:10.1002/hipo.450030307

Pettit, N. L., Yap, E. L., Greenberg, M. E., & Harvey, C. D. (2022). Fos ensembles encode and shape stable
spatial maps in the hippocampus. *Nature, 609*(7926), 327-334. doi:10.1038/s41586-022-05113-1

Qasim, S. E., Fried, I., & Jacobs, J. (2021). Phase precession in the human hippocampus and entorhinal cortex.
Cell, 184(12), 3242-3255 e3210. doi:10.1016/j.cell.2021.04.017

Schneider, M., Tzanou, A., Uran, C., & Vinck, M. (2023). Cell-type-specific propagation of visual flicker. *Cell
Rep, 42*(5), 112492. doi:10.1016/j.celrep.2023.112492

Schomburg, E. W., Fernandez-Ruiz, A., Mizuseki, K., Berenyi, A., Anastassiou, C. A., Koch, C., & Buzsaki, G.
(2014). Theta phase segregation of input-specific gamma patterns in entorhinal-hippocampal networks.
Neuron, 84(2), 470-485. doi:10.1016/j.neuron.2014.08.051

Senior, T. J., Huxter, J. R., Allen, K., O'Neill, J., & Csicsvari, J. (2008). Gamma oscillatory firing reveals
distinct populations of pyramidal cells in the CA1 region of the hippocampus. *Journal of Neuroscience,
28*(9), 2274-2286. doi:10.1523/JNEUROSCI.4669-07.2008

Sharif, F., Tayebi, B., Buzsaki, G., Royer, S., & Fernandez-Ruiz, A. (2021). Subcircuits of Deep and Superficial
CA1 Place Cells Support Efficient Spatial Coding across Heterogeneous Environments. *Neuron, 109*(2),
363-376 e366. doi:10.1016/j.neuron.2020.10.034

Skaggs, W. E., McNaughton, B. L., Gothard, K. M., & Markus, E. J. (1992). *An information-theoretic approach
to deciphering the hippocampal code*. Paper presented at the Proceedings of the 5th International
Conference on Neural Information Processing Systems.

Skaggs, W. E., McNaughton, B. L., Wilson, M. A., & Barnes, C. A. (1996). Theta phase precession in
hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus, 6*(2),
149-172. doi:10.1002/(SICI)1098-1063(1996)6:2<149::AID-HIPO6>3.0.CO;2-K

Tallon-Baudry, C., Bertrand, O., Delpuech, C., & Pernier, J. (1997). Oscillatory γ-band (30-70 Hz) activity
induced by a visual search task in humans. *The Journal of Neuroscience, 17*(2), 722-734.
doi:10.1523/jneurosci.17-02-00722.1997

Tanaka, K. Z., He, H., Tomar, A., Niisato, K., Huang, A. J. Y., & McHugh, T. J. (2018). The hippocampal
engram maps experience but not place. *Science, 361*(6400), 392-397. doi:10.1126/science.aat5397

Topolnik, L., & Tamboli, S. (2022). The role of inhibitory circuits in hippocampal memory processing. *Nat Rev
Neurosci, 23*(8), 476-492. doi:10.1038/s41583-022-00599-0

Vinck, M., Uran, C., Spyropoulos, G., Onorato, I., Broggini, A. C., Schneider, M., & Canales-Johnson, A.
(2023). Principles of large-scale neural interactions. *Neuron, 111*(7), 987-1002.
doi:10.1016/j.neuron.2023.03.015

Wallis, J. D. (2018). Decoding Cognitive Processes from Neural Ensembles. *Trends Cogn Sci, 22*(12), 1091-
1102. doi:10.1016/j.tics.2018.09.002

Wang, M., Foster, D. J., & Pfeiffer, B. E. (2020). Alternating sequences of future and past behavior encoded
within hippocampal theta oscillations. *Science, 370*(6513), 247-250. doi:10.1126/science.abb4151

Wang, Y., Wang, X., Wang, L., Zheng, L., Meng, S., Zhu, N., . . . Ming, D. (2024). Dynamic prediction of goal
location by coordinated representation of prefrontal-hippocampal theta sequences. *Curr Biol, 34*(9),

861   1866-1879 e1866. doi:10.1016/j.cub.2024.03.032

862   Wikenheiser, A. M., & Redish, A. D. (2015). Hippocampal theta sequences reflect current goals. *Nature*
863   *Neuroscience, 18*(2), 289-294. doi:10.1038/nn.3909

864   Zhang, L., Lee, J., Rozell, C., & Singer, A. C. (2019). Sub-second dynamics of theta-gamma coupling in
865   hippocampal CA1. *eLife, 8*, e44320. doi:10.7554/eLife.44320

866   Zheng, C., Bieri, K. W., Hsiao, Y. T., & Colgin, L. L. (2016). Spatial Sequence Coding Differs during Slow and
867   Fast Gamma Rhythms in the Hippocampus. *Neuron, 89*(2), 398-408. doi:10.1016/j.neuron.2015.12.005

868   Zheng, C., Hwaun, E., Loza, C. A., & Colgin, L. L. (2021). Hippocampal place cell sequences differ during
869   correct and error trials in a spatial memory task. *Nat Commun, 12*(1), 3373. doi:10.1038/s41467-021-
870   23765-x

871   Zhu, N., Zhang, Y., Xiao, X., Wang, Y., Yang, J., Colgin, L. L., & Zheng, C. (2023). Hippocampal oscillatory
872   dynamics in freely behaving rats during exploration of social and non-social stimuli. *Cognitive*
873   *Neurodynamics, 17*(2), 411-429. doi:10.1007/s11571-022-09829-8

FIGURE LEGENDS

Figure 1 | The schematic of a model for theta sequence development

Fast gamma rhythms coordinate a subgroup of place cells, as their spikes are

dominantly phase-locked to fast gamma rhythms (FG-cells). The development

process of theta sequences is disrupted by excluding these FG-cells. A possible

model would be that development of sweep-ahead structure of theta sequences is

dependent on the slow gamma modulation, i.e., slow gamma phase precession,

whereby spatial information could be highly compressed within theta cycles. FG-

cells are those cells exhibiting slow gamma phase precession within theta cycles

since from the early stage of sequence development. However, the NFG-cells only

excited slow gamma phase locking during late stage, which may unlikely contribute

to the sequence development.

Figure 2 | A subset of hippocampal place cells was modulated by fast gamma

rhythms during active exploration.

(A) An example of simultaneously recorded LFPs (raw LFP; theta: 4-12Hz; slow

gamma: 25-45Hz; fast gamma: 65-100 Hz) and spikes of place cells. The troughs of

theta were represented by dark gray vertical lines. Slow and fast gamma episodes

were marked by dashed boxes and gray blocks. (B) Firing characteristics of two

examples of FG-cells and NFG-cells, including turning curve, fast and slow gamma

phase distribution during detected episodes. Spikes were represented by red rasters

on top of the gamma traces. Mean vector length of each cell was shown above the

gamma phase distribution. (C) The normalized turning curves of place cells sorted

by the center of mass of their main place field. The proportion of 4 types of cells

identified according to different modulation by fast and slow gamma was shown on

the right. (D) Mean vector length of fast gamma phase distribution of FG-cells and

NFG-cells across successive running laps (n=12 sessions). Data are presented as

mean ± SEM. ***p<0.001.

Figure 3 | Theta sequences development was disrupted without FG-cells.

(A) Top, An example of color-coded decoded probability with time during a single lap.

The running trajectory of the animal was indicated as white dashed line. Bottom, 4–

12 Hz bandpass filtered theta rhythm, with cycles divided on trough indicated as red

bars. (B) Examples of detected theta sequences decoded by 3 different decoders,

all spikes from all cells (top, i.e. raw sequences), all spikes excluding those from FG-

cells (middle, i.e. exFG-sequences) and all spikes excluding those from NFG-cells

(bottom, i.e. exNFG-sequences). The white triangle indicates the animal's current

position. The white boxes indicate the center of sequences for calculating weighted

correlation. (C) Weighted correlations of sequences among three types of decoders

(n=12 session). (D) Averaged decoded probabilities for each lap, at a range of the

animal's current position +-63 cm and the mid-time point of theta sequence +-170 ms,

of a single recording session. (E) Weighted correlation of raw sequences was

significantly increased with running laps (n=12 sessions). (F) The running speeds

were maintained across laps. (G) The sweep-ahead structure of the exFG-

sequences was disrupted compared to that of exNFG-sequences (n=12 sessions).

(H) The effect of excluding either FG-cells or NFG-cells at early (left) and late (right)

development of theta sequences (n=24 laps). Data are presented as mean ± SEM.

*p<0.05, **p<0.01, ***p<0.001.

Figure 4 | Temporospatially compressed structure of theta sequences required

a relatively large proportion of FG-cells.

(A) Examples of detected theta sequences in early development. Each column is a pair of theta sequences decoded by 2 different decoders, i.e. exFG-decoder and exNFG-decoder, in a same theta cycle. The left 2 columns of sequences were detected from laps with a relatively small proportion (~10%) of FG-cells. The right 2 columns of sequences were detected from laps with a relatively large proportion (~35%) of FG-cells. Weighted correlation (W.Corr) of each sequence was shown. (B) Same as (A), but for late development of theta sequences. (C) The difference of weighted correlation between exFGsequences and exNFGsequences as a function of the proportion of excluded cells, in early development. Each scatter represents data from each recording session (n=12 sessions). The dashed line is the linear regression line. (D) Same as (C), but for late development of theta sequences.

Figure 5 | Comparison of FG-cells and NFG-cells firing characteristics.

(A) Spatial tuning curves of representative FG- and NFG-cells during exploration on the circular track. Each column of cells exhibited similar place field positions on the track. (B) Distribution of the place field centers of mass (COM) of FG-cells and NFG-cells as a function of position on the track (n=113 cells for FG-cells and downsampled n=113 cells for NFG-cells). (C) Cumulative distribution of FG-cells and NFG-cells' place field COM. (D) Mean firing rate, (E) Excluded spike counts in decoding, (F) Place field size, and (G) Spatial information of FG-cells and NFG-cells (n=12 sessions). Data are presented as mean ± SEM. *p<0.05

Figure 6 | Slow gamma phase precession during early and late development of theta sequences.

(A) Probability distributions of slow gamma phases of spikes across successive slow

gamma cycles within theta sequences. Gamma cycles within theta cycles were

ordered, centered at cycle 0 (i.e., gamma cycle with maximal spiking). Slow gamma

phases of spikes shifted systematically across successive slow gamma cycles. (B)

same as(A) but during early development of theta sequences. Spikes' slow gamma

phase did not significantly backward-shifted across first 2 slow gamma cycles. (C)

Same as (A), but during late development of theta sequences. The white dots

denote the peak probability of the histogram.

Figure 7 | FG-cells constantly exhibited slow gamma phase precession across

laps.

(A) Examples of slow gamma phase precession within theta cycles on single cell

level. The top row shows posterior probability for theta sequences. The middle row

shows the LFPs waveform in the same theta cycle of the theta sequence, with each

slow gamma cycle divided by black lines. The bottom row shows the slow gamma

phases of spikes from a representative cell activated in this theta cycle. The spikes

occurred at earlier slow gamma phase in the late slow gamma cycles. (B) The

averaged slow gamma phase shift across N active cycles were significantly negative

(N=2, 3 or ≥4). The red lines indicate the median and the black lines indicate the 25%

and 75% quantile of group data (n=12sessions for N=2 and N=3, n=9sessions for

N≥4). (C) Histogram of mean slow gamma phase shift of theta cycles across laps.

The histograms of real data are shown in blue with blue triangle indicating median.

The histograms of mock data are shown in grey with black lines indicating the 95th

confident interval of the their median. (D) The same as (C) but for histogram of

mean slow gamma phase shift of place cells across laps. (E) Histogram of mean

slow gamma phase shift of FG-cells (red) or NFG-cells (yellow) across laps. The red and yellow triangles are the median of phase shift in FG-cells and NFG-cells respectively. *p<0.05, **p<0.01, ***p<0.001.

Figure 8 | The sweep-ahead structure of FG-cell sequences was positively correlated with the intensity of slow gamma phase precession during both early and late sequence development.

(A) Scatter plot of averaged slow gamma phase shift as a function of weighted correlation of sequences during all laps. Left panel shows real data (blue) and right panel shows mock data (gray). (B) Probability distributions of sequences falling in 4 quadrants (Q1-Q4) with slow gamma phase shift (<0 or >0) and weighted correlation (<0 or >0). (C) The count of sequences falling in Q1 (blue triangle) was significantly higher than 95% quantile of the shuffled data (black line). The gray histogram indicates the relative probability of mock data falling in Q1 by 1000 times shuffling. (D) Scatter plot of averaged slow gamma phase shift as a function of weighted correlation of FG-cell sequences during early development (top) and late development (bottom). (E) Probability distributions of FG-cell sequences falling in 4 quadrants during early development (top) and late development (bottom). (F) The count of FG-cell sequences falling in Q1 (red triangle) was significantly higher than 95% quantile of the shuffled data (black line) during both early development (top) and late development (bottom) of sequences. (G-H) Same as (D-E), but for NFG-cell sequences. (I) The count of NFG-cell sequences falling in Q1 (red triangle) was significantly higher than 95% quantile of the shuffled data (black line) only during late (bottom) but not early (top) development of sequences. *p<0.05, ***p<0.001.

Fast gamma phase-locking?

FG-cells ✓

NFG-cells ✗

exFG-sequences

Disrupted development
of theta sequences

exNFG-sequences

Successful development
of theta sequences

Slow gamma phase precession?

✓

early

✗

✓

late

✓

Theta sequence development

A

Early development

exFG-sequences

Excluding ~10%

W. Corr=0.17

W. Corr=0.13

Excluding ~35%

W. Corr=0.05   W. Corr=0

exNFG-sequences

W. Corr=0.41   W. Corr=0.25

W. Corr=0.17   W. Corr=0.53

Time (ms)   Time (ms)

B

Late development

exFG-sequences

Excluding ~10%

W. Corr=0.28

W. Corr=0.64

Excluding ~35%

W. Corr=0   W. Corr=0.23

exNFG-sequences

W. Corr=0.68

W. Corr=0.65

W. Corr=0.43   W. Corr=0.39

Time (ms)   Time (ms)

C

Early development

Weighted Corr Difference (exNFG - exFG)

corr = 0.1
p = 0.7

Proportion of excluded cells /%

D

Late development

Weighted Corr Difference (exNFG - exFG)

corr = 0.6
p = 0.03

Proportion of excluded cells /%

A

FG-cell #1

FG-cell #2

NFG-cell #1

NFG-cell #2

Firing rate (Hz)

Position on the track (cm)

B

FG-cells   NFG-cells

Cell count

Position on the track (cm)

C

FG-cells
NFG-cells

cumulative distribution

n.s.

Place field COM (cm)

D

Mean firing rate (Hz)

*

FG   NFG
cells

E

×10³

Excluded spike counts

n.s.

FG   NFG
cells

F

Place field size (cm)

n.s.

FG   NFG
cells

G

Spatial information (bit/spike)

n.s.

FG   NFG
cells