Genetic diversity affects ecosystem functions across trophic levels as much as species diversity, but in an opposite direction

Laura Fargeot*[1,2], Camille Poesy[1], Maxim Lefort[1], Jérôme G. Prunier[1], Madoka Krick[1], Rik Verdonck[3], Charlotte Veyssière[2], Murielle Richard[1], Delphine Legrand[1], Géraldine Loot[2] & Simon Blanchet*[1]

[1] Centre National de la Recherche Scientifique (CNRS), Station d'Ecologie Théorique et Expérimentale, UAR 2029, 2 route du CNRS, F-09200 Moulis, France; laura.fargeot@live.fr (L.F.); camille.poesy@orange.fr (C.P.); maxim.lft@gmail.com (M.L.); jerome.prunier@gmail.com (J.G.P.); madokakrick@gmail.com (M.K.); murielle.richard@sete.cnrs.fr (M.R.); simon.blanchet@sete.cnrs.fr (S.B.); delphine.legrand@sete.cnrs.fr (D.L.)

[2] Université Paul Sabatier (UPS) Toulouse III, 118 route de Narbonne, F-31062 Toulouse CEDEX 4, France; geraldine.loot@univ-tlse3.fr (G.L.); veyssiere.charlotte@gmail.com (C.V.)

[3] Center for Environmental Sciences, Environmental Biology, Hasselt University, Diepenbeek, Belgium; rik.verdonck@uhasselt.be (R.V.)

Running title: Genetic vs. species biodiversity effects

Key words: genomic diversity, decomposition, productivity, trophic chains, aquatic ecosystems, fish, shredders, riparian trees, biodiversity-function relationships

Number of words in abstract: 148

Number of words in main text: 4979

Number of references: 44

Number of figures and tables: 7

* Correspondence:

Laura Fargeot and Simon Blanchet

Station d'Ecologie Théorique et Expérimentale

09200, Moulis, France

Phone number: +33 5 61 04 03 61

Laura Fargeot, laura.fargeot@live.fr

Simon Blanchet, simon.blanchet@sete.cnrs.fr

ABSTRACT

Understanding the relationships between biodiversity and ecosystem functioning stands as a

cornerstone in ecological research. Extensive evidence now underscores the profound

impact of species loss on the stability and dynamics of ecosystem functions. However, it

remains unclear whether the loss of genetic diversity within key species yield similar

consequences. Here, we delve into the intricate relationship between species diversity,

genetic diversity, and ecosystem functions across three trophic levels —primary producers,

primary consumers, and secondary consumers— in natural aquatic ecosystems. Our

investigation involves estimating species diversity and genome-wide diversity -gauged within

three pivotal species- within each trophic level, evaluating seven key ecosystem functions,

and analyzing the magnitude of the relationships between biodiversity and ecosystem

functions (BEFs). We found that, overall, the absolute effect size of genetic diversity on

ecosystem functions mirrors that of species diversity in natural ecosystems. We nonetheless

unveil a striking dichotomy: while genetic diversity was positively correlated with various

ecosystem functions, species diversity displays a negative correlation with these functions.

These intriguing antagonist effects of species and genetic diversity persists across the three

trophic levels (underscoring its systemic nature), but were apparent only when BEFs were

assessed within trophic levels rather than across them. This study reveals the complexity of

predicting the consequences of genetic and species diversity loss under natural conditions,

and emphasizes the need for further mechanistic models integrating these two facets of

biodiversity.

Introduction

Diversity *within* and *among* species are both important to ensure and stabilize ecosystem functions (Cardinale *et al.* 2012; Raffard *et al.* 2019). Studies on the links between biodiversity and ecosystem functioning (BEFs) have primarily focused on the interspecific (species) facet of biodiversity (Balvanera *et al.* 2006; Hooper *et al.* 2005). However, the intraspecific (genetic) facet of biodiversity has also recently been shown to have substantial effects on ecosystem functions (Crutsinger *et al.* 2006; Hughes & Stachowicz 2004; Reusch *et al.* 2005). Recent meta-analyses have shown that genetic diversity of plant and animal populations affect ecosystem functions, and that the magnitude (and shape) of intraspecific BEFs is similar to that of species diversity (Raffard *et al.* 2019; Wan *et al.* 2022).

Although natural assemblages encompass both intra- and interspecific diversity, most studies investigating BEFs are considering each biodiversity facet separately (but see, Fridley & Grime 2010; Prieto *et al.* 2015; Grele *et al.* 2024). This makes it difficult to differentiate the relative role of genetic and species diversity in ecosystem functions, impeding general predictions regarding the consequences of biodiversity loss as a whole on ecosystem functions (Blanchet *et al.* 2023). For instance, we are currently unaware whether the loss of genetic diversity within a few species in an assemblage is as detrimental for ecosystem functions as a species loss, or whether the combined loss of genetic and species diversity may have non-additive consequences for ecosystem dynamics. Although these biodiversity loss scenarios are realistic, our knowledge on the relative role of genetic *vs.* species diversity in ecosystem functions are still too scarce to provide reliable predictions.

The few studies investigating the combined effects of genetic and species diversity on ecosystem functions were all conducted experimentally by manipulating the genetic and species diversity of assemblages under controlled conditions (Fridley & Grime 2010; Hargrave *et al.* 2011; Prieto *et al.* 2015; but see Grele *et al.* 2024). Our understanding of genetic (intraspecific) and species (interspecific) BEFs therefore relies on simplified

ecosystems that often lack variation in other factors (including spatial scales, abiotic

factors…), and in which feedbacks between ecosystem functions and biodiversity are limited

(Duffy *et al.* 2017; Prunier *et al.* 2023). However, knowledge acquired from BEFs at the

interspecific level reveals that environmental variation can either reduce or enhance the

effects of biodiversity on ecosystem functions, hence generating large variance in the

magnitude and direction of BEFs measured in the wild (Hagan *et al.* 2021; Van Der Plas

2019). One can therefore predict that, under natural conditions, the relative influence of

genetic and species diversity on ecosystem functions may deviate from what has been

quantified under controlled conditions, although it is difficult to predict the direction of this

deviation as field studies (in particular for genetic BEFs) are too scarce to generate clear

predictions. Therefore, we need further realistic field studies of BEFs, embracing the whole

diversity of life forms (from genes to species) and across realistic environmental gradients to

test whether -under natural conditions- species and genetic BEFs are of similar magnitude.

BEF studies often consider a single trophic level, despite accumulating evidence that

biodiversity at a given trophic level can propagate across trophic levels, generating "multi-

trophic BEFs" (Lefcheck *et al.* 2015; Soliveres *et al.* 2016, Seibold *et al.* 2018). In particular,

studies testing the joint effects of genetic and species diversity on ecosystem functions have

mostly considered the effect of primary producer diversity on their own productivity ("*within-*

trophic level BEFs", e.g., Hargrave *et al.* 2011; Prieto *et al.* 2015). However, genetic and

species diversity within a given trophic level may have propagating effects on the ecosystem

at other trophic levels (hereafter, "*between-trophic level BEFs*"). Indeed, it is predicted that a

genetically-diverse predator population shares their resources more efficiently than a

genetically-poor predator population, which might permit a higher prey species coexistence

and hence a larger prey biomass (*between-trophic level* BEFs due to genetic diversity, e.g.,

Raffard *et al.* 2021). Alternatively, a species-rich community of primary producers likely

exhibits higher primary production, as organisms in species-rich communities share basal

resources more efficiently than in species-poor communities (*within-trophic level BEFs due*

to species diversity, Balvanera *et al.* 2006; Hooper *et al.* 2005). Similarly, the relative impact

of genetic and species diversity should be inconsistent across trophic. At higher trophic

levels (e.g., predators), species richness is generally lower, which should increase the

likelihood for genetic diversity (of a few species) to have strong effects on functions. A

simple prediction might therefore be that the relative impact of genetic diversity on

ecosystem functions should increase with increasing trophic levels (Blanchet *et al.* 2020).

Studies considering genetic and species BEFs under a realistic multitrophic scenario may

thus help understanding the trophic contexts under which either genetic or species diversity

is more impactful on ecosystem functions than the other, and to test whether genetic and

species effects can propagate across trophic levels or not (Seibold *et al.* 2018, Li *et al.* 2020,

Moi *et al.* 2021).

Here, we conducted a field study to test the relative importance of genetic and

species diversity for ecosystem functions across multiple trophic levels in a natural

landscape. We focused on three trophic levels from river ecosystems; riparian trees (primary

producers), macroinvertebrate shredders (primary consumers) and fish (secondary

consumers). For each trophic level, we quantified the species diversity of each community,

as well as the genetic diversity of a single target and dominant species (*Alnus glutinosa*,

Gammarus sp. and *Phoxinus dragarum* respectively). We further estimated several

ecosystem functions, including leave decomposition of riparian trees, biomass (as

productivity estimates) of each target species and total biomass of each community within

each trophic level. We relied on causal analyses, taking into account the direct and indirect

effects of the environment (through biodiversity) on ecosystem functions (Duffy *et al.* 2016)

to test i) whether BEFs measured at the genetic level (*genetic* BEFs) are similar in

magnitude and direction to BEFs measured at the species level (*species* BEFs); and ii)

whether *within-trophic level* BEFs are similar in magnitude than *between-trophic level* BEFs.

We also tested whether the relative effects of species and genetic diversity on ecosystem

functions (within or between trophic levels) are consistent across the three trophic levels

(primary producers, primary consumers and secondary consumers), in order to generalize

findings along the trophic chain. We predicted that -contrary to what has been observed

under controlled conditions- *genetic* BEFs and *species* BEFS will not be similar in

magnitude, especially because environmental variation may modulate each of them

differentially. We further expected that significant genetic and species BEFs will be observed

both *within*- and *between*-trophic levels, leading to *within*- and *between*-trophic levels of

similar magnitude. Finally, we predicted that the magnitude of *genetic* BEFs will be higher

(than that of *species* BEFs) at the highest trophic level (secondary consumers) than at the

lowest trophic level (primary producers), mainly because species richness at higher trophic

levels presents a lower gradient that at the lowest trophic levels.

Materials and methods

Sampling sites and trophic chain

We sampled 52 sites in Southern France from the Adour-Garonne watershed, and

distributed along an east-west gradient in the Pyrenees Mountains (**Figure 1a**). We acquired

data on species diversity, genetic diversity and ecosystem functions at three trophic levels

(primary producers, primary consumers and secondary consumers) (**Figure 1b**). Riparian

trees (57 species in the sampled area) provide organic matter in the form of fallen leaves as

a food source for decomposers. We selected the common alder *Alnus glutinosa* for

acquisition of genetic data due to its dominance at most sites and its functional relevance, as

its roots serve as shelters for many aquatic species and are involved in nitrogen fixation.

Macroinvertebrate shredders (101 genera in the sampled area) are primary consumers using

leaves as resources, and converting them into accessible organic matter for other species.

We focused on the most abundant Gammarid (Crustacean) species for genetic data

acquisition, referred to as *Gammarus* sp. This species has not yet been formally named

although it is phylogenetically distinct from its closest relative, *Gammarus fossarum*

(Carnevali 2022; Piscart, unpublished data). This species is particularly efficient at

decomposing tree leaves, in particular those from *Alnus* (Macneil *et al.* 1997). Fish (20

species in the sampled area) are secondary consumers feeding on invertebrates (amongst

others). We used the minnow *Phoxinus dragarum* as the fish target species as it is an

abundant and important predator strongly impacting invertebrate communities (Raffard *et al.*

2021).

Biodiversity estimates

Species datasets. At each site, we collected data on the abundance of all species within

each trophic level, at one occasion for trees (July-August 2021) and two occasions for

invertebrates (July and November 2020) and fishes (mid-July to mid-August 2020 and 2021),

to obtain accurate biodiversity estimates. We identified tree species along a 200 m transect

of each river bank, excluding trees with trunk smaller than 2 cm in diameter and more than

one meter away from the bank. The abundances of trees were estimated as the total number

of individuals per species and per site. For invertebrates, we identified shredders to the

genus level (or to the family level for some groups such as chironomids) sampled from two

types of standardized traps installed in four micro-habitats distributed along the 200m

transect used to identify trees: natural coconut brushes (15*5.5 cm, bristles length 7.5 cm)

recovered after 1.5 month of colonization, and litter bags (15*11 cm, 0.8 cm mesh size) filled

with senescent *Alnus* leaves from each site and recovered after nine days of colonization

(see below). We calculated abundances of each genus by summing the number of

individuals per genus found in the coco brushes and the litter bags, and we averaged the

abundances over the two sampling occasions to get a single estimate per genus per site.

For fish, we collected all specimens during single-pass electric fishing sessions over a mean

area of ~ 469.9 m^2 (± 174 m^2) distributed along the 200 m transect. We anesthetized,

identified and counted individuals at the species level. We calculated fish abundances as the

number of individuals per species and per m², and we averaged the abundances over the

two sampling occasions as for invertebrates. Fish species number varies from 1 to 11,

invertebrate genus number varies from 15 to 42 and the tree species number varies from 7

to 20 (see Fargeot *et al.* 2023 for details).

Genetic datasets. At each site, we collected tissue from up to 32 individuals of each of the

three target species. We collected fresh leaves of *A. glutinosa* in May 2020, specimens of

Gammarus sp. in February 2020, and a piece of pelvic fin from *P. dragarum* individuals in

summer 2020. The DNA of these samples was extracted using commercial kits for *Alnus*

and *Gammarus sp.* and a salt-extraction protocol for *P. dragarum* (see Fargeot *et al.* 2023

for details). For each specimen, DNA concentrations were measured using Qubit 3.0

fluorometer (Life Technologies®, USA). Sequencing was performed based on equimolar

pools of DNA ("pool-seq" approach, Schlötterer *et al.* 2014) from each population and each

species. For *Gammarus sp.*, we also obtained a ~600 bp mitochondrial sequence from the

COI mitochondrial gene from each individual to ensure identification and avoid mixing

individuals from different species. *Gammarus* sp. was found allopatric in most sites, but for a

few sites from the eastern part of the area in which two species were identified (Carnevali

2022). In this latter case, we conserve only the target species for creating the DNA pools.

We created one DNA pool per site per species (52 pools for *A. glutinosa*, 47 pools for

Gammarus sp. and 44 pools for *P. dragarum*) and performed double-digest restriction-site

associated DNA sequencing (ddRAD-seq) for *A. Glutinosa* and *Gammarus* sp. (respectively,

PstI/MseI and Pst/HindIII enzymes) and normalized Genotyping-by-Sequencing (nGBS) for

P. dragarum (MsII enzyme). Library preparation and pool-sequencing were executed by

LGC Genomics (Biosearch Technologies®, Germany) on an Illumina NovaSeq® (2x150 pb).

Data processing was performed following De Kort *et al.* (2018), except that read mapping

was performed on reference genomes. The genome of *A. glutinosa* was already available

(Griesmann *et al.* 2018), and we assembled reference genomes from Illumina short-read

sequencing and PacBio long-read sequencing for *Gammarus. sp.* (available upon request)

and *P. dragarum* (accession number on DDBJ/ENA/GenBank: JARPMJ000000000),

respectively. SNP calling was performed with (i) filtering of raw sequencing files; (ii) indexing

of reference genomes; (iii) mapping reads to the reference; (iv) filtering for unpaired and

badly/non-mapped reads; (v) assembling all read information in a single file per population

and per species and (vi) calculating SNP allelic frequencies (De Kort *et al.* 2018). The total

numbers of SNPs retrieved were 583 862 for *A. glutinosa*, 331 728 for *Gammarus* sp. and

414 213 for *P. dragarum* (see Fargeot *et al.* 2023 for details).

Species and genetic diversity estimates. We calculated ☐- diversity per site using the

Shannon entropy from the "hillR" R package for both species and genetic diversity. The

Shannon entropy is a metric of evenness that takes into account the distribution of allele or

species abundances within each site (Chao *et al.* 2014) by weighting each species/allele by

its proportional abundance (q = 1). Results were similar when using the Simpson's diversity

index (q = 2, results not shown).

Ecosystem function measurements

At each site, we measured seven ecosystem functions. We collected biomass

production data of all species at each trophic level (hereafter "total biomass") and the

biomass production of each target species as estimates of productivity, as well as the

decomposition rate of *Alnus* leaves. Productivity -as we quantified it- is obviously affected by

local environmental characteristics, and for this reason, we took into account these potential

environmental effects (see hereafter). For riparian tree biomass, we used the trunk diameter

of each single tree as a proxy of individual tree biomass, and we summed the trunk

diameters of all trees found along the transect (divided by the length of the transect) to

estimate the total tree biomass per site and per meter of bank. The same approach was

used to estimate *A. glutinosa* biomass. For macroinvertebrate shredders, we estimated the

total invertebrate biomass by drying all individuals for 24 h at 60 °C before weighing them

(10^{-4} g precision). The same procedure was used to estimate the biomass of *Gammarus* sp.

For both estimates, we averaged biomasses over the two sampling sessions. For fish,

(fresh) total fish biomass was estimated as the total weight of all individuals (0.01 g

precision) per site, whereas *P. dragarum* biomass was the mass of all *P. dragarum*

specimens per site. Fish biomasses were averaged over the two sampling sessions.

For the decomposition rate, we quantified leaf mass loss in litter bags placed in four

micro-habitats per site twice (July and November 2020). We gathered and dried senescent

leaves during fall 2019 from five *Alnus* trees per site to limit individual-specific effects on

decomposition. Litter bags were 15 cm x 11 cm pockets of plastic-wire mesh (mesh size; 8

mm to allow invertebrates colonization) in which we introduced 4g of dried leaves before

closing the bags with staples. We installed three bags per micro-habitat (12 per site) that we

removed sequentially after ~9 days, ~18 days and ~27 days respectively to estimate

decomposition rates. Bags were brought back to the laboratory, the remaining leaves were

cleaned, dried and weighed. Decomposition rate was estimated as the slope of leaf mass

loss over time (obtained from a linear model) that we averaged across replicates and

temporal sessions (Raffard *et al.* 2021).

Environmental data

A major challenge for inferring BEFs from empirical data is to take into account the

direct and indirect (through biodiversity) effects of environmental factors on ecosystem

functions (Duffy *et al.* 2016, 2017). Failing to do this may result in overestimated and/or

artefactual BEFs, especially if the same environmental factor simultaneously affects

biodiversity and ecosystem processes (Grace *et al.* 2016). For each site, we measured

thirteen variables related to river topography and physico-chemical characteristics that likely

influence biodiversity and ecosystem processes (Altermatt 2013). *River bed width* (m) was

averaged from five measurements per site. *Connectivity* was calculated as the "closeness

centrality", *i.e.,* the inverse of the sum of the distances of a node to all other nodes along the

shortest paths possible (Altermatt 2013), using QGIS and the "RiverDist" R package. *Altitude*

(m), *distance from the outlet* (m) and *east-west gradient* (longitudinal position along the

Pyrenees chain) were measured using QGIS; *oxygen concentration* (mg.L^{-1}), *oxygen*

saturation (%), *water temperature* (°C), *specific conductivity* (µS/cm) and *pH* were measured

(and averaged) in summers 2020 and 2021 using a multi-parameter probe (Aqua TROLL

500, In-Situ Inc.). Concentration of *NO$_3$-, NO$_2$-, NH$_4^+$* and *PO$_4^{3-}$* were estimated (and

averaged) during summers 2020 and 2021 from a filtered water volume (100 mL) using the

Alpkem Flow Solution Iv Autoanalyzer (OI Analytical®).

A Principal Component Analysis combining all thirteen variables was performed using

the R package "ade4" (Dray & Dufour 2007), and coordinates of each site on the two first

axes (38.03% of the total variance, see **Table 1**) were used as two synthetic environmental

variables for further analyses. We kept only these two first axes to avoid collinearity and

over-parameterization of subsequent models. The first axis is defined by a strong

contribution of (in decreasing order) oxygen concentration and altitude (**Table 1**). The

second axis is defined by a strong contribution of east-west gradient and connectivity (**Table**

1).

Statistical analyses

BEF relationships. To quantify the magnitude of association between biodiversity estimates

and ecosystem functions (BEFs), we performed piecewise Structural Path Models (pSEM,

"piecewiseSEM" package, Lefcheck 2016). pSEM allows modelling direct and indirect causal

relationships among a set of response variables and predictors (Shipley 2009). Further,

pSEM uses local estimates of each linear structural equation separately (*i.e.*, parameters are

estimated from a series of independent models forming a general causal graph), which

allows the inclusion of a large number of parameters despite modest sample sizes (Shipley

2009). We ran a pSEM for each ecosystem function separately (*i.e.*, seven pSEMs, see an

example in **Figure 2**). In each pSEM, the ecosystem function was the dependent variable

whereas the six biodiversity estimates (species and genetic diversity estimated for each

trophic level) and the two synthetic environmental variables were the predictors. In each

model, environmental predictors were allowed to explain each biodiversity estimate (indirect

effects of environmental variables through their influence on biodiversity, see **Figure 2**). For

some functions (in particular those associated with plant biomass), irrelevant biodiversity-

functions links were not included (e.g., the impact of fish or invertebrate diversity on tree

biomasses), which results in 34 BEFs (out of the 42 possible links) having been included in

the meta-regression (see hereafter).

From each pSEM model, we retrieved the local parameter (standardized estimate, an

equivalent to a coefficient of correlation) associated with the direct effect of each biodiversity

estimate (six per function, but for some functions for which ecology-irrelevant BEFs were

excluded) on the function (coloured arrows in **Figure 2**), which provides both the magnitude

and the direction of each BEF. To smoothen comparison, we calculated a standardized

effect size for each BEF by applying the Fisher's Z transformation (Zr) to the standardized

estimates. Positive Zr indicate positive associations between biodiversity and ecosystem

functions, whereas negative Zr indicate negative relationships. The higher the absolute value

of Zr, the higher the strength of the association. Zr therefore indicate both the direction

(positive or negative) and the magnitude of the associations. Our seven measures of

ecosystem functions were not correlated one to each other (all $r_{pearson} < |0.39|$).

Direction and magnitude of all types of BEFs. We used a linear mixed-model to test (i)

whether the magnitude and direction of *genetic* BEFs are similar to those of *species* BEFs,

and (ii) whether *within-trophic level* BEFs are similar in effect size to *between-trophic level*

BEFs. In this model, Zr (providing the direction and magnitude of each BEF, n=34) was the

dependent variable, and the predictors were the diversity facets used to measure

biodiversity (genetic or species diversity) and the type of BEF (*within-trophic* or *between*

trophic levels, triangles vs. dots in **Figure 2**). We included the two-term interaction between

diversity facet and type of BEF to test whether the magnitude and direction of *genetic* and

species BEFs are consistent across *within-trophic level* and *between-trophic level* BEFs. We

further included in this model the type of ecosystem function as a random term (to take into

account that each ecosystem function was associated with several biodiversity estimates) as

well as the inverse of the asymptotic variance (v_z=n-3) associated with each effect size as a

weighting parameter for each case study (Balvanera *et al.* 2006; Raffard *et al.* 2019).

We ran an additional linear mixed-model similar to the previous one, except that we

added as a fixed effect the trophic level at which biodiversity was measured to estimate

BEFs (primary producers, primary consumers or secondary consumers) as well as all

interaction terms. Interaction terms allow testing the consistency of major conclusions across

trophic levels, thereby determining the extent to which our findings can be generalized along

the trophic chain. Models were run using the lmer function ("lme4" package) and significance

of fixed effects was determined using type III ANOVA with Wald chi-square tests (function

Anova from the "car" R package, α=0.05).

Results

Details of causal models linking environmental parameters, species and genetic

diversity and ecosystem functions are graphically depicted in **Figure 3a-g**. The percentage

of variance in ecosystem function explained by the environment and biodiversity varies from

10% (invertebrate biomass, **Figure 3e**) to 55% (*Phoxinus* biomass, **Figure 3f**) and was

moderate overall. For all functions but the three biomass, part of the variance was (directly)

explained by at least one out of the two environmental PCA axes. For some functions (e.g.,

Phoxinus biomass, Figure 3f), there was a combined effect of several biodiversity estimates,

whereas for other functions (e.g., *Alnus* biomass, Litter decomposition, **Figure 3a, 3c**) the

effect of a single biodiversity estimate predominates. Overall, direct environmental effects on

ecosystem functions did not predominate, and environmental effect sizes were similar (in

strength) to that of biodiversity effects, showing the important role of biodiversity for

ecosystem functions in the wild.

Individual effect sizes (*Zr*) measured between biodiversity estimates and ecosystem

functions were weak to moderate, irrespectively of the considered ecosystem function and of

the type of BEFs (*genetic/species BEFs*, *within-trophic level/ between-trophic level* BEFs)

(**Figure 4a, 4b, Table S2**). As expected under natural conditions (Hagan *et al.* 2021), BEFs

ranged from negative to positive, and their distribution were centred around 0, although we

observed a slight tendency for *genetic* BEFs toward positive values (**Figure 3b**). Four out of

the 34 BEFs were strong and significant; two significant BEFS concerned *species* BEFs

(negative relationship between the biomass of *A. glutinosa* and the diversity of trees, *Zr* = -

0.446, 95% CI [-0.695, -0.143]; negative relationship between the biomass of *P. dragarum*

and the diversity of fish, *Zr* = -0.529, 95% CI [-0.802, -0.166]) and two concerned *genetic*

BEFs (negative relationship between the biomass of *P. dragarum* and the diversity of *A.*

glutinosa, *Zr* = -0.321, 95% CI [-0.602, -0.019]; positive relationship between the biomass of

Gammarus sp. and the diversity of *P. dragarum*, *Zr* = 0.446, 95% CI [0.001, 0.829]) (**Figure**

4a). Noteworthily, for *within-trophic* BEFs, most case studies fall into the category whereby

genetic BEFs tend to be positive and species BEFs tend to be negative (grey bottom-right

square in **Figure 4a**).

We confirmed this visual tendency by summarizing all individual *Zr* through a meta-

regression. Indeed, we found a significant interaction between the facet at which biodiversity

is measured (genetic or species diversity), and the type of BEF that was measured (within-

or between trophic levels; **Table 2**). This interaction indicates (i) that -overall- *within-trophic*

level BEFs were significantly negative when considering species diversity ($Zr_{\text{Within*Species}}$ = -

0.185, 95% CI [-0.343, -0.027]), whereas *within-trophic level* BEFs were significantly positive

when considering genetic diversity ($Zr_{\text{Within*Genetic}}$ = 0.168, 95% CI [0.010, 0.326], see **Figure**

5a), and (ii) that this pattern was not observed for *between-trophic levels* BEFs, where no

particular trend was observed (**Figure 5a**). Although most individual Zr were weak to

moderate (and not significant), their consistency (in term of magnitude and direction)

resulted in a significant pattern whereby species and genetic diversity have opposite effects

on ecosystem functions for *within-trophic level* BEFs; species diversity is negatively

associated, whereas genetic diversity is positively associated with ecosystem functions, but

only when the influence of biodiversity on ecosystem functions is measured within the same

trophic level.

When including the trophic level at which biodiversity is measured, we found no

significant interaction terms between trophic levels and other fixed effects nor any additive

effect of trophic levels (see **Table S1**). This indicates that our main findings were consistent

across trophic levels, *i.e.*, the respective negative and positive effects on ecosystem

functions of species and genetic diversity hold statistically true across all trophic levels

(**Figure 5b**).

Discussion

We provide empirical evidence that, in natural ecosystems, the effect sizes of genetic

and species diversity on multi-trophic ecosystem functions are of similar magnitude, but

operate in opposite directions. Indeed, for BEFs measured within the same trophic level, the

effects of species diversity across multiple ecosystem functions were moderately negative

on average, whereas the effects of genetic diversity were moderately positive. This suggests

an antagonistic effect between the genetic and the species components of biodiversity in the

modulation of ecosystem functions within one trophic level. This antagonistic effect was not

identified for BEFs measured across trophic levels, since in these cases the influence of

both genetic diversity and species diversity across multiple ecosystem functions was

generally not different from zero. These conclusions hold true across three trophic levels

(plants, invertebrates and fish), indicating that the relative effects of genetic and species

diversity on ecosystem functions are not limited to a specific trophic level.

Our study is one of the few field-based study revealing BEFs across an entire

(riverine) food chain spanning from primary producers to secondary consumers. Indeed,

most previous BEF studies in the field focused on a single trophic level, and predominantly

on terrestrial primary producers (Duffy *et al.* 2017; Van Der Plas 2019, but see e.g., Li et al.

2020, Moi et al. 2021). This permitted encompassing a broad range of ecosystem functions

that depict the overall functioning of a riverine ecosystem (rather than focusing on a single

compartment). Moreover, we focused both on the effects of genetic and species diversity on

these ecosystem functions, which has rarely (if not ever) been evaluated so far and which

provides an exhaustive understanding of BEFs in the wild. Our causal analyses also took

into account the direct (and indirect) effect of environmental factors on ecosystem functions,

which is a prerequisite to isolate biodiversity effects. Direct environmental effects on

ecosystem functions were (in average) not stronger in intensity than biodiversity effects,

which is coherent with previous syntheses on species BEFs in the wild (Duffy *et al.* 2017).

Furthermore, environmental factors used to describe sampling sites in this study were not

strong predictors of species and genetic biodiversity. Two non-exclusive hypotheses may

explain this observation: (i) using PCA axes to resume environmental gradients may blur

some specific environment-biodiversity links, and (ii) as shown and explained in a

companion paper (Fargeot *et al.* 2023), the East-West gradient used in this study (rather

than a classical upstream-downstream gradient) intrinsically limits the potential for strong

environmental effects on biodiversity (which was the purpose of this sampling design).

Nonetheless, after accounting for these environmental covariates, we found that most

individual BEFs (either *genetic* or *species*, *within-trophic levels* or *between-trophic levels*

BEFs) were weak to moderate in magnitude, and that they operated almost equally in both

direction (i.e., positive and negative association between biodiversity and ecosystem

functions). As such, the distribution of individual effect sizes was centred around 0, for both

genetic and *species* BEFs. Accordingly, there were only four individual BEFs that were significant, three out of them were negative and one was positive (**Table S2**). This general pattern (low to moderate BEFs with both positive and negative direction) is actually consistent with the most exhaustive meta-analysis having synthesized the magnitude and direction of *species* BEFs in the wild (Van Der Plas 2019) and with recent conceptual works (Hagan *et al.* 2021) concluding that strong and positive BEFs should not be the norm in natural ecosystems, but rather that a mix of positive, neutral and negative BEFs are expected. Our results confirm, using a dataset encompassing three trophic levels in a single ecosystem type, this conclusion.

We focused both on *within-trophic level* and *between-trophic level* BEFs, which likely encompasses a broad array of mechanisms sustaining potential associations between biodiversity and ecosystem functions. For instance, two out of the significant BEFs we reveal are negative association between species diversity (fish or tree species diversity respectively) and the biomass production of one of the target species (*Phoxinus sp.* and *Alnus sp.* respectively). These *within-trophic level* BEFs can -for instance- arise either because, if resources are limited, increased number of species within a patch limit the biomass production of each individual species, or because of a poorer competitive ability of the target species under some environmental conditions, which favors the settlement of additional species. Although teasing apart these two hypotheses is difficult, the former hypothesis implies a negative co-variation within the biomass of the focal species and that of the total pool of species, which was not observed, neither for fish, not for trees (results not shown). The other two significant BEFs concerned the association (either positive or negative) between the genetic diversity of a target species (*Alnus sp.* or *Phoxinus sp.*) with the biomass production of another target species (*Phoxinus sp.* or *Gammarus sp.,* respectively). These *between-trophic level* BEFs likely arise through indirect effects implying the diversity and availability of (prey) resources. An obvious limit of this field-based study is the impossibility to tease out these mechanisms. Another limit is associated with the fact

that, although environmental covariates were taken into account in causal models, they were synthetized by two PCA axes, and we cannot ensure that all potential environmental covariates have been taken into account. This can influence the actual estimates of BEFs in the wild (Duffy *et al*. 2017). Nonetheless, it is noteworthy that we have previously shown that -in this dataset- species and genetic diversity were not correlated one to each other and that each biodiversity facet was sustained by different environmental predictors (Fargeot *et al*. 2023, see also **Figure 3**). This implies that environmental and biodiversity effects inferred in this study are relatively robust (i.e., not distorted by strong collinearities) and can be interpreted independently one from each other (e.g., the positive effect *Phoxinus* genetic diversity on *Phoxinus* biomass is independent from the negative of fish species diversity as the two estimates of biodiversity do not co-vary, see **Figure 3f**).

Keeping limitations associated with field-based studies into account (see above), revealing associations between ecosystem functions and species and genetic biodiversity (or the lack of) in natural ecosystems is -as ecologists- an important step forward to set theoretical and experimental approaches aiming at understanding this complex biological reality. Beyond individual BEF case studies (that were not the main aim of this study), their aggregation across trophic levels and biodiversity facets revealed a clear (and statistically supported) pattern whereby, within trophic levels, genetic and species diversity display antagonistic association with ecosystem functions; the global effect of species diversity across multiple ecosystem functions was negative, whereas the global effect of genetic diversity was positive. This pattern emerges from the "cumulative" effects of weak to moderate associations between biodiversity and ecosystem functions that consistently point toward the same direction (positive for genetic diversity, negative for species diversity), emphasizing the meaningfulness of meta-regressions (and more generally approaches based on effect sizes rather than on p-values) to reveal biological patterns. We hereafter discuss the ecological relevance of this general pattern.

Our results confirm a previous meta-analysis demonstrating that genetic and species

diversity modulates ecosystem functions with a similar magnitude (Raffard *et al.* 2019), and

results from few experimental studies that manipulated both the genetic and species

components of biodiversity under controlled conditions (e.g., Jiang *et al.* 2022; Prieto *et al.*

2015). Indeed, within trophic levels, the absolute mean effect size of genetic and species

diversity across ecosystem functions were of the same magnitude ($|Zr|$ = 0.168 and 0.185 for

genetic and species diversity effects respectively), and slightly greater than the effect sizes

reported under controlled conditions ($|lnRR|$ = 0.132 and 0.134 for genetic and species

diversity effects respectively, Raffard *et al.* 2019) and those more generally reported for

species BEFs ($|Zr|$ = 0.101, Balvanera *et al.* 2006). This demonstrates for the first time that

under natural conditions, the effects of genetic and species components of biodiversity on

ecosystem functions are comparable. However, our study goes two steps further as (i) it

extends the conclusion made by Raffard *et al.* (2019) to multiple trophic levels and (ii) it

demonstrates that the effects of genetic and species BEFs can actually operate in opposite

directions.

As pointed out by Raffard *et al.* (2019), the vast majority (91% of 23 reviewed studies

by 2019, see also Wan *et al.* 2022) of studies investigating the effects of genetic diversity on

ecosystem functions have focused on primary producers, and all of them were based on

experiments, which is also the case for most studies manipulating both genetic and species

diversity. These trends strongly hamper any generalization. On the contrary, our findings

provide a strong support for broadening the conclusion that both genetic and species

diversity can influence ecosystem functions in the wild. More strikingly, our results

demonstrate that, although the absolute effect sizes of genetic and species BEFs are of

similar magnitude, for *within-trophic level* BEFs, the direction of their effects are opposite;

species diversity (in general) reduces the rate of ecosystem functions, whereas genetic

diversity enhances the same functions. For instance, all other things being equal, higher fish

species diversity is associated with a lower productivity (biomass) in *P. dragarum* (see above

for potential explanations), whereas its own genetic diversity tends to be associated with a

higher productivity (**Table S2**). As genetic and species diversity have been found to be

uncorrelated spatially in this landscape (Fargeot *et al.* 2023), covariation among diversity

estimates cannot explain these patterns. These antagonistic effects of genetic and species

diversity on ecosystem functions parallel previous experimental findings on plants (Hazard *et*

al. 2017; Tang *et al.* 2022). It is now essential to understand the mechanisms sustaining

these antagonistic effects as a step forward.

S*pecies* BEFs were on average negative (see **Table S2** for individual estimates),

which contrasts with the general view that species biodiversity favours ecosystem functions,

although it is not that surprising (Dee *et al.* 2023; Hagan *et al.* 2021). Indeed, the net effect

of species biodiversity on ecosystem functions results from the combined effects of both

negative factors, arising from antagonistic interactions such as negative complementarity or

negative selection effect, and positive factors, arising from beneficial interactions such as

niche complementarity or facilitation (Loreau & Hector 2001). It is likely that, in our case, the

net effect of interspecific interactions mostly results from negative complementarity among

species (or strong negative selection effect), whereas the net effect of intraspecific

interactions may result from facilitative interactions and/or improved niche complementarity

with increased genetic diversity. Intraspecific competition is generally stronger than

interspecific competition (Connell 1983), and intraspecific interactions could be expected to

lead more frequently to negative complementarity (and hence negative *genetic* BEFs) than

interspecific interactions. Since we observe the opposite, we can hypothesize that genetic

diversity is essential to increase niche complementarity within species (Bolnick *et al.* 2003)

and hence to reduce the pervasive effects of intraspecific interactions (Hughes *et al.* 2008;

Prunier *et al.* 2023). Given the empirical nature of our study, it is hard to tease apart

underlying mechanisms. Theoretical approaches, modelling simultaneously the genetic and

species components of biodiversity, would be extremely useful to reveal the mechanisms

sustaining opposite effects of intra- and interspecific diversity on ecosystem functions.

These antagonistic effects were observed only for BEFs measured *within* trophic

levels, not for those measured *between* trophic levels. An overall *between-trophic level* BEF

not different from zero suggests that biodiversity at a trophic level has only limited impact on

ecosystem functions at another trophic level. For example, the biomass of *P. dragarum* was

primarily influenced by genetic and species diversity in fishes, rather than the diversity of

their preys (**Table S2**). However, for both genetic and species estimates of biodiversity,

there was a substantial variation in effect sizes for *between-trophic level* BEFs that ranged

from negative to positive BEFs (**Figure 5**). This suggests that biodiversity effects across

trophic levels may be more variable in their direction than within-trophic level BEFs, which

appear as more constrained. Variability in the magnitude and direction of effect sizes for

between-trophic level BEFs likely blur a more general trend, but this variation is actually

expected under natural conditions in which interactions involve multiple prey and predator

species, fostering co-adaptation among communities from different trophic levels (Aubree *et*

al. 2020; Poisot *et al.* 2013). In these cases, trophic complementarity between two trophic

levels (i.e., the originality of a species based on the identity of the species it interacts with)

might be a stronger determinant of ecosystem functions than complementarity measured at

either one of the two trophic levels (Poisot *et al.* 2013). Quantifying trophic complementarity

among our three target species (and communities) using stable isotope or gut content

analyses for instance would be extremely valuable to assess whether this complexity can

better explain BEFs between trophic levels than diversity measured at one of the trophic

level (Aubree *et al.* 2020).

The empirical patterns we revealed here were all extremely consistent across the

three trophic levels, hence allowing generalization. It is noteworthy that, although statistically

strong and consistent, these patterns must be interpreted with care as field-based

approaches are limited in properly taking into account the environmental heterogeneity of

natural ecosystems (Hagan *et al.* 2021). BEFs were not particularly stronger at any specific

trophic level and the relative effects of genetic and species diversity were not dependent on

the trophic level at which the function was estimated. We may have expected a stronger top-

down regulation (i.e., biodiversity of predators has more effects than biodiversity of preys) of

ecosystem functions since previous studies showed that biodiversity loss should have

greater consequences for multi-functionality when it occurs at higher trophic levels (Lefcheck

et al. 2015, Seibold *et al.* 2018). For instance, increased genetic diversity within a predatory

fish species has experimentally been shown to indirectly increase the rate of litter

decomposition by increasing the diversity of shredders (Raffard *et al.* 2021). Similarly, the

relative effects of genetic and species diversity on functions may have varied among trophic

levels, and in particular the relative importance of genetic diversity may have been higher for

species-poor trophic levels (i.e., fish community) because of a "compensatory effect". We

found no evidence for these potential trophic-level dependencies, but instead found

extremely consistent patterns, which, from a broader perspective, reveal the importance of

integrating both multi-trophic and multi-faceted approaches in predicting the overall

consequences of biodiversity loss on ecosystem functioning.

To conclude, we found that the genetic (intraspecific) and species (interspecific)

facets of biodiversity are both important drivers of multiple ecosystem functions in a natural

and multi-trophic context. In the wild, these two facets of biodiversity can, as expected,

generate low to moderately high impacts on ecosystem functions measured across three

trophic levels, and they can operate in opposite directions. This demonstrates the

importance for managers to develop integrative conservation plans spanning the entire

diversity of life (from genes to species). For instance, genetic diversity loss often precedes

species loss, and our results suggest that -in mountain streams- losing genes may actually

be particularly detrimental for the performance of ecosystem functions. As such, it appears

essential to maintain populations with high levels of genetic diversity in these ecosystems.

Future studies should (i) extend these findings to other ecosystems and by quantifying

natural genetic variation in more than a single species per trophic level, (ii) generate

theoretical predictions regarding the mechanisms sustaining the antagonistic effects of

genetic and species diversity on functions we revealed, and (iii) use a broader integrative

approach for estimating biodiversity across facets (*inclusive* biodiversity) by using either a

trait-based approach or a genetic-based approach as recently proposed by Blanchet *et al.*

(2023) and Loreau *et al.* (2023).

AUTHOR CONTRIBUTION

Laura Fargeot: Conceptualization (Supporting); Methodology (Equal); Software

(Equal); Validation (Equal); Formal analysis (Lead); Investigation (Lead); Data curation

(Lead); Writing - original draft (Lead); Writing - review & editing (Equal); Visualization (Lead);

Supervision (Equal); Project administration (Lead). **Camille Poesy:** Methodology (Equal);

Validation (Equal); Investigation (Lead); Data curation (Supporting); Supervision (Equal);

Project administration (Supporting). **Maxim Lefort:** Methodology (Supporting); Investigation

(Lead); Data curation (Supporting); Supervision (Equal); Project administration (Supporting).

Jérôme G. Prunier: Methodology (Supporting); Software (Equal); Formal analysis

(Supporting); Investigation (Supporting); Resources (Equal); Data curation (Supporting);

Writing - original draft (Supporting); Writing - review & editing (Supporting). **Madoka Krick:**

Methodology (Supporting); Investigation (Equal). **Rik Verdonck**: Methodology (Supporting);

Software (Equal); Investigation (Supporting); Writing - review & editing (Supporting).

Charlotte Veyssière: Methodology (Supporting); Investigation (Equal). **Murielle Richard:**

Methodology (Supporting); Investigation (Supporting); Writing - review & editing

(Supporting). **Delphine Legrand:** Validation (Equal); Writing - review & editing (Equal);

Visualization (Supporting). **Géraldine Loot:** Validation (Equal); Investigation (Supporting);

Writing - review & editing (Supporting); Visualization (Supporting). **Simon Blanchet:**

Conceptualization (Lead); Methodology (Lead); Software (Supporting); Validation (Equal);

Formal analysis (Supporting); Investigation (Lead); Resources (Equal); Data curation

(Supporting); Writing - original draft (Supporting); Writing - review & editing (Lead);

Visualization (Lead); Supervision (Lead); Project administration (Lead); Funding acquisition

(Lead).

REFERENCES

Altermatt, F. (2013). Diversity in riverine metacommunities: a network perspective. *Aquatic*
Ecology, 47, 365–377.

Aubree, F., David, P., Jarne, P., Loreau, M., Mouquet, N. & Calcagno, V. (2020). How
community adaptation affects biodiversity–ecosystem functioning relationships. *Ecol*
Lett, 23, 1263–1275.

Balvanera, P., Pfisterer, A.B., Buchmann, N., He, J.-S., Nakashizuka, T., Raffaelli, D., *et al.*
(2006). Quantifying the evidence for biodiversity effects on ecosystem functioning
and services: Biodiversity and ecosystem functioning/services. *Ecol Lett*, 9, 1146–
1156.

Blanchet, S., Fargeot, L. & Raffard, A. (2023). Phylogenetically conserved candidate genes
unify biodiversity–ecosystem function relationships and eco evolutionary dynamics
across biological scales. *Mol Ecol*, 32, 4467–4481.

Blanchet, S., Prunier, J.G., Paz Vinas, I., Saint Pé, K., Rey, O., Raffard, A., *et al.* (2020). A
river runs through it: The causes, consequences, and management of intraspecific
diversity in river networks. *Evol Appl*, 13, 1195–1213.

Bolnick, D.I., Svanback, R., Fordyce, J.A., Yang, L.H., Davis, J.M., Hulsey, C.D., *et al.*
(2003). The ecology of individuals: Incidence and implications of individual
specialization. *Am Nat*, 161, 1–28.

Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., *et al.*
(2012). Biodiversity loss and its impact on humanity. *Nature*, 486, 59–67.

Carnevali, M. (2022). Etude phylogénétique des populations de Gammares (Gammarus sp.)
du piedmont Pyrénéen. Master thesis. University of Toulouse, France.

Chao, A., Chiu, C.-H. & Jost, L. (2014). Unifying Species Diversity, Phylogenetic Diversity,
Functional Diversity, and Related Similarity and Differentiation Measures Through Hill
Numbers. *Annu Rev Ecol Evol Syst*, 45, 297–324.

Connell, J.H. (1983). On the prevalence and relative importance of interspecific
competition : evidence from field experiments. *Am Nat*, 122, 661–696.

Crutsinger, G.M., Collins, MD, Fordyce, J.A., Gompert, Z., Nice, C. & Sanders, N.J. (2006).
Plant Genotypic Diversity Predicts Community Structure and Governs an Ecosystem
Process. *Science*, 313, 966–968.

De Kort, H., Baguette, M., Prunier, J.G., Tessier, M., Monsimet, J., Turlure, C., *et al.* (2018).
Genetic costructure in a meta-community under threat of habitat fragmentation. *Mol*
Ecol, 27, 2193–2203.

Dee, L.E., Ferraro, P.J., Severen, C.N., Kimmel, K.A., Borer, E.T., Byrnes, J.E.K., *et al.*
(2023). Clarifying the effect of biodiversity on productivity in natural ecosystems with
longitudinal data and methods for causal inference. *Nat Commun*, 14, 2607.

Dray, S. & Dufour, A.-B. (2007). The **ade4** Package: Implementing the Duality Diagram for
Ecologists. *J Stat Soft*, 22.

Duffy, J.E., Godwin, C.M. & Cardinale, B.J. (2017). Biodiversity effects in the wild are
common and as strong as key drivers of productivity. *Nature*, 549, 261–264.

Duffy, J.E., Lefcheck, J.S., Stuart-Smith, R.D., Navarrete, S.A. & Edgar, G.J. (2016).
Biodiversity enhances reef fish biomass and resistance to climate change. *P Natl*
Acad Sci, 113, 6230–6235.

Fargeot, L., Poesy, C., Lefort, M., Prunier, J.G., Krick, M., Veyssière, C., *et al.* (2023).
Genomic and species diversity patterns across multiple trophic levels in riverscapes.
Oikos, 2023, e10015.

Fridley, J.D. & Grime, J.P. (2010). Community and ecosystem effects of intraspecific genetic
diversity in grassland microcosms of varying species diversity. *Ecology*, 91, 2272–
2283.

Grace, J.B., Anderson, T.M., Seabloom, E.W., Borer, E.T., Adler, P.B., Harpole, W.S., *et al.*
(2016). Integrative modelling reveals mechanisms linking productivity and plant
species richness. *Nature*, 529, 390–393.

Griesmann, M., Chang, Y., Liu, X., Song, Y., Haberer, G., Crook, M.B., *et al.* (2018).

Phylogenomics reveals multiple losses of nitrogen-fixing root nodule symbiosis.

Science, 361, eaat1743.

Grele, A., Massad, T.J., Uckele, K.A., Dyer, L.A, Antonini, Y., Braga, L., Forister, M.L., Sulca, L., Kato,

M., Lopez, H.G., Nascimento, A.R., Parchman, T., Simbaña, W.R., Smilanich, A.M., Stireman,

J.O., Tepe, E.J., Walla, T. & Richards, L.A. (2024). Intra- and interspecific diversity in a tropical

plant clade alter herbivory and ecosystem resilience. *eLife*, 12, RP86988

Hagan, J.G., Vanschoenwinkel, B. & Gamfeldt, L. (2021). We should not necessarily expect

positive relationships between biodiversity and ecosystem functioning in

observational field data. *Ecol Lett*, 24, 2537–2548.

Hargrave, C.W., Hambright, K.D. & Weider, L.J. (2011). Variation in resource consumption

across a gradient of increasing intra- and interspecific richness. *Ecology*, 92, 1226–

1235.

Hazard, C., Kruitbos, L., Davidson, H., Taylor, A.F.S. & Johnson, D. (2017). Contrasting

effects of intra- and interspecific identity and richness of ectomycorrhizal fungi on

host plants, nutrient retention and multifunctionality. *New Phytologist*, 213, 852–863.

Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., *et al.* (2005).

Effects of biodiversity on ecosystem functioning: A consensus of current knowledge.

Ecol Monogr, 75, 3–35.

Hughes, A.R., Inouye, B.D., Johnson, M.T.J., Underwood, N. & Vellend, M. (2008).

Ecological consequences of genetic diversity: Ecological effects of genetic diversity.

Ecol Lett, 11, 609–623.

Hughes, A.R. & Stachowicz, J.J. (2004). Genetic diversity enhances the resistance of a

seagrass ecosystem to disturbance. *Proc Natl Acad Sci U.S.A.*, 101, 8998–9002.

Jiang, M., He, L., Fan, B., Wang, T., Yang, N., Liu, Y., *et al.* (2022). Intraspecific more than

interspecific diversity plays an important role on Inner Mongolia grassland ecosystem

functions: A microcosm experiment. *Sci Total Env* 826, 154134.

Lefcheck, J.S. (2016). PIECEWISESEM□: Piecewise structural equation modelling in R for

ecology, evolution, and systematics. *Methods Ecol Evol*, 7, 573–579.

Lefcheck, J.S., Byrnes, J.E.K., Isbell, F., Gamfeldt, L., Griffin, J.N., Eisenhauer, N., *et al.*

(2015). Biodiversity enhances ecosystem multifunctionality across trophic levels and

habitats. *Nat Commun*, 6.

Li, F., Altermatt, F., Yang, J., An, S., Li, A., & Zhang, X. (2020). Human activities' fingerprint

on multitrophic biodiversity and ecosystem functions across a major river catchment

in China. Global change biology, 26, 6867-6879.

Loreau, M. & Hector, A. (2001). Partitioning selection and complementarity in biodiversity

experiments. *Nature*, 412, 72–76.

Loreau, M., Jarne, P. & Martiny, J.B.H. (2023). Opportunities to advance the synthesis of

ecology and evolution. *Ecol Lett*, 26.

Macneil, C., Dick, J.T.A. & Elwood, R.W. (1997). THE TROPHIC ECOLOGY OF

FRESHWATER *GAMMARUS* SPP. (CRUSTACEA:AMPHIPODA): PROBLEMS AND

PERSPECTIVES   CONCERNING   THE   FUNCTIONAL   FEEDING   GROUP

CONCEPT. *Biol Rev*, 72, 349–364.

Moi, D. A., Romero, G. Q., Antiqueira, P. A., Mormul, R. P., Teixeira de Mello, F., &

Bonecker, C. C. (2021). Multitrophic richness enhances ecosystem multifunctionality

of tropical shallow lakes. *Funct Ecol*, 35, 942-954.

Poisot, T., Mouquet, N. & Gravel, D. (2013). Trophic complementarity drives the

biodiversity–ecosystem functioning relationship in food webs. *Ecol Lett*, 16, 853–861.

Prieto, I., Violle, C., Barre, P., Durand, J.-L., Ghesquiere, M. & Litrico, I. (2015).

Complementary effects of species and genetic diversity on productivity and stability

of sown grasslands. *Nat Plants*, 1, 15033.

Prunier, J.G., Chevalier, M., Raffard, A., Loot, G., Poulet, N. & Blanchet, S. (2023). Genetic

erosion reduces biomass temporal stability in wild fish populations. *Nat Commun*, 14,

4362.

Raffard, A., Cucherousset, J., Montoya, J.M., Richard, M., Acoca-Pidolle, S., Poésy, C., *et*

al. (2021). Intraspecific diversity loss in a predator species alters prey community

structure and ecosystem functions. *PLoS Biol*, 19, e3001145.

Raffard, A., Santoul, F., Cucherousset, J. & Blanchet, S. (2019). The community and

ecosystem consequences of intraspecific diversity: a meta-analysis: The ecological

effects of intraspecific diversity. *Biol Rev*, 94, 648–661.

Reusch, T.B.H., Ehlers, A., Hämmerli, A. & Worm, B. (2005). Ecosystem recovery after

climatic extremes enhanced by genotypic diversity. *Proc Natl Acad Sci U.S.A.*, 102,

2826–2831.

Schlötterer, C., Tobler, R., Kofler, R. & Nolte, V. (2014). Sequencing pools of individuals —

mining genome-wide polymorphism data without big funding. *Nat Rev Genet*, 15,

749–763.

Seibold, S., Cadotte, M. W., MacIvor, J. S., Thorn, S., & Müller, J. (2018). The necessity of

multitrophic approaches in community ecology. *Trends Ecol Evol*, 33, 754-764.

Shipley, B. (2009). Confirmatory path analysis in a generalized multilevel context. *Ecology*,

90, 363–368.

Soliveres, S., van der Plas, F., Manning, P., Prati, D., Gossner, M.M., Renner, S.C., *et al.*

(2016). Biodiversity at multiple trophic levels is needed for ecosystem

multifunctionality. *Nature*, 536, 456–459.

Tang, T., Zhang, N., Bongers, F.J., Staab, M., Schuldt, A., Fornoff, F., *et al.* (2022). Tree

species and genetic diversity increase productivity via functional diversity and trophic

feedbacks. *eLife*, 11, e78703.

Van Der Plas, F. (2019). Biodiversity and ecosystem functioning in naturally assembled

communities. *Biological Reviews*, 94, 1220–1245.

Wan, N.-F., Fu, L., Dainese, M., Hu, Y.-Q., Pødenphant Kiær, L., Isbell, F., *et al.* (2022).

Plant genetic diversity affects multiple trophic levels and trophic interactions. *Nat*

Commun, 13, 7312.

Table 1. Characteristics of the two first principal components identified by the Principal Component Analysis (PCA) ran on the thirteen environmental variables. The part of the total environmental variance (%) and the contribution of each variable on each component are shown. The variables that contributed significantly to the axis are highlighted in bold.

	Component 1	Component 2
Part of total Variance (%)	21.66	16.37
River width	0.596	0.320
Connectivity	-0.155	**0.646**
Altitude	**0.648**	-0.385
Distance from outlet	0.528	-0.331
East-west gradient	0.105	**-0.795**
Oxygen concentration	**0.738**	0.343
Oxygen saturation	0.266	-0.509
Water temperature	-0.594	-0.012
Specific conductivity	-0.463	-0.045
pH	0.573	0.398
Concentration in $NO_3^+NO_2$	-0.369	-0.286
Concentration in NH_4+	-0.019	-0.207
Concentration in PO_4^{3-}	-0.279	0.236
Global characteristic	Low altitude, poorly oxygenated site - High altitude, highly oxygenated sites	Poorly connected east site - Highly connected west site

761   **Table 2.** ANOVA table for the linear mixed model testing whether the relationships between
762   biodiversity and ecosystem functions measured in a riverine trophic chain differ between the
763   biodiversity facets (species or genetic diversity) and the types of BEF (*within-* or *between-*
764   *trophic levels*). A Wald chi-quare test is used to test the significance of each fixed effect.

	Degree of freedom	Chisq-value	P-value
(Intercept)	1	0.287	0.595
Biodiversity facet	1	0.232	0.630
Type of BEF	1	5.393	**0.020**
Biodiversity facet*Type of BEF	1	5.567	**0.018**

Figure captions

Figure 1. Location of sampling sites and illustration of the trophic chain. **(a)** Distribution of the 52 sampling sites (black dots) spanning an East-West gradient at the foothills of the Pyrenees Mountains (France). Each site is denoted by a six-letter code, with the three first letters indicating the river name and the three last letters indicating the closest city or village. **(b)** Our study focused on a tri-trophic food chain commonly found in mountain rivers, consisting of riparian trees, macroinvertebrate shredders and fishes (from bottom to top). Within each trophic level, we measured two facets of biodiversity: genetic diversity in a single target species within each trophic level (specifically *Phoxinus dragarum*, *Gammarus* sp. and *Alnus glutinosa*), and species diversity from communities.

Figure 2. Example of one of the seven causal models used to quantify the relationships between (species and genetic) diversity and ecosystem functions. We focused on seven ecosystem functions associated with genetic and species diversity at three trophic levels (green for primary producer, orange for primary consumer and blue for secondary consumer). Each relationship between biodiversity and ecosystem functions (n = 6 values per function, but for some functions for which irrelevant links were not considered, see the text, n = 32 values in total) was measured at the same trophic level (triangles) or at another trophic level (dots).

Figure 3. Details of the seven causal models linking abiotic parameters, species and genetic diversity and ecosystem functions. Each causal graph (a to g) represents a simplified illustration of the relationships between the two PCA axes synthetizing the environmental parameters of each sampling site (Environmental axis 1 and 2), the species diversity estimated at each trophic level (boxes "Fish", "Shredders" and "Trees"), the genomic diversity estimated from each focal species at each trophic level (boxes "*Phoxinus*", "*Gammarus*" and "*Alnus*"), and each ecosystem function (one model per function). Only the relationships for which the p-value was inferior to 0.20 are indicated for visual simplification. Full arrows indicated positive effects, whereas dotted arrows indicated negative effects. The width of the arrows is proportional to the size of their effects. The percentage of variance explained by environmental and biodiversity effects on ecosystem functions (r^2) is indicated for each function.

Figure 4. General description of individual effect sizes measured between biodiversity estimates and ecosystem functions (BEFs) in a riverine trophic chain. (a) The magnitude and direction of individual effect sizes (*Zr*) of biodiversity is shown for each ecosystem functions as a biplot between *Zr* associated with genetic diversity (y-axis, *genetic* BEFs) measured for one of three target species (*Alnus glutinosa*, *Gammarus sp.* and *Phoxinus dragarum*) and *Zr* associated with species diversity (x-axis, *species* BEFs) measured for one of three trophic levels (trees, invertebrates and fish). For each ecosystem function (but the biomass of trees and of *A. glutinosa*), a total of six *Zr* are depicted in the biplot; four of them are associated with biodiversity measured at another trophic level than the one of the target functions (red symbols, e.g., effect of fish diversity on invertebrate biomass) and two of them are associated with biodiversity measured at the same trophic level than the one of the target functions (blue symbols, e.g., effect of fish diversity on fish biomass). The arrows indicate significant *Zr* (95% confidence intervals excluded 0, see Table S2); vertical arrows are for significant *genetic* BEFs, horizontal arrows are for *species* BEFs. White quadrats stand for situation in which genetic and species BEFs are in the same direction, whereas grey quadrats indicate situation in which genetic and species BEFs are in the opposite direction. Within each quadrat, sub-quadrats indicate the relative magnitude of BEFs, i.e., whether *genetic* BEFs are stronger, weaker or equal in magnitude than *species*

BEFs. (b) Density plots displaying the distribution of individual *Zr* for *species-* and *genetic*
BEFs (dotted and full lines respectively).

Figure 5. Magnitude and direction of the mean effects sizes estimated from the
relationships between biodiversity and ecosystem functions (BEFs) measured in a
riverine trophic chain. (a) The magnitude and direction of BEFs are expressed as effect
sizes (*Zr*) and are displayed according to the facet used to measured biodiversity (genetic or
species diversity, light grey and white boxplots respectively) and to the type of BEFs (*within-*
trophic level BEFs or *between-trophic level* BEFs, triangles and dots respectively). Red
colour and stars indicate global effect sizes that are significantly different from zero (p-value
<0.05). Large symbols are mean ± 95%IC. (b) Same representation as (a) but with details at
each trophic level (mean ± 95%IC, green for primary producer, orange for primary consumer
and blue for secondary consumer). The trophic level at which BEFs are measured is
coherent across all trophic levels.

Figure 1

835
836

837 **Figure 2**

Figure 3

(a) Alnus biomass

(b) Tree biomass

(c) Litter decomposition

(d) Gammarus biomass

845 **Figure 3 (continued)**

(e) Invertebrate biomass

(f) Phoxinus biomass

(g) Fish biomass

846
847

Figure 4

 Figure 5

SUPPLEMENTARY MATERIALS

Table S1. ANOVA table for the linear mixed model testing whether the relationships
between biodiversity and ecosystem functions measured in a riverine trophic chain differ
between the biodiversity facets (species or genetic diversity), the types of BEF (*within-* or
between-trophic levels) and the trophic levels at which BEFs are estimated (primary
producers, primary consumers or secondary consumers). A Wald chi-quare test is used to
test the significance of each fixed effect.

	Degree of freedom	Chisq-value	P-value
(Intercept)	1	1.453	0.228
Biodiversity facet	1	1.293	0.255
Type of BEF	1	7.498	**0.006**
Trophic level	2	1.976	0.372
Biodiversity facet*Type of BEF	1	7.884	**0.005**
Biodiversity facet*Trophic level	2	3.520	0.172
Trophic level*Type of BEF	2	2.901	0.234
Biodiversity facet*Type of BEF*Trophic level	1	2.994	0.224

867 **Table S2 – Estimates of individual effect sizes of BEFs (Zr, n=34) for each ecosystem**
868 **function, each biodiversity facet (genetic or species diversity) and each type of BEF**
869 **(within- or between-trophic levels).** 95% confidence intervals are provided together with
870 the estimate of each BEF. BEFs are considered as significant when the 95%CI does not
871 overlap 0. P-values estimated from t-test are also provided.

Ecosystem function	Predictor	Biodiversity facet	Type of BEF	Zr	negative 95% CI	positive 95% CI	P values
Phoxinus biomass	Fish species diversity	Species diversity	Whithin-trophic levels	-0.529	-0.803	-0.166	0.006
Alnus biomass	Tree species diversity	Species diversity	Whithin-trophic levels	-0.447	-0.695	-0.143	0.006
Phoxinus biomass	*Alnus* genetic diversity	Genetic diversity	Between-trophic levels	-0.321	-0.602	-0.020	0.045
Litter decomposition	Tree species diversity	Species diversity	Whithin-trophic levels	-0.293	-0.630	0.060	0.115
Gammarus biomass	Invertebrate species diversity	Species diversity	Whithin-trophic levels	-0.273	-0.661	0.128	0.194
Litter decomposition	Fish species diversity	Species diversity	Between-trophic levels	-0.210	-0.625	0.210	0.337
Tree biomass	Tree species diversity	Species diversity	Whithin-trophic levels	-0.183	-0.485	0.122	0.250
Fish biomass	Invertebrate species diversity	Species diversity	Between-trophic levels	-0.138	-0.428	0.153	0.506
Phoxinus biomass	Invertebrate species diversity	Species diversity	Between-trophic levels	-0.129	-0.471	0.215	0.362
Invertebrate biomass	Invertebrate species diversity	Species diversity	Whithin-trophic levels	-0.051	-0.381	0.279	0.766
Fish biomass	*Gammarus* genetic diversity	Genetic diversity	Between-trophic levels	-0.047	-0.457	0.364	0.825
Gammarus biomass	*Alnus* genetic diversity	Genetic diversity	Between-trophic levels	-0.043	-0.438	0.352	0.833
Litter decomposition	Invertebrate species diversity	Species diversity	Between-trophic levels	-0.020	-0.401	0.361	0.918
Litter decomposition	*Alnus* genetic diversity	Genetic diversity	Whithin-trophic levels	-0.012	-0.394	0.370	0.950
Litter decomposition	*Phoxinus* genetic diversity	Genetic diversity	Between-trophic levels	0.018	-0.379	0.416	0.930
Invertebrate biomass	*Alnus* genetic diversity	Genetic diversity	Between-trophic levels	0.025	-0.306	0.356	0.885
Invertebrate biomass	Tree species diversity	Species diversity	Between-trophic levels	0.063	-0.236	0.362	0.682
Invertebrate biomass	*Phoxinus* genetic diversity	Genetic diversity	Between-trophic levels	0.085	-0.260	0.429	0.634
Fish biomass	Tree species diversity	Species diversity	Between-trophic levels	0.087	-0.276	0.449	0.644
Invertebrate biomass	*Gammarus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.097	-0.242	0.435	0.580
Litter decomposition	*Gammarus* genetic diversity	Genetic diversity	Between-trophic levels	0.110	-0.281	0.500	0.586
Alnus biomass	*Alnus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.128	-0.179	0.433	0.422
Invertebrate biomass	Fish species diversity	Species diversity	Between-trophic levels	0.144	-0.219	0.504	0.444
Fish biomass	*Phoxinus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.159	-0.260	0.575	0.464
Gammarus biomass	Tree species diversity	Species diversity	Between-trophic levels	0.161	-0.198	0.516	0.388
Phoxinus biomass	Tree species diversity	Species diversity	Between-trophic levels	0.165	-0.100	0.426	0.233
Gammarus biomass	Fish species diversity	Species diversity	Between-trophic levels	0.185	-0.249	0.614	0.413
Fish biomass	*Alnus* genetic diversity	Genetic diversity	Between-trophic levels	0.193	-0.211	0.592	0.360
Phoxinus biomass	*Gammarus* genetic diversity	Genetic diversity	Between-trophic levels	0.228	-0.074	0.522	0.151
Phoxinus biomass	*Phoxinus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.244	-0.064	0.542	0.133
Tree biomass	*Alnus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.280	-0.062	0.609	0.121
Gammarus biomass	*Gammarus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.292	-0.120	0.688	0.178
Fish biomass	Fish species diversity	Species diversity	Whithin-trophic levels	0.326	-0.123	0.753	0.169
Gammarus biomass	*Phoxinus* genetic	Genetic diversity	Between-trophic levels	0.446	0.007	0.830	0.047