Genetic diversity affects ecosystem functions across trophic levels as much as species diversity, but in an opposite direction

Laura Fargeot*[1,2], Camille Poesy[1], Maxim Lefort[1], Jérôme G. Prunier[1], Madoka Krick[1], Rik Verdonck[3], Charlotte Veyssière[2], Murielle Richard[1], Delphine Legrand[1], Géraldine Loot[2] & Simon Blanchet*[1]

[1] Centre National de la Recherche Scientifique (CNRS), Station d'Ecologie Théorique et Expérimentale, UAR 2029, 2 route du CNRS, F-09200 Moulis, France; laura.fargeot@live.fr (L.F.); camille.poesy@orange.fr (C.P.); maxim.lft@gmail.com (M.L.); jerome.prunier@gmail.com (J.G.P.); madokakrick@gmail.com (M.K.); murielle.richard@sete.cnrs.fr (M.R.); simon.blanchet@sete.cnrs.fr (S.B.); delphine.legrand@sete.cnrs.fr (D.L.)

[2] Université Paul Sabatier (UPS) Toulouse III, 118 route de Narbonne, F-31062 Toulouse CEDEX 4, France; geraldine.loot@univ-tlse3.fr (G.L.); veyssiere.charlotte@gmail.com (C.V.)

[3] Center for Environmental Sciences, Environmental Biology, Hasselt University, Diepenbeek, Belgium; rik.verdonck@uhasselt.be (R.V.)

Running title: Genetic vs. species biodiversity effects

Key words: genomic diversity, decomposition, productivity, trophic chains, aquatic ecosystems, fish, shredders, riparian trees, biodiversity-function relationships

Number of words in abstract: 148

Number of words in main text: 4979

Number of references: 44

Number of figures and tables: 7

* Correspondence:

Laura Fargeot and Simon Blanchet

Station d'Ecologie Théorique et Expérimentale

09200, Moulis, France

Phone number: +33 5 61 04 03 61

Laura Fargeot, laura.fargeot@live.fr

Simon Blanchet, simon.blanchet@sete.cnrs.fr

ABSTRACT

Understanding the relationships between biodiversity and ecosystem functioning stands as a

cornerstone in ecological research. Extensive evidence now underscores the profound impact

of species loss on the stability and dynamics of ecosystem functions. However, it remains

unclear whether the loss of genetic diversity within key species yield similar consequences.

Here, we delve into the intricate relationship between species diversity, genetic diversity, and

ecosystem functions across three trophic levels —primary producers, primary consumers, and

secondary consumers— in natural aquatic ecosystems. Our investigation involves estimating

species diversity and genome-wide diversity -gauged within three pivotal species- within each

trophic level, evaluating seven key ecosystem functions, and analyzing the magnitude of the

relationships between biodiversity and ecosystem functions (BEFs). We found that, overall,

the absolute effect size of genetic diversity on ecosystem functions mirrors that of species

diversity in natural ecosystems. We nonetheless unveil a striking dichotomy: while genetic

diversity was positively correlated with various ecosystem functions, species diversity displays

a negative correlation with these functions. These intriguing antagonist effects of species and

genetic diversity persists across the three trophic levels (underscoring its systemic nature),

but were apparent only when BEFs were assessed within trophic levels rather than across

them. This study reveals the complexity of predicting the consequences of genetic and species

diversity loss under natural conditions, and emphasizes the need for further mechanistic

models integrating these two facets of biodiversity.

Introduction

Diversity *within* and *among* species are both important to ensure and stabilize ecosystem functions (Cardinale *et al.* 2012; Raffard *et al.* 2019). Studies on the links between biodiversity and ecosystem functioning (BEFs) have primarily focused on the interspecific (species) facet of biodiversity (Balvanera *et al.* 2006; Hooper *et al.* 2005). However, the intraspecific (genetic) facet of biodiversity has also recently been shown to have substantial effects on ecosystem functions (Crutsinger *et al.* 2006; Hughes & Stachowicz 2004; Reusch *et al.* 2005). Recent meta-analyses have shown that genetic diversity of plant and animal populations affect ecosystem functions, and that the magnitude (and shape) of intraspecific BEFs is similar to that of species diversity (Raffard *et al.* 2019; Wan *et al.* 2022).

Although natural assemblages encompass both intra- and interspecific diversity, most studies investigating BEFs are considering each biodiversity facet separately (but see, Fridley & Grime 2010; Prieto *et al.* 2015; Grele *et al.* 2024). This makes it difficult to differentiate the relative role of genetic and species diversity in ecosystem functions, impeding general predictions regarding the consequences of biodiversity loss as a whole on ecosystem functions (Blanchet *et al.* 2023). For instance, we are currently unaware whether the loss of genetic diversity within a few species in an assemblage is as detrimental for ecosystem functions as a species loss, or whether the combined loss of genetic and species diversity may have non-additive consequences for ecosystem dynamics. Although these biodiversity loss scenarios are realistic, our knowledge on the relative role of genetic *vs*. species diversity in ecosystem functions are still too scarce to provide reliable predictions.

The few studies investigating the combined effects of genetic and species diversity on ecosystem functions were all conducted experimentally by manipulating the genetic and species diversity of assemblages under controlled conditions (Fridley & Grime 2010; Hargrave *et al.* 2011; Prieto *et al.* 2015; but see Grele *et al.* 2024). Our understanding of genetic (intraspecific) and species (interspecific) BEFs therefore relies on simplified ecosystems that often lack variation in other factors (including spatial scales, abiotic factors…), and in which

feedbacks between ecosystem functions and biodiversity are limited (Duffy *et al.* 2017; Prunier

et al. 2023). However, knowledge acquired from BEFs at the interspecific level reveals that

environmental variation can either reduce or enhance the effects of biodiversity on ecosystem

functions, hence generating large variance in the magnitude and direction of BEFs measured

in the wild (Hagan *et al.* 2021; Van Der Plas 2019). One can therefore predict that, under

natural conditions, the relative influence of genetic and species diversity on ecosystem

functions may deviate from what has been quantified under controlled conditions, although it

is difficult to predict the direction of this deviation as field studies (in particular for genetic

BEFs) are too scarce to generate clear predictions. Therefore, we need further realistic field

studies of BEFs, embracing the whole diversity of life forms (from genes to species) and

across realistic environmental gradients to test whether -under natural conditions- species and

genetic BEFs are of similar magnitude.

BEF studies often consider a single trophic level, despite accumulating evidence that

biodiversity at a given trophic level can propagate across trophic levels, generating "multi-

trophic BEFs" (Lefcheck *et al.* 2015; Soliveres *et al.* 2016, Seibold *et al*. 2018). In particular,

studies testing the joint effects of genetic and species diversity on ecosystem functions have

mostly considered the effect of primary producer diversity on their own productivity ("*within-*

trophic level BEFs", e.g., Hargrave *et al.* 2011; Prieto *et al.* 2015). However, genetic and

species diversity within a given trophic level may have propagating effects on the ecosystem

at other trophic levels (hereafter, "*between-trophic level BEFs*"). Indeed, it is predicted that a

genetically-diverse predator population shares their resources more efficiently than a

genetically-poor predator population, which might permit a higher prey species coexistence

and hence a larger prey biomass (*between-trophic level* BEFs due to genetic diversity, e.g.,

Raffard *et al.* 2021). Alternatively, a species-rich community of primary producers likely

exhibits higher primary production, as organisms in species-rich communities share basal

resources more efficiently than in species-poor communities (*within-trophic level BEFs due to*

species diversity, Balvanera *et al.* 2006; Hooper *et al.* 2005). Similarly, the relative impact of

genetic and species diversity should be inconsistent across trophic. At higher trophic levels

(e.g., predators), species richness is generally lower, which should increase the likelihood for

genetic diversity (of a few species) to have strong effects on functions. A simple prediction

might therefore be that the relative impact of genetic diversity on ecosystem functions should

increase with increasing trophic levels (Blanchet *et al.* 2020). Studies considering genetic and

species BEFs under a realistic multitrophic scenario may thus help understanding the trophic

contexts under which either genetic or species diversity is more impactful on ecosystem

functions than the other, and to test whether genetic and species effects can propagate across

trophic levels or not (Seibold *et al.* 2018, Li *et al.* 2020, Moi *et al.* 2021).

Here, we conducted a field study to test the relative importance of genetic and species

diversity for ecosystem functions across multiple trophic levels in a natural landscape. We

focused on three trophic levels from river ecosystems; riparian trees (primary producers),

macroinvertebrate shredders (primary consumers) and fish (secondary consumers). For each

trophic level, we quantified the species diversity of each community, as well as the genetic

diversity of a single target and dominant species (*Alnus glutinosa*, *Gammarus* sp. and

Phoxinus dragarum respectively). We further estimated several ecosystem functions,

including leave decomposition of riparian trees, biomass (as productivity estimates) of each

target species and total biomass of each community within each trophic level. We relied on

causal analyses, taking into account the direct and indirect effects of the environment (through

biodiversity) on ecosystem functions (Duffy *et al.* 2016) to test i) whether BEFs measured at

the genetic level (*genetic* BEFs) are similar in magnitude and direction to BEFs measured at

the species level (*species* BEFs); and ii) whether *within-trophic level* BEFs are similar in

magnitude than *between-trophic level* BEFs. We also tested whether the relative effects of

species and genetic diversity on ecosystem functions (within or between trophic levels) are

consistent across the three trophic levels (primary producers, primary consumers and

secondary consumers), in order to generalize findings along the trophic chain. We predicted

that -contrary to what has been observed under controlled conditions- *genetic* BEFs and

species BEFS will not be similar in magnitude, especially because environmental variation

may modulate each of them differentially. We further expected that significant genetic and

species BEFs will be observed both *within-* and *between*-trophic levels, leading to *within-* and

between-trophic levels of similar magnitude. Finally, we predicted that the magnitude of

genetic BEFs will be higher (than that of *species* BEFs) at the highest trophic level (secondary

consumers) than at the lowest trophic level (primary producers), mainly because species

richness at higher trophic levels presents a lower gradient that at the lowest trophic levels.

Materials and methods

Sampling sites and trophic chain

We sampled 52 sites in Southern France from the Adour-Garonne watershed, and

distributed along an east-west gradient in the Pyrenees Mountains (**Figure 1a**). We acquired

data on species diversity, genetic diversity and ecosystem functions at three trophic levels

(primary producers, primary consumers and secondary consumers) (**Figure 1b**). Riparian

trees (57 species in the sampled area) provide organic matter in the form of fallen leaves as

a food source for decomposers. We selected the common alder *Alnus glutinosa* for acquisition

of genetic data due to its dominance at most sites and its functional relevance, as its roots

serve as shelters for many aquatic species and are involved in nitrogen fixation.

Macroinvertebrate shredders (101 genera in the sampled area) are primary consumers using

leaves as resources, and converting them into accessible organic matter for other species.

We focused on the most abundant Gammarid (Crustacean) species for genetic data

acquisition, referred to as *Gammarus* sp. This species has not yet been formally named

although it is phylogenetically distinct from its closest relative, *Gammarus fossarum* (Carnevali

2022; Piscart, unpublished data). This species is particularly efficient at decomposing tree

leaves, in particular those from *Alnus* (Macneil *et al.* 1997). Fish (20 species in the sampled

area) are secondary consumers feeding on invertebrates (amongst others). We used the

minnow *Phoxinus dragarum* as the fish target species as it is an abundant and important

predator strongly impacting invertebrate communities (Raffard *et al.* 2021).

Biodiversity estimates

Species datasets. At each site, we collected data on the abundance of all species within each

trophic level, at one occasion for trees (July-August 2021) and two occasions for invertebrates

(July and November 2020) and fishes (mid-July to mid-August 2020 and 2021), to obtain

accurate biodiversity estimates. We identified tree species along a 200 m transect of each

river bank, excluding trees with trunk smaller than 2 cm in diameter and more than one meter

away from the bank. The abundances of trees were estimated as the total number of

individuals per species and per site. For invertebrates, we identified shredders to the genus

level (or to the family level for some groups such as chironomids) sampled from two types of

standardized traps installed in four micro-habitats distributed along the 200m transect used to

identify trees: natural coconut brushes (15*5.5 cm, bristles length 7.5 cm) recovered after 1.5

month of colonization, and litter bags (15*11 cm, 0.8 cm mesh size) filled with senescent *Alnus*

leaves from each site and recovered after nine days of colonization (see below). We calculated

abundances of each genus by summing the number of individuals per genus found in the coco

brushes and the litter bags, and we averaged the abundances over the two sampling

occasions to get a single estimate per genus per site. For fish, we collected all specimens

during single-pass electric fishing sessions over a mean area of ~ 469.9 m^2 (± 174 m^2)

distributed along the 200 m transect. We anesthetized, identified and counted individuals at

the species level. We calculated fish abundances as the number of individuals per species

and per m², and we averaged the abundances over the two sampling occasions as for

invertebrates. Fish species number varies from 1 to 11, invertebrate genus number varies

from 15 to 42 and the tree species number varies from 7 to 20 (see Fargeot *et al.* 2023 for

details).

Genetic datasets. At each site, we collected tissue from up to 32 individuals of each of the

three target species, a sample size having found sufficient for estimating the genomic diversity

of populations (Hale *et al*. 2012). We collected fresh leaves of *A. glutinosa* in May 2020,

specimens of *Gammarus* sp. in February 2020, and a piece of pelvic fin from *P. dragarum*

individuals in summer 2020. The DNA of these samples was extracted using commercial kits

for *Alnus* and *Gammarus sp.* and a salt-extraction protocol for *P. dragarum* (see Fargeot *et al.*

2023 for details). For each specimen, DNA concentrations were measured using Qubit 3.0

fluorometer (Life Technologies®, USA). Sequencing was performed based on equimolar pools

of DNA ("pool-seq" approach, Schlötterer *et al.* 2014) from each population and each species.

For *Gammarus sp.*, we also obtained a ~600 bp mitochondrial sequence from the COI

mitochondrial gene from each individual to ensure identification and avoid mixing individuals

from different species. *Gammarus* sp. was found allopatric in most sites, but for a few sites

from the eastern part of the area in which two species were identified (Carnevali 2022). In this

latter case, we conserve only the target species for creating the DNA pools. We created one

DNA pool per site per species (52 pools for *A. glutinosa*, 47 pools for *Gammarus* sp. and 44

pools for *P. dragarum*) and performed double-digest restriction-site associated DNA

sequencing (ddRAD-seq) for *A. Glutinosa* and *Gammarus* sp. (respectively, PstI/MseI and

Pst/HindIII enzymes) and normalized Genotyping-by-Sequencing (nGBS) for *P. dragarum*

(MsII enzyme). Library preparation and pool-sequencing were executed by LGC Genomics

(Biosearch Technologies®, Germany) on an Illumina NovaSeq® (2x150 pb). Data processing

was performed following De Kort *et al.* (2018), except that read mapping was performed on

reference genomes. The genome of *A. glutinosa* was already available (Griesmann *et al.*

2018), and we assembled reference genomes from Illumina short-read sequencing and

PacBio long-read sequencing for *Gammarus. sp.* (available upon request) and *P. dragarum*

(accession number on DDBJ/ENA/GenBank: JARPMJ000000000), respectively. SNP calling

was performed with (i) filtering of raw sequencing files; (ii) indexing of reference genomes; (iii)

mapping reads to the reference; (iv) filtering for unpaired and badly/non-mapped reads; (v)

assembling all read information in a single file per population and per species and (vi)

calculating SNP allelic frequencies (De Kort *et al.* 2018). The total numbers of SNPs retrieved

were 583 862 for *A. glutinosa*, 331 728 for *Gammarus* sp. and 414 213 for *P. dragarum* (see

Fargeot *et al.* 2023 for details).

Species and genetic diversity estimates. We calculated α- diversity per site using the Shannon

entropy from the "hillR" R package for both species and genetic diversity. The Shannon

entropy is a metric of evenness that takes into account the distribution of allele or species

abundances within each site (Chao *et al.* 2014) by weighting each species/allele by its

proportional abundance (q = 1). Results were similar when using the Simpson's diversity index

(q = 2, results not shown). It is noteworthy that -given the spatial extent of the sampling area

and the number of sampling sites- genetic and species diversity estimated in this study

constitutes a fair representation of the biodiversity found in the rivers from the Pyrenean

Piedmont, covering a wide range of biological complexity.

Ecosystem function measurements

At each site, we measured seven ecosystem functions. We collected biomass

production data of all species at each trophic level (hereafter "total biomass") and the biomass

production of each target species as estimates of productivity, as well as the decomposition

rate of *Alnus* leaves. Productivity -as we quantified it- is obviously affected by local

environmental characteristics, and for this reason, we took into account these potential

environmental effects (see hereafter). For riparian tree biomass, we used the trunk diameter

of each single tree as a proxy of individual tree biomass, and we summed the trunk diameters

of all trees found along the transect (divided by the length of the transect) to estimate the total

tree biomass per site and per meter of bank. The same approach was used to estimate *A.*

glutinosa biomass. For macroinvertebrate shredders, we estimated the total invertebrate

biomass by drying all individuals for 24 h at 60 °C before weighing them (10^{-4} g precision). The

same procedure was used to estimate the biomass of *Gammarus* sp. For both estimates, we

averaged biomasses over the two sampling sessions. For fish, (fresh) total fish biomass was

estimated as the total weight of all individuals (0.01 g precision) per site, whereas *P. dragarum*

biomass was the mass of all *P. dragarum* specimens per site. Fish biomasses were averaged

over the two sampling sessions.

For the decomposition rate, we quantified leaf mass loss in litter bags placed in four

micro-habitats per site twice (July and November 2020). We gathered and dried senescent

leaves during fall 2019 from five *Alnus* trees per site to limit individual-specific effects on

decomposition. Litter bags were 15 cm x 11 cm pockets of plastic-wire mesh (mesh size; 8

mm to allow invertebrates colonization) in which we introduced 4g of dried leaves before

closing the bags with staples. We installed three bags per micro-habitat (12 per site) that we

removed sequentially after ~9 days, ~18 days and ~27 days respectively to estimate

decomposition rates. Bags were brought back to the laboratory, the remaining leaves were

cleaned, dried and weighed. Decomposition rate was estimated as the slope of leaf mass loss

over time (obtained from a linear model) that we averaged across replicates and temporal

sessions (Raffard *et al.* 2021).

Environmental data

A major challenge for inferring BEFs from empirical data is to take into account the

direct and indirect (through biodiversity) effects of environmental factors on ecosystem

functions (Duffy *et al.* 2016, 2017). Failing to do this may result in overestimated and/or

artefactual BEFs, especially if the same environmental factor simultaneously affects

biodiversity and ecosystem processes (Grace *et al.* 2016). For each site, we measured

thirteen variables related to river topography and physico-chemical characteristics that likely

influence biodiversity and ecosystem processes (Altermatt 2013). *River bed width* (m) was

averaged from five measurements per site. *Connectivity* was calculated as the "closeness

centrality", *i.e.,* the inverse of the sum of the distances of a node to all other nodes along the

shortest paths possible (Altermatt 2013), using QGIS and the "RiverDist" R package. *Altitude*

(m), *distance from the outlet* (m) and *east-west gradient* (longitudinal position along the

Pyrenees chain) were measured using QGIS; *oxygen concentration* (mg.L^{-1}), *oxygen*

saturation (%), *water temperature* (°C), *specific conductivity* (μS/cm) and *pH* were measured

(and averaged) in summers 2020 and 2021 using a multi-parameter probe (Aqua TROLL 500,

In-Situ Inc.). Concentration of NO_{3-}, NO_{2-}, NH_4^+ and PO_4^{3-} were estimated (and averaged)

during summers 2020 and 2021 from a filtered water volume (100 mL) using the Alpkem Flow

Solution Iv Autoanalyzer (OI Analytical®).

A Principal Component Analysis combining all thirteen variables was performed using

the R package "ade4" (Dray & Dufour 2007), and coordinates of each site on the two first axes

(38.03% of the total variance, see **Table 1**) were used as two synthetic environmental

variables for further analyses. We kept only these two first axes to avoid collinearity and over-

parameterization of subsequent models. The first axis is defined by a strong contribution of (in

decreasing order) oxygen concentration and altitude (**Table 1**). The second axis is defined by

a strong contribution of east-west gradient and connectivity (**Table 1**).

Statistical analyses

BEF relationships. To quantify the magnitude of association between biodiversity estimates

and ecosystem functions (BEFs), we performed piecewise Structural Path Models (pSEM,

"piecewiseSEM" package, Lefcheck 2016). pSEM allows modelling direct and indirect causal

relationships among a set of response variables and predictors (Shipley 2009). Further, pSEM

uses local estimates of each linear structural equation separately (*i.e.*, parameters are

estimated from a series of independent models forming a general causal graph), which allows

the inclusion of a large number of parameters despite modest sample sizes (Shipley 2009).

We ran a pSEM for each ecosystem function separately (*i.e.*, seven pSEMs, see an example

in **Figure 2**). In each pSEM, the ecosystem function was the dependent variable whereas the

six biodiversity estimates (species and genetic diversity estimated for each trophic level) and

the two synthetic environmental variables were the predictors. In each model, environmental

predictors were allowed to explain each biodiversity estimate (indirect effects of environmental

variables through their influence on biodiversity, see **Figure 2**). For some functions (in

particular those associated with plant biomass), irrelevant biodiversity-functions links were not

included (e.g., the impact of fish or invertebrate diversity on tree biomasses), which results in

34 BEFs (out of the 42 possible links) having been included in the meta-regression (see

hereafter).

From each pSEM model, we retrieved the local parameter (standardized estimate, an

equivalent to a coefficient of correlation) associated with the direct effect of each biodiversity

estimate (six per function, but for some functions for which ecology-irrelevant BEFs were

excluded) on the function (coloured arrows in **Figure 2**), which provides both the magnitude

and the direction of each BEF. To smoothen comparison, we calculated a standardized effect

size for each BEF by applying the Fisher's Z transformation (*Zr*) to the standardized estimates.

Positive *Zr* indicate positive associations between biodiversity and ecosystem functions,

whereas negative *Zr* indicate negative relationships. The higher the absolute value of Zr, the

higher the strength of the association. *Zr* therefore indicate both the direction (positive or

negative) and the magnitude of the associations. Our seven measures of ecosystem functions

were not correlated one to each other (all $r_{pearson}<|0.39|$).

Direction and magnitude of all types of BEFs. We used a linear mixed-model to test (i) whether

the magnitude and direction of *genetic* BEFs are similar to those of *species* BEFs, and (ii)

whether *within-trophic level* BEFs are similar in effect size to *between-trophic level* BEFs. In

this model, *Zr* (providing the direction and magnitude of each BEF, n=34) was the dependent

variable, and the predictors were the diversity facets used to measure biodiversity (genetic or

species diversity) and the type of BEF (*within-trophic* or *between trophic* levels, triangles vs.

dots in **Figure 2**). We included the two-term interaction between diversity facet and type of

BEF to test whether the magnitude and direction of *genetic* and *species* BEFs are consistent

across *within-trophic level* and *between-trophic level* BEFs. We further included in this model

the type of ecosystem function as a random term (to take into account that each ecosystem

function was associated with several biodiversity estimates) as well as the inverse of the

asymptotic variance (v_z=n-3) associated with each effect size as a weighting parameter for

each case study (Balvanera *et al.* 2006; Raffard *et al.* 2019).

We ran an additional linear mixed-model similar to the previous one, except that we

added as a fixed effect the trophic level at which biodiversity was measured to estimate BEFs

(primary producers, primary consumers or secondary consumers) as well as all interaction

terms. Interaction terms allow testing the consistency of major conclusions across trophic

levels, thereby determining the extent to which our findings can be generalized along the

trophic chain. Models were run using the lmer function ("lme4" package) and significance of

fixed effects was determined using type III ANOVA with Wald chi-square tests (function Anova

from the "car" R package, α=0.05).

Results

Details of causal models linking environmental parameters, species and genetic

diversity and ecosystem functions are graphically depicted in **Figure 3a-g**. Note that only

relationships for which p-values were below 0.20 are shown on these graphs. This threshold

was chosen arbitrarily to provide readable causal graphs and to highlight only on the most

biologically relevant relationships.

The percentage of variance in ecosystem function explained by the environment and

biodiversity varies from 10% (invertebrate biomass, **Figure 3e**) to 55% (*Phoxinus* biomass,

Figure 3f) and was moderate overall. For all functions but the three biomass, part of the

variance was (directly) explained by at least one out of the two environmental PCA axes. For

some functions (e.g., *Phoxinus* biomass, Figure 3f), there was a combined effect of several

biodiversity estimates, whereas for other functions (e.g., *Alnus* biomass, Litter decomposition,

Figure 3a, 3c) the effect of a single biodiversity estimate predominates. Overall, direct

environmental effects on ecosystem functions did not predominate, and environmental effect

sizes were similar (in strength) to that of biodiversity effects, showing the non-negligible role

of biodiversity for ecosystem functions in the wild.

Individual effect sizes (*Zr*) measured between biodiversity estimates and ecosystem

functions were weak to moderate, irrespectively of the considered ecosystem function and of

the type of BEFs (*genetic/species BEFs*, *within-trophic level/ between-trophic level* BEFs)

(**Figure 4a, 4b, Table S2**). As expected under natural conditions (Hagan *et al.* 2021), BEFs

ranged from negative to positive, and their distribution were centred around 0, although we

observed a slight tendency for *genetic* BEFs toward positive values (**Figure 3b**). Only four out

of the 34 BEFs were strong and significant; two significant BEFS concerned *species* BEFs

(negative relationship between the biomass of *A. glutinosa* and the diversity of trees, Zr = -

0.446, 95% CI [-0.695, -0.143]; negative relationship between the biomass of *P. dragarum* and

the diversity of fish, Zr = -0.529, 95% CI [-0.802, -0.166]) and two concerned *genetic* BEFs

(negative relationship between the biomass of *P. dragarum* and the diversity of *A. glutinosa*,

Zr = -0.321, 95% CI [-0.602, -0.019]; positive relationship between the biomass of *Gammarus*

sp. and the diversity of *P. dragarum*, Zr = 0.446, 95% CI [0.001, 0.829]) (**Figure 4a**).

Noteworthily, for *within-trophic* BEFs, most case studies fall into the category whereby genetic

BEFs tend to be positive and species BEFs tend to be negative (grey bottom-right square in

Figure 4a).

We confirmed this visual tendency by summarizing all individual Zr through a meta-

regression. Indeed, we found a significant interaction between the facet at which biodiversity

is measured (genetic or species diversity), and the type of BEF that was measured (within- or

between trophic levels; **Table 2**). This interaction indicates (i) that -overall- *within-trophic level*

BEFs were significantly negative when considering species diversity ($Zr_{\text{Within*Species}}$ = -0.185,

95% CI [-0.343, -0.027]), whereas *within-trophic level* BEFs were significantly positive when

considering genetic diversity ($Zr_{Within*Genetic}$ = 0.168, 95% CI [0.010, 0.326], see **Figure 5a**), and

(ii) that this pattern was not observed for *between-trophic levels* BEFs, where no particular

trend was observed (**Figure 5a**). Although most individual Zr were weak to moderate (and not

significant), their consistency (in term of magnitude and direction) resulted in a significant

pattern whereby species and genetic diversity have opposite effects on ecosystem functions

for *within-trophic level* BEFs; species diversity is negatively associated, whereas genetic

diversity is positively associated with ecosystem functions, but only when the influence of

biodiversity on ecosystem functions is measured within the same trophic level.

When including the trophic level at which biodiversity is measured, we found no

significant interaction terms between trophic levels and other fixed effects nor any additive

effect of trophic levels (see **Table S1**). This indicates that our main findings were consistent

across trophic levels, *i.e.*, the respective negative and positive effects on ecosystem functions

of species and genetic diversity hold statistically true across all trophic levels (**Figure 5b**).

Discussion

We provide empirical evidence that, in natural ecosystems, the effect sizes of genetic

and species diversity on multi-trophic ecosystem functions are of similar magnitude, but

operate in opposite directions. Indeed, for BEFs measured within the same trophic level, the

effects of species diversity across multiple ecosystem functions were moderately negative on

average, whereas the effects of genetic diversity were moderately positive. This suggests an

antagonistic effect between the genetic and the species components of biodiversity in the

modulation of ecosystem functions within one trophic level. This antagonistic effect was not

identified for BEFs measured across trophic levels, since in these cases the influence of both

genetic diversity and species diversity across multiple ecosystem functions was generally not

different from zero. These conclusions hold true across three trophic levels (plants,

invertebrates and fish), indicating that the relative effects of genetic and species diversity on

ecosystem functions are not limited to a specific trophic level.

Our study is one of the few field-based study revealing BEFs across an entire (riverine)

food chain spanning from primary producers to secondary consumers. Indeed, most previous

BEF studies in the field focused on a single trophic level, and predominantly on terrestrial

primary producers (Duffy *et al.* 2017; Van Der Plas 2019, but see e.g., Li et al. 2020, Moi et

al. 2021). This permitted encompassing a broad range of ecosystem functions that depict the

overall functioning of a riverine ecosystem (rather than focusing on a single compartment).

Moreover, we focused both on the effects of genetic and species diversity on these ecosystem

functions, which has rarely (if not ever) been evaluated so far and which provides an

exhaustive overview of BEFs in the wild. Our causal analyses also statistically took into

account the direct (and indirect) effect of environmental factors on ecosystem functions, which

is a prerequisite to isolate biodiversity effects. Nonetheless, causal relationships obtained from

observational data (rather than from experimental data) are notoriously difficult to infer and

must therefore be interpreted with care (Duffy *et al*. 2017). As a result, the BEFs we estimated

display strong variability (ranging from negative to positive values) and a very few of them (4

out of 34) were statistically significant according to conventional tresholds. Although the

statistical inferences made in this study are based on a large sample size, it is noteworthy that

the general patterns we will describe hereafter (and their interpretation) have to be considered

with care, as we can not rule out the possibility that some patterns might arise because of

statistical biases rather than biological reality. Nonetheless, we -as ecologists- feel important

to provide such a general picture from field data (even if partially distort by statistical limits),

as this represents basic patterns that we have to understand.

We revealed that direct environmental effects on ecosystem functions were (in

average) not stronger in intensity than biodiversity effects, which is coherent with previous

syntheses on species BEFs in the wild (Duffy *et al*. 2017). Furthermore, environmental factors

used to describe sampling sites in this study were not strong predictors of species and genetic

biodiversity. Two non-exclusive hypotheses may explain this observation: (i) using PCA axes

to resume environmental gradients may blur some specific environment-biodiversity links, and

(ii) as shown and explained in a companion paper (Fargeot *et al*. 2023), the East-West

gradient used in this study (rather than a classical upstream-downstream gradient) intrinsically

limits the potential for strong environmental effects on biodiversity (which was the purpose of

this sampling design). Nonetheless, after accounting for these environmental covariates, we

found that most individual BEFs (either *genetic* or *species*, *within-trophic levels* or *between-*

trophic levels BEFs) were weak to moderate in magnitude, and that they operated almost

equally in both direction (i.e., positive and negative association between biodiversity and

ecosystem functions). As such, the distribution of individual effect sizes was centred around

0, for both *genetic* and *species* BEFs. Accordingly, there were only four individual BEFs that

were significant, three out of them were negative and one was positive (**Table S2**). This

general pattern (low to moderate BEFs with both positive and negative direction) is actually

consistent with the most exhaustive meta-analysis having synthesized the magnitude and

direction of *species* BEFs in the wild (Van Der Plas 2019) and with recent conceptual works

(Hagan *et al.* 2021) concluding that strong and positive BEFs should not be the norm in natural

ecosystems, but rather that a mix of positive, neutral and negative BEFs are expected. Our

empirical findings are consistent with this conclusion.

We focused both on *within-trophic level* and *between-trophic level* BEFs, which likely

encompasses a broad array of mechanisms sustaining potential associations between

biodiversity and ecosystem functions. For instance, two out of the four significant BEFs we

reveal are negative association between species diversity (fish or tree species diversity

respectively) and the biomass production of one of the target species (*Phoxinus sp*. and *Alnus*

sp. respectively). These *within-trophic level* BEFs can -for instance- arise either because, if

resources are limited, increased number of species within a patch limit the biomass production

of each individual species, or because of a poorer competitive ability of the target species

under some environmental conditions, which favors the settlement of additional species.

Teasing apart these two hypotheses is difficult and further studies are needed to isolate

underlying mechanisms. The other two significant BEFs concerned the association (either

positive or negative) between the genetic diversity of a target species (*Alnus sp.* or *Phoxinus*

sp.) with the biomass production of another target species (*Phoxinus sp.* or *Gammarus sp.,*

respectively). These *between-trophic level* BEFs likely arise through indirect effects implying

the diversity and availability of (prey) resources. An obvious limit of this field-based study is

the impossibility to tease out these mechanisms. Another limit is associated with the fact that,

although environmental covariates were taken into account in causal models, they were

synthetized by two PCA axes, and we cannot ensure that all potential environmental

covariates have been taken into account (see above). This can influence the actual estimates

of BEFs in the wild (Duffy *et al*. 2017). Nonetheless, it is noteworthy that we have previously

shown that -in this dataset- species and genetic diversity were not correlated one to each other

and that each biodiversity facet was sustained by different environmental predictors (Fargeot

et al. 2023, see also **Figure 3**). This implies that environmental and biodiversity effects inferred

in this study should not be strongly distorted by collinearities and can -in theory- be interpreted

independently one from each other (e.g., the positive effect *Phoxinus* genetic diversity on

Phoxinus biomass is independent from the negative of fish species diversity as the two

estimates of biodiversity do not co-vary, see **Figure 3f**).

Keeping limitations associated with field-based studies into account (see above),

revealing associations between ecosystem functions and species and genetic biodiversity (or

the lack of) in natural ecosystems is an important step forward to set theoretical and

experimental approaches aiming at understanding this complex biological reality. Beyond

individual BEF case studies (that were not the main aim of this study), their aggregation across

trophic levels and biodiversity facets revealed a clear (and statistically supported) pattern

whereby, within trophic levels, genetic and species diversity display antagonistic association

with ecosystem functions; the global effect of species diversity across multiple ecosystem

functions was negative, whereas the global effect of genetic diversity was positive. This pattern

emerges from the "cumulative" effects of weak to moderate associations between biodiversity

and ecosystem functions that consistently point toward the same direction (positive for genetic

diversity, negative for species diversity), emphasizing the meaningfulness of meta-regressions

(and more generally approaches based on effect sizes rather than on p-values) to reveal

biological patterns. We hereafter discuss the ecological relevance of this general pattern.

Our results confirm a previous meta-analysis demonstrating that genetic and species

diversity modulates ecosystem functions with a similar magnitude (Raffard *et al.* 2019), and

results from few experimental studies that manipulated both the genetic and species

components of biodiversity under controlled conditions (e.g., Jiang *et al.* 2022; Prieto *et al.*

2015). Indeed, within trophic levels, the absolute mean effect size of genetic and species

diversity across ecosystem functions were of the same magnitude ($|Zr|$ = 0.168 and 0.185 for

genetic and species diversity effects respectively), and slightly greater than the effect sizes

reported under controlled conditions ($|lnRR|$ = 0.132 and 0.134 for genetic and species

diversity effects respectively, Raffard *et al.* 2019) and those more generally reported for

species BEFs ($|Zr|$ = 0.101, Balvanera *et al.* 2006). Although comparing effect sizes among

studies that strikingly differ in their spatial coverage (small or large spatial scale), their

taxonomic focus (e.g., primary producers *vs*. predators, species *vs*. genetic diversity…) and/or

their approaches (experimental *vs*. observational studies) is questionable (especially given

the non-linear nature of BEFs), our findings suggests for the first time that under natural

conditions, the effects of genetic and species components of biodiversity on ecosystem

functions are comparable. However, our study goes two steps further as (i) it extends the

conclusion made by Raffard *et al.* (2019) to multiple trophic levels and (ii) it suggests that the

effects of genetic and species BEFs can actually operate in opposite directions.

As pointed out by Raffard *et al.* (2019), the vast majority (91% of 23 reviewed studies

by 2019, see also Wan *et al.* 2022) of studies investigating the effects of genetic diversity on

ecosystem functions have focused on primary producers, and all of them were based on

experiments, which is also the case for most studies manipulating both genetic and species

diversity. These trends strongly hamper any generalization. On the contrary, our findings

provide a solid support for broadening the conclusion that both genetic and species diversity

can influence ecosystem functions in the wild. More strikingly, our results suggests that,

although the absolute effect sizes of genetic and species BEFs are of similar magnitude, for

within-trophic level BEFs, the direction of their effects are opposite; species diversity (in

general) reduces the rate of ecosystem functions, whereas genetic diversity enhances the

same functions. For instance, all other things being equal, higher fish species diversity is

associated with a lower productivity (biomass) in *P. dragarum* (see above for potential

explanations), whereas its own genetic diversity tends to be associated with a higher

productivity (**Table S2**). In this specific case, genetic and species diversity of the same trophic

group (fish) tended to have opposite effects on the same function (productivity of *P. dragarum*).

However, in most cases this was not the case as genetic diversity was positively associated

with some functions, whereas species was negatively associated with other functions. The

distinction between these two patterns is important as in the latter case (genetic and species

diversity are associated with different functions) managing/conserving the intra- and

interspecific diversity of a single trophic group (e.g., trees) can alter more than one ecosystem

function, and sometimes functions that are even not directly associated to the managed trophic

group. Moreover (and importantly), as genetic and species diversity have been found to be

uncorrelated spatially in this landscape (Fargeot *et al.* 2023), covariation among diversity

estimates cannot explain these patterns. These antagonistic effects of genetic and species

diversity on ecosystem functions parallel previous experimental findings on plants (Hazard *et*

al. 2017; Tang *et al.* 2022). It is now essential to understand the mechanisms sustaining these

antagonistic effects as a step forward.

S*pecies* BEFs were on average negative (see **Table S2** for individual estimates), which

contrasts with the general view that species biodiversity favours ecosystem functions,

although it is not that surprising (Dee *et al.* 2023; Hagan *et al.* 2021). Indeed, the net effect of

species biodiversity on ecosystem functions results from the combined effects of both negative

factors, arising from antagonistic interactions such as negative complementarity or negative selection effect, and positive factors, arising from beneficial interactions such as niche complementarity or facilitation (Loreau & Hector 2001). We can speculate that, in our case, the net effect of interspecific interactions mostly results from negative complementarity among species (or strong negative selection effect), whereas the net effect of intraspecific interactions may result from facilitative interactions and/or improved niche complementarity with increased genetic diversity. Intraspecific competition is generally stronger than interspecific competition (Connell 1983), and intraspecific interactions could be expected to lead more frequently to negative complementarity (and hence negative *genetic* BEFs) than interspecific interactions. Since we observe the opposite, we can hypothesize that genetic diversity is essential to increase niche complementarity within species (Bolnick *et al.* 2003) and hence to reduce the pervasive effects of intraspecific interactions (Hughes *et al.* 2008; Prunier *et al.* 2023). Given the empirical nature of our study and the fact that our meta-regressive approach includes several types of BEFs (e.g., species richness acting either on the biomass of a single focal species or on the biomass of an entire focal community), it is hard to tease apart specific and underlying mechanisms. Theoretical approaches, modelling simultaneously the genetic and species components of biodiversity, would be extremely useful to reveal the mechanisms sustaining opposite effects of intra- and interspecific diversity on ecosystem functions.

These antagonistic effects were observed only for BEFs measured *within* trophic levels, not for those measured *between* trophic levels. An overall *between-trophic level* BEF not different from zero suggests that biodiversity at a trophic level has only limited impact on ecosystem functions at another trophic level. For example, the biomass of *P. dragarum* was primarily influenced by genetic and species diversity in fishes, rather than the diversity of their preys (**Table S2**). However, for both genetic and species estimates of biodiversity, there was a substantial variation in effect sizes for *between-trophic level* BEFs that ranged from negative to positive BEFs (**Figure 5**). This suggests that biodiversity effects across trophic levels may be more variable in their direction than within-trophic level BEFs, which appear as more

constrained. Variability in the magnitude and direction of effect sizes for *between-trophic level*

BEFs likely blur a more general trend, but this variation is actually expected under natural

conditions in which interactions involve multiple prey and predator species, fostering co-

adaptation among communities from different trophic levels (Aubree *et al.* 2020; Poisot *et al.*

2013). In these cases, trophic complementarity between two trophic levels (i.e., the originality

of a species based on the identity of the species it interacts with) might be a stronger

determinant of ecosystem functions than complementarity measured at either one of the two

trophic levels (Poisot *et al.* 2013). Quantifying trophic complementarity among our three target

species (and communities) using stable isotope or gut content analyses for instance would be

extremely valuable to assess whether this complexity can better explain BEFs between trophic

levels than diversity measured at one of the trophic level (Aubree *et al.* 2020).

The empirical patterns we revealed here were all extremely consistent across the three

trophic levels, hence allowing generalization. It is noteworthy that, although statistically strong

and consistent, these patterns must be interpreted with care as field-based approaches are

limited in properly taking into account the environmental heterogeneity of natural ecosystems

(Hagan *et al.* 2021). BEFs were not particularly stronger at any specific trophic level and the

relative effects of genetic and species diversity were not dependent on the trophic level at

which the function was estimated. We may have expected a stronger top-down regulation (i.e.,

biodiversity of predators has more effects than biodiversity of preys) of ecosystem functions

since previous studies showed that biodiversity loss should have greater consequences for

multi-functionality when it occurs at higher trophic levels (Lefcheck *et al.* 2015, Seibold *et al.*

2018). For instance, increased genetic diversity within a predatory fish species has

experimentally been shown to indirectly increase the rate of litter decomposition by increasing

the diversity of shredders (Raffard *et al.* 2021). Similarly, the relative effects of genetic and

species diversity on functions may have varied among trophic levels, and in particular the

relative importance of genetic diversity may have been higher for species-poor trophic levels

(i.e., fish community) because of a "compensatory effect". We found no evidence for these

potential trophic-level dependencies, but instead found extremely consistent patterns, which,

from a broader perspective, reveal the importance of integrating both multi-trophic and multi-

faceted approaches in predicting the overall consequences of biodiversity loss on ecosystem

functioning.

To conclude, we found that the genetic (intraspecific) and species (interspecific) facets

of biodiversity are both important drivers of multiple ecosystem functions in a natural and multi-

trophic context. In the wild, these two facets of biodiversity can, as expected, generate low to

moderately high impacts on ecosystem functions measured across three trophic levels, and

they can operate in opposite directions (but on different functions; genetic diversity is positively

associated with some functions, species diversity is negatively associated with other

functions). This shows the importance for managers to develop integrative conservation plans

spanning the entire diversity of life (from genes to species). For instance, genetic diversity loss

often precedes species loss, and our results suggest that -in mountain streams- losing genes

may actually be particularly detrimental for the performance of ecosystem functions. As such,

it appears essential to maintain populations with high levels of genetic diversity in these

ecosystems. Future studies should (i) extend these findings to other ecosystems and by

quantifying natural genetic variation in more than a single species per trophic level, (ii)

generate theoretical predictions regarding the mechanisms sustaining the antagonistic effects

of genetic and species diversity on functions we revealed, and (iii) use a broader integrative

approach for estimating biodiversity across facets (*inclusive* biodiversity) by using either a

trait-based approach or a genetic-based approach as recently proposed by Blanchet *et al.*

(2023) and Loreau *et al.* (2023).

AUTHOR CONTRIBUTION

Laura Fargeot: Conceptualization (Supporting); Methodology (Equal); Software (Equal); Validation (Equal); Formal analysis (Lead); Investigation (Lead); Data curation (Lead); Writing - original draft (Lead); Writing - review & editing (Equal); Visualization (Lead); Supervision (Equal); Project administration (Lead). **Camille Poesy:** Methodology (Equal); Validation (Equal); Investigation (Lead); Data curation (Supporting); Supervision (Equal); Project administration (Supporting). **Maxim Lefort:** Methodology (Supporting); Investigation (Lead); Data curation (Supporting); Supervision (Equal); Project administration (Supporting). **Jérôme G. Prunier:** Methodology (Supporting); Software (Equal); Formal analysis (Supporting); Investigation (Supporting); Resources (Equal); Data curation (Supporting); Writing - original draft (Supporting); Writing - review & editing (Supporting). **Madoka Krick:** Methodology (Supporting); Investigation (Equal). **Rik Verdonck**: Methodology (Supporting); Software (Equal); Investigation (Supporting); Writing - review & editing (Supporting). **Charlotte Veyssière:** Methodology (Supporting); Investigation (Equal). **Murielle Richard:** Methodology (Supporting); Investigation (Supporting); Writing - review & editing (Supporting). **Delphine Legrand:** Validation (Equal); Writing - review & editing (Equal); Visualization (Supporting). **Géraldine Loot:** Validation (Equal); Investigation (Supporting); Writing - review & editing (Supporting); Visualization (Supporting). **Simon Blanchet:** Conceptualization (Lead); Methodology (Lead); Software (Supporting); Validation (Equal); Formal analysis (Supporting); Investigation (Lead); Resources (Equal); Data curation (Supporting); Writing - original draft (Supporting); Writing - review & editing (Lead); Visualization (Lead); Supervision (Lead); Project administration (Lead); Funding acquisition (Lead).

REFERENCES

Altermatt, F. (2013). Diversity in riverine metacommunities: a network perspective. *Aquatic Ecology*, 47, 365–377.

Aubree, F., David, P., Jarne, P., Loreau, M., Mouquet, N. & Calcagno, V. (2020). How community adaptation affects biodiversity–ecosystem functioning relationships. *Ecol Lett*, 23, 1263–1275.

Balvanera, P., Pfisterer, A.B., Buchmann, N., He, J.-S., Nakashizuka, T., Raffaelli, D., *et al.* (2006). Quantifying the evidence for biodiversity effects on ecosystem functioning and services: Biodiversity and ecosystem functioning/services. *Ecol Lett*, 9, 1146–1156.

Blanchet, S., Fargeot, L. & Raffard, A. (2023). Phylogenetically-conserved candidate genes unify biodiversity–ecosystem function relationships and eco-evolutionary dynamics across biological scales. *Mol Ecol*, 32, 4467–4481.

Blanchet, S., Prunier, J.G., Paz-Vinas, I., Saint-Pé, K., Rey, O., Raffard, A., *et al.* (2020). A river runs through it: The causes, consequences, and management of intraspecific diversity in river networks. *Evol Appl*, 13, 1195–1213.

Bolnick, D.I., Svanback, R., Fordyce, J.A., Yang, L.H., Davis, J.M., Hulsey, C.D., *et al.* (2003). The ecology of individuals: Incidence and implications of individual specialization. *Am Nat*, 161, 1–28.

Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., *et al.* (2012). Biodiversity loss and its impact on humanity. *Nature*, 486, 59–67.

Carnevali, M. (2022). Etude phylogénétique des populations de Gammares (Gammarus sp.) du piedmont Pyrénéen. Master thesis. University of Toulouse, France.

Chao, A., Chiu, C.-H. & Jost, L. (2014). Unifying Species Diversity, Phylogenetic Diversity, Functional Diversity, and Related Similarity and Differentiation Measures Through Hill Numbers. *Annu Rev Ecol Evol Syst*, 45, 297–324.

Connell, J.H. (1983). On the prevalence and relative importance of interspecific competition :

evidence from field experiments. *Am Nat*, 122, 661–696.

Crutsinger, G.M., Collins, MD, Fordyce, J.A., Gompert, Z., Nice, C. & Sanders, N.J. (2006).

Plant Genotypic Diversity Predicts Community Structure and Governs an Ecosystem

Process. *Science*, 313, 966–968.

De Kort, H., Baguette, M., Prunier, J.G., Tessier, M., Monsimet, J., Turlure, C., *et al.* (2018).

Genetic costructure in a meta-community under threat of habitat fragmentation. *Mol*

Ecol, 27, 2193–2203.

Dee, L.E., Ferraro, P.J., Severen, C.N., Kimmel, K.A., Borer, E.T., Byrnes, J.E.K., *et al.* (2023).

Clarifying the effect of biodiversity on productivity in natural ecosystems with

longitudinal data and methods for causal inference. *Nat Commun*, 14, 2607.

Dray, S. & Dufour, A.-B. (2007). The **ade4** Package: Implementing the Duality Diagram for

Ecologists. *J Stat Soft*, 22.

Duffy, J.E., Godwin, C.M. & Cardinale, B.J. (2017). Biodiversity effects in the wild are common

and as strong as key drivers of productivity. *Nature*, 549, 261–264.

Duffy, J.E., Lefcheck, J.S., Stuart-Smith, R.D., Navarrete, S.A. & Edgar, G.J. (2016).

Biodiversity enhances reef fish biomass and resistance to climate change. *P Natl Acad*

Sci, 113, 6230–6235.

Fargeot, L., Poesy, C., Lefort, M., Prunier, J.G., Krick, M., Veyssière, C., *et al.* (2023).

Genomic and species diversity patterns across multiple trophic levels in riverscapes.

Oikos, 2023, e10015.

Fridley, J.D. & Grime, J.P. (2010). Community and ecosystem effects of intraspecific genetic

diversity in grassland microcosms of varying species diversity. *Ecology*, 91, 2272–

2283.

Grace, J.B., Anderson, T.M., Seabloom, E.W., Borer, E.T., Adler, P.B., Harpole, W.S., *et al.*

(2016). Integrative modelling reveals mechanisms linking productivity and plant

species richness. *Nature*, 529, 390–393.

Griesmann, M., Chang, Y., Liu, X., Song, Y., Haberer, G., Crook, M.B., *et al.* (2018).
Phylogenomics reveals multiple losses of nitrogen-fixing root nodule symbiosis.
Science, 361, eaat1743.

Grele, A., Massad, T.J., Uckele, K.A., Dyer, L.A, Antonini, Y., Braga, L., Forister, M.L., Sulca, L., Kato, M.,
Lopez, H.G., Nascimento, A.R., Parchman, T., Simbaña, W.R., Smilanich, A.M., Stireman, J.O.,
Tepe, E.J., Walla, T. & Richards, L.A. (2024). Intra- and interspecific diversity in a tropical plant
clade alter herbivory and ecosystem resilience. *eLife*, 12, RP86988

Hale, M.L., Burg, T.M. & Steeves, T.E. (2012). Sampling for microsatellite-based population genetic
studies: 25 to 30 individual per population is enough to accurately estimate allele frequencies.
PloS ONE, 7, e45170

Hagan, J.G., Vanschoenwinkel, B. & Gamfeldt, L. (2021). We should not necessarily expect
positive relationships between biodiversity and ecosystem functioning in observational
field data. *Ecol Lett*, 24, 2537–2548.

Hargrave, C.W., Hambright, K.D. & Weider, L.J. (2011). Variation in resource consumption
across a gradient of increasing intra- and interspecific richness. *Ecology*, 92, 1226–
1235.

Hazard, C., Kruitbos, L., Davidson, H., Taylor, A.F.S. & Johnson, D. (2017). Contrasting
effects of intra- and interspecific identity and richness of ectomycorrhizal fungi on host
plants, nutrient retention and multifunctionality. *New Phytologist*, 213, 852–863.

Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., *et al.* (2005).
Effects of biodiversity on ecosystem functioning: A consensus of current knowledge.
Ecol Monogr, 75, 3–35.

Hughes, A.R., Inouye, B.D., Johnson, M.T.J., Underwood, N. & Vellend, M. (2008). Ecological
consequences of genetic diversity: Ecological effects of genetic diversity. *Ecol Lett*, 11,
609–623.

Hughes, A.R. & Stachowicz, J.J. (2004). Genetic diversity enhances the resistance of a
seagrass ecosystem to disturbance. *Proc Natl Acad Sci U.S.A.*, 101, 8998–9002.

Jiang, M., He, L., Fan, B., Wang, T., Yang, N., Liu, Y., *et al.* (2022). Intraspecific more than

interspecific diversity plays an important role on Inner Mongolia grassland ecosystem

functions: A microcosm experiment. *Sci Total Env* 826, 154134.

Lefcheck, J.S. (2016). PIECEWISESEM : Piecewise structural equation modelling in R for

ecology, evolution, and systematics. *Methods Ecol Evol*, 7, 573–579.

Lefcheck, J.S., Byrnes, J.E.K., Isbell, F., Gamfeldt, L., Griffin, J.N., Eisenhauer, N., *et al.*

(2015). Biodiversity enhances ecosystem multifunctionality across trophic levels and

habitats. *Nat Commun*, 6.

Li, F., Altermatt, F., Yang, J., An, S., Li, A., & Zhang, X. (2020). Human activities' fingerprint

on multitrophic biodiversity and ecosystem functions across a major river catchment in

China. Global change biology, 26, 6867-6879.

Loreau, M. & Hector, A. (2001). Partitioning selection and complementarity in biodiversity

experiments. *Nature*, 412, 72–76.

Loreau, M., Jarne, P. & Martiny, J.B.H. (2023). Opportunities to advance the synthesis of

ecology and evolution. *Ecol Lett*, 26.

Macneil, C., Dick, J.T.A. & Elwood, R.W. (1997). THE TROPHIC ECOLOGY OF

FRESHWATER *GAMMARUS* SPP. (CRUSTACEA:AMPHIPODA): PROBLEMS AND

PERSPECTIVES CONCERNING THE FUNCTIONAL FEEDING GROUP CONCEPT.

Biol Rev, 72, 349–364.

Moi, D. A., Romero, G. Q., Antiqueira, P. A., Mormul, R. P., Teixeira de Mello, F., & Bonecker,

C. C. (2021). Multitrophic richness enhances ecosystem multifunctionality of tropical

shallow lakes. *Funct Ecol*, 35, 942-954.

Poisot, T., Mouquet, N. & Gravel, D. (2013). Trophic complementarity drives the biodiversity–

ecosystem functioning relationship in food webs. *Ecol Lett*, 16, 853–861.

Prieto, I., Violle, C., Barre, P., Durand, J.-L., Ghesquiere, M. & Litrico, I. (2015).

Complementary effects of species and genetic diversity on productivity and stability of

sown grasslands. *Nat Plants*, 1, 15033.

Prunier, J.G., Chevalier, M., Raffard, A., Loot, G., Poulet, N. & Blanchet, S. (2023). Genetic

erosion reduces biomass temporal stability in wild fish populations. *Nat Commun*, 14,

4362.

Raffard, A., Cucherousset, J., Montoya, J.M., Richard, M., Acoca-Pidolle, S., Poésy, C., *et al.*

(2021). Intraspecific diversity loss in a predator species alters prey community

structure and ecosystem functions. *PLoS Biol*, 19, e3001145.

Raffard, A., Santoul, F., Cucherousset, J. & Blanchet, S. (2019). The community and

ecosystem consequences of intraspecific diversity: a meta-analysis: The ecological

effects of intraspecific diversity. *Biol Rev*, 94, 648–661.

Reusch, T.B.H., Ehlers, A., Hämmerli, A. & Worm, B. (2005). Ecosystem recovery after

climatic extremes enhanced by genotypic diversity. *Proc Natl Acad Sci U.S.A.*, 102,

2826–2831.

Schlötterer, C., Tobler, R., Kofler, R. & Nolte, V. (2014). Sequencing pools of individuals —

mining genome-wide polymorphism data without big funding. *Nat Rev Genet*, 15, 749–

763.

Seibold, S., Cadotte, M. W., MacIvor, J. S., Thorn, S., & Müller, J. (2018). The necessity of

multitrophic approaches in community ecology. *Trends Ecol Evol*, 33, 754-764.

Shipley, B. (2009). Confirmatory path analysis in a generalized multilevel context. *Ecology*,

90, 363–368.

Soliveres, S., van der Plas, F., Manning, P., Prati, D., Gossner, M.M., Renner, S.C., *et al.*

(2016). Biodiversity at multiple trophic levels is needed for ecosystem

multifunctionality. *Nature*, 536, 456–459.

Tang, T., Zhang, N., Bongers, F.J., Staab, M., Schuldt, A., Fornoff, F., *et al.* (2022). Tree

species and genetic diversity increase productivity via functional diversity and trophic

feedbacks. *eLife*, 11, e78703.

Van Der Plas, F. (2019). Biodiversity and ecosystem functioning in naturally assembled

communities. *Biological Reviews*, 94, 1220–1245.

Wan, N.-F., Fu, L., Dainese, M., Hu, Y.-Q., Pødenphant Kiær, L., Isbell, F., *et al.* (2022). Plant

genetic diversity affects multiple trophic levels and trophic interactions. *Nat Commun*,

13, 7312.

Table 1. Characteristics of the two first principal components identified by the Principal
Component Analysis (PCA) ran on the thirteen environmental variables. The part of the
total environmental variance (%) and the contribution of each variable on each component are
shown. The variables that contributed significantly to the axis are highlighted in bold.

	Component 1	Component 2
Part of total Variance (%)	21.66	16.37
River width	0.596	0.320
Connectivity	-0.155	**0.646**
Altitude	**0.648**	-0.385
Distance from outlet	0.528	-0.331
East-west gradient	0.105	**-0.795**
Oxygen concentration	**0.738**	0.343
Oxygen saturation	0.266	-0.509
Water temperature	-0.594	-0.012
Specific conductivity	-0.463	-0.045
pH	0.573	0.398
Concentration in $NO_3^+NO_2$	-0.369	-0.286
Concentration in NH_4+	-0.019	-0.207
Concentration in PO_4^{3-}	-0.279	0.236
Global characteristic	Low altitude, poorly oxygenated site - High altitude, highly oxygenated sites	Poorly connected east site - Highly connected west site

779 **Table 2.** ANOVA table for the linear mixed model testing whether the relationships between
780 biodiversity and ecosystem functions measured in a riverine trophic chain differ between the
781 biodiversity facets (species or genetic diversity) and the types of BEF (*within-* or *between-*
782 *trophic levels*). A Wald chi-quare test is used to test the significance of each fixed effect.

	Degree of freedom	Chisq-value	P-value
(Intercept)	1	0.287	0.595
Biodiversity facet	1	0.232	0.630
Type of BEF	1	5.393	**0.020**
Biodiversity facet*Type of BEF	1	5.567	**0.018**

 Figure captions

Figure 1. Location of sampling sites and illustration of the trophic chain. **(a)** Distribution of the 52 sampling sites (black dots) spanning an East-West gradient at the foothills of the Pyrenees Mountains (France). Each site is denoted by a six-letter code, with the three first letters indicating the river name and the three last letters indicating the closest city or village. **(b)** Our study focused on a tri-trophic food chain commonly found in mountain rivers, consisting of riparian trees, macroinvertebrate shredders and fishes (from bottom to top). Within each trophic level, we measured two facets of biodiversity: genetic diversity in a single target species within each trophic level (specifically *Phoxinus dragarum*, *Gammarus* sp. and *Alnus glutinosa*), and species diversity from communities.

Figure 2. Example of one of the seven causal models used to quantify the relationships between (species and genetic) diversity and ecosystem functions. We focused on seven ecosystem functions associated with genetic and species diversity at three trophic levels (green for primary producer, orange for primary consumer and blue for secondary consumer). Each relationship between biodiversity and ecosystem functions (n = 6 values per function, but for some functions for which irrelevant links were not considered, see the text, n = 32 values in total) was measured at the same trophic level (triangles) or at another trophic level (dots).

Figure 3. Details of the seven causal models linking abiotic parameters, species and genetic diversity and ecosystem functions. Each causal graph (a to g) represents a simplified illustration of the relationships between the two PCA axes synthetizing the environmental parameters of each sampling site (Environmental axis 1 and 2), the species diversity estimated at each trophic level (boxes "Fish", "Shredders" and "Trees"), the genomic diversity estimated from each focal species at each trophic level (boxes "*Phoxinus*", "*Gammarus*" and "*Alnus*"), and each ecosystem function (one model per function). Only the relationships for which the p-value was inferior to 0.20 are indicated for visual simplification. Full arrows indicated positive effects, whereas dotted arrows indicated negative effects. The width of the arrows is proportional to the size of their effects. The percentage of variance explained by environmental and biodiversity effects on ecosystem functions (r^2) is indicated for each function.

Figure 4. General description of individual effect sizes measured between biodiversity estimates and ecosystem functions (BEFs) in a riverine trophic chain. (a) The magnitude and direction of individual effect sizes (*Zr*) of biodiversity is shown for each ecosystem functions as a biplot between *Zr* associated with genetic diversity (y-axis, *genetic* BEFs) measured for one of three target species (*Alnus glutinosa*, *Gammarus sp.* and *Phoxinus dragarum*) and *Zr* associated with species diversity (x-axis, *species* BEFs) measured for one of three trophic levels (trees, invertebrates and fish). For each ecosystem function (but the biomass of trees and of *A. glutinosa*), a total of six *Zr* are depicted in the biplot; four of them are associated with biodiversity measured at another trophic level than the one of the target functions (red symbols, e.g., effect of fish diversity on invertebrate biomass) and two of them are associated with biodiversity measured at the same trophic level than the one of the target functions (blue symbols, e.g., effect of fish diversity on fish biomass). The arrows indicate significant *Zr* (95% confidence intervals excluded 0, see Table S2); vertical arrows are for significant *genetic* BEFs, horizontal arrows are for *species* BEFs. White quadrats stand for situation in which genetic and species BEFs are in the same direction, whereas grey quadrats indicate situation in which genetic and species BEFs are in the opposite direction. Within each quadrat, sub-quadrats indicate the relative magnitude of BEFs, i.e., whether *genetic* BEFs are stronger, weaker or equal in magnitude than *species* BEFs. (b) Density plots displaying the distribution of individual *Zr* for *species*- and *genetic* BEFs (dotted and full lines respectively).

Figure 5. Magnitude and direction of the mean effects sizes estimated from the
relationships between biodiversity and ecosystem functions (BEFs) measured in a
riverine trophic chain. (a) The magnitude and direction of BEFs are expressed as effect
sizes (*Zr*) and are displayed according to the facet used to measured biodiversity (genetic or
species diversity, light grey and white boxplots respectively) and to the type of BEFs (*within-*
trophic level BEFs or *between-trophic level* BEFs, triangles and dots respectively). Red colour
and stars indicate global effect sizes that are significantly different from zero (p-value <0.05).
Large symbols are mean ± 95%IC estimated as marginal effects from the meta-regressions.
Small symbols are raw estimates. (b) Same representation as (a) but with details at each
trophic level (mean ± 95%IC estimated as marginal effects from the meta-regressions, green
for primary producer, orange for primary consumer and blue for secondary consumer). The
trophic level at which BEFs are measured is coherent across all trophic levels.

Figure 1

854   **Figure 2**

859   **Figure 3**

(a) Alnus biomass

(b) Tree biomass

(c) Litter decomposition

(d) Gammarus biomass

 Figure 3 (continued)

(e) Invertebrate biomass

(f) Phoxinus biomass

(g) Fish biomass

Figure 4

Figure 5

SUPPLEMENTARY MATERIALS

Table S1. ANOVA table for the linear mixed model testing whether the relationships between
biodiversity and ecosystem functions measured in a riverine trophic chain differ between the
biodiversity facets (species or genetic diversity), the types of BEF (*within-* or *between-trophic*
levels) and the trophic levels at which BEFs are estimated (primary producers, primary
consumers or secondary consumers). A Wald chi-quare test is used to test the significance of
each fixed effect.

	Degree of freedom	Chisq-value	P-value
(Intercept)	1	1.453	0.228
Biodiversity facet	1	1.293	0.255
Type of BEF	1	7.498	**0.006**
Trophic level	2	1.976	0.372
Biodiversity facet*Type of BEF	1	7.884	**0.005**
Biodiversity facet*Trophic level	2	3.520	0.172
Trophic level*Type of BEF	2	2.901	0.234
Biodiversity facet*Type of BEF*Trophic level	1	2.994	0.224

Table S2 – Estimates of individual effect sizes of BEFs (Zr, n=34) for each ecosystem function, each biodiversity facet (genetic or species diversity) and each type of BEF (within- or between-trophic levels). 95% confidence intervals are provided together with the estimate of each BEF. BEFs are considered as significant when the 95%CI does not overlap 0. P-values estimated from t-test are also provided.

Ecosystem function	Predictor	Biodiversity facet	Type of BEF	Zr	negative 95% CI	positive 95% CI	P values
Phoxinus biomass	**Fish species diversity**	**Species diversity**	**Whithin-trophic levels**	**-0.529**	**-0.803**	**-0.166**	**0.006**
Alnus biomass	**Tree species diversity**	**Species diversity**	**Whithin-trophic levels**	**-0.447**	**-0.695**	**-0.143**	**0.006**
Phoxinus biomass	*Alnus* genetic diversity	**Genetic diversity**	**Between-trophic levels**	**-0.321**	**-0.602**	**-0.020**	**0.045**
Litter decomposition	Tree species diversity	Species diversity	Whithin-trophic levels	-0.293	-0.630	0.060	0.115
Gammarus biomass	Invertebrate species diversity	Species diversity	Whithin-trophic levels	-0.273	-0.661	0.128	0.194
Litter decomposition	Fish species diversity	Species diversity	Between-trophic levels	-0.210	-0.625	0.210	0.337
Tree biomass	Tree species diversity	Species diversity	Whithin-trophic levels	-0.183	-0.485	0.122	0.250
Fish biomass	Invertebrate species diversity	Species diversity	Between-trophic levels	-0.138	-0.428	0.153	0.506
Phoxinus biomass	Invertebrate species diversity	Species diversity	Between-trophic levels	-0.129	-0.471	0.215	0.362
Invertebrate biomass	Invertebrate species diversity	Species diversity	Whithin-trophic levels	-0.051	-0.381	0.279	0.766
Fish biomass	*Gammarus* genetic diversity	Genetic diversity	Between-trophic levels	-0.047	-0.457	0.364	0.825
Gammarus biomass	*Alnus* genetic diversity	Genetic diversity	Between-trophic levels	-0.043	-0.438	0.352	0.833
Litter decomposition	Invertebrate species diversity	Species diversity	Between-trophic levels	-0.020	-0.401	0.361	0.918
Litter decomposition	*Alnus* genetic diversity	Genetic diversity	Whithin-trophic levels	-0.012	-0.394	0.370	0.950
Litter decomposition	*Phoxinus* genetic diversity	Genetic diversity	Between-trophic levels	0.018	-0.379	0.416	0.930
Invertebrate biomass	*Alnus* genetic diversity	Genetic diversity	Between-trophic levels	0.025	-0.306	0.356	0.885
Invertebrate biomass	Tree species diversity	Species diversity	Between-trophic levels	0.063	-0.236	0.362	0.682
Invertebrate biomass	*Phoxinus* genetic diversity	Genetic diversity	Between-trophic levels	0.085	-0.260	0.429	0.634
Fish biomass	Tree species diversity	Species diversity	Between-trophic levels	0.087	-0.276	0.449	0.644
Invertebrate biomass	*Gammarus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.097	-0.242	0.435	0.580
Litter decomposition	*Gammarus* genetic diversity	Genetic diversity	Between-trophic levels	0.110	-0.281	0.500	0.586
Alnus biomass	*Alnus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.128	-0.179	0.433	0.422
Invertebrate biomass	Fish species diversity	Species diversity	Between-trophic levels	0.144	-0.219	0.504	0.444
Fish biomass	*Phoxinus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.159	-0.260	0.575	0.464
Gammarus biomass	Tree species diversity	Species diversity	Between-trophic levels	0.161	-0.198	0.516	0.388
Phoxinus biomass	Tree species diversity	Species diversity	Between-trophic levels	0.165	-0.100	0.426	0.233
Gammarus biomass	Fish species diversity	Species diversity	Between-trophic levels	0.185	-0.249	0.614	0.413
Fish biomass	*Alnus* genetic diversity	Genetic diversity	Between-trophic levels	0.193	-0.211	0.592	0.360
Phoxinus biomass	*Gammarus* genetic diversity	Genetic diversity	Between-trophic levels	0.228	-0.074	0.522	0.151
Phoxinus biomass	*Phoxinus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.244	-0.064	0.542	0.133
Tree biomass	*Alnus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.280	-0.062	0.609	0.121
Gammarus biomass	*Gammarus* genetic diversity	Genetic diversity	Whithin-trophic levels	0.292	-0.120	0.688	0.178
Fish biomass	Fish species diversity	Species diversity	Whithin-trophic levels	0.326	-0.123	0.753	0.169

| Gammarus biomass | Phoxinus genetic diversity | Genetic diversity | Between-trophic levels | 0.446 | 0.007 | 0.830 | 0.047 |

43