Mitogenomics, phylogenetic position, and updated distribution of *Ergasilus kandti*, an ergasilid copepod
parasitizing African cichlid fishes

Dries Jansen[1], Maarten P. M. Vanhove[1], Lawrence Makasa[2], Jiří Vorel[3], Nikol Kmentová[1,4*], Armando J. Cruz-
Laufer[1,5*]

[1]UHasselt – Hasselt University, Centre for Environmental Sciences, Research Group Zoology: Biodiversity &
Toxicology, Hasselt, Belgium; [2]Lake Tanganyika Research Unit, Department of Fisheries, Ministry of Fisheries
and Livestock, Mpulungu, Zambia; [3]CESNET z.s.p.o, Prague, the Czech Republic; [4]Aquatic and Terrestrial
Ecology, Operational Directorate Natural Environment, Royal Belgian Institute of Natural Sciences, Brussels,
Belgium; [5]Systems Ecology and Resource Management Research Unit (SERM), Université Libre de Bruxelles-
ULB, Brussels, Belgium.

*these authors contributed equally

Corresponding author: Armando J. Cruz-Laufer, Email: armando.cruzlaufer@uhasselt.be

Abstract

Ergasilidae are a family of globally distributed copepods parasitizing freshwater fish. Despite their widespread

occurrence, their phylogeographic patterns are poorly understood, specifically in the African Great Lakes. Here,

we aim to provide an update on distribution of *Ergasilus kandti,* a copepod species infecting *Tylochromis*

polylepis, an endemic cichlid fish species in Lake Tanganyika, and the phylogenetic relationship of African

ergasilids. We present the first record of *E. kandti* parasitizing the gills of *T. polylepis* in Lake Tanganyika

proper, identified through light microcopy and, for the first time for any ergasilid, confocal laser scanning

microscopy. We suggest that this technique adds spatial context to characters and are hardly visible while using

light microscopy. Phylogenetic analyses based on ribosomal DNA fragments suggest two monophyletic groups

of African ergasilids. However, the phylogenetic relationships of *Ergasilus* remain unresolved, possibly because

of the insufficient resolution of these widely used phylogenetic markers and low taxonomic coverage. A

comparison of ergasilid mitochondrial genomes highlights traits found in other parasite lineages including

genome shrinkage and low evolutionary rates of the *cox1* gene. This study presents the most extensive

molecular characterization of any ergasilid species to date.

Keywords: Lake Tanganyika; phylogenetics; NGS; confocal microscopy; *Tylochromis polylepis*

Introduction

Copepods are among the most abundant multicellular organisms on Earth (Humes, 1994; Walter & Boxshall, 2023). Parasitic copepods with their free-living larval stages constitute a major proportion of this abundance. Among these, Ergasilidae Burmeister, 1835 is one of most species-rich taxa containing 275 described species infecting mostly teleost fishes (Walter & Boxshall, 2023). Ergasilid copepods pursue a unique life cycle among copepods: they complete their first developmental stages (nauplius to adult) as plankton and after mating, only fertilized females infest a host for the final parasitic phase of the life cycle (Boxshall & Defaye, 2008). As fish is a crucial component of the human food industry, understanding their parasites is key for sustainable food production. Copepods can be serious pests in fish, e.g. *Lernaea cyprinacea* Linnaeus, 1758 and *Ergasilus sieboldi* von Nordmann, 1832, both of which may cause substantial mortality and commercial losses in fish aquaculture (Boxshall & Defaye, 2008). This highlights the importance of studying copepods, and specifically parasitic copepods, yet, despite this relevance, the diversity of parasitic copepods remains understudied (Bernot et al., 2021).

Ancient lakes are biodiversity hotspots, including a high species richness of copepods. Together with Lake Baikal with over 120 species (Boxshall et al., 1993), Lake Tanganyika in East Africa is a prime example with 69 copepod species reported to date (including 34 endemic species) (Boxshall & Strong, 2006). Lake Tanganyika is the longest (660 km) and the second deepest (1436 m) lake in the world (O'Reilly et al., 2003) with an exceptional level of endemism (~95%) of e.g. fishes, snails, and ostracods (Salzburger et al., 2014). These species-rich communities have often resulted from adaptive radiations, a rapid diversification event associated with adaptations to various available ecological niches (Schluter, 2000; Salzburger et al., 2014). The best-studied example of such a community in Lake Tanganyika are cichlid fishes (Cichlidae Bonaparte, 1835), which have become a model system for studying evolution and speciation (Salzburger et al., 2014; Salzburger, 2018; Ronco et al., 2020). Concerning parasitological research in Lake Tanganyika, the majority of studies focus on monogenean flatworm parasites infecting the gills of these cichlids (Vanhove et al., 2011; Rahmouni et al., 2017; Kmentová et al., 2021). Parasitic copepods occupy some of the same host organs as monogenean flatworms (e.g. fish gills) yet receive far less attention. Hitherto, eight species of parasitic copepods belonging to Ergasilidae have been reported in the lake, all of which belong to *Ergasilus* von Nordmann, 1832 (Míč et al., 2023). Notably, copepods dominate the lake's zooplankton community whereas in most other lakes in East Africa cladoceran crustaceans are dominant, highlighting the importance of copepods to Lake Tanganyika's nutrient cycle and ecosystem functioning (Sarvala et al., 1999). Given the combination of free-living and parasitic life stages in ergasilid copepods, knowledge on fish-copepod interactions, including parasitic relationships, are an essential component for lake-wide ecosystem dynamics. However, phylogeographic relationships of parasitic copepods are generally underexplored as only 3% of all copepod species and only seven out of 30 accepted genera of Ergasilidae have been examined in a molecular phylogenetic context (Bernot et al., 2021; Míč et al., 2023). Most of the molecular data on species of *Ergasilus* were published by Song et al. (2008), in a first phylogenetic analysis of the family with most specimens sampled in China. The first molecular data of African ergasilids have been released for five species of the eight so far reported in Lake Tanganyika (Míč et al., 2023) and one from Madagascar (Míč et al., 2024).

The cichlids of Lake Tanganyika all evolved from a common ancestor after the formation of the lake 9–12 Mya,

except for two species: *Tylochromis polylepis* Boulenger, 1900 and *Oreochromis tanganicae* Günther, 1894.

These last two colonized the lake only recently (510,000 years ago for *T. polylepis*), establishing themselves in

an already mature adaptive radiation occupied by species with fine-tuned niche segregation (Koch et al., 2007).

Tylochromis polylepis is less specialized and more of a generalist in ecological terms: The species is a bottom

feeder, usually found above muddy and sandy bottoms close to the shore and river mouths (Koch et al., 2007)

and feeds on gastropods, insect larvae, and plant material (Lamboj, 2004). As the lakes' only tylochromine

cichlid, *T. polylepis* contrasts with the many other cichlid lineages that have undergone adaptive radiation. Its

parasites can, therefore, serve as a model of understanding how the absence of ecological differentiation of host

species affects their parasite fauna. To date, only four parasite species including three monogenean flatworm

species [*Cichlidogyrus mulimbwai* Muterezi Bukinga, Vanhove, Van Steenberge & Pariselle, 2012, *C.*

muzumanii Muterezi Bukinga, Vanhove, Van Steenberge & Pariselle, 2012, and *C. sergemorandi* Rahmouni,

Vanhove & Šimková, 2018] (Muterezi Bukinga et al., 2012; Rahmouni et al., 2018) and a single copepod

species (*Ergasilus kandti* van Douwe, 1912) have been reported to infect *T. polylepis*.

This study investigates the copepod *E. kandti*, a widely distributed species (Fig. 1, Supplementary Table S1)

infecting a specimen of *T. polylepis* in Lake Tanganyika. The species has previously been reported from Lake

Tanganyika parasitizing *Limnotilapia dardennii* Boulenger, 1899, *Lamprologus lemairii* Boulenger, 1899,

Plecodus paradoxus Boulenger, 1898, *Pseudosimochromis curvifrons* Poll, 1942, *Oreochromis* sp. (cited as

Tilapia sp.), and *O. tanganicae* (Capart, 1944; Fryer, 1965). Furthermore, *E. kandti* has also been reported from

T. polylepis in the Malagarasi Delta (Tanzania) that borders the lake (Fryer, 1967). Therefore, our study

provides the first record of *E. kandti* on *T. polylepis* in Lake Tanganyika proper. The species has also been

reported infecting *Lates niloticus* (Linnaeus, 1758) and *Bagrus bajad* (Fabricius, 1775) in Lake Albert (Fryer,

1965; Thurston, 1970), *Pterochromis congicus* (Boulenger, 1897) in Lake Tumba (Fryer, 1964), *Tylochromis*

bangwelensis Regan, 1920 and *T. mylodon* Regan, 1920 in Lake Mweru (Fryer, 1967), *Citharinus citharus*

(Geoffroy Saint-Hilaire, 1808), *Hemisynodontis* sp., *L. niloticus*, and *Synodontis membranaceus* (Geoffroy

Saint-Hilaire, 1809) in the Black and White Volta confluence (Ghana) (Paperna, 1969), and *L. niloticus* in the

River Niger (Capart, 1956; Fryer, 1968) and the Lower Nile (Fryer, 1968). Here, we provide a morphological

comparison with other ergasilid species through light and confocal laser microscopy, the latter of which is

applied to ergasilids for the first time. We conduct phylogenetic analyses of all species of Ergasilidae of which

molecular data are currently available and assess the relationship of African representatives. We also provide the

first ergasilid whole-genome sequencing reads [but see previous studies employing Sanger sequencing to obtain

complete mitochondrial genome sequences, see Feng et al. (2016), He et al. (2023), and Hua et al. (2023)], the

first complete mitogenome sequence and ribosomal operon of any African ergasilid species, and the first

quantitative comparison of mitochondrial genome features of ergasilid copepods.

Materials and methods

Morphological analysis

Copepod specimens (n = 23) were collected from the gills of the single specimen of *T. polylepis* purchased from

a fisherman at the market in Mpulungu, Zambia (-08.767, 031.117) on September 20th, 2019. Specimens were

stored in 70% ethanol at 4°C. The samples were stained with Congo Red (1 mg/ml aqueous solution) for a

minimum of 24 h at room temperature. The dye proved to be stable compared to other fluorescence dyes

(Michels & Büntzow, 2010). Following the staining procedure, the samples were rinsed in deionized water for

30 min. The specimens were mounted on concave microscope slides (to prevent crushing by the cover clip) in

glycerin. Copepod specimens were imaged with a confocal laser microscope (ZEISS LSM900) with an

excitation of 561 nm, an emission at 570–670 nm, pinhole size of 1 AU (airy unit); laser intensity of 0,2%, 400x

magnification; resolution of 512 x 512 px, bidirectional scanning, eight-fold averaging, 30 tiles, and Z-stack

with interval of 1–1.12 µm. Tile images were stitched with the software ZEISS ZEN version 3.1 (Carl Zeiss,

Jena, Germany). Images were further processed in Fiji/ImageJ v1.54d (Schindelin et al., 2012). To adjust the

contrast of dimmed regions of the image, the *StackContrastAdjustment* plugin was used (Čapek et al., 2006),

followed by maximum-intensity projections for two-dimensional representation of the images and denoising.

Specimens were identified to species level using a brightfield light microscope (Leica DM2500) based on the

spine-seta formulae provided by (Capart, 1944) and additional characteristics listed in previous studies

(Oldewage & van As, 1988; Schlebusch, 2014).

DNA extraction and PCRs

DNA extractions were performed in a UV cabinet and all utensils were sterilized and UV–irradiated before

usage. The extractions were completed with a slightly modified version of Kmentová et al. (2021), applying an

extended enzymatic digestion period (55°C, 400 rpm, three hours) to digest the soft tissues of the specimen.

Exoskeletons were removed after this digestion step. The exoskeletons were stained with Congo Red as

mentioned above but mounted in glycerin on standard (non-concave) microscopic slides. For permanent storage,

specimens were either mounted in glycerin in sealed 1.5-ml tubes or in Euparal on microscopic slides. Voucher

material was deposited in the collection of Hasselt University under accession numbers HU XXIII.1.41–HU

XXIII.1.50, HU XXIII.2.01, and HU T-I.1–HU T-I.11.

The PCR reaction targeting a portion of the 28S rDNA region contained forward primer 28SF (5' - ACA ACT

GTG ATG CCC TTA - 3') and reverse primer 28SR (5' - TGG TCC GTG TTT CAA GAC - 3') (Song et al.,

2008). The PCR targeting a portion of the 18S rDNA region contained forward primer 18SF (5' - AAG GTG

TGM CCT ATC AAC - 3') and reverse primer 18SR (5'- TTA CTT CCT CTA AAC GCT - 3') (Song et al.,

2008). An identical master mix was used for both PCRs. It contained 1x PCR buffer, 0.2mM dNTPs, 0.5µM of

each primer, 0.5U of Q5 polymerase, 2µl of DNA and ddH2O was added to bring the total volume to 25µl. The

PCR conditions were set as follows: 5 min at 94°C; 35 cycles of 30 s at 94°C, 1 min at 54°C, 1 min at 72°C; a

final elongation of 10 min at 72°C. Gel electrophoresis was used to check the PCR products for successful

amplification. The PCR products were purified using a GeneJET PCR Purification Kit (Thermo Fisher

Scientific, Waltham, MA, USA) and Sanger sequencing was outsourced to Macrogen Europe (Amsterdam, The

Netherlands). Sequence quality was checked visually through the sequencing chromatograms. The 28S rDNA

and 18S rDNA sequences were uploaded to GenBank and can be found under accession numbers PQ249839 and

PQ249841X–PQ249843, respectively. We also attempted the amplification of a fragment of the cytochrome *c*

oxidase subunit I gene (*cox1*) using a set of general invertebrate primers (Folmer et al., 1994; Nadler et al.,

2006) previously applied successfully to ergasilids (Santacruz et al., 2020; Fikiye et al., 2023). However, none

of these reactions were successful at the amplification stage, possibly due to age of the samples and the low
primer specificity.

Whole-genome sequencing and assembly of ribosomal operon and mitogenome

The extracted genomic DNA of a single copepod individual was subjected to the generation of a genomic read
pool. Library preparation and sequencing on the Illumina NovaSeq6000 platform (short-insert paired-end
sequencing, 2x 150 bp configuration) were outsourced to Novogene Europe (Cambridge, UK). For the assembly
of the ribosomal operon, raw reads were trimmed through Trimmomatic v0.39 (Bolger et al., 2014) using a
sliding window approach (settings: *SLIDINGWINDOW:4:28 HEADCROP:5 MINLEN:100*
ILLUMINACLIP:TruSeq3-PE.fa:2:30:10:2:True). The quality of filtered reads was checked in FastQC v0.11.8
(Andrews, 2018). Reads of the ribosomal operon were baited through complete sequences of the full ribosomal
operon of other copepod species (Ki et al., 2011; Weydmann et al., 2017) using *mirabait* in Mira V5rc1
(Chevreux B & Suhai, 1999). The operon was assembled *de novo* through Spades v3.15.2 (Prjibelski et al.,
2020) with a range of Kmer lengths (21, 33, 55, 77, 99, and 127). The resulting assembly graphs were visualized
in Bandage v0.8.1 (Wick et al., 2015). The resulting contigs were subjected to a BLAST search on NCBI
GenBank to identify those belonging to a copepod. The selected contig sequence was then realigned to the bait
sequences (see above) using MAFFT v.7.409 to locate the boundaries of the rDNA and transcribed spacer
regions.

The mitogenome of *E. kandti* was assembled using raw reads and part of the cytochrome *c* oxidase subunit I
mitochondrial DNA (*cox1*) of *Ergasilus* sp. (GenBank accession number KU557411) as a seed in NOVOPlasty
v4.3.1 with a k-mer length of 21, read length of 151 and insert size of 300. Results from the MITOS web server
(code: invertebrate mitochondrial) (Bernt et al., 2013) were used for mitogenome annotation combined with the
tRNAscan-SE (Lowe & Chan, 2016) and RNAfold (Lorenz et al., 2011) web servers to verify the tRNAcoding
regions. The annotation was verified via a visualization of open reading frames and aligned with available
ergasilid mitogenomes [*Ergasilus tumidus* Markevich, 1940 - OQ596537, NC_073502; *Sinergasilus major*
(Markevich, 1940) – OQ160840, NC_081952; *Sinergasilus polycolpus* (Markevich, 1940) – KR263117,
EU621723, NC_028085; *Sinergasilus undulatus* (Markevich, 1940) – MW080644, NC_054173] in Geneious
Prime 2023.2.1 (Auckland, New Zealand). The annotated mitochondrial genome of *E. kandti* was visualized
using CGView (Grant et al., 2023) including the information on GC content and GC skew+/-. Protein coding
rDNA regions were aligned separately using Clustal Omega v 1.2.2 and subsequently concatenated in Geneious
Prime 2023.2.1. The sliding window analysis and dN/dS ratio across all 13 PCRs was inferred through DnaSP
v5 (Librado & Rozas, 2009) to summarize nucleotide polymorphisms across all ergasilid copepod species of
which the mitochondrial genomes were assembled to date (see above). The assembled nuclear ribosomal and
mitogenome DNA sequences of *E. kandti* are available in the GenBank Nucleotide Database under the
accession numbers PQ249840 and PQ276880. Raw Illumina reads were submitted to SRA (accession number:
SRR30471034) under BioProject accession PRJNA1153390.

Phylogenetic reconstruction

To infer the evolutionary history of *E. kandti*, we assembled multiple sequence alignments of partial 18S and
28S nuclear ribosomal DNA (18S and 28S rDNA) and *cox1* combining sequences generated during this study
with sequences accessed on NCBI GenBank (Table 1). Due to the close relationship of species belonging to
Lernaeidae and Ergasilidae (Song et al., 2008; Kvach et al., 2021; Abdel-Radi et al., 2022), three representatives
of Lernaeidae were chosen as outgroups. Sequences of three species of Mytilicolidae Bocquet & Stock, 1957
and two species of Myicolidae Yamaguti, 1936 were also added to the alignment because previous studies
indicate close ties of these families with Ergasilidae (Khodami et al., 2019; Oliveira et al., 2021). We aligned
18S and 28S rDNA sequences using *L-INS-i* in MAFFT v.7.409 (Katoh & Standley, 2013) and removed poorly
aligned positions and divergent regions using the options for less stringent parameters (options: -
b2=<*minimum*> -b3=10 -b4=5 -b5=h) in Gblocks v.0.91b (Talavera & Castresana, 2007). For *cox1*, sequences
were aligned in MACSE v2.06 (Ranwez et al., 2011, 2018) using the algorithm *trimNonHomologousFragments*,
alignSequences, and *TrimSequenceAlignment* under default parameters and with the Invertebrate Mitochondrial
Code.

We estimated tree topologies through maximum likelihood (ML), Bayesian inference (BI), and maximum
parsimony (MP) methods using IQ-Tree v2.2.26 (Minh et al., 2020), MrBayes v3.2.7 (Ronquist et al., 2012),
and TNT v1.6 (Goloboff et al., 2008; Goloboff & Morales, 2023), respectively. Because of the minimal overlap
in terms of species and specimens between the nuclear and mitochondrial loci, we opted to concatenate only the
18S and 28S rDNA and analyze *cox1* separately. The alignments were partitioned by locus and, for *cox1*, by
codon position. For ML and BI, we selected the substitution models for each partition according to the Bayesian
information criterion (BIC) as suggested by ModelFinder in IQ-Tree (Kalyaanamoorthy et al., 2017) using
partition merging (Chernomor et al., 2016) (Table 2). For ML analyses, we estimated branch support values
using both ultrafast bootstrap approximation (UF-Boot) (Hoang et al., 2018) and Shimodaira–Hasegawa-like
approximate likelihood ratio tests (SH-aLRT) (Guindon et al., 2010) with 10,000 replicates. To minimise false
positives, UF-Boot was estimated by resampling partitions as well as sites within resampled partitions
(Gadagkar et al., 2005; Seo et al., 2005). For BI analyses, model selection in ModelFinder was restricted to
models implemented in MrBayes (Table 2). We used two runs and four chains of Metropolis-coupled Markov
chain Monte Carlo (MCMC) iterations with calculation conducted in parallel to optimize runtime (Altekar et al.,
2004). We ran 20 million generations with a burn-in fraction of 0.25 and sampled the trees every 1000th
generation. We checked convergence criteria by assessing the average standard deviation of split frequencies
(<0.01 in all datasets) and the effective sample size (>200) using Tracer v1.7 (Rambaut et al., 2018).

For analyses in TNT, we used extended implied weighting (command *xpiwe*) (Goloboff, 2014) for different
values of the concavity constant k (20, 21, 23, 26, 30, 35, 41, 48, 56). Extended implied weighting minimizes
the impact of characters with missing data that are weighted artificially high in the original implied weighting
method (Goloboff, 1993). For the rDNA data, we explored two weighting schemes proposed by Mirande
(2019): all characters weighted separately (SEP) and all characters weighted according to the average
homoplasy of the marker (BLK). For *cox1*, we tested an additional scheme: characters weighted according to the
average homoplasy of their codon position (POS). Similar to Mirande (2009), we selected the k-value and
weighting scheme that produced the most stable topology to compute a strict consensus tree. The distortion
coefficient and subtree pruning and regrafting (SPR) distance were used as selection criteria by computing the

similarity of each consensus tree to the others obtained under the different parameters (commands *tcomp* and
sprdiff). MP tree searches involved rounds of tree fusing, sectorial searches, and tree drifting (Goloboff, 1999)
under default settings and each round was stopped following three hits of the same optimum. Gaps were treated
as missing data. As bootstrapping and jackknifing are reported to be distorted by differently weighted characters
(Goloboff, 2003), branch support was estimated through symmetrical resampling (probability of change of 0.33)
and values expressed as differences in frequencies (GC, "Groups present/Contradicted"). We considered a BI
posterior probability ≥ 0.95, UF-Boot ≥ 95 and SH-aLRT statistic ≥ 80 (Hoang et al., 2018), and GC ≥ 0 as
well-supported. Nodes without support above these thresholds in any of the phylogenetic trees were collapsed
into polytomies for visualization. Trees were visualized through R v4.3.2 (R Core Team, 2023) packages
ggplot2 v3.4.4 (Wickham, 2016) and *ggtree* v3.10.0 (Yu et al., 2017, 2018).

Results

Imaging and species identification

The collected specimens were identified as *E. kandti* based on characters observed through light and laser
confocal microscopy (Fig. 2, 3). Congo red showed no signs of fading or bleaching several months after the
initial staining and the multiple imaging sessions of up to 12 hours. All specimens presented characters
associated with the genus *Ergasilus*, i.e. a short, drop-shaped body (Fig. 2a, 3a, 3c) with a sclerotized terminal
segment of the second antennae and a single point (Kuchta et al., 2018). Furthermore, the specimens all
presented a two-segmented exopod on the fourth leg (in contrast with the rest of the legs that all have three
segmented exopods), which is also characteristic of species of *Ergasilus* (see Fryer, 1965). All of the specimens
examined in detail (n = 10) presented a characteristic combination of the following morphological features
specific to *E. kandti*: a fold/groove shaped like an inverted T on the carapace posterior to the eyespot (Fig. 3a),
and a circular cephalic structure anterior to the fold/groove (Fig. 3a), a pair of fifth legs reduced to a single
segment with four setae (Fig. 3b), a pair of modified large grasper-like antennae with a tooth-like projection on
the distal inner side of the third segment (Fig. 2b, 3d), a pair of five-segmented antennules with four setae on the
distal segments (Fig. 2c and 3e), a pair of caudal rami that terminates in two long setae and two short setae (Fig.
2d). Specifically, the tooth-like projection associated with the large grasper-like antennae has been suggested to
be a unique character of the species among species of *Ergasilus* from Lake Tanganyika and the rest of Africa
(Oldewage & van As, 1988; Míč et al., 2023).

Concerning the spine-seta formula, seven out of ten specimens were identical to the original description (Capart,
1944). Occasionally, deviations in the numbers of setae were observed, i.e. on the second segment of the
endopodite of the third leg for a single specimen (see Supplementary Table S2) compared to two setae
mentioned in the original description. However, these likely broke off during the mounting process as all species
of *Ergasilus* have one or two setae on this leg segment. Apart from the abovementioned characters, African
congeneric species were ruled out through the comparison of their respective spine-seta formulas. *Ergasilus*
cunningtoni Capart, 1944 and *E. macrodactylus* (Sars, 1909) have four (Sars, 1909; Capart, 1944) and *E.*
nodosus Wilson, 1924 five setae on the third segment of the exopodite of the third leg (Wilson, 1924) compared
to the six setae present in *E. kandti* (Fig. 2b) (van Douwe, 1912). Furthermore, *E. cunningtoni, E. lamellifer*
Fryer, 1961*, E. latus* Fryer, 1960*, E. macrodactylus, E. megacheir* Sars, 1909*, E. mirabilis* Oldewage & van As,

1987, *E. caparti* Míč, Řehulková & Seifertová, 2023, *E. parvus* Míč, Řehulková & Seifertová, 2023, *E.*

parasarsi Míč, Řehulková & Seifertová, 2023, and *E. sarsi* Capart, 1944 have a six-segmented antennule

compared to a five-segmented one present in *E. kandti* (Fig. 2a, 3d). *Ergasilus kandti* differs from *E. flaccidus*

Fryer, 1965 and *E. nodosus* by having a single segmented fifth leg, and from *E. inflatipes* Cressey, 1970 by not

having plumose setae on the fifth leg.

DNA sequence assemblies

Through PCRs and Sanger sequencing, we obtained overall three rDNA sequences of three specimens of *E.*

kandti (Table 1), the first from this species. Genomic DNA sequencing yielded 38,115,846 indexed paired-end

reads. The complete ribosomal operon is 3,616 bp long (56.6% GC). Overall, for 28S rDNA (629–3838 bp) and

18S rDNA (1047–1849 bp), the sequences showed 99.8–99.9% similarity (uncorrected distance) for both

markers. The complete mitochondrial genome of *E. kandti* is 14,639 bp long (35.8% GC) and includes 13

intron-free protein coding genes (PCGs), 22 tRNA genes, and two genes coding for the large and small subunits

of the mitochondrial rRNA (Fig. 4, Table 3). The usage of T- and TA-abbreviated stop codon in the PCGs are

three and two out of 13 PCGs, respectively. A comparison of available mitogenomes with other representatives

of Ergasilidae highlighted differences concerning the gene order. *Ergasilus kandti* differs in the mutual position

of *trnS1* and *trnS2* with *E. tumidus* (GenBank accession number NC_073502). Furthermore, we found that the

mutual position of multiple tRNAs (*trnR, trnI, trnA* and *trnH*) differed between *S. polycolpus* and the four other

ergasilid species, of which the mitochondrial genome sequences are available (*trnC, trnI, trnA, trnH, trnR*, and

trnT in *S polycolpus*, and *trnC, trnR, trnI, trnA trnH*, and *trnT* in the rest). The existence of a control region or

AT-loop region between *trnT* and *rrnL* was suggested by Hua et al. (2021) and Feng et al. (2016) for *S. majo*r

(597 bp) and *S. polycolpus* (156 bp), respectively. In *E. kandti*, a non-coding region of 443 bp between *trnT* and

rrnL (see Fig. 4) resembles this putative control region. The AT content in this non-coding region of *E. kandti* is

59.4% compared to 52.6% in *S. major* and 81.4% in *S. polycolpus*. We also observed that the length of rrnL in

S. polycolpus (Feng et al., 2016) appears to be shorter compared to the other species including *E. kandti* (668 bp

vs. 878 bp).

The sliding window analysis indicated an overall higher nucleotide diversity in genes coding for dehydrogenase

subunits compared to other protein coding regions of cytochromes and ATP synthases (see Fig. 5). The lowest

values were found for the *cox1* and *cox2* genes, respectively. The dN/dS ratios follow a similar trend with the

highest values reported for nad6 and the lowest for *cox1* (see Fig. 6).

Phylogenetic relationships

We applied maximum likelihood (ML), Bayesian inference (BI), and maximum parsimony (MP) analyses to a

concatenated alignment of partial 18S rDNA and 28S rDNA sequences of ergasilids (1500 bp, 212 informative

sites). These different analyses resulted in phylogenetic trees with congruent topologies. Therefore, only the ML

tree is shown in Fig. 7. The resulting tree topology supports four clades within Ergasilidae including the three

clades (Clade I–III) reported in previous studies (see Song et al., 2008). Furthermore, a fourth clade (Clade IV)

includes all sequenced African species of *Ergasilus*, namely *E. caparti, E. parvus, E. parasarsi, E.*

macrodactylus, *E. mirabilis*, *E. megacheir*, and *E. kandti* as well as the Malagasy *Dermoergasilus*

madagascarensis Míč, Řehulková, Šimková, Razanabolana & Seifertová, 2024 and the non-African species *E.*
sieboldi and *E. yaluzangbus* Kuang & Qian, 1985. Clade IV is subdivided into two well-supported groups: the
African species of *Ergasilus* and the Asian species *E. yaluzangbus* on one hand and *D. madagascarensis* and *E.*
sieboldi on the other. Our results also indicate that *Ergasilus* is rendered paraphyletic by *Dermoergasilus* Ho &
Do, 1982*, Neoergasilus* Yin, 1956, *Paraergasilus* Markevich, 1937, and *Sinergasilus* Yin, 1949. Phylogenetic
analyses (ML, BI, MP) of *cox1* deviated from the rDNA-based topology (Supplementary Fig. S3), e.g. the
myicolid and mytilicolid copepods appear to be nested inside Ergasilidae and the monophyly of *Sinergasilus* has
low support. Generally, support values were lower throughout the *cox1* phylogeny, leading to many deeper
nodes being collapsed into polytomies.

Discussion

Morphology and host repertoire of *E. kandti*

We investigated parasitic copepods infecting a specimen of *Tylochromis polylepis* from Lake Tanganyika in
East Africa, one of the world's biodiversity hotspots, and with a so far understudied copepod diversity. A
morphological examination based on light and confocal laser microscopy revealed that *T. polylepis* was infected
by *Ergasilus kandti* and morphological features of the parasites largely matched those listed in previous
characterizations (van Douwe, 1912; Capart, 1944). However, we failed to observe the variation of short setae
associated with the caudal rami reported by Capart (1944). Our samples only presented two of these setae as
reported by the original description (van Douwe, 1912).

Previous studies on ergasilid copepods have mostly applied light microscopy to highlight systematically
informative characters. In recent decades, other imaging techniques have found their way into ergasilid research.
For instance, scanning electron microscopy was employed to visualize the pathological alterations in host gill
filaments and the feeding apparatus (Kilian & Avenant☐Oldewage, 2013; Dos Santos et al., 2021; Avenant-
Oldewage et al., 2024) as well as systematically informative characters on the legs (Murray et al., 2016).
Confocal laser microscopy has been applied successfully to caligids, tisbids, calanids, centropagids, cyclopids,
and peltidiids (Buttino et al., 2003; Michels, 2007; Zirbel et al., 2007; Fitzer et al., 2012; Kaji et al., 2012), but
this technique has remained absent from the ergasilid literature. Here, we applied confocal laser microscopy to
ergasilid copepods for the first time using Congo Red as a fluorescent marker for the exoskeleton. Confocal
laser microscopy revealed no additional features compared to traditional light microscopy and was more time-
intensive. However, it provides a three-dimensional (i.e. more natural) view of the organism and requires no
clearing agent. Therefore, we suggest that this technique adds spatial context to characters that require flattening
to be observed in a single plane through light microscopy.

Lake Tanganyika contains several examples of adaptive radiations, i.e. explosive speciation, including snails,
fishes (Salzburger et al., 2014), and flatworms (Vanhove et al., 2015; Brand, 2023). The best-known examples
are cichlid fishes with around 250 species (Ronco et al., 2020). However, *T. polylepis*, the host of *E. kandti*, is
only a recent colonizer of Lake Tanganyika (Koch et al., 2007). The parasites of *T. polylepis* are, hence, of
particular interest in terms of studying the effects of adaptive radiations on symbiotic relationships. Prior studies
suggest that ergasilid copepods often infect a broad array of host species, e.g., *E. sieboldi* is reported from 15

fish families (Walter & Boxshall, 2023). *Ergasilus kandti* appears to follow this trend. The species has
previously been reported from Lake Tanganyika parasitizing both the cichlid radiation flock (Fryer, 1965) as
well as a late arrival [see also *E. kandti* infecting *Oreochromis tanganicae* (Fryer, 1965)]. In contrast, no
monogenean flatworms shared between *T. polylepis* and other cichlid species in the lake were reported so far
(Muterezi Bukinga et al., 2012; Rahmouni et al., 2018; Cruz☐Laufer et al., 2022). The distribution range of *E.*
kandti is not limited to Lake Tanganyika (Fig. 1). This wide geographical range and host repertoire (Fig. 1,
Supplementary Table S1) aligns with the low environmental specificity reported in other ergasilid copepods
(Mathews et al., 2017; Fikiye et al., 2023).

Ergasilidae: clades and nomenclature

Due to the persistent lack of DNA sequence data, inferring any biogeographical patterns in the evolution of
ergasilid copepods has long been a challenge. As noted by Míč et al. (2023) and mentioned above, these data
have a pronounced geographical bias as all molecular data until 2019 were from specimens collected in Asia.
The phylogenetic relationships of ergasilid copepods remain generally poorly understood as only 11% (31, see
Table 1) of described ergasilid species [275, see Walter & Boxshall (2023)] have been sequenced to date. Here,
the phylogenetic analyses suggest that ergasilid species reported from Lake Tanganyika including *E. kandti*
form a monophyletic group (part of Clade IV, Fig. 7). No DNA sequences of other African species have been
published, bar *E. mirabilis* (Fikiye et al., 2023) as well as the recently described *D. madagascarensis*. The latter
species is reported from Madagascar (Míč et al., 2024) and is more closely related to *E. sieboldi*, a species from
Eurasia, suggesting that at least two lineages of ergasilids are present in Africa. Nonetheless, low taxonomic
coverage alongside the high evolutionary rates of parasitic organisms (Bromham et al., 2013; Vanhove et al.,
2013; Gao et al., 2022) might also explain the difficulty of resolving the deeper nodes of the ergasilid
phylogeny, especially in the case of the *cox1* phylogeny, which involves sequences of fewer species than the
ribosomal phylogeny.

From a systematic point of view, morphological characters widely used to distinguish lineages of Ergasilidae
appear to result in mixed success. Several genera, each with their own set of diagnostic characters, including
Neoergasilus, *Paraergasilus*, and *Sinergasilus*, are repeatedly confirmed as monophyletic groups (Song et al.,
2008; Santacruz et al., 2020; Kvach et al., 2021; Míč et al., 2023, 2024). At the same time, these genera render
Ergasilus a paraphyletic taxon as suggested in previous studies (Song et al., 2008; Santacruz et al., 2020; Kvach
et al., 2021; Míč et al., 2023, 2024). Some species including *E. anchoratus* Markevich, 1946 and *E. wilsoni*
Markevich, 1933 appear as sister taxa to *Sinergasilus* and *Paraergasilus* (Fig. 7), yet they were not placed in
these genera by the authors erecting these genera, respectively. The nomenclatural history of the ergasilids
further highlights the difficulty of creating generic diagnoses that allow to effectively distinguish groups.
Pseudergasilus Yamaguti, 1936, *Trigasilus* Fryer, 1956, and *Ergasiloides* Sars, 1909 are three genera that were
synonymized, *Pseudergasilus* and *Ergasiloides* with *Ergasilus*, and *Trigasilus* with *Paraergasilus*, after authors
of subsequent publications found the diagnoses to be inadequate. *Pseudergasilus* was differentiated by having
"thoracic segments that are distinctly segmented from each other" (Yamaguti, 1936), but Kim & Choi (2003)
suggested that this difference was merely an artifact of the growth state of the copepods. *Trigasilus* was created
by Fryer (1956). The author later admitted to not being aware of the original diagnosis of *Paraergasilus*

(Markewitsch, 1937) but also remarked on the difference in the attachment sites (gill chamber *vs.* gills) and it

should be noted that *Paraergasilus minutus* (Fryer, 1956) (synonym: *Trigasilus minutus*) is the only African

freshwater representative of this genus (but see *Paraergasilus lagoonaris* Paperna, 1969, which is reported from

brackish water fishes in Ghana).

Remarkable in the context of the present study is *Ergasiloides*. This genus was created to encompass three

species from Lake Tanganyika (*E. megacheir*, *E. macrodactylus*, and *E. brevimanus*), the two former of which

form part of the Lake Tanganyika group in the present study (highlighted in bold as part of Clade IV, Fig. 4).

The genus was synonymized with *Ergasilus* because Sars (1909) based his diagnosis on immature specimens

according to Fryer (1965). However, Fryer (1965) also noted that species considered as members of

Ergasiloides by Sars (1909) differed from other species of *Ergasilus* because adult females have two instead of

three abdominal somites (or 'urosomites') but opted to emend the generic diagnosis of *Ergasilus* to incorporate

both this feature as well as the five-segmented antennule of *E. kandti* as he considered these characters mere

'anomalies' and variations to be expected for this group of parasites. The abdominal somites are indeed not

fused in several species from Lake Tanganyika, including *E. kandti* (van Douwe, 1912), *E. parasarsi*, and *E.*

parvus (Míč et al., 2023), and a recent re-description also refutes this feature in *E. macrodactylus* (Míč et al.,

2023). Nonetheless, our phylogenetic analysis (Fig. 7) suggests that if the genus *Ergasiloides* were to be re-

established, it could be applied as the name of the Lake Tanganyika group (part of Clade IV) as it was

previously assigned to several of these species. However, this step would require the provision of a consistent

generic diagnosis through a systematic revision.

Mitochondrial genomes of ergasilids: tRNA rearrangement and nucleotide polymorphism

In general, crustaceans have high rates of gene order rearrangement in mitochondrial genomes (Sterling-

Montealegre & Prada, 2024). Given the general scarcity of ergasilid DNA sequences, it comes as no surprise

that full mitochondrial genomes are available for even fewer species. Nonetheless, a comparison of ergasilids

and other copepods can provide some insights into the interspecific variation of these sequences and the gene

order. While the order of mitochondrial rDNA and rRNA regions is stable across ergasilid copepods, we

detected several instances of rearrangement between different tRNAs in the mitochondrial genome.

Furthermore, we report the presence of the TA-abbreviated stop codon (15.38%) in the protein coding genes

being higher than the average usage (2.01%) reported mitochondrial regions at the level of Copepoda (He et al.,

2023). In comparison to free-living copepods (Ki et al., 2009; Minxiao et al., 2011), ergasilid copepods show the

complete absence of repetitive regions and a relatively short non-coding region in their mitochondria (in those

assembled so far).

We also observed similar patterns found in other parasites when comparing nucleotide polymorphism and

dN/dS ratios. Nucleotide polymorphism across the mitochondrial rrDNA regions of ergasilid copepods was

disproportionately higher in regions coding for dehydrogenase subunits compared to cytochrome and ATP

coding regions (see Figs. 5 and 6). Our results on the dN/dS ratio further suggest that the *cox1* gene is under the

strongest selection pressure. This pattern aligns with the results of previous studies highlighting low

evolutionary rates of the *cox1* compared to other mitochondrial protein coding regions, e.g., in calanoids (Zhang

et al., 2021), but also other copepods (He et al., 2023; Huang et al 2024). Similar patterns are also present in

monogenean flatworms infecting cichlid (Kmentová et al., 2021) and clupeid fishes (Kmentová et al., 2023) in
Lake Tanganyika. These parasites share not only the gill habitat and host lineages (i.e. cichlids) with ergasilid
copepods, but also both have a single-host life cycle, However, the limited number of ergasilid copepods of
which the mitochondrial genomes has been assembled today (n = 5, including the present study) in comparison
to the estimated worldwide diversity of 275 species prevents any general conclusion and more studies are
needed to unravel (mito-)genomic patterns underlying diversification processes of ergasilid copepods.

Conclusions

The East African lakes and, specifically Lake Tanganyika, are a biodiversity hotspot for many different
vertebrate and invertebrate taxa. Despite this knowledge, copepods—specifically parasitic species—are one of
the most overlooked groups in these lakes. A major knowledge gap also remains associated with the lack of
molecular data across the geographical ranges of ergasilid species. We suggest that to understand the drivers of
evolution of ergasilid copepods, these parasites need to be sampled across their geographical range. Specifically,
E. kandti should be sampled in different water bodies and, specifically, hosts inside and outside of Lake
Tanganyika. Furthermore, ergasilid copepods were suggested as bioindicators being sensitive to eutrophication
and other processes derived from ecosystem degradation (Sasal et al., 2007). Therefore, studying temporal
dynamics of ergasilid copepods, species combining both free-living and parasitic life stages, could be
informative for ecosystem health assessment.

Our study provides an insight of how upscaling both morphological characterization and molecular analyses
using technological solutions applied for other copepods and metazoan parasites can expand our knowledge on
this key taxon. While the use of next-generation sequencing and confocal imaging is new to ergasilids and other
copepods in East Africa, protocols can be transferred without facing major challenges. We encourage
researchers to incorporate these tools into future studies on ergasilids worldwide.

Data availability. Parasite voucher material was deposited in the collection of Hasselt University under
accession numbers HU XXIII.1.41–HU XXIII.1.50, HU XXIII.2.01, and HU T-I.1–HU T-I.11. The ribosomal
DNA sequences were uploaded to GenBank and can be found under accession numbers PQ249839–PQ249843.
The mitogenome DNA sequence is available in the GenBank Nucleotide Database under the accession number
PQ276880. Raw Illumina reads were submitted to SRA (accession number: SRR30471034). All sequences are
linked under BioProject accession PRJNA1153390. Image files and phylogenetic tree files can be accessed at
MorphoBank (http://www.morphobank.org/, Project 5467).

Acknowledgements. We would like to dedicate this manuscript to Geoffrey Fryer (1927–2024), whose works
proved essential for the present study. We would like to thank Stephan Koblmüller, Holger Zimmermann,
Simona Georgieva, Gyrhaiss Kapepula Kasembele, Cyprian Katongo and Taylor Banda, for their help in
organising and conducting field work. We further thank Natascha Steffanie and Martijn Heleven for assistance
concerning laboratory work. We would also like to thank the anonymous reviewers for the valuable feedback on
this manuscript.

Author's contribution. DJ, NK, and AJCL conceptualized the study. NK led the sampling campaign with the
assistance of JV and LM. LM collected the fish host specimen in the field. AJCL collected parasite specimens
off the host gills. DJ prepared samples and conducted imaging and molecular analyses under supervision of NK
and AJCL. NK and AJCL analyzed next-generation sequencing data. DJ and AJCL performed phylogenetic
analyses. DJ, NK, and AJCL wrote the article with input from MPMV, JV and LM. All co-authors approved the
final version of the manuscript.

Financial support. The study was supported by the Czech Science Foundation (GACR) standard project GA19-
13573S, research grant 1513419N of the Research Foundation—Flanders (FWO-Vlaanderen). AJCL
(BOF19OWB02), NK (BOF21PD01) and MPMV (BOF20TT06) received support from the Special Research
Fund of Hasselt University. Part of the research leading to results presented in this publication was carried out
with infrastructure funded by the European Marine Biological Research Centre (EMBRC) Belgium, FWO-
Vlaanderen project GOH3817N. NK is currently funded by the Belgian Science Policy Office (BELSPO)
(AfroWetMap project). AJCL is currently funded by the Research Foundation—Flanders (FWO-Vlaanderen)
(12APB24N). Parts of the resources and services used in this work were provided by the VSC (Flemish
Supercomputer Center) funded by the Research Foundation – Flanders (FWO) and the Flemish Government.
Computational resources were provided by the e-INFRA CZ project (ID:90254), supported by the Ministry of
Education, Youth and Sports of the Czech Republic and the ELIXIR-CZ project (ID:90255), part of the
international ELIXIR infrastructure.

Competing interests. The authors declare that there are no conflicts of interest.

Ethical standards. Fieldwork was carried out with the approval of the competent local authorities under the
permission of the Fisheries Department of Zambia and under a study permit issued by the government of
Zambia (SP 008732).

References

Abdel-Radi, S., M. M. Rashad, G. E. Ali, A. E. Eissa, M. Abdelsalam, & M. Abou-Okada, 2022. Molecular
characterization and phylogenetic analysis of parasitic Copepoda; *Ergasilus sieboldi* isolated from cultured
gilthead sea bream (*Sparus aurata*) in Egypt, associated with analysis of oxidative stress biomarkers. Journal of
Parasitic Diseases 46: 1080–1089.

Altekar, G., S. Dwarkadas, J. P. Huelsenbeck, & F. Ronquist, 2004. Parallel Metropolis coupled Markov chain
Monte Carlo for Bayesian phylogenetic inference. Bioinformatics 20: 407–415.

Andrews, S., 2018. FastQC. Accessed May 7, 2023. http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Avenant-Oldewage, A., K. Nagasawa, Q. M. Dos Santos, & W. H. Oldewage, 2024. Pathology caused by
introduced *Neoergasilus japonicus* (Copepoda: Ergasilidae) to the skin of indigenous *Tilapia sparrmanii* in
South Africa and scanning electron microscopy study of wound-inflicting structures. Journal of Fish Diseases
47: e13867.

Baek, S. Y., K. H. Jang, E. H. Choi, S. H. Ryu, S. K. Kim, J. H. Lee, Y. J. Lim, J. Lee, J. Jun, M. Kwak, Y. S.
Lee, J. S. Hwang, B. A. V. Maran, C. Y. Chang, I. H. Kim, & U. W. Hwang, 2016. DNA barcoding of metazoan
zooplankton copepods from South Korea. PLoS ONE 11: e015307.

Bernot, J. P., G. A. Boxshall, & K. A. Crandall, 2021. A synthesis tree of the Copepoda: integrating
phylogenetic and taxonomic data reveals multiple origins of parasitism. PeerJ 9: e12034.

Bernt, M., A. Donath, F. Jühling, F. Externbrink, C. Florentz, G. Fritzsch, J. Pütz, M. Middendorf, & P. F.
Stadler, 2013. MITOS: improved *de novo* metazoan mitochondrial genome annotation. Molecular Phylogenetics
and Evolution 69: 313–319.

Bolger, A. M., M. Lohse, & B. Usadel, 2014. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics 30: 2114–2120.

Boxshall, G. A., & D. Defaye, 2008. Global diversity of copepods (Crustacea: Copepoda) in freshwater.
Hydrobiologia 595: 195–207.

Boxshall, G. A., T. D. Evstigneeva, & P. F. Clark, 1993. A new interstitial cyclopoid copepod from a sandy
beach on the western shore of Lake Baikal, Siberia. Hydrobiologia 268: 99–107.

Boxshall, G. A., & E. E. Strong, 2006. An extraordinary shift in life habit within a genus of cyclopid copepods
in Lake Tanganyika. Zoological Journal of the Linnean Society 146: 275–285.

Brand, J. N., 2023. Support for a radiation of free-living flatworms in the African Great Lakes region and the
description of five new *Macrostomum* species. Frontiers in Zoology 20: 31.

Bromham, L., P. F. Cowman, & R. Lanfear, 2013. Parasitic plants have increased rates of molecular evolution
across all three genomes. BMC Evolutionary Biology 13: 126.

Buttino, I., A. Ianora, Y. Carotenuto, V. Zupo, & A. Miralto, 2003. Use of the confocal laser scanning
microscope in studies on the developmental biology of marine crustaceans. Microscopy Research and
Technique 60: 458–464.

Capart, A., 1944. Notes sur les copépodes parasites. III. Copépodes parasites des poissons déau douce du Congo
Belge. Bulletin du Musée royal d'Histoire naturelle de Belgique 20: 1–24.

Capart, A., 1956. Quelques copépodes parasites de poissons du Niger (Gourao) récoltés par Th. Monod. Bulletin
de l'Institut Fondamental d'Afrique Noire 18A: 485–494.

Čapek, M., J. Janáček, & L. Kubínová, 2006. Methods for compensation of the light attenuation with depth of
images captured by a confocal microscope. Microscopy Research and Technique 69: 624–635.

Chernomor, O., A. von Haeseler, & B. Q. Minh, 2016. Terrace aware data structure for phylogenomic inference
from supermatrices. Systematic Biology 65: 997–1008.

Chevreux B, & S. Suhai, 1999. Genome sequence assembly using trace signals and additional sequence
information In Wingender, E. (ed), Proceedings of the German Conference on Bioinformatics, Hannover,
Germany, 4–6 October 1999. Gesellschaft für Biotechnologische Forschung, Department of Bioinformatics,
Braunschweig, Germany: 45–56.

Cruz☐Laufer, A. J., T. Artois, S. Koblmüller, A. Pariselle, K. Smeets, M. Van Steenberge, & M. P. M.
Vanhove, 2022. Explosive networking: the role of adaptive host radiations and ecological opportunity in a
species☐rich host–parasite assembly. Ecology Letters 25: 1795–1812.

Dos Santos, Q. M., A. Avenant-Oldewage, W. Piasecki, K. Molnár, B. Sellyei, & C. Székely, 2021. An alien
parasite affects local fauna—confirmation of *Sinergasilus major* (Copepoda: Ergasilidae) switching hosts and
infecting native *Silurus glanis* (Actinopterygii: Siluridae) in Hungary. International Journal for Parasitology:
Parasites and Wildlife 15: 127–131.

Elsner, N. O., S. Jacobsen, D. W. Thieltges, & K. Reise, 2011. Alien parasitic copepods in mussels and oysters
of the Wadden Sea. Helgoland Marine Research 65: 299–307.

Feng, H. L., L. X. Wang, J. Huang, J. Jiang, D. Tang, R. Fang, & Y. B. Su, 2016. Complete mitochondrial
genome of *Sinergasilus polycolpus* (Copepoda: Poecilostomatoida). Mitochondrial DNA 27: 2960–2962.

Fikiye, P. P., N. J. Smit, L. L. Van As, M. Truter, & K. A. Hadfield, 2023. Integrative morphological and
genetic characterisation of the fish parasitic copepod *Ergasilus mirabilis* Oldewage & van As, 1987: insights
into host specificity and distribution in southern Africa. Diversity 15: 965.

Fitzer, S. C., J. D. D. Bishop, G. S. Caldwell, A. S. Clare, R. C. Upstill-Goddard, & M. G. Bentley, 2012.
Visualisation of the copepod female reproductive system using confocal laser scanning microscopy and two-
photon microscopy. Journal of Crustacean Biology 32: 685–692.

Folmer, O., M. Black, W. Hoeh, R. Lutz, & R. Vrijenhoek, 1994. DNA primers for amplification of
mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology
and Biotechnology 3: 294–299.

Fryer, G., 1956. A report on the parasitic Copepoda and Branchiura of the fishes of Lake Nyasa. Proceedings of
the Zoological Society of London 127: 293–344.

Fryer, G., 1964. Further studies on the parasitic Crustacea of African freshwater fishes. Proceedings of the
Zoological Society of London 143: 79–102.

Fryer, G., 1965. Crustacean parasites of African freshwater fishes, mostly collected during the expeditions to
Lake Tanganyika and to Lakes Kivu, Edward and Albert by the Institut Royal des Sciences Naturelles de
Belgique. Bulletin of the Royal Belgian Institute of Natural Sciences 41: 1–22.

Fryer, G., 1967. Parasitic copepods from African cichlid fishes in the Musée Royal de l'Afrique Centrale. Revue
de Zoologie et de Botanique africaine 76: 357–363.

Fryer, G., 1968. The parasitic Crustacea of African freshwater fishes; their biology and distribution. Journal of
Zoology 156: 45–95.

Gadagkar, S. R., M. S. Rosenberg, & S. Kumar, 2005. Inferring species phylogenies from multiple genes:
Concatenated sequence tree versus consensus gene tree. Journal of Experimental Zoology Part B: Molecular and
Developmental Evolution 304B: 64–74.

Gao, J. W., X. P. Yuan, H. Wu, C. Y. Xiang, M. Xie, R. Song, Z. Y. Chen, Y. A. Wu, & D. S. Ou, 2022.
Mitochondrial phylogenomics of Acanthocephala: nucleotide alignments produce long-branch attraction
artefacts. Parasites and Vectors 15: 376.

Goloboff, P. A., 1993. Estimating character weights during tree search. Cladistics 9: 83–91.

Goloboff, P. A., 1999. Analyzing large data sets in reasonable times: solutions for composite optima. Cladistics
15: 415–428.

Goloboff, P. A., 2003. Improvements to resampling measures of group support. Cladistics 19: 324–332.

Goloboff, P. A., 2014. Extended implied weighting. Cladistics 30: 260–272.

Goloboff, P. A., J. S. Farris, & K. C. Nixon, 2008. TNT, a free program for phylogenetic analysis. Cladistics 24:
774–786.

Goloboff, P. A., & M. E. Morales, 2023. TNT version 1.6, with a graphical interface for MacOS and Linux,
including new routines in parallel. Cladistics 39: 144–153.

Grant, J. R., E. Enns, E. Marinier, A. Mandal, E. K. Herman, C. Y. Chen, M. Graham, G. Van Domselaar, & P.
Stothard, 2023. Proksee: in-depth characterization and visualization of bacterial genomes. Nucleic Acids
Research 51: W484–W492.

Guindon, S., J.-F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk, & O. Gascuel, 2010. New algorithms and
methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Systematic
Biology 59: 307–321.

He, J., Z. Zhou, Y. Huang, J. Feng, W. Li, G. Wang, & C. Hua, 2023. Evolutionary rates, divergence rates, and
performance of individual mitochondrial genes based on phylogenetic analysis of Copepoda. Genes 14: 1496.

Hoang, D. T., O. Chernomor, A. von Haeseler, B. Q. Minh, & S. Le Vinh, 2018. UFBoot2: improving the
ultrafast bootstrap approximation. Molecular Biology and Evolution 35: 518–522.

Hua, C. jie, M. yan Su, Z. wei Sun, Y. hong Lu, & J. mei Feng, 2021. Complete mitochondrial genome of the
copepod *Sinergasilus undulates* [sic] (Copepoda: Poecilostomatoida). Mitochondrial DNA Part B: Resources 6:
1226–1228.

Hua, C.-J., Y. Huang, H.-Q. Li, J.-M. Feng, J.-L. Qiao, S. Hu, & W. Liu, 2023. Characterization of the complete
mitochondrial genome of *Ergasilus anchoratus* and phylogeny of Copepoda [preprint]. Research Square DOI:
10.21203/rs.3.rs-3390559/v1.

Humes, A. G., 1994. How many copepods?. Hydrobiologia 292–293: 1–7.

Huys, R., J. Llewellyn-Hughes, P. D. Olson, & K. Nagasawa, 2006. Small subunit rDNA and Bayesian
inference reveal *Pectenophilus ornatus* (Copepoda *incertae sedis*) as highly transformed Mytilicolidae, and
support assignment of Chondracanthidae and Xarifiidae to Lichomolgoidea (Cyclopoida). Biological Journal of
the Linnean Society 87: 403–425.

Kaji, T., B. A. Venmathi Maran, Y. Kondoh, S. Ohtsuka, G. A. Boxshall, & A. Tsukagoshi, 2012. The lunule of
caligid copepods: an evolutionarily novel structure. Evolution and Development 14: 465–475.

Kalyaanamoorthy, S., B. Q. Minh, T. K. F. Wong, A. von Haeseler, & L. S. Jermiin, 2017. ModelFinder: fast
model selection for accurate phylogenetic estimates. Nature Methods 14: 587–589.

Katoh, K., & D. M. Standley, 2013. MAFFT multiple sequence alignment software version 7: improvements in
performance and usability. Molecular Biology and Evolution 30: 772–780.

Khodami, S., N. F. Mercado-Salas, D. Tang, & P. Martinez Arbizu, 2019. Molecular evidence for the retention
of the Thaumatopsyllidae in the order Cyclopoida (Copepoda) and establishment of four suborders and two
families within the Cyclopoida. Molecular Phylogenetics and Evolution 138: 43–52.

Ki, J. S., H. G. Park, & J. S. Lee, 2009. The complete mitochondrial genome of the cyclopoid copepod
Paracyclopina nana: a highly divergent genome with novel gene order and atypical gene numbers. Gene 435:
13–22.

Ki, J. S., H. G. Park, & J. S. Lee, 2011. Extensive analysis of nuclear cistron rDNA sequence of *Paracyclopina*
nana (Cyclopoida: Cyclopettidae). Hydrobiologia 666: 3–9.

Kilian, E., & A. Avenant☐Oldewage, 2013. Observations on the pathology of *Ergasilus* (Crustacea: Copepoda)
from Lake Tanganyika on the gills of *Lamprichtys tanganicanus*. Suid-Afrikaanse Tydskrif vir
Natuurwetenskap en Tegnologie 32: 803.

Kim, I.-H., & S.-K. Choi, 2003. Copepod parasites (Crustacea) of freshwater fishes in Korea. Korean Journal of
Systematic Zoology 19: 57–93.

Kmentová, N., C. Hahn, S. Koblmüller, H. Zimmermann, J. Vorel, T. Artois, M. Gelnar, & M. P. M. M.
Vanhove, 2021. Contrasting host-parasite population structure: morphology and mitogenomics of a parasitic
flatworm on pelagic deepwater cichlid fishes from Lake Tanganyika. Biology 10: 797.

Kmentová, N., K. Thys, C. Hahn, J. Vorel, S. Koblmüller, M. Van Steenberge, A. Chocha Manda, L. Makasa,
T. Mulimbwa N'sibula, P. Masilya Mulungula, & M. P. M. Vanhove, 2023. Comparative population
mitogenomics of fish parasites reveals contrasting geographic pattern in the pelagic zone of Lake Tanganyika
[PREPRINT]. Authorea 10.22541/au.167274604.41657501/v1.

Koch, M., S. Koblmüller, K. M. Sefc, N. Duftner, C. Katongo, & C. Sturmbauer, 2007. Evolutionary history of
the endemic Lake Tanganyika cichlid fish *Tylochromis polylepis*: a recent intruder to a mature adaptive
radiation. Journal of Zoological Systematics and Evolutionary Research 45: 64–71.

Kuchta, R., L. Basson, C. Cook, I. Fiala, P. Bartošová-Sojková, & E. Řehulková, 2018. A systematic survey of
the parasites of freshwater fishes in Africa In Scholz, T., M. P. M. Vanhove, N. Smit, Z. Jayasundera, & M.
Gelnar (eds), A guide to the parasites of African freshwater fishes. CEBioS, Royal Belgian Institute of Natural
Sciences, Brussels, Belgium: 137–402.

Kvach, Y., M. Y. Tkachenko, M. Seifertová, & M. Ondračková, 2021. Insights into the diversity, distribution
and phylogeny of three ergasilid copepods (Hexanauplia: Ergasilidae) in lentic water bodies of the Morava river
basin, Czech Republic. Limnologica 91: 125922.

Lamboj, A., 2004. The cichlid fishes of Western Africa. Schmettkamp, Bornheim, Germany.

Librado, P., & J. Rozas, 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data.
Bioinformatics 25: 1451–1452.

Lorenz, R., S. H. Bernhart, C. Höner zu Siederdissen, H. Tafer, C. Flamm, P. F. Stadler, & I. L. Hofacker, 2011.
ViennaRNA Package 2.0. Algorithms for Molecular Biology 6: 26.

Lowe, T. M., & P. P. Chan, 2016. tRNAscan-SE On-line: integrating search and context for analysis of transfer
RNA genes. Nucleic Acids Research 44: W54–W57.

Markewitsch, A. P., 1937. Copepoda parasitica prisnikh vod SRSR. Copepoda parasitica der Binnengewässer
der USSR. Institut Zoologii i Biologii, Akademiya Nauk Ukrainii SSR, Kyiv, USSR.

Mathews, P. D., A. C. M. F. Patta, G. S. Gama, & O. Mertins, 2017. Infestation by *Ergasilus coatiarus*
(Copepoda: Ergasilidae) in two Amazonian cichlids with new host record from Peru: an ectoparasites natural
control approach. Comptes Rendus. Biologies 341: 16–19.

Oldewage, W. H., & J. G. van As, 1988. A key for the identification of African piscine parasitic Ergasilidae
(Copepoda: Poecilostomatoida). South African Journal of Zoology 23: 42–46.

Míč, R., E. Řehulková, & M. Seifertová, 2023. Species of *Ergasilus* von Nordmann, 1832 (Copepoda:
Ergasilidae) from cichlid fishes in Lake Tanganyika. Parasitology 150: 579–598.

Míč, R., E. Řehulková, A. Šimková, J. R. Razanabolana, & M. Seifertová, 2024. New species of
Dermoergasilus Ho & Do, 1982 (Copepoda: Cyclopoida: Ergasilidae) parasitizing endemic cichlid *Paretroplus*
polyactis (Bleeker) in Madagascar. Parasitology 151: 319–336.

Michels, J., 2007. Confocal laser scanning microscopy: using cuticular autofluorescence for high resolution
morphological imaging in small crustaceans. Journal of Microscopy 227: 1–7.

Michels, J., & M. Büntzow, 2010. Assessment of Congo red as a fluorescence marker for the exoskeleton of
small crustaceans and the cuticle of polychaetes. Journal of Microscopy 238: 95–101.

Minh, B. Q., H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. Von Haeseler, R. Lanfear, & E.
Teeling, 2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era.
Molecular Biology and Evolution 37: 1530–1534.

Minxiao, W., S. Song, L. Chaolun, & S. Xin, 2011. Distinctive mitochondrial genome of calanoid copepod
Calanus sinicus with multiple large non-coding regions and reshuffled gene order: useful molecular markers for
phylogenetic and population studies. BMC Genomics 12: 73.

Mirande, J. M., 2009. Weighted parsimony phylogeny of the family Characidae (Teleostei: Characiformes).
Cladistics 25: 574–613.

Mirande, J. M., 2019. Morphology, molecules and the phylogeny of Characidae (Teleostei, Characiformes).
Cladistics 35: 282–300.

Murray, H. M., S. J. Hill, & K. P. Ang, 2016. The external morphology of adult female *Egrasilus* [sic] *labracis*
as shown using hexamethyldisilazane treated, uncoated specimens for scanning electron microscopy.
Microscopy Research and Technique 79: 657–663.

Muterezi Bukinga, F., M. P. M. Vanhove, M. Van Steenberge, & A. Pariselle, 2012. Ancyrocephalidae
(Monogenea) of Lake Tanganyika: III: *Cichlidogyrus* infecting the world's biggest cichlid and the non-endemic
tribes Haplochromini, Oreochromini and Tylochromini (Teleostei, Cichlidae). Parasitology Research 111:
2049–2061.

Nadler, S. A., E. Bolotin, & S. P. Stock, 2006. Phylogenetic relationships of *Steinernema* Travassos, 1927
(Nematoda: Cephalobina: Steinernematidae) based on nuclear, mitochondrial and morphological data.
Systematic Parasitology 63: 161–181.

Oliveira, M. S. B., L. L. Corrêa, E. A. Adriano, & M. Tavares-Dias, 2021. Integrative taxonomy of a new
species of *Therodamas* (Ergasilidae) infecting the Amazonian freshwater fish *Leporinus fasciatus*
(Anostomidae). Parasitology Research 120: 3137–3147.

Ondračková, M., J. Fojtů, M. Seifertová, Y. Kvach, & P. Jurajda, 2019. Non-native parasitic copepod
Neoergasilus japonicus (Harada, 1930) utilizes non-native fish host *Lepomis gibbosus* (L.) in the floodplain of
the River Dyje (Danube basin). Parasitology Research 118: 57–62.

O'Reilly, C. M., S. R. Alin, P. D. Piisnier, A. S. Cohen, & B. A. McKee, 2003. Climate change decreases
aquatic ecosystem productivity of Lake Tanganyika, Africa. Nature 424: 766–768.

Paperna, I., 1969. Parasitic crustacea from fishes of the Volta Basin and South Ghana. Revue de Zoologie et de
Botanique Africaine 53: 208–216.

Prjibelski, A., D. Antipov, D. Meleshko, A. Lapidus, & A. Korobeynikov, 2020. Using SPAdes *de novo*
assembler. Current Protocols in Bioinformatics 70: e102.

R Core Team, 2023. R: a language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria, Accessed February 10, 2023. https://www.R-project.org/.

Rahmouni, C., M. P. M. Vanhove, & A. Šimková, 2017. Underexplored diversity of gill monogeneans in
cichlids from Lake Tanganyika: eight new species of *Cichlidogyrus* Paperna, 1960 (Monogenea:
Dactylogyridae) from the northern basin of the lake, with remarks on the vagina and the heel of the male
copulatory organ. Parasites & Vectors 10: 591.

Rahmouni, C., M. P. M. Vanhove, & A. Šimková, 2018. Seven new species of *Cichlidogyrus* Paperna, 1960
(Monogenea: Dactylogyridae) parasitizing the gills of Congolese cichlids from northern Lake Tanganyika. PeerJ
6: e5604.

Rambaut, A., A. J. Drummond, D. Xie, G. Baele, & M. A. Suchard, 2018. Posterior summarization in Bayesian
phylogenetics using Tracer 1.7. Systematic Biology 67: 901–904.

Ranwez, V., E. J. P. Douzery, C. Cambon, N. Chantret, & F. Delsuc, 2018. MACSE v2: Toolkit for the
alignment of coding sequences accounting for frameshifts and stop codons. Molecular Biology and Evolution
35: 2582–2584.

Ranwez, V., S. Harispe, F. Delsuc, & E. J. P. Douzery, 2011. MACSE: Multiple alignment of coding SEquences
accounting for frameshifts and stop codons. PLoS ONE 6: e22594.

Rindoria, N. M., Z. Gichana, G. N. Morara, C. van Wyk, W. J. Smit, N. J. Smit, & W. J. Luus-Powell, 2023.
Scanning electron microscopy and first molecular data of two species of *Lamproglena* (Copepoda: Lernaeidae)
from *Labeo victorianus* (Cyprinidae) and *Clarias gariepinus* (Clariidae) in Kenya. Pathogens 12: 980.

Ronco, F., H. H. Büscher, A. Indermaur, & W. Salzburger, 2020. The taxonomic diversity of the cichlid fish
fauna of ancient Lake Tanganyika, East Africa. Journal of Great Lakes Research 46: 1067–1078.

Ronquist, F., M. Teslenko, P. Van Der Mark, D. L. Ayres, A. Darling, S. Höhna, B. Larget, L. Liu, M. A.
Suchard, & J. P. Huelsenbeck, 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice
across a large model space. Systematic Biology 61: 539–542.

Salzburger, W., 2018. Understanding explosive diversification through cichlid fish genomics. Nature Reviews
Genetics 19: 705–717.

Salzburger, W., B. van Bocxlaer, & A. S. Cohen, 2014. Ecology and evolution of the African Great Lakes and
their faunas. Annual Review of Ecology, Evolution, and Systematics 45: 519–545.

Santacruz, A., F. N. Morales-Serna, M. Leal-Cardín, M. Barluenga, & G. Pérez-Ponce de León, 2020. *Acusicola*
margulisae n. sp. (Copepoda: Ergasilidae) from freshwater fishes in a Nicaraguan crater lake based on
morphological and molecular evidence. Systematic Parasitology 97: 165–177.

Sars, G. O., 1909. Zoological results of the third Tanganyika expedition, conducted by Dr. W. A. Cunnington,
F.Z.S., 1904–1905.—Report on the Copepoda. Proceedings of the Zoological Society of London 79: 31–77.

Sarvala, J., K. Salonen, M. Järvinen, E. Aro, T. Huttula, P. Kotilainen, H. Kurki, V. Langenberg, P. Mannini, A.
Peltonen, P.-D. Plisnier, I. Vuorinen, H. Mölsä, & Ossi, & V. Lindqvist, 1999. Trophic structure of Lake
Tanganyika: carbon flows in the pelagic food web. Hydrobiologia 407: 149–173.

Sasal, P., D. Mouillot, R. Fichez, S. Chifflet, & M. Kulbicki, 2007. The use of fish parasites as biological
indicators of anthropogenic influences in coral-reef lagoons: a case study of Apogonidae parasites in New-
Caledonia. Marine Pollution Bulletin 54: 1697–1706.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S.
Saalfeld, B. Schmid, J. Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, & A. Cardona, 2012.
Fiji: an open-source platform for biological-image analysis. Nature Methods 9: 676–682.

Schlebusch, R., 2014. Phylogeny of the African genus *Ergasilus* (Copepoda: Poecilostomatoida). Master's
thesis, University of the Free State, Bloemfontein, South Africa.

Schluter, D., 2000. The ecology of adaptive radiation. Oxford Series in Ecology and Evolution. Oxford Series in
Ecology and Evolution, Oxford, UK.

Seo, T.-K., H. Kishino, & J. L. Thorne, 2005. Incorporating gene-specific variation when inferring and
evaluating optimal evolutionary tree topologies from multilocus sequence data. Proceedings of the National
Academy of Sciences 102: 4436–4441.

Song, Y., G. T. Wang, W. J. Yao, Q. Gao, & P. Nie, 2008. Phylogeny of freshwater parasitic copepods in the
Ergasilidae (Copepoda: Poecilostomatoida) based on 18S and 28S rDNA sequences. Parasitology Research 102:
299–306.

Sterling-Montealegre, R. A., & C. F. Prada, 2024. Variability and evolution of gene order rearrangement in
mitochondrial genomes of arthropods (except Hexapoda). Gene 892: 147926.

Talavera, G., & J. Castresana, 2007. Improvement of phylogenies after removing divergent and ambiguously
aligned blocks from protein sequence alignments. Systematic Biology 56: 564–577.

Thieme, M. L., R. Abell, M. L. J. Stiassny, P. Skelton, B. Lehner, G. G. Teugels, E. Dinerstein, A. K. Toham,
N. Burgess, & D. Olson, 2005. Freshwater ecoregions of Africa and Madagascar: a conservation assessment.
Island Press, Washington, USA.

Thurston, J. P., 1970. The incidence of Monogenea and parasitic Crustacea on the gills of fish in Uganda. Revue
de Zoologie et de Botanique Africaines 82: 111–130.

van Douwe, C., 1912. Copepoden des ostafrikanischen Seengebietes Wissenschaftliche Ergebnisse der
deutschen Zentral-AfriKa-Expedition, 1907-1908□: unter Führung Adolf Friedrichs, Herzogs zu Mecklenburg.
Klinkhardt & Biermann, Leipizig, Germany: 487–496.

Vanhove, M. P. M., A. Pariselle, M. Van Steenberge, J. A. M. Raeymaekers, P. I. Hablützel, C. Gillardin, B.
Hellemans, F. C. Breman, S. Koblmüller, C. Sturmbauer, J. Snoeks, F. A. M. Volckaert, & T. Huyse, 2015.
Hidden biodiversity in an ancient lake: phylogenetic congruence between Lake Tanganyika tropheine cichlids
and their monogenean flatworm parasites. Scientific Reports 5: 13669.

Vanhove, M. P. M., J. Snoeks, F. A. M. Volckaert, & T. Huyse, 2011. First description of monogenean parasites
in Lake Tanganyika: the cichlid *Simochromis diagramma* (Teleostei, Cichlidae) harbours a high diversity of
Gyrodactylus species (Platyhelminthes, Monogenea). Parasitology 138: 364–380.

Vanhove, M. P. M., B. Tessens, C. Schoelinck, U. Jondelius, D. T. J. Littlewood, T. Artois, & T. Huyse, 2013.
Problematic barcoding in flatworms: a case-study on monogeneans and rhabdocoels (Platyhelminthes).
ZooKeys 365: 355–379.

Vasquez, A. A., B. L. Bonnici, D. R. Kashian, J. Trejo-Martinez, C. J. Miller, & J. L. Ram, 2022. The
biodiversity of freshwater crustaceans revealed by taxonomy and mitochondrial DNA barcodes In Zupo, V. (ed),
Crustaceans. Endocrinology, biology and aquaculture. CRC Press, Boca Raton, FL, USA: 166–181.

Walter, T. C., & G. Boxshall, 2023. World of Copepods Database. In O. Bánki, Y. Roskov, M. Döring, G.
Ower, D. R. Hernández Robles, C. A. Plata Corredor, T. Stjernegaard Jeppesen, A. Örn, L. Vandepitte, D.
Hobern, P. Schalk, R. E. DeWalt, K. Ma, J. Miller, T. Orrell, R. Aalbu, J. Abbott, R. Adlard, E. M.
Adriaenssens, et al. Catalogue of Life Checklist. Accessed March 10, 2024. https://doi.org/10.48580/dfgnm-3g4.

Weydmann, A., A. Przyłucka, M. Lubośny, K. S. Walczyńska, E. A. Serrão, G. A. Pearson, & A. Burzyński,
2017. Mitochondrial genomes of the key zooplankton copepods Arctic *Calanus glacialis* and North Atlantic
Calanus finmarchicus with the longest crustacean non-coding regions. Scientific Reports 7: 13702.

Wick, R. R., M. B. Schultz, J. Zobel, & K. E. Holt, 2015. Bandage: interactive visualization of de novo genome
assemblies. Bioinformatics 31: 3350–3352.

Wickham, H., 2016. ggplot2: elegant graphics for data analysis. Springer, New York, USA.

Wilson, C. B., 1924. Parasitic copepods from the White Nile and the Red Sea In Jägerskiöld, L. A. (ed), Results
of the Swedish Zoological Expedition to Egypt and the White Nile 1901. Part 5. Library of the Royal University
of Uppsala, Uppsala, Sweden: 1–17.

Yamaguti, S., 1936. Parasitic copepods from fishes of Japan. Part I. Cyclopoda, 1. Kyoto Imperial University,
Kyoto, Japan.

Yu, G., T. T.-Y. Lam, H. Zhu, & Y. Guan, 2018. Two methods for mapping and visualizing associated data on
phylogeny using ggtree. Molecular Biology and Evolution 35: 3041–3043.

Yu, G., D. K. Smith, H. Zhu, Y. Guan, & T. T.-Y. Lam, 2017. ggtree: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. Methods in Ecology and
Evolution 8: 28–36.

Zhang, X. L., P. Liu, S. L. Xu, E. Z. Rizo, Q. Zhang, H. J. Dumont, & B. P. Han, 2021. Geographic variation of
Phyllodiaptomus tunguidus Mmtogenomes: genetic differentiation and phylogeny. Frontiers in Genetics 12:
711992.

Zirbel, M. J., C. B. Miller, & H. P. Batchelder, 2007. Staging egg development of marine copepods with DAPI
and PicoGreen®. Limnology and Oceanography: Methods 5: 106–110.

 Tables

772 **Table 1**: Specimen data of copepods used for phylogenetic analyses including host species, GenBank accession numbers, and reference study.

				GenBank accession numbers			
Parasite species	**Host species**	**Family**	**Isolate**	**18S rDNA**	**28S rDNA**	*cox1*	**Reference**
Acusicola margulisae Santacruz, Morales-Serna, Leal-Cadín, Barluenga & Pérez-Ponce de León, 2020	*Amphilophus citrinellus* (Günther, 1864)	Cichlidae	513AcitAsL			MN854841.1	Santacruz et al. (2020)
A. margulisae	*A. citrinellus*	Cichlidae	770AcitAsL			MN854869.1	Santacruz et al. (2020)
A. margulisae	Not specified		824N	MN852694	MN852851.1		Santacruz et al. (2020)
A. margulisae	*Oreochromis* sp.	Cichlidae	496TilaAsL			MN854838.1	Santacruz et al. (2020)
A. margulisae	*Oreochromis* sp.	Cichlidae	580TilaAsL			MN854846.1	Santacruz et al. (2020)
A. margulisae	*Oreochromis* sp.	Cichlidae	752TilaAsL			MN854855.1	Santacruz et al. (2020)
Dermoergasilus madagascarensis Míč, Řehulková, Šimková, Razanabolana & Seifertová, 2024	*Paretroplus polyactis* Bleeker, 1878	Cichlidae		PP115568.1	PP115569.1		Míč et al. (2024)
Ergasilus anchoratus Markevich, 1946	*Tachysurus fulvidraco* (Richardson, 1846)	Bagridae	EAH	DQ107564	DQ107528		Song et al. (2008)
E. anchoratus	*T. fulvidraco*	Bagridae				MZ557411	Hua et al. (2023)
E. briani Markevich, 1933	*Misgurnus anguillicaudatus* (Cantor, 1842)	Cobitidae	EBN	DQ107572	DQ107532		Song et al. (2008)
E. caparti Míč, Řehulková & Seifertová, 2023	*Spathodus erythrodon* Boulenger, 1900	Cichlidae	SPER_67	OQ407469	OQ407474		Míč et al. (2023)

E. hypomesi Yamaguti, 1936	*Acanthogobius hasta* (Temminck & Schlegel, 1845)	Gobiidae	EHX	DQ107573	DQ107539		Song et al. (2008)
E. kandti van Douwe, 1912	*Tylochromis polylepis* (Boulenger, 1900)	Cichlidae	TYPO_C5	PQ249840	PQ249840	PQ276880	This study
E. kandti	*T. polylepis*	Cichlidae	TYPO_C6	PQ249839	PQ249841		This study
E. kandti	*T. polylepis*	Cichlidae	TYPO_C13		PQ249843		This study
E. macrodactylus (Sars G.O., 1909)	*Gnathochromis permaxillaris* (David, 1936)	Cichlidae	GNPE_159	OQ407465.1	OQ407470		Míč et al. (2023)
E. megacheir (Sars G.O., 1909)	*Simochromis diagramma* (Günther, 1893)	Cichlidae	SIDI_EM2	OQ407466.1	OQ407471		Míč et al. (2023)
E. mirabilis Oldewage & van As, 1987	*Clarias gariepinus* Burchell, 1822	Clariidae	P38K-38VR	OR449753.1	OR449755.1	OR448769	Fikiye et al. 2023
E. mirabilis	*C. gariepinus*	Clariidae	P49K-72ZR	OR449754.1	OR449756.1	OR448770	Fikiye et al. 2023
E. parasarsi Míč, Řehulková & Seifertová, 2023	*Ophthalmotilapia nasuta* (Poll & Matthes, 1962)	Cichlidae	OPNA_99	OQ407467	OQ407473		Míč et al. (2023)
E. parasiluri (Yamaguti, 1936)	*Silurus asotus* Linnaeus, 1758	Siluridae	PPN	DQ107567	DQ107537		Song et al. (2008)
E. parasiluri	*T. fulvidraco*	Bagridae	PPH	DQ107568	DQ107536		Song et al. (2008)
E. parvus Míč, Řehulková & Seifertová, 2023	*S. erythrodon*	Cichlidae	SPER_70	OQ407468	OQ407472		Míč et al. (2023)
E. peregrinus Heller, 1865	*Siniperca chuatsi* (Basilewsky, 1855)	Sinipercidae	EPG	DQ107577	DQ107531		Song et al. (2008)
E. scalaris Markevich, 1940	*Pseudobagrus vachellii* (Richardson, 1846)	Bagridae	ESW	DQ107565			Song et al. (2008)
E. scalaris	*Tachysurus dumerili* (Bleeker, 1864)	Bagridae	ESC	DQ107566	DQ107538		Song et al. (2008)

E. sieboldi von Nordmann, 1832	*Perca fluviatilis* Linnaeus, 1758	Percidae	U_Jezu_9Erg	MW810238	MW810242		Kvach et al. (2021)
E. tumidus Markevich, 1940	*Acheilognathus taenianalis* (Günther, 1873)	Cyprinidae	ETP	DQ107569	DQ107534		Song et al. (2008)
E. tumidus	*A. taenianalis*	Cyprinidae	EXK	DQ107570	DQ107533		Song et al. (2008)
E. tumidus	*A. taenianalis*	Cyprinidae	EXM	DQ107571	DQ107535		Song et al. (2008)
E. wilsoni Markevich, 1933	plankton sampling		LEGO-POE014	KR048765.1	KR048843.1	KR049036.1	Baek et al. 2016
E. yaluzangbus Kuang & Qian, 1985	*Oxygymnocypris stewartii* (Lloyd, 1908)	Cyprinidae	EYJ	DQ107578	DQ107540		Song et al. (2008)
Lamproglena chinensis Yü, 1937	*Channa argus* (Cantor, 1842)	Channidae	LCW	DQ107553	DQ107545		Song et al. (2008)
L. clariae Fryer, 1956	*C. gariepinus*	Clariidae	UL241	OR242503.1	OR338196.1	OR232208.1	Rindoria et al. (2023)
L. clariae	*C. gariepinus*	Clariidae	UL242	OR242504.1		OR232209.1	Rindoria et al. (2023)
L. cleopatra Humes, 1957	*Labeo victorianus* Boulenger, 1901	Cyprinidae	UL237	OR242502.1	OR338170.1	OR232207.1	Rindoria et al. (2023)
L. orientalis Markevich, 1936	*Chanodichthys dabryi* (Bleeker, 1871)	Cyprinidae	LOQ	DQ107549	DQ107542		Song et al. (2008)
L. orientalis	*C. erythropterus* (Basilewsky, 1855)	Cyprinidae	LOH	DQ107551	DQ107541		Song et al. (2008)
L. orientalis	*C. mongolicus* (Basilewsky, 1855)	Cyprinidae	LOM	DQ107550	DQ107543		Song et al. (2008)
L. orientalis	*Squaliobarbus curriculus* (Richardson, 1846)	Cyprinidae	LOC	DQ107552	DQ107544		Song et al. (2008)
Lernaea cyprinacea Linnaeus, 1758	*C. erythropterus*	Cyprinidae	LCC	DQ107556	DQ107547		Song et al. (2008)

Species	Host	Family	Code				Reference
L. cyprinacea	*Cyprinus carpio* Linnaeus, 1758	Cyprinidae	LCE	DQ107555			Song et al. (2008)
L. cyprinacea	*Hemiculter leucisculus* (Basilewsky, 1855)	Cyprinidae	LCH	DQ107554	DQ107546		Song et al. (2008)
L. cyprinacea	*Opsariichthys bidens* Günther, 1873	Cyprinidae	LCM	DQ107557	DQ107548		Song et al. (2008)
Mytilicola intestinalis Steuer, 1902	*Mytilus edulis* Linnaeus, 1758	Mytilidae	Mi2	HM775187		HM775191.1	Elsner et al. (2011)
M. intestinalis	*M. edulis*	Mytilidae	Mo6	HM775188		HM775192.1	Elsner et al. (2011)
M. orientalis Mori, 1935	*Magallana gigas* (Thunberg, 1793)	Ostreidae	Mo10	HM775190		HM775196.1	Elsner et al. (2011)
M. orientalis	*M. gigas*	Ostreidae	Mo9	HM775189		HM775195.1	Elsner et al. (2011)
Neoergasilus japonicus (Harada, 1930)	*Lepomis gibbosus* (Linnaeus, 1758)	Centrarchidae	Babice2_3Neoerg	MW810236	MW810240		Kvach et al. (2021)
N. japonicus	*L. gibbosus*	Centrarchidae	Hvezda	MH167969	MH167967		Ondračková et al. (2019)
N. japonicus	*L. gibbosus*	Centrarchidae	Rohlik	MH167970	MH167968		Ondračková et al. (2019)
N. japonicus	*L. gibbosus*	Centrarchidae	U_Jezu_4_5Neoerg	MW810237	MW810241		Kvach et al. (2021)
N. japonicus	plankton sampling		LE080317-626			MZ964938	Vasquez et al. (2022)
N. japonicus	plankton sampling		LE080317-727			MZ964935	Vasquez et al. (2022)
N. japonicus	plankton sampling		LE080317-828			MZ964936	Vasquez et al. (2022)
N. japonicus	plankton sampling		LE080317-929			MZ964937	Vasquez et al. (2022)

Species	Host	Family	Code				Reference
N. japonicus	plankton sampling		LEGO-POE015	KR048752.1	KR048823.1	KR049037.1	Baek et al. (2016)
Ostrincola japonica Tanaka, 1961	plankton sampling		LEGO-POE023			KR049041.1	Baek et al. (2016)
Paraergasilus brevidigitus Yin, 1954	*C. carpio*	Cyprinidae	PBL	DQ107576	DQ107530		Song et al. (2008)
P. longidigitus Yin, 1954	*Abramis brama* (Linnaeus, 1758), *P. fluviatilis*, *Scardinius erythrophthalmus* (Linnaeus, 1758)	Cyprinidae, Percidae	Pahrbek_10_11_12_13Paraerg	MW810239	MW810243		Kvach et al. (2021)
P. medius Yin, 1956	*Ctenopharyngodon idella* (Valenciennes in Cuvier & Valenciennes, 1844)	Cyprinidae	PMC	DQ107574	DQ107529		Song et al. (2008)
P. medius	*Mylopharyngodon piceus* (Richardson, 1846)	Cyprinidae	PMQ	DQ107575			Song et al. (2008)
Pseudomyicola spinosus (Raffaele & Monticelli, 1885)	plankton sampling		LEGO-POE025	KR048751	KR048822.1	KT030282.1	Baek et al. (2016)
Sinergasilus major (Markevich, 1940)	*C. idella*	Cyprinidae	H1	OP076956.1		OQ160840.1	He et al. (2023)
S. major	*C. idella*	Cyprinidae	SMC	DQ107558	DQ107524		Song et al. (2008)
S. major	*Elopichthys bambusa* (Richardson, 1845)	Cyprinidae	SMG	DQ107560			Song et al. (2008)
S. major	*Silurus glanis* Linnaeus, 1758	Siluridae	PE2 - PE8	MZ047814	MZ047815		Dos Santos et al. (2021)
S. major	*Squaliobarbus curriculus* (Richardson, 1846)	Cyprinidae	SMCH	DQ107559			Song et al. (2008)
S. polycolpus (Markevich, 1940)	*Hypophthalmichthys molitrix* (Valenciennes, 1844)	Cyprinidae	SPL	DQ107563	DQ107525		Song et al. (2008)

S. polycolpus	*H. molitrix*	Cyprinidae				NC_028085.1	Feng et al. (2016)
S. undulatus (Markevich, 1940)	*Carassius auratus* (Linnaeus, 1758)	Cyprinidae	SUJ	DQ107562	DQ107527		Song et al. (2008)
S. undulatus	*C. carpio*	Cyprinidae	SUL	DQ107561	DQ107526		Song et al. (2008)
S. undulatus	*C. carpio*	Cyprinidae				MW080644	Hua et al. (2021
Therodamas longicollum Oliveira, Correa, Adriano & Tavares-Dias, 2021	*Leporinus fasciatus* (Bloch, 1794)	Anostomidae	Jarilandia	MW652731			Oliveira et al. (2021)
Trochicola enterica Dollfus, 1914	*Calliostoma zizyphinum* (Linneaus, 1758)	Calliostomatidae		AY627006			Huys et al. (2006)

773   **Table 2**: Best-fitting nucleotide substitution models for alignments of 18S rDNA, 28S rDNA, and each codon
774   position of COI mDNA selected by ModelFinder for use in IQ-Tree and MrBayes.

Alignment	Base pairs	Model IQ-Tree	Model MrBayes
18S rDNA	1500	TIM3e+I+R2	SYM+I+G4
28S rDNA	754	GTR+F+I+G4	GTR+F+I+G4
Cox1	1551	TIM2+F+G4	GTR+F+G4
		K3Pu+F+G4	F81+F+G4
		HKY+F+I+R2	HKY+F+G4

776   **Table 3**: Annotation table of the mitochondrial genome of *Ergasilus kandti*.

Region	Position	Length (bp)	Start/stop codon	Anticodon
Non-coding region	1 -> 215, 14,412 -> 14,639	443		
16S	1,093 -> 216	878		
atp6	1,968 -> 1,285	684	ATG/TAG	
atp8	2,044 -> 1,961	84	ATA/TAG	
trnK	2,115 -> 2,054	62		TTT
trnL1	2,243 -> 2,185	59		CAG
trnG	2,243 -> 2,185	59		TCC
nad4L	2,574 -> 2,243	332	ATG/TA	
nad6	3,046 -> 2,567	480	ATA/TAG	
12S	3,667 -> 3,047	621		
trnV	3,724 -> 3,667	58		TAC
cox3	4,514 -> 3,737	780	ATG/T	
cox1	4,527 -> 6,075	1549	ATA/T	
trnM	6,077 -> 6,139	63		CAT
trnP	6,140 -> 6,202	63		TGG
trnD	6,199 -> 6,256	58		GTC
trnE	6,255 -> 6,316	63		TTC
trnS1	6,366 -> 6,309	58		TCT
trnS2	6,367 -> 6,424	58		AGT
trnN	6,482 -> 6,421	62		GTT
nad3	6,826 -> 6,483	339	ATA/TA	
cox2	7,529 -> 6,833	697	TTG/T	
trnW	7,586 -> 7,527	60		TCA
nad1	7,529 -> 6,833	924	ATA/TAA	
trnF	8,566 -> 8,509	58		GAA
nad5	8,568 -> 10,268	1701	ATA/TAA	
trnQ	10,322 -> 10,261	62		TTG
nad2	14,358 -> 14,411	960	ATA/TAA	
trnY	11,286 -> 11,343	58		GTA
cytb	11,344 -> 12,480	1137	ATC/TAA	
trnL2	12,545 -> 12,485	61		TAA

nad4	12,564 -> 13,892	1329	ATA/TAG	
trnC	13,948 -> 13,891	58		GCA
trnR	14,002 -> 13,948	55		TCG
trnI	14,063 -> 14,003	61		GAT
trnA	14,122 -> 14,065	58		TGC
trnH	14,180 -> 14,123	58		GTG
trnT	14,358 -> 14411	54		ACA

List of figure captions

Fig. 1. Recorded locations of *Ergasilus kandti*. Geographical coordinates were estimated based on reports in the literature (i.e. geographical centers of lakes, approximate locations indicated in figures etc.) (see Supplementary Table S1). Borders indicate limits of freshwater ecoregions according to Thieme et al. (2005).

Fig. 2. Light microscope images of *Ergasilus kandti*. a: whole specimen, ventral view. b: detail tooth-like projection on third segment of second antenna. c: detail of the five-segmented antennule. d: caudal rami that terminates in two long (1, 3) and two short (2, 4) setae.

Fig. 3: Confocal laser microscopic images of *Ergasilus kandti*. a: dorsal view with (1) eyespot (circular cephalic structure) and (2) characteristic groove in the shape of an inverse T. b: detail of the reduced fifth leg, characteristic of species of *Ergasilus*. c: ventral view. d: detail of second antenna with lightly stained tooth-like projection (see arrow, greyscale was used for better contrast). e: detail of the five-segmented antennule with four setae at distal end.

Fig. 4: Mitochondrial genome of *Ergasilus kandti* with GC content (in black), positive (in green) and negative GC Skew (light orange) values.

Fig. 5: Sliding window analysis of the alignment across 13 protein coding regions among ergasilid copepods (*Ergasilus kandti, E. tumidus, Sinergasilus major, S. polycolpus* and *S. undulatus*). The line shows the value of nucleotide diversity (π) with a window size of 200 bp and a step size of 20; the value is inserted at its mid-point.

Fig. 6: Graphical representation of the ratios between the total number of non-synonymous (dN) and synonymous (dS) nucleotide substitutions per mitochondrial protein coding region across ergasilid copepods (*Ergasilus kandti, E. tumidus, Sinergasilus major, S. polycolpus*, and *S. undulatus*).

Fig. 7: Maximum-likelihood phylogram of ergasilid copepods based on partial 18S and 28S rDNA sequences also including mytilicolid, myicolid, and lernaeid specimens as outgroups (used for rooting). Support is indicated at nodes with ultrafast bootstrap values (UFBoot) and Shimodaira–Hasegawa-like approximate likelihood ratios (SH-aLRT), followed by Bayesian posterior probabilities (PP), and GC values inferred through symmetric resampling (P□=□0.33) in parsimony analysis; asterisk (*) indicates low or moderate support below the threshold (UFBoot□<□95, SH-aLRT□<□80, PP□<□95, and GC < 0); black dots, internal nodes with strong support across all support values; labels in bold are species occurring in Lake Tanganyika; labels in grey refer to isolate numbers in Table 1. Clade numbering (I-III) follows Song et al. (2008), with Clade IV added in the present study. Scale bar: estimated number of substitutions per site.

List of supplementary material.

Supplementary Table S1. Recorded locations of *Ergasilus kandti*. Geographical coordinates were estimated

based on reports in the literature (i.e. geographical centers of lakes, approximate locations indicated in figures

etc.)

Supplementary Table S2. Counts of spines and setae of specimens of *Ergasilus kandti* collected in the present

study. Empty cells indicate missing (unobserved) data. Cells highlighted in yellow indicate deviations from the

spine-seta formula of the species.

Supplementary Fig. S3. Maximum-likelihood phylogram of ergasilids copepods based on partial *cox1*

sequences. Support is indicated at nodes with ultrafast bootstrap values (UFBoot) and Shimodaira–Hasegawa-

like approximate likelihood ratios (SH-aLRT), followed by Bayesian posterior probabilities (PP), and GC values

inferred through symmetric resampling (P = 0.33) in parsimony analysis; asterisk (*) indicates low or

moderate support below the threshold (UFBoot < 95, SH-aLRT < 80, PP < 95, and GC < 0); black dots,

internal nodes with strong support across all support values. Labels in grey refer to isolate numbers in Table 1.

Scale bar: estimated number of substitutions per site.

Ergasilus kandti

◆ This study
● Previous studies

0   250   500 km