Outer membrane vesicles can contribute to cellulose degradation in *Teredinibacter turnerae,* a cultivable intracellular endosymbiont of shipworms

Mark T. Gasser[a]# | Annie Liu[a] | Marvin Altamia[b] | Bryan R. Brensinger[a] | Sarah L. Brewer[a] | Ron

Flatau[b] | Eric R. Hancock[a] | Sarah P. Preheim[c] | Claire Marie Filone[a] | Dan L. Distel[b]#

[a]Johns Hopkins University Applied Physics Laboratory, Laurel, Maryland, USA 20723

[b]Ocean Genome Legacy Center, Northeastern University, Nahant, Massachusetts, USA 01908

[c]Johns Hopkins University, Baltimore, Maryland, USA 21218

Running title: Cellulolytic Outer Membrane Vesicles from *T. turnerae*

Keywords: carbohydrate-active enzymes, CAZymes, lignocellulose degradation, OMVs, protein

secretion, symbiosis, *tonB*-dependent receptors

MTG: mark.gasser@jhuapl.edu (ORCID: 0000-0002-9396-4954) #Co-corresponding author

AL: annie.liu@jhuapl.edu (ORCID: 0009-0002-3224-6008)

MAA: marvin.altamia@gmail.com (ORCID: 0000-0002-8625-767X)

BRB: bryan.brensinger@jhuapl.edu (ORCID: 0009-0002-0327-021X)

SLB: sarah.brewer@jhuapl.edu (ORCID: 0009-0007-9039-0784)

RF: r.flatau@northeastern.edu (ORCID: 0000-0002-4357-5870)

ERH: eric.hancock@jhuapl.edu (ORCID: 0009-0002-3041-2148)

SPP: sprehei1@jhu.edu (ORCID: 0000-0001-7365-2382)

CMF: claire.marie.filone@jhuapl.edu (ORCID: 0000-0002-4041-3948)

DLD: d.distel@northeastern.edu (ORCID: 0000-0002-3860-194X) #Co-corresponding author

Data availability

28   The authors confirm that the data supporting this study's findings are available within the article

29   and its supplementary materials. The mass spectrometric raw files are accessible

30   at https://massive.ucsd.edu under accession MassIVE MSV000094250 and at

31   https://www.proteomexchange.org under accession PXD050419.

Funding statement

34   Research reported in this publication was supported by the following awards to DLD: National

35   Oceanic and Atmospheric Administration (NA19OAR0110303), Gordon and Betty Moore

36   Foundation (GBMF 9339), National Institutes of Health (1R01AI162943-01A1, subaward:

37   10062083-NE), and JHU/APL Internal Research and Development. The National Science

38   Foundation (DBI 1722553) also funded some equipment used in this research.

Conflict of interest statement

The authors report no conflict of interest. The funders had no role in study design, data

collection, interpretation, or the decision to submit the work for publication.

Abstract

Teredinibacter turnerae is a cultivable cellulolytic Gammaproeteobacterium (Cellvibrionaceae)

that commonly occurs as an intracellular endosymbiont in the gills of wood-eating bivalves of

the family Teredinidae (shipworms). The genome of *T. turnerae* encodes a broad range of

enzymes that deconstruct cellulose, hemicellulose, and pectin and contribute to lignocellulose

digestion in the shipworm gut. However, the mechanism by which symbiont-made enzymes are

secreted by *T. turnerae* and subsequently transported to the site of lignocellulose digestion in

the shipworm gut is incompletely understood. Here, we show that *T. turnerae* cultures grown on

carboxymethyl cellulose (CMC) produce outer membrane vesicles (OMVs) that contain a variety

of proteins identified by LC-MS/MS as carbohydrate-active enzymes with predicted activities

against cellulose, hemicellulose, and pectin. Reducing sugar assays and zymography confirm

that these OMVs retain cellulolytic activity, as evidenced by hydrolysis of CMC. Additionally,

these OMVs were enriched with *TonB*-dependent receptors, which are essential to carbohydrate

and iron acquisition by free-living bacteria. These observations suggest potential roles for OMVs

in lignocellulose utilization by *T. turnerae* in the free-living state, in enzyme transport and host

interaction during symbiotic association, and in commercial applications such as lignocellulosic

biomass conversion.

Introduction

The biological degradation of lignocellulose, a primary component of wood and all plant biomass, is a critical process in the global carbon cycle and has potential applications in renewable energy and chemical production through biomass conversion (Ragauskas *et al.*, 2014; Østby *et al.*, 2020). Lignocellulose is a complex composite material composed primarily of microfibrils of cellulose, the most abundant organic polymer on Earth (Dahmen *et al.*, 2019), embedded in a matrix of hemicellulose and lignin. Pectins may also be present in smaller quantities. However, due to its complexity, lignocellulose degradation requires a suite of specialized enzymes to convert its component polymers into accessible nutrients. This complexity precludes most organisms from utilizing lignocellulose and creates obstacles to bioconversion to renewable fuels or fine chemicals (Cragg *et al.*, 2015).

One of the unique biological systems where efficient wood digestion occurs is found within shipworms, a group of marine bivalve mollusks of the family Teredinidae and the primary degraders of woody plant material in mangrove forests and in association with driftwood and marine woodfalls (Voight, 2015; Cragg *et al.*, 2020). Shipworms harbor endosymbiotic bacteria in bacteriocytes confined to the gland of Deshayes, a tissue located within their gills (Distel *et al.*, 1991). The genomes of these bacteria, primarily from the genus *Teredinibacter*, encode a wide variety of enzymes targeting lignocellulose (Distel *et al.*, 2002; Yang *et al.*, 2009; O'Connor *et al.*, 2014; Altamia, Shipway, *et al.*, 2020; Altamia *et al.*, 2021) and express them within this gland (O'Connor *et al.*, 2014; Sabbadin *et al.*, 2018). The resulting bacterial enzymes are also found in the shipworm's cecum (O'Connor *et al.*, 2014; Sabbadin *et al.*, 2018; Altamia and Distel, 2022), a specialized organ that is the primary location of wood digestion for the shipworm and is nearly devoid of microbes (Betcher *et al.*, 2012). This physical separation

between the bacterial symbionts in the gill and wood digestion in the gut requires a transport

mechanism for symbiont-made cellulolytic enzymes. Recently, a system of ducts called the ducts

of Deshayes was shown to serve as an extracellular transport path for bacterial cellulases from

the gills to the mouth of shipworms (Altamia and Distel, 2022). However, the mechanism by

which enzymes produced by the intracellular bacteria move across multiple membranes to an

external transport path is still unknown.

Bacteria use a variety of mechanisms to degrade lignocellulosic substrates in their

environments. For example, polysaccharide-degrading bacteria, including the common rumen

bacteria *Bacteroides*, *Clostridia*, and *Fibrobacter,* use the type IX secretion pathway (T9SS) to

secrete large molecular-weight proteins, including multidomain carbohydrate-active enzymes

(CAZymes) (Gharechahi *et al.*, 2023). These enzymes may be released into the environment or

bound to the cell envelope (McGavin *et al.*, 1990; Cai *et al.*, 1999; Yan and Wu, 2013). In

addition to the secretion of soluble or membrane-bound enzymes, bacteria may export

CAZymes by producing outer membrane vesicles (OMVs). OMVs are spherical buds that

originate from the bacterial outer membrane and can contain various cellular components such

as lipopolysaccharides, proteins, small molecules, and nucleic acids (Schwechheimer and Kuehn,

2015; Huang *et al.*, 2022).

Recent studies have shown that OMVs can be highly enriched in CAZymes and retain

activity capable of degrading cellulose and other plant biomass (Arntzen *et al.*, 2017; Ichikawa

et al., 2019; Salvachúa *et al.*, 2020)*.* The majority of research on OMVs has been done on

pathogenic bacteria due to the capacity of OMVs to carry and deliver virulence factors and

toxins into host cells (O'Donoghue and Krachler, 2016; Lynch and Alegado, 2017; Caruana and

Walper, 2020). However, the role of OMVs in marine and other environments has been

increasingly recognized, likely playing crucial roles in nutrient processing and ecological

interactions (Frias *et al.*, 2010; Biller *et al.*, 2014, 2022; Fischer *et al.*, 2019; Fadeev *et al.*, 2023).

Teredinibacter turnerae, unlike obligate intracellular symbionts, is also capable of free-living

growth (Waterbury *et al.*, 1983) and may directly acquire carbon from cellulose or other plant

biomass in the marine environment (Naka and Haygood, 2023). Investigating the functionality of

OMVs in *T. turnerae* could be pivotal to understanding their complex symbiotic interaction and

the ecological impact of their dual lifestyle.

Here, we purify and characterize the protein composition of OMVs produced by *T.*

turnerae during growth on water-soluble carboxymethyl cellulose (CMC). Proteins putatively

involved in interactions with lignocellulose were identified by LC-MS/MS analysis and

comparison to the Carbohydrate Active Enzyme Database (CAZy). Additionally, activity assays

were used to show that OMVs produced by *T. turnerae* are capable of cellulose hydrolysis.

These observations suggest that OMVs contribute to cellulose utilization by *T. turnerae* and may

play essential roles in their metabolism in free-living and symbiotic states.

Experimental Procedures
Strain isolation, selection, and growth conditions
T. turnerae is an endosymbiont species with widespread occurrence among shipworm

species (Distel *et al.*, 2002; Altamia *et al.*, 2014; Altamia, Lin, *et al.*, 2020). Two strains were

selected: *T. turnerae* T7901 (ATCC 39867), which was previously isolated from *Bankia gouldi* and

was the first shipworm symbiont brought into pure culture (Waterbury *et al.*, 1983; Distel *et al.*,

2002; Yang *et al.*, 2009); and *T. turnerae* SR01903, a closely related strain of *T. turnerae* (Altamia

et al., 2014). *T. turnerae* SR01903 was isolated from the gills of a single specimen of *Lyrodus*

pedicellatus found in naturally occurring driftwood collected by hand in shallow water in the

Indian River Lagoon, Merit Island, FL. (N 28.40605 W 80.66034) on January 24, 2020. Gills were

removed by dissection and homogenized in 1.0 ml of SBM medium (Waterbury *et al.*, 1983) in

an autoclave-sterilized glass dounce homogenizer. Homogenate was streaked onto a culture

plate containing 1.0% Bacto agar shipworm basal medium (SBM) at pH 8.0 supplemented with

0.2% w/v powdered cellulose (Sigmacell Type 101; Sigma-Aldrich) and 0.025% NH_4Cl. Plates

were incubated at 30 °C until individual colonies could be observed. An individual colony was

then picked and subjected to multiple rounds of restreaking from single colonies to ensure

clonality. To identify the isolate, the 16S rRNA gene was PCR amplified, sequenced, and

compared to published *T. turnerae* genomes. The complete genome of *T. turnerae* SR01903 was

then sequenced (SRA number SRR28421271) and submitted to Genbank (Submission number

SUB14332655, Bioproject PRJNA1090931, assembly accession number pending; assembly and

annotation files provided in additional files for review but not for publication). For the

experiments described herein, strains were initially propagated in 6 mL cultures in shipworm

basal medium (Waterbury *et al.*, 1983) supplemented with 0.025% NH_4Cl and 0.2%

carboxymethyl cellulose for 4 days before being diluted 1/250 in fresh media and harvested

after 2 days (OD600 0.2-0.3). All cultures were incubated in a shaker incubator at 30 °C and 100

rpm. An overview of the procedures conducted in this research is presented in Figure 1.

Figure 1. Methods used to isolate and characterize outer membrane vesicles from T. turnerae. Diagram showing: (A) bacterial culture, (B) OMV isolation based on differential and density gradient separation, (C) OMV visualization and size by TEM and NTA, proteome analysis by LC-MS/MS, and (D) detection of cellulase activity in purified OMV preparations by zymography and reducing sugar (DNS) assay. Created with BioRender.com

Isolation of Outer Membrane Vesicles (OMVs)

Bacterial cells were separated from the culture supernatant by centrifugation at 5,000 x

g for 20 minutes at 4 °C. The supernatant was then transferred to a fresh tube, and

centrifugation was repeated to remove residual bacterial cells. The final supernatant was

carefully collected without disturbing the remaining pellet and filtered through a 0.22 μm

polyethersulfone filter as an additional precaution to remove cells. The putative OMVs were

then pelleted from the filtrate by ultracentrifugation at 120,000 x g (T-647.5 rotor, Sorvall) for 90

minutes at 4 °C. For purification, the resulting pellet was resuspended in 0.1 M phosphate-

buffered saline (PBS) and fractionated by bottom-up density gradient ultracentrifugation using

Optiprep™ (iodixanol density gradient medium) as follows. The resuspended OMV-containing

pellet was mixed with 60% iodixanol solution to a final density of 45% (w/v) and placed at the

bottom of an ultracentrifuge tube. A discontinuous density gradient was generated using a

syringe and G21 needle to deposit layers of 40%, 35%, 30%, 20%, and 10% iodixanol with a final

layer of 0.25 mL 0.1 M PBS on top. The resulting gradient was then subjected to centrifugation

for 16 hours at 150,000 x g and 4 °C (SW55 Ti rotor, Beckman-Coulter). Sample banding was

visually observed at the interface of the 30% and 20% fractions (Supporting Information: Figure

S1). All fractions were collected, and iodixanol was removed by passive diffusion dialysis (1,000

kDa MWCO) in exchanging buffer of 50 mM ammonium bicarbonate pH 8.3 at 4 °C. OMV

samples were removed from the dialysis bag and concentrated under vacuum to 100 μL

(SPD121P SpeedVac, Thermo Scientific). Particle visualization and characterization proceeded

with the single 30% fraction.

Transmission electron microscopy (TEM)
Purified OMVs were diluted and absorbed onto 200 mesh formvar-treated and carbon-

coated copper grids for 60 seconds. Samples were then fixed for 5 minutes in 4% glutaraldehyde

in 0.1 M sodium cacodylate, and grids were stained with 1% aqueous uranyl acetate for 60

seconds and left to dry. OMVs were imaged in an FEI Tecnai T12 (Thermo Fisher) transmission

electron microscope at 80 KV with an AMT bottom-mount camera.

Nanoparticle tracking analysis (NTA)
NTA was performed using the ZetaView® PMX-220 Twin (Particle Metrix) configured with

488 nm and 640 nm lasers with long wave-pass cut-off filters (500 nm and 660 nm, respectively)

and a sensitive CMOS camera 640 x 480 pixels. Samples were diluted in 2 mL of 0.1 μm filtered

deionized water (18 MΩ/cm) to obtain a particle concentration between 1×10^7 and 1×10^8

particles/mL. The instrument was set to a sensitivity of 80, a shutter speed of 100, and a frame

rate of 30 frames per second. Each sample was measured at 11 different positions throughout

the sample cell, with 1 cycle of reading at each position to have a minimum of 1,000 traces. If

the number of traces was below 1,000 counts, some additional sample was flushed inside the

sample cell, and the acquisition was repeated. Post-acquisition parameters were set to a

minimum brightness of 20, a maximum size area of 1,000 pixels, a minimum size area of 10

pixels, and tracelength of 15 frames. Automated cell quality control was checked using high-

quality deionized water. Camera alignment and focus optimization were performed using

polystyrene Nanosphere™ 100 nm size standard beads. Data analysis was performed with

ZetaView® 8.05.14 software provided by the manufacturer. Automated reports of the particle

recordings across the 11 positions were manually checked, and any outlier position was

removed to calculate particle concentration and distribution.

Proteomic analysis
197   OMVs were lysed and denatured in a solution containing 5% sodium dodecyl sulfate

(SDS), 100 mM Tris (pH=8), 20 mM chloroacetamide, and 10 mM tris (2-carboxyethyl) phosphine

hydrochloride and incubated at 90 °C for 10 minutes. Proteins were aggregated and isolated

using Sera-Mag™ carboxylate-modified SpeedBeads (SP3 beads) and digested in 100 mM

NH_4HCO_3 containing Trypsin (final concentration 6 ng/μL) overnight at 37 °C and 1,200 rpm.

Digested samples were loaded onto C18 tips, and peptides were separated online using an 88-

minute LC gradient (Evosep LC system). MS analysis was performed on a Q-Exactive HF-X Hybrid

Orbitrap mass spectrometer (Thermo Scientific). One full high-resolution MS spectrum was

acquired with a resolution of 45,000, an AGC target of 3 x 10^6 with a maximum ion time of 45

ms, and a scan range of 400-1,500 m/z, followed by 20 HCD MS/MS scans with a resolution of

1,5000, an AGC target of 1 x 10^5 with a maximum ion time of 120 ms, NCE of 27, and isolation

window of 2 m/z. The resulting MS/MS spectra were searched against the strain-specific and

universal contaminant protein databases using Sequest within Proteome Discoverer 1.4.

Identifications were filtered to include only high-confidence peptides based on a better than 1%

false discovery rate (FDR) against a decoy database. Proteins were linked to peptides, and only

proteins with >2 peptide spectral matches (PSM) with two unique peptides were kept

(Supporting Information: Dataset S1).

Functional characterization of OMV proteins
Identified proteins were annotated against the Clusters of Orthologous Groups (COG) of

proteins database (Galperin *et al.*, 2021), and their subcellular locations were predicted using

CELLO v2.5 (http://cello.life.nctu.edu.tw/) (Yu *et al.*, 2006) and PSORTdb v4.0

(https://db.psort.org/) (Lau *et al.*, 2021). Enrichment and functional associations between

proteins were determined by STRINGdb v11 (https://string-db.org/) (Szklarczyk *et al.*, 2019) and

ShinyGO v0.8 (http://bioinformatics.sdstate.edu/go/) (Ge *et al.*, 2020). Settings were selected to

use all coding sequences in the T7901 genome as background and a minimum FDR stringency of

1×10^{-5}. We then established which OMV proteins are predicted to exhibit carbohydrate-active

properties using the annotated T7901 genome in the Carbohydrate Active Enzymes Database

(CAZy, https://cazy.org).

Enzymatic activity assays
The cellulolytic activity of proteins in OMV preparations was visualized and size

fractionated by denaturing polyacrylamide gel electrophoresis (SDS-PAGE, 1.5 mm thickness, 9%

acrylamide, and 0.2% CMC final concentration). For zymographic analysis, 14 µg of protein

sample determined by Pierce 660 assay (Thermo Scientific) using bovine serum albumin as

standard were heat denatured by boiling for 3 minutes in SDS without a reducing agent to

facilitate recovery of activity after refolding and added to each lane. After electrophoresis, the

SDS-PAGE gel was transferred to 500 mL of refolding buffer (20% isopropanol, 0.1% Triton X-100

in 1x PBS, pH 7.4) and gently shaken for 1 hour at room temperature. The gel was then

incubated in fresh 1x PBS buffer, pH 7.4, for 16 hours at room temperature before being stained

with 0.1% Congo Red in 1x PBS buffer, pH 7.4 for 1 hour. The zymogram was destained with 1 M

NaCl overnight to visualize regions where cellulase activity removed CMC. Additionally, the

production of reducing ends was measured by the colorimetric 3,5-dinitrosalicylic acid (DNS)

method (Ghose, 1987). Reactions were carried out in 96-well plates containing 50 μL 1% CMC,

30 μL 50 mM citrate buffer, and 20 μL purified OMV suspension and incubated for 4 hours at 37

°C. Next, 900 μL of DNS was added to each reaction, and reactions were heated at 99 °C for 10

minutes. Absorbance was measured at 540 nm, and sugar concentration was calculated by

comparison to a glucose standard curve. The enzyme activity unit was defined as the amount of

enzyme that liberated 1 μmol of reducing sugar per minute.

Results

OMV isolation and characterization

OMVs were isolated from cell-free supernatants of two closely related strains, T7901 and

SR01903, by density gradient ultracentrifugation. The presence of OMVs was confirmed by NTA

and TEM (Figure 2 A-D). The mean concentration of particles and average particle diameter size

determined by NTA were 1.67 x 10^9 particles/mL and 162.5 $\pm$ 49.7 nm for T7901 and 1.30 x 10^9

particles/mL and 206.6 $\pm$ 61.7 nm for SR01903. TEM images (Figure 2 C-D) show that the

putative OMV-enriched fractions from strains T7901 and SR01903 contain particles with size

distribution and morphology typical of bacterial OMVs.

Figure 2. Size, concentration, and visualization of OMVs isolated from T. turnerae culture supernatants after growth on carboxymethyl cellulose (CMC). (A-B) Size distribution and particle concentration for strains T7901 and SR01903, respectively. OMVs from SR01903 were slightly larger in average size than OMVs from T7901. (C-D) Isolated OMVs from strains T7901 and SR01903 were imaged using transmission electron microscopy. Scale bar = 100 nm.

OMV proteome analysis

The protein contents of OMVs from both strains were analyzed by LC-MS/MS. Peptide-generated data were used to search against T7901 and SR01903 annotated protein sequences to identify OMV proteins and their relative abundance. A cut-off of two or more unique

predicted peptides per identified protein was used, resulting in the identification of 472 and 569

proteins for T7901 and SR01903, respectively. The Top 100 most abundant proteins identified in

T7901 and SR01903 represented 81% and 74% of the total peptide content, respectively. Among

those proteins, 71 were identified in both strains, indicating that the two strains produce a

common core set of OMV proteins when grown under the examined culture condition. After

protein identification, PSORTdb v4.0 (Lau *et al.*, 2021) and CELLO v2.5 (Yu *et al.*, 2006) were

used to predict the subcellular location of all identified proteins. Outer membrane proteins

comprised most (~75% and ~60%, respectively) of the OMV protein content of both T7901 and

SR01903 (Figure 3 A). OMV proteins were clustered into orthologous groups (COG); 65% and

60% of OMV proteins for T7901 and SR01903 were assigned a single cluster, respectively. Most

OMV proteins assigned to clusters in both strains were categorized as Cell

Wall/Membrane/Envelope Biogenesis, Inorganic Ion Transport and Metabolism, and

Carbohydrate Transport and Metabolism (Figure 3 B).

Figure 3. T. turnerae OMV proteome profiles. (A) Relative protein abundance grouped by predicted subcellular location and (B) Clusters of Orthologous Groups (COG) functional categories of identified proteins. (C) Top 10 STRINGdb pathways (ranked by FDR) found in OMVs of T7901. Similar top pathways were found in OMVs of SR01903.

TonB-dependent receptors and glycosyl hydrolase activity are enriched in OMVs

To highlight functional associations, OMV protein content was analyzed using STRING

(Szklarczyk *et al.*, 2019) enrichment. We found significant enrichment of pathways involving

TonB-dependent receptors (TBDR) and glycosyl hydrolase activity (Figure 4 C). The most

abundant proteins were TBDRs, which constituted ~47% and ~44% of the OMV protein content

in T7901 and SR01903, respectively. TBDRs are known to be deployed for the uptake of sugars

from complex carbohydrates (Blanvillain *et al.*, 2007; Pollet *et al.*, 2021). Additionally, *TonB*

knockouts of *T. turnerae* were shown to have lost the ability to grow on cellulose (Naka and

Haygood, 2023), which suggests that the *TonB* system in *T. turnerae* is critical to cellulose

utilization.

Efficient utilization of lignocellulose for growth often requires a cocktail of multiple

enzymes targeting inner (endo-acting) and outer (exo-acting) polysaccharide glycosidic bonds

and various bonds found in heteroxylans. In total, 39 and 48 predicted CAZymes were present,

representing 30 and 29 different catalytic CAZy families and 8.3% and 8.4% of identified OMV

proteins in T7901 and SR01903, respectively. Most identified CAZy modules belong to glycoside

hydrolase families (GH) predicted to be involved in cellulose and hemicellulose degradation

(Figure 4 A) and that include several hydrolytic activities: endo-glucanases (GH5, GH8, GH9,

GH10, GH11, GH26, GH44, GH45, GH51, GH62), exo-glucanases (GH5, GH9, GH26GH43GH26,

GH43), and beta-glucosidases (GH2, GH3, GH5, GH39). In addition, multiple enzymes were

predicted to include non-catalytic carbohydrate-binding modules (CBMs), which can enhance

enzyme activity by increasing accessibility to the substrate (Hervé *et al.*, 2010).

Cellulase activity
Many OMV proteins were predicted to have or be associated with cellulolytic activity.

We confirmed these by testing OMV suspensions for hydrolytic activity against CMC by

zymography and reducing sugar DNS assay. In zymograms, clear zones indicating endoglucanase

activity (Teather and Wood, 1982) were visualized between 95-130 kDa for both strains after 16-

hour incubation (Figure 4 B). The production of reducing sugars from CMC was measured using

DNS assay, and when normalized to total protein content, OMVs produced from SR01903 had

the equivalent cellulolytic activity as a commercial soluble cellulase enzyme cocktail. Notably,

OMVs produced from T7901 showed ~10x the activity of the commercial cellulase cocktail

(Figure 4 C).

Figure 4. Carbohydrate-active modules and observed cellulase activity of T. turnerae OMVs. (A) Heatmap profile of the relative abundance of the predicted carbohydrate-active enzymes found in OMVs. (B) Denaturing SDS-PAGE (left) and zymogram (right; CMC substrate) of isolated OMVs. (C) Histogram showing cellulase activity (CMC substrate) in Units/mg (as determined by DNS assay) for isolated OMVs and a purified cellulase enzyme cocktail (Cellulase R-10, Goldbio CAS# 9012-54-8). Bars indicate means (error bars: standard deviations of three replicates).

Discussion

T. turnerae secretes a broad array of carbohydrate-active enzymes within the host's gills

and when grown in pure culture (Yang *et al.*, 2009; O'Connor *et al.*, 2014; Sabbadin *et al.*, 2018).

Genome sequence suggests these enzymes may be translocated across the inner membrane by

Sec and Sec-independent (twin-arginine, Tat) pathways and through the outer membrane by the

Type II (T2SS) generalized secretory pathway. The *T. turnerae* genome also encodes the

complete Type VI secretion system (T6SS), which mediates many types of bacteria-bacteria and

bacteria-host interactions (Yang *et al.*, 2009). However, the role of OMVs in the transport of

cellulolytic enzymes by *T. turnerae* has not been investigated. Here, we show that *T. turnerae*

strains T7901 and SR01903 produce OMVs that contain diverse carbohydrate-active enzymes,

including glycoside hydrolases, carbohydrate esterases, and polysaccharide lyase with predicted

activity against cellulose, hemicellulose, and pectin. We further show that these OMV

preparations can hydrolyze carboxymethyl cellulose (CMC) with specific activities comparable to

or greater than that observed for a commercial purified cellulase enzyme cocktail (Figure 4 C).

In purified OMVs isolated from *T. turnerae* strains T7901 and SR01903, we detected

representatives of 11 carbohydrate-binding module (CBM) families (CBM2, 5, 9, 10, 22, 35, 48,

50, 57, 85, and 91). Each of these families is known to bind cellulose or hemicellulose

components, except for CBM50, which typically binds peptidoglycan or chitin. Additionally,

catalytic modules representing 23 glycoside hydrolase (GH) families, four carbohydrate esterase

(CE) families, and one polysaccharide lyase (PL) were detected. All of these GH families (GH2, 3,

5, 8, 9, 10, 11, 13, 23, 26, 28, 31, 35, 39, 43, 44, 45, 51, 62, 67, 94, 105, and 115) target bonds

found within cellulose, hemicellulose or pectin, except GH23, which primarily acts on

peptidoglycan. Finally, the four carbohydrate esterase (CE) families detected (CE1, 2, 15, and 20)

target bonds found in hemicellulose, or bonds covalently linking hemicellulose to lignin. This

combination of modules indicates a complete lignocellulose deconstruction system, including

potential endo- and exo-activities against cellulose and hemicellulose backbones, debranching

of hemicellulose sidechains, and hydrolysis of the resulting oligomers and monomers.

The fact that this wide array of activities was found in OMVs produced by cells of *T.*

turnerae strains T7901 and SR01903 grown with cellulose as a sole carbon source indicates that

the expression of genes active against other lignocellulose components was either constitutive

or co-regulated with expression of genes targeting cellulose. The ability of *T. turnerae* to express

these activities in the absence of induction by their specific target substrates is consistent with

the observation that shipworm endosymbionts express a wide range of lignocellulolytic

enzymes when growing within the gill bacteriocytes of the host where they have no direct

contact with lignocellulose (O'Connor *et al.*, 2014; Sabbadin *et al.*, 2018). Notably, six of the

catalytic module families detected in *T. turnerae* OMVs (CE11, 15, and GH9, 10, 11, and 45) were

also detected in the cecum content of *Bankia setacea* in a previous investigation (O'Connor *et*

al., 2014); however, it should be noted that this shipworm species hosts four *Teredinibacter*

species, *T. waterburyi* (Altamia, Shipway, *et al.*, 2020), *T. franksiae*, *T. haidensis*, and *T. purpureus*

(Altamia *et al.*, 2021) but does not harbor *T. turnerae* (O'Connor *et al.*, 2014).

While diverse CAZymes are present in the OMV proteome of *T. turnerae*, they comprise

less than 10% of the total protein content of OMVs produced under the conditions examined

here. Interestingly, the most abundant proteins observed were identified as TonB-dependent

receptors (TBDR), accounting for nearly half of the proteins detected. TBDR enrichment has

been previously reported in OMVs of several Gram-negative bacteria (Veith *et al.*, 2015;

Zakharzhevskaya *et al.*, 2017; Dhurve *et al.*, 2022; Fadeev *et al.*, 2023). TBDRs are outer

membrane-associated proteins that bind and mediate the energy-dependent movement of

siderophores and various nutrients, including carbohydrates that are too large to be taken up

via transmembrane diffusion across the outer membrane (Silale and Van Den Berg, 2023). For

example, in the marine bacterium *Alteromonas macleodii*, TBDR genes were selectively

expressed in carbon- and iron-limiting conditions (Manck *et al.*, 2020) and during

polysaccharide utilization (Neumann *et al.*, 2015). Recently, it was shown that two of *T.*

turnerae's four *TonB* genes are essential for growth in iron-limiting conditions and for growth

with cellulose as a sole carbon source. This dependence indicates that the *TonB* system is

essential for both iron acquisition and cellulose catabolism by *T. turnerae* (Naka and Haygood,

2023).

Interestingly, in a proteomic examination of the shipworm *Banka setacea*, a TBDR was

the only major symbiont-derived protein observed in the cecum content that was not

associated with the decomposition of lignocellulose (O'Connor *et al.*, 2014). This finding

suggests the presence of symbiont outer membranes in the cecum. However, the cecum of

shipworms has been shown by microscopic (Betcher *et al.*, 2012; Pesante *et al.*, 2021),

transcriptomic (Sabbadin et al., 2018), and proteomic (O'Connor *et al.*, 2014) methods to be

nearly devoid of intact bacterial cells. These observations argue against the transport of whole

bacterial cells from the gill to the cecum as suggested in (Pesante *et al.*, 2021), but might

suggest a role for OMVs in transporting lignocellulose degrading enzymes from the gill to the

gut in shipworms.

Outer membrane vesicles are critical in bacterial interactions with many animal and

plant hosts (Berleman and Auer, 2013; Lynch and Alegado, 2017). While most research has

focused on pathogenic interactions, OMVs may also contribute to beneficial host-symbiont

associations. For example, OMVs have been shown to mimic whole cells of *Vibrio fisheri* in the

induction of specific aspects of light organ development in the Hawaiian bobtail squid *Euprymna*

scolopes (Aschtgen *et al.*, 2016)*.* OMVs have also been shown to be produced by

chemoautotrophic symbionts that occur in the trophosome of flatworms of the genus

Paracatenula, where they play a critical role in provisioning the host with carbohydrates

synthesized by the symbionts (Jäckle *et al.*, 2019)*.* In the human gut microbiome, members of

Bacteroides have been shown to tailor OMV content to specific polysaccharides (Sartorio *et al.*,

2023), and OMVs produced by human and ruminant gut bacteria, including Bacteroides,

Fibrobacter, and Clostridium, communally degrade polysaccharides into nutrients available to

the hosts (Rakoff-Nahoum *et al.*, 2014; Arntzen *et al.*, 2017; Gharechahi *et al.*, 2023).

Outer membrane vesicles can also transport proteins across the plasma membranes of

eukaryotic organisms (O'Donoghue and Krachler, 2016; Schorey *et al.*, 2021; Toyofuku *et al.*,

2023). While previous research has focused mainly on the role of OMVs in introducing bacterial

molecules into eukaryotic cells, OMVs may also transport molecules produced by intracellular

bacteria out of their eukaryotic host. For example, *Salmonella enterica,* which grows in vesicles

within their host's cells, produces toxin-loaded OMVs that escape the infected host cells and

deliver their toxic cargo to uninfected neighboring cells (O'Donoghue and Krachler, 2016).

Similarly, OMVs produced by *Mycobacterium tuberculosis* are released from infected

macrophages, exporting Mycobacterial lipoproteins and lipoglycans that affect the immune

functions of neighboring uninfected host cells (Athman *et al.*, 2015). Thus, OMV-mediated

mechanisms could potentially explain the transport of cellulolytic proteins produced by *T.*

turnerae in their intracellular location within the host's bacteriocytes to an extracellular

location, allowing subsequent transport to the host's gut.

Packaging enzymes within OMVs or bound to OMV surfaces can have distinct advantages

over their secretion as soluble proteins or proteins bound to the cell surface. For example,

OMVs may protect proteins from degradation in the environment (Bonnington and Kuehn,

2014; Alves *et al.*, 2016; Zingl *et al.*, 2021), selectively concentrate proteins with specific

functions (Orench-Rivera and Kuehn, 2021; Sartorio *et al.*, 2023), and deliver proteins to remote

locations in sufficient quantity to produce a desired effect without the dilution that would be

experienced by soluble proteins (Toyofuku *et al.*, 2023). These factors may be especially

important for the degradation of complex substrates like lignocellulose, where the simultaneous

delivery of specific sets of proteins in the correct proportions and spatial orientation can yield

significant synergistic interactions that enhance efficiency (Park *et al.*, 2014).

However, little is known about OMV production and its function in environmental

lignocellulose degradation. Several investigations of lignocellulose-degrading microorganisms

from soil suggest OMVs play critical roles in terrestrial plant biomass degradation. For example,

when grown in the presence of lignin, *Pseudomonas putida* produces OMVs containing enzymes

functionally active against lignin aromatic components (Salvachúa *et al.*, 2020). Similarly,

Trichoderma reesei, a filamentous fungus, produces OMVs containing a variety of cellulases

when grown in the presence of cellulose (De Paula *et al.*, 2019).

In addition to their potential roles in shipworm symbiosis and the environmental

turnover of lignocellulose in terrestrial and aquatic environments, cellulolytic OMVs are of

significant interest in converting plant biomass into renewable liquid fuels or fine chemicals

(Thakur *et al.*, 2023). A significant challenge in biomass conversion is understanding how

cooperation among cellulases and associated enzymes can improve the design of efficient

enzyme cocktails tailored to individual feedstocks (Østby *et al.*, 2020). OMVs produced by *T.*

turnerae and other lignocellulolytic organisms may represent nature-based solutions that reveal

specific combinations, concentrations, and spatial organization of cooperating enzymes that

significantly enhance lignocellulose degradation in nature and so might inspire the design of

engineered lignocellulose deconstruction systems.

Figure legends

Figure 1. Methods used to isolate and characterize outer membrane vesicles from *T. turnerae*.

Diagram showing: (A) bacterial culture, (B) OMV isolation based on differential and

density gradient separation, (C) OMV visualization and size by TEM and NTA, proteome

analysis by LC-MS/MS, and (D) detection of cellulase activity in purified OMV

preparations by zymography and reducing sugar (DNS) assay. Created with

BioRender.com.

Figure 2. Size, concentration, and visualization of OMVs isolated from *T. turnerae* culture

supernatants after growth on carboxymethyl cellulose (CMC). (A-B) Size distribution

and particle concentration for strains T7901 and SR01903, respectively. OMVs from

SR01903 were slightly larger in average size than OMVs from T7901. (C-D) Isolated

OMVs from strains T7901 and SR01903 were imaged using transmission electron

microscopy. Scale bar = 100 nm.

Figure 3. T. turnerae OMV proteome profiles. (A) Relative protein abundance grouped by

predicted subcellular location and (B) Clusters of Orthologous Groups (COG) functional

categories of identified proteins. (C) Top 10 STRINGdb pathways (ranked by FDR) found

in OMVs of T7901. Similar top pathways were found in OMVs of SR01903.

Figure 4. Carbohydrate-active modules and observed cellulase activity of *T. turnerae* OMVs. (A)

Heatmap profile of the relative abundance of the predicted carbohydrate-active

enzymes found in OMVs. (B) Denaturing SDS-PAGE (left) and zymogram (right; CMC

substrate) of isolated OMVs. (C) Histogram showing cellulase activity (CMC substrate)

in Units/mg (as determined by DNS assay) for isolated OMVs and a purified cellulase

enzyme cocktail (Cellulase R-10, Goldbio CAS# 9012-54-8). Bars indicate means (error

bars: standard deviations of three replicates).

Author Contribution Statement
Mark T. Gasser: Conceptualization; investigation; formal analysis; methodology; writing –

original draft preparation. **Annie Liu:** Investigation; methodology. **Marvin A. Altamia:**

Investigation; methodology. **Bryan R. Brensinger:** Investigation; methodology. **Sarah L. Brewer:**

Investigation; methodology. **Ron Flatau:** Investigation; methodology. **Eric R. Hancock:**

Investigation; methodology. **Sarah P. Preheim:** Supervision; writing – review and editing. **Claire**

Marie Filone: Funding acquisition; supervision; writing – review and editing. **Dan L. Distel:**

Funding acquisition; Conceptualization; supervision; writing – original draft preparation; writing

– review and editing.

Acknowledgments
We would like to thank the PennVet Extracellular Vesicle Core SCR_022444 (Philadelphia,

Pennsylvania) for work and assistance with OMV isolation and NTA analysis; the University of

Maryland School of Medicine's and School of Dentistry's Electron Microscopy Core Imaging

Facility - EMCIF (Baltimore, Maryland) for work and assistance with TEM imaging; and the NYU

Langone's Proteomics Laboratory SCR_017926 (New York, New York) for work and assistance

with mass spectrometry analysis.

Supplemental Information
Figure_S1_SuppInfo.docx

Dataset_S1_SuppInfo.xlsx

References

Altamia, M.A. and Distel, D.L. (2022) Transport of symbiont-encoded cellulases from the gill to
the gut of shipworms via the enigmatic ducts of Deshayes: a 174-year mystery solved.
Proc R Soc B **289**: 20221478.
Altamia, M.A., Lin, Z., Trindade-Silva, A.E., Uy, I.D., Shipway, J.R., Wilke, D.V., et al. (2020)
Secondary Metabolism in the Gill Microbiota of Shipworms (Teredinidae) as Revealed by
Comparison of Metagenomes and Nearly Complete Symbiont Genomes. *mSystems* **5**:
e00261-20.
Altamia, M.A., Shipway, J.R., Stein, D., Betcher, M.A., Fung, J.M., Jospin, G., et al. (2021)
Teredinibacter haidensis sp. nov., Teredinibacter purpureus sp. nov. and Teredinibacter
franksiae sp. nov., marine, cellulolytic endosymbiotic bacteria isolated from the gills of
the wood-boring mollusc Bankia setacea (Bivalvia: Teredinidae) and emended
description of the genus Teredinibacter. *International Journal of Systematic and
Evolutionary Microbiology* **71**:.
Altamia, M.A., Shipway, J.R., Stein, D., Betcher, M.A., Fung, J.M., Jospin, G., et al. (2020)
Teredinibacter waterburyi sp. nov., a marine, cellulolytic endosymbiotic bacterium
isolated from the gills of the wood-boring mollusc Bankia setacea (Bivalvia: Teredinidae)
and emended description of the genus Teredinibacter. *International Journal of
Systematic and Evolutionary Microbiology* **70**: 2388–2394.
Altamia, M.A., Wood, N., Fung, J.M., Dedrick, S., Linton, E.W., Concepcion, G.P., et al. (2014)
Genetic differentiation among isolates of *Teredinibacter turnerae,* a widely occurring
intracellular endosymbiont of shipworms. *Molecular Ecology* **23**: 1418–1432.
Alves, N.J., Turner, K.B., Medintz, I.L., and Walper, S.A. (2016) Protecting enzymatic function
through directed packaging into bacterial outer membrane vesicles. *Sci Rep* **6**: 24866.
Arntzen, M.Ø., Várnai, A., Mackie, R.I., Eijsink, V.G.H., and Pope, P.B. (2017) Outer membrane
vesicles from *Fibrobacter succinogenes* S85 contain an array of carbohydrate-active
enzymes with versatile polysaccharide-degrading capacity. *Environmental Microbiology*
19: 2701–2714.
Aschtgen, M.-S., Wetzel, K., Goldman, W., McFall-Ngai, M., and Ruby, E. (2016) *Vibrio fischeri* -
derived outer membrane vesicles trigger host development: OMV deliver signals in the
squid/vibrio symbiosis. *Cellular Microbiology* **18**: 488–499.
Athman, J.J., Wang, Y., McDonald, D.J., Boom, W.H., Harding, C.V., and Wearsch, P.A. (2015)
Bacterial Membrane Vesicles Mediate the Release of *Mycobacterium tuberculosis*
Lipoglycans and Lipoproteins from Infected Macrophages. *The Journal of Immunology*
195: 1044–1053.

Berleman, J. and Auer, M. (2013) The role of bacterial outer membrane vesicles for intra- and
interspecies delivery. *Environmental Microbiology* **15**: 347–354.

Betcher, M.A., Fung, J.M., Han, A.W., O'Connor, R., Seronay, R., Concepcion, G.P., et al. (2012)
Microbial Distribution and Abundance in the Digestive System of Five Shipworm Species
(Bivalvia: Teredinidae). *PLoS ONE* **7**: e45309.

Biller, S.J., Coe, A., Arellano, A.A., Dooley, K., Gong, J.S., Yeager, E.A., et al. (2022) Environmental
and taxonomic drivers of bacterial extracellular vesicle production in marine ecosystems,
Microbiology.

Biller, S.J., Schubotz, F., Roggensack, S.E., Thompson, A.W., Summons, R.E., and Chisholm, S.W.
(2014) Bacterial Vesicles in Marine Ecosystems. *Science* **343**: 183–186.

Blanvillain, S., Meyer, D., Boulanger, A., Lautier, M., Guynet, C., Denancé, N., et al. (2007) Plant
Carbohydrate Scavenging through TonB-Dependent Receptors: A Feature Shared by
Phytopathogenic and Aquatic Bacteria. *PLoS ONE* **2**: e224.

Bonnington, K.E. and Kuehn, M.J. (2014) Protein selection and export via outer membrane
vesicles. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **1843**: 1612–
1619.

Cai, Y.J., Chapman, S.J., Buswell, J.A., and Chang, S. (1999) Production and Distribution of
Endoglucanase, Cellobiohydrolase, and β-Glucosidase Components of the Cellulolytic
System of *Volvariella volvacea* , the Edible Straw Mushroom. *Appl Environ Microbiol* **65**:
553–559.

Caruana, J.C. and Walper, S.A. (2020) Bacterial Membrane Vesicles as Mediators of Microbe –
Microbe and Microbe – Host Community Interactions. *Front Microbiol* **11**: 432.

Cragg, S.M., Beckham, G.T., Bruce, N.C., Bugg, T.D., Distel, D.L., Dupree, P., et al. (2015)
Lignocellulose degradation mechanisms across the Tree of Life. *Current Opinion in
Chemical Biology* **29**: 108–119.

Cragg, S.M., Friess, D.A., Gillis, L.G., Trevathan-Tackett, S.M., Terrett, O.M., Watts, J.E.M., et al.
(2020) Vascular Plants Are Globally Significant Contributors to Marine Carbon Fluxes and
Sinks. *Annu Rev Mar Sci* **12**: 469–497.

Dahmen, N., Lewandowski, I., Zibek, S., and Weidtmann, A. (2019) Integrated lignocellulosic
value chains in a growing bioeconomy: Status quo and perspectives. *GCB Bioenergy* **11**:
107–117.

De Paula, R.G., Antoniêto, A.C.C., Nogueira, K.M.V., Ribeiro, L.F.C., Rocha, M.C., Malavazi, I., et al.
(2019) Extracellular vesicles carry cellulases in the industrial fungus Trichoderma reesei.
Biotechnol Biofuels **12**: 146.

Dhurve, G., Madikonda, A.K., Jagannadham, M.V., and Siddavattam, D. (2022) Outer Membrane
Vesicles of *Acinetobacter baumannii* DS002 Are Selectively Enriched with TonB-
Dependent Transporters and Play a Key Role in Iron Acquisition. *Microbiol Spectr* **10**:
e00293-22.

Distel, D.L., DeLong, E.F., and Waterbury, J.B. (1991) Phylogenetic characterization and in situ
localization of the bacterial symbiont of shipworms (Teredinidae: Bivalvia) by using 16S
rRNA sequence analysis and oligodeoxynucleotide probe hybridization. *Appl Environ
Microbiol* **57**: 2376–2382.

Distel, D.L., Morrill, W., MacLaren-Toussaint, N., Franks, D., and Waterbury, J. (2002)
Teredinibacter turnerae gen. nov., sp. nov., a dinitrogen-fixing, cellulolytic,

endosymbiotic gamma-proteobacterium isolated from the gills of wood-boring molluscs
(Bivalvia: Teredinidae). *International Journal of Systematic and Evolutionary*
Microbiology **52**: 2261–2269.
Fadeev, E., Carpaneto Bastos, C., Hennenfeind, J.H., Biller, S.J., Sher, D., Wietz, M., and Herndl,
G.J. (2023) Characterization of membrane vesicles in *Alteromonas macleodii* indicates
potential roles in their copiotrophic lifestyle. *microLife* **4**: uqac025.
Fischer, T., Schorb, M., Reintjes, G., Kolovou, A., Santarella-Mellwig, R., Markert, S., et al. (2019)
Biopearling of Interconnected Outer Membrane Vesicle Chains by a Marine
Flavobacterium. *Appl Environ Microbiol* **85**: e00829-19.
Frias, A., Manresa, A., De Oliveira, E., López-Iglesias, C., and Mercade, E. (2010) Membrane
Vesicles: A Common Feature in the Extracellular Matter of Cold-Adapted Antarctic
Bacteria. *Microb Ecol* **59**: 476–486.
Galperin, M.Y., Wolf, Y.I., Makarova, K.S., Vera Alvarez, R., Landsman, D., and Koonin, E.V. (2021)
COG database update: focus on microbial diversity, model organisms, and widespread
pathogens. *Nucleic Acids Research* **49**: D274–D281.
Ge, S.X., Jung, D., and Yao, R. (2020) ShinyGO: a graphical gene-set enrichment tool for animals
and plants. *Bioinformatics* **36**: 2628–2629.
Gharechahi, J., Vahidi, M.F., Sharifi, G., Ariaeenejad, S., Ding, X.-Z., Han, J.-L., and Salekdeh, G.H.
(2023) Lignocellulose degradation by rumen bacterial communities: New insights from
metagenome analyses. *Environmental Research* **229**: 115925.
Ghose, T.K. (1987) Measurement of cellulase activities. *Pure and Applied Chemistry* **59**: 257–
268.
Hervé, C., Rogowski, A., Blake, A.W., Marcus, S.E., Gilbert, H.J., and Knox, J.P. (2010)
Carbohydrate-binding modules promote the enzymatic deconstruction of intact plant
cell walls by targeting and proximity effects. *Proc Natl Acad Sci USA* **107**: 15293–15298.
Huang, Y., Nieh, M., Chen, W., and Lei, Y. (2022) Outer membrane vesicles (OMVs) enabled bio-
applications: A critical review. *Biotechnol Bioeng* **119**: 34–47.
Ichikawa, S., Ogawa, S., Nishida, A., Kobayashi, Y., Kurosawa, T., and Karita, S. (2019)
Cellulosomes localise on the surface of membrane vesicles from the cellulolytic
bacterium *Clostridium thermocellum*. *FEMS Microbiology Letters* **366**: fnz145.
Jäckle, O., Seah, B.K.B., Tietjen, M., Leisch, N., Liebeke, M., Kleiner, M., et al. (2019)
Chemosynthetic symbiont with a drastically reduced genome serves as primary energy
storage in the marine flatworm *Paracatenula*. *Proc Natl Acad Sci USA* **116**: 8505–8514.
Lau, W.Y.V., Hoad, G.R., Jin, V., Winsor, G.L., Madyan, A., Gray, K.L., et al. (2021) PSORTdb 4.0:
expanded and redesigned bacterial and archaeal protein subcellular localization
database incorporating new secondary localizations. *Nucleic Acids Research* **49**: D803–
D808.
Lynch, J.B. and Alegado, R.A. (2017) Spheres of Hope, Packets of Doom: the Good and Bad of
Outer Membrane Vesicles in Interspecies and Ecological Dynamics. *J Bacteriol* **199**:.
Manck, L.E., Espinoza, J.L., Dupont, C.L., and Barbeau, K.A. (2020) Transcriptomic Study of
Substrate-Specific Transport Mechanisms for Iron and Carbon in the Marine Copiotroph
Alteromonas macleodii. *mSystems* **5**: e00070-20.

McGavin, M., Lam, J., and Forsberg, C.W. (1990) Regulation and distribution of Fibrobacter
succinogenes subsp. succinogenes S85 endoglucanases. *Appl Environ Microbiol* **56**:
1235–1244.
Naka, H. and Haygood, M.G. (2023) The dual role of TonB genes in turnerbactin uptake and
carbohydrate utilization in the shipworm symbiont *Teredinibacter turnerae*. *Appl Environ*
Microbiol **89**: e00744-23.
Neumann, A.M., Balmonte, J.P., Berger, M., Giebel, H., Arnosti, C., Voget, S., et al. (2015)
Different utilization of alginate and other algal polysaccharides by marine *A lteromonas*
macleodii ecotypes. *Environmental Microbiology* **17**: 3857–3868.
O'Connor, R.M., Fung, J.M., Sharp, K.H., Benner, J.S., McClung, C., Cushing, S., et al. (2014) Gill
bacteria enable a novel digestive strategy in a wood-feeding mollusk. *Proc Natl Acad Sci*
USA **111**: E5096–E5104.
O'Donoghue, E.J. and Krachler, A.M. (2016) Mechanisms of outer membrane vesicle entry into
host cells: MicroReview - OMV entry into host cells. *Cellular Microbiology* **18**: 1508–
1517.
Orench-Rivera, N. and Kuehn, M.J. (2021) Differential Packaging Into Outer Membrane Vesicles
Upon Oxidative Stress Reveals a General Mechanism for Cargo Selectivity. *Front*
Microbiol **12**: 561863.
Østby, H., Hansen, L.D., Horn, S.J., Eijsink, V.G.H., and Várnai, A. (2020) Enzymatic processing of
lignocellulosic biomass: principles, recent advances and perspectives. *Journal of*
Industrial Microbiology and Biotechnology **47**: 623–657.
Park, M., Sun, Q., Liu, F., DeLisa, M.P., and Chen, W. (2014) Positional Assembly of Enzymes on
Bacterial Outer Membrane Vesicles for Cascade Reactions. *PLoS ONE* **9**: e97103.
Pesante, G., Sabbadin, F., Elias, L., Steele-King, C., Shipway, J.R., Dowle, A.A., et al. (2021)
Characterisation of the enzyme transport path between shipworms and their bacterial
symbionts. *BMC Biol* **19**: 233.
Pollet, R.M., Martin, L.M., and Koropatkin, N.M. (2021) TonB-dependent transporters in the
Bacteroidetes: Unique domain structures and potential functions. *Molecular*
Microbiology **115**: 490–501.
Ragauskas, A.J., Beckham, G.T., Biddy, M.J., Chandra, R., Chen, F., Davis, M.F., et al. (2014) Lignin
Valorization: Improving Lignin Processing in the Biorefinery. *Science* **344**: 1246843.
Rakoff-Nahoum, S., Coyne, M.J., and Comstock, L.E. (2014) An Ecological Network of
Polysaccharide Utilization among Human Intestinal Symbionts. *Current Biology* **24**: 40–
49.
Sabbadin, F., Pesante, G., Elias, L., Besser, K., Li, Y., Steele-King, C., et al. (2018) Uncovering the
molecular mechanisms of lignocellulose digestion in shipworms. *Biotechnol Biofuels* **11**:
59.
Salvachúa, D., Werner, A.Z., Pardo, I., Michalska, M., Black, B.A., Donohoe, B.S., et al. (2020)
Outer membrane vesicles catabolize lignin-derived aromatic compounds in
Pseudomonas putida KT2440. *Proc Natl Acad Sci USA* **117**: 9302–9310.
Sartorio, M.G., Pardue, E.J., Scott, N.E., and Feldman, M.F. (2023) Human gut bacteria tailor
extracellular vesicle cargo for the breakdown of diet- and host-derived glycans. *Proc Natl*
Acad Sci USA **120**: e2306314120.

Schorey, J.S., Cheng, Y., and McManus, W.R. (2021) Bacteria- and host-derived extracellular
vesicles – two sides of the same coin? *Journal of Cell Science* **134**: jcs256628.
Schwechheimer, C. and Kuehn, M.J. (2015) Outer-membrane vesicles from Gram-negative
bacteria: biogenesis and functions. *Nat Rev Microbiol* **13**: 605–619.
Silale, A. and Van Den Berg, B. (2023) TonB-Dependent Transport Across the Bacterial Outer
Membrane. *Annu Rev Microbiol* **77**: 67–88.
Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., et al. (2019) STRING
v11: protein–protein association networks with increased coverage, supporting
functional discovery in genome-wide experimental datasets. *Nucleic Acids Research* **47**:
D607–D613.
Teather, R.M. and Wood, P.J. (1982) Use of Congo red-polysaccharide interactions in
enumeration and characterization of cellulolytic bacteria from the bovine rumen. *Appl*
Environ Microbiol **43**: 777–780.
Thakur, M., Dean, S.N., Caruana, J.C., Walper, S.A., and Ellis, G.A. (2023) Bacterial Membrane
Vesicles for In Vitro Catalysis. *Bioengineering* **10**: 1099.
Toyofuku, M., Schild, S., Kaparakis-Liaskos, M., and Eberl, L. (2023) Composition and functions of
bacterial membrane vesicles. *Nat Rev Microbiol* **21**: 415–430.
Veith, P.D., Chen, Y.-Y., Chen, D., O'Brien-Simpson, N.M., Cecil, J.D., Holden, J.A., et al. (2015)
Tannerella forsythia Outer Membrane Vesicles Are Enriched with Substrates of the Type
IX Secretion System and TonB-Dependent Receptors. *J Proteome Res* **14**: 5355–5366.
Voight, J.R. (2015) Xylotrophic bivalves: aspects of their biology and the impacts of humans.
Journal of Molluscan Studies **81**: 175–186.
Waterbury, J.B., Calloway, C.B., and Turner, R.D. (1983) A Cellulolytic Nitrogen-Fixing Bacterium
Cultured from the Gland of Deshayes in Shipworms (Bivalvia: Teredinidae). *Science* **221**:
1401–1403.
Yan, S. and Wu, G. (2013) Secretory pathway of cellulase: a mini-review. *Biotechnol Biofuels* **6**:
177.
Yang, J.C., Madupu, R., Durkin, A.S., Ekborg, N.A., Pedamallu, C.S., Hostetler, J.B., et al. (2009)
The Complete Genome of Teredinibacter turnerae T7901: An Intracellular Endosymbiont
of Marine Wood-Boring Bivalves (Shipworms). *PLoS ONE* **4**: e6085.
Yu, C., Chen, Y., Lu, C., and Hwang, J. (2006) Prediction of protein subcellular localization.
Proteins **64**: 643–651.
Zakharzhevskaya, N.B., Vanyushkina, A.A., Altukhov, I.A., Shavarda, A.L., Butenko, I.O., Rakitina,
D.V., et al. (2017) Outer membrane vesicles secreted by pathogenic and nonpathogenic
Bacteroides fragilis represent different metabolic activities. *Sci Rep* **7**: 5008.
Zingl, F.G., Thapa, H.B., Scharf, M., Kohl, P., Müller, A.M., and Schild, S. (2021) Outer Membrane
Vesicles of Vibrio cholerae Protect and Deliver Active Cholera Toxin to Host Cells via
Porin-Dependent Uptake. *mBio* **12**: e00534-21.
685
686

A) Culture

30 °C

C) OMV Characterization

NTA   TEM   LC-MS/MS

D) Cellulase Activity

Zymogram

DNS Assay

B) Differential and Density Gradient Ultracentrifugation

Cells

OMV-containing secretome

Soluble protein fraction

Raw OMVs

10%
20%
30%
35%
40%
45%

Purified OMVs

A)

T7901

- Outer Membrane
- Periplasmic
- Cytoplasmic
- Extracellular
- Inner Membrane

SR01903

- Outer Membrane
- Periplasmic
- Cytoplasmic
- Extracellular
- Inner Membrane

B)

T7901

SR01903

- Not Assigned, Multiple Assigned, Unknown Function
- Cell Wall/Membrane/Envelope Biogenesis
- Inorganic Ion Transport and Metabolism
- Carbohydrate Transport and Metabolism
- Energy Production and Conversion
- Amino Acid Transport and Metabolism
- Post-Translational Modification, Protein Turnover, Chaperones
- Intracellular Trafficking, Secretion, and Vesicular Transport
- Translation, Ribosomal Structure, and Biogenesis
- Nucleotide Transport and Metabolism
- Coenzyme Transport and Metabolism
- Signal Transduction Mechanisms
- Cell Motility
- Cell Cycle Control, Cell Division, Chromosome Partitioning
- Transcription
- Secondary Metabolites Biosynthesis, Transport, and Catabolism
- Defense Mechanisms
- Replication, Recombination, and Repair

C)

TonB-dependent Receptor Plug Domain, and Tryptophan halogenase
Mixed, incl. tonb-dependent receptor plug domain, and protein transport
Mixed, incl. tonb-dependent receptor plug domain, and tryptophan halogenase
Mixed, incl. tonb-dependent receptor plug domain, and peptidase s9, prolyl oligopeptidase, catalytic domain
Bacterial TonB-dependent receptor, and Tryptophan halogenase
Hydrolase activity, hydrolyzing o-glycosyl compounds, and pentose and glucuronate interconversions
Hydrolase activity, hydrolyzing o-glycosyl compounds, and six-hairpin glycosidase superfamily
Mixed, incl. biopolymer transport protein exbd/tolr, and tetratricopeptide repeat
Mixed, incl. glycosyl hydrolases family 43, and xylan catabolic process
Mixed, incl. hydrolase activity, hydrolyzing o-glycosyl compounds, and pentose and glucuronate interconversions

N. of Genes
- 15
- 20
- 25
- 30
- 35
- 40

-log10(FDR)
- 6
- 7
- 8
- 9
- 10
- 11

Fold Enrichment

A) Glycoside Hydrolase families

T7901
SR01903

High
Low

CBM9
CBM6
CBM16
CBM50
CBM35
CBM67
CBM5/12
CBM48
CBM6
CBM81
CE1
CE2
CE4/6
GH2
GH3
GH5
GH6
GH8
GH9
GH10
GH11
GH13
GH16/7
GH26
GH27
GH28
GH39
GH43/51/55
GH44
GH48
GH51
GH67
GH94
GH105
GH115
GT2
GT4
PL31

B)

marker T7901 SR01903   T7901 SR01903

kDa

Denaturing   Zymogram
SDS-PAGE

C)

Activity (Units/mg)

Purified cellulase
enzyme cocktail

T7901   SR01903

Unlysed OMVs