Transposable elements contribute to the evolution of host shift-related genes in

cactophilic *Drosophila* species

Oliveira D. S.[1,2,3], Larue A.[2], Nunes W. V. B.[2], Sabot F.[4], Bodelón A.[5], García Guerreiro M. P.[5], Vieira C.[2,*], Carareto

C. M. A.[1,*]

[1] São Paulo State University (Unesp), Institute of Biosciences, Humanities and Exact Sciences, São José do Rio Preto, São

Paulo, 15054-000, Brazil.

[2] Universite Claude Bernard Lyon 1, Laboratoire de Biométrie et Biologie Évolutive, CNRS, UMR5558, Villeurbanne 69100,

France.

[3] Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14850

[4] DIADE, University of Montpellier, Cirad, IRD, 911 avenue Agropolis BP64501, 34394 Montpellier, France.

[5] Grup de Genòmica, Bioinformàtica i Biologia Evolutiva, Departament de Genètica i Microbiologia (Edifici C), Universitat

Autònoma de Barcelona, Barcelona, Spain

* Corresponding authors

Daniel Siqueira de Oliveira: ds2497@cornell.edu

Anaïs Larue: anais.larue@univ-lyon1.fr

William Vilas Boas Nunes: william.vilas-boas-nunes@univ-lyon1.fr

François Sabot: francois.sabot@ird.fr

Alejandra Bodelón: Alejandra.Bodelon@uab.cat

María Pilar García Guerreiro: MariaPilar.Garcia.Guerreiro@uab.cat

Cristina Vieira: cristina.vieira@univ.lyon1.fr

Claudia Marcia Aparecida Carareto: claudia.carareto@unesp.br

ABSTRACT

Host shifts in insects have been considered a key process with the potential to contribute to reproductive isolation and speciation. Both genomics and transcriptomics variation have been attributed to such a process, in which gene families with functions associated with host localization, acceptance, and usage have been proposed to evolve. In this context, cactophilic *Drosophila* species are an excellent model to study host shift evolution, since they use a wide range of cacti as hosts, and many species have different preferences. Transposable elements are engines of genetic novelty between populations and species, driving rapid adaptive evolution. However, the extent of TEs' contribution to host shift remains unexplored. We performed genomic and transcriptomic analyses in six genomes of cactophilic species/subspecies to investigate how TEs interact with genes associated with host shift. Our results revealed enrichment of TEs at promoter regions of host shift-related genes, with ~39% of the odorant receptors containing their transcription factor binding sites within TEs. We observed that ~50% of these TEs are *Helitrons*, demonstrating an unprecedented putative *cis*-regulatory role of *Helitrons* in *Drosophila*. Differential expression analysis between species with different preferred hosts revealed divergence in gene expression in head and larval tissues. Although TEs' presence does not affect overall gene expression, we observed 6.27% of the expressed genes generating gene-TE chimeric transcripts, including those with function affecting host preference. Our combined genomic and transcriptomic approaches provide evidence of TE-driven divergence between species,

44 highlighting the evolutionary role of TEs in the context of host shift, a key adaptive process

45 that can cause reproductive isolation.

46 **Keywords**: comparative transcriptomics, adaptation, chimeric transcripts.

Introduction

The molecular consequences of host shift evolution can ultimately cause specialization and ecological speciation (1). In many species, reproductive isolation arises as a result of successive adaptations driven by host shift, which results from divergent selection between different environments (1–4). Thus, natural selection increases the frequency of alleles that confer higher fitness benefits when a host shift occurs. *Drosophila* species use a wide range of necrotic host tissues as feeding and breeding sites, providing a suitable model to study the host shift process and its role in insect speciation. Among them, the *repleta* group comprises a clade with ~100 cactophilic *Drosophila* species (5). The specialization in cacti represents an important ecological challenge to the flies, mainly due to the presence of toxic compounds (6). In the *repleta* species, the host shift was initially to the *Opuntia sp.* cacti, and secondarily, several independent shifts happened to columnar cacti (6), considered a host with higher toxicity than *Opuntia sp.* (7). Such host shift is observed between sibling species with recent divergence in two clusters of the subgroup *mulleri*: cluster *buzzatii* and cluster *mojavensis*. In the cluster *buzzatii*, *Drosophila buzzatii* and *D. koepferae* are sibling species (divergence 4-5 Mya (8)) with different primary hosts. The former species preferentially use *Opuntia sp.* cacti, whereas the latter is a columnar cacti dweller (9). In the cluster *mojavensis*, *D. arizonae* and the three allopatric subspecies of *D. mojavensis* have different cacti preferences. *D. arizonae* uses both *Opuntia sp.* and columnar cacti, whereas *D. m. mojavensis* uses the columnar cacti *Ferocactus cylindraceus* as preferential host; *D. m. wrigleyi* uses only *Opuntia sp.*, *D. m. baja* uses the columnar cacti

Stenocereus gummosus, and *D. m. sonorensis* is also a columnar cacti dweller, using *S.*

thurberi (10). Therefore, this set of species from *buzzatii* and *mojavensis* clusters provides

an excellent model to investigate the molecular mechanisms underlying host shift

evolution.

The radiation of cactophilic *Drosophila* species across different hosts is an

evolutionary outcome that gave rise to specific adaptations (11). They can be summarized

into three steps: localization, acceptance, and host usage (12). In the localization step,

odorant-binding proteins (OBPs) and odorant receptors (ORs) are key proteins to

distinguish specific hosts in the environment, through the integration of the visual and

olfactory systems. Subsequently, in the acceptance step, the insect evaluates the nutritional

compounds present on the host, as well as the presence of competitors, predators,

pathogens and parasites (12). Several receptors are associated with this process, such as

gustatory receptors (GRs) (13) and ionotropic receptors (IRs) (14–16). Finally, the use of

metabolites derived from the host is essential for feeding and to complete the breeding

process. Although not always associated with detoxification (17), cactophilic flies must

overcome the presence of toxic substances from columnar cacti. Several gene families are

associated with detoxification, such as Cytochrome P450s (CYPs), Glutathione S-

Transferases (GSTs), UDP-Glycosyltransferases (UGTs), esterases (ESTs), and ATP binding-

cassette transporters (ABCs). Altogether, these nine gene families associated with the Host

Localization, Acceptance, and Usage will henceforth be referred to as HLAU.

Transposable elements (TEs) may play a role in environmental adaptation because of

their ability to generate mutations. In most cases, mutations caused by TEs are likely to be

deleterious or neutral. Throughout evolutionary time, TEs that remain in the genome tend

to be silenced by epigenetic control and/or small RNA pathways (18), accumulating

mutations, and losing their transposition ability (19). Despite the majority of TEs becoming

silenced, the remaining TE sequences may still contain regulatory motifs or protein

domains (20). These sequences can be co-opted by the cell machinery, modifying gene

expression or protein sequences of the nearby genes (reviewed in (21)). Depending on the

outcome of the cooption for the individual fitness, these TEs can increase their frequency in

the population, contributing to populational evolution and hence being determined as

adaptive insertions. For instance, in *D. melanogaster*, the gene *CHKov1* has a truncated

mRNA derived from a *DOC* TE located into the intron, resulting in resistance to viral

infection and insecticides (22). Such exaptation and domestication events can be often

identified by the occurrence of chimeric transcripts, which are mRNAs with both gene and

TE-derived sequences (23). A recent transcriptome-wide study has identified 327 genes of

D. melanogaster genes produce chimeric transcripts in different populations (24). Among all

genes, 76 generate chimeric transcripts from TE insertions that were present in one strain

but absent in another, highlighting the potential of TEs as a source of genetic novelty

between different *Drosophila* ecotypes.

Many aspects involving host shift adaptation in *repleta* species have been shown

previously, such as detoxification pathways (25), morphology (26), life history traits (27),

behavior (28), and genomic and transcriptomic differences (29–33). However, the

contribution of TEs to the evolution of HLAU genes has not yet been assessed. Here, we

aimed to uncover the extension of the genetic variability derived from TEs in cactophilic

Drosophila species, using both genome- and transcriptome-wide analysis. We tested the

hypothesis whether TEs have contributed to the evolution of HLAU genes, and

consequently to the host shift in cactophilic species.

Results

Genome assemblies and gene annotation

To investigate the potential role of TEs in host shift of cactophilic *Drosophila* species,

we performed Nanopore long-read sequencing on species that have different preferential

cacti as hosts: *D. buzzatii* (*Opuntia sp.*), *D. koepferae* (columnar cacti), *D. arizonae* (*Opuntia*

sp. or columnar cacti), *D. m. mojavensis* (*F. cylindraceus*), *D. m. wrigleyi* (*Opuntia sp.*), and *D.*

m. sonorensis (*S. thurberi*). We obtained high-resolution assemblies for all genomes, with an

average of 567 scaffolds, 13.8 Mb of N50 (Supplemental Table 1), and ~98% of BUSCO

genes (Supplemental Fig. S1). Furthermore, the gene annotation for these genomes

covered 98.63% of all reference genes in the *D. mojavensis* subspecies, and 95.98% in *D.*

arizonae. The *de novo* gene annotation in *D. buzzatii* and *D. koepferae* revealed a total of

18,050 and 17,848 coding genes, for which 63.77%, and 64.13% were successfully assigned

as one-to-one orthologs with *D. mojavensis*, respectively. These results represent the higher

gene repertoire identified for *D. buzzatii* and *D. koepferae* species, altogether with their

ortholog relationship with *D. mojavensis* and *D. arizonae*. Since analysis of HLAU genes is the focus of this study, their efficient annotation is relevant. They are represented by nine gene families associated with the three steps in the use of host resources: (1) Localization: OBPs, ORs; (2) Acceptance: GRs, IRs; and (3) Host usage: ABCs, ESTs, GSTs, UDPs, CYPs (Supplemental Table 2). In *D. arizonae* and *D. mojavensis* subspecies, we recovered ~98.21% of the total HLAU genes from the reference genome. In *D. buzzatii* and *D. koepferae*, the number of annotated HLAU genes covered ~95% in both species compared to *D. mojavensis* (Table 1).

Species/subspecies	Total genes	Host localization, acceptance and usage genes								
		Localization		*Acceptance*		*Host usage*				
		OBP	OR	GR	IR	ABC	EST	GST	UDP	CYP
Ref. *D. m. wrigleyi*	**15,045**	**25**	**59**	**45**	**22**	**27**	**17**	**28**	**21**	**59**
D. m. mojavensis	14,810	25	59	44	22	27	17	27	20	59
D. m. wrigleyi	14,948	25	59	45	22	27	17	27	20	59
D. m. sonorensis	14,808	25	57	44	22	27	17	27	20	58
D. arizonae	14,441	25	56	41	21	27	17	27	20	59
D. buzzatii	18,050	25	52	34	21	27	16	27	20	68
D. koepferae	17,848	25	45	34	24	27	13	25	23	62

Table 1. Total number of genes annotated in six *Drosophila* cactophilic species. The "Ref. *D. m. wrigleyi*" represents the annotation in the reference genome used to perform the gene annotation for our long-read Nanopore genomes. HLAU genes are separated according to their functions: Localization: odorant-binding proteins (OBP), odorant receptor (OR); Acceptance: gustatory receptor (GR), ionotropic receptor (IR); Host usage: ATP-binding cassette (ABC), carboxylesterase (EST), glutathione S-Transferase (GST), glycosyltransferase (UDP), and cytochrome P450 (CYP).

Minor effects of positive selection in HLAU genes

The use of *Opuntia sp.* has been proposed as the ancestral state in cactophilic

Drosophila species, whereas columnar cacti are the derived state (6). To investigate whether

species that prefer columnar cacti as hosts exhibit signs of positive selection in HLAU

genes, we employed a branch-site approach to compare their evolutionary rates with those

of species that use *Opuntia sp.* For this analysis, we removed *D. arizonae* due to its non-

preferential usage between columnar and *Opuntia sp.* cacti (6). Moreover, we added *D.*

navojoa, which is a basal species of the *mojavensis* cluster that uses solely *Opuntia* sp. (34).

Therefore, we carried out the branch-site selection test with *D. buzzatii*, *D. m. wrigleyi*, and

D. navojoa as *Opuntia*-related branches (background), and *D. m. mojavensis*, *D. sonorensis*,

and *D. koepferae* as columnar-related branches (foreground). It is important to highlight

that due to the requirement for one-to-one orthologous genes between the six genomes,

our analysis has reduced the total number of HLAU genes to 42.9% of the initial number

presented in Table 1. From 130 genes, we observed six genes under positive selection (FDR

< 0.05): *Obp56d*, *Obp19d*, *Gr5a*, *Ir25a*, *Cyp4s3*, and *esterase S*. This result suggests that

positive selection in HLAU genes is not a major evolutionary driving force in the species

under study.

Evolutionary dynamics of TEs in cactophilic Drosophila species

We used a combination of automatic and manual methods to build a curated TE

library for each species, based on the classification of TEs (35). The method was designed to

perform polishing steps in both TE consensus and copies to remove artifacts (see Methods).

The total number of TE consensuses (LTRs separated from internal sequences) ranged from

255 in *D. buzzatii* to 372 in *D. m. mojavensis*. The TE content in the genomes ranged from

15.04% in *D. buzzatii* to 22.01% in *D. m. mojavensis* (Fig. 1A). In addition, we observed a

positive correlation between TE content and genome size estimated from the assemblies

(*Pearson* correlation: p = 0.0082; R = 0.93) (Supplemental Fig. S2). Both species from the

buzzatii cluster had the lower genome sizes and TE contents compared to species of the

mojavensis cluster (Fig. 1A). Regarding the proportional load of TE orders in the genomes,

LTRs and Helitrons contribute on average with ~70% of total TE content in all species (Fig.

1B).

Figure 1. A) Genome size of the six cactophilic *Drosophila* species sequenced in this work, and their respective TE content. **B**) Proportional distribution of the total TE content in the genomes, in crescent order. **C**) Phylogenetic relationship among species (adapted from (25,37)), alongside TE order landscapes representing total copy number (y-axis), and the divergence of TE insertions compared to the consensuses (x-axis), based on Kimura 2-parameters (K2P) corrected with CpG. The copies located on the left of the graph have low divergence with their respective consensus, hence inferring that they might be

conserved/recent/active, whereas copies on the right side represent copies with high divergence due to the

accumulation of mutations, corresponding to old/inactive copies.

In the *mojavensis* cluster (*D. arizonae* and the three *D. mojavensis* subspecies), the

abundance of TEs in the genome varied substantially (Fig. 1B), despite their recent

divergence time of 1.5 Mya (36). *D. m. mojavensis* has the TE-richest genome (22.01%), with

the abundance following from the highest to the lowest: LTR > Helitron > LINE > DNA (Fig.

1B). The distribution of TEs in the other two *D. mojavensis* subspecies and *D. buzzatii* does

not follow such order. LTRs and Helitrons have the highest abundance, followed by DNA

and LINEs (Fig. 1B). In *D. arizonae* and *D. koepferae*, Helitrons are the most abundant,

following LTR, DNA, and LINEs (Fig. 1B). The observed differences are often associated with

the divergence in the evolutionary patterns of the TE orders between the genomes. We

used copy number as a proxy to measure TE expansion in the genomes, and divergence

Kimura 2-parameters (K2P) as age measurement. In the *mojavensis* cluster, we observed

that LTRs and Helitrons had a higher expansion in *D. m. mojavensis* compared to the other

genomes (Fig. 1C). The higher copy number in *D. m. mojavensis* is mostly from young TEs

(K2P 0-5), which might explain the higher TE content in this species. DNA elements have a

constant evolutionary dynamics in the genomes, except in *D. m. wrigleyi*. In this species, we

observed a peak in copy number with K2P between 16 and 20, considered old insertions.

LINEs had a similar evolutionary landscape in *D. mojavensis* subspecies, with higher copy

number from young copies compared to *D. arizonae*. In the *buzzatii* cluster, Helitrons show

higher copy number from young insertions in *D. koepferae* than in *D. buzzatii* (Fig. 1C),

whereas LINEs show a past transposition burst in *D. buzzatii* compared to *D. koepferae*. The

antagonistic patterns of TE dynamics observed in the six genomes suggest that TEs had

different successful lineages across these species.

HLAU genes are enriched in TEs within regulatory regions

217   TE insertions may be selected in the promoter of genes if they provide a benefit to

the individual fitness, through the spreading of histone marks, or *cis*-regulatory elements

(38,39). To investigate whether TEs are enriched in the promoters (2 Kb upstream of the

TSS) from the nine HLAU families, we performed permutation tests in the six genomes (see

Methods). To compare our findings with other non-host shift related genes, we also

included zinc-finger and serine/threonine kinases as two non-HLAU gene families. We

observed that ORs have enrichment of TEs at the promoters in all genomes from species of

the *mojavensis* cluster (p-value < 0.05), (Fig. 2A). For acceptance genes, *D. koepferae* had

enrichment of TEs in the promoters of GRs (Fig. 2A). Finally, the enrichment of TEs within

promoters of detoxification genes was observed in *D. arizonae* (CYPs), *D. buzzatii* (GSTs),

and *D. koepferae* (ABCs). The results depicting the histograms with frequency of genes with

TEs for each significant permutation test, and p-values are shown in the Supplemental Fig.

S3 and Supplemental Table 3. Overall, detoxification genes had z-scores towards depletion

of TEs on promoters, with significant depletion for UGTs in *D. buzzatii* (p-value = 0.033) and

D. arizonae (p-value = 0.0135), GSTs in *D. koepferae* (p-value = 0.0205) (Fig. 2A). In the non-

HLAU gene families, we observed the expected result by chance: either z-score indicated

the expected number of genes with TEs on promoters in a genome-wide comparison (z-

score near 0), or it indicated depletion of TEs, as observed for SER genes in *D. buzzatii*. This

result reinforces our findings, indicating the significant and evolutionary unexpected

enrichment of TEs in HLAU genes, especially on the promoters of ORs.

Figure 2. A) Enrichment of TEs in the promoter region (2kb upstream) of HLAU gene families. Z-score > 0

indicates enrichment, and Z-score < 0 indicates depletion. Red dots: p-value < 0.05, grey dots: p-value > 0.05.

B) The same enrichment analysis as depicted on A, but for non-HLAU gene families, represented by zinc-

finger proteins (ZF) and Serine/Threonine Kinases (SER). **C**) Proportion of TE orders located upstream and

downstream ORs. **D**) Transcription factor binding sites (TFBSs) frequency on TEs located upstream of OR

genes in all species under study. **E**) Proportional frequency of TE families with TFBSs embedded on their

sequences from TE copies located upstream of OR genes. **F**) Significant Fisher's exact test between OR genes

with OR-related TFBSs from TE insertions compared to the frequency observed in the non-HLAU gene families

Serine/Threonine Kinases (SER) and zinc-finger proteins (ZF). **G**) Proportion of TE orders carrying OR-related

TFBSs in non-HLAU gene families.

To obtain further insights into the potential regulatory role of TEs, we also analyzed

the enrichment at the 2 kb downstream region. Among all species, the only significant

enrichments were observed in ORs from species of the *mojavensis* cluster (Supplemental

Table 3), as we observed in the promoters. The TE enrichment upstream and downstream

of ORs indicates a possible coevolution of this gene family with TEs. Alternatively, it may

also suggest that ORs are located in TE-rich regions of the genomes. To investigate this

hypothesis, we identified TE-rich regions (Supplemental Fig. S4) and analyzed the frequency

of ORs. We observed a single OR in TE-rich regions in *D. arizonae*, *D. m. mojavensis*, and *D.*

m. wrigleyi. (Supplemental Table 4). Therefore, we rejected the hypothesis that ORs are

enriched with TEs in their vicinity due to their location in TE-rich regions. Other factors

might be involved in the evolutionary maintenance of TEs near ORs, such as the co-option

of regulatory motifs.

A few TE families are more prone to contribute to *cis*-elements than others, such as

retroelements in *D. melanogaster* (40). Therefore, we investigated whether the observed

enrichment of TEs in the HLAU gene families is TE-specific. Considering all species, 45.36%

and 50.36% of the insertions located upstream and downstream of ORs correspond to

Helitrons, respectively (Fig. 2C). We decided to analyze whether these *Helitron* insertions

upstream to HLAU genes, mainly ORs, could be part of the first exons from genes, since

such overlap has been observed previously in other species than *Drosophila* (41,42). By

intersecting exons from genes enriched with their respective upstream *Helitrons*, we

observed that *Or85c* (overlap 772 nt) and *Or22c* (52 nt) had overlap with *Helitrons* in *D. m.*

mojavensis, whereas *D. m. sonorensis* had only *Or22c* (58 nt). While the insertion in *Or85c*

suggests a subspecies-specific insertion, the insertion in *Or22c* is likely to be inherited from

the species' common ancestor (Supplemental Table 5). At the ORs' downstream region, we

observed four genes with *Helitrons* overlapping exons: *Or71a*, *Or10a*, *Or13a*, and *Or82a*.

Interestingly, the same *Helitron* in the *Or82a* was observed in the three *D. mojavensis*

subspecies, highlighting the inheritance from the common ancestor and maintenance of

the insertion over time. Taken together, our results rejected the hypothesis regarding the

possible role of *Helitrons* carrying HLAU gene fragments. However, these elements could

still act on gene regulation if they carry *cis*-elements as TFBSs.

TEs located near genes can modulate gene expression by donating TFBSs or polyA

signals to the ancestral regulatory motifs. To investigate the potential functional role of the

observed enrichment of TEs in ORs, we performed TFBS identification on all TE insertions

located in OR promoters. We used the transcription factors (TFs) previously described to

participate in OR's transcription in *D. melanogaster*: acj6, onecut, xbp1, fer1, and zf30C (43).

Taking all species together, the most frequent TFBS found was for the TF onecut (46.15%),

followed by zf30C (33.84%), and Xbp1 (7.69%) (Fig. 2D). *Helitrons* were the TEs with the

highest frequency, representing 50% of all insertions carrying ORs' TFBSs (Fig. 2E). Our

analysis revealed that 39.28% of the ORs with TEs in promoters have the TFBS located

within the insertion (Supplemental Table 6). To test whether these results are specific to

ORs, we also performed the same detection of TFBSs and calculation of TEs frequency in

the non-HLAU families ZF and SER. Our results show that ZF and SER have, on average,

15.55% and 17.14% of the genes with OR-related TFBSs, respectively. The detection of

these TFBSs in ZF and SER is likely due to chance, given the short length of the motifs.

Comparing these proportions with the ones found in ORs, we observed significantly more

TFBSs compared to ZF (*Fisher's exact test: p-value = 0.0004341*) and SER (*Fisher's exact test:*

p-value = 0.0008519) (Fig. 2F and Supplemental Table 7). In addition, *Helitrons* also had

higher prevalence than other TEs upstream of SER and ZF genes (Fig. 2G), demonstrating a

potential insertion preference for promoter regions. Although similar *Helitron* frequencies

between OR, SER, and ZF genes, our data suggest that *Helitrons* might have been retained

in OR promoters due to the presence of functional TFBSs (Fig. 2F).

HLAU genes are differentially expressed between *Opuntia sp.* and columnar cacti

dwellers

Differential expression of genes has been associated with adaptation to new hosts in

insects (32). Here, aiming to test whether gene expression divergence is associated with

adaptation to different cactus hosts, we performed differential expression analysis between

species with *Opuntia sp.* preference vs columnar cacti preference, in head and larval tissues.

Since our fly stocks were reared in the lab with corn media, the observed transcriptome

response is constitutive rather than triggered by the native cacti. Overall, the principal

component analysis from HLAU gene expression (normalized read counts) revealed

remarkable differences between species in both tissues (Fig. 3A). The variance of gene

expression resembles the phylogenetic divergence of the species, with the highest variance

between the *buzzatii* and *mojavensis* clusters (PC1), followed by divergence between *D.*

buzzatii and *D. koepferae* (PC2). The species of the *mojavensis* cluster were clustered

together with a slight separation between *D. arizonae* and the subspecies of *D. mojavensis*.

Figure 3: **A**) Principal component analysis of HLAU genes' expression in head and larval tissues. Both tissues

had HLAU genes' variance representing the phylogenetic distance between species: *D. buzzatii* and *D.*

koepferae, compared to the *mojavensis* cluster corresponds to the PC1 (45%), and 26% between them in PC2.

Species from the *mojavensis* cluster had strong similarity, with *D. arizonae* slightly separated from *D.*

mojavensis. **B**) HLAU genes' expression in the *Opuntia sp.* dweller *D. m. wrigleyi* versus its sister species *D.*

arizonae, which does not have preference between *Opuntia sp.* and columnar cacti. **C**) HLAU expression in

head: differential expression analysis of HLAU genes from head in the *Opuntia sp.* dweller *D. m. wrigleyi*

versus the three columnar dwellers *D. m. mojavensis,* and *D. buzzatii* versus *D. koepferae.* In A and B: Filled

dots represent HLAU genes with log2 fold-change > |1| and FDR < 0.05; transparent dots are non-significant

differential expression. **D**) HLAU expression in larvae: The same pairwise species as in A with *Opuntia sp.*

dweller vs columnar dwellers, but from larvae tissue.

We performed pairwise analysis of HLAU gene expression between species with

Opuntia sp. preference (*D. m. wrigleyi* and *D. buzzatii*) versus species with preference for

columnar cacti, and *D. arizonae* as non-preferential cacti. All differentially expressed HLAU

genes, as well as the presence of TEs in their promoters, can be accessed in (Supplemental

Table 8). In head, *D. m. wrigleyi* had 27 and 19 HLAU genes with higher expression

compared to *D. m. mojavensis* and *D. m. sonorensis,* respectively (Fig. 3A). Four genes were

found with higher expression in the head of *D. m. wrigleyi* compared to the other two

subspecies: *UDP-Ugt4, Gst-theta3, Abc-a3,* and *Cyp6a21.* In the subspecies with columnar

cacti preference, we observed, on average, 26 HLAU genes with higher expression

compared to *D. m. wrigleyi.* Among them, seven were recurrent in the two subspecies:

Ir21a, EstB1, UDP2, UDP-Ugt5, Cyp9b2, UDP-Ugt5-1, and *Cyp12a2.* Since they were

observed in the two subspecies with columnar cacti preference, we propose that they are

candidates to be related to the adaptation to columnar cacti. Finally, the same analysis on

larvae revealed that *D. m. wrigleyi* has three genes with recurrent higher expression compared to the other subspecies: *Or82a*, *Obp59a*, and *Ir21a*; whereas in the other two *D. mojavensis* subspecies we observed 17 common up-regulated genes compared to *D. m. wrigleyi*, which comprise seven CYP genes, three GSTs, four UDPs, one Esterase and one OBP (Fig. 3B). Regarding *D. buzzatii* and *D. koepferae*, we observed 27 HLAU genes in head and 54 in larvae with higher expression in *D. buzzatii*, and 63 in head and 29 with higher expression in *D. koepferae*. We also compared the head expression of HLAU genes between *D. m. wrigleyi* and its sister species without cacti host preference - *D. arizonae*. In *D. m. wrigleyi*, we found 31 HLAU genes with higher expression compared to its counterpart *D. arizonae*. In larvae, *D. m. wrigleyi* had 43 HLAU genes with higher expression compared to *D. arizonae* (Fig. 3D). This result demonstrates that HLAU genes have divergent differential expression in both head and larval tissues.

To further investigate whether differential expression of HLAU genes has a higher prevalence compared to non-HLAU genes, we compared the proportion of differentially expressed genes associated with localization, acceptance, and usage with the proportion of differentially expressed genes from the non-HLAU gene families Ser/Thr Kinases and Zinc-finger. In head, our analysis revealed that localization and usage have higher differential expression than non-HLAU genes between *D. buzzatii* and *D. koepferae* (Supplemental Fig. S5). In *D. mojavensis* subspecies, we observed higher differential expression for localization genes between *D. m. wrigleyi* and *D. m. mojavensis*. In larvae, we did not observe any significant difference between non-HLAU and HLAU genes, but the pairwise differential

expression between *D. m. wrigleyi* and two subspecies revealed a higher prevalence of

gene families associated with usage (detoxification) compared to non-HLAU (Supplemental

Fig. S5). In *D. m. wrigleyi*, we also observed higher differential expression of localization

genes than in non-HLAU. Finally, the differential expression between *D. m. wrigleyi* and *D.*

arizonae demonstrated significantly higher proportion of genes associated with localization

and usage compared to non-HLAU in larvae, and higher prevalence of differentially

expressed usage genes in head (Supplemental Fig. S5). Taken together, the quantitative

results suggest that adaptations that have allowed the shift from *Opuntia sp* to columnar

cacti may arise from the evolution of gene expression.

The enrichment of TEs on HLAU genes is not associated with changes in gene

expression

Since we observed enrichment of TEs on the regulatory regions of a few host shift-

related families, mainly ORs, and we also found differentially expressed HLAU genes

between species, we tested whether the presence of TE insertions is associated with

differential expression. To do so, we selected expressed HLAU genes (see Methods) and

performed X^2 independence test, splitting them into two variables with two categories

each: 1) expression level: differentially expressed, and non-differentially; 2) TE presence:

with TEs at 2kb upstream, and without TE at 2kb upstream. Considering all pairwise analysis

(Fig. 3A,B-D), the X^2 results demonstrated that the differential expression of HLAU genes is

not associated with the presence of TEs on the promoter region (Supplemental Table 9).

Since we did not observe overall effects of TEs on the expression of HLAU genes together,

we investigated the association of TEs with differential expression only for HLAU gene

families that we found TE enrichment (Fig. 2A). None of the enriched gene families had a

significant association between the presence of TEs and their expression level

(Supplemental Table 9). Thus, a global enrichment of TEs in the regulatory region is not

likely to be associated with the differential expression of HLAU genes.

Make or break: gene-TE chimeras contribute either predominantly or negligibly to

gene expression

Although highly deleterious, a small fraction of TEs generating chimeric transcripts

can be positively selected, giving rise to domestication and exaptation events (44). Aiming

to identify the extent of the transcriptome variability generated by gene-TE chimeric

transcripts, we analyzed the transcriptome of head and larval tissues in the six cactophilic

Drosophila species. The method employed in this work considered paired-end reads

spanning from TEs to exons. Although systematically found in all replicates, many chimeras

are likely to be products of pervasive transcription, generating ectopic transcripts whose

expression levels are insufficient to play a functional role. Therefore, we developed an

automatic method to assemble chimeric transcript isoforms and compute their expression

contribution relative to the total gene expression. This method is implemented in the

pipeline *ChimeraTE v2.0* (24) as an optional downstream analysis to the total chimeras

detection.

Our results revealed that, on average, 715 and 574 genes produce chimeric

transcripts across all species, corresponding to 7.51% and 5.04% of the expressed genes in

head and larvae, respectively. In both tissues, TE-exonized transcripts represented the

highest frequency, with 87.59% in head and 89.09% in larvae (Fig. 4A), followed by TE-

terminated transcripts with 10.63% and 9.45%, and TE-initiated transcripts with 1.79% and

1.51%. Interestingly, all species have a similar number of expressed genes (head: $\bar{x}$ = 9,508;

σ = 300; larvae: $\bar{x}$ = 11,375; σ = 117), but we observed on average 11-fold fewer chimeric

transcripts in *D. buzzatii* and *D. koepferae* compared to *D. arizonae* and *D. mojavensis*

subspecies. This result suggests that species from the *mojavensis* cluster are more prone to

generate chimeras, or that *D. buzzatii* and *D. koepferae* have a strong purifying selection

against gene-TE chimeric transcripts.

Figure 4: **A**) Proportion of each chimeric transcript category based on the position of TE insertions relative to the gene structure. **B**) Histograms with contribution of chimeric transcript isoform relative to total gene expression (x-axis). Chimeric transcripts without assembled sequence or contribution < 25% were removed (grey bars). **C**) Number of common and unique chimeric transcripts from HLAU genes detected in head and larvae. **D**) *Eato/LOC6578120* (ABC transporter) gene has a conserved TE-exonized transcript with a LINE/R1 element located in its 7th intron. The depth in exons, including the 7th intron, was assessed with RNA-seq from head tissue.

In head and larvae tissues, ~49% of the detected chimeric transcript isoforms were not assembled or had contribution to gene expression lower than 25% (Fig. 4B). Among these discarded chimeras, the vast majority was not detected by transcriptome assembly

approach (0% contribution) (Fig. 4B), which is likely to be associated with negligible

expression. Considering chimeric transcripts with contribution >= 25%, more than 95%

have contribution equal to 100% (Fig. 4B). This result indicates that almost all genes

producing chimeras are dependent on the TE sequence to constitute their complete mRNA

sequence. Since ~88% of all chimeras detected with > 25% contribution are from TE-

exonized transcripts, for which 72.35% are from TEs embedded in exons, a high frequency

of chimeras with total gene expression could be expected. Notably, this pattern is also

observed in TE-initiated and TE-terminated transcripts (Supplemental Fig. S6), despite the

TEs being located upstream and downstream of the gene body. Our results reveal a clear

binomial trend: chimeric transcripts either have low expression and are unlikely to be

biologically relevant, or they account for the total gene expression, suggesting a functional

role.

HLAU genes generate chimeric transcripts

To investigate cases where TEs produce chimeric transcripts from HLAU genes, and

their possible role in the evolution of these genes, we identified all chimeras derived from

HLAU genes in head and larvae transcriptomes. We found a total of 28 (head) and 24

(larvae) HLAU genes producing chimeric transcripts. None of them are the six genes under

positive selection. Notably, ABCs, GSTs, and IRs were the most frequent gene families with

chimeric transcripts (Supplemental Table 10). Although _D. buzzatii_ had 11-fold fewer

chimeras than species from the _mojavensis_ cluster, we observed 29 gene-TE (a few genes

with more than one chimera) chimeras derived from HLAU genes in both tissues, the

highest number among all species. Its sister species *D. koepferae* had only four chimeras,

which are the same two ABC genes in head and larvae (Supplemental Table 10). In *D.*

buzzatii, ~93% of all chimeras are from GSTs, expressed in both larvae and head tissue.

Interestingly, all of them represent cases of *Helitron* exonizations (Supplemental Table 10).

We did not find any GST with chimeric transcripts in the other transcriptomes, revealing an

unprecedented recent evolutionary novelty in GSTs' evolution in *D. buzzatii*.

Chimeric transcripts from HLAU genes are represented by 14.81% TE-terminated

transcripts and 85.19% TE-exonized transcripts. We did not observe any TE-initiated

transcript from HLAU genes. Despite the compelling evidence of TEs enrichment in the

promoter of OR genes from species of the *mojavensis* cluster, we did not observe chimeric

transcripts from ORs at high frequency. Only *Or45b* (*LOC6578440*), *Or13a* (*LOC26527676*),

and *Or85a* (*LOC26528431*) had chimeric transcripts in *D. m. sonorensis*, *D. m. wrigleyi*, and

D. arizonae, respectively (Supplemental Table 10). This result reveals that TEs located in the

promoter of ORs are not likely to be co-opted as alternative transcription start sites. In the

species of the *mojavensis* cluster, we observed ABCs as the gene family with the highest

frequency of chimeras (Supplemental Table 10), despite not having enrichment (Fig. 2A).

We observed, on average, 5 ABC genes with exonization of multiple TE copies in this group

of species. This result indicates that the genomic distribution of TEs is not necessarily

associated with their transcriptional interaction with genes.

The quantitative analysis on chimeric transcripts revealed transcriptional

polymorphism in HLAU genes due to TEs. We investigated common chimeras present in

more than one species, highlighting their retention over evolutionary time, as well as

species-specific chimeras that may be associated with recent insertions. Our results

revealed that 94.44% and 88.57% of the chimeric transcripts derived from HLAU genes are

species-specific in head and larvae tissues, respectively (Fig. 4C). In head, the *Opuntia sp.*

dwellers *D. buzzatii* and *D. m. wrigleyi* were the ones with the highest number of HLAU

genes with species-specific chimeric transcripts (Fig. 4C). The 15 chimeras uniquely found in

D. buzzatii are derived from 12 genes (2 ABCs and 10 GSTs); whereas the 14 chimeras from

D. m. wrigleyi were observed from 10 genes (5 ABCs, 3 IRs, 1 CYP, and 1 OBP)

(Supplemental Table 10). In larvae, *D. buzzatii* follows a similar result as observed in head,

15 chimeras derived from 14 genes (2 ABCs and 12 GSTs), while *D. m. wrigleyi* had only four

unique chimeras (2 ORs, 1 IR, and 1 CYP).

Cases of common chimeric transcripts between populations/species may indicate

exaptation/domestication events. In the six transcriptomes analyzed here, we found 8

genes in the head and four in larvae present in more than one genome (Supplemental

Table 11). These common chimeric transcripts occurred only between *D. mojavensis*

subspecies, revealing that TEs interacting with HLAU genes are likely to be lost over higher

evolutionary scales. Their inclusion into the mRNA sequence and high contribution can be

observed. For instance, the ABC transporter (*Eato* gene) has an exonized LINE/R1, which is

located in the 7th intron (Fig. 4D), with an average contribution of 84.25% to the gene

expression. Taken together, these results demonstrate unprecedented transcriptional

interaction between TEs and HLAU genes in a tissue-specific manner.

Discussion

Insect host shift and its role in speciation have been the subject of research to

understand the evolution of reproductive isolating barriers and adaptation to local

environments. Indeed, many insects have finely tuned adaptations to locate, feed,

reproduce, and develop on the host tissue. Adaptations to new hosts can lead to

reproductive isolation, wherein ecological speciation arises as a consequence within

divergent natural populations inhabiting distinct environments. In this context, we

investigated whether TEs could play a role in contributing to the evolution of HLAU genes,

with the potential to facilitate rapid adaptive radiation to new hosts. To test this hypothesis,

we selected six *Drosophila* species/subspecies from the *repleta* group with different primary

cacti hosts.

Weak signal of positive selection in HLAU genes through branch-site model

The necrotic tissue of each cactus-host used by cactophilic *Drosophila* species differs

in its composition in terms of yeasts and alkaloids (45,46). Therefore, species have adapted

to feed and breed on different hosts, as demonstrated by behavioral and physiological

response when species feed and breed in non-preferential hosts. Positive selection in HLAU

genes have been proposed to allow the recognition of a broader spectrum of odors,

metabolize nutrients, and activate detoxification pathways (12). Indeed, specialist species

are likely to lose some of their genes related to sensory pathways, while the genes that are

retained are subject to positive selection (47,48). A preliminary genome-wide analysis of

signatures of positive selection in the four *D. mojavensis* subspecies revealed high ω rates

on genes associated with chemosensation, perception, immunity, behavior, detoxification,

and reproduction (49). Although a previous genome-wide analysis has identified 1,294

genes under positive selection (50), here we focused the analysis to investigate specifically

HLAU gene families. Our results with brach-site model had only six HLAU genes under

positive selection: *Obp19d, Obp56d, Gr5a, Ir25a, Cyp4s3, esterase S*. The unprecedent

method used in this work focused on the common differences of species adapted to

columnar cacti, rather than independent evolutionary events, making it more stringent. The

lack of genes with statistical support for positive selection may be also associated with

technical limitations, since we based our analysis on 1:1 orthologs, and a certain proportion

of the genes evolve too fast to be correctly assigned during orthologs identification (51).

Additionally, we observed ω values higher than 1.5, for instance as *Gr59d* with ω = 3.49.

Although we did not observe signficiant positive selection, this result indicates ongoing

divergence between species/subspecies.

OR promoters are enriched with TE insertions

The literature has recurrently demonstrated that TEs can create new regulatory

elements, including gene promoters (52–54), enhancers (55,56), and insulators (57,58). In

mammals, TE-derived sequences contribute up to 40% of genome-wide binding sites for

transcription factors located in TE-rich regions (59). It has been proposed that such co-

option of TE-regulatory elements for gene regulatory networks can provide substantial

modification of gene expression over short time scales, contributing to genetic variability at

the transcriptome level (60–62). The colonization of new habitats and hosts is accompanied

by behavioral changes, mainly through alterations in environmental perception (63,64).

Here, we observed that ORs are enriched by TEs in the four species from the *mojavensis*

cluster. In ants, ORs have also been found enriched in TEs. But in ants, TEs are associated

with tandem duplication by crossing over, allowing massive duplication events (65). Since

there is no expansion of ORs in cactophilic species, the enrichment in TEs is likely to have

another function, probably related to gene regulation due to its specific location in

promoters. Although we did not observe a significant enrichment of TEs in differentially

expressed genes, we argue that TEs might contribute to the regulation of specific HLAU

genes. In ORs, ~39% of the genes have OR's TFBSs located within TEs. We propose that this

subset of genes is not sufficient to have a significant signal when tested with ORs without

TEs, but they may have functional TE-derived *cis*-regulatory elements.

Helitrons have been extensively reported on the promoter of other eukaryotic

species (41,66–68), in which a few cases of embedded TFBSs were reported (69,70). Here,

we demonstrated that ~50% of the enrichment in TEs observed at the promoter region of

ORs might be due to the exaptation of TFBSs derived from *Helitrons*. In addition, the

recurrence of the enrichment in species of the *mojavensis* cluster might demonstrate

potential cases of TE co-option inherited by the common ancestor. Our genomic analysis

provides a robust list of potential candidate genes in which *Helitrons* might play a

regulatory role. This is an interesting hypothesis that can be determined through cutting-

edge technologies such as CRISPR methods to delete insertions with site-specific resolution

(71). Our results suggest a putative role of TEs in the regulation of HLAU genes across

multiple species/subspecies.

Head transcriptome reveals subspecies-specific patterns

The differential expression of HLAU genes in the head tissue revealed that

localization and host usage (detoxification) gene families have the highest divergence

across the subspecies, since we observed more differentially expressed genes for these

families. In *D. m. wrigleyi*, which is the only *D. mojavensis* subspecies with a preference for

Opuntia sp., we observed that CYP genes had on average 12 up-regulated genes compared

to other species. This result highlights that the divergence of the cacti preference between

D. mojavensis subspecies might be explained by modifications in gene expression of HLAU

genes in the head tissue. Although *Opuntia sp.* has lower toxicity than columnar cacti (72),

we found a similar number of up-regulated detoxification genes between *D. m. wrigleyi*

and the other subspecies. This suggests that *D. m. wrigleyi* has a constitutive detoxification

response that is independent of host toxicity. This is a plausible explanation since *Opuntia*

sp. produces fewer benzaldehyde volatiles than the columnar cacti *S. thurberi*, but both

cacti are lethal to *D. melanogaster* after 48 hours (72). Therefore, the adaptation to use

Opuntia sp. must have a constitutive gene expression of detoxification pathways in *D. m.*

wrigleyi.

Comparative transcriptomic studies have revealed divergence in gene expression

between *D. buzzatii* and *D. koepferae*, particularly in head tissues associated with sensory

perception and detoxification. For instance, Guillén et al. (2014) reported significant

differences in the expression of genes involved in olfaction and xenobiotic metabolism,

suggesting that these species have adapted to their respective cactus hosts (*Opuntia* for *D.*

buzzatii and columnar cacti for *D. koepferae*). Similarly, divergence in expression patterns

has been reported in these species in ORs, OBPs, and cytochrome P450 genes (73). These

reported differences are consistent with our results in head and larval tissue, reinforcing the

hypothesis that ecological specialization and divergence in head transcriptomes play a role

in host shift between *D. buzzatii* and *D. koepferae*.

Previous studies with *D. melanogaster* have shown that sequence divergence in ORs

and OBPs significantly modulates odor sensitivity (64,74), potentially leading to divergence

of host location. In addition, many ORs are involved in a broad range of ecological

interactions, influencing behaviors associated with oviposition and feeding (75). For

instance, OR evolution has mediated the herbivorous nature of leaf-mining specialist

Scaptomyza flava (76). Furthermore, transcriptomic analyses have identified 18 ORs that are

differentially expressed between heads of *D. m. wrigleyi* and *D. m. mojavensis*,

demonstrating that modifications in the sensory transcriptome have evolved between these

species (77). In this study, we expanded the understanding of this divergence by comparing

species with a preference for *Opuntia sp.* and columnar cacti, and showed that each species

has a distinct set of differentially expressed HLAU genes.

Gene expression in larvae is one of the factors shaping the adaptation to new hosts

586   Differences in larval behavior and survival have been observed when *Opuntia*-related

species are reared in columnar cacti (78,79). A previous study showed that *D. m. mojavensis*,

which primarily uses *S. thurberi* as its host, has a 60% reduction in larval viability when

reared on *S. gummosus*-based media (80). The authors also proposed that 21% of the

genes are differentially expressed between larvae reared in the primary and alternative cacti

hosts. Many genes were reported with detoxification, energy production, among others.

Another similar study with larvae transcriptome, but with *D. buzzatii* and *D. koepferae*,

proposed that *D. buzzatii* flies reared in columnar cacti media have a stronger

detoxification response in comparison with *D. koepferae* flies, suggesting that *D. koepferae*

has a canalized transcriptome response to the toxic alkaloids from columnar cacti (81).

596   Here, we observed that the majority of differentially expressed HLAU genes are from

gene families associated with detoxification. Since *D. m. wrigleyi* and *D. buzzatii* use

primarily *Opuntia sp.*, we expected to find a higher number of detoxification genes in the

other species/subspecies with columnar cacti preference. However, we observed a similar

number of differentially expressed detoxification genes regardless of the host preference.

CYPs were overrepresented in our results, but they represent a diverse gene family that can

have functions other than detoxification, such as those related to developmental pathways

in larvae (82,83). Thus, further analysis of the biological role of up-regulated CYP genes in

D. m. wrigleyi and *D. buzzatii* might be assessed. Alternatively, as we observed in the head

transcriptome, although *Opuntia sp.* has lower toxic compounds compared to columnar

cacti, *D. m. wrigleyi* and *D. buzzatii* may also have a resistance to toxicity (72). Thus, any

cactophilic species is expected to have evolved a certain level of detoxification ability, which

explains our observations. In this study, we propose a set of common HLAU genes in

columnar cacti dweller species associated with detoxification. We observed that the three

D. mojavensis subspecies have the same seven genes with higher expression compared to

D. m. wrigleyi. Most of them are associated with detoxification (*EstB1*, *UDP2*, *UDP-Ugt5*,

Cyp9b2, *UDP-Ugt5-1*, *Cyp12a2*), except *Ir21a*. It is noteworthy that other evolutionary

forces than gene expression might be associated with host shift. For instance, gene

duplication events have been reported in CYPs and GSTs in cactophilic species (84). These

duplications are thought to enable flies' resistance to different toxic environments.

Similarly, *D. buzzatii* and *D. koepferae* show both copy number variation and expression

divergence in detoxification genes, indicating that gene duplication followed by regulatory

or coding sequence divergence contributes to host-specific physiological adaptations

(50,84). These findings support the view that gene duplication serves as a key evolutionary

mechanism facilitating ecological specialization in cactophilic *Drosophila*. Our results show

that divergence in the regulation of gene expression in larvae is one of the factors likely to

shape the adaptation to new hosts.

Transposable elements are a source of transcriptome variability between cactophilic

species

TEs contribute to genome and transcriptome evolution by rewiring regulatory networks, and by incorporating protein domains into gene transcripts. In both cases, TE copies that are inserted within genes can integrate their sequences into the mRNA, generating chimeric transcripts. A recent study with different ecotypes of *D. melanogaster* has shown that these chimeras have the potential to generate transcriptome novelties (24). These chimeras have also been demonstrated to be active in a tissue-specific manner (85). Although the production of chimeric transcripts derived from polymorphic TE insertions is well-known, there is no study addressing the question of how TEs contribute to the evolution of genes associated with host shift. Here, we took advantage of genomic and transcriptomic data to analyze the transcriptional interaction of TEs with HLAU genes through gene-TE chimeric transcripts. Our finding demonstrated that around half of the chimeras have low expression levels (0-5%), and are likely to represent pervasive transcription of TEs. Notably, over 95% of the chimeras overpassing the 5% contribution to gene expression constitute nearly the total gene expression.

Considering previous findings in different tissues of *D. melanogaster*, where 38% of all chimeric transcripts express only the chimeric isoform (85), our result shows a higher prevalence of chimeras in the repertoire of gene splicing. In tetrapods, the exonization of TE-derived transposases has been shown to constitute the unique gene isoform (86). In

HLAU genes, all chimeric transcripts represent TE-exonized transcripts. We suggest that TEs

in the promoters of ORs might still recruit transcription factors at their terminal sequences,

which would prevent their inclusion in the gene mRNA (chimera). In addition, the

maintenance of TEs in the promoters of ORs might also be due to epigenetic marks. TEs

can have chromatin marks (87,88), making them versatile regulatory motifs acting in the

regulation of nearby genes (89,90). Further experimental analysis, as ChIP-seq, may be

performed to test the TEs' function in OR's transcription. Furthermore, TE annotation

remains challenging in non-model organisms. Although we performed several steps to

remove potentially false TE copies from the genome, we can not rule out the possibility of

remaining annotation errors in the exons of HLAU genes.

A comparative analysis between generalist and specialist *Drosophila* species revealed

that niche amplitude is not likely to play a role in TE dynamics in the genomes (91).

Nevertheless, several studies have proposed that the successful adaptation to new

environments in invasive species might be associated with TE activity, despite reduced

genetic diversity caused by founder effects (92,93). Here, we demonstrate that TEs interact

with genes associated with HS, providing a transcriptomic variability in six cactophilic

species with different host preferences. Our results show that ABC transporters produce

chimeric transcripts in all cactophilic species, highlighting a compelling mechanism of

structural innovation. In insects specifically, MITEs (non-autonomous DNA transposons)

have been repeatedly identified within defensome genes, including ABC transporters, such

as in *Helicoverpa* spp., suggesting a potential causal link to insecticide resistance (94).

Functionally, exonized TEs are stably expressed and translated, often producing altered

protein domains or localization signals, thereby expanding proteomic repertoires while

providing raw material for adaptive selection (95). Taken together, our findings suggest

that TE exonization in ABC transporters may offer cactophilic *Drosophila* a versatile

genomic strategy to fine-tune transporter function, potentially under the selective

pressures of xenobiotics. Although we found a high contribution of the chimeric isoform to

total gene expression, future work should be conducted to elucidate which exonized TEs

are translated, and whether they modulate substrate specificity, protein stability, or cellular

localization, thereby affecting organismal fitness.

Our discovery of 13 GST genes in *D. buzzatii* harboring exonized Helitrons

underscores an unprecedented finding of TE-driven functional diversification in

detoxification enzymes. *Helitrons* have been shown to supply novel exons, splice sites, and

promoters across eukaryotes (41). This finding, unique to *D. buzzatii,* suggests a lineage-

specific co-option of *Helitrons* to expand and diversify GST isoforms. GSTs are central to

xenobiotic metabolism, and exonization of Helitron-derived sequences could introduce

novel domains or alter enzymatic properties, potentially conferring adaptive advantages

under environmental pressures such as exposure to cactus-derived toxins. Future work

should be performed to determine the function of these *Helitron*-derived motifs in the host

preference of *D. buzzatii.*

It is important to state that the identification of chimeric transcripts through

transcriptome analysis is an important, but limited step in terms of function. These mRNA

molecules might be degraded by surveillance pathways that control for aberrant mRNA

production, such as no-go decay (96), non-stop decay (97), and nonsense-mediated RNA

decay (98). Our study provides a set of HLAU genes producing chimeric transcripts, in

which part of them might have a phenotypic impact and ultimately be associated with host

preference. However, it is fundamental to perform comparative phenotypic studies of host

adaptation cues with mutant strains for the TEs interacting with HLAU genes identified in

this study. The application of techniques of RNAi aiming at chimeric transcripts, or deletion

of TE insertions related to chimeras with CRISPR/Cas9 (71) would be crucial to confirm the

predictions of our results.

Transposable elements driving host shift

Host specificity is a key mechanism of reproductive isolation in plant pathogens and

is regulated by the repertoire of effector genes within each pathogen. In *Phytophthora sp.*,

the host specificity is partly regulated by RXLR class effectors that facilitate host

exploitation. Notably, synthetic chimeras of a short interspersed element (SINE) linked to an

effector gene in *P. infestans* induced their silencing (99). This silencing likely occurs

naturally, as transcriptional inactivation of effectors is known to occur, and over half of

RXLR effectors in the *P. infestans* genome are located in TE-rich regions (100).

Consequently, TEs inserted near these genes may have influenced host shift in *P. infestans*.

Our findings demonstrate multiple interactions between genes and TEs, highlighting the

contribution of TEs to the genome and transcriptome evolution of cactophilic species.

Although our data did not allow the quantification of the TE insertions frequency in natural

populations, our results demonstrate the first transcriptome-wide evidence of TE co-option

in cactophilic species. The genes producing chimeras with TEs must be further studied,

since the lack of gene-phenotype association between HLAU genes is not fully reported,

preventing us from pinpointing a specific relation between chimeras and possibly host

preference. In addition, our study focused exclusively on nine gene families that have been

shown to evolve during host shift events (12). However, host shift is undoubtedly a complex

trait to study, and many other genes can be associated with the adaptation to new hosts.

For instance, differential expression of genes associated with development and

neurological processes has been identified in cactophilic species reared in different cacti

media (101). Behavioral changes are also based on host shift evolution and have been

reported in many systems (102). Despite focusing on nine relevant gene families, we

suggest further studies including other aspects of host shift than localization, acceptance,

and host usage.

The existence of reproductive barriers between populations that undergone a host

shift is the key process to ecological speciation. In our model, with species from the *buzzatii*

and *mojavensis* clusters, incipient reproductive isolation has been reported between *D. m.*

sonorensis and *D. m. baja* (26,103,104). The premating isolation between them occurs due

to the difference in cuticular hydrocarbons, which participate in the signaling pathway of

mate recognition (105). Importantly, artificially shifting the cactus host was the main cause

of changes in the hydrocarbon profiles. Thus, the source of reproductive barriers is

dependent on the cacti species on which the flies feed and breed (106,107). The close

relationship between HLAU genes to the host shift process, and the observed contribution

of TEs to the evolution of these genes, supports a possible role of TEs in the host

preference of cactophilic species.

Methods

Fly stocks and genome sequencing

Most of the strains were obtained from the UC San Diego *Drosophila* Stock

Center: [*D. m. mojavensis*: Anza (01), Anza Borrego Desert, California, USA, Stock Center

n°: 15081-1352.01; *D. m. wrigleyi*: CI (22), Catalina Island, California, USA, Stock Center

n°: 15081-1352.22; and *D. m. sonorensis*: AG (26), Agiabampo Bay, Sonora, México, Stock

Center n°: 15081-1352.26] and *D. arizonae* [HI (17), collected in Metztitlan, Hidalgo, Mexico,

Stock Center n°: 15081-1271.17]. *D. buzzatii* (Bu28) and *D. koepferae* (Ko2) stocks

correspond to two laboratory lines used in previous works (108,109). Both stocks were

maintained by brother-sister mating for more than a decade and then kept by mass

culturing.

We used the same protocols of DNA extraction and genome sequencing for all

samples. DNA was extracted from 10 males and 10 females from each species using the

Qiagen DNeasy Blood&Tissue kit. Then, we evaluated the genomic DNA amount and

quality with NanoDropTM One UV-Vis spectrophotometer (Thermo Fisher Scientific,

Waltham, MA, USA) and Qubit ® 1.0 Fluorometer (Invitrogen, Carlsbad, CA, USA). Three

micrograms of DNA were repaired using the NEBNext FFPE DNA Repair Mix (NEB M6630).

We performed end repair and dA-tailing using the NEBNext End repair/dA-tailing Module

(E7546, NEB). Ligation was then assessed with the Ligation Sequencing Kit 1D. MinION

sequencing was performed according to the manufacturer's guidelines using R9.4.1 flow

cells (FLO-MIN106, ONT) and a Nanopore MinION Mk1b sequencer (ONT) controlled by

ONT MinKNOW v.18.3.1. Base calling was performed after sequencing using *Guppy v.4.0.5*

in high accuracy mode for isogenic wild-type strains *v.3.1.5.*

Genome assemblies and gene annotation

The quality control for Nanopore reads was performed with *Nanoplot* v.1.10.2

(https://github.com/wdecoster/NanoPlot). Reads with quality lower than 7 were removed

from downstream analysis. In order to assemble contigs, *Flye v.2.8* (110) was used with

default parameters, except –plasmids. All contigs were aligned with *minimap2* v2.16 (111),

with –x map-ont option. The alignment was used to perform the polishing of contigs with

four rounds of *RACON* v1.3.2 (112), default parameters. The assembly quality metrics was

assessed with *Assembly-Stats* v1.0.1 (https://github.com/sanger-pathogens/assembly-stats).

Assembly incongruences were manually visualized with *D-genies* v1.2.0 (113) and corrected

with *samtools* v1.9.0. (114), function *faidx*, and *Gepard* v1.4.0(115) for determination of

breaking points. The super scaffolding of all corrected assemblies was performed with

RaGOO v.1.1 (116), with –s and –t 4 parameters, using the respective reference genome

assembly for each species. Subsequently, the chromosome-scale assemblies were

submitted to a Benchmarking Universal Single-Copy Orhologs (BUSCO) v.5.4.3 analysis

(117) with lineage *diptera_odb10* (-*l* parameter) with nucleotide sequences. The gene

annotations for *D. mojavensis* subspecies and *D. arizonae* were performed with the

software *Liftoff v.1.6.3* (118), with the reference genome of *D. mojavensis*

(GCA_018153725.1) (119). We have used the parameters –*exclude_partial; -a 0.5* and –*s 0.75*

to keep annotated genes with at least 75% of the reference gene length. We removed from

our gene annotation HLAU genes flagged as "low quality protein" and "unestablished gene

function" in the reference *D. mojavensis* genome.

D. buzzatii and *D. koepferae* genomes were annotated with a de novo strategy with

the software *BRAKER3 v.3.0.8* (120). To optimize the prediction of gene structures and

repertoire of isoforms, we used a mix of RNA-seq data from different tissues publicly

available and produced in this work (Supplemental Table 12), in addition to protein

sequences from *D. mojavensis* as reference. Ortholog genes between *D. buzzatii, D.*

koepferae, D. mojavensis, and *D. arizonae* were obtained with *Orthofinder v.2.5.4* (121) using

DIAMOND v.0.9.14 (122) as search engine and default parameters. One-to-one orthologs

were obtained directly from *OrthoFinder v.2.5.4,* and remaining proteins without an

ortholog pair were analyzed with *blastp v.2.5.0+* (123), for which we considered as one-to-

one proteins only those with a single homologous with parameters -*qcov_hsp_perc 95* and

identity >= 90%. Since *Orthofinder* performs its search with protein sequences, the

identification of orthologs from non-coding RNA genes was assessed with *Liftoff v.1.6.3*

(124) using all non-coding RNA genes from *D. mojavensis* as reference to maximize our

homology-based gene annotation.

Molecular evolution analysis

To identify HLAU genes with signatures of positive selection in species that prefer

columnar cacti compared to those that prefer *Opuntia sp.*, we selected the three *D.*

mojavensis subspecies, *D. navojoa*, *D. koepferae*, and *D. buzzatii*. *D. arizonae* species was

removed from this analysis due to its generalist preference for both columnar and *Opuntia*

cacti. The *D. navojoa* genome was retrieved from NCBI (GCF_001654015.2) (125). For the

other Nanopore genomes sequenced in this work, we performed a polishing step with

Illumina paired-end reads. We used the software *NextPolish* (126) to fix base errors in the

assembled genomes. The three *D. mojavensis* subspecies were polished with genomic

Illumina reads sequenced previously (127), whereas for *D. koepferae* and *D. buzzatii,* we

used publicly available genomic Illumina reads (128) and RNA-seq reads (50), respectively.

In the latter, to increase the read coverage and optimize the polishing to the highest

number of genes and exons as possible, we merged RNA-seq reads from adult males and

females (50). All genomes were corrected using two rounds of polishing, as recommended

(126). Then, we extracted the longest CDS sequences for all genes combining AGAT (129)

agat_extract_sequences.pl parameters *-t mrna*, *samtools faidx* (130), and *seqtk*

(https://github.com/lh3/seqtk). Posteriorly, we intersected the occurrence of orthologous

genes between the six genomes, recovering a list of genes that were present in all of them.

The sequences were aligned using *MAFFT v.7.487* [67] and the non-aligned codons were

removed using *pal2nal* (131). To obtain a strongly supported tree for this analysis, we

inferred the species phylogeny tree with 2,043 shared BUSCO genes. The phylogeny was

estimated using the concatenated alignment (3,730,914 sites) with *IQ-TREE v. 2.2.2.6* (132),

using the best model for each gene, 10,000 replicates of ultrafast bootstrap and 1000

replicates for Approximate Likelihood-Ratio Test (ALRT) (parameters *-m MFP -bb 10000 -*

alrt 1000). Finally, we carried out an analysis to detect signatures of positive selection for

each gene individually with *CODEML* from *PAML v.4.10.7* (133), using *branch-site* approach.

In order to find signatures of positive selection in species that use columnar cacti, we

marked the respective branches of *D. koepferae*, *D. m. mojavensis,* and *D. m. sonorensis* with

the same label (#1), to compare them with the background species that use *Opuntia sp.* (*D.*

buzzatii, D. m. wrigleyi, and *D. navojoa*). For this, we tested two hypotheses: (i) H0 – Null

hypothesis, in which all branches evolved under negative selection (model = 2, Nssites = 2,

fix_omega = 1, omega = 1); (ii) H1 – Alternative hypothesis, in which the labeled branches

(foreground) have sites with signatures of positive selection (model = 2, NSsites = 2,

fix_omega = 0, omega = 1.5). The hypotheses were tested through a chi-square test,

calculated by the log-likelihood (lnL) ratio between H1 and H0 (2ΔlnL) and degrees of

freedom obtained from the difference between the number of estimated parameters for H1

and H0 (Δnp). Sites under positive selection were identified by Bayes Empirical Bayes and

posterior probability, implemented in *CODEML*.

RNA extraction and sequencing

We performed RNA-seq for larvae and head tissues from _D. buzzatii, D. koepferae_,

D. arizonae, and the four _D. mojavensis_ subspecies. The RNA extraction was assessed with

the RNeasy Plus Kit for 10 female heads (10 days old) and 15 larvae, with three technical

replicates. The library preparation was done with stranded mRNA-seq/standard quantity

(ligation), and 100 bp paired-end reads were produced in an Illumina HiSeq 4000. Illumina

adaptors and low-quality reads were removed for downstream analysis with _Trimommatic_

_v.0.39_ (134), parameters: _LEADING:3; TRAILING:3; SLIDINGWINDOW:4:15; HEADCROP:10;_

CROP:90.

TE annotation

For each sequenced genome, we developed a novel pipeline combining automatic

and manual approaches to construct TE consensuses. First, we used _Earl Grey v.2.2_ (135) to

build de novo TE libraries with _RepeatModeler2 v.2.0.4_ (136), which uses _RECON v.1.08_ (137),

and _RepeatScout v.1.0.6_ (138). We used the following parameters: -r 32281 (NCBI taxon ID

for _Drosophila_), -c yes (cluster redundant consensus), -m yes (remove consensus with less

than 100 nt), and -e yes (enable _HELIANO v1.2.1_ (139) to identify Helitrons). In short,

EarlGrey automatically retrieves TE consensus, and performs an extension of each sequence

using a method named "Blast, Extract, Extend" (BEE) (140). Such method extracts the

sequences of the 20 full-length (or longest) TE insertions for each consensus, and then it

retrieves the 1 kb flanking regions from them. Posteriorly, insertions altogether with

flanking regions are aligned with *MAFFT v.7.487* (141), and the alignment is trimmed with

trimAl v.1.4 (142), parameters *–gt 0.6* and *–cons 60*. Subsequently, the alignment is analyzed

to check whether the flanking regions from the full-length insertions have high quality,

meaning that they are part of the TE rather than genomic regions. Finally, the consensus

sequence for each TE family is updated with *EMBOSS v.6.6.0.0* (143). The BEE method is

repeated up to 10 times for each consensus.

Once the extended TE libraries were obtained for the six genomes with EarlGrey, we

implemented our pipeline into eight steps. Firstly, we removed all sequences other than TEs

from EarlGrey's library: satellites, low complexity repeats, tRNA, and rRNA. At step 2, we

checked the presence of redundant consensus respecting the 80-80 (35) rule with all-vs-all

blastn (-qcov_hsp_perc 80 and -perc_identity 80). Surprisingly, ~45% of the consensuses

were duplicated. Some of them were identical, or nearly identical, but in opposite strands.

We selected the longest consensus from the ones considered redundant. At step 3, we

used Tandem Repeat Finder (144) to identify consensuses that were had over 50%

annotated as tandem repeats. Consensus falling in this cutoff were removed for being

indistinguishable from a tandem repeat. At step 4, we removed all consensus with high

similarity with *D. mojavensis wrigleyi* CDSs using blastn, parameters *-qcov_hsp_perc 80 and -*

perc_identity 80. TE consensuses that had passed our filtering thresholds were then

submitted to the classification analysis. At step 5 of the pipeline, TE consensus will be

classified at the family level based on similarity analysis of non-reference consensus with a

database provided by the user (https://github.com/OliveiraDS-hub/Pipelines-Cactophilic-

<u>Drosophila-Species</u>). In our case, we used all consensuses from *Drosophila* species available

on Dfam v3.7 (145). Due to the "curated" Dfam library containing only *D. melanogaster*

consensus, we used the "uncurated" library from Dfam. We extracted *Drosophila* TEs with

famdb.py v.0.4.3, parameters -i Dfam.h5 families --include-class-in-name -f fasta_name -ad

'Diptera'. Then, we filtered out non-Drosophila sequences and "Unknown" TEs with the Unix

command line: grep -i 'drosophila' file.fa | grep -v -i 'unknown', obtaining 49,916 out of

344,562 consensus. The pipeline firstly classifies at the family level by doing a blastn (-

qcov_hsp_perc 80 and -perc_identity 80) between the *Drosophila* Dfam consensus and the

provided de novo consensus. Approximately 50% of the identified consensus were

classified on the family level based on this blastn. The consensus without results from this

stringent blastn and smaller than 200 nt were filtered out. Due to the stringency of the

matches with blastn, we also masked the remaining unclassified consensus with the

Drosophila Dfam consensus with RepeatMasker .v4.1.8 (146). Based on RepeatMasker

output, we computed the length of all matches from a given family in the unclassified

consensus, and how much it has been covered based on the unclassified length. If the

coverage is > 80% of the total length, we assume that the unclassified consensus has

diverged from that one of the database, but it still classified as the same matched family.

This step classified ~25% of our consensuses. Finally, the remaining unclassified sequences

that could not be classified at the family level are classified at the class/order level. We used

a cutoff of 50% of covered matches in the unknown consensus to classify their classes. As

the family name, we substituted the pattern from RepeatModeler2 as (i.e. rnd-1_family-1)

to a pattern representing: "species preffix", "author initials", "homology percentage" (i.e.

Dari_DSO_86.0275p#DNA/TcMar-Tc1). In this example, "Dari" stands for *D. arizonae*, "DSO"

is the author's initials, and "86.0275p" represents the percentage of the total length that

this consensus has been matched with a DNA/TcMar-Tc1. The remaining putative TE

consensuses with less than 50% of coverage with known *Drosophila* TEs were removed. At

the end of this step, we end up with polished TE consensus.

In step 6, we checked the strandness of TE consensus sequences based on stranded

RNA-seq data. During the clustering of similar sequences and removal of redundancy, we

noticed many consensuses with nearly identical sequences, but in different senses. To

correct TE strandness, we used our stranded RNA-seq data. We annotated TE insertions in

the genome with *RepeatMasker v.4.1.4* (122), parameters *-cutoff 250*, *-s*, and *-norna*. Then,

we mapped the reads from a mixed library from head and larvae tissues with STAR v2.7.11b

(147). To be certain about the strandness origin of the reads, we removed all insertions that

have any overlap with other TEs and/or exons. Then, for each consensus, the pipeline

identifies the three copies in the genome with higher expression. Based on these copies,

the number of uniquely mapped reads to the same strand of the copy is quantified. If >

50% of the reads align to the same strand of the copies, we assume the consensus

strandness was right. Otherwise, the pipeline converts the consensus sequence to its

reverse complement. This step will add a strandness tag to the consensus ID. For instance,

if 75% of the reads support forward strand, a tag "75st" will be added to the family name.

This is useful to provide a metric of certainty about the strandness of each consensus. Such

information can be valuable when working with piRNAs, where alignment strandness is

crucial to interpret the results. Then, we used *RepeatCraft* (148) with *LTR Finder v.1.2* results

to merge LTRs and their respective internal regions. Subsequently, we removed all TE

insertions shorter than 80 nucleotides, following the classification rule 80-80-80 (35).

Finally, we performed a last filtering step in the TE annotation to remove insertions

composed of simple sequence repeats. We used *TRF v.4.09* (144) in all TE insertions from

each genome, with parameters *2, 5, 6, 75, 20, 50, 500, -m, -h*, to remove TEs with masked

repeats over more than 50% of their sequences. This method has been previously used to

remove TE copies that are indistinguishable from simple repeats (24,149).

TE enrichment at regulatory regions

To identify TE insertions within and nearby genes for each genome, we created *bed*

files with the genomic positions of all insertions. Then, we created *bed* files for the gene

regions corresponding to the 2 Kb upstream of the transcription start-site (TSS). To test

whether the prevalence of TEs in each gene region was significantly higher than expected

(enriched) compared to random genomic distributions of TEs in the same region of other

genes, we performed permutation tests with the R package *regioneR* (150), parameter

randomization = resampleRegions, with 2,000 random samples. For each gene family with *n*

genes, the number of genes with TEs was compared to other random *n* genes 2,000 times.

This analysis allowed us to infer whether there are HLAU gene families with the observed

frequency of genes with TEs at the promoter region higher than expected compared to the

rest of the genome. We also selected zinc-finger proteins and serine/threonine kinases as

two non-HLAU gene families to compare our findings. Taking into account that TEs might

harbor regulatory and/or coding motifs in both strands (62), our analysis considered TEs

within the investigated gene regions regardless of the strand.

Identification of TE-rich genomic regions

We split the genomes into bins of 200 kb, generating a bed file with such intervals

across the genomes. Then, we used the function *intersect* from *bedtools v.2.30.0* (151) to

identify the number of TE insertions located in each genomic bin. Posteriorly, we merged all

overlapped TE insertions into single ranges. These ranges were summed to obtain the total

TE content in bp for each 200kb bin. By plotting the TE content of each genome, we

noticed that bins with more than 50kb of TE content were outliers of the normal

distribution. Therefore, we defined TE-rich regions as all bins with > 50kb of TE content,

which represents 25% of the bin size. We used the same function from *bedtools* to identify

whether OR genes were present in the TE-rich regions, or the TE-poor (bins with less than

50kb of TE content).

Transcription factor binding sites on TE insertions

We assessed the identification of transcription factor binding sites (TFBSs) with *FIMO*

(152), from the *MEME v. 5.0.5* suite tools (153). We extracted the sequences of all TE

insertions located 2kb upstream of ORs with *bedtools v. 2.30.0*, function *getfasta*, using

parameter *-s* (151). We obtained the TFBS motifs described for *D. melanogaster* from Jaspar

database (154), for the transcription factors (TFs) acj6, zf30c, fer1, onecut, and xbp1, which

have been demonstrated to activate ORs (43). To compare our findings with OR genes, we

performed the same analysis with Serine/Threonine Kinases and zinc-finger as non-HLAU

gene families. Only TFBSs with p-value < 0.01 were considered. The enrichment of OR

TFBSs in OR genes compared to non-HLAU gene families was assessed with the four

frequencies: 1) ORs with TFBSs; 2) ORs without TFBSs; 3) non-HLAU with TFBSs; and 4) non-

HLAU without TFBSs.

Differential expression of genes

We aligned the RNA-seq reads with *STAR v.2.7.3* (147), using the assembled

genomes and annotations of their respective species. The alignments were used to create

the count matrix reporting the expression of each gene, with *htseq v.0.13.5* (155),

parameters *–order pos*; *-s reverse*; *–i gene*; *-t exon*; the latter indicates that the count for

each gene corresponds to the sum of all reads aligned into exons (CDSs and UTRs). We

merged the counts from species of the *buzzatii* and *mojavensis* clusters separately,

maintaining only ortholog genes that have at least 90% of the length of the annotated

gene in the reference genome. This threshold allowed us to avoid misinterpretation of the

expression level by comparing genes with different lengths. Then, we performed

independent statistical analysis for species of the *buzzatii* and *mojavensis* clusters with

DEseq2 v.1.28.1 (156), using a single-factor experimental design, which considers only the

factor that different species may have differentially expressed genes. Due to the

phylogenetic distance between species of the *buzzatii* and *mojavensis* clusters, we

performed the differential expression analysis separately for each group. We only

considered genes with adjusted *p-value* < 0.05 and |*log2 fold-change*| > 1 as differentially

expressed. To test for association between TE enrichment and differential expression, we

selected HLAU genes with the mean of normalized counts between replicates >= 100.

Then, we build the contingency table to apply the X^2 independence test, splitting them into

four categories: differentially expressed, non-differentially expressed, with TEs at the

promoter region, and without TEs at the promoter region. We carried out the test for all

pairwise analyses between cactophilic species with preference for *Opuntia sp* and columnar

cacti. X^2 with p-value < 0.05 was considered significant. The proportional differences

between localization, acceptance, and host use genes compared to non-HLAU genes

(Ser/Thr Kinases and zinc-finger gene families) were assessed with Fisher's exact test, with a

contingency table containing the average number of genes per group: 1) non-HLAU

differentially expressed genes; 2) non-HLAU non-differentially expressed genes; 3) HLAU

differentially expressed genes; and 4) HLAU non-differentially expressed genes. *Fisher's*

Exact test < 0.05 was considered significant.

985 *Identification of gene-TE chimeric transcripts*

We detected gene-TE chimeric transcripts with *ChimeraTE v.2.0.0* Mode 1 (24). Each

species was analyzed using its respective genome, gene annotation, and TE annotation.

Shortly, *ChimeraTE* identifies chimeras based on paired-end reads spanning from exons to

TE sequences. Based on the TE position compared to the gene (upstream, exon, intron,

downstream), chimeric transcripts are classified into categories. We have used default

parameters, except *--replicate 3*, which considers true chimeras only those identified in the

three RNA-seq replicates, and *–coverage 10* to obtain chimeras with at least 10 chimeric

reads on average between replicates. To calculate the contribution of the chimeric

transcript expression relative to total gene expression, we developed a new module for

ChimeraTE. This module can be used downstream to ChimeraTE mode 1 analysis. In the first

step, all RNA-seq replicates will be merged into a single fastq library. Then, the merged

reads are aligned to the genome with STAR (147). Next, Stringtie2 (157) is used to assess

transcript expression, including both reference and non-reference isoforms. All TEs

generating chimeras are intersected with exons predicted by Stringtie with pyranges (158).

Only overlaps with 80% of the TE within the exon are considered. Finally, the expression of

the TE-containing isoforms is compared to the total gene expression. In our analysis, we

removed all chimeric transcripts for which Stringtie was not able to build the chimeric

isoform. In addition, aiming to keep chimeras with putative function, we selected only the

ones with contribution >= 25%. This analysis can be reproduced with any output from

ChimeraTE with the code "mode1_contrib_chimeras.py" (https://github.com/OliveiraDS-

hub/ChimeraTE/blob/main/scripts/mode1_contrib_chimeras.py).

Data access

The quality control data from ONT and RNA-seq data, the genome assemblies, and their

respective gene and TE annotation (libraries and gtf files) are available in the Zenodo

repository: https://zenodo.org/records/16501018. The codes to reproduce differential

expression analysis, positive selection tests, gene annotation, permutation tests, TE

annotation, and TFBS analysis are available on the GitHub repository:

https://github.com/OliveiraDS-hub/Pipelines-Cactophilic-Drosophila-Species.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We thank Séverine Chambeyron for useful discussions and advice. This work was performed

using the computing facilities of the CC LBBE/PRABI. This study was funded by Brazilian

agencies FAPESP- The São Paulo Research Foundation [grant number 2020/06238-2] and

from the National Council for Scientific and Technological Development (308020/2021-9)

to C. M. A. C; the Eiffel Fellowship Campus France [P769649C] and CAPES – PrInt

[88887.716810/2022-00] to D. S. O.; by Ministerio de Ciencia e Innovación (Spain) [grant

number PID2021-127107NB-I00], and Generalitat de Catalunya (Spain) [grant number 2021

SGR 00526] to M. P. G. G.; and by Udl-FAPESP ANR-IDEX-0005 to C. V. This study was

financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior –

Brasil (CAPES) – Finance Code 001 to DSO.

Authors' contributions: A. L., A. B, M. P. G. G, performed experiments; F.S., Performed

genome assemblies; W. V. B. N. performed phylogenetic analysis; D. S. O performed

computational analysis related to genomics, transcriptomics, and statistical tests; D. S. O.

wrote and edited the paper; C. V. and C. M. A. C. designed the study, and reviewed the

manuscript. All authors read, corrected, and approved the final manuscript.

References

1.   Whiteman NK, Pierce NE. Delicious poison: genetics of *Drosophila* host plant

preference. Trends in Ecology & Evolution. 2008;23(9):473–8.

2.   Forbes AA, Devine SN, Hippee AC, Tvedte ES, Ward AKG, Widmayer HA, et al.

Revisiting the particular role of host shifts in initiating insect speciation. Evolution. 2017

May;71(5):1126–37.

3.   Rundle HD, Nosil P. Ecological speciation. Ecology Letters. 2005;8(3):336–52.

4.   Howard DJ, Berlocher SH. Endless Forms: Species and Speciation. Oxford University

Press; 1998. 486 p.

5.   O'Grady PM, Markow TA. Rapid morphological, behavioral, and ecological evolution

in *Drosophila*: comparisons between the endemic Hawaiian *Drosophila* and the cactophilic

repleta species group. In: Singh RS, Xu J, Kulathinal RJ, editors. Rapidly Evolving Genes and

Genetic   Systems.   Oxford   University   Press;   2012.   Available   from:

https://doi.org/10.1093/acprof:oso/9780199642274.003.0018

6.   Oliveira DCSG, Almeida FC, O'Grady PM, Armella MA, DeSalle R, Etges WJ.

Monophyly, divergence times, and evolution of host plant use inferred from a revised

phylogeny of the *Drosophila repleta* species group. Molecular Phylogenetics and Evolution.

2012;64(3):533–44.

7.   Stintzing FC, Carle R. Cactus stems (*Opuntia spp.*): A review on their chemistry,

technology, and uses. Molecular Nutrition & Food Research. 2005;49(2):175–94.

8.   Gomez GA. Transpecific Polymorphisms in an Inversion Linked Esterase Locus in

Drosophila buzzatii. Molecular Biology and Evolution. 2003;20(3):410–23.

9.   Hasson E, Naveira H, Fontdevila A. The breeding sites of the Argentinian species of

the *Drosophila mulleri* complex (subgenus *Drosophila-repleta* group). Rev Chil Hist Nat.

1992;65:319–26.

10.   Newby BD, Etges WJ. Host Preference Among Populations of *Drosophila mojavensis*

(Diptera: Drosophilidae) that Use Different Host Cacti. Journal of Insect Behavior.

1998;11(5):691–712.

11.   Rane RV, Clarke DF, Pearce SL, Zhang G, Hoffmann AA, Oakeshott JG. Detoxification

Genes Differ Between Cactus-, Fruit-, and Flower-Feeding *Drosophila*. Journal of Heredity.

2019;110(1):80–91.

12.   Markow TA. Host use and host shifts in *Drosophila*. Current Opinion in Insect

Science. 2019;31:139–45.

13.   Montell C. A taste of the *Drosophila* gustatory receptors. Current Opinion in

Neurobiology. 2009;19(4):345–53.

14.   Ni L. The Structure and Function of Ionotropic Receptors in *Drosophila*. Front Mol

Neurosci. 2021;13:638839.

15.   Chen Y, Amrein H. Ionotropic Receptors Mediate *Drosophila* Oviposition Preference

through Sour Gustatory Receptor Neurons. Current Biology. 2017;27(18):2741-2750.e4.

16.   Gomez-Diaz C, Martin F, Garcia-Fernandez JM, Alcorta E. The Two Main Olfactory

Receptor Families in *Drosophila*, ORs and IRs: A Comparative Approach. Front Cell Neurosci.

2018;12:253.

17.   Lang M, Murat S, Clark AG, Gouppil G, Blais C, Matzkin LM, et al. Mutations in the

neverland Gene Turned *Drosophila pachea* into an Obligate Specialist Species. Science.

2012;337(6102):1658–61.

18.   Aravin AA, Sachidanandam R, Girard A, Fejes-Toth K, Hannon GJ. Developmentally

Regulated piRNA Clusters Implicate MILI in Transposon Control. Science.

2007;316(5825):744–7.

19.   Fedoroff NV. Transposable Elements, Epigenetics, and Genome Evolution. Science.

2012;338(6108):758–67.

20.   Capy P. Taming, Domestication and Exaptation: Trajectories of Transposable

Elements in Genomes. Cells. 2021;10(12):3590.

21.   Drongitis D, Aniello F, Fucci L, Donizetti A. Roles of Transposable Elements in the

Different Layers of Gene Expression Regulation. IJMS. 2019;20(22):5755.

22.   Magwire MM, Bayer F, Webster CL, Cao C, Jiggins FM. Successive Increases in the

Resistance of *Drosophila* to Viral Infection through a Transposon Insertion Followed by a

Duplication. Begun DJ, editor. PLoS Genet. 2011;7(10):e1002337.

23.   Lipatov M, Lenkov K, Petrov DA, Bergman CM. Paucity of chimeric gene-

transposable element transcripts in the *Drosophila melanogaster* genome. BMC Biol.

2005;3(1):24.

24.   Oliveira DS, Fablet M, Larue A, Vallier A, Carareto CMA, Rebollo R, et al. ChimeraTE: a

pipeline to detect chimeric transcripts derived from genes and transposable elements.

Nucleic Acids Research. 2023;51(18):9764–84.

25.   Matzkin LM. The Molecular Basis of Host Adaptation in Cactophilic *Drosophila*:

Molecular Evolution of a Glutathione *S* -Transferase Gene (*GstD1*) in *Drosophila*

mojavensis. Genetics. 2008;178(2):1073–83.

26.   Pfeiler E, Castrezana S, Reed LK, Markow TA. Genetic, ecological and morphological

differences among populations of the cactophilic *Drosophila mojavensis* from southwestern

USA and northwestern Mexico, with descriptions of two new subspecies. Journal of Natural

History. 2009;43(15–16):923–38.

27.   Etges WJ. Genetics of Host-Cactus Response and Life-History Evolution Among

Ancestral and Derived Populations of Cactophilic *Drosophila mojavensis*. Evolution.

1993;47(3):750–67.

28.   Crowley-Gall A, Date P, Han C, Rhodes N, Andolfatto P, Layne JE, et al. Population

differences in olfaction accompany host shift in *Drosophila mojavensis*. Proc R Soc B. 2016

Aug 31;283(1837):20161562.

29.   Benowitz KM, Coleman JM, Allan CW, Matzkin LM. Contributions of cis- and trans-

Regulatory Evolution to Transcriptomic Divergence across Populations in the *Drosophila*

mojavensis Larval Brain. Genome Biology and Evolution. 2020;12(8):1407–18.

30.   Rajpurohit S, Oliveira CC, Etges WJ, Gibbs AG. Functional genomic and phenotypic

responses to desiccation in natural populations of a desert drosophilid. Molecular Ecology.

2013;22(10):2698–715.

31.   De Panis D, Dopazo H, Bongcam-Rudloff E, Conesa A, Hasson E. Transcriptional

responses are oriented towards different components of the rearing environment in two

Drosophila sibling species. BMC Genomics. 2022;23(1):515.

32.   Matzkin L, Markow TA. Transcriptional Differentiation Across the Four Subspecies of

Drosophila mojavensis. In: Speciation: Natural Processes, Genetics and Biodiversity. New

York: Nova Scientific Publishers; 2013. p. 119–35.

33.   Benowitz KM, Allan CW, Jaworski CC, Sanderson MJ, Diaz F, Chen X, et al.

Chromosome-length genome assemblies of cactophilic *Drosophila* illuminate links between

structural   and   sequence   evolution.   bioRxiv;   2022.   Available   from:

https://www.biorxiv.org/content/10.1101/2022.10.16.512445v1

34.   Oliveira DCSG, Almeida FC, O'Grady PM, Armella MA, DeSalle R, Etges WJ.

Monophyly, divergence times, and evolution of host plant use inferred from a revised

phylogeny of the *Drosophila* repleta species group. Molecular Phylogenetics and Evolution.

2012;64(3):533–44.

35.   Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified

classification system for eukaryotic transposable elements. Nat Rev Genet. 2007;8(12):973–

82.

36.   Sanchez-Flores A, Peñaloza F, Carpinteyro-Ponce J, Nazario-Yepiz N, Abreu-Goodger

C, Machado CA, et al. Genome Evolution in Three Species of Cactophilic *Drosophila*. G3

Genes|Genomes|Genetics. 2016;6(10):3097–105.

37.   Smith G, Lohse K, Etges WJ, Ritchie MG. Model-based comparisons of

phylogeographic scenarios resolve the intraspecific divergence of cactophilic *Drosophila*

mojavensis: Phylogeography of *Drosophila mojavensis*. Molecular Ecology.

2012;21(13):3293–307.

38.   Guio L, Vieira C, González J. Stress affects the epigenetic marks added by natural

transposable element insertions in *Drosophila melanogaster*. Sci Rep. 2018;8(1):12197.

39.   Daborn PJ, Yen JL, Bogwitz MR, Le Goff G, Feil E, Jeffers S, et al. A Single P450 Allele

Associated with Insecticide Resistance in *Drosophila*. Science. 2002;297(5590):2253–6.

40.   Chang NC, Rovira Q, Wells J, Feschotte C, Vaquerizas JM. Zebrafish transposable

elements show extensive diversification in age, genomic distribution, and developmental

expression. Genome Res. 2022;32(7):1408–23.

41.   Thomas J, Phillips CD, Baker RJ, Pritham EJ. Rolling-Circle Transposons Catalyze

Genomic Innovation in a Mammalian Lineage. Genome Biology and Evolution.

2014;6(10):2595–610.

42.   Pritham EJ, Feschotte C. Massive amplification of rolling-circle transposons in the

lineage of the bat *Myotis lucifugus*. Proceedings of the National Academy of Sciences.

2007;104(6):1895–900.

43.   Jafari S, Alkhori L, Schleiffer A, Brochtrup A, Hummel T, Alenius M. Combinatorial

activation and repression by seven transcription factors specify *Drosophila* odorant receptor

expression. PLoS Biol. 2012;10(3):e1001280.

44.   Capy P. Taming, Domestication and Exaptation: Trajectories of Transposable

Elements in Genomes. Cells. 2021;10(12):3590.

45.   Fogleman JC, Danielson PB. Chemical Interactions in the Cactus-Microorganism-

Drosophila Model System of the Sonoran Desert. American Zoologist. 2001;41(4):877–89.

46.   Fogleman JC, Heed WB, Kircher HW. *Drosophila mettleri* and senita cactus alkaloids:

Fitness measurements and their ecological significance. Comparative Biochemistry and

Physiology Part A: Physiology. 1982;71(3):413–7.

47.   McBride CS. Rapid evolution of smell and taste receptor genes during host

specialization in *Drosophila sechellia*. Proceedings of the National Academy of Sciences.

2007;104(12):4996–5001.

48.   McBride CS, Arguello JR. Five *Drosophila* Genomes Reveal Nonneutral Evolution and

the Signature of Host Specialization in the Chemoreceptor Superfamily. Genetics.

2007;177(3):1395–416.

49.   Allan CW, Matzkin LM. Genomic analysis of the four ecologically distinct cactus host

populations of *Drosophila mojavensis*. BMC Genomics. 2019;20(1):732.

50.   Guillén Y, Rius N, Delprat A, Williford A, Muyas F, Puig M, et al. Genomics of

Ecological Adaptation in Cactophilic *Drosophila*. Genome Biology and Evolution.

2015;7(1):349–66.

51.   Kristensen DM, Wolf YI, Mushegian AR, Koonin EV. Computational methods for Gene

Orthology inference. Briefings in Bioinformatics. 2011;12(5):379–91.

52.   Faulkner GJ, Kimura Y, Daub CO, Wani S, Plessy C, Irvine KM, et al. The regulated

retrotransposon transcriptome of mammalian cells. Nat Genet. 2009;41(5):563–71.

53.   Kapusta A, Kronenberg Z, Lynch VJ, Zhuo X, Ramsay L, Bourque G, et al.

Transposable Elements Are Major Contributors to the Origin, Diversification, and

Regulation of Vertebrate Long Noncoding RNAs. Hoekstra HE, editor. PLoS Genet.

2013;9(4):e1003470.

54.   Emera D, Casola C, Lynch VJ, Wildman DE, Agnew D, Wagner GP. Convergent

Evolution of Endometrial Prolactin Expression in Primates, Mice, and Elephants Through the

Independent Recruitment of Transposable Elements. Molecular Biology and Evolution.

2012;29(1):239–47.

55.   Ye M, Goudot C, Hoyler T, Lemoine B, Amigorena S, Zueva E. Specific subfamilies of

transposable elements contribute to different domains of T lymphocyte enhancers. Proc

Natl Acad Sci USA. 2020;117(14):7905–16.

56.   Xie M, Hong C, Zhang B, Lowdon RF, Xing X, Li D, et al. DNA hypomethylation within

specific transposable element families associates with tissue-specific enhancer landscape.

Nat Genet. 2013;45(7):836–41.

57.   Schmidt D, Schwalie PC, Wilson MD, Ballester B, Gonçalves Â, Kutter C, et al. Waves

of Retrotransposon Expansion Remodel Genome Organization and CTCF Binding in

Multiple Mammalian Lineages. Cell. 2012;148(1–2):335–48.

58.   Wang J, Lunyak VV, Jordan IK. Genome-wide prediction and analysis of human

chromatin boundary elements. Nucleic Acids Research. 2012;40(2):511–29.

59.   Sundaram V, Cheng Y, Ma Z, Li D, Xing X, Edge P, et al. Widespread contribution of

transposable elements to the innovation of gene regulatory networks. Genome Res.

2014;24(12):1963–76.

60.   Gill RA, Scossa F, King GJ, Golicz AA, Tong C, Snowdon RJ, et al. On the Role of

Transposable Elements in the Regulation of Gene Expression and Subgenomic Interactions

in Crop Genomes. Critical Reviews in Plant Sciences. 2021;40(2):157–89.

61.   Hof AE van't, Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, et al. The

industrial melanism mutation in British peppered moths is a transposable element. Nature.

2016;534(7605):102–5.

62.   Mateo L, Ullastres A, González J. A Transposable Element Insertion Confers

Xenobiotic Resistance in *Drosophila*. PLoS Genet. 2014;10(8):e1004560.

63.   Dekker T, Ibba I, Siju KP, Stensmyr MC, Hansson BS. Olfactory Shifts Parallel

Superspecialism for Toxic Fruit in *Drosophila melanogaster* Sibling, *D. sechellia*. Current

Biology. 2006;16(1):101–9.

64.   Matsuo T, Sugaya S, Yasukawa J, Aigaki T, Fuyama Y. Odorant-Binding Proteins

OBP57d and OBP57e Affect Taste Perception and Host-Plant Preference in *Drosophila*

sechellia. PLoS Biol. 2007;5(5):e118.

65.   McKenzie SK, Kronauer DJC. The genomic architecture and molecular evolution of

ant odorant receptors. Genome Res. 2018;28(11):1757–65.

66.   Cultrone A, Domínguez YR, Drevet C, Scazzocchio C, Fernández-Martín R. The tightly

regulated promoter of the *xanA* gene of *Aspergillus nidulans* is included in a helitron: A

fungal helitron includes a regulated promoter. Molecular Microbiology. 2007;63(6):1577–87.

67.   Gupta S, Gallavotti A, Stryker GA, Schmidt RJ, Lal SK. A novel class of Helitron-

related transposable elements in maize contain portions of multiple pseudogenes. Plant

Mol Biol. 2005;57(1):115–27.

68.   Huang K, Li CF, Wu J, Wei JH, Zou Y, Han MJ, et al. Enhancer activity of *Helitron* in

sericin-1 gene promoter from *Bombyx mori*: Enhancer activity of Helitrons in sericin-1.

Insect Science. 2016;23(3):396–405.

69.   Batista RA, Moreno-Romero J, Qiu Y, van Boven J, Santos-González J, Figueiredo DD,

et al. The MADS-box transcription factor PHERES1 controls imprinting in the endosperm by

binding to domesticated transposons. eLife. 2019;8:e50541.

70.   Carareto CMA, Hernandez EH, Vieira C. Genomic regions harboring insecticide

resistance-associated *Cyp* genes are enriched by transposable element fragments carrying

putative transcription factor binding sites in two sibling *Drosophila* species. Gene.

2014;537(1):93–9.

71. Merenciano M, Aguilera L, González J. Two-step CRISPR-Cas9 protocol for

transposable element deletion in *D. melanogaster* natural populations. STAR Protocols.

2023;4(3):102501.

72. Wright CR, William S. Characterization of Volatiles of Necrotic *Stenocereus thurberi*

and *Opuntia littoralis* and Toxicity and Olfactory Preference of *Drosophila melanogster, D.*

mojavensis wrigleyi, and *D. mojavensis sonorensis* to Necrotic Cactus Volatiles. Natural

Product Communications. 2014;9(6):1934578X1400900833.

73. Hasson E, De Panis D, Hurtado J, Mensch J. Host Plant Adaptation in Cactophilic

Species of the *Drosophila buzzatii* Cluster: Fitness and Transcriptomics. Journal of Heredity.

2019;110(1):46–57.

74. Hickner PV, Rivaldi CL, Johnson CM, Siddappaji M, Raster GJ, Syed Z. The making of

a pest: Insights from the evolution of chemosensory receptor families in a pestiferous and

invasive fly, *Drosophila suzukii*. BMC Genomics. 2016;17(1):648.

75. Mansourian S, Stensmyr MC. The chemical ecology of the fly. Current Opinion in

Neurobiology. 2015;34:95–102.

76.   Goldman-Huertas B, Mitchell RF, Lapoint RT, Faucher CP, Hildebrand JG, Whiteman

NK. Evolution of herbivory in Drosophilidae linked to loss of behaviors, antennal responses,

odorant receptors, and ancestral diet. Proc Natl Acad Sci USA. 2015;112(10):3026–31.

77.   Crowley-Gall A, Date P, Han C, Rhodes N, Andolfatto P, Layne JE, et al. Population

differences in olfaction accompany host shift in *Drosophila mojavensis*. Proc R Soc B.

2016;283(1837):20161562.

78.   Ruiz A, Heed WB. Host-Plant Specificity in the Cactophilic *Drosophila mulleri* Species

Complex. The Journal of Animal Ecology. 1988;57(1):237.

79.   Mateus RP, Nazario-Yepiz NO, Ibarra-Laclette E, Ramirez Loustalot-Laclette M,

Markow TA. Developmental and Transcriptomal Responses to Seasonal Dietary Shifts in the

Cactophilic *Drosophila mojavensis* of North America. Journal of Heredity. 2019;110(1):58–67.

80.   Matzkin LM. Population transcriptomics of cactus host shifts in *Drosophila*

mojavensis: Population Transcriptomics of *D. mojavensis*. Molecular Ecology.

2012;21(10):2428–39.

81.   De Panis D, Dopazo H, Bongcam-Rudloff E, Conesa A, Hasson E. Transcriptional

responses are oriented towards different components of the rearing environment in two

Drosophila sibling species. BMC Genomics. 2022;23(1):515.

82.   Feyereisen R. Insect CYP Genes and P450 Enzymes. In: Insect Molecular Biology and

Biochemistry.   Elsevier;   2012.   p.   236–316.   Available   from:

https://linkinghub.elsevier.com/retrieve/pii/B978012384747810008X

83.   Tijet N, Helvig C, Feyereisen R. The cytochrome P450 gene superfamily in *Drosophila*

melanogaster: Annotation, intron-exon organization and phylogeny. Gene. 2001;262(1–

2):189–98.

84.   Rane RV, Pearce SL, Li F, Coppin C, Schiffer M, Shirriffs J, et al. Genomic changes

associated with adaptation to arid environments in cactophilic *Drosophila* species. BMC

Genomics. 2019;20(1):52.

85.   Coronado-Zamora M, González J. Transposons contribute to the functional

diversification of the head, gut, and ovary transcriptomes across *Drosophila* natural strains.

Genome Res. 2023;33(9):1541–53.

86.   Cosby RL, Judd J, Zhang R, Zhong A, Garry N, Pritham EJ, et al. Recurrent evolution

of   vertebrate   transcription   factors   by   transposase   capture.   Science.

2021;371(6531):eabc6405.

87.   Pal D, Patel M, Boulet F, Sundarraj J, Grant OA, Branco MR, et al. H4K16ac activates

the transcription of transposable elements and contributes to their cis-regulatory function.

Nat Struct Mol Biol. 2023;30(7):935–47.

88.   Lippman Z, Gendrel AV, Black M, Vaughn MW, Dedhia N, Richard McCombie W, et

al. Role of transposable elements in heterochromatin and epigenetic control. Nature.

2004;430(6998):471–6.

89.   Guio L, Vieira C, González J. Stress affects the epigenetic marks added by natural

transposable element insertions in *Drosophila melanogaster*. Sci Rep. 2018 Aug

15;8(1):12197.

90.   Coronado-Zamora M, González J. The epigenetics effects of transposable elements

are context dependent and not restricted to gene silencing. bioRxiv; 2023. Available from:

https://www.biorxiv.org/content/10.1101/2023.11.27.568862v1

91.   Fonseca PM, Moura RD, Wallau GL, Loreto ELS. The mobilome of *Drosophila*

incompta, a flower-breeding species: comparison of transposable element landscapes

among generalist and specialist flies. Chromosome Res. 2019;27(3):203–19.

92.   Mérel V, Gibert P, Buch I, Rodriguez Rada V, Estoup A, Gautier M, et al. The

Worldwide Invasion of *Drosophila suzukii* Is Accompanied by a Large Increase of

Transposable Element Load and a Small Number of Putatively Adaptive Insertions.

Molecular Biology and Evolution. 2021;38(10):4252–67.

93.   Marin P, Jaquet A, Picarle J, Fablet M, Merel V, Delignette-Muller ML, et al.

Phenotypic and Transcriptomic Responses to Stress Differ According to Population

Geography in an Invasive Species. Genome Biology and Evolution. 2021;13(9):evab208.

94. Zidi M, Klai K, Confais J, Chénais B, Caruso A, Denis F, et al. Genome-Wide Screening

of Transposable Elements in the Whitefly, *Bemisia tabaci* (Hemiptera: Aleyrodidae),

Revealed Insertions with Potential Insecticide Resistance Implications. Insects.

2022;13(5):396.

95. Arribas YA, Baudon B, Rotival M, Suárez G, Bonté PE, Casas V, et al. Transposable

element exonization generates a reservoir of evolving and functional protein isoforms. Cell.

2024;187(26):7603-7620.e22.

96. Harigaya Y, Parker R. No‐go decay: a quality control mechanism for RNA in

translation. WIREs RNA. 2010;1(1):132–41.

97. Vasudevan S, Peltz SW, Wilusz CJ. Non‐stop decay—a new mRNA surveillance

pathway. BioEssays. 2002;24(9):785–8.

98. Hug N, Longman D, Cáceres JF. Mechanism and regulation of the nonsense-

mediated decay pathway. Nucleic Acids Res. 2016;44(4):1483–95.

99. Whisson SC, Vetukuri RR, Avrova AO, Dixelius C. Can silencing of transposons

contribute to variation in effector gene expression in Phytophthora infestans? Mobile

Genetic Elements. 2012;2(2):110–4.

100. Raffaele S, Farrer RA, Cano LM, Studholme DJ, MacLean D, Thines M, et al. Genome

Evolution Following Host Jumps in the Irish Potato Famine Pathogen Lineage. Science.

2010;330(6010):1540–3.

101.   De Panis DN, Padró J, Furió-Tarí P, Tarazona S, Milla Carmona PS, Soto IM, et al.

Transcriptome modulation during host shift is driven by secondary metabolites in desert

Drosophila. Molecular Ecology. 2016;25(18):4534–50.

102.   Knolhoff LM, Heckel DG. Behavioral Assays for Studies of Host Plant Choice and

Adaptation in Herbivorous Insects. Annual Review of Entomology. 2014;59(Volume 59,

2014):263–78.

103.   Zouros E, d'Entremont CJ. Sexual Isolation Among Populations of *Drosophila*

mojavensis: Response to Pressure from a Related Species. Evolution. 1980;34(3):421–30.

104.   Markow TA. Sexual Isolation among Populations of *Drosophila mojavensis*. Evolution.

1991;45(6):1525–9.

105.   Howard RW, Blomquist GJ. Ecological, behavioral, and biochemical aspects of the

insect hydrocarbons. Annual Review of Entomology. 2005;50:371–93.

106.   Stennett MD, Etges WJ. Premating Isolation Is Determined by Larval Rearing

Substrates in Cactophilic *Drosophila mojavensis*. III. Epicuticular Hydrocarbon Variation Is

Determined by Use of Different Host Plants in *Drosophila mojavensis* and *Drosophila*

arizonae. J Chem Ecol. 1997;23(12):2803–24.

107.   Havens JA, Etges WJ. Premating isolation is determined by larval rearing substrates

in cactophilic *Drosophila mojavensis*. IX. Host plant and population specific epicuticular

hydrocarbon expression influences mate choice and sexual selection. Journal of

Evolutionary Biology. 2013;26(3):562–76.

108.   Romero-Soriano V, Burlet N, Vela D, Fontdevila A, Vieira C, García Guerreiro MP.

Drosophila Females Undergo Genome Expansion after Interspecific Hybridization. Genome

Biol Evol. 2016;8(3):556–61.

109.   Bodelón A, Fablet M, Veber P, Vieira C, García Guerreiro MP. High Stability of the

Epigenome in *Drosophila* Interspecific Hybrids. Genome Biology and Evolution.

2022;14(2):evac024.

110.   Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using

repeat graphs. Nat Biotechnol. 2019;37(5):540–6.

111.   Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.

2018;34(18):3094–100.

112.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. arXiv:13033997 [q-bio]. 2013; Available from: http://arxiv.org/abs/1303.3997

113.   Cabanettes F, Klopp C. D-GENIES: dot plot large genomes in an interactive, efficient

and simple way. PeerJ. 2018;6:e4958.

114.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078–9.

115.   Krumsiek J, Arnold R, Rattei T. Gepard: a rapid and sensitive tool for creating

dotplots on genome scale. Bioinformatics. 2007;23(8):1026–8.

116.   Alonge M, Soyk S, Ramakrishnan S, Wang X, Goodwin S, Sedlazeck FJ, et al. RaGOO:

fast and accurate reference-guided scaffolding of draft genomes. Genome Biol.

2019;20(1):224.

117.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics. 2015;31(19):3210–2.

118.   Shumate A, Salzberg SL. Liftoff: accurate mapping of gene annotations. Valencia A,

editor. Bioinformatics. 2021;37(12):1639–43.

119.   Kim BY, Wang JR, Miller DE, Barmina O, Delaney E, Thompson A, et al. Highly

contiguous assemblies of 101 drosophilid genomes. eLife. 2021;10(e66405):33.

120.   Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, et al. BRAKER3: Fully

automated genome annotation using RNA-seq and protein evidence with GeneMark-ETP,

AUGUSTUS, and TSEBRA. Genome Research. 2024;34(5):769-777.

121.   Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biol. 2019;20(1):238.

122.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND.

Nat Methods. 2015;12(1):59–60.

123.   Altschul S. Gapped BLAST and PSI-BLAST: a new generation of protein database

search programs. Nucleic Acids Research. 1997;25(17):3389–402.

124.   Shumate A, Salzberg SL. Liftoff: accurate mapping of gene annotations. Valencia A,

editor. Bioinformatics. 2021;37(12):1639–43.

125.   Vanderlinde T, Dupim EG, Nazario-Yepiz NO, Carvalho AB. An Improved Genome

Assembly for *Drosophila navojoa*, the Basal Species in the mojavensis Cluster. Journal of

Heredity. 2019;110(1):118–23.

126.   Hu J, Fan J, Sun Z, Liu S. NextPolish: a fast and efficient genome polishing tool for

long-read assembly. Berger B, editor. Bioinformatics. 2020;36(7):2253–5.

127.   Banho CA, Mérel V, Oliveira TYK, Carareto CMA, Vieira C. Comparative

transcriptomics between *Drosophila mojavensis* and *D. arizonae* reveals transgressive gene

expression and underexpression of spermatogenesis-related genes in hybrid testes. Sci

Rep. 2021;11(1):9844.

128.   Moreyra NN, Almeida FC, Allan C, Frankel N, Matzkin LM, Hasson E. Phylogenomics

provides insights into the evolution of cactophily and host plant shifts in *Drosophila*.

Molecular Phylogenetics and Evolution. 2023;178:107653.

129.   Dainat J. AGAT: Another Gff Analysis Toolkit to handle annotations in any GTF/GFF

format. Zenodo. 2023; Available from: https://zenodo.org/records/8178877

130.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078–9.

131.   Suyama M, Torrents D, Bork P. PAL2NAL: robust conversion of protein sequence

alignments   into   the   corresponding   codon   alignments.   Nucleic   Acids   Research.

2006;34(2):W609–12.

132.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,

et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the

Genomic Era. Molecular Biology and Evolution. 2020;37(5):1530–4.

133.   Yang Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Molecular Biology

and Evolution. 2007;24(8):1586–91.

134.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014;30(15):2114–20.

135.   Baril TJ, Galbraith JD, Hayward A. Earl Grey: a fully automated user-friendly

transposable element annotation and analysis pipeline. Molecular Biology and Evolution.

2024;41(4)msae068.

136.   Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2

for automated genomic discovery of transposable element families. Proc Natl Acad Sci USA.

2020;117(17):9451–7.

137.   Bao Z, Eddy SR. Automated De Novo Identification of Repeat Sequence Families in

Sequenced Genomes. Genome Res. 2002;12(8):1269–76.

138.   Price AL, Jones NC, Pevzner PA. De novo identification of repeat families in large

genomes. Bioinformatics. 2005;21(Suppl 1):i351–8.

139.   Li Z, Gilbert C, Peng H, Pollet N. Discovery of numerous novel *Helitron* -like elements

in eukaryote genomes using HELIANO. Nucleic Acids Research. 2024;52(17):e79–e79.

140.   Platt RN, Blanco-Berdugo L, Ray DA. Accurate Transposable Element Annotation Is

Vital When Analyzing New Genome Assemblies. Genome Biol Evol. 2016;8(2):403–10.

141.   Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Molecular Biology and Evolution.

2013;30(4):772–80.

142.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25(15):1972–

3.

143.   Rice P, Longden I, Bleasby A. EMBOSS: The European Molecular Biology Open

Software Suite. Trends in Genetics. 2000;16(6):276–7.

144.   Benson G. Tandem repeats finder: a program to analyze DNA sequences. Nucleic

Acids Research. 1999;27(2):573–80.

145.   Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. The Dfam community resource of

transposable element families, sequence models, and genome annotations. Mobile DNA.

2021;12(1):2.

146.   SMIT, Arian FA. RepeatMasker Open 3.0. 2004.

147.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics. 2013;29(1):15–21.

148.   Wong WY, Simakov O. RepeatCraft: a meta-pipeline for repetitive element de-

fragmentation and annotation. Hancock J, editor. Bioinformatics. 2019;35(6):1051–2.

149.   Flutre T, Permal E, Quesneville H. Transposable Element Annotation in Completely

Sequenced Eukaryote Genomes. In: Grandbastien MA, Casacuberta JM, editors. Plant

Transposable Elements. Berlin, Heidelberg: Springer Berlin Heidelberg; 2012. p. 17–39.

(Topics in Current Genetics; vol. 24). Available from: https://link.springer.com/10.1007/978-

3-642-31842-9_2

150.   Gel B, Díez-Villanueva A, Serra E, Buschbeck M, Peinado MA, Malinverni R. regioneR:

an R/Bioconductor package for the association analysis of genomic regions based on

permutation tests. Bioinformatics. 2015;32(2):289-291.

151.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 2010;26(6):841–2.

152.   Grant CE, Bailey TL, Noble WS. FIMO: scanning for occurrences of a given motif.

Bioinformatics. 2011;27(7):1017–8.

153.   Bailey TL, Johnson J, Grant CE, Noble WS. The MEME Suite. Nucleic Acids Research.

2015;43(W1):W39–49.

154.   Fornes O, Castro-Mondragon JA, Khan A, van der Lee R, Zhang X, Richmond PA, et

al. JASPAR 2020: update of the open-access database of transcription factor binding

profiles. Nucleic Acids Research. 2019;48(D1):87-92.

155.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-

throughput sequencing data. Bioinformatics. 2015;31(2):166–9.

156.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

157.   Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M. Transcriptome

assembly from long-read RNA-seq alignments with StringTie2. Genome Biol.

2019;20(1):278.

158.   Stovner EB, Sætrom P. PyRanges: efficient comparison of genomic intervals in

Python. Bioinformatics. 2020;36(3):918–9.

1461