Cross-feeding maintains diversification in an experimentally evolved microbial community

Zahraa Al-Tameemi[1], Alejandra Rodriguez-Verdugo[1*]

[1] Department of Ecology and Evolutionary Biology, University of California, Irvine, Irvine, CA.

* Corresponding author:

Alejandra Rodríguez-Verdugo

e-mail: alejanr1@uci.edu

Competing interests

The authors declare that they have no conflict of interest.

Running title: Bacterial diversification persists only in a community.

Abstract

Microbial communities are incredibly diverse. Yet, the eco-evolutionary processes originating

and maintaining this diversity remain understudied. Here, we investigate the effect of

interspecies interactions on the generation and maintenance of diversity in microbial

communities. We studied a commensal interaction in a community of *Acinetobacter johnsonii*

and *Pseudomonas putida* where the latter species cross-feeds on leaked resources produced by *A.*

johnsonii. By evolving four experimental replicates in monoculture and co-culture for 200

generations, we observed the diversification of *P. putida* into two distinct morphotypes that

differed from their ancestor by single-point mutations. One of the most prominent mutations hit

the *fleQ* gene encoding the master regulator of flagella and biofilm formation. We experimentally

confirmed that *fleQ* mutants were unable to swim and formed less biofilm than their ancestor, but

they also produced higher yields. Interestingly, the *fleQ* genotype swept to fixation in

monocultures but not in co-cultures. In co-cultures, the two lineages stably coexisted for

approximately 150 generations. We hypothesized that *A. johnsonii* modulates the coexistence of

the two lineages through frequency-dependent selection. However, invasion experiments in

monoculture and co-culture did not support this hypothesis, suggesting other mechanisms

maintain coexistence. Overall, our study shows that interspecies interactions play an important

role in maintaining the diversity of newly formed lineages in microbial communities.

Importance

In nature, bacteria live in microbial communities and interact with other species, for example,

through the exchange of resources leaked into the external environment (i.e., cross-feeding

interactions). The role that these cross-feeding interactions play in the formation of new bacterial

'species' remains unexplored. Using a simple bacterial system in which one species cross-feeds

resources to a second species (commensal species), we showed that the commensal species

diversified into two subpopulations that persisted only when the cross-feeder partner was

present. We further observed loss-of-function mutations in flagellar genes in both monocultures

and co-cultures. Instead, mutants that retained the ability to swim and form biofilms were

exclusively observed in co-cultures. These findings suggest that motility and surface attachment

traits are important during cross-feeding, even in a well-shaken environment. Given that nutrient

leakage is pervasive in microbial communities, the findings from this study have the potential to

extend beyond our specific bacterial system.

Keywords: interspecies interactions, cross-feeding, evolutionary tradeoffs, maintenance of

diversity, experimental evolution, synthetic communities

Introduction

Long-term experimental evolution studies have demonstrated that microbial populations can undergo significant adaptation and diversification over thousands of generations (1). Such adaptation involves the acquisition of new genetic mutations that enhance survival and reproduction in specific environments. Often, these adaptive mutations sweep in the population, resulting in a loss of genetic polymorphism (2, 3). However, polymorphisms are sometimes maintained, leading to the formation of new lineages. For example, in the bacterium *Escherichia coli*, acetate specialists evolved from a clonal ancestor that uses glucose and releases acetate as a waste product, resulting in a polymorphic population (4, 5). Spatial structure can also drive microbial diversification by creating distinct ecological niches, which leads to more genetic polymorphism within populations (6–8). Additionally, ecological mechanisms such as localized dispersal and interactions can maintain diversity by decreasing negative interactions (9). In these previous examples of diversification, polymorphism is maintained by frequency-dependent selection, which is often linked to fitness tradeoffs (10). During frequency-dependent selection, a rare mutant rises in frequency in a resident population, but it loses its fitness advantage when it becomes common (11).

Despite the numerous studies on microbial diversification, the role that other species in a community play in diversification has been largely understudied. When interspecies interactions have been studied in the context of diversification, they have mostly focused on negative interactions such as those between predators and bacteria (12, 13). However, interspecific microbial interactions can also be positive. For example, during cross-feeding, microbes exchange metabolites that benefit each other. This phenomenon is prevalent across different

systems (14). However, it remains unclear how positive interspecific interactions such as cross-

feeding influence the generation and maintenance of biodiversity.

To address this question, we are using a synthetic microbial community composed of

Acinetobacter johnsonii C6 and *Pseudomonas putida* KT2440. These bacterial species have been

isolated from a creosote-polluted aquifer and have naturally evolved to biodegrade aromatic

compounds that make up a significant portion of environmental pollutants (15, 16). In addition to

its importance for bioremediation, this synthetic community is considered a model system for

studying metabolic and commensal interactions (17–19). The commensalism is based on a

unidirectional cross-feeding interaction between *A. johnsonii* and *P. putida*. *A. johnsonii*

consumes benzyl alcohol and produces benzoate – an intermediate by-product of benzyl alcohol

oxidation – which leaks into the external environment where *P. putida* uses it as a source of

carbon and energy. This cross-feeding interaction has been shown to be stable over ecological

and evolutionary timescales (20, 21).

In this study, we build on a previous evolution experiment in which *P. putida* evolved

independently and in conjunction with *A. johnsonii* over 200 generations (21). Over the course of

this experiment, we observed that populations of *P. putida* diversified into two morphotypes, but

this diversification was maintained exclusively in the presence of *A. johnsonii*. To understand

how diversity is maintained in the presence of a commensal interaction, we used both genotypic

and phenotypic data to address the following questions: How does trait evolution differ in the

presence and absence of *A. johnsonii*? What types of mutations underpin the observed

phenotypic diversity? How do these mutations affect physiology and tradeoffs? Finally, what

mechanism underlies the persistence of this diversity in the co-culture conditions?

Materials and Methods

Growth conditions

All strains were cultured in 10 mL FAB minimal medium (1 mM $MgCl_2$, 0.1 mM $CaCl_2$, 0.003 mM $FeCl_3$, 15 mM $(NH_4)_2SO_4$, 33 mM Na_2HPO_4, 22 mM KH_2PO_4, and 50 mM NaCl) with either 0.6 mM of benzoate or 0.6 mM benzyl alcohol as the sole carbon source. The media was run through a 0.2 μm filter and stored in glass treated to remove assimilated organic carbon (AOC). AOC-free glassware was prepared as in Rodriguez-Verdugo et al. (2019) (20). Before running experiments, all strains were acclimated by inoculating 10 μl of frozen glycerol stocks in FAB minimal medium supplemented by either 0.6 mM benzoate or 0.6 mM benzyl alcohol and incubated for 24 hours at 26 °C with constant shaking (150 r.p.m.).

Evolution experiment

The evolution experiment had been previously described in Rodriguez-Verdugo and Ackermann (2021) (21). Briefly, the experiment consisted of evolving four replicates of *P. putida* in monoculture, four replicates of *A. johnsonii* in monoculture, and four replicates of *P. putida* with *A. johnsonii* (i.e., co-cultures) for 30 days, which equates to approximately 200 generations (~6.6 generations per day). All cultures were grown in batch culture with FAB minimal media and supplemented with 0.6 mM benzyl alcohol. The experimental lines were established from four ancestors of *P. putida* and *A. johnsonii*, isolated from single clones (Fig. S1). Thus, the ancestors diverged by a few generations when the evolution experiment was started. During the evolution experiment, the population densities for the monoculture and the co-culture from replicate #2 were estimated at the end of each day by plating cultures in LB agar

with streptomycin (64 µg/ml) and in LB agar with gentamycin (10 µg/ml) to track *A. johnsonii*

and *P. putida*, respectively (21).

Growth curves and resource use traits

To characterize the phenotypic diversification of *P. putida* in monoculture and co-culture,

we performed growth curve experiments on 30 single colonies isolated from each of the eight

experimental lines and from four ancestral populations. Briefly, bacteria were grown in FAB

medium supplemented with 0.6 mM benzoate for 24 hours and were then plated on Lysogeny

Broth (LB) agar and incubated for another 24 hours at 26 °C. We randomly selected 30 colonies

off the plate and resuspended them into wells filled with 200 µL 1% $MgSO_4$ in a 96-well plate.

We then transferred 2 µL of the diluted cells and transferred them to another 96-well plate

containing 198 µL of FAB supplemented with 0.6 mM benzoate. The plate was sealed with

vacuum grease to reduce evaporation. The plate was incubated in a photospectrometer plate

reader (Epoch2™, Agilent™) at 30 °C and continuous linear shaking. Optical density (OD)

measurements were taken every 10 minutes for 24 hours. From these growth curves, we

estimated four quantitative traits involved in resource use: the maximum growth rate in

exponential phase (μ_{max}), the yield (Y) defined as the amount of biomass produced per unit of

resource, the maximum uptake rate (V) defined as the maximum rate of resource consumption

per hour and the half-saturation constant (K) defined as the resource concentration supporting

half-maximum uptake rate. Parameters were estimated using the same method outlined in

Rodriguez-Verdugo et al. (2019) (20) using MATLAB (version R2017a, MathWorks). To

quantify the evolution of phenotypic traits, we compared the trait distributions from ancestral

populations (n = 30) to those from evolved populations (n = 30) using a two-tailed *t*-test (R,

version 4.3).

Genome resequencing and variant calling

We extracted DNA from the ancestors' clones, monoculture and co-culture populations,

and single clones from large and small morphotypes of all four experimental lines. Single clones

were revived from glycerol stocks in 3 mL of LB broth and grown overnight at 26 °C with

constant shaking (150 r.p.m.). The overnight cultures from monoculture and co-cultures whole

populations were grown FAB with 0.6 mM benzoate supplemented with gentamycin (10

µg/mL). We isolated and purified genomic DNA from 1 mL of the overnight cultures using the

Wizard DNA purification kit (Promega). Short-read sequencing (2 x 150bp paired-end reads)

was conducted on a NovaSeq6000 S4 flow cell (UCI Genomics High-Throughput Facility).

These sequences can be accessed at the NCBI Sequence Read Archive under Bioproject ID

number PRJNA623337.

All sequences were aligned to the reference genome of *P. putida* KT2440 (reference

NC_002947) using the computational pipeline *breseq* (version 0.36.1) and utility program

GDtools (22). We used consensus mode and polymorphism mode to call variants for clones and

whole populations, respectively. Each sample was sequenced to a depth of ~600X coverage.

Single-nucleotide polymorphisms in genic regions at or above 10% and passing the Fisher's

exact test for biased strand distribution were considered. *De novo* non-synonymous mutations

were defined as uniquely observed in the evolved sample and not the ancestral clone.

Physiological tradeoffs

Biofilm formation - We measured biofilm formation in the *P. putida* ancestors and the

evolved morphotypes using the method described in Martínez-García et al. (2014) (23). Samples

were inoculated into FAB media supplemented with 0.6 mM benzoate and incubated for 24

hours at 26 ℃ with constant shaking (150 r.p.m.). The tubes were then vortexed to homogenize

the culture and the OD_{600} was adjusted to 0.5. We added 2 µL of the adjusted acclimated culture

to a 96-well plate containing 198 µL of FAB and 0.6 mM benzoate and was incubated at 26 ℃

with no shaking. Each sample was inoculated into 5 wells that serve as technical replicates. After

24 hours, the plate was removed from the incubator, and the OD_{600} was measured using the plate

reader. The media with planktonic cells was discarded by gently tapping the plate and was

washed 3 times with water. The cells attached to the plate were stained by adding 200 µL of

0.1% (w/v) crystal violet into each well and were incubated on the bench top for 30 minutes. We

then disposed of the crystal violet and gently washed the plate 2-3 times with deionized water.

The plate was left to completely dry. We added 200 µL of 33% (w/v) acetic acid to each well,

pipetting up and down until it was completely homogenized. The OD_{595} was measured, and an

index of biofilm formation was calculated, which was the crystal violet stained biofilm

(OD_{595}) divided by the initial culture with planktonic cells (OD_{600}). The higher the biofilm index

value, the greater the biofilm formation. These experiments were replicated three times. We

conducted a two-tailed *t*-test to determine if there were significant differences in biofilm

formation in the evolved morphotypes (n = 3) compared to the ancestor (n = 3).

Motility - We performed a swimming assay on the ancestors and evolved morphotypes of

P. putida using 0.3% (w/v) soft agar (23, 24). We inoculated 2 µL of an acclimated culture onto

FAB soft agar supplemented with 0.6 mM benzoate and incubated the plates at 30 ℃. We

measured the halo around the initial inoculum after 24 hours to determine whether the evolved

morphotypes' swimming abilities differed from the ancestors' swimming abilities. A larger halo

indicates higher motility, while the absence of a halo suggests a lack of flagellar motility.

Reciprocal invasion assays

To test if frequency-dependent selection is a possible mechanism for maintenance of the

emergent diversity, we performed reciprocal invasion experiments. These experiments consisted

of culturing two morphotypes (large and small) at a frequency of 1 to 9 to see if they can invade

when rare in the presence and absence of *A. johnsonii.*

The two morphotypes of *P. putida* and *A. johnsonii* were revived from a freezer stock of

replicate 2 and were allowed to acclimate for 24 hours. The OD was then adjusted to 1.0 for

both morphotypes to standardize the starting ratio. For the condition without *A. johnsonii* (i.e.,

monoculture), 10 µL of the large morphotype and 90 µL of the small morphotype were

inoculated into 9.9 mL of fresh media supplemented with 0.6 mM benzyl alcohol and 0.06 mM

benzoate. The concentration of supplemented benzoate was selected based on the amount of

benzoate excreted to the external environment by *A. johnsonii* and measured by high-

performance liquid chromatography[20]. The starting volumes were reversed when the small

morphotypes started from a rare frequency. For the condition with 100 µl of *A. johnsonii* (i.e. co-

culture)*,* no external benzoate was added to the medium, and the large and small morphotypes

were inoculated at 1:9 and 9:1 ratios as previously described. Co-cultures and monocultures were

grown at 26°C with constant shaking (150 r.p.m.). After 24 hours, 1% of culture was transferred

into fresh media and these daily transfers were continued for three days. We counted colony-

forming units (CFU) at the start of the experiment (T0) and the end of each transfer cycle by

plating cultures onto LB agar plates supplemented with gentamicin (10 µg mL^{-1}). We included

two technical replicates per experiment, and each experiment was replicated three times.

The Malthusian parameters for the rare and common morphotypes were calculated from

cell densities at the initial and final time points, $m = \ln(N_f/N_0)$, with N_0 and N_f the initial and

final density after 24 hours, respectively. From the Malthusian parameters, we calculated the

relative growth rate of a morphotype against its competitor when starting from rare. For example,

$S_{i\text{-}rare} = m_{i\text{-}rare} - m_j$, for a pair of competitors i and j as in Zhao et al. (2016) (25). If the value of

S_{rare} for a given morphotype was significantly larger than zero, the morphotype could invade

when rare. If both the large and small morphotypes had a positive S_{rare} value, the pair was

considered to stably coexist through frequency-dependent selection.

Results

The evolution experiment was started with four experimental lines (replicates) of *P.*

putida in monoculture and four lines in co-culture with *A. johnsonii* grown in minimal growth

media supplemented with benzyl alcohol as the sole carbon source (Fig. S1). Following this

experiment, we conducted whole genome sequencing and identified *de novo* mutations in the

evolved lines. The effect of *de novo* mutations on yield, motility, and biofilm formation was then

evaluated by pairwise comparisons of the evolved lines to their ancestors. Finally, we tested if

frequency-dependent selection was a possible mechanism for maintaining diversity.

Diversification evolved in co-culture but not in monoculture.

After 200 generations of evolution, we observed diversification of *P. putida* in terms of

colony morphology relative to the ancestor (Fig. 1A). All four replicates were bimodal for colony

size and consisted of one cluster of strains that formed larger colonies on LB agar plates

compared to the ancestor (referred as 'large morphotype'), and one cluster that formed colonies

the size of the ancestor (referred as 'small morphotype'). This diversification in colony size was

only observed when *P. putida* was evolved in co-culture with *A. johnsonii*. In monocultures,

populations only formed larger colonies than the ancestor.

Next, we quantitively characterized this diversification in four traits involved in resource

use. In all monoculture and co-culture replicates, the phenotypic variance for each trait changed

compared to the variance of the ancestral population (Figs. S2 and S3). Changes in yield were

the most pronounced. We found that *P. putida* grown in co-culture with *A. johnsonii* had a

different distribution of yields compared to monoculture (Fig. S2). For example, co-cultures of

replicate #2 showed a bimodal distribution, with one subpopulation having a similar yield to the

ancestor and the other having a higher yield to the ancestor (Fig. 1B, right panel). On the other

hand, the monoculture had a distribution shifted towards higher yield (Fig. 1B, left panel). We

confirmed that the large morphotypes previously observed on LB agar plates had higher yields

compared to their ancestors, and the small morphotypes had similar yields to their ancestors. For

example, large morphotypes from replicate #2 had, on average, a 3.4-fold higher yield than the

ancestor (0.068 ± 0.001 and 0.020 ± 0.001 OD per mM for the large and ancestral morphotypes,

respectively; t-test, $p < 2.2*10^{-16}$). The small morphotypes achieved a yield of 0.023 ± 0.002 OD

per mM, which is comparable to that of the ancestor (t-test, $p = 0.122$).

Finally, leveraging the fact that large morphotypes with higher yield formed larger colonies

on LB agar plates that could be distinguished from the ancestral morphotype, we tracked the

evolution of phenotypic diversification for one of the replicates (replicate #2) during our

experiment. After approximately 40 and 70 generations of evolution, we observed the emergence

of a large morphotype in monoculture and co-culture, respectively (Fig. 2). In co-culture, the

large and small morphotypes coexisted for the remaining 130 generations. Instead, in

monoculture, the large morphotype rapidly outcompeted the ancestral morphotype, driving it to

extinction after 140 generations. Thus, diversification only persisted in co-culture.

Fig. 1. *P. putida* population evolved in monoculture is monomorphic, while the population

evolved in co-culture diversified into two sub-populations after 200 generations. A) Agar

plates illustrating the difference in colony morphologies between the evolved population in

monoculture (left panel) and the evolved population in co-culture (right panel) for replicate #2.

B) In monoculture, *P. putida*'s population from replicate #2 evolved higher yield (t-test, $p <$

$2.2*10^{-16}$). In co-culture, *P. putida*'s population evolved bimodality (Hartigans' dip test for

multimodality, $p < 2.2*10^{-16}$). The results from the other replicates and parameters are available

in Fig. S2 and Fig. S3.

Fig. 2. Diversification was maintained in co-culture but not in monoculture. The large

morphotype appears in the monoculture after approximately 50 generations and outcompetes the

ancestral morphotype. In co-culture, both morphotypes coexist until the end of the experiment.

Population density (CFU mL^{-1}) trajectories from replicate #2 are shown over 200 generations. *A.*

johnsonii's densities are from (21). Each circle represents the final density of each species and

morphotypes after daily cycles of growth; lines connecting circles help visualization and do not

have any biological meaning.

Morphological diversification in *P. putida* is based on genetic diversification

To determine if the observed phenotypic diversity (Fig. 1) reflected specific mutations,

we performed whole population sequencing and identified *de novo* nonsynonymous mutations in

the evolved lines (Table S1). We detected single substitution mutations in the *fleQ* gene at

relatively high frequencies in evolved co-cultures from replicates #1 and #2 (Table 1). This gene

codes for flagellar transcriptional regulator FleQ, which has been shown to affect motility and

biofilm formation (26–28). Sequencing of single clones isolated from these populations

confirmed that the *fleQ* mutations were associated with a large morphotype. Instead, the small

morphotypes had mutations in genes other than flagellar genes in three out of four replicates

(Table S1), but most of these genes were unannotated. One exception was the small morphotype

from replicate #3, which had a 1 bp deletion in the *oplA* gene and was present at a 12.9%

frequency in the population (Table S1). The *oplA* gene codes for 5-oxoprolinase A, which has

been associated with the catabolism of amino acid derivatives (29, 30), but its functional

relevance in our experimental conditions is unclear.

The evolved monocultures also had mutations in flagellar genes. For example, replicate

#2 had a mutation in the *fleQ* gene at 100% frequency in the population (Table 1). This is

consistent with our observation that the large morphotype carrying the *fleQ* mutation was fixed

in the population (Fig. 2). Finally, we detected other flagellar mutations, including a mutation in

the *flgH* gene, coding for flagellar basal body L-ring protein FlgH in monoculture replicate #4

(Table 1). Thus, we infer that mutations in either *fleQ* or *flgH* lead to the enlarged morphotype.

Table 1. Frequency of *de novo* flagellar mutations in the populations of *P. putida* in

monoculture and co-culture after 200 generations.

Experimental populations	Gene	Mutation	Frequency
Co-culture 1	*fleQ*	A226V (GCC→GTC) *	23.1%
Co-culture 2	*fleQ*	L353Q (CTG→CAG) *	35.6%
Co-culture 3	-		-
Co-culture 4	-		-
Monoculture 1	-		-
Monoculture 2	*fleQ*	2 bp→ C †	100%

Monoculture 3	-		-
Monoculture 4	*flgH*	+C	100%

*Same as found in single clones of the large morphotype.
† Deletion of first base and substitution of second base with C.

Mutations in the large morphotype led to loss of motility and biofilm formation

We previously confirmed that the large morphotypes - commonly associated with *fleQ*

mutations - had significantly higher yields than their ancestors as well as changes in other

resource use traits (Figs. S2 and S3). To explore whether *fleQ* mutations affected other traits, we

explored the mutants' abilities to swim and form biofilms. The motility assay showed a complete

lack of motility in the *fleQ* mutants compared to their ancestors (Fig. 3). Other large

morphotypes not carrying the *fleQ* mutations (e.g. from monoculture replicate #4) had a

reduction in swimming, possibly due to a mutation in the flagellar component *flgH*. Replicate #4

also had a small morphotype colony that lacked motility, but the cause for the lack of motility

remains unclear.

The *fleQ* mutation also resulted in a significant reduction in biofilm formation for the

large morphotype (Fig. 4). The large morphotype, isolated from co-cultures #1 and 2, and

monoculture #2, exhibited significant reductions in biofilm formation compared to their

ancestors (t-test, *p*-values of 0.003, 0.023, and 0.008, respectively). The *flgH* mutations (large

morphotype from monoculture #4) also had reduced biofilm formation but the difference was

marginally significant (t-test, $p = 0.068$).

Fig. 3. **Swimming assay shows the loss of flagellar motility in the large morphotypes with flagellar mutations.** All samples were spotted on 0.3% (w/v) FAB soft agar. The absence of a halo around the initial inoculum indicates the loss of flagellum-driven motility.

Fig. 4. Biofilm assay shows that *fleQ* mutants produce less biofilm than the ancestor. Biofilm index (CV/OD_{600}) after a 24-hour incubation period. All *fleQ* mutants (i.e. large morphotypes of co-cultures 1 and 2 and monoculture 2) had a significant reduction in the biofilm index compared to their respective ancestors. Note: $* = 0.01 < p < 0.05$, $** = p < 0.01$.

Frequency-dependent selection does not explain the coexistence of morphotypes

We previously observed that the large and small morphotypes stably coexisted for 150 generations in co-culture from replicate#2 (Fig. 2). To test if frequency-dependent selection is a possible mechanism for the stable coexistence of these two morphotypes and if the presence/absence of *A. johnsonii* influence this selection, we performed a reciprocal invasion experiment. The reciprocal invasion assay did not show significant differences in the rare morphotype's ability to invade depending on whether *A. johnsonii* was present (Table 2). One exception was the large morphotype, which had a significantly positive relative growth rate when rare (S_{rare}) in co-culture, meaning it could invade when rare (i.e., at 10% frequency). None of the other morphotypes invaded from rare (Table 2). We confirmed that this was the case by tracking the frequencies of the large and small morphotypes over two additional days (Fig. S4). Overall, these results suggest that frequency-dependent selection is not the mechanism driving coexistence.

Table 2. The relative growth rate when rare was not affected by growth conditions (monoculture or co-culture). The values ± standard deviations were obtained from four to six replicates.

Strain	Condition	Relative growth rate when rare (S_{rare})	p-value*
Small morphotype	Monoculture	0.65 ± 0.26	0.470
	Co-culture	0.19 ± 0.24	0.468
Large morphotype	Monoculture	0.58 ± 0.52	0.326
	Co-culture	0.94 ± 0.18	**0.019**

* One sample t-test testing whether S_{rare} is different than zero.

Discussion

The results from this study show that diversity can arise in commensal communities, even

in a homogeneous environment with low resources. Diversity arose in both monoculture and co-

culture conditions; however, it persisted only in co-culture conditions. In monoculture

conditions, the ancestral morphotype was outcompeted by a new mutant. This suggests that the

presence of *A. johnsonii* and, hence, positive interspecific interactions (cross-feeding) play a role

in the maintenance of the emerging diversity.

We found that the phenotypic diversity of *P. putida* is based on genetic diversity. A

mutation in *fleQ* gene was detected in two out of the four cocultures and one monoculture that

corresponds to the large morphotype. The *fleQ* mutation led to an inability to swim and a

significant decrease in biofilm formation. Mutants also had an increase in yield compared to their

respective ancestors, which explains why they form larger colonies than the ancestor when plated

in LB agar. This is unsurprising as the loss of flagellar function is a common response to shaking

environments (31–35). The loss of flagellar function alleviates a large metabolic burden that can

be invested in biomass production while still being able to passively move due to the continuous

shaking of our culture conditions (23). The small morphotype also accumulated mutations that

are at low frequency in the population (Table S1). However, these mutations are in genes that are

not annotated, and their physiological effects on our culture conditions are unknown.

Even though our system lacked spatial structure initially, cross-feeding can lead to niche

construction. The excretion or leakage of nutritional by-products leads to the construction of new

niches, which can affect the generation and maintenance of biodiversity (36, 37). We hypothesize

that *A. johnsonii* creates transient patches of resources, and this spatiotemporal heterogeneity

maintains the diversity of newly formed lineages through a 'yield-foraging' tradeoff. That is, by

reducing the burden of flagellar motility, the *fleQ* mutants (large morphotypes) achieve higher

yields and have an advantage over the small morphotypes in monoculture (i.e., 'yield' strategy).

Instead, in co-culture, the small morphotypes retain flagellar motility and biofilm formation

abilities (Fig. 3 and 4), which may increase their abilities to cross-feed on the benzoate excreted

by *A. johnsonii* (i.e., 'foraging' strategy). A previous study showed that motility increases the

encounter rate between cross-feeding partners and cell-cell adhesion in a well-mixed

environment (38), supporting our hypothesis that *A. johnsonii* plays a key role in maintaining the

coexistence of *P. putida*'s lineages. Nevertheless, our findings did not support frequency-

dependent selection in the presence of *A. johnsonii* as the mechanism for maintaining lineages'

coexistence. It is possible that our representative clones did not capture the stable coexistence

dynamics. That is, maintenance of polymorphism through negative frequency-dependent

selection may only occur when considering the entire genetic diversity in *P. putida* populations.

Furthermore, other ecological and evolutionary processes could potentially maintain lineages

coexistence. One possibility is that small fitness differences between the two morphs could result

in neither of them able to invade, leading to neutral coexistence (39). Another possible

mechanism is multiple niche selection leading to the maintenance of stable polymorphisms in the

population (11, 40). An interesting avenue of research would be to explore further the

mechanisms by which *A. johnsonii* mediate the coexistence of newly formed lineages.

Our study has provided valuable insights into the emergence of diversity in commensal

microbial communities and suggests that interspecific interactions play a key role in the

diversification and maintenance of diversity. However, it is important to acknowledge certain

limitations in interpreting the results. First, while 200 generations was enough time to observe

diversification, we refrain from making a definitive conclusion about whether the diversity is

transient (short-lived) or if it can be sustained long-term. Another limitation in interpreting these

results is the lack of control for the benzoate excreted by *A. johnsonii*. Although benzoate was

not externally supplemented to the *P. putida* monocultures, they relied on residues of organic

carbon present in the culture medium, i.e., assimilable organic carbon or AOC (21). Thus, an

alternative hypothesis from the yield-foraging hypothesis is that diversification in the presence of

A. johnsonii was maintained through resource partitioning. To discern between these two

hypotheses, future evolution experiments should add an external source of benzoate to the

monocultures. Such experiments would allow us to determine how important it is that benzoate

is produced by *A. johnsonii* (i.e., a biogenic resource supply) versus being externally supplied to

the medium.

Data Availability

The genome sequences have been deposited at the National Center for Biotechnology

Information (NCBI) Sequence Read Archive (SRA) under BioProject ID number

PRJNA623337. All the other data generated or analyzed during this study are included in this

published article and its supplemental materials.

Acknowledgments

We thank Jennifer Martiny, Brandon Gaut, and the Rodriguez-Verdugo lab members for

providing useful feedback during this research. This work was supported by the National Science

Foundation under Grant No. DEB-2234627. ARV was supported by UC Irvine's School of

Biological Sciences.

References

1. **Lenski, RE, Travisano, M**. 1994. Dynamics of adaptation and diversification: a 10,000-generation experiment with bacterial populations. *Proc Natl Acad Sci U S A* 91:6808–6814.

2. **Atwood, KC, Schneider, LK, Ryan, FJ**. 1951. Periodic selection in Escherichia coli. *Proc Natl Acad Sci U S A* 37:146–155.

3. **Sniegowski, PD, Gerrish, PJ**. 2010. Beneficial mutations and the dynamics of adaptation in asexual populations. *Philos Trans R Soc Lond B Biol Sci* 365:1255–1263.

4. **Rosenzweig, RF, Sharp, RR, Treves, DS, Adams, J**. 1994. Microbial evolution in a simple unstructured environment: genetic differentiation in Escherichia coli. *Genetics* 137:903–917.

5. **Friesen, ML, Saxer, G, Travisano, M, Doebeli, M**. 2004. Experimental evidence for sympatric ecological diversification due to frequency-dependent competition in Escherichia coli. *Evolution* 58:245–260.

6. **Lind, PA, Farr, AD, Rainey, PB**. 2015. Experimental evolution reveals hidden diversity in evolutionary pathways. *Elife* 4:e07074.

7. **Rainey, PB, Travisano, M**. 1998. Adaptive radiation in a heterogeneous environment. *Nature* 394:69-72.

8. **Korona, R, Nakatsu, CH, Forney, LJ, Lenski, RE**. 1994. Evidence for multiple adaptive peaks from populations of bacteria evolving in a structured habitat. *Proceedings of the National Academy of Sciences* 91:9037–9041.

9. **Kerr, B, Riley, MA, Feldman, MW, Bohannan, BJM**. 2002. Local dispersal promotes biodiversity in a real-life game of rock–paper–scissors. *Nature* 418:171-174.

10. **Winter, C, Bouvier, T, Weinbauer, MG, Thingstad, TF**. 2010. Trade-Offs between

Competition and Defense Specialists among Unicellular Planktonic Organisms: the

"Killing the Winner" Hypothesis Revisited. *Microbiology and Molecular Biology Reviews*

74:42–57.

11.   **Brisson, D**. 2018. Negative Frequency-Dependent Selection Is Frequently Confounding.

Front Ecol Evol 6:10.

12.   **Betts, A, Kaltz, O, Hochberg, ME**. 2014. Contrasted coevolutionary dynamics between a

bacterial pathogen and its bacteriophages. *Proc Natl Acad Sci U S A* 111:11109–11114.

13.   **Gallet, R, Alizon, S, Comte, PA, Gutierrez, A, Depaulis, F, van Baalen, M, Michel, E,**

Müller-Graf, CD. 2007. Predation and disturbance interact to shape prey species diversity.

Am Nat 170:143–154.

14.   **D'Souza, G, Shitut, S, Preussger, D, Yousif, G, Waschina, S, Kost, C**. 2018. Ecology

and evolution of metabolic cross-feeding interactions in bacteria. *Nat Prod Rep* 35:455–

488.

15.   **Flyvbjerg, J, Arvin, E, Jensen, BK, Olsen, SK**. 1993. Microbial degradation of phenols

and aromatic hydrocarbons in creosote-contaminated groundwater under nitrate-reducing

conditions. *Journal of Contaminant Hydrology* 12:133-150.

16.   **Møller, S, Sternberg, C, Andersen, JB, Christensen, BB, Ramos, JL, Givskov, M,**

Molin, S. 1998. In situ gene expression in mixed-culture biofilms: evidence of metabolic

interactions between community members. *Appl Environ Microbiol* 64:721–732.

17.   **Hansen, SK, Rainey, PB, Haagensen, JA, Molin, S**. 2007. Evolution of species

interactions in a biofilm community. *Nature* 445:533–536.

18.   **Christensen, BB, Haagensen, JA, Heydorn, A, Molin, S**. 2002. Metabolic commensalism

and competition in a two-species microbial consortium. *Appl Environ Microbiol* 68:2495–

2502.

19. **Rodríguez-Verdugo, A**. 2021. Evolving Interactions and Emergent Functions in Microbial

Consortia. *mSystems* 6:e0077421.

20. **Rodríguez-Verdugo, A, Vulin, C, Ackermann, M**. 2019. The rate of environmental

fluctuations shapes ecological dynamics in a two-species microbial system. *Ecol Lett*

22:838–846.

21. **Rodríguez-Verdugo, A, Ackermann, M**. 2021. Rapid evolution destabilizes species

interactions in a fluctuating environment. *ISME J* 15:450–460.

22. **Deatherage, DE, Barrick, JE**. 2014. Identification of mutations in laboratory-evolved

microbes from next-generation sequencing data using breseq. *Methods Mol Biol* 1151:165–

188.

23. **Martínez-García, E, Nikel, PI, Chavarría, M, de Lorenzo, V**. 2014. The metabolic cost

of flagellar motion in Pseudomonas putida KT2440. *Environ Microbiol* 16:291–303.

24. **Ritchings, BW, Almira, EC, Lory, S, Ramphal, R**. 1995. Cloning and phenotypic

characterization of fleS and fleR, new response regulators of Pseudomonas aeruginosa

which regulate motility and adhesion to mucin. *Infection and Immunity* 63:4868–4876.

25. **Zhao, L, Zhang, Q, Zhang, D**. 2016. Evolution alters ecological mechanisms of

coexistence in experimental microcosms. *Functional Ecology* 30:1440–1446.

26. **Molina-Henares, MA, Ramos-González, MI, Daddaoua, A, Fernández-Escamilla, AM,**

Espinosa-Urgel, M. 2017. FleQ of Pseudomonas putida KT2440 is a multimeric cyclic

diguanylate binding protein that differentially regulates expression of biofilm matrix

components. *Res Microbiol* 168:36–45.

27. **Xiao, Y, Nie, H, Liu, H, Luo, X, Chen, W, Huang, Q**. 2016. C-di-GMP regulates the

expression of lapA and bcs operons via FleQ in Pseudomonas putida KT2440. *Environ*

Microbiol Rep 8:659–666.

28. **Jiménez-Fernández, A, López-Sánchez, A, Jiménez-Díaz, L, Navarrete, B, Calero, P,**

Platero, AI, Govantes, F. 2016. Complex Interplay between FleQ, Cyclic Diguanylate and

Multiple σ Factors Coordinately Regulates Flagellar Motility and Biofilm Development in

Pseudomonas putida. *PLoS One* 11:e0163142.

29. **Thompson, MG, Valencia, LE, Blake-Hedges, JM, Cruz-Morales, P, Velasquez, AE,**

Pearson, AN, Sermeno, LN, Sharpless, WA, Benites, VT, Chen, Y, Baidoo, EEK,

Petzold, CJ, Deutschbauer, AM, Keasling, JD. 2019. Omics-driven identification and

elimination of valerolactam catabolism in Pseudomonas putida KT2440 for increased

product titer. *Metab Eng Commun* 9:e00098.

30. **Niehaus, TD, Elbadawi-Sidhu, M, de Crécy-Lagard, V, Fiehn, O, Hanson, AD**. 2017.

Discovery of a widespread prokaryotic 5-oxoprolinase that was hiding in plain sight. *J Biol*

Chem 292:16360–16367.

31. **Koskiniemi, S, Sun, S, Berg, OG, Andersson, DI**. 2012. Selection-driven gene loss in

bacteria. *PLoS Genet* 8:e1002787.

32. **Kümmerli, R, Santorelli, LA, Granato, ET, Dumas, Z, Dobay, A, Griffin, AS, West,**

SA. 2015. Co-evolutionary dynamics between public good producers and cheats in the

bacterium Pseudomonas aeruginosa. *J Evol Biol* 28:2264–2274.

33. **Rodríguez-Verdugo, A, Tenaillon, O, Gaut, BS**. 2016. First-Step Mutations during

Adaptation Restore the Expression of Hundreds of Genes. *Mol Biol Evol* 33:25–39.

34. **Knöppel, A, Knopp, M, Albrecht, LM, Lundin, E, Lustig, U, Näsvall, J, Andersson,**

DI. 2018. Genetic Adaptation to Growth Under Laboratory Conditions in Escherichia coli

and Salmonella enterica. *Front Microbiol* 9:756.

35.   **Mohamed, ET, Werner, AZ, Salvachúa, D, Singer, CA, Szostkiewicz, K, Rafael**

Jiménez-Díaz, M, Eng, T, Radi, MS, Simmons, BA, Mukhopadhyay, A, Herrgård, MJ,

Singer, SW, Beckham, GT, Feist, AM. 2020. Adaptive laboratory evolution of

Pseudomonas putida KT2440 improves p-coumaric and ferulic acid catabolism and

tolerance. *Metab Eng Commun* 11:e00143.

36.   **Callahan, BJ, Fukami, T, Fisher, DS**. 2014. Rapid evolution of adaptive niche

construction in experimental microbial populations. *Evolution* 68:3307–3316.

37.   **Roman, MS, Wagner, A**. 2018. An enormous potential for niche construction through

bacterial cross-feeding in a homogeneous environment. *PLoS computational biology*

14:e1006340.

38.   **Scarinci, G, Sourjik, V**. 2023. Impact of direct physical association and motility on fitness

of a synthetic interkingdom microbial community. *ISME J* 17:371–381.

39.   **Grainger, TN, Letten, AD, Gilbert, B, Fukami, T**. 2019. Applying modern coexistence

theory to priority effects. *Proc Natl Acad Sci U S A* 116:6205–6210.

40.   **Chesson, P**. 2000. Mechanisms of Maintenance of Species Diversity. *Annual Review of*

Ecology, Evolution and Systematics 31:343–366.