Membrane curvature association of amphipathic helix 8 drives constitutive GPCR endocytosis

Rasmus Herlo*[1], Jan Hendrik Schmidt*[1], Joscha Rombach[1], Andreas Haahr Larsen[1], Mikkel Stoklund[1], Mathias Perslev[1], Tommas Theiss Ehlers Nielsen[1], Keenie Ayla Andersen[1], Carmen Klein Herenbrink[1], Matthew D. Lycas[1], Aske Lykke Ejdrup[1], Nikolaj Riis Christensen[1], Jan P. Christensen[2], Mootaz Salman[3], Freja Herborg[1], Ulrik Gether[1], Alexander Sebastian Hauser[4], Patricia Bassereau[5], David Perrais[6], Kenneth Lindegaard Madsen[1]

[1]Molecular Neuropharmacology and Genetics Laboratory, Department of Neuroscience, Faculty of Health and Medical Sciences; University of Copenhagen, 2200 Copenhagen N, Denmark.

[2]Department of Immunology and Microbiology, Faculty of Health and Medical Sciences; University of Copenhagen, 2200 Copenhagen N, Denmark.

[3]Department of Physiology, Anatomy and Genetics, Wolfson College, University of Oxford, Oxford OX1 3QU, UK

[4]Department of Drug Design and Pharmacology, University of Copenhagen, 2100 Copenhagen, Denmark

[5]Institut Curie, Université PSL, Sorbonne Université, CNRS UMR168, Laboratoire Physico-Chimie Curie, 75005 Paris, France

[6]Univ. Bordeaux, CNRS, Interdisciplinary Institute for Neuroscience; IINS, UMR 5297, 33000 Bordeaux, France.

*These authors contributed equally to this work

Correspondance: kennethma@sund.ku.dk

1 One-Sentence Summary

Receptor proteins navigate cellular membranes by interacting with their curvature using an evolutionary conserved mechanism that complements direct coupling to the endocytic protein machinery.

5 Abstract

Cellular signalling relies on the activity of transmembrane receptors and their presentation on the cellular surface. Their continuous insertion in the plasma membrane is balanced by constitutive and activity dependent internalization, which is orchestrated by adaptor proteins recognizing semi-specific motifs within the receptors' intracellular regions. Using multiple of fluorescence-based cellular assays, we discover a complementary and evolutionary conserved trafficking mechanism for G-protein coupled receptors, which relies on the insertion of their amphipathic helix 8 (H8) into the inner leaflet of lipid membranes, orthogonal to the transmembrane helices. These amphipathic helices autonomously drive endocytosis of receptors by cooperative assembly and association with areas of high membrane curvature. The strength of H8 membrane insertion propensity quantitatively predicts the rate of constitutive internalization of G-proteins coupled receptors.

17 Introduction

Cellular trafficking and sorting processes direct transmembrane proteins (TMPs) to their proper destination within the cell. Consequently, these processes are of fundamental importance for cellular homeostasis, metabolism, and communication. Malfunction results in diseases including

cancer, metabolic syndroms, neuropathies, neuro-degenerative disease and psychiatric

disorders [1].

Vesicular membrane carriers are responsible for the transport of TMPs between different

cellular compartments and TMP cargo selection is typically mediated by semi-specific protein-

protein interactions between compartment specific adaptor proteins (e.g. AP1-5 and GGAs) and

short linear motifs within the cargo TMPs (e.g. FXNPXY, tyrosine- and dileucine-based signals) [2,3].

The initial budding and formation of these carriers involve generation of high membrane

curvature which serves to recruit specific scaffolding and regulatory cytosolic proteins [4,5]. This

mechanism, termed membrane curvature sensing (MCS), is commonly mediated by

amphipathic helices (AHs) that insert partially into exposed lipid packing defects in the convex

leaflet of curved membranes [6,7]. We hypothesized that AH motifs located in the intracellular

termini of TMPs, positioned at the membrane-cytoplasm interface, serve to partition TMPs into

areas of high curvature during formation of vesicular carriers. Using G protein-coupled receptors

(GPCRs) as example, our data demonstrates that AHs direct constitutive trafficking of TMPs with

rates quantitatively predicted by their membrane insertion propensity (MIP). We demonstrate

that this new trafficking mechanism for TMPs, relying on their direct association with highly

curved membranes in budding membrane carriers, is fundamentally distinct from conventional

adaptor-based TMP trafficking processes, such as that underlying agonist-induced

internalization. Moreover, we present data suggesting that it is evolutionary conserved across

GPCR families and animal species. We refer to this mechanism as Trafficking of Transmembrane

proteins by Amphipathic Motifs (ToTAM).

Results

Amphipathic helices drive constitutive internalization of transmembrane proteins

G protein-coupled receptors (GPCRs) constitute the largest family of TMPs in the human genome, they control vital physiological functions, and represent targets for roughly a third of all approved drugs [8]. GPCRs consist of 7 transmembrane ⍺-helices and an amphipathic 8th helix (H8) that can insert into the cytosolic leaflet of the membrane orthogonal to the transmembrane helices (Figure 1A) [9–12]. Interactions between H8 and cytosolic adaptor proteins are known to mediate various trafficking processes, particularly agonist-induced endocytosis, but also constitutive internalization [13]. We tested a putative autonomous function of H8 in endocytosis in a membrane-anchored context, by fusing peptide sequences harboring H8 (17-18 residues) from a number of well-characterized class A GPCRs (rhodopsin (Rho), β2 adrenergic receptor (β2), substance P receptor (NK1)), and dopamine D2 receptor (D2) to the cytosolic C-terminus of the internalization inert transmembrane interleukin-2 receptor a-subunit (IL-2a), hereafter referred to as Tac (Figure 1A) [14,15].

The fusion of AHs increased internalization of Tac as determined by quantitative confocal microscopy of antibody feeding experiments in HEK293 cells (up to 10-fold for Tac-D2) (Figure 1B-C). Disrupting the hydrophobic wedge of the amphipathic helices by introducing negative charges (Figure 1A) abrogated this effect (Figure 1B-C). We recapitulated these findings in HeLa cells (Figure S1). Alanine substitution of cysteines, previously shown to modify trafficking of GPCRs through acylation [16], did not compromise internalization (Figure S2).

After confirming the relative internalization rates with high-throughput quantitative flow

cytometry [17] (Figure S3), we showed that five additional H8s with distinct amphipathic nature,

i.e., with high hydrophobic moment, also increased Tac internalization (Figure 1D). These were

H8 from trace amine receptor 1 and 6 (TA1 and TA6), ☐ opioid receptor (☐), dopamine D1

receptor (D1) and angiotensin 2 type 1 receptor (AT1) (biochemical parameters outlined in

Figure S4). To consolidate and broaden these findings, we also tested non-GPCR helices. Fusion

of a hydrophilic helix from troponin [18] or the hydrophobic helix from citrate [19] did not increase

Tac internalization, whereas AHs of several membrane-associated proteins involved in

membrane trafficking, including Amphipathic Lipid Packing Sensor (ALPS), Sar1 and Endophilin

A1, but not Epsin-1 [20,21] facilitated internalization (Figure 1E). Similarly, exogenous AHs, including

the bee venom Melittin, the bacterial protein SpoVM and the artificial AH Hecate, strongly

facilitated internalization. Together these findings demonstrate that AHs, diverse in origin and

devoid of known adaptor motifs, autonomously increase endocytosis of a transmembrane

protein. We define this mechanism as Trafficking of Transmembrane proteins by Amphipathic

Motifs (ToTAM).

To quantify the propensity of the individual AHs to associate with positive membrane curvature,

we adjusted the calculated hydrophobic moment of AHs by their intrinsic helical propensity (see

methods) [22]. The resulting value, membrane insertion propensity (MIP), correlated with

empirical measures of membrane association strength as determined by liposome binding to

peptides on a SPOT array (Figure S5). Furthermore, plotting internalization of the AH-fused Tac-

constructs as a function of MIP yielded a remarkably strong correlation irrespective of the origin

of the AHs (Figure 1F), demonstrating that AHs drive internalization of TMPs in a highly

predictive manner. Comparatively, net charge and hydrophobicity demonstrated less predictive

power on their own (Figure S6). In absolute terms, introduction of AHs increased internalization

from a basal level of 20% of Tac over a 30-minute period to a complete (100%) turnover of Tac-

AHs for the strongest AHs (Figure S7).

Finally, to address the endocytic function of H8 in context of full-length receptors, we focused

on β2 and AT1, that contain H8s with pronounced differences in their autonomous endocytic

drive (Figure 1F). In concordance, AT1 displayed significantly higher constitutive internalization

than β2, and this pattern was reversed by swapping their helices (Figure 1G-H and S8).

Furthermore, the introduction of the potent AH Hecate gave rise to an even higher

internalization rate than that of AT1. In contrast, internalization in presence of agonist was

robust for all constructs irrespective of AH (Figure 1H and S8). Further, while constitutive

internalization scaled linearly with expression, agonist induced internalization for the H8

containing constructs deviated from linearity consistent with saturation of adaptor-based

internalization machinery (Figure S8). This suggests that MIP of H8 drives constitutive, but not

agonist-induced, internalization of GPCRs in a mechanistically novel fashion.

AH driven Internalization is temporally dissociated from CME

Next, we chose a series of five Tac-AHs, spanning from low to high MIP values and assessed

their differential association with endocytic machineries using a proximity biotinylation

approach followed by quantitative label-free LC-MS² (Figure 2A) [23]. Importantly, the BioID2-

fusion constructs recapitulated ToTAM driven internalization (Figure 2B and S9A-B). The Tac-AHs

conferred low enrichment of biotinylated cytosolic proteins relative to control cells not

expressing the BioID2 construct, whereas labeling of plasma membrane proteins was strongly

enriched (Figure 2C and S9E). This enrichment of plasma membrane proteins was gradually

reduced with increasing MIP of the residing AHs, consistent with an increased internalization

and/or reduced maturation of these constructs (Figure 2D).

The MIP of the expressed Tac-AHs did not significantly affect the enrichment of actin, (Figure 2C

and S9F) consistent with a reduced impact of treatment with the actin polymerization inhibitor

Latrunculin A on internalization of Tac-AHs with high MIP (Figure S10A-C). Dynamin showed

slight enrichment with increasing MIP of the AHs and all constructs showed reduced

internalization upon Dyngo-4A treatment, indicating at least a partial dependence on dynamin

(Figure S10A-C). Likewise, ATP-depletion reduced internalization for all constructs albeit

relatively less for Tac-AHs with high MIP (Figure S10D-F).

Overall, key proteins of clathrin-mediated endocytosis (CME)[24], and in particular clathrin and

EPS15, were enriched for all AHs. On the other hand, identified components of the GLIC/GEEC

pathway, as well as caveolae[25,26], were consistently less enriched with increasing MIP of the AHs

(Figure 2E). Finally, enrichment of proteins involved in alternative endocytic pathways, including

Arf6, Rac1, Endophilin A2, and Flotillins (FLOT1 and FLOT2 [27]), were relatively independent of

MIP. Grouping all identified CME associated proteins according to temporally defined modules

[28], we observed no marked increase in association with nucleation and assembly, neck

constriction or scission and movement modules, while proteins involved in uncoating as well as

endosome fusion were enriched for all AHs (Figure S9G). Notably, proteins of the nucleation and

formation module that showed an increased abundance, including clathrin and EPS15,

represented a group that has been shown to be continuously associated with CME events until

uncoating [24]. Together, this could indicate that ToTAM may accelerate maturation of CME

structures or substitute for several of the proteins involved in maturation, including, e.g., BAR

domain proteins (Figure S9H). EPS15 is known to recruit ubiquitinated cargo to endocytic sites

[29]. In accordance, we detected minor increases in ubiquitin binding proteins of the ESCRT

machinery and E3 ubiquitin ligases [30], however, this could simply reflect the localization of these

components to intracellular membranes (Figure S9I-J). Taken together, ToTAM associates with

CME rather than GLIC/GEEC pathway, caveolae or alternative pathways, and displays a MIP

dependent association with dynamin. Notably, we found no association between Tac-AHs and

any classical or putative novel adapter proteins.

To directly evaluate the trafficking dynamics of Tac-AH constructs and the CME structures they

associate with, we turned to pulsed pH (ppH) TIRFM assay [31] using super ecliptic pHluorin (SEP)

tagged constructs co-expressed with mCherry-tagged clathrin light chain (CLC) (Figure 2G-H and

S11). Corroborating the biochemical findings, the number of endocytic events increased by

almost an order of magnitude with increased MIP of the AHs, approaching the rate of the

transferrin receptor (TfR) for the strongest helices (H8 from AT1, and Hecate) (Figure 2I).

Interestingly, exocytosis increased correspondingly, suggesting efficient intracellular sorting to

recycling endosomes (Figure S12). CLC fluorescence intensity traces associated with TfR

endocytic events displayed a typical profile with gradual accumulation until point of scission

followed by a rapid decrease after scission (Figure 2J). This distinct profile was gradually lost

with increasing MIP of the Tac-AH constructs indicating desynchronization with the CME

machinery. Individual normalized traces displayed a range of CLC recruitment profiles (Figure

S13). Principal component analysis (PCA) revealed a decreasing fraction of traces similar to the

typical TfR-associated CLC profile for Tac-AHs with increasing MIP (Figure S14). Conversely, an

increased fraction of events displayed accumulation of CLC immediate post scission. In

summary, increased MIP directs ToTAM towards bulk association with CME core machinery at

the cost of other endocytic machineries, while simultaneously decreasing the temporal

synchronization and coordination with clathrin-dynamics.

Membrane curvature association facilitates internalization

Next, we analyzed the recruitment dynamics of the SEP-Tac-AH constructs prior to scission

(Figure 3A). While TfR accumulates slowly during vesicle formation [32], the AH-linked profiles

showed accelerating kinetics with increasing MIP of AHs. Fitting these profiles with a logistic

equation showed increasingly cooperative clustering mechanism consistent with an auto-

nucleating function (Figure 3B). This effect implies an increased rate of maturation of endocytic

structures, which was supported by the observed MIP dependent effect on Tf-uptake in trans

(Figure 3C). Taken together, we find that ToTAM is characterized by numerous, fast and

cooperative endocytic events with the potential to accelerate overall trafficking rates.

The desynchronization with canonical endocytic machinery, the cooperative recruitment and

the inductive properties together indicate an underlying autonomous and recursive mechanism,

resulting from concurrent membrane curvature sensing and induction, a common characteristic

of AH association with lipid membranes. We therefore investigated the MCS properties of

Hecate (the AH with highest tested MIP) in a buckled membrane system using molecular

dynamics simulation. This simulation predicted a curvature sensitive membrane sorting (Figure

3D-E), akin to ALPS [33]. We experimentally confirmed this finding by pulling tethers from a GUV

preincubated with OG488-Hecate, observing a significant enrichment in areas of high curvature

(peptide on tether relative to GUV) (Figure 3F-G). Reducing helical propensity and MIP of the

helix, while keeping charge constant, disrupted binding to GUVs (Figure S15A-B). Consistent

with the cooperative clustering observed in the ppH-TIRF experiments, the Hecate peptide

occasionally formed microdomains on GUVs in the absence of manipulation (Figure S15C).

To assess the size and curvature of the endocytic budding structures prior to scission, we

antibody-labelled surface expressed Tac-AHs for 3D-dSTORM imaging and identified protein

clusters using DBSCAN analysis (Figure 3H and S16). While the median radius of clusters was 100

nm for Tac, consistent with the size of clathrin-coated pits (CCPs), cluster sizes were consistently

smaller with increasing MIP of AHs (Figure 3I). In accordance, cargo load of pH 5.5 events at

point of scission in ppH-TIRFM experiments was decreased with increasing MIP of AHs (Figure

S17). Protein densities of individual budding structures obtained by 3D-dSTORM gradually

approached a linear dependence on size with increasing MIP of the AHs (Figure 3J-K), as

previously modelled for insertion-based membrane curvature sensing [6]. Estimates from relative

protein density and quenching rates of vesicular carriers immediately post-scission obtained

from ppH-TIRFM experiments supported increased protein density in endocytic structures

containing Tac-AHs with high MIP (Figure S17). In summary, we propose that ToTAM

mechanistically serves both to localize cargo to endocytic pits as well as to facilitate the

membrane curvature induction and possibly scission of these pits to drive endocytosis.

Membrane insertion propensity is evolutionary conserved across distant organisms

Next, given the conservation of amphipathic periodicity in H8 across GPCR families [34] (Figure

S18A), we upscaled the liposome binding SPOT assay (Figure S5) to a large-scale chip array

format (Figure 4A-B). This allowed us to scan through the full range of human GPCR C-termini,

without compromising assay performance (Figure S18B). Binding profiles demonstrated a

confined membrane binding region overlapping with the H8 region (consecutive high MIP-

values) as illustrated by the C-terminal regions of rhodopsin, β2 and AT1 (Figure 4C). Averaged

membrane binding for each of the GPCR families confirmed this pattern for rhodopsin

(disregarding olfactory receptors), secretin and frizzled families, albeit with membrane binding

extending slightly further away from the TM region for the latter. Membrane binding was less

pronounced for the remaining families and absent from the glutamate receptor family (Figure

S18C). UMAP analysis of the binding profiles revealed a group of GPCRs (cluster 1) without

membrane binding, while clusters 2-5 showed membrane binding profiles consistent with

membrane proximal H8, albeit with increasing length and distance from TM7 from cluster 2 to 5

(Figure 4D-E and Figure S19). In contrast, cluster 6 contained a wide range of different

membrane binding profiles, which are distant from TM7 and/or contain multiple membrane

binding sites. We did not observe an enrichment of individual clusters within GPCR receptor

families (Figure S18D), specific tissues (Figure S18E) or Gene Ontology (GO) terms (data not

shown).

The experimentally derived membrane binding profiles enabled us to calculate MIP values for

the regions corresponding to bona fide H8s (clusters 2-5, Figure S20), while the lack of

membrane binding in cluster 1, precluded calculation of MIP values. We previously

characterized the constitutive internalization of a range of pharmacologically relevant GPCRs [35]

and here asked if this internalization could be explained by MIP values of the regions defined by

their membrane binding profiles. Constitutive internalization significantly correlated with MIP

(cluster 2-5) (Figure 4F), however, traditional biophysical measures, alternative to MIP, failed to

explain constitutive internalization (Figure S21A-C). The complex membrane binding profiles

cluster 6 precluded extraction of MIP values predictive of constitutive internalization (Figure

S21D). Consistent with the lack of effects of H8 substitution on agonist induced internalization

(Figure 1H), MIP did not correlate with agonist-induced internalization (Figure S21E). Together,

these results indicate that the MIP of H8 selectively predicts constitutive internalization across

this functionally important and structurally well-characterized subset of GPCRs.

Given the functional importance of constitutive internalization for dynamic surface expression

of GPCRs, we utilized MIP for H8 as a proxy to map predicted internalization rates across all non-

odorant GPCRs belonging to cluster 2-5 (Figure 4G and S21F). High MIP values are

predominantly observed as phylogenetically related clusters within the rhodopsin family of

receptors, including receptors for biogenic amines, chemokine receptors, and MAS-related

receptors as well as free fatty acid receptors and individual receptors such as NPBW2 and

C5AR1. We also note that Secretin and Frizzled receptors largely contained a H8 predicted or

found to be longer than 20 residues and the functional significance of this remains to be

explored. Notably, MIP is not significantly related to β-arrestin recruitment and reported G-

protein coupling (Figure S22A-B).

To assess functional conservation of high MIP of H8, we used data from UK biobank on naturally

occurring variants in human GPCRs. While the cost of individual amino acid substitutions was

similar, the frequency of non-synonymous natural variants was significantly lower within H8s

compared to the remaining C-terminus (Figure S23A-B). Moreover, the observed natural

variants give rise to more conserved MIP than potential substitutions not observed (Figure

S23C). Finally, the frequency of natural variants within H8 for individual receptors declines with

increasing MIP, emphasizing the functional conservation of high MIP (Figure S23D). This

functional selection is further evident from the multimodal distribution of H8 MIP values across

human GPCRs, significantly different from a normal distribution (Figure 4H, p <0.0001, Shapiro-

Wilk). Sub-groups of GPCRs shown to be directed to the Fast Endophilin-Mediated Endocytosis

(FEME) pathway, to mediate endosomal signaling or to function as mechanosensitive GPCRs

display H8 MIP value distributions similar to this overall distribution. GPCRs associated with

primary cilia, which exhibit inverse curvature topology to endocytic structures, on the other

hand, display H8 sequences with significantly lower MIPs. Conversely, GPCRs expressed by

viruses, which are known to elicit high constitutive internalization [36], mostly have high MIP

values of their H8 (Figure 4H and S22C).

GPCR C-termini from evolutionary distant organisms, including C. elegans, D. Melanogaster and

S. Cerevisiae, show membrane binding profiles akin to H8 in human GPCRs (cluster 2 and 3) as

well as non-binding (cluster 1) and more complex binding profiles (cluster 4 and 5) (Figure S24).

This enabled us to address evolutionary conservation across species by aligning H8 sequences

defined in human across species back to Ecdysozoa (Figure 4H). Individual receptors show

strong conservation of MIP throughout evolution, and the relative variation across species is

lowest for H8s with high MIP (Figure 4H, top). Additionally, the fraction of receptors with high

MIP of H8 (> 0.3) gradually increase in higher organisms indicating functional significance

(Figure 4H). Together, this demonstrates both evolutionary selection and functional

conservation of H8 in GPCRs with high MIP.

Discussion

Here we show that AHs enable TMPs to associate with areas of high membrane curvature

similar to what has been demonstrated for cytosolic proteins. For TMPs, this association

accelerates their constitutive trafficking in a manner that correlates with the MIP of the AH.

While this paper has exclusively addressed endocytosis and recycling, we hypothesize that this

mechanism, due to the basic biophysical principle, will generalize to trafficking events

throughout the endomembrane system involving highly curved membrane topologies. We term

this mechanism ToTAM.

Mechanistically, ToTAM is characterized by fast, cooperative cargo recruitment into small,

spherical, high-density carriers and thus ToTAM displays elements of both membrane curvature

sensing and facilitation of the cargo/carrier complex formation. In fact, ToTAM increases

endocytosis of the transferrin receptor in trans, implying an active impact on endocytic

processes. During this process, partial biochemical association with CME was observed,

however, actual CLC accumulation prior to scission was low and increasingly asynchronous with

higher MIP. Together, our data suggest that ToTAM might hijack nucleation involving CLC and

EPS15 to shortcut classical CME in a relatively autonomous fashion that is complementary to

the established adaptor based endocytic machinery.

This specific impact of high MIP H8 on constitutive, and not agonist-induced, internalization of

full length GPCRs, was causally demonstrated by experiments with chimeric receptors. Using

large-scale liposome-binding data to exclude GPCRs with C-termini that did not display

membrane-binding or complex membrane binding enabled us to correlate H8 MIP with full-

length constitutive internalization across all human GPCRs. Highest constitutive internalization

was bioinformatically predicted for the groups of chemokine receptors, biogenic amine

receptors, MAS-related receptors, and free fatty acid receptors. It will be interesting to elucidate

the functional implication of the predicted high constitutive turnover with e.g. ligand

scavenging, membrane modulating ligands or continuous receptor availability.

Another recently described membrane-associated function of H8 is its role in conferring

mechanosensitivity to GPCRs [37]. Several of these receptors include a H8 with high MIP, including

H1, AT1, and S1P1, and a putative interplay between mechanosensation and high constitutive

internalization due to membrane curvature sensing might reflect two aspects of the same

biophysical principle and functional implications of these might unveil themselves perhaps in

the context of the cardiovascular system. Moreover, GPCRs implicated in viral entry, as well as

virally expressed GPCRs displaying high constitutive internalization, all except mGluR2 show high

MIP [38], potentially advocating a novel molecular target for anti-viral therapy. Finally, the simple

molecular nature of ToTAM is compatible with an early evolutionary origin, dating back prior to

the yeast GPCRs characterized here, conceivably even prior to adaptor based trafficking.

Ultimately, functional studies will be needed to settle the evolutionary relation between these

two complementary trafficking mechanisms.

References

1.   McMahon, H. T. & Boucrot, E. Molecular mechanism and physiological functions of clathrin-
mediated endocytosis. *Nature Reviews Molecular Cell Biology* vol. 12 517–533 Preprint at
https://doi.org/10.1038/nrm3151 (2011).

2.   Bonifacino, J. S. & Traub, L. M. Signals for sorting of transmembrane proteins to endosomes and
lysosomes. *Annual Review of Biochemistry* vol. 72 395–447 Preprint at
https://doi.org/10.1146/annurev.biochem.72.121801.161800 (2003).

3.   Sanger, A., Hirst, J., Davies, A. K. & Robinson, M. S. Adaptor protein complexes and disease at a
glance. *J Cell Sci* **132**, (2019).

4.   McMahon, H. T. & Boucrot, E. Membrane curvature at a glance. *J Cell Sci* **128**, 1065–1070 (2015).

5.   Antonny, B. Mechanisms of Membrane Curvature Sensing. *Annu Rev Biochem* **80**, 101–123
(2011).

6.   Hatzakis, N. S. *et al.* How curved membranes recruit amphipathic helices and protein anchoring
motifs. *Nat Chem Biol* **5**, 835–841 (2009).

7.   Drin, G. & Antonny, B. Amphipathic helices and membrane curvature. *FEBS Lett* **584**, 1840–1847
(2010).

8.   Hauser, A. S., Attwood, M. M., Rask-Andersen, M., Schiöth, H. B. & Gloriam, D. E. Trends in GPCR
drug discovery: New agents, targets and indications. *Nat Rev Drug Discov* **16**, 829–842 (2017).

9.   Palczewski, K. *et al.* Crystal Structure of Rhodopsin : A G Protein – Coupled Receptor. **289**, 739–
746 (2000).

10.   Isberg, V. *et al.* GPCRdb: An information system for G protein-coupled receptors. *Nucleic Acids*
Res **44**, D356–D364 (2016).

11.   Rosenbaum, D. M., Rasmussen, S. G. F. & Kobilka, B. K. The structure and function of G-protein-
coupled receptors. *Nature* vol. 459 356–363 Preprint at https://doi.org/10.1038/nature08144 (2009).

12.   Zhang, H. *et al.* Structure of the Angiotensin Receptor Revealed by Serial Femtosecond
Crystallography Article Structure of the Angiotensin Receptor Revealed by Serial Femtosecond
Crystallography. 833–844 (2015) doi:10.1016/j.cell.2015.04.011.

13.   Huynh, J., Thomas, W. G., Aguilar, M. I. & Pattenden, L. K. Role of helix 8 in G protein-coupled
receptors based on structure-function studies on the type 1 angiotensin receptor. *Mol Cell Endocrinol*
302, 118–127 (2009).

14.   Bonifacino, J. S., Cosson, P. & Klausner, R. D. Colocalized transmembrane determinants for ER
degradation and subunit assembly explain the intracellular fate of TCR chains. *Cell* (1990)
doi:10.1016/0092-8674(90)90447-M.

15.   Tan, P. K., Waites, C., Liu, Y., Krantz, D. E. & Edwards, R. H. A leucine-based motif mediates the
endocytosis of vesicular monoamine and acetylcholine transporters. *Journal of Biological Chemistry*
(1998) doi:10.1074/jbc.273.28.17351.

16.   Neve, K. A., Qanbar, R. & Bouvier, M. Role of palmitoylation/depalmitoylation reactions in G-protein-coupled receptor function. *Pharmacology and Therapeutics* vol. 97 1–33 Preprint at https://doi.org/10.1016/S0163-7258(02)00300-5 (2003).

17.   Mostafapour, S., Kobilka, B. K. & von Zastrow, M. Pharmacological sequestration of a chimeric beta 3/beta 2 adrenergic receptor occurs without a corresponding amount of receptor internalization. *Recept Signal Transduct* **6**, 151–163 (1996).

18.   Heidorn, D. B. & Trewhell, J. Comparison of the Crystal and Solution Structures of Calmodulin and Troponin C. *Biochemistry* **27**, 909–915 (1988).

19.   Introduction to Protein Structure, by Branden, 2nd Edition: Carl Branden,John Tooze: 9788153230505: Amazon.com: Books. https://www.amazon.com/Introduction-Protein-Structure-Branden-2nd/dp/8153230506.

20.   Magdeleine, M. *et al.* A filter at the entrance of the Golgi that selects vesicles according to size and bulk lipid composition. *Elife* (2016) doi:10.7554/eLife.16988.

21.   Mesmin, B. *et al.* Two lipid-packing sensor motifs contribute to the sensitivity of ArfGAP1 to membrane curvature. *Biochemistry* **46**, 1779–1790 (2007).

22.   Høie, M. H. *et al.* NetSurfP-3.0: accurate and fast prediction of protein structural features by protein language models and deep learning. *Nucleic Acids Res* **50**, W510–W515 (2022).

23.   Kim, D. I. *et al.* An improved smaller biotin ligase for BioID proximity labeling. *Mol Biol Cell* **27**, 1188–1196 (2016).

24.   Kaksonen, M. & Roux, A. Mechanisms of clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol* **19**, 313–326 (2018).

25.   Shafaq-Zadah, M., Dransart, E. & Johannes, L. Clathrin-independent endocytosis, retrograde trafficking, and cell polarity. *Current Opinion in Cell Biology* vol. 65 112–121 Preprint at https://doi.org/10.1016/j.ceb.2020.05.009 (2020).

26.   Parton, R. G. Caveolae: Structure, Function, and Relationship to Disease. *Annu Rev Cell Dev Biol* **34**, 111–136 (2018).

27.   Casamento, A. & Boucrot, E. Molecular mechanism of Fast Endophilin-Mediated Endocytosis. *Biochemical Journal* vol. 477 2327–2345 Preprint at https://doi.org/10.1042/BCJ20190342 (2020).

28.   Traub, L. M. Regarding the Amazing Choreography of Clathrin Coats. *PLoS Biol* **9**, e1001037 (2011).

29.   Polo, S. *et al.* A single motif responsible for ubiquitin recognition and monoubiquitination in endocytic proteins. *Nature* **416**, 451–455 (2002).

30.   Christ, L., Raiborg, C., Wenzel, E. M., Campsteijn, C. & Stenmark, H. Cellular Functions and Molecular Mechanisms of the ESCRT Membrane-Scission Machinery. *Trends in Biochemical Sciences* vol. 42 42–56 Preprint at https://doi.org/10.1016/j.tibs.2016.08.016 (2017).

31.   Merrifield, C. J., Perrais, D. & Zenisek, D. Coupling between clathrin-coated-pit invagination,
cortactin recruitment, and membrane scission observed in live cells. *Cell* **121**, 593–606 (2005).

32.   Taylor, M. J., Perrais, D. & Merrifield, C. J. A High Precision Survey of the Molecular Dynamics of
Mammalian Clathrin-Mediated Endocytosis. *PLoS Biol* **9**, e1000604 (2011).

33.   Stroh, K. S. & Risselada, H. J. Quantifying Membrane Curvature Sensing of Peripheral Proteins by
Simulated Buckling and Umbrella Sampling. *J Chem Theory Comput* **17**, 5276–5286 (2021).

34.   Dijkman, P. M. *et al.* Conformational dynamics of a G protein–coupled receptor helix 8 in lipid
membranes. *Sci Adv* **6**, (2020).

35.   Schmidt, J. H. *et al.* Constitutive internalization across therapeutically targeted GPCRs correlates
with constitutive activity. *Basic Clin Pharmacol Toxicol* **126**, 116–121 (2020).

36.   Rosenkilde, M. M., Tsutsumi, N., Knerr, J. M., Kildedal, D. F. & Garcia, K. C. Viral G Protein–
Coupled Receptors Encoded by β- and γ-Herpesviruses. *https://doi.org/10.1146/annurev-virology-*
100220-113942 **9**, 329–351 (2022).

37.   Erdogmus, S. *et al.* Helix 8 is the essential structural motif of mechanosensitive GPCRs. *Nat*
Commun **10**, 1–15 (2019).

38.   Wang, G. *et al.* The G Protein-Coupled Receptor FFAR2 Promotes Internalization during Influenza
A Virus Entry. *J Virol* **94**, (2020).

39.   Gautier, R., Douguet, D., Antonny, B. & Drin, G. HELIQUEST: a web server to screen sequences
with specific -helical properties. *Bioinformatics* **24**, 2101–2102 (2008).

40.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–
589 (2021).

Figures and legends

Figure 1. **Amphipathic helices in transmembrane proteins autonomously drive endocytosis**. **a**

Illustration of GPCR H8 sequences transferred to the single transmembrane protein Tac for

determination of their autonomous endocytic drive alongside mutational disruption of

amphipathicity by introduction of negative charges in the hydrophobic face of H8. **b**

Representative images of HEK293 cells transfected with Tac-H8 constructs and subjected to

antibody feeding (illustrated in insert, lower left). **c** Quantification of internalization (Red/Green

fluorescence ratio), relative to Tac, from confocal images (every dot represents a cell, data

compiled from three independent experiments). **d** Internalization of Tac-H8 constructs assessed

by flow cytometry and normalized to Tac. Mean $\pm$ SEM, N=3 **e** Internalization of Tac with AHs

from cytosolic proteins or exogenous AHs assessed by flow cytometry. Mean $\pm$ SEM, N=3. **f**

Linear correlation (R^2 = 0.67 p <0.0001) between Membrane Insertion Propensity (MIP) of Tac-

AH constructs and their respective endocytic rate (relative to Tac). Mean $\pm$ SEM, N=3. **g**

Illustration of chimeric H8 GPCRs with swapped part highlighted in helical wheel

representations using HeliQuest [39]. (H) Constitutive and agonist-induced (10 µM agonists; ISO =

Isoproterenol and ANGII = angiotensin II) internalization rates of chimeric FLAG-tagged GPCR

constructs with swapped H8 sequences relative to Tac. Mean $\pm$ SEM, N=3.

Figure 2. **ToTAM associates with CME machinery in a non-canonical manner. a** Schematic of

proximity biotinylation assay with BioID2. **b** Quantification of internalization of Flag-Tac AH

BioID2 constructs from confocal images (every dot represents a cell, compiled from three

independent experiments). **c** Differences (⬚⬚ in label-free quantification (LFQ) values for

indicated constructs versus basal levels (pcDNA) normalized by Tac compiled for all cytosolic and

membrane proteins, as defined by a STRING localization score of 5 with no other localization

above 4.5. **d** Western blot (M1 ab) of Flag-Tac AH-BioID2 constructs transfected in HEK293 cells

showing mature proteins (high MW band) and immature protein (low MW band). **e** Difference

in LFQ values versus endogenously biotinylated protein for identified actin and dynamin

subunits, respectively. **f** Absolute difference in LFQ values (normalized to Tac) of selected

identified proteins associated with CME, CLIG/GEEC, Caveolae and other endocytic pathways. **g**

Schematic of (ppH-TIRFM) protocol enabling definition of time of scission for individual

endocytic events. **h** Images of representative endocytic event from ppHTIRFM experiments of

Super Ecliptic pHluorin (SEP) tagged transferrin receptor (TfR) together with clathrin light chain

(CLC)-mCherry. **i** Endocytic event rate for the 5 different SEP-TacAH constructs compared to TfR-

SEP. Statistical analysis using Kruskal-Wallis multiple comparisons test. **j** (Upper) Average of

individually normalized CLC-mCherry traces for TfR-SEP and 5 SEP-TacAH constructs. (Lower)

Heatmaps of normalized individual CLC-mCherry traces pre- and post-scission. N = 5-10 cells

(846-2797 events) per condition.

Figure 3. **Amphipathic helices in transmembrane proteins promote endocytosis through**

association with curved membranes. a (Upper) Average of individually normalized pH 7.4 TfR-

SEP and SEP-TacAH fluorescence intensity traces building up to point of scission (Time = 0).

(Lower) Heatmaps of normalized individual SEP-fluorescence profiles at pH 7.4 of TfR-SEP and

SEP-TacAH constructs prior to scission. N = 5-10 cells per condition, n = 518-2367 total traces. **b**

Non-linear regression fits of averaged normalized pHluorin traces up to point of scission. H is

the Hill coefficient. **c** Effect of TacAH constructs on AF647-labelled transferrin endocytosis in

trans, by flow cytometry. **d** Relative protein density (Hecate peptide) as a function of tubule

diameter determined in buckled membrane MD analysis indicating a strong preference for

positive curvature. **e** Free energy from (**d**) as a function of membrane curvature. **f** Tethers pulled

from GUVs (red) pre-incubated with Oregon Green labelled Hecate peptide (green). Insert 1 and

3, zooms of GUV and summed intensity profile. Insert 2 and 4, zooms of tether and summed

intensity profile. **g** Quantification of the relative sorting to the tether as a function of relative

size of the tether (Tetherred/GUVred). **h** Representative 3D-dSTORM images of FLAG-Tac and 5

FLAG-TacAH constructs after surface labelling with Alexa Fluor 647-conjugated aFLAG antibody,

scale bar 5 μm. Inserts show representative single clusters. **i** Average size of clusters for the

individual constructs, median $\pm$ 95% CI, N=6. **j** Logarithmic relationship of cluster size and cargo

density (FLAG-Tac-AH). Black bars show binned averages $\pm$ SEM and the blue line represents a

linear fit of the data. Better adherence of the fit to the binned averages indicates curvature

dependent sorting for Tac-AH constructs with high MIP **k** Selectivity slope (i.e., curvature

sensitivity) derived for linear fits with error on log-log plots of the data in (**j**) (see Figure S16).

Figure 4. **Proteome-wide characterization of C-terminal membrane interactions in GPCRs and**

their role in constitutive endocytosis of therapeutically relevant GPCRs. a Schematic of chip

array for scanning of liposome binding strength throughout the full range of human GPCR C-

termini at dual AA resolution. Individual peptides are randomly positioned on the chip. **b**

Representative image of fluorescent signal from bound liposomes on zoomed in area of array.

Regular pattern in top left corner is control Hecate peptide. **c** Compiled binding profile of the C-

termini of Rhodopsin, β2AR and ATR along with corresponding MIP and net charge values

displayed below. **d** UMAP-based clustering of all human GPCR C-termini based on position and

strength of liposome binding. **e** Heatmaps of all individual GPCR C-terminal liposome binding

profiles within the six clusters and profile plots of the median fluorescence intensity (line above)

(see S19 from detailed view). **f** Linear correlation (R^2 = 0.83, p <0.0001) between MIP of H8 from

clusters 2-5, based on sequences (Figure S20) in most pharmacologically relevant GPCRs and

their respective endocytic rate (relative to Tac) determined by flow cytometry [35]. Mean ± SEM,

N=3. **g** Circular phylogenetic tree of human GPCRs based on sequence homology and divided

into GRAFS families and clusters, alongside the corresponding calculated MIP based on H8 from

either refined representative structures as annotated on GPCRdb (purple) [10], AlphaFold2

prediction (magenta) [40]. Receptors used in (**f**) are highlighted by green and receptors described

as mechanosensitive [37] are highlighted by asterisks (name). GPCRs from cluster 1 (no binding) in

grey. Receptors with a structural H8 predicted longer than 20 amino acids marked with asterisk

(bar). **h** Evolutionary tree of GPCRs displaying MIP of H8 sequences identified in Human

receptors and aligned across species (left). The ratio of variance vs. mean of all MIP decrease

with increasing MIP (upper, left). (Upper, right) MIP values for viral GPCRs (Table S2). (Lower

right) Density distribution of MIP values for individual species shift to the right during evolution.

473