Neurophysiology of effortful listening:
Decoupling motivational modulation from task demands

Frauke Kraus [1,2], Bernhard Ross [3,4], Björn Herrmann [3,5]*, & Jonas Obleser [1,2] *

[1] Department of Psychology, University of Lübeck, Lübeck, Germany

[2] Center of Brain, Behavior, and Metabolism, University of Lübeck, Lübeck, Germany

[3] Rotman Research Institute, Baycrest Academy for Research and Education, Toronto, ON, Canada

[4] Department of Medical Biophysics, University of Toronto, Toronto, ON, Canada

[5] Department of Psychology, University of Toronto, Toronto, ON, Canada

joint senior authorship

Correspondence concerning this article should be addressed to Björn Herrmann, Rotman Research Institute, Baycrest, 3560 14 Bathurst St, North York, ON, M6A 2E1, Canada, bherrmann@research.baycrest.org or to Jonas Obleser, Department of Psychology, University of Lübeck, Maria-Goeppert-Str. 9a, 23562 Lübeck, Germany, jonas.obleser@uni-luebeck.de.

Word count: Abstract (174), Introduction (778), Discussion (1773)

Number of Figures: 8

Keywords: Cognitive demand, pupil dilation, gap detection, motivation intensity theory, reward prospect, event-related field, neural alpha power

Conflict of interest statement:
The authors declare no competing financial interests.

Acknowledgments:
We thank Gloria Lai and Tazeen Atif for their assistance with data collection. This work was supported by Deutsche Forschungsgemeinschaft (DFG; grant number HE 7857/1-1) awarded to BH. BH is supported by the Natural Sciences and Engineering Research Council of Canada (Discovery Grant: RGPIN-2021-02602) and the Canada Research Chair program (CRC-2019-00156).

Author contributions:
F.K., B.H. and J.O. designed research, F.K. performed research, B.R. provided resources, F.K., J.O., and B.H. analyzed data, F.K. visualized the data, F.K. wrote first draft of the manuscript, F.K., B.R., B.H., and J.O. edited the manuscript.

Abstract

In demanding listening situations, a listener's motivational state may affect their cognitive investment. Here, we aim to delineate how domain-specific sensory processing, domain-general neural alpha power, and pupil size as a proxy for cognitive investment encode influences of motivational state under demanding listening. Participants performed an auditory gap-detection task while pupil size and the magnetoencephalogram (MEG) were simultaneously recorded. Task demand and a listener's motivational state were orthogonally manipulated through changes in gap duration and monetary-reward prospect, respectively. Whereas task difficulty impaired performance, reward prospect enhanced it. Pupil size reliably indicated the modulatory impact of an individual's motivational state. At the neural level, the motivational state did not affect auditory sensory processing directly but impacted attentional post-processing of an auditory event as reflected in the late evoked-response field and alpha power change. Both pre-gap pupil dilation and higher parietal alpha power predicted better performance at the single-trial level. The current data support a framework wherein the motivational state acts as an attentional top-down neural means of post-processing the auditory input in challenging listening situations.

Significance Statement

How does an individual's motivational state affect cognitive investment during effortful listening? In this simultaneous pupillometry and MEG study, participants performed an auditory gap-detection task while their motivational state was manipulated through varying prospect of a monetary reward. The pupil size directly mirrored this motivational modulation of the listening demand. The individual's motivational state also enhanced top-down attentional post-processing of the auditory

50 event but did neither change auditory sensory processing nor pre-gap parietal alpha power. These

51 data suggest that a listener's motivational state acts as a late attentional top-down effect on auditory

52 neural processes in challenging listening situations.

53   **Introduction**

The motivational state of a person influences whether they are willing to invest cognitively in

demanding listening contexts (Brehm & Self, 1989; Kraus et al., 2023a; Richter et al., 2016). That is, a

person who sees no intrinsic (e.g., enjoyment) or extrinsic (e.g., financial) value in listening, and is

thus in a low motivational state, may be more likely to disengage from listening than someone who

is highly motivated, especially under highly demanding conditions (Brehm & Self, 1989; Herrmann &

Johnsrude, 2020; Pichora-Fuller et al., 2016; Richter et al., 2016). When listening conditions are less

demanding, a person may still engage in listening despite being little motivated (Brehm & Self, 1989;

Richter et al., 2016). Although this interacting influence of task demands and motivation (see Fig. 1C)

is established theoretically and empirically, for example, for cognitive control tasks (Parro et al., 2018;

Yee & Braver, 2018), little research has been conducted to investigate the impact of task demands

and motivation on listening under challenges and to characterize the underlying neurophysiological

mechanisms.

66   The noradrenergic (NE) modulation emerging from locus coeruleus (LC) in the brainstem is

one driver of pupil dilation (Joshi et al., 2016; Joshi & Gold, 2020). The LC–NE system is sensitive to

attention (Vazey et al., 2018) and has a role in optimizing task performance and task engagement

(Aston-Jones & Cohen, 2005a). Accordingly, pupil dilation as a proxy of LC–NE activity has often been

used as an indicator of cognitive investment: The pupil dilates as a task becomes more difficult

(Kadem et al., 2020; Kahneman & Beatty, 1966; Koelewijn et al., 2012; Ohlenforst et al., 2018; Wendt

et al., 2016; Winn et al., 2015; Zekveld et al., 2010). Moreover, pupil size dynamics during high

listening demands are known to be modulated by an individual's motivational state (Alfandari et al.,

2023; Bijleveld et al., 2009; Zhang et al., 2019); recently confirmed by us using highly-controlled noise

stimuli (Kraus et al., 2023a) that will also be used the present study. The motivation-driven additional

cognitive investment indexed by pupil dilation predicted better behavioral outcomes (Kraus et al.,

2023a).

Neural oscillatory alpha power is sensitive to listening demands, likely reflecting top-down

attentional modulation (Dimitrijevic et al., 2017; Herrmann et al., 2023; Kraus et al., 2023b; Obleser et

al., 2012; Paul et al., 2021; Petersen et al., 2015; Weisz et al., 2011; Wöstmann et al., 2015). However,

in studies using highly controlled stimulus designs, in which task difficulty was manipulated without

changing stimulus acoustics, an increase in alpha power has only been observed sometimes, for

example, when relevant sound information occurs shortly after stimulus onset (Herrmann et al.,

2023), but not when relevant sound information occurs late in a stimulus and when visual stimuli are

presented concurrently (Kraus et al., 2023b). Manipulating motivation orthogonally to task demands

may help identifying the neural correlates of cognitive investment during listening. That is, neural

activity in brain regions that mediate top-down control of attention (e.g., cingulate cortex) is

modulated by motivation (Small et al., 2005). Consistently, occipital alpha power in a visual search

task has been sensitive to monetary rewards (Sawaki et al., 2015), raising the possibility that alpha

power may be modulated by a person's motivational state during listening, and, in turn, providing

an opportunity to identify the neural mechanisms underlying challenging listening.

Apart from neural signals possibly indexing attention networks directly, neural activity elicited

by auditory stimuli may provide additional insights into motivation influences on neural processing

during listening. Auditory sensory responses around 100 ms after stimulus onset are enhanced as

the saliency of a sensory input increases (for a review: Näätänen & Picton, 1987) and when individuals

attend to, compared to ignore, auditory stimuli (Hillyard et al., 1973) but effects of motivation have

not been observed (Goldstein et al., 2006; Krebs et al., 2013). Motivation rather seems to modulate

neural signals about 300 ms after stimulus onset (Baines et al., 2011; Goldstein et al., 2006; van den

Berg et al., 2014).

The current study will use magnetoencephalography (MEG) to investigate the

neurophysiological changes associated with listening demands under different motivational states

(using a reward manipulation). We will use pupil-size data as a benchmark for modulatory influences

of motivation on cognitive investment during listening (Kraus et al., 2023a). Using MEG, we will

investigate whether reward prospect impacts auditory sensory processing and how reward prospect

influences attentional processing under varying task demands (alpha power). In detail, we expect

increased attentional processing under high motivation when listening is hard, but no additional

motivational boast when listening is easy (see Fig. 1C). Additionally, we will investigate the

relationship among changes in pupil size, sensory responses, and neural alpha power and to

behavioral listening outcome.

Methods

Participants

Thirty-seven adults aged between 18 and 34 participated in the current study (mean = 23.43 years;

SD = 4.05 years; 11 males and 26 females; all right-handed). None of the participants reported having

a history of neural disorders or hearing problems. Participants gave written informed consent before

participation and received an honorarium of $15 CAD per hour. Their motivation was manipulated

through financial rewards. Participants could earn an additional $15 CAD depending on their

behavioral performance. The study was conducted at the Rotman Research Institute at the Baycrest

Academy for Research and Education in Toronto, Ontario, Canada. The study protocols are in

accordance with the Declaration of Helsinki and were approved by the local ethics committee of the

Rotman Research Institute at the Baycrest Academy for Research and Education in Toronto, Ontario,

Canada.

Experimental environment

MEG data were recorded with a 275-channel axial gradiometer MEG device (CTF-MEG, Coquitlam,

BC, Canada) in an electromagnetically shielded room. Participants were placed in a seating position

with their head centered in the MEG helmet. A back-projection screen was placed about 70 cm in

front of participants. An eye-tracking camera (Eyelink 1000 Plus, SR Research) was mounted to the

screen. The experimental stimulation was controlled by a laptop (Windows 10) running Psychtoolbox

(version 3.0.14) in MATLAB. Sounds were presented via an external sound card (RME Fireface UCx ii)

and delivered binaurally via MEG-compatible insert earphones (EARTONE 3A, 3M). A button response

box was used to record participants' behavioral responses. After the MEG experiment, the location

of the three fiducials (left tragus, right tragus, nasion) and the shape of the head were digitized using

a 3-D digitizer (Polhemus Fastrak).

Experimental design

The experimental design was similar to the one used in Kraus et al. (2023a). An auditory gap-

detection task was chosen because the timing and degree of listening demand can be more tightly

controlled than for speech materials that have been used in previous studies (Kraus et al., 2023a). In

fact, our previous work shows effects of motivation on behavior and pupil size, and a critical

interaction in line with theoretical frameworks (Brehm & Self, 1989; Richter et al., 2016; Figure 1C),

which studies using speech materials have not been able to observe. The experimental control

afforded by the auditory gap-detection task (see also Henry et al., 2014, 2017; Henry & Obleser,

2012; Herrmann et al., 2023; Kraus et al., 2023a, 2023b) facilitates the analysis of the interaction of

motivation, listening demand and listening outcome (Kraus et al., 2023a).

Participants were presented with 5.2-s white-noise sounds that each contained one gap (Fig.

1) and pressed a button with the right index finger on the response box as soon as they detected

the gap. The gap could occur at one of 70 randomly selected time points between 2.2 and 4.2 s after

noise onset (linearly spaced). Auditory stimuli were presented at 45 dB above the participant's

sensation level, which was determined for each ear prior to the experiment using an audiometer.

There was no visual stimulation during the presentation of the white noise. Participants were asked

to keep their gaze on the black screen and not close their eyes.

The 2 × 2 design included the factors of task difficulty (easy, hard) and reward prospect

(reward-irrelevant, reward-relevant). Task difficulty was manipulated via the duration of the gap in

the noise sound. In the hard condition, gap duration was individually titrated to about 65% gap-

detection performance in training blocks prior to the main experimental blocks (4–6 training blocks

of 2 min each). Gaps in the easy condition were twice as long (Kraus et al., 2023a, 2023b).

Figure 1. Experimental design. A: Auditory gap-detection task: Participants' task was to detect a gap within 5.2 s white-noise sound. The gap occurred randomly between 2.2–4.2 s post noise onset (gap window, marked by the gray line). **B**: Two-by-two design: Task difficulty was determined by the gap duration (titrated to 65% detection performance for the hard condition and twice as long in easy trials). An auditory cue 1 s before each trial indicated whether the upcoming trial was reward-relevant or not. **C**: Hypothesis: According to the Motivation Intensity Theory (gray lines; Brehm & Self, 1989; Richter, 2016), participants should invest cognitively under the hard listening condition only when motivated to succeed (solid line), but should give up investing resources when they are less motivated (dashed line). The colored dots show our hypothesis. Figure design follows Kraus et al., (2023a).

Half of the trials were paired with the possibility of receiving an additional financial reward

based on the participant's performance on these trials (reward-relevant trials). In contrast, the other

half of the trials were not paired with any reward (reward-irrelevant trials). Specifically, after the

experiment, three trials for each of the two difficulty levels were chosen from the reward-relevant

trials (Cole et al., 2022; Kraus et al., 2023a; Teoh et al., 2020; Tusche & Hutcherson, 2018). Participants

could gain $15 CAD in addition to their hourly compensation rate if their average performance across

these six trials was above 80%. All of the participants reached this threshold and got the reward.

Trials were presented in 14 blocks, each containing 20 trials. In half of the blocks, task difficulty

was easy, whereas task difficulty was hard in the other half. Easy and hard blocks alternated. The

difficulty of the starting task was counterbalanced across participants. At the beginning of each block,

participants received written information about the task difficulty (easy or hard) of the upcoming

block. Hence, participants had prior knowledge about whether detecting the gap would be easy or

hard. In random order, half of the 20 trials per block were reward-relevant trials, whereas the other

half were reward-irrelevant trials.

An auditory cue consisting of a guitar or a flute sound occurred before each white noise

sound, indicating whether a trial was reward-relevant or reward-irrelevant. The pairings of guitar and

flute sound to reward-relevant and reward-irrelevant trials were counterbalanced across participants.

Participants familiarized themselves with the cue-reward association during training trials before the

experiment. Overall, the experiment contained 70 white-noise sounds per Task difficulty (easy, hard)

× Reward prospect (reward-irrelevant, reward-relevant) condition, resulting in 280 trials.

After the experiment, participants completed a questionnaire regarding their use of the

reward cues. They first indicated which auditory cue was associated with reward-relevant trials. We

then asked them to rate the following statement: "I used the auditory cues (guitar and flute) to

distinguish between important and unimportant conditions" on a 6-point scale ranging from

"strongly disagree" over "disagree", "somewhat disagree", "somewhat agree", "agree" to "strongly

agree". Participants rated this statement twice, separately for the easy and the hard conditions.

Analysis of behavioral data

Any button press within 0.1 to 1.5 s after gap onset was defined as a hit (coded 1). Trials for which

no button was pressed within this time window were considered a miss (coded 0). The time between

the gap onset and the button press was calculated as the response time. The time at which the gap

occurred within the sound was included in the statistical modeling (see Statistical Analysis) to account

and test for an expected hazard effect (higher accuracy and faster response times for later compared

to early gaps relative to white-noise onset; Herrmann et al., 2023; Niemi & Näätänen, 1981; Nobre

et al., 2007). Response-time data were log-transformed for statistical analysis to obtain values closer

to a normal distribution.

Pupil data recording and preprocessing

Eye movements and pupil size of the right eye were continuously recorded using an Eyelink 1000

Plus eye tracker (SR research). Data was recorded as an additional channel of the MEG data

acquisition. MATLAB (MathWorks, Inc.) was used for the preprocessing and analyses of the data. To

detect eye blinks, we determined the threshold in the pupil channel above which the eye-tracker had

lost the pupil in case of a blink. All datapoints above this threshold were coded as invalid (NaN).

Additionally, 100 ms before and 200 ms after above threshold time points were also coded as NaN.

NaN-coded data points were linearly interpolated. Pupil data were subsequently low-pass filtered at

4 Hz (Butterworth, 4th order) and divided into epochs ranging from -2 to 6.2 s time-locked to white-

noise onset. An epoch was excluded if more than 40% of the trial had been NaN-coded prior to

interpolation. If more than 50% of the trials in any condition were excluded, the full pupil dataset of

the respective participant was excluded from analysis (N=12). The high number of excluded pupil

datasets is in part due to the technical challenge combining the concurrent recording of pupil size

and MEG. Pupil-size data were downsampled to 50 Hz. The polarity of the pupil data was inverted at

the analysis stage because the pupil recordings through the MEG were inverted (we determined this

in a brief black vs. white screen experiment to elicit dilation vs constriction, respectively). For each

trial, the mean pupil size in the -1.5 to -1.1 s time window was subtracted from the pupil size at every

time point of the epoch (baseline correction). This baseline time window was chosen to avoid

contamination of the auditory cue which was presented at -1 s. For each participant, single-trial time

courses were averaged, separately for each condition.

Pupil size analysis

For the statistical analysis (described in the Statistical analysis section) of the effects of task difficulty

and reward prospect, pupil-size data were averaged across the time window ranging from 2.2 s

(onset of gap window) to 6.2 s (end of trial).

Similar to our previous approach (Kraus et al., 2023a), we investigated whether a smaller pupil

size is associated with an increased probability that a participant misses a gap or responds slower to

the gap. Therefore, pupil-size data (baseline-corrected to -1.5 to -1.1 s to noise-onset) were time-

locked to the gap onset. For statistical analysis, pupil size was averaged across the -.5 to 0 s time

window, time-locked to gap onset.

To illustrate the association between pupil size and response time (Fig. 3E), we created two

groups of trials: fast and slow trials. Fast trials were defined as those with response times equal to or

faster than 0.75 SD of the mean. Slow trials were defined as those with response times equal to or

slower than 0.75 SD of the mean. The threshold was chosen as a good compromise between too

many and too few trials per group. The grouping was done separately for reward-relevant and

reward-irrelevant trials to ensure the same number of trials were used for each group and reward

condition. Additionally, we included only trials into the group of fast trials if their gap-onset time

matched with one trial of the slow trials and vice versa to ensure that there was no difference in gap-

onset times between slow and fast trials. The pupil size of the averaged slow versus fast trials was

compared using a one-sample t-test per time point. To account for multiple comparisons, p-values

were corrected across time points using a false discovery rate (FDR) of q = 5% (Benjamini &

Hochberg, 1995).

MEG recording and preprocessing

Magnetoencephalography (MEG) data were recorded using a CTF-MEG (Coquitlam, BC, Canada) with

275 axial gradiometers at a sampling frequency of 1200 Hz.

MEG data analysis was performed with the Fieldtrip MATLAB toolbox (2019-09-20;

Oostenveld et al., 2011). To avoid numerical issues during processing due to very small numbers

associated with MEG recordings in the Tesla range, we multiplied all MEG data with 10^{12}, leading to

data in the picotesla range. Data were filtered with a 100 Hz low-pass (Hann window, 89 points), a

0.7 Hz high-pass (Hann window, 2869 points), and a 60 Hz elliptic band-stop filter to suppress line

noise. Data were cut into 10.2-s trials ranging from -3 to 7.2 s time-locked to noise onset. For the

calculation of independent component analysis (ICA), these trials were divided into 1-s snippets.

Components containing blinks, lateral eye movement, and heart-related activity were identified

through visual inspection. The continuous data (filtered) were projected to ICA space using the

unmixing matrix. The previously identified components containing artifacts were removed from the

continuous data and the data were then back-projected to the original 275 sensors using the mixing

matrix. Data were then low-pass filtered at 30 Hz (Hann window, 111 points) and divided into trials

of 10.2 s (-3 to 7.2 s time-locked to noise onset). Data were downsampled to 600 Hz, and trials that

exceeded a signal change of more than 4 picotesla in any of the MEG channels were excluded from

analysis.

Analysis of event-related fields for gap onset and button response

Data were transformed from axial to planar gradiometers (Vrba & Robinson, 2001). Planar

gradiometers show the strongest sensitivity to sources that originate from directly below them

(Hämäläinen, 1995; Vrba & Robinson, 2001), which makes the interpretation of topographical

distributions more intuitive. The signal at the two planar gradiometers forming one pair were

combined by calculating the sum. This resulted in signals at 275 sensors.

For the analysis of event-related fields (ERFs), we focused on hit trials and each trial was time-

locked to the respective gap-onset time. To investigate evoked auditory sensory processing, data

were averaged across temporal sensors (MRF67, MRT13, MRT14, MRT15, MRT23, MRT24, MRT25,

MRT2, MRT35, MRT36, MLF67, MLT13, MLT14, MLT15, MLT23, MLT24, MLT25, MLT26, MLT35,

MLT36). For illustrative purposes, trials were averaged separately for each of the four conditions and

baseline-corrected by subtracting the mean amplitude in the time window before gap onset (-.5 to

0 s) from each time point. For the statistical analysis, data were averaged across the time window of

the M100-component of the ERF (0.09 to 0.13 s; Näätänen & Picton, 1987). The M100 time window

was selected to investigate possible reward-prospect influences on auditory sensory processing.

For an exploratory analysis of the late ERF-component with a parietal topography (Figure 4A

& C), hit trials were time-locked to a person's response time in the respective trial and averaged

across parietal sensors (MLC16, MLC17, MLC24, MLC25, MLC31, MLC32, MLC41, MLC42, MLC53,

MLC54, MLC55, MLC61, MLC62, MLC63, MRC16, MRC17, MRC24, MRC25, MRC31, MRC32, MRC41,

MRC42, MRC53, MRC54, MRC55, MRC61, MRC62, MRC63, MLP11, MLP12, MLP21, MLP22, MLP23,

MLP31, MLP32, MLP33, MLP34, MLP35, MLP41, MLP42, MLP43, MLP44, MLP45, MLP54, MLP55,

MLP56, MLP57, MRP11, MRP12, MRP21, MRP22, MRP23, MRP31, MRP32, MRP33, MRP34, MRP35,

MRP41, MRP42, MRP43, MRP44, MRP45, MRP54, MRP55, MRP56, MRP57, MZC03, MZC04, MZP01).

For illustrative purposes, data were baseline-corrected using the same time window as for the gap-

locked data (-.5 to 0 s time-locked to gap onset). For statistical analysis, single-trial response-locked

data were averaged across the 0.1 to 0.6 s time window (time-locked to button response) to

investigate task difficulty and reward-prospect effects on the response-related component (see Fig.

4D).

Analysis of time-frequency power

In order to analyze oscillatory activity in the alpha frequency range (8 to 12 Hz; Jensen & Mazaheri,

2010; Klimesch et al., 2007; Weisz et al., 2011), single-trial time-domain data (after planar gradiometer

calculation) were convolved with Morlet wavelets. Complex wavelet coefficients were calculated for

frequencies ranging from 8–12 Hz (1 Hz steps) and time points ranging from –2 s to 6.2 s (0.04 s

steps) time-locked to noise onset, separately for each trial, sensor, and participant. For the

visualization of time-frequency data across a wider frequency range, calculations were also done for

frequencies ranging from 1–20 Hz in steps of 1 Hz and in time steps of 0.16 s. We analyzed data first

time-locked to noise onset and then time-locked to the gap onset.

For the analysis of data time-locked to noise onset (Fig. 5), power was calculated by squaring

the magnitude of the complex wavelet coefficients separately for each trial, sensor, and time-

frequency bin. Data for the two corresponding planar gradiometers of a pair were combined by

calculating the sum, which resulted in power data for 275 sensors. For visualization purposes, time-

frequency power per condition was averaged across trials, and baseline-corrected to dB power

change: Data at each time point were divided by the mean power in the baseline time window (-1.5

to -1.1 time-locked to noise onset), and subsequently log10 transformed. To visualize the difference

in power between hit and miss trials, trials were categorized as hit and miss trials and averaged across

parietal sensors. For statistical analysis, parietal alpha power in the 1.7 to 2.2 s time window (time-

locked to noise onset) was averaged. This time window was chosen because it precedes the time

window during which a gap could occur (2.2–4.2 s).

For the investigation of alpha activity changes related to gap detection, we time-locked the

data to the gap onset and focused on hit trials only. This was done on single-trial complex wavelet

coefficients, and power calculation and gradiometer combination were performed in the same way

as for the noise-locked data above. For visualization purposes, condition-averaged data were

baseline-corrected to dB power change using the same time window as for the noise-locked data (-

1.5 to -1.1 s time-locked to noise onset) and separately averaged across parietal or temporal sensors

(see section above). For the statistical analysis, pre-gap alpha power for each trial was averaged

across the -0.5 to 0 s time window (time-locked to gap onset).

Trials in which participants detected the gap elicited an alpha-power suppression (Fig. 4 D).

We thus also investigated possible effects of task difficulty and reward prospect on the post-gap

alpha-power suppression. To this end, we calculated the latency of minimal alpha power between 0

and 1 s post-gap for each condition and participant. For the statistical analysis of the alpha power at

its minimum, we averaged the data ±40 ms around the estimated latency for each trial.

Source localization

For source localization, the MRI image of the standard brain and skull in Fieldtrip toolbox was non-

linearly warped for each participant to fit their head-shape data from the Polhemus digitization. An

automatic freesurfer's approach was used to segment the MRI and extract the cortical mesh. This

procedure ensured that the cortical mesh consisted of comparable grid-point locations across

participants (Alavash et al., 2021) in accordance with the Human Connectome Project (HCP) standard

atlas template (Glasser et al., 2017; Keitel & Gross, 2016). This cortical mesh served as the source

model for each participant. Fieldtrip's 'singleshell' method was used to calculate the inner skull

surface that was used as a volume conductor model. Subsequently, leadfields were calculated

separately for each experimental block to account for the slightly different head positions relative to

the sensors in the different MEG recording blocks.

For each participant and block, a multi-taper (±2Hz spectral smoothing) fast Fourier

transform (FFT) was calculated. The complex coefficients at 10 Hz that resulted from the FFT were

used to calculate a cross-spectral density matrix per block. Dynamic imaging of coherent sources

was used for source localization (DICS, Gross et al., 2001). For each source location, a spatial filter

was calculated using a participant's cross-spectral density matrix (dipole orientation: axis of most

variance using singular value decomposition). For each trial, spatial filter weights were multiplied

time point per time point with the complex wavelet coefficients from the analysis that used data

time-locked to the gap onset (8–12 Hz frequency range). Power was calculated by squaring the

magnitude of the complex wavelet coefficients that were projected through the spatial filter. Time-

frequency source power was averaged across the 8 to 12 Hz frequencies.

For the statistical analysis, we focused on two predefined regions of interest (ROI) according

to functional parcels defined by the Glasser atlas as available in the Human Connectome Project

workspace (Glasser et al., 2017; Keitel & Gross, 2016). We included an auditory ROI due to the

auditory nature of the task and because auditory stimulation is known to elicit sensory alpha activity

(Herrmann et al., 2023; Kraus et al., 2023b; Mazaheri et al., 2014; Wöstmann et al., 2017). We also

included a parietal ROI, because the parietal cortex is part of the attentional networks and is known

to elicit alpha activity under task load (Banerjee et al., 2011; Behrmann et al., 2004; Herrmann et al.,

2023; Kraus et al., 2023b; Rushworth et al., 2001). To analyze post-gap alpha-power suppression in

source space, we calculated the latency of minimal alpha power between 0 and 1s post-gap per

condition, participant, and ROI. For each trial, we then averaged the data ±40 ms around the

estimated latency.

For visualization only, hit trials were averaged for each condition and participant. The dB

power change was calculated relative to the original baseline time window (-1.5 to -1.1 s time-locked

to noise onset).

Statistical analysis

A paired sample t-test was used to analyze the subjective rating of cue use. The effect size was

reported as Cohen's d (J. Cohen, 1988).

A variety of linear mixed-effect models in R (v4.1.2), with the packages lme4 and sjPlot, were

used for the statistical analyses. All linear mixed-effect models included participant-specific random

intercepts to account for individual differences in the respective dependent variable. Single-trial

accuracy data were binary. Hence, for accuracy data, a generalized linear mixed-effect model (GLMM)

with a binomial distribution and a logit link function was used (Kraus et al., 2023a; Tune et al., 2021).

All other models included continuous data as dependent variables, and we thus used a linear mixed-

effect model (LMM) with a Gaussian distribution and an identity link function (Kraus et al., 2023a;

Tune et al., 2021). All continuous variables were z-scored. All models predicting behavioral outcomes

included the gap time as a regressor to account for hazard-rate effects (Herrmann et al., 2023; Niemi

& Näätänen, 1981; Nobre et al., 2007). For all statistical models using neurophysiological data as

dependent variable, data averaged across the baseline time window were included in the model as

a separate regressor instead of using baseline-corrected data as the dependent variable (Alday,

2019).

First, linear mixed-effect models were calculated to analyze the influence of the two

experimental factors on behavioral outcomes, pupil size, M100-component, and alpha power. The

models included effects of task difficulty (easy, hard), reward prospect (reward-irrelevant, reward-

relevant), and task difficulty × reward prospect interaction. Task difficulty and reward prospect were

categorical predictors and were, therefore, deviation-coded (i.e., -0.5 [easy, no-reward] and 0.5 [hard,

reward]).

Second, linear mixed-effect models were calculated to analyze the influence of pupil size or

parietal alpha power on accuracy or response time. For these analyses, only hard trials were used

because of the large variance in the behavioral measures between easy and hard conditions and the

ceiling performance for the easy condition. Pupil size and parietal alpha power were used as

independent variables in these models; therefore, baseline correction was needed before inclusion

into the respective statistical model. Including the baseline data as an additional regressor (as

described for the prediction of pupil/alpha data) would lead to collinearity issues. For pupil-size data,

we used subtractive baselining (see Pupil size recording and preprocessing), which can be used on

single trials (Mathôt et al., 2018). However, dB-power-baselining should not be used on single-trial

data (M. X. Cohen, 2014). To account for this, we calculated a model with pre-gap alpha power as

the dependent variable and alpha power within the baseline time window as the only independent

variable. The residuals of this model served as the alpha power regressor in the models to predict

behavioral outcomes (Alday, 2019).

To disentangle associations of pupil size/alpha power and behavior at the trial-by-trial state

level (i.e., within-participant) from associations at the trait level (i.e., between-participants), we

included two separate regressors associated with changes in pupil size/alpha power. The between-

participants regressor contained trial-averaged pupil size/alpha power per individual, whereas the

within-participant regressor contained the single-trial pupil size/alpha power relative to the

individual mean (Bell et al., 2019; Kraus et al., 2023a, 2023b; Tune et al., 2021). Reward prospect and

gap time were included as regressors to account for their potential influence on behavior. A random

slope for the within-participant effect was included whenever it improved the respective model and

did not lead to missing convergence of the model.

Third, a linear mixed-effect model was calculated to analyze the relationship between the

M100 amplitude and the post-gap alpha power. Reward prospect, task difficulty, and their interaction

were included as regressors to account for their potential influence on post-gap alpha power. A

within-participant and a between-participants regressor were included for the M100-amplitude

regressor as well as a random slope for the within-participant effect.

Fourth, linear mixed-effect models were calculated to analyze the relationship between the

pre-gap pupil size and the M100-amplitude or the post-gap alpha power. Reward prospect, task

difficulty and their interaction were included as regressors to account for their potential influence on

the M100-amplitude or post-gap alpha power. A within-participant and a between-participants

regressor were included for the pupil size regressor, as well as a random slope for the within-

participant.

For the interpretation of effects analyzed with the different linear mixed-effect models, we

calculated Bayes Factors (BF) as log(BF) = [BIC(H0)-BIC(H1)]/2, with BIC being the Bayes–Schwartz

information criterion (Alavash & Obleser, 2024; Wagenmakers, 2007). To this end, we calculated the

BIC for the full model, including the regressor of interest (H1), and for the reduced model, excluding

the regressor of interest (H0). A log(BF) larger than 1 provides evidence for the presence of an effect

of the regressor of interest, whereas a log(BF) value smaller than –1 suggests the absence of an effect

of the regressor of interest (Dienes, 2014).

Data availability

408   All data and analysis scripts are available at https://osf.io/prwb7/.

<center>**Results**</center>

Reward prospect improves performance

As intended, performance was better in the easy compared to the hard condition for both accuracy

and response time (accuracy: GLMM; odds ratio (OR) = .05, std. error (SE) = .004, $p = 1.74 \times 10^{-238}$,

log(BF) = Inf; response time: LMM; β = 0.35, SE = .007, $p < 1.74 \times 10^{-238}$, log(BF) = Inf; Fig. 2).

Furthermore, participants were more accurate and faster for the reward-relevant compared to the

reward-irrelevant condition (accuracy: OR = 1.31, SE = .117, $p = 3.54 \times 10^{-3}$, log(BF) = -.15; response

time: β = -.03, SE = .007, $p = 8.87 \times 10^{-5}$, log(BF) = 3.35). The interaction was not significant (accuracy:

odds ratio (OR) = 0.98, SE = .175, p > .9, log(BF) = -4.6; response time: β = -.02, SE = .014, p > .2,

log(BF) = -3.75). We found a hazard-rate effect, such that individuals were more accurate and faster

when the gap occurred later rather than earlier (accuracy: OR = 1.21, SE = .036, p = 3.19 × 10^{-10},

log(BF) = 15.8; response time: β = -.08, SE = .003, p = 2.44 × 10^{-118}, log(BF) = 263.5).

Participants rated their use of the auditory cue to distinguish between reward-relevant and

reward-irrelevant higher in the hard compared to the easy condition (t_{36} =3.54, p = 1.11 × 10^{-3}, d =

.58).

Figure 2. Behavioral results. A: Accuracy. Proportion correct was better for the easy compared to the hard condition and for the reward-relevant compared to the reward-irrelevant condition. Insets: 45-degree scatter plots showing the task difficulty (left) and the reward prospect effect (right) from linear mixed-model analysis. Difference plots (y minus x axis) are shown in the upper right corners). **B**: Response time. Participants were faster for the easy compared to the hard condition and for the reward-relevant compared to the reward-irrelevant condition. Insets: 45-degree scatter plots showing the task difficulty (left) and the reward prospect effect (right) from linear mixed-model analysis. Difference plots (y minus x axis) are shown in the upper right corners).

Reward prospect modulates the task difficulty effect on pupil size

To investigate the influence of task difficulty and reward prospect on pupil size, we averaged pupil

size over the time window ranging from 2.2 to 6.2 s (Fig. 3B; time-locked to noise-onset). Pupil size

was larger for hard compared to easy trials (β = .05, SE = .009, p = 5.21 × 10^{-7}, log(BF) = 9.1) and

larger for reward-relevant compared to reward-irrelevant trials (β = .03, SE = .009, p = 3.32 × 10^{-4},

log(BF) = 2.6). Importantly, we observed a significant task difficulty × reward prospect interaction,

indicating a larger increase in pupil size for reward-relevant than reward-irrelevant trials when task

difficulty was hard compared to easy (β = .04, SE = .018, p = 3.08 × 10^{-2}, log(BF) = -1.9; Fig. 3C).

432   Next, we investigated whether a larger pupil size during the interval preceding the gap is

associated with behavioral performance. This analysis was conducted using the hard trials only and

the information about the reward conditions was included as a regressor into the model. With this

approach we controlled for the effects of reward prospect and task difficulty on both pupil size and

response time. To this end, pupil-size data for hard trials were aligned to gap-onset times, and the

pupil size was averaged across the -0.5 s to 0 s pre-gap time window. Pre-gap pupil size did not

predict behavioral accuracy, neither at the trial-by-trial state level (within-participant: OR = 1.07, SE

= .043, p > .1, log(BF) = -2.7) nor at the trait level (between-participants: OR = 1.04, SE = .158, p > .8

, log(BF) = -1.6). However, at the trial-by-trial state level response times were shorter when pre-gap

pupil size was larger (within-participant: β = -.02, SE = .008, p = 9.03 × 10^{-3}, log(BF) = 0), but at the

trait level there was no association between a participant's mean pupil size and mean response time

(between-participants: β = -.01, SE = .038, p > .8, log(BF) = -1.6; Fig. 3F). To illustrate this effect, we

grouped the hard trials into fast and slow response trials, separately for the reward-relevant and the

reward-irrelevant condition (trials were selected such that the mean gap time for fast and slow

response trials did not differ; see Methods section). Pupil size was significantly larger for fast trials

compared to slow trials already more than 1 s before the gap onset (Fig. 3E).

In addition, we conducted a temporal response function analysis (Crosse et al., 2016) to

ensure that the observed interaction of task difficulty and reward prospect is not resulting from a

temporal shift in the pupil-size response due to response-time differences between conditions (see

Fig. 2B). The time points of the noise onset, gap onset, and response time of the respective trial

served as separate regressors to model the temporal response functions (TRF) to the different events.

The TRF to the gap onset (Fig. 3G left) but not to the button response (Fig. 3G right) shows temporal

variability between the different experimental conditions. However, importantly, we observed the

interaction of task difficulty and reward for both the TRF to gap onset (1.82 s to 2.56 s) and to the

button response (1.38 s to 2.22 s). Additionally, the pupil-size TRF was larger for hard compared to

easy trials (TRF to gap onset: -2 s to -0.88 s and 1.36 s to 3.14 s; TRF to button response (0.96s to

2.54 s).

Figure 3. Pupil size results. A: Averaged pupil-size time courses across participants per condition. Error bands reflect the within-participants error. Gray areas indicate time window during which a gap could occur, from 2.2 to 4.2 s. **B**: Averaged data for 2.2–6.2 s time window. Error bars indicate the standard error of the mean. **C**: 45-degree scatter plots illustrate the interaction. Left: Data from the easy condition. Right: Data from the hard condition. Colored dots show averaged pupil data per task-difficulty level, separately for each participant. The 45-degree line indicates no difference between conditions. Crosshairs indicate the 95% confidence interval (CI). Difference plots (y- minus x-axis) are shown in upper right corners. **D**: Pupil-size time courses (time-locked to gap onset) for the hard condition, separately for each reward-prospect condition (light vs. dark) and for hit and miss trials (solid vs. dashed lines). Error bands reflect the within-participants error. **E**: Pupil-size data grouped into trials with slow (dashed) and fast response times (solid). The histogram shows the distribution of response times for each group. Gray area indicates the time window

for the linear mixed-effect model analysis in panel F. Black line indicates the time window in which the pupil size was significantly larger for fast-response trials compared to slow-response trials (after FDR-correction). Error bands reflect the within-participants error. **F**: Effect of pupil size on response time in a linear mixed-effect model analysis. A larger pupil size is associated with a faster response time on a within-participant level (not for the between-participants effect). Participant-specific slope for pupil-size did not improve the model, but we show it here for illustrative purposes. **G:** TRF-approach of the pupil-size response to the gap onset (left) and button press (right). Black line indicates the time window in which the pupil size increased stronger for reward-relevant than reward-irrelevant trials when task difficulty was hard compared to easy. Dashed black line indicates the time window in which the pupil size was significantly larger in hard compared to easy trials. All shown significant effects are based on FDR-correction.

Task difficulty, but not reward prospect, affects the auditory sensory response

We tested whether task difficulty and reward prospect influenced sensory responses, that is, responses in the auditory cortex. To this end, we focused on the M100 component in the 0.09 to 0.13 s time window after gap onset in temporal sensors. The M100 amplitude was larger for easy (i.e., longer gap duration) compared to hard trials (i.e., shorter gap duration; $\beta = -.40$, SE $= .013$, p $= 1.33 \times 10^{-200}$, log(BF) $= 210.5$; Fig. 4A & B). The effect for reward prospect ($\beta = -.00$, SE $= .017$, p $> .9$, log(BF) $= -4.5$), and the interaction were not significant ($\beta = -.01$, SE $= .034$, p $> .9$, log(BF) $= -4.5$).

Task difficulty and reward prospect modulate the response-related evoked field

We also explored the late ERF component from 0.5 s onwards (Fig. 4A). Because the topography indicates the responses originate from the parietal cortex, the analysis of this component focused on parietal sensors. Sorting the gap-locked trials according to their response times (Fig. 4C) shows that this component is time-locked to the response rather than to the gap-onset (at 0 s in Fig. 4C), because the negative deflection succeeds the response time (dashed line). Hence, we analyzed this component time-locked to the button response (Fig. 4D) instead of time-locked to the gap-onset. Topographies for the time windows before, during, and after the button response are shown in Figure 4D & E, indicating that the response-related component is similar to the component we observed in

475   the gap-locked analysis (Fig. 4C). The response-related component was stronger deflected for hard

476   than easy trials (β = -.18, SE = .013, p = 1.54 × 10^{-39}, log(BF) = 83), and for reward-relevant than

477   reward-irrelevant trials (β = -.04, SE = .013, p = 1.33 × 10^{-3}, log(BF) = 1; Fig. 4E). The interaction was

478   not significant (β = -.05, SE = .026, p > .05, log(BF) = -2.5).

Figure 4. Event-related field to gap onset and to button response. A: Event-related field to gap onset. Data are averaged across temporal sensors marked in inset. Error bands reflect the within-participants error. Gray areas indicate time windows for analysis in B and C. **B**: Effect on M100. Average from .09 to .13 s. Top:

Topography for the average across all for conditions. Bottom: Statistical results are calculated using a linear mixed-effect model. Main effect is significant for difficulty but not for reward prospect. **C**: Late ERF-component. Top: Topography for the average across all four conditions from .5 to 1 s. Bottom: Trials sorted by response time and time-locked to the gap-onset. Average across parietal sensors that are marked in the inset in panel D. Data are from all conditions. Black dashed line indicates response time. **D:** Event-related field to button response. Data are averaged across parietal sensors marked in the inset. Error bands reflect the within-participants error. Gray areas indicate time windows for topographies on the right, and in panel E. Topographies for time window before button response and around button response show main activity around left motor areas. **E:** After-response-component. Top: Topography for the average across all four conditions from .1 to .6 s. Bottom: Statistical analysis of time window .1 to .6 s after button response using a linear mixed-effect model. Stronger deflection for hard compared to easy trials and for reward-relevant compared to reward-irrelevant trials. All significant effects are based on FDR-correction.

Neither reward prospect nor task difficulty modulate pre-gap parietal alpha power

The time-frequency analysis shows that parietal alpha power in the 8-12 Hz frequency band

increased over time after noise onset, up to when the gap occurred (Fig. 5C). However, alpha power

in the 1.7 to 2.2 s time window after noise onset (i.e., 0.5 s before any gap could occur) showed no

effect of task difficulty (β = .00, SE = .016, p > 0.9, log(BF) = -4.5), reward prospect (β = -.01, SE =

.016, p > 0.9, log(BF) = -4.5) nor the interaction: β = -.02, SE = .032, p > 0.9, log(BF) = -4.5). Grouping

trials according to hits and misses (irrespective of condition) showed the pre-gap alpha power

increase for hits but not for miss trials. For hit trials, alpha power also decreased after gap onset (Fig.

5D).

Alpha power time courses were time-locked to gap onset for an analysis of the exact time

window before gap onset. We investigated whether alpha power in the pre-gap time window (-0.5

to 0 s) was affected by our experimental manipulations. However, neither the effect of reward

prospect (β = -.01, SE = .016, p > .5 , log(BF) = -4.5), task difficulty (β = -.02, SE = .016, p > .5, , log(BF)

= -4), nor the interaction were significant (β = -.03, SE = .032, p > 0.5, , log(BF) = -4.5).

Although pre-gap alpha power was not modulated by task difficulty or reward prospect, we

found a positive correlation between pre-gap alpha power and behavior. The probability of detecting

a gap (for hard trials) was higher when pre-gap alpha power in parietal sensors was larger (Fig. 5E).

This effect was present on the within-participant level (OR = 1.12, SE = .037, p = 7.81 × 10⁻⁴, log(BF)

= 1.7), but not on the between-participants level (OR = .90, SE = .143, p > .5 , log(BF) = -1.6). There

were no significant alpha-power effects on response time (within-participant: β = .03, SE = .029, p >

0.2, log(BF) = -3.5; between-participants: β = .19, SE = .142, p > 0.2, log(BF) = -1.1).

Figure 5. Pre-gap alpha power analysis. All data are averaged across parietal sensors marked in the inset in panel C. **A**: Time-frequency power measured in dB relative to a pre-stimulus time interval and averaged across all conditions. **B:** Topography reflects alpha power in the 500 ms (1.7-2.2 s) before the time window when a gap could occur (averaged across all conditions; gray area in C). **C:** Alpha power averaged for all four conditions separately. Error bands reflect the within-participants error. Effects of task difficulty and reward

prospect were not significant in the 500 ms before a gap could occur (gray area). **D:** Alpha power for hit trials (top) and miss trials (bottom) sorted by gap onset. Black dashed lines indicate the gap time. Alpha power increased towards the gap but was suppressed after gap occurrence (for hit trials). **E:** Effect of pre-gap parietal alpha power on accuracy in a linear mixed-effect model analysis. Larger pre-gap alpha power is associated with better performance on a within-participant level only. Reported effects are based on FDR correction.

Reward prospect modulates post-gap alpha-power suppression

We further explored the alpha-power suppression that followed the gap. Figure 6A suggests the

alpha suppression is related to the gap rather than the button press, as illustrated by relation of the

two dashed lines to the alpha-power suppression in Figure 6A. Task difficulty led to a post-gap alpha-

power suppression that was only marginally significant for temporal sensors (β = -.04 SE = .018, p =

4.03×10^{-2}, log(BF) = -2, Fig. 6C) and not significant for parietal sensors (β = -.03, SE = .018, p > .2 ,

log(BF) = -3.5, Fig. 6D). However, alpha power was more suppressed in reward-relevant compared

to reward-irrelevant trials for temporal sensors (β = -.07, SE = .017, p = 3.33×10^{-4}, log(BF) = 3) and

parietal sensors (β = -.05, SE = .018, p = 1.53×10^{-2}, log(BF) = -.5). For neither of the two sensor

clusters was the interaction significant (temporal sensors: β = -.06, SE = .035, p > .1 , log(BF) = -3;

parietal sensors: β = -.03, SE = .036, p > .4 , log(BF) = -4). Projecting gap-locked alpha power data

into source space revealed qualitatively similar results to the results in sensor space, although not all

effects reached statistical significance in source space (see Fig. 7).

Correlating the M100-amplitude with the post-gap alpha-power suppression revealed a

positive relationship. When the M100-amplitude across temporal sensors was enhanced, subsequent

alpha power was less suppressed in both temporal (within-participant: β = .11, SE = .020, p = 2.74 ×

10^{-8}, log(BF) = 108; between-participants: β = .37, SE = .049, p = 6.26×10^{-13}, log(BF) = 12.8) and

517   parietal sensors (within-participant: β = .09, SE = .016, p = 2.74 × 10⁻⁸, log(BF) = 47; between-

518   participants: β = .20, SE = .076, p = 1.36 × 10⁻², log(BF) = 1.2).

Figure 6. Post-gap alpha power analysis. A: Gap-locked alpha power trials sorted by response time. Data is averaged across parietal sensors see inset in D. Black lines represent the gap onset and the response time. Alpha-power suppression is rather time-locked to the gap-onset than to the response time. **B:** Left: Grand average topography for -.5 to 0s time-locked to gap onset. Right: Grand average topography for maximal alpha-power suppression after gap onset. **C:** Temporal sensors. Left: Alpha power averaged per condition across participants and temporal sensors (see inset). Error bands reflect the within-participants error. Right: linear mixed-effect model results of alpha-power suppression. Alpha-power suppression was larger for hard compared to easy and for reward-relevant compared to reward-irrelevant trials. **D:** Parietal sensors. Left: Alpha power averaged per condition across participants and parietal sensors (see inset). Error bands reflect the within-participants error. Right: linear mixed-effect model results of alpha-power suppression. Alpha-power suppression was larger for reward-relevant compared to reward-irrelevant trials.

Figure 7. Source localized alpha power, time-locked to gap onset. A: Left: Grand average source plot for -.5 to 0 s time-locked to gap onset. Right: Grand average source plot for maximal alpha-power suppression after gap onset. **B:** Auditory ROI. Left: Source-projected alpha power for each condition. Inset shows the auditory ROI. Error bands reflect the within-participants error. Right: Results from a linear mixed-effect model predicting alpha-power suppression. Alpha-power suppression was larger for hard compared to easy trials (β = -.07, SE = .018, p = 9.64 × 10⁻⁴, log(BF) = 2.5). The effect of reward (β = -.02, SE = .018, p > .6, log(BF) = -4) and the difficulty × reward interaction (β = .02, SE = .036, p > .7, log(BF) = -4.5) were not significant. Direction of the effects is same as in sensor space. **C:** Same as in panel B for a parietal ROI. No significant effect was found (task difficulty: β = -.03, SE = 0.016, p > .7, log(BF) = -2.5; reward prospect: β = -.02, SE = .016, p > .1, log(BF) = -3.5; interaction: β = -.06, SE = .031, p > .1, log(BF) = -3) Direction of the effects is the same as in sensor space.

Larger pre-gap pupil size covaries with lower pre-gap parietal alpha power

To investigate the relationship between pupil size and brain measures, we calculated different linear

mixed-effect models (Fig. 8). First, we tested whether the pre-gap pupil size is associated with

auditory cortex responses to the gap (M100 amplitude; averaged across temporal sensors). There

was no within-participant effect (β = -.01, SE = .018, p > .6, log(BF) = -6) nor a between-participants

effect (β = -.07, SE = .053, p > .4, log(BF) = -1.73).

Second, we investigated whether the pre-gap pupil size is associated with pre-gap parietal

alpha power (averaged across parietal sensors). A larger pupil size was associated with lower parietal

alpha power on a within-participant level (β = -.05, SE = 0.02, p = 1.25 × 10^{-2}, log(BF) = 5.5; Fig. 8 B).

There was no significant effect on a between-participants-level (β = -.06, SE = .104, p > .6, log(BF) =

-1.64). Importantly, controlling for possible changes of alpha power over time via including the trial

number across the experiment as an additional regressor into the model had no effect on the results.

Figure 8. Relation between pupil size and brain measures. A: Relation between pre-gap pupil size and M100-amplitude. Linear mixed-effect modeling controlled for experimental conditions and was done on hit trials only. Pupil size was averaged across 0.5 s before gap-onset and baseline-corrected to the baseline window before the trial. M100 data was averaged across temporal sensors. No within-participant nor between-participants effect of pupil size on the M100-amplitude was found. **B:** Relation between pre-gap pupil size and pre-gap parietal alpha power. Linear mixed-effect modeling controlled for experimental conditions and was done on hit trials only. Pupil size was averaged across 0.5 s before gap-onset and baseline-corrected to the baseline window before the trial. Alpha power data was averaged across the same 0.5 s time window and across parietal sensors. A significant negative relationship between pupil size and alpha power was found on a within-participant level.

532   **Discussion**

In the current study, we investigated a listener's motivation affecting behavior, pupil size, and neural

activity during a challenging auditory detection task. Participants performed better, and pupil size

increased when listeners were more motivated compared to when they were less motivated by

reward prospects. Neural oscillatory activity in parietal regions increased throughout the

presentation of the noise sound, but this power increase was independent of task difficulty and

reward prospect. Auditory sensory responses to auditory target stimuli were not modulated by

reward prospect. However, both task difficulty and reward prospect independently enhanced post-

gap activity. In sum, pupil-size changes index cognitive investment during the interplay of listening

demand and motivation, whereas motivation impacts neural indices post-stimulus rather than

preparatory attentional modulation. The current study thus shows the complex and distinct impacts

of motivation and listening demands on different neurophysiological systems.

Motivational state moderates the effect of listening demand on pupil size

Pupil size increased as listening demands increased, as expected (Kadem et al., 2020; Kahneman &

Beatty, 1966; Koelewijn et al., 2012; Kraus et al., 2023b; Ohlenforst et al., 2018; Wendt et al., 2016;

Winn et al., 2015; Zekveld et al., 2010). Previous work has also shown that the pupil size is sensitive

to individuals giving up listening under impossible listening conditions (Herrmann & Ryan, 2024;

Ohlenforst et al., 2017; Zekveld & Kramer, 2014). The observation of the current study that a low

motivational state is associated with a smaller pupil size, especially under high listening demands

(see Fig. 3B), is consistent with the literature on 'giving up' listening. People may invest less

cognitively when they are little motivated (Brehm & Self, 1989; Herrmann & Johnsrude, 2020;

Koelewijn et al., 2018; Kraus et al., 2023a; Pichora-Fuller et al., 2016). A consideration in the current

study may be that response times differed between conditions and that response time temporally

aligned with the pupil dilation. However, the moderating effect of the motivational state on the task

difficulty effect on pupil size remains qualitatively the same when controlling for the potential

confound of the different response times (see Fig. 3 G).

558   Critically, the behavioral results by themselves suggest that the influence of motivation is

independent of the influence of listening demand (two main effects; Fig. 2). However, the pupil-size

results indicate, consistently with cognitive control frameworks (Brehm & Self, 1989; Parro et al., 2018;

Yee & Braver, 2018), that motivation is most impactful under difficult compared to easy listening

demands. Consequently, based on motivational intensity theory (Brehm & Self, 1989; Richter, 2016)

the individual motivational state is another important determinant of the amount of effort allocated

during listening besides the level of listening demand. The here and previously found increases in

pupil size and response speed with heightened motivation (Kraus et al., 2023a) highlights the

importance of taking listeners' motivation into account when assessing their invested effort. Our

pupil size data thus provide an important confirmation to examine motivational impacts on the

neural level.

Expectations about different levels of listening demand are not sufficient to modulate alpha

power

Parietal alpha power increased throughout the presentation of the noise up to when individuals

detected the gap (Fig. 5C). Alpha power thus showed sensitivity to when in time attention is allocated,

followed by a post-stimulus suppression (Herrmann et al., 2023). However, in contrast to previous

work (Obleser et al., 2012; Petersen et al., 2015; Wisniewski et al., 2017; Wöstmann et al., 2015) and

our hypothesis, we did not observe an increase in parietal alpha power with increasing task demand

nor when listeners could expect a reward for performing well. One important difference between our

and these former studies is that different tasks and time windows of interest were employed. Most

of the studies that report demand-related effects on parietal alpha power during listening observed

them in a memory task, during the retention period when participants held a stimulus in memory to

compare it with a later probe stimulus (Obleser et al., 2012; Petersen et al., 2015; Wisniewski et al.,

2017; Wöstmann et al., 2015). In the present study, we investigated alpha power changes in a period

– leading up to a behaviorally relevant auditory event – during which our experimental conditions

were acoustically identical (as opposed to some of the previous studies). It thus appears that top-

down knowledge about the difficulty of an upcoming event is insufficient to drive changes in parietal

alpha power (but also see Herrmann et al. (2023), indicating additional complexities). Palva & Palva

(2011) suggest that distinguishing between endogenous vs. exogenous task demands is critical to

understanding alpha-power dynamics. Manipulation of task demand via the saliency of a stimulus is

an exogenous manipulation (Obleser et al., 2012; Petersen et al., 2015; Wisniewski et al., 2017;

Wöstmann et al., 2015), whereas adjusting expectations about demand level is an endogenous

manipulation (present study). Therefore, we suggest that changes in alpha power due to listening

demand may only be observed under specific task and stimulus conditions but do not generally vary

with different levels of cognitive resource recruitment during listening.

Not auditory sensory processing but post-stimulus alpha power is modulated by motivational

state

Auditory sensory responses (M100 amplitude) were enhanced for a long compared to a short gap

duration – and thus for the more salient target – but not for variations in motivation. The M100 and

the N100, the electric equivalent, are known to respond most strongly to changes in physical stimulus

properties (Frank et al., 2020; Hansen & Hillyard, 1980; Näätänen & Picton, 1987; Paiva et al., 2016).

Nevertheless, attentional modulations of the M100/N100 have been observed previously but often

limited to spatial tasks (Ding & Simon, 2012; Hillyard et al., 1973; Kraus et al., 2021; O'Sullivan et al.,

2015; Orf et al., 2023; Woldorff & Hillyard, 1991). It appears, however, that the attentional boost

through motivation is insufficient to affect auditory sensory responses.

In contrast to the auditory sensory processing, we observed a response-related neural signal

that scaled with task difficulty and reward prospect. Research on the impact of reward on stimulus

processing often focuses on the feedback-related negativity (FRN). The FRN is a component that

peaks around 200 ms after the feedback about an earned or lost reward (Glazer et al., 2018; Holroyd

& Coles, 2002). The FRN is thought to scale with the degree to which the actual feedback deviates

from the expected feedback (Frömer et al., 2021; Sambrook & Goslin, 2015) and is modulated by

reward expectation (M. X. Cohen et al., 2007). In the present study, participants knew whether they

were in a reward-irrelevant or a reward-relevant trial and how difficult target detection was.

Moreover, the response participants made upon gap detection was the only feedback that they

received about possibly gaining a reward. Detecting a gap in hard trials may have led to a more

unexpected feedback signal (i.e., positive surprise) than in easy trials because participants knew that

it would be less likely for them to detect the gap in hard trials. This feedback signal may be greater

in the reward-relevant than the reward-irrelevant condition. We, therefore, interpret the observed

effect of motivation as a positive surprise related to the detection of the gap.

617   Although there was no change in pre-gap alpha power nor in auditory sensory processing

(M100) due to changes in motivational state, the post-gap alpha-power suppression was stronger

during high compared to low motivation, independently of task demand. The suppression of alpha

power is thought to be a sign of the facilitation, and therefore gating, of the processing of sensory

input (Jensen & Mazaheri, 2010; Klimesch et al., 2007; Palva & Palva, 2011; Pfurtscheller, 2003). In

visual attention tasks, reward prospect enhanced post-target encoding (Hall-McMaster et al., 2019)

and neural processing in attention networks (Small et al., 2005), suggesting motivation influences of

attentional top-down mechanisms on sensory processing. Both the response-related neural signal

and the post-stimulus alpha-power suppression suggest motivation affects top-down mechanisms

during effortful listening.

Discrepancy between changes in pupil size and alpha power

Both parietal alpha power and pupil size increased over the trial time course toward the gap, which

proved beneficial for behavior. However, the degree to which both measures were affected by our

experimental manipulations differed. Pupil size was greater for higher task demands and modulated

by motivation, whereas parietal alpha power prior to the gap did not change for either manipulation.

632   Research investigating the relations between both metrics is rare. The few studies that

observed simultaneous demand-driven dynamics in pupil size and alpha power manipulated the

demand via the stimulus itself, for example, by adding different levels of background noise to speech

or using noise-vocoded speech at different levels (Ala et al., 2020; McMahon et al., 2016; Miles et al.,

2017). As outlined above, the present study had no acoustic differences in the pre-gap analysis

window, which may explain the absence of alpha-power changes. In a recent study using the same

auditory paradigm in an audiovisual dual-task, we also found that changes driven by listening

demands were more prominent in pupil size than in parietal alpha power (Kraus et al., 2023b).

The current study also showed that pre-gap pupil size and pre-gap parietal alpha power are

negatively related (see also Kraus et al., 2023b). Although pupil size and alpha power were differently

sensitive to listening demand and motivation, a common underlying mechanism seems to affect

both pupil size and parietal alpha power but in opposite directions. Noradrenergic dynamics

originating from the locus coeruleus may be one candidate. Pupil size variations are correlated with

activity in the locus coeruleus (Aston-Jones & Cohen, 2005b), and the pupil dilates in response to

LC-stimulation (Liu et al., 2017). Increased noradrenergic activity emerging from LC is linked to

optimizing task performance and task engagement (Aston-Jones & Cohen, 2005a). The

noradrenergic influence on thalamic modulations (McCormick, 1998) and the thalamic connection to

cortical oscillatory activity (Steriade, 2000) provide evidence for a neural mechanism affecting both

pupil size and neural oscillatory alpha power (Dahl et al., 2022).

651   **Conclusion**

The present study confirms that pupil size robustly indexes cognitive investment during listening and

provides an integrated read-out of an individual's motivational state. However, the neural

mechanisms that encode listening demand and motivational state reveal a more nuanced picture:

Expectation of a more demanding or more important (i.e., reward-relevant) sensory input does not

suffice to elicit variations in neural alpha power. Auditory sensory processing was also not affected

by a listener's motivational state. Importantly, we observed that the known alpha-power suppression

in the wake of an auditory target event appears amplified under top-down motivational regulation.

In sum, while pupil size poses an integrated read-out of listening demand and motivation, both

dimensions selectively affect sensory and attentional post-processing aspects of auditory

neurophysiology.

662   **References**

Ala, T. S., Graversen, C., Wendt, D., Alickovic, E., Whitme, W. M., & Lunner, T. (2020). An exploratory
study of EEG alpha oscillation and pupil dilation in hearing-aid users during effortful listening
to continuous speech. *PLoS ONE*, *15*, 1–15. https://doi.org/10.1371/journal.pone.0235782
Alavash, M., & Obleser, J. (2024). Brain Network Interconnectivity Dynamics Explain Metacognitive
Differences in Listening Behavior. *The Journal of Neuroscience*, *44*(31),
https://doi.org/10.1523/JNEUROSCI.2322-23.2024
Alavash, M., Tune, S., & Obleser, J. (2021). Dynamic large-scale connectivity of intrinsic cortical
oscillations supports adaptive listening in challenging conditions. In *PLoS Biology* (Vol. 19,
Issue 10). https://doi.org/10.1371/journal.pbio.3001410
Alday, P. M. (2019). How much baseline correction do we need in ERP research? Extended GLM
model can replace baseline correction while lifting its limits. *Psychophysiology*, *56*(12), 1–14.
https://doi.org/10.1111/psyp.13451
Alfandari, D., Richter, M., Wendt, D., Fiedler, L., & Naylor, G. (2023). Previous Mental Load and
Incentives Influence Anticipatory Arousal as Indexed by the Baseline Pupil Diameter in a
Speech-in-Noise Task. *Trends in Hearing*, *27*. https://doi.org/10.1177/23312165231196520
Aston-Jones, G., & Cohen, J. D. (2005a). Adaptive gain and the role of the locus coeruleus-
norepinephrine system in optimal performance. *Journal of Comparative Neurology*, *493*(1), 99–
110. https://doi.org/10.1002/cne.20723
Aston-Jones, G., & Cohen, J. D. (2005b). An integrative theory of locus coeruleus-norepinephrine
function: Adaptive gain and optimal performance. *Annual Review of Neuroscience*, *28*, 403–
450. https://doi.org/10.1146/annurev.neuro.28.061604.135709
Baines, S., Ruz, M., Rao, A., Denison, R., & Nobre, A. C. (2011). Modulation of neural activity by
motivational and spatial biases. *Neuropsychologia*, *49*(9), 2489–2497.
https://doi.org/10.1016/j.neuropsychologia.2011.04.029
Banerjee, S., Snyder, A. C., Molholm, S., & Foxe, J. J. (2011). Oscillatory alpha-band mechanisms and
the deployment of spatial attention to anticipated auditory and visual targetlocations:
Supramodal or sensory-specific control mechanisms? *Journal of Neuroscience*, *31*(27), 9923–
9932. https://doi.org/10.1523/JNEUROSCI.4660-10.2011
Behrmann, M., Geng, J. J., & Shomstein, S. (2004). Parietal cortex and attention. *Current Opinion in*

Neurobiology, *14*(2), 212–217. https://doi.org/10.1016/j.conb.2004.03.012

Bell, A., Fairbrother, M., & Jones, K. (2019). Fixed and random effects models: making an informed

choice. *Quality and Quantity*, *53*(2), 1051–1074. https://doi.org/10.1007/s11135-018-0802-x

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)*,

57(1), 289–300.

Bijleveld, E., Custers, R., & Aarts, H. (2009). The Unconscious Eye Opener. *Psychological Science*,

20(11), 1313–1315. https://doi.org/10.1111/j.1467-9280.2009.02443.x

Brehm, J. W., & Self, E. A. (1989). The intensity of motivation. *Annual Review of Psychology*, *40*, 109–

131. https://doi.org/10.1146/annurev.ps.40.020189.000545

Cohen, J. (1988). Statistical power analysis for the behavioral sciences. Lawrence Erlbaum

Associates. *Hillsdale, NJ*, 20–26.

Cohen, M. X. (2014). Analyzing neural time series data: theory and practice. In *MIT press*.

https://doi.org/10.7551/mitpress/9609.001.0001

Cohen, M. X., Elger, C. E., & Ranganath, C. (2007). Reward expectation modulates feedback-related

negativity and EEG spectra. *NeuroImage*, *35*(2), 968–978.

https://doi.org/10.1016/j.neuroimage.2006.11.056

Cole, L., Lightman, S., Clark, R., & Gilchrist, I. D. (2022). Tonic and phasic effects of reward on the

pupil: implications for locus coeruleus function. *Proceedings of the Royal Society B: Biological*

Sciences, *289*. https://doi.org/10.1098/rspb.2022.1545

Crosse, M. J., Di Liberto, G. M., Bednar, A., & Lalor, E. C. (2016). The multivariate temporal response

function (mTRF) toolbox: A MATLAB toolbox for relating neural signals to continuous stimuli.

Frontiers in Human Neuroscience, *10*(NOV2016), 1–14.

https://doi.org/10.3389/fnhum.2016.00604

Dahl, M. J., Mather, M., & Werkle-Bergner, M. (2022). Noradrenergic modulation of rhythmic neural

activity shapes selective attention. *Trends in Cognitive Sciences*, *26*(1), 38–52.

https://doi.org/https://doi.org/10.1016/j.tics.2021.10.009

Dienes, Z. (2014). Using Bayes to get the most out of non-significant results. *Frontiers in*

Psychology, *5*, 1–17. https://doi.org/10.3389/fpsyg.2014.00781

Dimitrijevic, A., Smith, M. L., Kadis, D. S., & Moore, D. R. (2017). Cortical alpha oscillations predict

speech intelligibility. *Frontiers in Human Neuroscience*, *11*.

https://doi.org/10.3389/fnhum.2017.00088

Ding, N., & Simon, J. Z. (2012). Neural coding of continuous speech in auditory cortex during

monaural and dichotic listening. *Journal of Neurophysiology*, *107*(1), 78–89.

https://doi.org/10.1152/jn.00297.2011

Frank, M., Muhlack, B., Zebe, F., & Scharinger, M. (2020). Contributions of pitch and spectral

information to cortical vowel categorization. *Journal of Phonetics*, *79*, 100963.

https://doi.org/10.1016/j.wocn.2020.100963

Frömer, R., Nassar, M. R., Bruckner, R., Stürmer, B., Sommer, W., & Yeung, N. (2021). Response-

based outcome predictions and confidence regulate feedback processing and learning. *ELife*,

10, 1–29. https://doi.org/10.7554/ELIFE.62825

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Yacoub, E., Ugurbil, K., Andersson, J.,

Beckmann, C. F., Jenkinson, M., Smith, S. M., & Essen, D. C. Van. (2017). Manuscripts A multi-

modal parcellation of human cerebral cortex. *Nature*, *536*(7615), 171–178.
https://doi.org/10.1038/nature18933.A

Glazer, J. E., Kelley, N. J., Pornpattananangkul, N., Mittal, V. A., & Nusslock, R. (2018). Beyond the
FRN: Broadening the time-course of EEG and ERP components implicated in reward
processing. *International Journal of Psychophysiology*, *132*, 184–202.
https://doi.org/10.1016/j.ijpsycho.2018.02.002

Goldstein, R. Z., Cottone, L. A., Jia, Z., Maloney, T., Volkow, N. D., & Squires, N. K. (2006). The effect
of graded monetary reward on cognitive event-related potentials and behavior in young
healthy adults. *International Journal of Psychophysiology*, *62*(2), 272–279.
https://doi.org/10.1016/j.ijpsycho.2006.05.006

Gross, J., Kujala, J., Hämäläinen, M., Timmermann, L., Schnitzler, A., & Salmelin, R. (2001). Dynamic
imaging of coherent sources: Studying neural interactions in the human brain. *Proceedings of
the National Academy of Sciences of the United States of America*, *98*(2), 694–699.
https://doi.org/10.1073/pnas.98.2.694

Hall-McMaster, S., Muhle-Karbe, P. S., Myers, N. E., & Stokes, M. G. (2019). Reward boosts neural
coding of task rules to optimize cognitive flexibility. *Journal of Neuroscience*, *39*(43), 8549–
8561. https://doi.org/10.1523/JNEUROSCI.0631-19.2019

Hämäläinen, M. (1995). Functional localization based on measurements with a whole-head
magnetometer system. *Brain Topography*, *7*(4), 283–289. https://doi.org/10.1007/BF01195254

Hansen, J. C., & Hillyard, S. A. (1980). Endogeneous brain potentials associated with selective
auditory attention. *Electroencephalography and Clinical Neurophysiology*, *49*, 277–290.
https://doi.org/10.1016/0013-4694(80)90222-9

Henry, M. J., Herrmann, B., Kunke, D., & Obleser, J. (2017). Aging affects the balance of neural
entrainment and top-down neural modulation in the listening brain. *Nature Communications*,
8. https://doi.org/10.1038/ncomms15801

Henry, M. J., Herrmann, B., & Obleser, J. (2014). Entrained neural oscillations in multiple frequency
bands comodulate behavior. *Proceedings of the National Academy of Sciences of the United
States of America*, *111*(41), 14935–14940. https://doi.org/10.1073/pnas.1408741111

Henry, M. J., & Obleser, J. (2012). Frequency modulation entrains slow neural oscillations and
optimizes human listening behavior. *Proceedings of the National Academy of Sciences of the
United States of America*, *109*(49), 20095–20100. https://doi.org/10.1073/pnas.1213390109

Herrmann, B., & Johnsrude, I. S. (2020). A model of listening engagement (MoLE). *Hearing Research*,
397. https://doi.org/10.1016/j.heares.2020.108016

Herrmann, B., Maess, B., Henry, M. J., Obleser, J., & Johnsrude, I. S. (2023). Neural signatures of task-
related fluctuations in auditory attention and age-related changes. *NeuroImage*, *268*, 119883.
https://doi.org/10.1016/j.neuroimage.2023.119883

Herrmann, B., & Ryan, J. D. (2024). Pupil Size and Eye Movements Differently Index Effort in Both
Younger and Older Adults. *Journal of Cognitive Neuroscience*, 1–16.
https://doi.org/10.1162/jocn_a_02172

Hillyard, S. A., Hink, R. F., Schwent, V. L., & Picton, T. W. (1973). Electrical Signs of Selective Attention
in the Human Brain. *Science*, *182*, 177–180. https://doi.org/10.1126/science.182.4108.177

Holroyd, C. B., & Coles, M. G. H. (2002). The neural basis of human error processing: Reinforcement
learning, dopamine, and the error-related negativity. *Psychological Review*, *109*(4), 679–709.

https://doi.org/10.1037/0033-295X.109.4.679

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity:

Gating by inhibition. *Frontiers in Human Neuroscience*, *4*, 1–8.

https://doi.org/10.3389/fnhum.2010.00186

Joshi, S., & Gold, J. I. (2020). Pupil Size as a Window on Neural Substrates of Cognition. *Trends in*

Cognitive Sciences, *24*(6), 466–480. https://doi.org/10.1016/j.tics.2020.03.005

Joshi, S., Li, Y., Kalwani, R. M., & Gold, J. I. (2016). Relationships between Pupil Diameter and

Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron*, *89*(1), 221–

234. https://doi.org/10.1016/j.neuron.2015.11.028

Kadem, M., Herrmann, B., Rodd, J. M., & Johnsrude, I. S. (2020). Pupil Dilation Is Sensitive to

Semantic Ambiguity and Acoustic Degradation. *Trends in Hearing*, *24*.

https://doi.org/10.1177/2331216520964068

Kahneman, D., & Beatty, J. (1966). Pupil Diameter and Load on Memory. *Science*, 1583–1586.

Keitel, A., & Gross, J. (2016). Individual Human Brain Areas Can Be Identified from Their

Characteristic Spectral Activation Fingerprints. *PLoS Biology*, *14*(6), 1–22.

https://doi.org/10.1371/journal.pbio.1002498

Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-timing

hypothesis. *Brain Research Reviews*, *53*(1), 63–88.

https://doi.org/10.1016/j.brainresrev.2006.06.003

Koelewijn, T., Zekveld, A. A., Festen, J. M., & Kramer, S. E. (2012). Pupil dilation uncovers extra

listening effort in the presence of a single-talker masker. *Ear and Hearing*, *33*(2), 291–300.

https://doi.org/10.1097/AUD.0b013e3182310019

Koelewijn, T., Zekveld, A. A., Lunner, T., & Kramer, S. E. (2018). The effect of reward on listening

effort as reflected by the pupil dilation response. *Hearing Research*, *367*, 106–112.

https://doi.org/10.1016/j.heares.2018.07.011

Kraus, F., Obleser, J., & Herrmann, B. (2023a). Pupil Size Sensitivity to Listening Demand Depends

on Motivational State. *ENeuro*, *10*(12), 1–12. https://doi.org/10.1523/ENEURO.0288-23.2023

Kraus, F., Tune, S., Obleser, J., & Herrmann, B. (2023b). Neural α Oscillations and Pupil Size

Differentially Index Cognitive Demand under Competing Audiovisual Task Conditions. *The*

Journal of Neuroscience, *43*(23), 4352 LP – 4364. https://doi.org/10.1523/JNEUROSCI.2181-

22.2023

Kraus, F., Tune, S., Ruhe, A., Obleser, J., & Wöstmann, M. (2021). Unilateral Acoustic Degradation

Delays Attentional Separation of Competing Speech. *Trends in Hearing*, *25*.

https://doi.org/10.1177/23312165211013242

Krebs, R. M., Boehler, C. N., Appelbaum, L. G., & Woldorff, M. G. (2013). Reward Associations

Reduce Behavioral Interference by Changing the Temporal Dynamics of Conflict Processing.

PLoS ONE, *8*(1). https://doi.org/10.1371/journal.pone.0053894

Liu, Y., Rodenkirch, C., Moskowitz, N., Schriver, B., & Wang, Q. (2017). Dynamic Lateralization of

Pupil Dilation Evoked by Locus Coeruleus Activation Results from Sympathetic, Not

Parasympathetic, Contributions. *Cell Reports*, *20*(13), 3099–3112.

https://doi.org/10.1016/j.celrep.2017.08.094

Mathôt, S., Fabius, J., Van Heusden, E., & Van der Stigchel, S. (2018). Safe and sensible

preprocessing and baseline correction of pupil-size data. *Behavior Research Methods*, *50*(1),

94–106. https://doi.org/10.3758/s13428-017-1007-2

Mazaheri, A., van Schouwenburg, M. R., Dimitrijevic, A., Denys, D., Cools, R., & Jensen, O. (2014).

Region-specific modulations in oscillatory alpha activity serve to facilitate processing in the

visual and auditory modalities. *NeuroImage*, *87*, 356–362.

https://doi.org/10.1016/j.neuroimage.2013.10.052

McCormick, D. A. (1998). Cholinergic and noradrenergic modulation of thalamocortical processing.

Trends in Neurosciences, *12*, 215–221.

McMahon, C. M., Boisvert, I., de Lissa, P., Granger, L., Ibrahim, R., Lo, C. Y., Miles, K., & Graham, P. L.

(2016). Monitoring alpha oscillations and pupil dilation across a performance-intensity

function. *Frontiers in Psychology*, *7*, 1–12. https://doi.org/10.3389/fpsyg.2016.00745

Miles, K., McMahon, C., Boisvert, I., Ibrahim, R., de Lissa, P., Graham, P., & Lyxell, B. (2017). Objective

Assessment of Listening Effort: Coregistration of Pupillometry and EEG. *Trends in Hearing*, *21*,

1–13. https://doi.org/10.1177/2331216517706396

Näätänen, R., & Picton, T. (1987). The N1 Wave of the Human Electric and Magnetic Response to

Sound: A Review and an Analysis of the Component Struture. *Psychophysiology*, *24 (4)*.

Niemi, P., & Näätänen, R. (1981). Foreperiod and simple reaction time. *Psychological Bulletin*, *89*(1),

133–162. https://doi.org/10.1037/0033-2909.89.1.133

Nobre, A., Correa, A., & Coull, J. (2007). The hazards of time. *Current Opinion in Neurobiology*, *17*(4),

465–470. https://doi.org/10.1016/j.conb.2007.07.006

O'Sullivan, J. A., Power, A. J., Mesgarani, N., Rajaram, S., Foxe, J. J., Shinn-Cunningham, B. G., Slaney,

M., Shamma, S. A., & Lalor, E. C. (2015). Attentional Selection in a Cocktail Party Environment

Can Be Decoded from Single-Trial EEG. *Cerebral Cortex*, *25*(7), 1697–1706.

https://doi.org/10.1093/cercor/bht355

Obleser, J., Wöstmann, M., Hellbernd, N., Wilsch, A., & Maess, B. (2012). Adverse listening

conditions and memory load drive a common alpha oscillatory network. *Journal of

Neuroscience*, *32*(36), 12376–12383. https://doi.org/10.1523/JNEUROSCI.4908-11.2012

Ohlenforst, B., Wendt, D., Kramer, S. E., Naylor, G., Zekveld, A. A., & Lunner, T. (2018). Impact of

SNR, masker type and noise reduction processing on sentence recognition performance and

listening effort as indicated by the pupil dilation response. *Hearing Research*, *365*, 90–99.

https://doi.org/10.1016/j.heares.2018.05.003

Ohlenforst, B., Zekveld, A. A., Lunner, T., Wendt, D., Naylor, G., Wang, Y., Versfeld, N. J., & Kramer, S.

E. (2017). Impact of stimulus-related factors and hearing impairment on listening effort as

indicated by pupil dilation. *Hearing Research*, *351*, 68–79.

https://doi.org/10.1016/j.heares.2017.05.012

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for

advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational

Intelligence and Neuroscience*, *2011*. https://doi.org/10.1155/2011/156869

Orf, M., Wöstmann, M., Hannemann, R., & Obleser, J. (2023). Target enhancement but not distractor

suppression in auditory neural tracking during continuous speech. *IScience*, *26*(6).

https://doi.org/10.1016/j.isci.2023.106849

Paiva, T. O., Almeida, P. R., Ferreira-Santos, F., Vieira, J. B., Silveira, C., Chaves, P. L., Barbosa, F., &

Marques-Teixeira, J. (2016). Similar sound intensity dependence of the N1 and P2 components

of the auditory ERP: Averaged and single trial evidence. *Clinical Neurophysiology*, *127*(1), 499–

508. https://doi.org/10.1016/j.clinph.2015.06.016

Palva, S., & Palva, J. M. (2011). Functional roles of alpha-band phase synchronization in local and
large-scale cortical networks. *Frontiers in Psychology*, *2*.
https://doi.org/10.3389/fpsyg.2011.00204

Parro, C., Dixon, M. L., & Christoff, K. (2018). The neural basis of motivational influences on
cognitive control. *Human Brain Mapping*, *39*(12), 5097–5111.
https://doi.org/10.1002/hbm.24348

Paul, B. T., Chen, J., Le, T., Lin, V., & Dimitrijevic, A. (2021). Cortical alpha oscillations in cochlear
implant users reflect subjective listening effort during speech-in-noise perception. *PLoS ONE*,
16, 1–22. https://doi.org/10.1371/journal.pone.0254162

Petersen, E. B., Wöstmann, M., Obleser, J., Stenfelt, S., & Lunner, T. (2015). Hearing loss impacts
neural alpha oscillations under adverse listening conditions. *Frontiers in Psychology*, *6*, 1–11.
https://doi.org/10.3389/fpsyg.2015.00177

Pfurtscheller, G. (2003). Induced Oscillations in the Alpha Band: Functional Meaning. *Epilepsia*, *44*,
2–8. https://doi.org/10.1111/j.0013-9580.2003.12001.x

Pichora-Fuller, M. K., Kramer, S. E., Eckert, M. A., Edwards, B., Hornsby, B. W. Y., Humes, L. E., Lemke,
U., Lunner, T., Matthen, M., Mackersie, C. L., Naylor, G., Phillips, N. A., Richter, M., Rudner, M.,
Sommers, M. S., Tremblay, K. L., & Wingfield, A. (2016). Hearing impairment and cognitive
energy: The framework for understanding effortful listening (FUEL). *Ear and Hearing*, *37*, 5S-
27S. https://doi.org/10.1097/AUD.0000000000000312

Richter, M. (2016). The moderating effect of success importance on the relationship between
listening demand and listening effort. *Ear and Hearing*, *37*, 111S-117S.
https://doi.org/10.1097/AUD.0000000000000295

Richter, M., Gendolla, G. H. E., & Wright, R. A. (2016). Three Decades of Research on Motivational
Intensity Theory. In A. J. Elliot (Ed.), *Advances in Motivation Science* (pp. 149–186). Elsevier.
https://doi.org/10.1016/bs.adms.2016.02.001

Rushworth, M. F. S., Paus, T., & Sipila, P. K. (2001). Attention systems and the organization of the
human parietal cortex. *Journal of Neuroscience*, *21*(14), 5262–5271.
https://doi.org/10.1523/jneurosci.21-14-05262.2001

Sambrook, T. D., & Goslin, J. (2015). A Neural reward prediction error revealed by a meta-analysis
of ERPs using great grand averages. *Psychological Bulletin*, *141*(1), 213–235.
https://doi.org/10.1037/bul0000006

Sawaki, R., Luck, S. J., & Raymond, J. E. (2015). How Attention Changes in Response to Incentives.
Journal of Cognitive Neuroscience, *27*(11), 2229–2239. https://doi.org/10.1162/jocn_a_00847

Small, D. M., Gitelman, D., Simmons, K., Bloise, S. M., Parrish, T., & Mesulam, M. M. (2005). Monetary
incentives enhance processing in brain regions mediating top-down control of attention.
Cerebral Cortex, *15*(12), 1855–1865. https://doi.org/10.1093/cercor/bhi063

Steriade, M. (2000). Corticothalamic resonance, states of vigilance and mentation. *Neuroscience*,
101(2), 243–276. https://doi.org/10.1016/S0306-4522(00)00353-5

Teoh, Y. Y., Yao, Z., Cunningham, W. A., & Hutcherson, C. A. (2020). Attentional priorities drive
effects of time pressure on altruistic choice. *Nature Communications*, *11*(1).
https://doi.org/10.1038/s41467-020-17326-x

Tune, S., Alavash, M., Fiedler, L., & Obleser, J. (2021). Neural attentional-filter mechanisms of

listening success in middle-aged and older individuals. *Nature Communications*, *12*(1), 1–14.
https://doi.org/10.1038/s41467-021-24771-9

Tusche, A., & Hutcherson, C. A. (2018). Cognitive regulation alters social and dietary choice by
changing attribute representations in domain-general and domain-specific brain circuits. *ELife*,
7, 1–35. https://doi.org/10.7554/eLife.31185

van den Berg, B., Krebs, R. M., Lorist, M. M., & Woldorff, M. G. (2014). Utilization of reward-prospect
enhances preparatory attention and reduces stimulus conflict. *Cognitive, Affective and*
Behavioral Neuroscience, *14*(2), 561–577. https://doi.org/10.3758/s13415-014-0281-z

Vazey, E. M., Moorman, D. E., & Aston-Jones, G. (2018). Phasic locus coeruleus activity regulates
cortical encoding of salience information. *Proceedings of the National Academy of Sciences of*
the United States of America, *115*(40), E9439–E9448. https://doi.org/10.1073/pnas.1803716115

Vrba, J., & Robinson, S. E. (2001). Signal processing in magnetoencephalography. *Methods*, *25*(2),
249–271. https://doi.org/10.1006/meth.2001.1238

Wagenmakers, E. J. (2007). A practical solution to the pervasive problems of p values. *Psychonomic*
Bulletin and Review, *14*(5), 779–804. https://doi.org/10.3758/BF03194105

Weisz, N., Hartmann, T., Müller, N., Lorenz, I., & Obleser, J. (2011). Alpha rhythms in audition:
Cognitive and clinical perspectives. *Frontiers in Psychology*, *2*, 1–15.
https://doi.org/10.3389/fpsyg.2011.00073

Wendt, D., Dau, T., & Hjortkjær, J. (2016). Impact of background noise and sentence complexity on
processing demands during sentence comprehension. *Frontiers in Psychology*, *7*, 1–12.
https://doi.org/10.3389/fpsyg.2016.00345

Winn, M. B., Edwards, J. R., & Litovsky, R. Y. (2015). The Impact of Auditory Spectral Resolution on
Listening Effort Revealed by Pupil Dilation. *Ear and Hearing*, *36*(4), 1–31.
https://doi.org/10.1097/AUD.0000000000000145.The

Wisniewski, M. G., Thompson, E. R., & Iyer, N. (2017). Theta- and alpha-power enhancements in the
electroencephalogram as an auditory delayed match-to-sample task becomes impossibly
difficult. *Psychophysiology*, *54*(12), 1916–1928. https://doi.org/10.1111/psyp.12968

Woldorff, M. G., & Hillyard, S. A. (1991). Modulation of early auditory processing during selective
listening to rapidly presented tones. *Electroencephalography and Clinical Neurophysiology*, *79*,
170–191.

Wöstmann, M., Herrmann, B., Wilsch, A., & Obleser, J. (2015). Neural alpha dynamics in younger
and older listeners reflect acoustic challenges and predictive benefits. *Journal of Neuroscience*,
35(4), 1458–1467. https://doi.org/10.1523/JNEUROSCI.3250-14.2015

Wöstmann, M., Lim, S. J., & Obleser, J. (2017). The Human Neural Alpha Response to Speech is a
Proxy of Attentional Control. *Cerebral Cortex*, *27*(6), 3307–3317.
https://doi.org/10.1093/cercor/bhx074

Yee, D. M., & Braver, T. S. (2018). Interactions of motivation and cognitive control. *Current Opinion*
in Behavioral Sciences, *19*, 83–90. https://doi.org/10.1016/j.cobeha.2017.11.009

Zekveld, A. A., & Kramer, S. E. (2014). Cognitive processing load across a wide range of listening
conditions: Insights from pupillometry. *Psychophysiology*, *51*(3), 277–284.
https://doi.org/10.1111/psyp.12151

Zekveld, A. A., Kramer, S. E., & Festen, J. M. (2010). Pupil response as an indication of effortful
listening: The influence of sentence intelligibility. *Ear and Hearing*, *31*(4), 480–490.

https://doi.org/10.1097/AUD.0b013e3181d4f251

Zhang, M., Siegle, G. J., McNeil, M. R., Pratt, S. R., & Palmer, C. (2019). The role of reward and task

demand in value-based strategic allocation of auditory comprehension effort. *Hearing*

Research, *381*. https://doi.org/10.1016/j.heares.2019.107775

954