Assessment of Plant- and Microbial-Derived Protein Hydrolysates as Sustainable for Fetal Bovine Serum in Seafood Cell Culture Media

Arian Amirvaresi[a], Reza Ovissipour[a]

[a]Department of Food Science and Technology, Texas A&M University, College Station, TX 77843, USA

*Corresponding author: Reza Ovissipour, Email: ovissi@tamu.edu

Abstract

This study seeks to explore alternatives by substituting or reducing the conventional 10% serum concentration in Zebrafish embryonic stem cell (ESC) growth media with protein hydrolysates sourced from peas, mushrooms, yeast, and algae. Notably, algae exhibited the highest protein content, optimal amino acid balance, and favorable functional properties. When applied at concentrations ranging from 1 to 10 mg/mL, all protein hydrolysates demonstrated pro-apoptotic effects and inhibited cell growth, particularly when used in conjunction with 10% serum. However, concentrations ranging from 0.001 to 0.1 mg/mL displayed anti-apoptotic properties and promoted cell proliferation. The study found that media containing 1% or 2.5% serum, along with 0.01 mg/mL of protein hydrolysates, supported cell growth effectively. Lactate Dehydrogenase (LDH) Activity served as an indicator of cell health and integrity under the specified conditions of protein hydrolysate supplementation. Cells cultured in serum-free media exhibited significantly decreased cell membrane integrity ($P < 0.05$) compared to those in regular media or media containing low serum (1% and 2.5%) along with low concentrations (0.01 mg/mL) of protein hydrolysates. Furthermore, analysis of Greenhouse Gas emissions (GHG) suggested that media formulations containing 1% serum combined with low concentrations of protein hydrolysates present a sustainable approach for cell-based seafood production.

Keywords: Cultivated meat, cell-based seafood, serum-free media, protein hydrolysate, peptides, greenhouse gases.

1. Introduction

By 2050, the world population is projected to reach 10 billion, posing a significant challenge in meeting the global demand for food. According to the Food and Agriculture Organization (FAO), a 70% increase in meat production is required to address this growing need (FAO, 2020). Among the various animal-based meats, aquatic food products play a crucial role as an essential commodity, constituting approximately 20% of the animal protein consumed globally.

Cultivated meat involves the cultivation of cells or tissues in a carefully formulated medium that supports cell proliferation, metabolism, and differentiation. This innovative technique provides a promising solution to the challenges associated with traditional meat production. Cultivated meat reduces reliance on fisheries and aquaculture, mitigating environmental impacts, addressing animal welfare concerns, and reducing public health risks. It offers a sustainable and ethical solution to meeting global protein demand while overcoming challenges in traditional aquatic food production.

A culture medium plays a vital role in supporting the proliferation and growth of cells or tissues in the process of cultivated meat production (Yao and Asayama, 2017). The cost of cell culture media constitutes a significant portion of the total expenses involved in cultivated meat production, accounting for more than 99% of the total cost (Stout et al., 2022). One potential approach to reduce the overall cost of cultivated meat is to develop less expensive culture media. Therefore, finding alternative, cost-effective sources or formulations for these components could have a substantial impact on reducing the cost of cultivated meat production.

Fetal bovine serum (FBS) plays a crucial role in cell culture as it provides essential nutrients, growth factors, and hormones necessary for the proliferation and viability of cells. It has long been the gold standard in cell culture media due to its ability to support robust cell growth and differentiation (Stout et al., 2022). However, using FBS arouses several challenges due to its high cost presenting a significant barrier, impacting the scalability and commercial viability of cultivated meat production. Additionally, concerns about the ethical sourcing of FBS and the potential risk of disease transmission have prompted researchers to explore alternative, sustainable, and cost-effective options (Freshney, 2015). Developing suitable substitutes for FBS is essential to overcome these challenges and advance the field of cultivated meat towards a more scalable, affordable, and ethically sustainable future.

Protein hydrolysates, derived from protein cleavage, can be used as supplements in cell culture media to provide essential nutrients and growth factors for cells. Protein hydrolysates derived from various plant and animal sources have been utilized by researchers for cell growth and cultivation purposes (Batish et al., 2022). Peptides, derived from protein hydrolysates or synthesized chemically, play a role in cell signaling and can be incorporated into media to modulate cellular processes. The application of these peptides has gained significant attention due to their ability to provide essential nutrients and growth factors required for cell proliferation and maintenance. Researchers have explored the use of soy-derived peptides, among others, to support the growth of different cell types *in vitro* (Andreassen et al., 2022). These peptides offer a rich source of amino acids, which serve as building blocks for protein synthesis and cellular functions (Burteau et al., 2003; Spearman et al., 2014; Batish et al., 2022; Timoneda et al., 2024). Serum-free media can benefit from peptides in terms of cell growth, cell performance, differentiation and sustainability (Kim et al., 2011; Logarušić et al., 2021; Nikkhah et al., 2023; van der Valk et al., 2010).While, incorporating protein hydrolysates into the cell culture media, enhanced cell growth, proliferation and viability in many researches, the negative and pro-apoptosis properties of protein hydrolysates have been reported when applied at high concentrations between 1 to 10 mg/mL (Lee et al., 2008; Batish et al., 2022; Hsieh et al., 2022). In addition, sustainability and greenhouse gases (GHG) emissions have not been used to justify the application of protein hydrolysates in cell culture media. Thus, this research aims to evaluate the impact of different concentrations of serum, alone and in conjunction with different concentrations of various protein hydrolysates on fish cell line growth, morphology, viability, and performance. GHG emissions also is provided for different conditions.

2. Materials and methods

2.1. Materials and chemicals

The algae and pea protein were provided from Mountain Rose Herbs (Eugene, Oregon, USA) and NorCals organic (Crescent City, CA, USA), respectively. The white mushrooms (*Agaricus bisporus*) were obtained from the local market as fresh. Yeast hydrolysate, Bacto™ TC Yeastolate, serum, Lebovitz 15 media, HEPES and sodium bicarbonate were acquired from Thermo-Fisher scientific (Waltham, MA, USA). Alcalase®, a commercially available

endoprotease enzyme (2.4 AU/g) derived from *Bacillus licheniformis* was obtained from Sigma-Aldrich Inc. (St. Louis, MO, USA). The Zebrafish Embryonic Stem Cells (ZESCs), ZEM2S CRL-2147™, were obtained from the American Type Culture Collection (ATCC). The antibiotics which were initially used in cell culture were obtained from Cytiva (Marlborough, MA, USA).

2.2. Protein hydrolysates production

The hydrolysates were prepared in at least six replicates, based on the optimum conditions (Table 1). The enzymatic hydrolysis of different substrates was conducted with minor adjustments tailored to each specific substrate, as described before (Batish et al., 2022, 2020). Briefly, the raw materials were homogeneously mixed with water at a predetermined ratio as specified in Table 1. Subsequently, the mixture was subjected to enzymatic hydrolysis at 60°C for 4 h. To stop the enzymatic reaction, the samples were subjected to 90°C for 10 min. The samples were centrifuged to collect the protein hydrolysates, which were then freeze-dried. The dry yield (%) was determined using equations 1. The lyophilized protein hydrolysates were stored at -20°C until further use.

$$\text{Yield (\%)} = \frac{\text{Weight of dried protein hydrolysate (g)}}{\text{Weight of raw material used (g)}} \times 100 \qquad \text{eq.1}$$

2.3. Amino acid analysis, protein content and degree of hydrolysis

The total amino acid analysis was performed following the guidelines specified in the Association of Official Agricultural Chemists (AOAC) 982.30 E (a,b,c) (AOAC, 2006). Upon complete sample digestion with 6N HCl, an ion exchange chromatography method with post-column ninhydrin derivatization and quantitation was employed. The determination of crude protein content was carried out using the AOAC standard method, Kjeldahl. To calculate the protein content, the crude nitrogen content was multiplied by the appropriate nitrogen to protein conversion factor (*Kp*) specific to each sample type. For algae, a *Kp* value of 6.35 (Safi et al., 2013), for pea, a *Kp* value of 5.36 (Mariotti et al., 2008), and for mushroom, a *Kp* value of 4.7 were applied (Mattila et al., 2002).

The protein quality assessment was conducted using Digestible Indispensable Amino Acid Score (DIAAS) as recommended by FAO (FAO, 2011) using equation 2:

$$\text{DIAAS (\%)} = 100 \times \text{lowest value} \left(\frac{\text{mg of DIAA in 1g of protein}}{\text{mg of DIAA in 1g of reference protein}} \right) \qquad \text{eq.2}$$

Degree of hydrolysis was measured using formal titration method (Taylor, 1957).

2.4. Oil holding capacity (OHC)

The evaluation of the oil holding capacity (OHC) was carried out for each protein

hydrolysate (Shahidi et al., 1995). Briefly, 500 mg of protein hydrolysate was combined with 10

mL of pure canola oil. The mixture was then left at room temperature for 30 min, gentle mixing

every 10 min. Subsequently, centrifugation was performed at 2500*g* for 10 min at room

temperature. The OHC was determined by utilizing equation 3.

164
$$\text{Oil holding capacity} = \frac{\text{Mass of sample (with oil)} - \text{Mass of dry sample(g)}}{\text{Mass of dry sample(g)}} \qquad \text{eq. 3}$$
165

166 *2.5. Emulsifying capacity (EC)*

167   The evaluation of the EC involved the mixing of 500 mg of protein hydrolysates with 50 mL

168 of a 0.1 M sodium chloride solution in a 250 mL conical flask, conducted at ambient temperature

169 (Yasumatsu et al., 1972). The EC was determined by employing equation 4.

170
$$\text{Emulsyfying capacity} \left(\text{EC(\%)} \right) = \frac{\text{Total volume} - \text{Aqueous volume}}{\text{Total volume}} \times 100 \qquad \text{eq. 4}$$
171

172 *2.6. Foaming capacity (FC)*

173   The foaming capacity (FC) was determined based on the methodology described by

174 Pacheco-Aguilar et al. (2008). In this procedure, 750 mg of protein hydrolysates were mixed with

175 25 mL of distilled water for a duration of 10 minutes. Subsequently, the mixture was subjected to

176 homogenization for two minutes. The FC was then calculated utilizing equation 5.

177
$$\text{Foaming capacity} \left(\text{FC(\%)} \right) = \frac{\text{Aerated volume} - \text{Normal volume}}{\text{Normal volume}} \times 100 \qquad \text{eq. 5}$$
178

2.7. Cell culture and maintenance

For the initial cell culture, a combination of Lebovitz-L-15 media (L-15), Dulbecco's

Modified Eagle Media (DMEM), and Ham's F12 Media (F-12 media) was utilized in a ratio of

15:50:35, respectively. The culture media were supplemented with buffering agents, including 20

mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) and 0.18 g/L sodium

bicarbonate, and supplemented with 10% FBS. The cells in ampoules were thawed at a temperature

of 28°C and subsequently resuspended in 9 mL of media containing 5% fetal FBS within a T-25

cm^2 flask. The cells were allowed to settle for 30 min, following which additional FBS was added

to achieve a final concentration of 10% FBS. The subculturing procedure was conducted when the

cells reached 80-85% confluence. This was accomplished by rinsing the cells with phosphate-

buffered saline (PBS) and subsequently treating them with Tryple Express, a cell detachment

reagent, to detach the cells from the surface of the flask. The flask containing the detached cells

was maintained at a temperature of 28°C for a duration of 5 minutes to ensure complete

detachment. Subsequently, the cells were transferred to a 15 mL Falcon tube. To neutralize the

Tryple reagent, serum-free media (L-15 media) was used, and the cell suspension was then

subjected to centrifugation at 130g for 8.5 minutes. The supernatant was carefully discarded, and

the resulting cell pellet was resuspended in media containing 10% serum. The cell number and

viability were assessed using an automatic cell counter, and cell splitting was performed based on

the obtained results. The cells reached confluency within one week, with a media change

performed once on the third day.

2.8. Cell performance in reduced or serum free media

To evaluate the potential capability of protein hydrolysates in reducing serum usage, various

protein hydrolysates were applied at different concentrations in two different experiments. First in

Experiment I, the impact of different concentrations of serum (0, 1, 2.5, 5, and 10%) was evaluated

on the cell performance. In Experiment II, various concentrations of protein hydrolysates (0.001,

0.01, 0.1, 1, and 10 mg/mL) were studied in combination with media containing 0, 5, and 10%

serum. In experiment III, lower concentrations of protein hydrolysates (0.001, 0.01, and 0.1

mg/mL) in combination with media containing lower concentrations of serum (0, 1, 2.5, and 10%)

were studied. The protein hydrolysate samples were prepared by combining them with sterile water

that had been filtered through a 0.22 µm filter. The cells were initially seeded at a density of 50,000

cells/mL and incubated for 24 hours at 28°C in a medium containing 10% serum to facilitate cell

attachment to the culture plates.

Cell numbers and morphology were documented at 24-hour intervals for a total of three days

using the CKX-CCSW Confluence Checker, which captured three images per well. Each

hydrolysate condition was tested in four biological replicates, with each replicate having three

technical replicates. The growth rate was determined using equation 6, while the doubling time

was calculated using equation 7.

$$\text{Growth rate (hr}^{-1}) = \frac{\text{Ln} \frac{\text{(viable cells at passaging)}}{\text{viable cells at seeding}}}{\text{Time (hrs)}}$$ eq. 6

$$\text{Doubling time (hr)} = \frac{\text{Ln (2)}}{\text{grwoth rate (hr}^{-1})}$$ eq. 7

Cell viability was assessed utilizing PrestoBlue analysis. After 72 hours of cell seeding, 100 µL of

PrestoBlue reagent was added to the wells, and the plate was incubated at 28°C for 2 hours.

Subsequently, the absorbances at 570 nm and 600 nm were measured using a microplate reader.

The reduction of the dye was determined by applying equation 8 for calculation.

223
$$\text{Reagent reduction (\%)} = \frac{(117216 \times A1) - (80586 \times A2)}{(155677 \times N2) - (14652 \times N1)} \times 100$$ eq. 8
224

In the equation, A1 represents the absorbance of the sample wells at 570 nm, while A2 corresponds

to the absorbance of the sample wells at 600 nm. Similarly, N1 denotes the absorbance of the

media-only wells at 570 nm, and N2 represents the absorbance of the media-only wells at 600 nm.

2.9. Fluorescent imaging

Based on the results obtained from cell growth assays, optimal concentrations of protein

hydrolysates were selected for further image analysis. To prepare the cells for imaging, they were

first fixed with 4% paraformaldehyde for 10 min, followed by two rinses with PBS to remove the

paraformaldehyde. To facilitate permeabilization, the cells were then incubated in 0.1% Tween

solution for 10 min, followed by another round of rinsing with PBS. For visualizing the cell nuclei,

Hoechst dye was diluted in PBS at a ratio of 1:2000, added to the cells, and allowed to incubate in

the dark for 10 minutes. After incubation, the cells were washed twice with PBS, and 1 mL of PBS

was added to each well. Subsequently, a second dye for visualizing the cytoskeleton, actin green

dye, was added to each well, and the cells were incubated for 30 minutes before being washed

twice with PBS. To optimize the image quality, the cells were then immersed in a live cell imaging

solution. Finally, the cells were observed using a fluorescent microscope, with UV

excitation/emission wavelengths of 361/486 nm for Hoechst dye and blue-cyan light with

excitation/emission wavelengths of 495/518 nm for actin green dye.

2.10.   Lactate dehydrogenase activity

The activity of lactate dehydrogenase (LDH) was assessed in different concentrations of

selected protein hydrolysates supplemented with either 1% or 2.5% serum, which were found to

be optimal for supporting cell growth in experiment III. The cells were cultured under these

optimal conditions for a duration of three days. Following the three-day cultivation period, the

supernatant was collected and transferred to 96-well plates for the LDH activity assay. The assay

was performed according to the manufacturer's instructions. To ensure statistical reproducibility,

four biological replicates were included, with each biological replicate having three technical

replicates.

2.11.   Greenhouse gas emissions (GHG)

Greenhouse gas emissions associated with serum, intact proteins, and enzymatically

hydrolyzed proteins were sourced from various inventories. The functional unit (FU) for this study

is based on 1 kg of the products. Algae, mushroom, and pea were acquired as dried materials and

were subjected to enzymatic hydrolysis in our laboratory. However, for yeast, protein hydrolysates

were procured directly from the vendor. Thus, GHG emission data for yeast peptone (protein

hydrolysates) was obtained from inventory. For algae, GHG emissions ranged from 1.5 to 150 kg

CO_2 eq./kg, contingent upon the production method and carbon sequestration data. Production

method were obtained from the vendor website, and GHG emissions were selected based on a

review of multiple articles (Clarens et al., 2010; Handler et al., 2012; Nette et al., 2016; "The

Climate Intelligence Platform - CarbonCloud).

2.12.   Statistical analysis

The statistical analysis was carried out using JMP 16 software. To ensure the validity of the

analysis, the data was assessed for normality and homoscedasticity, confirming a normal

distribution. Normality was tested using the Shapiro-Wilk test and Normal Quantile Plots (NQP),

while the homogeneity of variability was assessed using Levene's test. If the data exhibited a

normal distribution, One-Way ANOVA and Tukey's HSD (Honestly Significant Difference) tests

were performed. However, if the data did not meet the criteria for normality, the

Wilcoxon/Kruskal-Wallis test and Steel-Dwass/Wilcoxon pair comparisons were employed, along

with their respective control conditions. Statistical significance was determined by considering

differences with *P < 0.05*.

3. Results and discussion

3.1. Protein hydrolysates properties

For producing hydrolysates that properly support animal cell culture growth, the whole protein-rich substrate was used instead of protein isolate or concentrate. The yield of protein hydrolysates were examined to determine their potential for large-scale manufacturing (Table 1). As shown later in this investigation, the concentration of protein hydrolysate required for animal cellular culture applications is between 0.001 mg/mL to 0.1 mg/mL, and the yield of hydrolysates reported in this research appear to be suitable for cultivated meat scaling up. Producing cell-based meat requires the cultivation of billions of cells using limited space, time, and resources. One 2 L batch of any protein hydrolysate would be sufficient to supply the maximum capacity of a stirred bioreactor used for animal cell culture which has a capacity of 2000 L to produce 10 to 100 kg cell-based meat (Bellani et al., 2020).

Table 1. Specific reaction conditions for enzymatic hydrolysis of substrates and the dry yield

Substrate	Water to substrate ratio (v/v)	Enzyme (%)	Hydrolysis time (h)	Dry yield (%)
Algae	10:1	2%	4	23.8 ± 1.48
Pea	20:1	4%	4	39.58 ± 1.74
Mushroom	10:1	2.5%	4	5.78 ± 0.35

3.2. Amino acid analysis, protein content and protein quality

The amino acid composition, protein content, DIAAS score and protein quality are presented in Table 2. All protein hydrolysates contained a high concentration of amino acids with cell-growth promoting properties, such as alanine, glycine, proline, and aspartic acid (Lu and Zhao, 2018). All protein hydrolysates examined in this study demonstrated a notable abundance of glutamic acid, a crucial component in animal cell culture. Glutamic acid serves as a significant nitrogen source for in vivo processes and contributes to the synthesis of amino acids, proteins, and nucleic acids. Furthermore, glutamic acid acts as a substantial supplier of metabolites in the tricarboxylic acid (TCA) cycle, supplying carbon and nitrogen for biosynthesis (Hosios et al., 2016). The hydrolysates were also found containing significant amounts of hydrophobic amino acids, including glycine, alanine, valine, and leucine. Furthermore, both yeast and algae protein

hydrolysates exhibited a high protein quality, as evidenced by their DIAAS content of essential amino acids.

Table 2. Amino acid and protein contents, DIAAS score, and quality of all hydrolysates

Amino acid (g/100g of sample)	Yeast	Algae	Pea	Mushroom
Phenylalanine	2.71	3.01	4.24	0.83
Valine	3.87	4.72	4.32	1.68
Threonine	2.78	3.69	2.94	1.62
Tryptophan	0.84	0.80	0.65	0.38
Methionine	1.05	1.56	0.75	0.31
Leucine	4.68	6.16	6.76	1.66
Isoleucine	3.42	2.81	4.03	1.36
Lysine	4.90	5.16	6.43	1.62
Histidine	1.31	1.44	2.05	0.63
Taurine	0.02	0.03	0.03	0.03
Hydroxyproline	0.00	0.23	0.05	0.28
Aspartic Acid	6.01	7.17	9.92	2.33
Serine	2.46	2.87	3.99	1.23
Glutamic Acid	11.29	9.73	15.62	2.24
Proline	2.29	3.78	3.44	0.74
Lanthionine	0.11	0.13	0.21	0.00
Glycine	2.93	4.42	3.28	0.98
Alanine	4.58	6.59	3.57	1.84
Cysteine	0.71	0.79	0.55	0.19
Tyrosine	0.82	2.53	2.84	0.53
Hydroxylysine	0.15	0.00	0.00	0.00
Ornithine	0.49	0.08	0.07	0.88
Arginine	3.02	4.82	7.44	1.20
Total amino acid content	60.4	72.48	83.14	22.61
Protein content	69.7	83.03	74.05	25.71
DIAAS score (%)	93.2	86.72	65.02	72.03
Protein quality	Good	Good	Low	Low

3.3. Degree of hydrolysis and techno-functional properties

The results of the degree of hydrolysis and functional properties are presented in Fig. 1A-D.

Degree of hydrolysis of were 15, 20, 40, and 46%, for pea, algae, yeast and mushroom,

respectively. Yeast and mushroom hydrolysate showed the highest degree of hydrolysis, 40% and

46.1%, respectively. Degree of hydrolysis can influence the level of peptides releasing, the size,

conformation, and amino acid sequence impacts various functional and bioactive properties to the

hydrolysate product (Hall et al., 2017; Monaya et al., 2022).

Fig. 1. Techno-functional properties of protein hydrolysates. (A) Degree of hydrolysis; (B)

foaming capacity; (C) Oil holding capacity; (D) emulsifying capacity.

Algae protein hydrolysates showed the highest foam capacity, followed by pea and yeast protein

hydrolysates, while, mushroom protein hydrolysates showed the lowest foam capacity. Foaming

properties are related to partially denaturing a protein and exposing as its hydrophobic regions.

These regions can efficiently absorb air-water interfaces and form a lower interfacial tension, thus

enhancing foaming capability. However, extensive denaturation will reduce foaming capability of

the proteins (Mauer, 2003). Foaming in cell culture in bioreactor environment, may cause issues,

such as inhibiting cell development by limiting the surface area contact between the growth media

and the bioreactor headspace and decreasing oxygen transfer rates. Therefore, ideal hydrolysates

are expected not to increase the foaming that occurs spontaneously in a bioreactor when added to

media. The highest oil holding and emulsifying capacities were related to algae and pea protein

hydrolysates, followed by yeast and mushroom hydrolysates.

3.4. Cell morphology, growth, and viability

In this study, protein hydrolysates from different sources were applied to reduce or fully

replace serum. The following results demonstrate the changes in cellular morphology, growth, and

viability of zebrafish embryonic stem cells growing under different media conditions with or with

serum and at various concentrations of protein hydrolysates.

3.5. Impact of serum concentration

For culturing zebrafish ESC, 10% serum is the gold standard culture condition for proper

growth of cells. In Experiment I, different concentrations of serum, 0, 1, 2.5, 5 and 10%, were

evaluated for cell growth and doubling time. The impact of serum at various concentrations on the

parameters of cell growth and viability is demonstrated in Fig. 2.

After 24 hours of incubation in media containing 10 % serum, the test media were changed into

media containing 0, 1, 2.5, 5, and 10% serum, then the growth of the cells was studied on a daily

basis for three days using a phase-contrast microscope equipped with image analysis software.

Doubling time was increased from 14.85 h in 10% serum media to 46 h after completely removing

serum. As the serum concentration decreased, the cell density decreased significantly ($P < 0.05$).

By decreasing the serum from 10% to 5%, while cell numbers reduced from 1.4×10^6 to 1×10^6

cell/mL, the cell morphology has not changed (Fig. 2A). At a serum-free media condition, there

was a dramatic decrease in cell number from 1.4×10^6 to 1.8×10^5 cell/mL, and morphological

changes with an increase in rounded cells indicating the cell death (Fig. 2). Complete serum

removal causes cell starvation, apoptosis, and cell morphology changes which is in agreement with

other researchers findings (Pirkmajer and Chibalin, 2011; Terra et al., 2011).

In Experiment II, we evaluated the impact of different protein hydrolysates concentrations from

0.001 to 10 mg/mL in combination with different concentrations of serum (0, 5, and 10%) on cells

performance including cell growth (Fig. 3A, C, E, G, I) and cell viability (Fig. 3B, D, F, H, J).

Overall, the cell density and cell viability results indicated that 1 and 10 mg/mL protein hydrolysates significantly reduced the cell growth and cell viability in media containing high concentrations of protein hydrolysates ($P < 0.05$). Serum-free media containing 10 mg/mL of mushroom and yeast hydrolysates, indicated higher cell growth compared to serum containing media supplemented with 10 mg/mL of protein hydrolysates. Providing protein hydrolysates at high concentrations for serum containing media (5 and 10%), also reduced cell viability and cell growth.

Fig. 2. Cell growth parameters of ZEM2S cells at various serum concentrations. (A) micrographs of ZEM2S cells obtained on day 3; pictures in **the** right column are 20X enlarged from the inserted boxes, indicating the cell density and morphology, which significantly changes after removing serum; (B) doubling time indicating that completely removing serum will significantly increase the doubling time; (C) Cell viability results from PrestoBlue cell viability tests, indicating significantly higher cell activity in cells which were cultured in 10% serum.

Within 24 hours of adding a medium containing high concentrations of protein hydrolysate, the
cells lost their fibroblast-like morphology, and cell death was observed. In contrast, the effects of
low concentrations of protein hydrolysates (0.001, 0,001, and 0.1 mg/mL) on cells growth, and
viability illustrated that the lower concentrations of protein hydrolysates supported cell growth and
viability (Fig. 3). At lower concentrations (0.001, 0.01 and 0.1 mg/mL) of protein hydrolysates,
serum-free media containing protein hydrolysates improved cell growth and viability compared to
the serum-free media control. The results suggested that algae, mushroom, and pea protein
hydrolysates could potentially support cell growth in a serum-free environment. When serum was
reduced to 5%, these protein hydrolysates successfully increased cell number more than the media
containing 10% serum. Protein hydrolysates have been demonstrated to have more than a
nutritional effect on cells, including protective, antiapoptotic, and growth-promoting properties.
In Experiment III, the impacts of lower concentrations of protein hydrolysates (0.001, 0.01, and
0.1 mg/mL) in combination with low concentrations of serum (1 and 2.5%) on cell growth,
viability (Fig. 4), and LDH activity (Fig. 5B) were evaluated.

Fig. 3. ZEM2S cells growth in media containing different serum concentrations (0, 5 and 10%) supplementing with various protein hydrolysates including C: Controls (no protein hydrolysates; AL: Algae; MS: Mushroom; Pea; YS: Yeast, at different concentrations including: (A, B) 10 mg/mL; (C, D) 1 mg/mL; (E, F) 0.1 mg/mL; (G, H) 0.01 mg/mL; (I, J) 0.001 mg/mL. Pink: 10% Serum; Blue: 5% serum; Green: 0% serum. Note: In experiments with 10 mg/mL protein, pea protein was not dissolved in media. Thus, the data was not collected for high concentrations of pea. Cell growth (A, C, E, G, I); Cell viability (B, D, F, H, J).

The results indicated that providing protein hydrolysates at 0.001, 0.01 and 0.01 mg/mL in combination with 1 and 2.5% serum, will increase cell growth and cell viability more than control groups as serum-free, and media containing 1, 2.5, and 10% serum. These results indicate that for fish cell lines such as zebrafish fibroblast cells, serum in media could be reduced up to 90% without compromising cell growth, and cell viability, when protein hydrolysates are provided at relatively low concentrations. No significant differences were observed among the protein hydrolysates in terms of the cell growth and cell viability.

A study using several plant peptones (protein hydrolysates) in CHO cell lines revealed that adding these hydrolysates did not increase the nutritive value of proteins, but dramatically boosted cell growth (Burteau et al., 2003). Similar effects were reported when CHO cells were grown with 10% serum and algal extracts (Ng et al., 2020). On CHO-K1 cell lines with cottonseed hydrolysate, this synergistic impact of protein hydrolysates with other medium components was demonstrated by increased cell biomass (Babcock and Antosh, 2012). In addition, providing rapeseed protein hydrolysates reduced cell death rate and increase culture time up to 160 h instead of 80 h in CHO cells (Farges-Haddani et al., 2006). Soy hydrolysates promoted CHO cells growth without increasing protein productivity, while, yeast hydrolysate reduced cell growth, but increased the level of protein productivity glycosylation profile (Spearman et al., 2014).

While, lower concentrations of protein hydrolysates support cell growth, higher concentrations, (4 to 6 mg/mL) of protein hydrolysates, including soy (Lee et al., 2008) and fish gelatin (Hsieh et al., 2022), significantly decreased cell proliferation. Our previous study results also indicated that oyster, mussel, lugworm, cricket, and black soldier fly larvae protein hydrolysates reduce cell proliferation at high (1 and 10 mg/mL) concentrations when combined with higher concentrations of serum (10%) (Batish et al., 2022; Timoneda et al., 2024). In addition, wheat gluten hydrolysate between 6 and 12 mg/mL reduced cell proliferation (Radošević et al., 2016). Protein hydrolysates derived from various wastes, such as eggshells and carcasses, at 10 mg/mL had a similar dose-

dependent cytotoxic effect on bovine stem cells (Andreassen et al., 2020). This might be related

to the nutrient balance due to high amino acid or oligopeptide concentrations (Chun et al., 2007;

Hsieh et al., 2022; Zhang et al., 1994). On the other hand, low concentrations of whey protein

hydrolysate, *i.e.* 0.01-0.5 mg/mL, boosted osteoblastic cell proliferation, viability, and alkaline

phosphatase activity (Jo et al., 2020). Similar results were observed in bovine cell culture with

algal extract and various other industrial byproduct hydrolysates; cell growth increased compared

to serum-free conditions, however less than in serum-rich conditions (Andreassen et al., 2020). It

has been shown that the cell growth in the presences of protein hydrolysates at different

concentrations of serum depends on the degree of hydrolysis (Girón-Calle et al., 2010). However,

in this study, no correlation was observed between degree of hydrolysis (15-46%, Fig. 1A) and

cell growth. This could be explained by the fact that different sources of biomass were applied in

this study, while, other researchers only used one specific biomass with different degree of

hydrolysis (Girón-Calle et al., 2010).

Overall, protein hydrolysates from different sources could potentially enhance cell growth in low-

serum or serum-free media. Protein hydrolysates provide several benefits for cells, due to their

free amino acids, chemical compositions, peptides structures, bioactive properties such as anti-

apoptotic effects (Spearman et al., 2014), anti-oxidation properties (Ovissipour et al., 2013),

immunomodulatory effects, and antibacterial properties (Spearman et al., 2014). However, using

protein hydrolysates may have several drawbacks including pro-apoptotic properties at higher

concentrations, being undefined, leading to batch-to-batch quality challenges, being contaminated

with other by-products from enzymatic hydrolysis process (Spearman et al., 2014). For example,

microbial protein hydrolysates from yeast may contain specific peptides from the cell wall, or

bacteria hydrolysates may have high level of lipopolysaccharide (LPS), an endotoxin which is

difficult to completely remove from the final product (Lobo-Alfonso et al., 2010). In addition,

accumulation of certain amino acids such as tryptophan in the culture, may inhibit cell growth

(Alden et al., 2020). The rapidly growing cultivated meat industry requires innovative, sustainable,

and cost-efficient media components. Protein hydrolysates sourced from affordable and eco-

friendly biomass materials hold promise in bolstering this sector. Nonetheless, further

investigation into emerging biomass conversion methods, the influence of hydrolysis degree on

cellular function, nutritional content, flavor profiles, and the metabolomic implications of peptides

on cellular pathways is imperative.

Fig. 4. ZEM2S cells growth (A, C, E) and cell viability (B, D, F) in media containing different serum concentrations (0, 1, 2.5 and 10%) supplementing with various protein hydrolysates including C: Controls (no protein hydrolysates; AL: Algae; MS: Mushroom; Pea; YS: Yeast, at different concentrations including: 0.1 mg/mL; 0.01 mg/mL; 0.001 mg/mL. Pink: 10% Serum; Blue: 2.5% Serum; Orang: 1% Serum; Green: 0% serum.

3.6. Fluorescent staining and LDH activity

The results of the combined Hoechst and actin green staining for cells in media with different concentrations of serum (0, 1, 2.5, and 10%) and also media containing 1% serum with different protein hydrolysates are presented in Fig. 5A. The results showed that serum deprivation has direct effects on the cytoskeleton, although there was no discernible effect on the nucleus other than a reduction in numbers. At 10% serum control, a high number of nuclei and abundant actin staining were observed. At 2.5% serum, there was no visible effect on the nucleus other than a reduction in cell number. However, the actin filaments densities reduced, indicating a decrease in the cell number. At 1% and serum free media conditions, the cell number reduced significantly.

Fig. 5. (A) Staining images from cells grown at different concentrations (0, 1, 2.5, and 10%) of serum, and cells grown only in 1% serum containing 0.01 mg/mL of different protein hydrolysates; (B) LDH activity for cells in serum-free media (0%), 10% serum, 1 and 2.5% serum supplemented with 0.01 mg/mL protein hydrolysates.

Overall, the actin staining decreased along with the serum concentration, and subsequently altered the cell morphology. A similar result was reported and demonstrated that these was a decrease in actin protein staining under serum-starved circumstances (Wallenstein et al., 2010). The media with 1% serum and different protein hydrolysates, showed that the cells morphology was similar to the cell morphology in 10% serum containing media, indicating that at lower concentrations of serum, providing protein hydrolysates is critical to maintain cell performance.

The results of the LDH activity of the cells which were grown in serum-free media (0%), media containing 10% serum, 1% serum with protein hydrolysates, and 2.5% serum with protein hydrolysates are presented in Fig. 5B. Cells grown in serum-free media, showed significantly higher LDH activity compared to other conditions, indicating damage to the plasma membrane due to the starvation and lack of available serum. In contrast, cell culture at a serum concentration of 10% had the lowest LDH activity. These results suggest that the integrity of the cell membrane was compromised due to serum starvation, resulting in the release of LDH into the medium, which indicates apoptosis and necrosis. All other groups had significantly greater LDH activity compared to 10% serum control group. Cells were grown in 1% serum and protein hydrolysates had higher LDH activity compared to the 2.5% serum with protein hydrolysates. These results showed that protein hydrolysates have a critical role in maintaining the cell integrity and performance when serum is reduced or fully replaced.

3.7. GHG emissions

The results of the GHG emissions for different proteins, protein hydrolysates, FBS, and FBS in media at 10% and 1% concentrations, as well as 0.01 mg/mL of protein hydrolysates are presented in Table 3. Overall, the GHG emissions for algae was higher compared to other sources including FBS. During the enzymatic hydrolysis, the GHG emissions for water, electricity and enzymes were added to the protein sources. Protein hydrolysates from all the resources used in this study, had significantly higher GHG emissions, compared to FBS per FU. However, due to their low concentrations (0.01 mg/mL/10 mg/L) applications in media formulation, the GHG emission based on one liter media preparation is very small compared to serum.

Table 3. GHG emissions (kg CO_2 eq.) for different biomass resources based on the literature data and calculated data in our laboratory[1]

| Biomass | Production | Enzymatic hydrolysis process | | | Total (kg CO_2 eq./kg product) | GHG for serum in a regular media (10%) | GHG for serum (1) and protein hydrolysates (0.01 mg/mL) in reduced serum media |
		Enzyme	Water	Energy			
Algae	4.4	0.08	2.98	0.58	17.9	-	1.8×10^{-4}
Pea protein	0.94	0.16	5.96	1.2	8.26	-	8.26×10^{-5}
Mushroom	1.54	0.1	2.98	0.58	5.20	-	5.2×10^{-5}
Yeast peptone	-	-	-	-	3.2	-	3.2×10^{-5}
FBS	1.8	-	-	-	1.8	0.18	0.018

[1] Production GHG was collected from literature data: Algae (Clarens et al., 2010; Handler et al. 2012); Pea protein (Nette et al., 2016); Mushroom (American Mushroom, 2024); Yeast peptone (Carbon Cloud, 2024); FBS (Nikkhah et al., 2023). The enzymatic hydrolysis GHG was calculated based on the available literature, water, and electricity consumption during the processing. Yeast peptone was purchased directly from the provider, and no enzymatic hydrolysis was conducted in our research.

4. Conclusion

In this study, we assessed the impact of different concentrations of serum, both individually and in conjunction with varied concentrations of protein hydrolysates sourced from algae, mushrooms, peas, and yeast, on cell growth and viability over a short period (3 days). Reducing serum concentrations in the media, resulted in cell starvation, cell death, and significantly lower cell viability. When combined with higher concentrations of protein hydrolysates (1 and 10 mg/mL), cell apoptosis was observed in media containing 10% serum, indicating that providing higher concentrations of protein hydrolysates in conjunction with serum, will reduce the cell growth. While, lower concentrations of serum (1 and 2.5%) and protein hydrolysates (0.001, 0.01, and 0.1 mg/mL) enhanced cell growth, and cell viability, more than media containing 10% serum. Additionally, cellular fluorescence imaging analysis has shown that protein hydrolysates supplementation with reduced serum content in cell culture can improve cytoskeleton density to a similar extent as 10% serum control. GHG data showed that by applying protein hydrolysates at low concentrations, the GHG will be reduced by 90% compared to conventional media containing 10% serum.

Declaration of Competing Interest

The authors declared that they have no conflicts of interest to this work.

Data availability

Data will be made available on request.

Ethical statement - studies in humans and animals

This work did not involve the use of human and animal subjects.

Acknowledgment

This research was financially supported by the Agriculture and Food Research Initiative (AFRI) Sustainable Agricultural Systems program, Grant No. 2021-699012-35978 from the USDA National Institute of Food and Agriculture and Texas A&M AgriLife Research.

References

Alden, N., Raju, R., McElearney, K., Lambropoulos, J., Kshirsagar, R., Gilbert, A., Lee, K., 2020. Using metabolomics to identify cell line-independent indicators of growth inhibition for Chinese Hamster Ovary Cell-based bioprocesses. Metab. 2020, Vol. 10, Page 199 10, 199. https://doi.org/10.3390/METABO10050199

Andreassen, R.C., Pedersen, M.E., Kristoffersen, K.A., Beate Rønning, S., 2020. Screening of by-products from the food industry as growth promoting agents in serum-free media for skeletal muscle cell culture. Food Funct. 11, 2477–2488. https://doi.org/10.1039/C9FO02690H

Andreassen, R.C., Rønning, S.B., Solberg, N.T., Grønlien, K.G., Kristoffersen, K.A., Høst, V., Kolset, S.O., Pedersen, M.E., 2022. Production of food-grade microcarriers based on by-products from the food industry to facilitate the expansion of bovine skeletal muscle satellite cells for cultured meat production. Biomaterials 286, 121602. https://doi.org/10.1016/J.BIOMATERIALS.2022.121602

AOAC, 2006. Method 982.30, Official Methods of Analysis.

Babcock, J.F., Antosh, A., 2012. Partial replacement of chemically defined CHO media with plant-derived protein hydrolysates. Proc. 21st Annu. Meet. Eur. Soc. Anim. Cell Technol. (ESACT), Dublin, Ireland, June 7-10, 2009 295–298. https://doi.org/10.1007/978-94-007-0884-6_44

Batish, I., Brits, D., Valencia, P., Miyai, C., Rafeeq, S., Xu, Y., Galanopoulos, M., Sismour, E., Ovissipour, R., 2020. Effects of enzymatic hydrolysis on the functional properties, antioxidant activity and protein structure of black soldier fly (*Hermetia illucens*) protein. Insects 2020, Vol. 11, Page 876 11, 876. https://doi.org/10.3390/INSECTS11120876

Batish, I., Zarei, M., Nitin, N., Ovissipour, R., 2022. Evaluating the potential of marine
invertebrate and insect protein hydrolysates to reduce fetal bovine serum in cell culture Media
for cultivated fish production. Biomolecules 12. https://doi.org/10.3390/BIOM12111697

Bellani, C.F., Ajeian, J., Duffy, L., Miotto, M., Groenewegen, L., Connon, C.J., 2020. Scale-Up
Technologies for the Manufacture of Adherent Cells. Front. Nutr. 7, 575146.
https://doi.org/10.3389/FNUT.2020.575146/BIBTEX

Burteau, C.C., Verhoeye, F.R., Molsl, J.F., Ballez, J.-S., Agathos, S.N., Schneider, Y.-J., 2003.
Fortification of a protein-free cell culture medium with plant peptones improves cultivation
and productivity of an interferon-γ-producing CHO cell line. Vitr. Cell. Dev. Biol. 39, 291–
296.

Chun, B.H., Kim, J.H., Lee, H.J., Chung, N., 2007. Usability of size-excluded fractions of soy
protein hydrolysates for growth and viability of Chinese hamster ovary cells in protein-free
suspension culture. Bioresour. Technol. 98, 1000–1005.
https://doi.org/10.1016/J.BIORTECH.2006.04.012

Clarens, A.F., Resurreccion, E.P., White, M.A., Colosi, L.M., 2010. Environmental life cycle
comparison of algae to other bioenergy feedstocks. Environ. Sci. Technol. 44, 1813–1819.
https://doi.org/10.1021/ES902838N/SUPPL_FILE/ES902838N_SI_001.PDF

Ding, D., Yu, T., Du, B., Huang, Y., 2019. Collagen hydrolysate from Thunnus orientalis bone
induces osteoblast proliferation and differentiation. Chem. Eng. Sci. 205, 143–150.
https://doi.org/10.1016/J.CES.2019.04.040

FAO, 2020. The state of world fisheries and aquaculture 2020. 1–244.

FAO, 2011. Dietary protein quality evaluation in human nutrition. FAO Food Nutr. Pap 92, 1–66.

Farges-Haddani, B., Tessier, B., Chenu, S., Chevalot, I., Harscoat, C., Marc, I., Goergen, J.L.,
Marc, A., 2006. Peptide fractions of rapeseed hydrolysates as an alternative to animal proteins
in CHO cell culture media. Process Biochem. 41, 2297–2304.
https://doi.org/10.1016/J.PROCBIO.2006.06.002

Freshney, R.I., 2015. Culture of animal cells: a manual of basic technique and specialized
applications. John Wiley & Sons.

Girón-Calle, J., Alaiz, M., Vioque, J., 2010. Effect of chickpea protein hydrolysates on cell
proliferation and in vitro bioavailability. Food Res. Int. 43, 1365–1370.
https://doi.org/10.1016/J.FOODRES.2010.03.020

Hall, F.G., Jones, O.G., O'Haire, M.E., Liceaga, A.M., 2017. Functional properties of tropical banded cricket (Gryllodes sigillatus) protein hydrolysates. Food Chem. 224, 414–422. https://doi.org/10.1016/J.FOODCHEM.2016.11.138

Handler, R.M., Canter, C.E., Kalnes, T.N., Lupton, F.S., Kholiqov, O., Shonnard, D.R., Blowers, P., 2012. Evaluation of environmental impacts from microalgae cultivation in open-air raceway ponds: Analysis of the prior literature and investigation of wide variance in predicted impacts. Algal Res. 1, 83–92. https://doi.org/10.1016/J.ALGAL.2012.02.003

Hosios, A.M., Hecht, V.C., Danai, L. V, Steinhauser, M.L., Manalis, S.R., Vander, M.G., Correspondence, H., 2016. Amino Acids Rather than Glucose Account for the Majority of Cell Mass in Proliferating Mammalian Cells. Dev. Cell 36, 540–549. https://doi.org/10.1016/j.devcel.2016.02.012

Hsieh, C.H., Wang, T.Y., Tung, B.C., Liu, H.P., Yeh, L. Te, Hsu, K.C., 2022. The Hydrolytic Peptides of Soybean Protein Induce Cell Cycle Arrest and Apoptosis on Human Oral Cancer Cell Line HSC-3. Molecules 27, 2839. https://doi.org/10.3390/MOLECULES27092839/S1

Jo, K., Hong, K.B., Suh, H.J., 2020. Effects of the Whey Protein Hydrolysates of Various Protein Enzymes on the Proliferation and Differentiation of 3T3-E1 Osteoblasts. Prev. Nutr. Food Sci. 25, 71. https://doi.org/10.3746/PNF.2020.25.1.71

Kim, J.Y., Kim, Y.G., Han, Y.K., Choi, H.S., Kim, Y.H., Lee, G.M., 2011. Proteomic understanding of intracellular responses of recombinant Chinese hamster ovary cells cultivated in serum-free medium supplemented with hydrolysates. Appl. Microbiol. Biotechnol. 89, 1917–1928. https://doi.org/10.1007/S00253-011-3106-9/FIGURES/5

Lee, Y.K., Kim, S.Y., Kim, K.H., Chun, B.H., Lee, K.H., Oh, D.J., Chung, N., 2008. Use of soybean protein hydrolysates for promoting proliferation of human keratinocytes in serum-free medium. Biotechnol. Lett. 30, 1931–1936. https://doi.org/10.1007/S10529-008-9796-0/FIGURES/4

Lobo-Alfonso, J., Price, P., Jayme, D., 2010. Benefits and limitations of protein hydrolysates as components of serum-free media for animal cell culture applications protein hydrolysates in serum free media. Protein Hydrolysates Biotechnol. 55–78. https://doi.org/10.1007/978-1-4020-6674-0_4/FIGURES/6

Logarušić, M., Slivac, I., Srček, V.G., Berljavac, S., Pavunc, A.L., Radošević, K., 2021. Protein Hydrolysates from Flaxseed Oil Cake as a Media Supplement in CHO Cell Culture. Resour.

2021, Vol. 10, Page 59 10, 59. https://doi.org/10.3390/RESOURCES10060059

Lu, M., Zhao, X.H., 2018. The Growth Proliferation, Apoptotic Prevention, and Differentiation
Induction of the Gelatin Hydrolysates from Three Sources to Human Fetal Osteoblasts (hFOB
1.19   Cells).   Mol.   2018,   Vol.   23,   Page   1287   23,   1287.
https://doi.org/10.3390/MOLECULES23061287

Mariotti, F., Tomé, D., Patureau Mirand, P., Ois Mariotti, F., Tométom, D., 2008. Converting
Nitrogen into Protein—Beyond 6.25 and Jones' Factors. Crit. Rev. Food Sci. Nutr. 48, 177–
184. https://doi.org/10.1080/10408390701279749

Mattila, P., Salo-Va¨a¨na¨nen, P., Va¨a, V., Va¨a¨na, V., Va¨a¨na¨nen, V., Ko¨nko¨, Karoliina,
Ko¨nko, K., Ko¨nko¨, Ko¨nko¨, Aro, H., Jalava, T., 2002. Basic Composition and Amino Acid
Contents of Mushrooms Cultivated in Finland. J. Agric. Food Chem. 50, 6419–6422.
https://doi.org/10.1021/JF020608M

Mauer, L., 2003. PROTEIN | Heat Treatment for Food Proteins. Encycl. Food Sci. Nutr. 4868–
4872. https://doi.org/10.1016/B0-12-227055-X/00988-3

Monaya, K.J.M., Gomez, H.L.R., Tejano, L.A., Peralta, J.P., 2022. Production and
Characterization of Protein Isolates and Hydrolysates from Slipper Cupped Oyster
(Crassostrea iredalei). Philipp. J. Sci. 151, 853–861. https://doi.org/10.56899/151.03.06

Nette, A., Wolf, P., Schlüter, O., Meyer-Aurich, A., 2016. A Comparison of Carbon Footprint and
Production Cost of Different Pasta Products Based on Whole Egg and Pea Flour. Foods 2016,
Vol. 5, Page 17 5, 17. https://doi.org/10.3390/FOODS5010017

Ng, J.Y., Chua, M.L., Zhang, C., Hong, S., Kumar, Y., Gokhale, R., Ee, P.L.R., 2020. Chlorella
vulgaris Extract as a Serum Replacement That Enhances Mammalian Cell Growth and
Protein   Expression.   Front.   Bioeng.   Biotechnol.   8,   564667.
https://doi.org/10.3389/FBIOE.2020.564667/BIBTEX

Nikkhah, A., Rohani, A., Zarei, M., Kulkarni, A., Batarseh, F.A., Blackstone, N.T., Ovissipour,
R., 2023. Toward sustainable culture media: Using artificial intelligence to optimize reduced-
serum   formulations   for   cultivated   meat.   Sci.   Total   Environ.   894,   164988.
https://doi.org/10.1016/J.SCITOTENV.2023.164988

Ovissipour, M., Rasco, B., Shiroodi, S.G., Modanlow, M., Gholami, S., Nemati, M., 2013.
Antioxidant activity of protein hydrolysates from whole anchovy sprat (Clupeonella
engrauliformis) prepared using endogenous enzymes and commercial proteases. J. Sci. Food

Agric. 93, 1718–1726. https://doi.org/10.1002/JSFA.5957

Pacheco-Aguilar, R., Mazorra-Manzano, M.A., Ramírez-Suárez, J.C., 2008. Functional properties

of fish protein hydrolysates from Pacific whiting(Merluccius productus) muscle produced by

a   commercial   protease.   Food   Chem.   109,   782–789.

https://doi.org/10.1016/J.FOODCHEM.2008.01.047

Pirkmajer, S., Chibalin, A. V, 2011. Serum starvation: caveat emptor. Am. J. Physiol. Physiol.

301, C272–C279.

Radošević, K., Dukić, B., Andlar, M., Slivac, I., Gaurina Srček, V., 2016. Adaptation and

cultivation of permanent fish cell line CCO in serum-free medium and influence of protein

hydrolysates   on   growth   performance.   Cytotechnology   68,   115–121.

https://doi.org/10.1007/S10616-014-9760-X/FIGURES/4

Safi, C., Charton, M., Pignolet, O., Silvestre, F., Vaca-Garcia, C., Pontalier, P.Y., 2013. Influence

of microalgae cell wall characteristics on protein extractability and determination of nitrogen-

to-protein conversion factors. J. Appl. Phycol. 25, 523–529. https://doi.org/10.1007/S10811-

012-9886-1/FIGURES/1

Shahidi, F., Han, X.Q., Synowiecki, J., 1995. Production and characteristics of protein

hydrolysates   from   capelin   (Mallotus   villosus).   Food   Chem.   53,   285–293.

https://doi.org/10.1016/0308-8146(95)93934-J

Spearman, M., Lodewyks, C., Richmond, M., Butler, M., 2014. The bioactivity and fractionation

of peptide hydrolysates in cultures of CHO cells. Biotechnol. Prog. 30, 584–593.

https://doi.org/10.1002/BTPR.1930

Stout, A.J., Mirliani, A.B., Rittenberg, M.L., Shub, M., White, E.C., Yuen, J.S.K., Kaplan, D.L.,

2022. Simple and effective serum-free medium for sustained expansion of bovine satellite

cells for cell cultured meat. Commun. Biol. 2022 51 5, 1–13. https://doi.org/10.1038/s42003-

022-03423-8

Taylor, W.H., 1957. Formol titration: An evaluation of its various modifications. Analyst 488–

498. https://doi.org/10.1039/AN9578200488

Terra, L.F., Garay-Malpartida, M.H., Wailemann, R.A.M., Sogayar, M.C., Labriola, L., 2011.

Recombinant human prolactin promotes human beta cell survival via inhibition of extrinsic

and   intrinsic   apoptosis   pathways.   Diabetologia   54,   1388–1397.

https://doi.org/10.1007/S00125-011-2102-Z/FIGURES/4

The Climate Intelligence Platform - CarbonCloud [WWW Document],. URL
https://carboncloud.com/ (accessed 3.14.24).

Timoneda, A., Amirvaresi, A., Ovissipour, R., 2024. Black Soldier Fly Bioconversion to
Cultivated Meat Media Components Using Blue Catfish Gut Microbiome. bioRxiv
2024.01.20.576372. https://doi.org/10.1101/2024.01.20.576372

van der Valk, J., Brunner, D., De Smet, K., Fex Svenningsen, Å., Honegger, P., Knudsen, L.E.,
Lindl, T., Noraberg, J., Price, A., Scarino, M.L., Gstraunthaler, G., 2010. Optimization of
chemically defined cell culture media – Replacing fetal bovine serum in mammalian in vitro
methods. Toxicol. Vitr. 24, 1053–1063. https://doi.org/10.1016/J.TIV.2010.03.016

Wallenstein, E.J., Barminko, J., Schloss, R.S., Yarmush, M.L., 2010. Serum starvation improves
transient transfection efficiency in differentiating embryonic stem cells. Biotechnol. Prog. 26,
1714–1723. https://doi.org/10.1002/BTPR.472

Yao, T., Asayama, Y., 2017. Animal-cell culture media: history, characteristics, and current issues.
Reprod. Med. Biol. 16, 99–117.

Yasumatsu, K., Sawada, K., Moritaka, S., Misaki, M., Toda, J., Wada, T., Ishii, K., 1972.
Whipping and emulsifying properties of soybean products. Agric. Biol. Chem. 36, 719–727.
https://doi.org/10.1080/00021369.1972.10860321

Zhang, Y., Zhou, Y., Yu, J., 1994. Effects of peptone on hybridoma growth and monoclonal
antibody formation. Cytotechnology 16, 147–150.
https://doi.org/10.1007/BF00749901/METRICS