Research disturbance negatively impacts incubation behaviour of female Great Tits

Léanne Clemencin[1], Emilio Barba[2] & David Diez-Méndez[3,4]*

[1] Université Claude Bernard Lyon 1, 69622 Villeurbanne Cedex, France

[2] "Cavanilles" Institute of Biodiversity and Evolutionary Biology, University of Valencia, Valencia,

Spain

[3] Biology Centre of the Czech Academy of Sciences, Institute of Entomology, České Budějovice,

Czech Republic.

[4] University of South Bohemia, Faculty of Science, České Budějovice, Czech Republic.

* Corresponding author

Abstract

Human-induced disturbance is perceived by avian species as a predation risk. However, the anti-

predatory behaviour triggered by these non-lethal events can have negative impacts on

reproduction and offspring survival. Research on breeding birds often imply visiting their nests and

likely disrupt parental behaviour, but nest visits where females are not directly handled have

previously been overlooked as important disturbance events. This study focuses on the impacts of

short visits to the nest of incubating Great tit (*Parus major*) females. We investigated how long they

stay away from the nest (off-bout) after a disturbance, their possible compensatory behaviour once

they resume incubation (longer on-bouts), and the effects on daily incubation rhythms. We used

three years of data from two breeding populations to assess the consequences of disturbance in

two scenarios: when the female is present in the nest and flushed out, and when the female is

absent. We found that following a disturbance, the immediate off-bout was longer, both when the

female was present and absent, with the magnitude of the disturbance being greater when females

were flushed. The females did not compensate by longer on-bouts afterwards, therefore research

disturbance altered daily incubation behaviour by decreasing the total time spent on the nest as a

function of the number of daily disturbance events. Females that alter their behaviour in response

to the perception of predation risk would perform longer incubation periods that result in lower

hatching rates. These effects of research on female behaviour should be considered when planning

field experiments.

Keywords: Incubation behaviour, anti-predator response, off-bouts, *Parus major*, human

disturbance

Introduction

Human-induced disturbance events are perceived similarly to predation risk ("risk-disturbance hypothesis", Frid & Dill, 2002), although they are usually non-lethal by nature (Cresswell, 2008). In response, animals may adjust their behaviour (Beale & Monaghan, 2004; Perona et al., 2019), physiology (Eggers et al., 2006; Coslovsky & Richner, 2011) or life-history traits (Hutfluss & Dingemanse, 2019) to the magnitude of perceived predation risk (Lima & Bednekoff, 1999). Anti-predatory behavioural responses can have a negative impact on reproduction and offspring survival, even if they contribute to a breeding individual not being predated. These effects can be both indirect, by facilitating nest predation in the absence of parents (Stien & Ims, 2016) or via maternal effects on offspring (Coslovsky & Richner, 2011), and direct, by altering incubation behaviour (Baudains & Lloyd, 2007) or offspring feeding (Verhulst et al., 2001). Direct effects may impair offspring growth (Remacha et al., 2016) and influence their survival and subsequent recruitment (Blas et al., 2007). Such non-consumptive impacts are often associated with human recreation, although research activities during the breeding season may have analogous effects.

Nest disturbance is inevitably associated with avian reproductive research (Lenington, 1979). It can be assumed that research activities are more disruptive than recreational activities, as they usually involve handling individuals, visiting nests or manipulating eggs, but only temporarily. Attracting predators to nest sites has traditionally been the main impact that researchers have been concerned about, although an early review showed no clear patterns overall (Götmark, 1992). Later studies not only found no measurable negative effects (Schaub et al., 1992; Lloyd et al., 2000; Weidinger, 2008; Hu et al., 2020) but even positive effects of research activities (Ibáñez-Álamo & Soler, 2010, as the presence of humans near nests may deter mammal predators rather than attracting them (reviewed in Ibáñez-Álamo et al., 2012). However, as with recreational activities, nest predation is only one possible indirect effect, and other direct effects of research activities on breeding individuals may also be at play.

Breeding passerines that are the subject of research activities are regularly visited to record their

reproductive phases from nest building to offspring feeding (e.g., Cresswell & McCleery, 2003;

Andreasson et al., 2018; Arct et al., 2022). Nest visits while incubating have been dismissed as

important disturbance events as there are almost no previous assessments of their impact. The

incubating period is of particular interest because females, in gyneparental species, spend up to

80% of their active time in the nest to incubate (on-bouts), leaving it only briefly for foraging or other

self-care activities (off-bouts) (e.g., Ardia et al., 2010; Johnson et al., 2013; Ospina et al., 2018;

Diez-Méndez et al., 2021a). Regular incubation rhythms facilitate researchers to encounter females

in the nest. Females are constraint to maintain an adequate clutch temperature for embryonic

development (Nord & Nilsson, 2011; DuRant et al., 2013), but show certain plasticity adjusting the

number and length of their bouts to external conditions (Conway & Martin, 2000a; Diez-Méndez et

al., 2021a). Incubating females undergo high energy costs (Nord & Williams, 2015) and may

abandon their nest when predation risk is high, especially in the first half of the period when,

according to life-history theory, the trade-off between current and future reproduction favours the

latter (Clutton-Brock, 1991; Stearns, 1992). For this reason, assessments of research activity in this

period have only been conducted when visiting a nest implies high desertion rates, like after

handling incubating females (up to 41% of desertions, Kania, 1989; Kilgas et al., 2007; Dubiec,

2011) or after experimental drastic reductions in clutch size (e.g., simulating partial predation,

Verboven & Tinbergen, 2002; Johnston, 2011; Zieliński, 2020).

Visiting a nest and flushing the female to determine clutch size, a common practise, may also affect

incubation behaviour, but to a lesser extent than the above-mentioned disturbances. Simply visiting

a nest is perceived by females as a predation risk and triggers anti-predatory behaviours such as

hissing (Krams et al., 2014; Zub et al., 2017; Dutour et al., 2020; Van Iersel et al., 2023). Following

previous studies using stuffed predators, we may expect an effect on incubation behaviour if

females reduce their activity in the nest. Nevertheless, lengthening both off- and on-bouts, ends up

with having no negative impacts in nest attendance (Conway & Martin, 2000b; Ghalambor & Martin,

2002; Kovařík & Pavel, 2011; Basso & Richner, 2015a; Tkaczyk et al., 2023, but see Basso &

Richner, 2015b). However, similar predation experiments resulted in females reducing their activity

by returning later after being exposed to the predator (longer off-bouts), and limiting the time spent

in the nest (shorter on-bouts), resulting in an overall decrease in nest attendance (Zanette et al.,

2011; Ibáñez-Álamo & Soler, 2012). Repeated prolonged absences of the incubating female may

have dramatic consequences on embryonic development due to increased temperature variability

(Olson et al., 2006) and longer incubation period that lead to reduced egg hatchability (Nord &

Nilsson, 2011; Macdonald et al., 2013; Diez-Méndez et al., 2020) and survival rates after fledging

(Hepp et al., 2006; Berntsen & Bech, 2016; Ospina et al., 2018, reviewed in DuRant et al., 2013).

Here we investigated the effects of research disturbance during the incubation period in two Great

tit (*Parus major*) breeding populations. We assessed the effects of disturbance and potential

recovery time in two common scenarios when conducting nest visits: 1) the incubating female is

present in the nest (an incubating on-bout) and is consequently flushed, and 2) the female is absent

(during an off-bout). Female flushing has been dismissed as an important disturbance because, to

our knowledge, its effects have not been considered so far, despite Great tits are one of the most

common model species in the wild. We hypothesized that visiting a nest-box would have a negative

effect, albeit limited, on incubation behaviour in both scenarios, i.e., either when the female is

present and flushed or when she is absent. A limited negative effect would mean that females

lengthen the immediate off-bout after a disturbance event, followed by a longer on-bout to

compensate. This limited effect would not cause overall changes in daily incubation behaviour. We

hypothesised that the effect of research disturbance in the absence of a female would still elicit a

noticeable behavioural response because foraging territories are small (Naef-Daenzer & Keller,

1999) and females can still detect intruders. We also expected that flushing a female triggers a

stronger response than when females are absent as it is perceived as a riskier encounter.

Material and Methods

 Study area and species

 Great tits are small (12.5 – 15 cm) secondary hole-nesting passerines that naturally breed in cavities

 but also readily accept nest-boxes (Perrins, 1979). They lay one egg per day and usually start

 incubating when the clutch is complete, although there is a great variation depending on external

 conditions (Diez-Méndez et al., 2021b). They are assisted gyneparental incubators (Nord & Williams,

 2015), which means that only the females incubate and the male may assist by bringing food to the

 nest (Bambini et al., 2019), yet the female regularly leaves the nest to forage and perform other self-

 care activities (intermittent incubation).

 Breeding data was collected in two Mediterranean nest-box populations during three consecutive

 breeding seasons (2015-2017) as part of ongoing long-term projects (e.g., (Álvarez et al., 2013)):

 Sagunto (39.70°N, 0.25°W) is part of an extensive monoculture of orange trees (*Citrus aurantium*)

 at sea level (30 m a.s.l.), while Pina (40.02°N, 0.63°W) is part of a mountain range (1200 m a.s.l.)

 which contains a mixed forest dominated by maritime pines (*Pinus pinaster*) with scattered stands

 of Portuguese oaks (*Quercus faginea*).

 Field procedures

 We monitored the first clutches of breeding Great tits in both populations to determine when the

 first egg was laid in each nest and to initiate the protocol for measuring incubation behaviour (see

 Diez-Méndez et al., 2021b, 2021a for further details). In brief, after the first egg was laid, we attached

 a probe (30-gauge wire, accuracy ± 1.0°C; Onset Computer Corporation, USA) to the centre of the

 clutch, which was connected to a thermocouple data logger (HOBO UX100-014M Single Channel,

 Onset Computer Corporation, USA), and recorded the temperature every 10s until the first egg

 hatched. The probe was inserted into a plastic egg (Factory Direct Craft Supply, USA), filled with a

 lubricant (Clear Glide, Ideal industries, USA), that was similar in shape and colour to the Great tit

 eggs but slightly smaller (15.9 × 11.5 mm versus 18.1 × 13.3 mm; average egg size from Sagunto;

 Encabo et al. 2001). In addition, we attached a Thermochron iButton data logger (accuracy ± 0.5°C,

model DS1922L-F5, Maxim Integrated) to the upper inner part of each nest-box wall to record the

local ambient temperature every 520 s.

We visited each nest-box at least three times during the incubation period: 1) when we first found

an incubating female or warm eggs during a regular morning check, 2) five days into the incubation

period, when we visited the nest to check the probe and clutch size, and 3) eleven days into the

incubation period, when we began monitoring egg hatching. Each time we visited a nest, we noted

the time and whether the female was present or absent from the nest. If the female was present,

we always flushed her out to check the contents of the nest and experimental materials. The nest-

boxes of these populations allow the front wall to be removed during checkings, so the female

usually left the nest as soon as we opened the nest-box. For the few more stubborn females we

carefully push them out. The incubating females were never handled in order to be as unobtrusive

as possible. The standard visits to the nest-boxes lasted less than one minute (i.e., a short

disturbance), but depending on the year and ongoing experiments (e.g., measuring and weighing

individual eggs), these visits lasted longer (i.e., 1-5 min, long disturbance events).

Incubation behaviour and disturbance measurements

Incubation behaviour of Great Tit females was assessed using recorded nest-cup temperatures in

each nest during the incubation period, We defined the onset of incubation as the first time nest

attentiveness exceeded 50% of the active day (incubation day 1), and the incubation period lasted

to the day before the first egg hatched (following Cresswell & McCleery, 2003; Simmonds et al.,

2017; Diez-Méndez et al., 2021b). A female's active day was considered as the period between the

first off-bout in the early morning and the last on-bout in the evening to overnight.

To identify off- and on-bouts, we used the software *Rhythm* (Cooper & Mills, 2005) and followed

the criteria of setting off-bouts to last at least 2 min and indicating a temperature drop of 2.0°C. We

then visualized the Rhythm output in Raven (Cornell Laboratory of Ornithology) and modified off-

bouts when we detected a misidentification or visualized clear off-bouts that did not meet the

criteria. Only off-bouts with a duration of less than 120 minutes were taken into account, as longer

off-bouts are rare and considered outliers (~1.5%, Diez-Méndez et al., 2021a). A total of 46 nests,

29 from Sagunto and 17 from Pina, were suitable for assessing the effects of research visits on

female incubation behaviour. We recorded 273 disturbances and analysed the on- and off-bout

duration before, during and after an event, totalling 1,638 bouts (see below, Suppl. Mat. Table S1).

Statistical analyses

We performed statistical analyses with the software R 4.3.1 (R Core Team, 2023). Six linear (LMMs)

and one generalised linear mixed-effects model (GLMMs) were created using the package *lme4*

(Bates et al., 2015). We built a model to assess the off-bout duration when a female was absent

during a nest-box visit (disturbance) using the off-bout temporal sequence (before, during and after

disturbance) as a fixed factor together with day of incubation, ambient temperature (quadratic term,

poly- function), time of the day (quadratic term, poly- function), calendar date, population and year,

including nest-box as a random effect. A second model was built to assess the off-bout duration

when a female was present in the nest during a visit, using off-bout categorization (before,

immediately after the disturbance, and the next one) as a fixed factor. The third model looked at

the duration of the on-bouts when a female was absent, using a fixed factor categorizing on-bouts

before (one) and after the disturbance (two on-bouts). The fourth model dealt with the duration of

the on-bouts when the incubating female was present during a visit to the nest, using a fixed effect

to categorize on-bouts before (one) and after the disturbance (two on-bouts). We did not consider

the on-bout that abruptly finished with the disturbance because its duration was shortened by the

visit itself. Bout duration was measured in minutes and log transformed in every model. Two more

LMMs were built to determine whether the duration of the disturbance influenced the subsequent

off-bout duration. Off-bout duration when a female was present or absent was used as the

dependent variable, and disturbance duration was an independent factor with two levels: short and

long. The rest of the fixed and random factors were similar to the previous LMMs. Our final model

(GLMM, binomial family) addressed daily nest attentiveness and added disturbance intensity as a

fixed factor. The number of disturbance events per day could be up to two if the experiments

required it. For each day on which at least one disturbance event occurred, a preceding or

subsequent day on which no disturbance occurred was also selected. In addition, day of incubation,

mean daily ambient temperature (quadratic, poly- function), calendar date, population and year

were included as fixed factors, and nest- box as a random factor in the model. All numerical

variables were centred. Normality, homogeneity of variance and collinearity were visually checked

for the LMMs models using the *performance* package (Lüdecke et al., 2021). The resulting figures

were created with *ggplot2* (Wickham, 2016) after calculating the estimates $\pm$ 95%CI with the

package *Effects* (Fox & Hong, 2009).

Results

Our data showed the general pattern of average off-bouts being three times shorter than on-bouts

(~10 vs. ~30 min, Supp. Mat. Table S1). When the female was absent from the nest during a nest-

box visit, the ongoing off-bout lasted longer than both the previous and following off-bouts (Table

1, Fig. 1a). However, we found no detectable effect on the first on-bout after a disturbance

compared to the previous and the following on-bouts (Table S2, Fig. 1c). When the female was

present in the nest during a nest-box visit, the immediate off-bout also lasted longer than the

previous and following off-bouts (Table 1, fig 1b), and the two on-bouts following the disturbance

event tended to be longer than before, but the effect was not clear (Table S2, Fig. 1d).

Finding a female in the nest, and flushing her out, led to a subsequent longer off-bout compared to

off-bouts from females that were out of the nest. While in the latter females extended their off-bout

by ~35%, the former performed ~75% longer off-bouts (Table S1). The duration of the disturbance

also affected how long a female stayed off the nest, but only if she was present in the nest during

the visit: a long disturbance caused a longer off-bout (Table 2, see Table S3 for absent females).

Moreover, only when females were absent, the off-bouts after a disturbance event tended to be

shorter in the Sagunto population (an anthropized orange plantation) than in Pina (a mixed forest)

(Table S3).

Disturbances in the nest were reflected in daily nest attentiveness. Females spent less time

incubating when they underwent two disturbance events in a single day (estimate = 67.4, CI = 64.1

– 70.6) than when there was only one (estimate = 71.3, CI = 68.7 – 73.7), and less time when there

was only one daily event than when there was no disturbance at all (estimate = 76.1, CI = 74.0 –

78.1) (Table 3. Fig 2).

We also found in our data subset clear effects of both the incubation day and ambient temperature.

As incubation progressed, females, either absent or present in the nest, shortened their off-bouts

(Table 1) and increased their daily incubation effort (Table 3). Females also lengthened their off-

bouts at increasing ambient temperatures, but only if they were present in the nest during a visit

(Table 1 & 2). Lastly, females decreased their daily nest attendance at increasing daily ambient

temperatures (Table 3).

Discussion

In this study, we found evidence that research disturbance negatively affects Great tit incubation

behaviour. Our initial hypothesis was that females reacted to a disturbance event by staying away

from the nest longer than expected given their incubation rhythms, but we also expected that this

anti-predatory behaviour, observed in previous studies (Kovařík & Pavel, 2011; Ibáñez-Álamo &

Soler, 2012; Croston et al., 2021; Tkaczyk et al., 2023), was compensated with longer on-bouts that

ended up balancing any negative effect on nest attendance. However, we found that 1) the

response of females was restricted to the ongoing (absent females) or first (present females) off-

bout after a disturbance, i.e., a short recovery period; 2) females did not clearly compensate for

their prolonged absence from the nest by lengthening subsequent on-bouts; and consequently,

and despite the limited anti-predator response, 3) daily incubation rhythms were negatively affected

by the disturbance events (i.e., lower nest attentiveness).

Breeding females are flexible in their response to different perceived predation risks (Lima, 2009;

Martin & Briskie, 2009). We found that females that were present in the nest at the time of a

disturbance stayed away from the nest longer than absent females, presumably because they

perceived being flushed as a greater risk than simply recognising an intruder from a distance.

Flushed females also stayed away from the nest longer if the researcher performed additional

activities during a visit (e.g., measuring eggs), in contrast to absent females, which seemed

unaffected by prolonged visits. Experiments simulating the presence of predators near the nest

generally result in incubating females leaving the nest for longer periods (Kovařík & Pavel, 2011;

Ibáñez-Álamo & Soler, 2012; Croston et al., 2021; Tkaczyk et al., 2023), although the duration and

magnitude of the response is species-specific (Ghalambor & Martin, 2002; Zanette et al., 2011;

Ghalambor et al., 2013) and/or dependent on predator type (Basso & Richner, 2015a). Absent

females appeared to evaluate the disturbance caused by the researchers differently. While flushed

females remained near the nest-box, called and displayed mobbing behaviour before flying away

(DDM pers. obs.), absent females may encounter a researcher in their nest when they return to

incubate or while already foraging nearby. Flushed females may refrain from foraging or other self-

care activities during part of their off-bout to increase vigilance (Morosinotto et al., 2013; Croston

et al., 2021), at least until the researcher has left the nest, which could partly explain their longer

off-bouts. An unexpected trend also emerged between populations when females were absent from

the nest, although more data would be needed to assess a clear effect. In Sagunto, off-bouts of

absent females tended to be shorter than in Pina. Given the anthropogenic nature of Sagunto, an

extensive orange tree plantation, Great tit females could either be partially habituated to humans

near the tree in which they built the nest, or the structure of the trees (short trees with crowns that

generally reach the ground) would partly conceal the researcher from the foraging the bird.

Females were still constrained by a clutch that requires a suitable thermal range for embryo

development (Drent, 1975; DuRant et al., 2013). In elicited long absences from the nest, after being

flushed, higher air temperature allowed females to stay off the nest for longer, pointing towards

thermal constraints in the magnitude of the anti-predatory response, a trade-off between their own

risk of being predated and the embryo survival that air temperature ameliorates. This is in line with

a milder response when females were absent, where no association between air temperature and

off-bout duration was found. Additionally, the general relationship between day of incubation and

off-bout duration was expected (Diez-Méndez et al., 2021a) because when embryos develop their

circulatory system, eggs cooling rates increase, causing a shortening in off-bouts (Cooper & Voss,

2013).

Reduced activity near the nest, either by prolonged or less frequent off-bouts is the expected

default behaviour when predation risk increases (Skutch, 1957; Conway & Martin, 2000b). However,

longer absences are usually compensated by longer incubation bouts, levelling out overall nest

attendance (Kovařík & Pavel, 2011; Ibáñez-Álamo & Soler, 2012; Basso & Richner, 2015b; Tkaczyk

et al., 2023). In our study, despite the limited females' response to a disturbance event, their daily

incubation rhythms were altered, reducing their time on the nest. Reduced nest attendance may

lengthen incubation period (Nord & Nilsson, 2012) and reduce egg hatchability (Diez-Méndez et al.,

2020) in the short term and reduce offspring survival in the long-term (Berntsen & Bech, 2016;

Ospina et al., 2018). Furthermore, Great tit females did not show a habituation response (see also

Croston et al., 2021), and a second disturbance event within a day maintained the decline in daily

nest attentiveness. Based on our results, it can be assumed that field experiments requiring regular,

repeated visits during the incubation period may result in at least a decrease in egg hatchability

compared to completely undisturbed nests. However, we were unable to link offspring productivity

and egg hatchability to disturbance in our population, as all the nests were involved in different

experiments and each of them was visited at least three times during the incubation period.

Our models showed a certain amount of variance that was not explained by the variables we measured. Physiological conditions at the individual level, which were not considered in this study, may explain some of the observed variability. For example, prolactin concentration following a disturbance may regulate nest attentiveness (Hope et al., 2020), and poorer body condition could lead to higher nest desertion rates (Dubiec, 2011). Female personality could also explain the observed variability, as shy individuals take longer to return to the nest after being exposed to a novel object (Cole & Quinn, 2014), although a long-term study on the effects of human recreation on Great tit breeding parameters showed no measurable within-population differences given their risk-taking personality (Hutfluss & Dingemanse, 2019). Overall, Great tits are a resilient species that tolerates high degree of manipulation in the nest compared to other passerines (e.g., shrikes Tryjanowski & Kuźniak, 1999; Golawski & Zduniak, 2022). However, we must not lose sight of the importance of reducing disturbance from nest building to egg incubation. Visiting a nest while the female is absent is desirable, although less likely if it is not intentional, because incubating Great tit females sit on the nest for about 75% of their active day (Bueno-Enciso et al., 2017; Schöll et al., 2019; Diez-Méndez et al., 2021a). Handling females in the nest during incubation can lead to nest abandonment (Dubiec, 2011) but nest visits also generate disturbances that disrupt incubation rhythms, a problem that should be taken into account when planning experimental approaches in the breeding populations.

Acknowledgments

We would like to thank all the collaborators and students for their assistance during the breeding seasons 2015-2017 in Sagunto and Pina, and specially José Verdejo for his help in Pina. This study was supported by projects CGL2013-48001-C2-1-P, CGL2016-79568-C3-1-P (former Spanish Ministry of Economy and Competitiveness, MINECO) and PID2021-122171NB-I00/ AEI/10.13039/501100011033/ FEDER, UE (Spanish Ministry of Science and Innovation). LC benefitted from a BRMIE (French regional grant for international student mobility), DDM benefitted

332 from an FPI fellowship (BES-2014-069191, MINECO) at the time, and is currently supported by the

333 ERC (European Research Council) grant BABE 805189 and a GAČR Star grant (22-17593M).

Data availability statement

336 Data will be available from the Dryad Digital Repository

References

Álvarez E, Belda EJ, Verdejo J, Barba E (2013) Variation in Great Tit nest mass and composition
and its breeding consequences: a comparative study in four Mediterranean habitats. Avian
Biol Res 6: 39–46. doi: 10.3184/175815513X13609517587237

Andreasson F, Nord A, Nilsson J-Å (2018) Experimentally increased nest temperature affects body
temperature, growth and apparent survival in blue tit nestlings. J Avian Biol 49: jav-01620.
doi: 10.1111/jav.01620

Arct A, Martyka R, Drobniak SM, Oleś W, Dubiec A, Gustafsson L (2022) Effects of elevated nest
box temperature on incubation behaviour and offspring fitness-related traits in the Collared
Flycatcher *Ficedula albicollis*. J Ornithol 163: 263–272. doi: 10.1007/s10336-021-01944-3

Ardia DR, Pérez JH, Clotfelter ED (2010) Experimental cooling during incubation leads to reduced
innate immunity and body condition in nestling tree swallows. Proc R Soc B 277: 1881–
1888. doi: 10.1098/rspb.2009.2138

Bambini G, Schlicht E, Kempenaers B (2019) Patterns of female nest attendance and male feeding
throughout the incubation period in Blue Tits *Cyanistes caeruleus*. Ibis 161: 50–65. doi:
10.1111/ibi.12614

Basso A, Richner H (2015a) Predator-specific effects on incubation behaviour and offspring growth
in Great Tits. PLoS ONE 10: 1–17. doi: 10.1371/journal.pone.0121088

Basso A, Richner H (2015b) Effects of nest predation risk on female incubation behavior and
offspring growth in great tits. Behav Ecol Sociobiol 69: 977–989. doi: 10.1007/s00265-015-
1910-4

Bates D, Maechler M, Bolker BM, Walker S (2015) Fitting linear mixed-effects models using lme4.
J Stat Softw 67: 1–48. doi: 10.18637/jss.v067.i01

Baudains TP, Lloyd P (2007) Habituation and habitat changes can moderate the impacts of human
disturbance on shorebird breeding performance. Anim Conserv 10: 400–407. doi:
10.1111/j.1469-1795.2007.00126.x

Beale CM, Monaghan P (2004) Human disturbance: people as predation-free predators? J Appl
Ecology 41: 335–343. doi: 10.1111/j.0021-8901.2004.00900.x

Berntsen HH, Bech C (2016) Incubation temperature influences survival in a small passerine bird. J
Avian Biol 47: 141–145. doi: 10.1111/jav.00688

Blas J, Bortolotti GR, Tella JL, Baos R, Marchant TA (2007) Stress response during development
predicts fitness in a wild, long lived vertebrate. Proc Natl Acad Sci USA 104: 8880–8884.
doi: 10.1073/pnas.0700232104

Bueno-Enciso J, Barrientos R, Sanz JJ (2017) Incubation behaviour of Blue *Cyanistes caeruleus*
and Great Tits *Parus major* in a Mediterranean habitat. Acta Ornithol 52: 21–34. doi:
10.3161/00016454AO2017.52.1.003

Clutton-Brock TH (1991) *The Evolution of Parental Care*. Vol. 64. Princeton University Press.

Cole EF, Quinn JL (2014) Shy birds play it safe: personality in captivity predicts risk responsiveness
during reproduction in the wild. Biol Lett 10: 20140178. doi: 10.1098/rsbl.2014.0178

Conway CJ, Martin TE (2000a) Effects of ambient temperature on avian incubation behavior. Behav
Ecol 11: 178–188. doi: 10.1093/beheco/11.2.178

Conway CJ, Martin TE (2000b) Evolution of passerine incubation behavior: influence of food,
temperature, and nest predation. Evolution 54: 670–685. doi: 10.1554/0014-3820(2000)054

Cooper CB, Mills H (2005) New software for quantifying incubation behavior from time-series
recordings. J Field Ornithol 76: 352–356. doi: 10.1648/0273-8570-76.4.352

Cooper CB, Voss MA (2013) Avian incubation patterns reflect temporal changes in developing
clutches. PLoS ONE 8: e65521. doi: 10.1371/journal.pone.0065521

Coslovsky M, Richner H (2011) Predation risk affects offspring growth via maternal effects. Funct
Ecol 25: 878–888. doi: 10.1111/j.1365-2435.2011.01834.x

Cresswell W (2008) Non-lethal effects of predation in birds. Ibis 150: 3–17. doi: 10.1111/j.1474-
919X.2007.00793.x

Cresswell W, McCleery RH (2003) How Great Tits maintain synchronization of their hatch date with
food supply in response to long-term variability in temperature. J Anim Ecol 72: 356–366.
doi: 10.1046/j.1365-2656.2003.00701.x

Croston R, Hartman CA, Herzog MP, Peterson SH, Kohl JD, Overton CT, Feldheim CL, Casazza M.
L., Ackerman J. T. 2021. Interrupted incubation: How dabbling ducks respond when flushed
from the nest. Ecol Evol 11: 2862–2872. doi: 10.1002/ece3.7245

Diez-Méndez D, Cooper CB, Sanz JJ, Verdejo J, Barba E (2021a) Deconstructing incubation
behaviour in response to ambient temperature over different timescales. J Avian Biol
jav.02781. doi: 10.1111/jav.02781

Diez-Méndez D, Rodríguez S, Álvarez E, Barba E (2020) The role of partial incubation and egg
repositioning within the clutch in hatching asynchrony and subsequent effects on breeding
success. Ibis 162: 63–74. doi: 10.1111/ibi.12730

Diez-Méndez D, Sanz JJ, Barba E (2021b) Impacts of ambient temperature and clutch size on
incubation behaviour onset in a female-only incubator songbird. Ibis 163: 1056–1071. doi:
10.1111/ibi.12937

Drent R (1975) Incubation. In: Farmer D., King J. (eds). Avian Biology. Academic Press, pp. 333–
420.

Dubiec A (2011) Condition-dependent clutch desertion in Great Tit (*Parus major*) females subjected to human disturbance. J Ornithol 152: 743–749. doi: 10.1007/s10336-011-0647-8

DuRant SE, Hopkins WA, Hepp GR, Walters JR (2013) Ecological, evolutionary, and conservation implications of incubation temperature-dependent phenotypes in birds. Biol Rev 88: 499–509. doi: 10.1111/brv.12015

Dutour M, Lévy L, Lengagne T, Holveck MJ, Crochet PA, Perret P, Doutrelant C, Grégoire A (2020) Hissing like a snake: bird hisses are similar to snake hisses and prompt similar anxiety behavior in a mammalian model. Behav Ecol Sociobiol 74: 1. doi: 10.1007/s00265-019-2778-5

Eggers S, Griesser M, Nystrand M, Ekman J (2006) Predation risk induces changes in nest-site selection and clutch size in the Siberian jay. Proc. R. Soc. B. 273: 701–706. doi: 10.1098/rspb.2005.3373

Encabo SI, Monrós JS, Barba E (2001) Egg size variation in a mediterranean Great tit *Parus major* populations. Ardeola 48:63-70. https://www.ardeola.org/en/volumes/481/articles/63-70/

Fox J, Hong J (2009) Effect displays in R for multinomial and proportional-odds logit models: extensions to the effects package. J Stat Softw 32: 1–24. doi: 10.2307/271037

Frid A, Dill LM (2002) Human-caused disturbance stimuli as a form of predation Risk. Conserv Ecol 6: art11. doi: 10.5751/ES-00404-060111

Ghalambor CK, Martin TE (2002) Comparative manipulation of predation risk in incubating birds reveals variability in the plasticity of responses. Behav Ecol 13: 101–108. doi: 10.1093/beheco/13.1.101

Ghalambor CK, Peluc SI, Martin TE (2013) Plasticity of parental care under the risk of predation: how much should parents reduce care? Biol Lett 9: 20130154. doi: 10.1098/rsbl.2013.0154

Golawski A, Zduniak P (2022) Influence of researcher experience and fieldwork intensity on the probability of brood losses in sensitive species: The case of the Red-backed Shrike *Lanius collurio*. J Nat Conserv 69: 126249. doi: 10.1016/j.jnc.2022.126249

Götmark F (1992) The effects of investigator disturbance on nesting birds. In: Power D. M. (ed). Current Ornithology. Plenum Press, pp. 63–104.

Hepp GR, Kennamer RA, Johnson MH (2006) Maternal effects in Wood Ducks: incubation temperature influences incubation period and neonate phenotype. Funct Ecol 20: 308–314. doi: 10.1111/j.1365-2435.2006.01108.x

Hope SF, DuRant SE, Angelier F, Hallagan JJ, Moore IT, Parenteau C, Kennamer RA, Hopkins WA (2020) Prolactin is related to incubation constancy and egg temperature following a disturbance in a precocial bird. Gen Comp Endocrinol 295: 113489. doi: 10.1016/j.ygcen.2020.113489

Hu Q, Wen Y, Yu G, Yin J, Guan H, Lv L, Wang P, Xu J, Wang Y, Zhang Z, Li J (2020) Research activity does not affect nest predation rates of the Silver-throated Tit, a passerine bird building domed nests. Avian Res 11: 28. doi: 10.1186/s40657-020-00214-9

Hutfluss A, Dingemanse NJ (2019) Human recreation reduces clutch size in great tits *Parus major*
regardless of risk-taking personality. Behav Ecol 30: 1751–1760. doi:
10.1093/beheco/arz145

Ibáñez-Álamo JD, Sanllorente O, Soler M (2012) The impact of researcher disturbance on nest
predation rates: a meta-analysis. Ibis 154: 5–14. doi: 10.1111/j.1474-919X.2011.01186.x

Ibáñez-Álamo JD, Soler M (2010) Investigator activities reduce nest predation in blackbirds *Turdus*
merula. J Avian Biol 41: 208–212. doi: 10.1111/j.1600-048X.2009.04805.x

Ibáñez-Álamo JD, Soler M (2012) Predator-induced female behavior in the absence of male
incubation feeding: an experimental study. Behav Ecol Sociobiol 66: 1067–1073. doi:
10.1007/s00265-012-1357-9

Johnson LS, Napolillo FM, Kozlovsky DY, Hebert RM, Allen A (2013) Variation in incubation effort
during egg laying in the Mountain Bluebirds and its association with hatching asynchrony. J
Field Ornithol 84: 242–254. doi: 10.1111/jofo.12023

Johnston JC (2011) Effects of stage in incubation, time in season, and proportion of original clutch
remaining on nest desertion by house sparrows, *Passer domesticus*. Behav 148: 1121–1135.
doi: 10.1163/000579511X596589

Kania W (1989) Brood desertion by great tits *Parus major* caught at the nest. Acta Ornithol 25: 77–
105. doi: https://www.academia.edu/48562768

Kilgas P, Tilgar V, MM, Mänd R (2007) Physiological condition of incubating and brood rearing
female Great Tits *Parus major* in two Contrasting habitats. Acta Ornithol 42: 129–136. doi:
10.3161/068.042.0207

Kovařík P, Pavel V (2011) Does threat to the nest affect incubation rhythm in a small passerine?
Ethology 117: 181–187. doi: 10.1111/j.1439-0310.2010.01865.x

Krams I, Vrublevska J, Koosa K, Krama T, Mierauskas P, Rantala MJ, Tilgar V (2014) Hissing calls
improve survival in incubating female great tits (*Parus major*). Acta ethol 17: 83–88. doi:
10.1007/s10211-013-0163-3

Lenington S (1979) Predators and Blackbirds: The "Uncertainty Principle" in Field Biology. The Auk
96: 190–192. doi: 10.1093/auk/96.1.190

Lima SL (2009) Predators and the breeding bird: behavioral and reproductive flexibility under the
risk of predation. Biol Rev 84: 485–513. doi: 10.1111/j.1469-185X.2009.00085.x

Lima SL, Bednekoff PA (1999) Temporal variation in danger drives antipredator behavior: The
predation risk allocation hypothesis. Am Nat 153: 649–659. doi: 10.1086/303202

Lloyd P, Little RM, Crowe TM (2000) Investigator effects on the nesting success of arid-zone birds.
J Field Ornithol 71: 227–235. doi: 10.1648/0273-8570-71.2.227

Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D (2021) performance: An R Package
for Assessment, Comparison and Testing of Statistical Models. JOSS 6: 3139. doi:
10.21105/joss.03139

Macdonald EC, Camfield AF, Jankowski JE, Martin K (2013) Extended incubation recesses by
alpine-breeding Horned Larks: a strategy for dealing with inclement weather? J Field
Ornithol 84: 58–68. doi: 10.1111/jofo.12006

Martin TE, Briskie JV (2009) Predation on dependent offspring: A review of the consequences for
mean expression and phenotypic plasticity in avian life history traits. Ann NY Acad Sci 1168:
201–217. doi: 10.1111/j.1749-6632.2009.04577.x

Morosinotto C, Thomson RL, Korpimäki E (2013) Plasticity in incubation behaviour under
experimentally prolonged vulnerability to nest predation. Behav 150: 1767–1786. doi:
10.1163/1568539X-00003119

Naef-Daenzer B, Keller LF (1999) The foraging performance of great and blue tits (*Parus major* and
P. caeruleus) in relation to caterpillar development, and its consequences for nestling growth
and fledging weight. J Anim Ecol 68: 708–718. doi: 10.1046/j.1365-2656.1999.00318.x

Nord A, Nilsson J-Å (2011) Incubation temperature affects growth and energy metabolism in Blue
Tit nestlings. Am Nat 178: 639–651. doi: 10.1086/662172

Nord A, Nilsson J-Å (2012) Context-dependent costs of incubation in the Pied Flycatcher. Anim.
Behav. 84: 427–436. doi: 10.1016/j.anbehav.2012.05.017

Nord A, Williams JB (2015) The energetic costs of incubation. In: Deeming D. C., Reynolds S. J.
(eds). Nests, Eggs, and Incubation: new ideas about avian reproduction. Oxford University
Press, pp. 152–170.

Olson CR, Vleck CM, Vleck D (2006) Periodic cooling of bird eggs reduces embryonic growth
efficiency. Physiological and Biochemical Zoology 79: 927–936. doi: 10.1086/506003

Ospina EA, Merrill L, Benson TJ (2018) Incubation temperature impacts nestling growth and survival
in an open-cup nesting passerine. Ecol Evol 8: 3270–3279. doi: 10.1002/ece3.3911

Perona AM, Urios V, López-López P (2019) Holidays? Not for all. Eagles have larger home ranges
on holidays as a consequence of human disturbance. Biol Conserv 231: 59–66. doi:
10.1016/j.biocon.2019.01.010

Perrins C. M. 1979. *British tits*. HarperCollins.

R Core Team. 2023. R: A language and environment for statistical computing.

Remacha C, Delgado JA, Bulaic M, Pérez-Tris J (2016) Human disturbance during early life impairs
nestling growth in birds inhabiting a nature recreation area. PLoS ONE 11: e0166748. doi:
10.1371/journal.pone.0166748

Schaub R, Mumme RL, Woolfenden' GE (1992) Predation on the eggs and nestlings of Florida Scrub
Jays. The Auk 109: 585–593. doi: 10.1093/auk/109.3.585

Schöll EM, Aparisi MP, Hille SM (2019) Diurnal patterns of ambient temperature but not precipitation
influence incubation behavior in Great Tits. J Ornithol 161: 529–538. doi: 10.1007/s10336-
019-01737-9

Simmonds EG, Sheldon BC, Coulson T, Cole EF (2017) Incubation behavior adjustments, driven by
ambient temperature variation, improve synchrony between hatch dates and caterpillar peak
in a wild bird population. Ecol Evol 7: 9415–9425. doi: 10.1002/ece3.3446

Skutch AF (1957) The incubation patterns of birds. Ibis 99: 69–93. doi: 10.1111/j.1474-
919X.1957.tb01934.x

Stearns SC (1992) *The Evolution of Life Histories*. Oxford University Press.

Stien J, Ims RA (2016) Absence from the nest due to human disturbance induces higher nest
predation risk than natural recesses in Common Eiders *Somateria mollissima*. Ibis 158: 249–
260. doi: 10.1111/ibi.12338

Tkaczyk SP, Chivers DP, Wiebe KL (2023) Shifting incubation rhythms in response to predation risk
and the length of the response in mountain bluebirds. Behav. 1–19. doi: 10.1163/1568539X-
bja10250

Tryjanowski P, Kuźniak S (1999) Effect of research activity on the success of Red-backed Shrike
Lanius collurio nests. OrnisFenn76:41–43. https://ornisfennica.journal.fi/article/view/133497

Van Iersel R, Boiten G, Pinxten R, Eens M (2023) Untangling behaviours: independent expressions
of female–female aggression and snake-like hissing in the blue tit (*Cyanistes caeruleus*). Sci
Rep 13: 16346. doi: 10.1038/s41598-023-43652-3

Verboven N, Tinbergen JM (2002) Nest desertion: a trade-off between current and future
reproduction. Anim Behav 63: 951–958. doi: 10.1006/anbe.2001.1971

Verhulst S, Oosterbeek K, Ens BJ (2001) Experimental evidence for effects of human disturbance
on foraging and parental care in oystercatchers. Biol Conserv 101: 375–380. doi:
10.1016/S0006-3207(01)00084-2

Weidinger K (2008) Nest monitoring does not increase nest predation in open-nesting songbirds:
inference from continuous nest-survival data. Auk 125: 859–868. doi:
10.1525/auk.2008.07016

Wickham H (2016) *ggplot2: Elegant graphics for data analysis*. Springer-Verlag.

Zanette LY, White AF, Allen MC, Clinchy M (2011) Perceived predation risk reduces the number of
offspring songbirds produce per year. Science 334: 1398–1401. doi:
10.1126/science.1210908

Zieliński J (2020) Nest desertion by Blackcaps *Sylvia atricapilla* in response to experimental partial
clutch reduction. Ardea 108: 31–37. doi: 10.5253/arde.v108i1.a1

Zub K, Czeszczewik D, Ruczyński I, Kapusta A, Walankiewicz W (2017) Silence is not golden: the
hissing calls of tits affect the behaviour of a nest predator. Behav Ecol Sociobiol 71: 79. doi:
10.1007/s00265-017-2313-5

Table 1. Estimates of linear mixed-effect models analysing the effect of a disturbance event on the

ongoing off-bout (female absent in the nest-box) compared to the previous and next off-bouts, or

the immediate off-bout after a disturbance (female present and flushed out of the nest-box)

compared to the previous and the next off-bouts. Incubation day, ambient temperature, time of the

day, population and year were also considered. Nest-box is a random effect. Significant effects P

< 0.05 are highlighted in bold.

	Estimate	se	*t*	*P*
Off-bout (female absent)				
(R^2 fixed factors = 0.18, R^2 model = 0.23)				
Intercept	2.74	0.200	12.46	
Previous	**-0.27**	**0.098**	**-2.80**	**0.006**
Next	**-0.42**	**0.096**	**-4.40**	**<0.001**
Incubation day	**-0.21**	**0.059**	**-3.58**	**<0.001**
Temperature	0.87	0.806	1.07	0.284
Temperature^2	-0.49	0.736	-0.67	0.503
Time	0.27	0.803	0.33	0.739
Time^2	-0.71	0.857	-0.83	0.408
Date	0.01	0.130	0.05	0.963
Population: Sagunto	-0.24	0.244	-0.99	0.330
Year 2016	0.18	0.177	1.01	0.319
Year 2017	0.03	0.171	0.19	0.851
Off-bout (female present)				
(R^2 fixed factors = 0.20, R^2 model = 0.22)				
Intercept	2.75	0.207	13.32	
Previous	**-0.61**	**0.069**	**-8.84**	**<0.001**
Next	**-0.54**	**0.069**	**-7.84**	**<0.001**
Incubation day	**-0.11**	**0.039**	**-2.91**	**0.004**
Temperature	**2.72**	**0.856**	**3.18**	**0.002**
Temperature^2	0.52	0.718	0.73	0.466
Time	0.30	0.802	0.37	0.709
Time^2	0.13	0.807	0.16	0.872
Date	0.08	0.084	1.00	0.323
Population: Sagunto	-0.03	0.178	-1.17	0.867
Year 2016	0.24	0.152	1.59	0.117
Year 2017	0.12	0.149	0.83	0.409

560 **Table 2.** Estimates of a linear mixed-effect model analysing the effect of the duration of a

561 disturbance event (short vs. long disturbance) on the duration of the subsequent off-bout when a

562 female was present and flushed out of the nest-box. Incubation day, ambient temperature, time of

563 the day, population and year were also considered. Nest-box is a random effect. Significant effects

564 $P < 0.05$ are highlighted in bold.

	Estimate	se	t	P
Off-bout duration (female present) ($R^2_{\text{fixed factors}} = 0.16$, $R^2_{\text{model}} = 0.32$)				
Intercept	2.40	0.385	6.24	
Long disturbance	**0.37**	**0.138**	**2.69**	**0.008**
Incubation day	0.03	0.074	0.36	0.718
Temperature	**2.00**	**0.828**	**2.42**	**0.017**
Temperature^2	0.27	0.710	0.38	0.702
Time	0.69	0.755	0.92	0.359
Time^2	-0.26	0.811	-0.32	0.748
Date	0.20	0.161	1.21	0.234
Population: Sagunto	0.27	0.339	0.80	0.431
Year 2016	0.41	0.286	1.45	0.154
Year 2017	0.12	0.279	0.45	0.658

568   **Table 3.** Estimates of a generalized linear mixed-effect model analysing the effect of the number of

569   disturbance events on the daily nest attentiveness. Incubation day, daily ambient temperature, time

570   of the day, population and year were also considered. Nest-box is a random effect. Significant

571   effects P < 0.05 are highlighted in bold.

	Estimate	se	z	P
Daily nest attentiveness ($R^2_{\text{fixed factors}}$ = 0.33, R^2_{model} = 0.68)				
Intercept	1.03	0.234	4.40	
One event	**-0.25**	**0.047**	**-5.36**	**<0.001**
Two events	**-0.43**	**0.063**	**-6.81**	**<0.001**
Incubation day	**0.19**	**0.047**	**3.97**	**<0.001**
Temperature	**-0.91**	**0.340**	**-2.68**	**0.007**
Temperature^2	-0.31	0.318	-0.98	0.327
Time	-0.17	0.151	-1.10	0.273
Time^2	-0.02	0.316	-0.05	0.961
Date	0.15	0.129	1.20	0.229
Population: Sagunto	0.15	0.221	0.66	0.513
Year 2016	**1.03**	**0.234**	**4.40**	**<0.001**
Year 2017	**-0.25**	**0.047**	**-5.36**	**<0.001**

 Figure 1. Comparison of the duration of the ongoing/first bout after a disturbance event and the

 bout before the event and the next one. The incubating female may be present or absent during

 such events. Box-plots represent the median and 50% of the values, and upper and lower whiskers

 represent values > 75th and < 25th percentile. Semi-transparent circles represent the bout duration

 raw data. Full circles and associated bars show the model estimates and 95% confidence intervals.

Figure 2. Daily nest attentiveness (proportion of time a female spends incubating compared to the

time spent out of the nest on self-care related activities) depending on the number of disturbance

events in a day. Box-plots represent the median and 50% of the values, and upper and lower

whiskers represent values > 75th and < 25th percentile. Semi-transparent circles represent the bout

duration raw data. Full circles and associated bars show the model estimates and 95% confidence

intervals.