Full title of paper:

Human stem cell transplantation for Parkinson's disease: A systematic review of *in situ* survival and maturation of progenitors derived from human embryonic or induced stem cells in Parkinsonian models.

Authors:

Giulia Comini and Eilís Dowd

Authors' affiliations:

Pharmacology & Therapeutics and Galway Neuroscience Centre, University of Galway, Ireland.

Corresponding author:

Professor Eilís Dowd

eilis.dowd@universityofgalway.ie

Running title:

Survival and maturation of dopaminergic transplants

Keywords:

Parkinson's disease; Cell replacement therapy; Embryonic stem cells (ESCs); Induced pluripotent stem cells (iPSCs), Transplant; Graft; Survival; Differentiation.

Abstract

Stem cell-based brain repair is a promising emergent therapy for Parkinson's which is based on years of foundational research using human fetal donors as a cell source. Unlike current therapeutic options for patients, this approach has the potential to provide long-term stem cell-derived reconstruction and restoration of the dopaminergic input to denervated regions of the brain allowing for restoration of certain functions to patients. The ultimate clinical success of stem cell-derived brain repair will depend on both the safety and efficacy of the approach, and the latter is dependent on the ability of the transplanted cells to survive and differentiate into functional dopaminergic neurons in the Parkinsonian brain. Because the pre-clinical literature suggests that there is a considerable variability in survival and differentiation between studies, the aim of this systematic review was to assess these parameters in human stem-derived dopaminergic progenitor transplant studies in animal models of Parkinson's. To do so, a defined systematic search of the PubMed database was completed to identify relevant studies published up to March 2024. After screening, 76 articles were included in the analysis from which 178 separate transplant studies were identified. From these, graft survival could be assessed in 52 studies and differentiation in 129 studies. Overall, we found that graft survival ranged from <1% to 500% of cells transplanted, with a median of 51% of transplanted cells surviving in the brain; while dopaminergic differentiation of the cells ranged from 0% to 46% of cells transplanted with a median of 3%. This systematic review suggests that there is considerable scope for improvement in the differentiation of stem cell-derived dopaminergic progenitors in order to maximize the therapeutic potential of this approach for patients.

Introduction

The concept of human stem cell-derived brain repair for Parkinson's disease is built on decades of research using fetal cells as a source of healthy cells for transplantation [1]. The first studies in this field date back to the 1970s when groups in the US [2] and Sweden [3] pioneered this approach using rat-to-rat fetal transplants. These studies demonstrated that fetal dopamine neuron-rich tissue transplanted into the brain could survive and ameliorate motor dysfunction in dopamine-depleted rats. A decade or so of preclinical research followed leading to the first clinical trials of human fetal transplantation in patients with Parkinson's disease [4, 5].

Over the intervening decades, numerous preclinical studies and clinical trials provided proof-of-principle for the efficacy and safety of cellular brain repair, with fetal dopaminergic grafts demonstrating the ability to survive in the brain, reinnervate the dopamine-depleted striatum, and ameliorate dysfunction/symptoms in Parkinsonian models and patients [reviewed in 1, 6]. However, it also became apparent that this approach had serious limitations associated with the need for human fetal donors including supply, ethical concerns and logistical difficulties, all of which would ultimately preclude this translating to a therapy for patients.

In parallel with these efforts, embryonic stem cells (ECSs) were first cultured from mice [7, 8] and later humans [9], and protocols to convert these to a dopaminergic phenotype were later developed and refined [reviewed in 10]. This led to the first mouse-to-rat [11] and human-to-rat [12] studies of ESC-derived brain repair in Parkinsonian rats. With the discovery that somatic cells could be reprogrammed into induced pluripotent stem cells (iPSCs; [13-15], the era of stem cell-derived brain repair for Parkinson's disease was truly underway [16, 17]. In contrast to fetal transplantation, both ESCs and iPSCs offered the

possibility of a limitless supply of quality-controlled dopaminergic neurons for allogenic (donor-to-patient) transplantation, with the iPSCs offering the additional potential of autologous (patient-to-self) transplantation.

Over the last 20 years, numerous preclinical trials have been undertaken to determine the efficacy and safety of human stem cell-derived brain repair in Parkinsonian models, and the outcome of these has led to the initiation of 4 clinical trials that are currently underway (Kyoto Trial: UMIN000033564; BlueRock Trial: NCT04802733; STEM-PD: NCT05635409; S.Biomedics Trial: NCT05887466). Critically, the ultimate success of these and future clinical trials is highly dependent on the ability of the transplanted stem cell-derived dopaminergic progenitors to survive and differentiate into functional dopaminergic neurons in the Parkinsonian brain. Although *in situ* survival and differentiation of ESC and iPSC-derived dopaminergic progenitors is well established in rodent and primate models of Parkinson's disease, this is highly variable between, and even within, studies.

Therefore, because of the pivotal role that *in situ* survival and dopaminergic differentiation will play in the clinical success of stem cell-derived brain repair for Parkinson's, the aim of this review was to systematically assess these parameters in human stem-derived dopaminergic progenitor transplant studies in animal models.

Methods

Search Strategy

This study was completed in accordance with the PRISMA 2020 guidelines [18] and used a search strategy in the "PUBMED" database to find articles related to human stem-derived dopaminergic progenitor transplantation in animal models of Parkinson's. The specific search

string used was (stem AND (embryonic OR induced) AND Parkinson's AND dopaminergic AND (transplant or graft)). This search identified a total of 887 articles spanning 1997 to March 2024. These articles were then screened according to the search strategy outlined below and depicted in the PRISMA flow diagram (Fig. 1). This yielded 76 articles which were included in this systematic review. All the articles obtained were managed using Microsoft Excel and EndNote software.

Screening Strategy

The 887 articles identified were first screened by Title and Abstract and the following inclusion and exclusion criteria were applied. Inclusion criteria: 1) original research studies, 2) human ESC or iPSC-derived dopaminergic progenitors, 3) transplanted into the brain. Exclusion criteria: 1) review articles, 2) clinical trials, 3) not ESC or iPSC cells, 4) not human origin, 5) no transplantation, 6) not relevant for other miscellaneous reasons. This initial screen left 128 articles which were screened by full text. In this second screen, two additional exclusion criteria were used: 7) transplanted animals were not Parkinsonian, and 8) full text was not available. This left 72 articles for inclusion to which a further 4 were added that were identified from the cited publications in the selected articles.

Fig. 1 PRISMA flow chart. Flow chart based on PRISMA 2020 guidelines [18] detailing the screening strategy employed for the study selection in the present study.

Data Extraction

Variables manually extracted from the selected articles (Table 1 and references therein) included stem cell type (ECSs or iPSCs), cell line, day of differentiation at time of transplantation (for protocols using dual SMAD inhibition only), host species, Parkinsonian model, immunological status of the host (i.e. immunosuppressed vs. immunodeficient), week of sacrifice after transplantation, number of cells transplanted, number of cells surviving and number of cells differentiating into dopaminergic neurons. For cell survival, cell counts were only extracted where the marker used identified all transplanted cells (almost exclusively human nuclear markers) and not a specific subset of lineage restricted cells. For dopaminergic differentiation, cell counts extracted were based on tyrosine hydroxylase immunostaining or, in a limited number of studies, GFP expression driven by the dopaminergic *PITX3* promoter. In most articles, the cell numbers were provided directly by the authors, but in some articles, we estimated them from ranges and/or figures provided. Any estimated cell number is highlighted in Table 1 by underlining. Many articles employed transplant groups with various treatments, but for consistency and comparability between studies, only the unmanipulated/control cohorts are included here. Data extraction was completed independently by both authors, and cross-checked for accuracy.

Based on the numbers of cells transplanted, surviving and differentiating, three parameters were then calculated: 1) cell survival (as a percent of cells transplanted, 2) cell differentiation (as a percent of cells transplanted) and 3) cell differentiation (as a percent of cells surviving). The latter was not depicted in any graphical representations in the results section but is available in Table 1. In a small number of articles, these percentages were self-reported by the authors without any underpinning cell counts. These are included (in brackets) in Table 1 but are not included in any graphical depictions of the data.

We also determined which articles included behavioural evaluation of graft-derived motor recovery (Supplementary Table 1) and, if so, which behavioural test was used and whether or not there was functional recovery.

Limitations of this Review

This systematic review is intended to give an overview of survival and differentiation of human stem cell-derived dopaminergic progenitor transplantation in animal models of Parkinson's. However, there are numerous factors that can affect graft survival and differentiation, and it is beyond the scope of the review to consider all of these. These factors include, but are not limited to, quality of stem cell line used, dopaminergic differentiation protocol used, validation of differentiation protocol, stage of dopaminergic differentiation at time of transplantation, verification of dopaminergic progenitor status at time of transplantation, use of cryopreservation for storage, viability of cell suspension used for transplantation, host animals used (species, strain, sex, age), immunological status of host animals (immunosuppressed or immunodeficient; immunosuppressive drug and drug regime; immunodeficient strain etc.), model of Parkinson's induced, site of implantation, cell proliferation in the brain, time of sacrifice after transplantation, and the experience and expertise of the laboratories and individual researchers performing all of these tasks. Some, but not all, of these factors were recorded in Table 1. Another limitation is that many studies excluded animals from their analyses where there were no surviving grafts or no differentiated dopaminergic neurons. Therefore, the survival and differentiation data shown in the results section here may be an overrepresentation of the true figures.

Statistical analysis of data

General study characteristics are shown as pie charts. All data related to cell survival and/or differentiation is shown as scatter plots depicting individual studies with the median. Any data cited in the text of the results section represents the mean ± standard error of the mean (SEM) unless specified otherwise. Comparisons between 2 groups were completed using Student's *t*-test, comparisons between 3 or more groups on a single factor were completed using one-way ANOVA, and comparisons between groups on two factors were completed using two-way ANOVA. Where appropriate, *post-hoc* testing was applied, and the test used is specified in the corresponding figure. Simple linear regression was used to assess correlations in the data. Importantly, when considering the graphical depictions and the statistical analyses, it is important to bear in mind that there are many different variables (some of which are outlined above) that could underpin and affect the data shown.

Results

Extracted Data

From the 76 articles included in this review, 178 different transplant studies were extracted (Table 1).

Table 1. Studies included in this review. A total of 76 articles were included in this review from which 178 separate transplant studies were identified. The data extracted from these studies is depicted in this table. Any underlined value was estimated from a range or figure in the original article. Any number in brackets (~~like this~~) was self-reported by the authors without any underpinning cell counts, and these were omitted from the results section. NA indicates that the data is not available for various reasons including not done, not reported, not a pan human cellular marker, expressed per section or per unit area/volume. *Indicates that the data was extracted from Table S1 in Kirkeby et al., 2017 [19]. Abbreviations: 6OH: 6-hydroxydopamine; SYN: Synuclein; Cyclo: Cyclosporine; Tac: Tacrine.

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[12]	Ben-Hur	2004	hESC	HES-1	-	Rat-6OH	Cyclo	12	400,000	294,774	389	73.69	0.10	0.13
[20]	Park	2005	hESC	HSF-6	-	Rat-6OH	Cyclo	2	500,000	NA	0	NA	0.00	NA
[21]	Brederlau	2006	hESC	SA002.5	-	Rat-6OH	Cyclo	2	100,000	100,000	30	100.00	0.03	0.03
[21]	Brederlau	2006	hESC	SA002.5	-	Rat-6OH	Cyclo	13	100,000	500,000	30	500.00	0.03	0.01
[22]	Roy	2006	hESC	H1/H9	-	Rat-6OH	Cyclo	10	500,000	NA	NA	NA	NA	(22)
[23]	Martinat	2006	hESC	H9	-	Mouse-6OH	?	6	100,000	NA	NA	NA	NA	(45)
[24]	Sonntag	2007	hESC	H7	-	Rat-6OH	Cyclo	12	100,000	NA	160	NA	0.16	NA
[24]	Sonntag	2007	hESC	H9	-	Rat-6OH	Cyclo	12	100,000	NA	2	NA	0.00	NA
[25]	Iacovitti	2007	hESC	Various	-	Rat-6OH	?	2-3	350,000	NA	NA	NA	NA	NA
[26]	Ko	2007	hESC	HSF-6	-	Rat-6OH	Cyclo	8	1,500,000	301,327	201	20.09	0.01	0.07
[27]	Chiba	2008	hESC	BG01V	-	Rat-6OH	Cyclo	4	1,000,000	NA	100	NA	0.01	NA
[28]	Yang	2008	hESC	H9	-	Rat-6OH	Cyclo	1	200,000	25,919	53	12.96	0.03	0.20
[28]	Yang	2008	hESC	H9	-	Rat-6OH	Cyclo	4	200,000	237,184	361	118.59	0.18	0.15
[28]	Yang	2008	hESC	H9	-	Rat-6OH	Cyclo	20	200,000	570,559	1,273	285.28	0.64	0.22

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[29]	Geeta	2008	hESC	Relicell®	-	Rat-6OH	Cyclo	12-48	1,200,000	NA	NA	NA	NA	NA
[17]	Wernig	2008	(?)iPSC	O9	-	Rat-6OH	?	4	200,000	NA	15,250	NA	7.63	NA
[30]	Ko	2009	hESC	HSF-6	-	Rat-6OH	Cyclo	8	750,000	483,814	976	64.51	0.13	0.20
[16]	Hargus	2010	hiPSC	S1	-	Rat-6OH	Cyclo	16	400,000	NA	4,890	NA	1.22	NA
[31]	Cai	2010	hiPSC	IMR90	-	Rat-6OH	Cyclo	6	1,000,000	NA	NA	NA	NA	NA
[32]	Kriks	2011	hESC	H1/H9	D25	Mouse-6OH	SCID	18	150,000	NA	5,000	NA	3.33	NA
[32]	Kriks	2011	hESC	H1/H9	D25	Rat-6OH	Cyclo	20	250,000	240,000	15,000	96.00	6.00	6.25
[32]	Kriks	2011	hESC	H1/H9	D25	NHP-MPTP	Cyclo	4	7,500,000	NA	NA	NA	NA	NA
[33]	Cho	2011	hESC	H9	-	Rat-6OH	Cyclo	24	550,000	NA	NA	NA	NA	(2)
[34]	Kikuchi	2011	hiPSC	253G4	-	NHP-MPTP	Tac	24	2,400,000	NA	30,700	NA	1.28	NA
[34]	Kikuchi	2011	hiPSC	253G4	-	NHP-MPTP	Tac	24	2,400,000	NA	126,000	NA	5.25	NA
[35]	Rhee	2011	hiPSC	Pro-1	-	Rat-6OH	Cyclo	8	300,000	NA	26,882	NA	8.96	NA
[35]	Rhee	2011	hiPSC	Pro-1	-	Rat-6OH	Cyclo	8	300,000	NA	0	NA	0.00	NA
[35]	Rhee	2011	hiPSC	Pro-1	-	Rat-6OH	Cyclo	8	750,000	NA	54,418	NA	7.26	NA
[36]	Kirkeby	2012	hESC	H9	D16	Rat-6OH	Cyclo	6	300,000	NA	25,014	NA	8.34	(54)
[36]	*Kirkeby	2012	hESC	H9	D16	Rat-6OH	Cyclo	18	300,000	NA	59,628	NA	19.88	(0.6)
[37]	Doi	2012	hiPSC	KhES-1	-	NHP-MPTP	Tac	36	2,400,000	NA	NA	NA	NA	NA
[37]	Doi	2012	hiPSC	KhES-1	-	NHP-MPTP	Tac	48	2,400,000	NA	9,950	NA	NA	NA
[38]	Daadi	2012	hESC	H7/H9	-	NHP-MPTP	Cyclo	8	1,000,000	NA	NA	NA	NA	(10)

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[39]	Emborg	2013	hESC	H9	-	NHP-MPTP	Cyclo	12	2,750,000	NA	0	NA	0.00	NA
[40]	Kang	2014	hESC		-	Rat-6OH	Cyclo	16	400,000	NA	NA	NA	NA	NA
[41]	Wakeman	2014	hESC	H1	-	NHP-MPTP	Cyclo	6	5,375,000	NA	0	NA	0.00	NA
[42]	Grealish	2014	hESC	H9	D16	Rat-6OH	Cyclo	18	150,000	NA	986	NA	0.66	NA
[42]	*Grealish	2014	hESC	H9	D16	Rat-6OH	Nude	24	150,000	NA	13,890	NA	9.26	NA
[42]	*Grealish	2014	hESC	H9	D16	Rat-6OH	Nude	24	100,000	NA	4,007	NA	4.01	NA
[43]	Yang	2014	hESC	H9	-	Mouse-MPTP	?	4	150,000	NA	NA	NA	NA	NA
[43]	Yang	2014	hESC	H9	-	Mouse-6OH	?	4	200,000	NA	NA	NA	NA	NA
[44]	Doi	2014	hiPSC	404C2/836 B3	D28	Rat-6OH	?	16	400,000	NA	3,436	NA	0.86	NA
[44]	Doi	2014	hiPSC	404C2/836 B3	D28	Rat-6OH	?	16	400,000	NA	6,747	NA	1.69	(0.3)
[44]	Doi	2014	hiPSC	404C2/836 B3	D42	Rat-6OH	?	16	400,000	NA	1,800	NA	0.45	(18)
[44]	Doi	2014	hiPSC	404C2/836 B3	D42	Rat-6OH	?	16	400,000	NA	1,900	NA	0.48	NA
[45]	Steinbeck	2015	hESC	H9	D25	Mouse-6OH	NOD-SCID	16-24	200,000	NA	NA	NA	NA	NA
[46]	*Grealish	2015	hESC	H9	D16	Rat-6OH	Nude	24	300,000	NA	11,569	NA	3.86	NA
[46]	*Grealish	2015	hESC	H9	D16	Rat-6OH	Cyclo	6	200,000	NA	340	NA	0.17	NA
[46]	*Grealish	2015	hESC	H9	D16	Rat-6OH	Cyclo	18	200,000	NA	11,009	NA	5.50	NA
[47]	Han	2015	hiPSC		D14-21	Rat-6OH	Cyclo	16	500,000	NA	NA	NA	NA	NA
[48]	Chen	2016	hESC	WA09	D32	Mouse-6OH	SCID	24	200,000	NA	6,110	NA	3.06	NA

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[49]	Nishimura	2016	hiPSC	1039A1	D28	Rat-6OH	X-SCID	12-16	400,000	NA	NA	NA	NA	NA
[19]	Kirkeby	2017	hESC	RC17	D16	Rat-6OH	Nude	25	100,000	NA	3,716	NA	3.72	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	4	300,000	NA	925	NA	0.31	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	6	200,000	NA	388	NA	0.19	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	6	150,000	NA	910	NA	0.61	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	6	400,000	NA	36	NA	0.01	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	16	300,000	NA	3,979	NA	1.33	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	16	200,000	NA	1,936	NA	0.97	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	18	400,000	NA	30	NA	0.01	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	18	150,000	NA	10,581	NA	7.05	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	18	75,000	NA	3,363	NA	4.48	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	20	300,000	NA	15,600	NA	5.20	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Cyclo	21	300,000	NA	798	NA	0.27	NA
[19]	*Kirkeby	2017	hESC	H9	D16	Rat-6OH	Nude	24	100,000	NA	38	NA	0.04	NA
[50]	Niclis	2017	hESC	H9	D25	Rat-6OH	Nude	24	50,000	NA	NA	NA	NA	NA
[50]	Niclis	2017	hESC	H9	D25	Mouse-6OH	Nude	5	50,000	NA	NA	NA	NA	NA
[51]	Niclis	2017	hESC	H9	D19	Rat-6OH	Nude	4	100,000	NA	6,081	NA	6.08	NA
[51]	Niclis	2017	hESC	H9	D19	Rat-6OH	Nude	28	100,000	NA	5,268	NA	5.27	NA
[52]	Kikuchi	2017	hiPSC		-	Rat-6OH	Cyclo	16	400,000	NA	252	NA	0.06	NA

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[52]	Kikuchi	2017	hiPSC		-	Rat-6OH	Cyclo	16	400,000	NA	267	NA	0.07	NA
[52]	Kikuchi	2017	hiPSC		-	Rat-6OH	Cyclo	16	400,000	NA	227	NA	0.06	NA
[52]	Kikuchi	2017	hiPSC		-	Mouse-SYN	Cyclo	24	200,000	NA	NA	NA	NA	NA
[53]	Kikuchi	2017	hiPSC		-	NHP-MPTP	Tac	32-96	2,400,000	NA	64,000	NA	2.67	NA
[54]	Wakeman	2017	hiPSC		D33/D38	Rat-6OH	Cyclo	24	450,000	101,580	26,081	22.57	5.80	25.68
[54]	Wakeman	2017	hiPSC		D33/D38	NHP-MPTP	Tac	12	3,750,000	NA	NA	NA	NA	NA
[55]	Cardoso	2018	hESC	H9/RC17	D16	Rat-6OH	Cyclo	6-18	75,000	NA	NA	NA	NA	NA
[55]	Cardoso	2018	hESC	H9/RC17	D16	Rat-6OH	Nude	24	75,000	NA	NA	NA	NA	NA
[56]	Wang	2018	hESC	H9	-	NHP-MPTP	Cyclo	36	2,000,000	NA	40,000	NA	2.00	NA
[57]	Adler	2019	hESC	RC17	D16	Rat-6OH	Cyclo	16	112,500	NA	NA	NA	NA	NA
[57]	Adler	2019	hESC	RC17	D16	Rat-6OH	Cyclo	7	300,000	NA	NA	NA	NA	NA
[57]	Adler	2019	hESC	RC17	D16	Rat-6OH	Nude	24	112,500	NA	NA	NA	NA	NA
[58]	Yu	2019	hiPSC		D15	Rat-6OH	Cyclo	5	15,000,000	NA	NA	NA	NA	NA
[59]	Zygogianni	2019	hiPSC		-	Mouse-6OH	NOD-SCID	12	200,000	NA	NA	NA	NA	(1)
[60]	Gantner	2020	hESC	H9	D20	Rat-6OH	Nude	24	100,000	NA	4,092	NA	4.09	NA
[60]	Gantner	2020	hESC	H9	D20	Rat-6OH	Nude	26	100,000	NA	19,810	NA	19.81	(10)
[60]	Gantner	2020	hESC	H9	D20	Mouse-6OH	Nude	2	100,000	NA	NA	NA	NA	NA
[61]	Goggi	2020	hESC	HES-3	D25	Rat-6OH	Nude	24	400,000	350,000	NA	87.50	NA	(10)

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[62]	Hoban	2020	hESC	RC17	D16	Rat-SYN	Cyclo	6-18	300,000	NA	NA	NA	NA	NA
[62]	Hoban	2020	hESC	RC17	D16	Rat-6OH	Cyclo	6-18	300,000	NA	NA	NA	NA	NA
[63]	Tiklova	2020	hESC	H9/RC17	D16	Rat-6OH	Nude	24	300,000	NA	NA	NA	NA	NA
[63]	Tiklova	2020	hESC	H9/RC17	D16	Rat-6OH	Nude	36	150,000	NA	NA	NA	NA	NA
[64]	Doi	2020	hiPSC	MCB003	-	Rat-6OH	Nude	20	400,000	NA	2,835	NA	0.71	NA
[64]	Doi	2020	hiPSC	MCB003	-	NHP-MPTP	Cyclo	24	1,750,000	NA	NA	NA	NA	NA
[65]	Song	2020	hiPSC	C4	D28	Rat-6OH	Nude	16	100,000	NA	5,621	NA	5.62	NA
[65]	Song	2020	hiPSC	C4	D28	Rat-6OH	Nude	16	300,000	NA	16,863	NA	5.62	NA
[65]	Song	2020	hiPSC	C4	D28	Rat-6OH	Nude	26	100,000	NA	34,560	NA	34.56	NA
[65]	Song	2020	hiPSC	C4	D28	Rat-6OH	Nude	26	100,000	NA	46,094	NA	46.09	NA
[66]	Hiller	2021	hiPSC	iCell® Dopa	D38	Rat-6OH	Nude	36	450,000	60,226	19,861	13.38	4.41	32.98
[66]	Hiller	2021	hiPSC	iCell® Dopa	D33	Rat-6OH	Nude	36	450,000	70,511	12,612	15.67	2.80	17.89
[66]	Hiller	2021	hiPSC	iCell® Dopa	D33	Rat-6OH	Nude	36	450,000	30,479	7,774	6.77	1.73	25.51
[66]	Hiller	2021	hiPSC	iCell® Dopa	D33	Rat-6OH	Nude	36	450,000	25,605	6,356	5.69	1.41	24.82
[66]	Hiller	2021	hiPSC	iCell® Dopa	D26	Rat-6OH	Nude	36	450,000	210,929	17,430	46.87	3.87	8.26
[67]	Yoo	2021	hESC	H9	-	Rat-6OH	Cyclo	16	350,000	NA	NA	NA	NA	NA
[68]	Piao	2021	hESC	H9	D16	Rat-6OH	Nude	36	400,000	495,000	45,865	123.75	11.47	9.27
[69]	de Luzy	2021	hESC	H1	D19	Rat-6OH	Nude	5	125,000	NA	4,752	NA	3.80	NA

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[69]	de Luzy	2021	hESC	H1	D19	Rat-6OH	Nude	10	125,000	NA	4,635	NA	3.71	NA
[69]	de Luzy	2021	hESC	H1	D19	Rat-6OH	Nude	24	125,000	NA	5,204	NA	4.16	NA
[69]	de Luzy	2021	hESC	H1	D19	Rat-6OH	Nude	24	125,000	577,226	6,061	461.78	4.85	1.05
[70]	Xiong	2021	hESC	H9	D32	Mouse-6OH	SCID	24	50,000	NA	NA	NA	NA	(68)
[71]	Shrigley	2021	hESC	RC17	D16	Rat-6OH	Nude	7	300,000	NA	2,100	NA	0.70	NA
[71]	Shrigley	2021	hiPSC	AST18	D16	Rat-6OH	Nude	7	300,000	NA	2,300	NA	0.77	NA
[71]	Shrigley	2021	hESC	RC17	D16	Rat-6OH	Nude	24	300,000	NA	5,000	NA	1.67	NA
[71]	Shrigley	2021	hiPSC	AST18	D16	Rat-6OH	Nude	24	300,000	NA	4,500	NA	1.50	NA
[71]	Shrigley	2021	hESC	RC17	D16	Rat-6OH	Cyclo	8	300,000	NA	2,100	NA	0.70	NA
[71]	Shrigley	2021	hiPSC	AST18	D16	Rat-6OH	Cyclo	8	300,000	NA	2,300	NA	0.77	NA
[72]	Elabi	2022	hESC	RC17	D16	Rat-6OH	Cyclo	16	300,000	2,000	500	0.67	0.17	25.00
[72]	Elabi	2022	hESC	RC17	D16	Rat-6OH	Cyclo	16	500,000	75,000	12,000	15.00	2.40	16.00
[73]	Lane	2022	hESC	H9	D16	Rat-6OH	Tolerised	52	240,000	4,600	2,000	1.92	0.83	43.48
[74]	Aldrin-Kirk	2022	hESC	RC17	D16	Rat-6OH	Nude	26	150,000	NA	NA	NA	NA	NA
[74]	Aldrin-Kirk	2022	hESC	RC17	D16	Rat-6OH	Nude	26	75,000	NA	NA	NA	NA	NA
[75]	Hiller	2022	hiPSC		D17	Rat-6OH	Nude	24	450,000	304,303	79,061	67.62	17.57	25.98
[75]	Hiller	2022	hiPSC		D24	Rat-6OH	Nude	24	450,000	266,956	67,830	59.32	15.07	25.41
[75]	Hiller	2022	hiPSC		D37	Rat-6OH	Nude	24	450,000	52,623	9,318	11.69	2.07	17.71
[75]	Hiller	2022	hiPSC		D38	Rat-6OH	Nude	24	450,000	108,093	20,355	24.02	4.52	18.83

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[75]	Hiller	2022	hiPSC		D17	Rat-6OH	Nude	24	7,500	4,604	1,087	61.39	14.49	23.61
[75]	Hiller	2022	hiPSC		D17	Rat-6OH	Nude	24	30,000	36,848	6,400	122.83	21.33	17.37
[75]	Hiller	2022	hiPSC		D17	Rat-6OH	Nude	24	120,000	214,898	19,973	179.08	16.64	9.29
[75]	Hiller	2022	hiPSC		D17	Rat-6OH	Nude	24	450,000	611,588	59,929	135.91	13.32	9.80
[76]	de Luzy	2022	hESC	H9	D13	Mouse-6OH	Nude	24	100,000	177,929	11,156	177.93	11.16	6.27
[76]	de Luzy	2022	hESC	H9	D19	Mouse-6OH	Nude	24	100,000	255,364	10,000	255.36	10.00	3.92
[76]	de Luzy	2022	hESC	H9	D22	Mouse-6OH	Nude	24	100,000	261,899	7,282	261.90	7.28	2.78
[76]	de Luzy	2022	hESC	H9	D25	Mouse-6OH	Nude	24	100,000	255,000	3,000	255.00	3.00	1.18
[76]	de Luzy	2022	hESC	H9	D30	Mouse-6OH	Nude	24	100,000	110,000	679	110.00	0.68	0.62
[76]	de Luzy	2022	hiPSC	RM3.5	D13	Mouse-6OH	Nude	24	100,000	29,023	11,904	29.02	11.90	41.02
[76]	de Luzy	2022	hiPSC	RM3.5	D19	Mouse-6OH	Nude	24	100,000	30,605	7,997	30.61	8.00	26.13
[76]	de Luzy	2022	hiPSC	RM3.5	D22	Mouse-6OH	Nude	24	100,000	141,525	23,050	141.53	23.05	16.29
[76]	de Luzy	2022	hiPSC	RM3.5	D25	Mouse-6OH	Nude	24	100,000	110,000	15,000	110.00	15.00	13.64
[76]	de Luzy	2022	hiPSC	RM3.5	D30	Mouse-6OH	Nude	24	100,000	29,500	3,240	29.50	3.24	10.98
[76]	de Luzy	2022	hiPSC	RM3.5	D13	Mouse-6OH	Nude	24	100,000	NA	9,000	NA	9.00	NA
[76]	de Luzy	2022	hiPSC	RM3.5	D19	Mouse-6OH	Nude	24	100,000	NA	8,000	NA	8.00	NA
[76]	de Luzy	2022	hiPSC	RM3.5	D22	Mouse-6OH	Nude	24	100,000	NA	15,000	NA	15.00	NA
[76]	de Luzy	2022	hiPSC	RM3.5	D13	Mouse-6OH	Nude	24	100,000	NA	8,000	NA	8.00	NA
[76]	de Luzy	2022	hiPSC	RM3.5	D19	Mouse-6OH	Nude	24	100,000	NA	6,000	NA	6.00	NA

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[76]	de Luzy	2022	hiPSC	RM3.5	D22	Mouse-6OH	Nude	24	100,000	NA	12,000	NA	12.00	NA
[76]	de Luzy	2022	hiPSC	RM3.5	D13	Rat-6OH	Nude	24	100,000	NA	NA	NA	NA	NA
[76]	de Luzy	2022	hiPSC	RM3.5	D19	Rat-6OH	Nude	24	100,000	NA	NA	NA	NA	NA
[76]	de Luzy	2022	hiPSC	RM3.5	D22	Rat-6OH	Nude	24	100,000	NA	NA	NA	NA	NA
[77]	Moriarty	2022	hiPSC	RM3.5	D19	Mouse-6OH	Nude	24	100,000	NA	12,435	NA	12.44	NA
[78]	Brot	2022	hiPSC		-	Mouse-6OH	RAG2 KO	4	100,000	28,817	1,000	28.82	1.00	3.47
[78]	Brot	2022	hiPSC		-	Mouse-6OH	RAG2 KO	24	100,000	36,365	3,627	36.37	3.63	9.97
[78]	Brot	2022	hiPSC		-	Mouse-6OH	RAG2 KO	32	100,000	43,994	5,079	43.99	5.08	11.54
[78]	Brot	2022	hiPSC		-	Mouse-6OH	RAG2 KO	48	100,000	46,767	7,689	46.77	7.69	16.44
[79]	Kirkeby	2023	hESC	RC17	D16	Rat-6OH	Nude	24	30,000	NA	1,848	NA	6.16	NA
[79]	Kirkeby	2023	hESC	RC17	D16	Rat-6OH	Nude	24	60,000	NA	3,644	NA	6.07	NA
[79]	Kirkeby	2023	hESC	RC17	D16	Rat-6OH	Nude	24	120,000	NA	8,464	NA	7.05	NA
[79]	Kirkeby	2023	hESC	RC17	D16	Rat-6OH	Nude	24	300,000	NA	8,504	NA	2.83	NA
[79]	Kirkeby	2023	hESC	RC17	D16	Rat-6OH	Nude	26	300,000	NA	7,000	NA	2.33	NA
[79]	Kirkeby	2023	hESC	RC17	D16	Pig-6OH	Tac	44	1,250,000	NA	NA	NA	NA	NA
[80]	Maimaitili	2023	hESC	H9	D16	Rat-6OH	Nude	27	125,000	NA	5,312	NA	4.25	(14)
[81]	Naderi	2023	hESC	hE7	D16	Rat-6OH	Cyclo	24	450,000	NA	NA	NA	NA	(92)
[82]	López-Ornelas	2023	hESC	H9	D22	NHP-MPTP	Cyclo	40	8,000,000	NA	225,000	NA	2.81	NA

Reference	1st Author	Year	Cells	Line	Differentiation (Day)	Host	Immunological status	Sacrifice (Week)	Transplanted (# cells)	Survived (# cells)	Differentiated (# cells)	Survived vs. transplanted (%)	Differentiated vs Transplanted (%)	Differentiated vs. Survived (%)
[83]	Rajova	2023	hESC	RC17	D16	Rat-6OH	Nude	24-60	187,500	NA	NA	NA	NA	NA
[84]	Nakamura	2023	hiPSC	201B7	-	Mouse-6OH	SCID	16	200,000	NA	NA	NA	NA	(14)
[85]	Ramos-Acevedo	2023	hESC	H9	-	Rat-6OH	Cyclo	1-4	60,000	NA	NA	NA	NA	NA
[86]	Hills	2023	hiPSC	410	D18	Rat-6OH	Cyclo	24	500,000	80,000	5,000	16.00	1.00	6.25
[86]	Hills	2023	hiPSC	410	D25	Rat-6OH	Cyclo	24	500,000	50,000	2,000	10.00	0.40	4.00
[86]	Hills	2023	hiPSC	411	D18	Rat-6OH	Cyclo	24	500,000	15,000	2,000	3.00	0.40	13.33
[86]	Hills	2023	hiPSC	411	D25	Rat-6OH	Cyclo	24	500,000	15,000	1,500	3.00	0.30	10.00
[87]	Lyu	2023	hiPSC		D14	Rat-6OH	?	16	200,000	NA	NA	NA	NA	NA
[88]	Daadi	2024	hiPSC		-	NHP-MPTP	Tac	24	4,000,000	NA	NA	NA	NA	NA
[89]	Fu	2024	hiPSC		-	Mouse-6OH	?	16	50,000	NA	NA	NA	NA	(84)
[90]	Park	2024	hESC	SNU-hES32	-	Rat-6OH	Cyclo	24	5,000	2,718	759	54.36	15.18	27.92
[90]	Park	2024	hESC	SNU-hES32	-	Rat-6OH	Cyclo	24	10,000	7,805	1,478	78.05	14.78	18.94
[90]	Park	2024	hESC	SNU-hES32	-	Rat-6OH	Cyclo	24	25,000	8,294	2,820	33.18	11.28	34.00
[90]	Park	2024	hESC	SNU-hES32	-	Rat-6OH	Cyclo	24	100,000	126,646	17,598	126.65	17.60	13.90
[90]	Park	2024	hESC	SNU-hES32	-	Rat-6OH	Cyclo	24	350,000	NA	NA	NA	NA	NA
[91]	Comini	2024	hiPSC	NAS2	D16	Rat-6OH	Nude	26	300,000	3,330	153	1.11	0.05	4.59

Study Characteristics

Of the 178 different transplant studies included in this review, 103 used human ESCs for transplantation while 75 used human iPSCs (Fig 2A). The vast majority of transplants were into 6-hydroxydopamine-lesioned rodents (128 rat studies and 32 mouse studies), with an additional 14 studies transplanting into MPTP-lesioned non-human primates and 4 into other models (1 rat α-synuclein model, 1 mouse α-synuclein model, 1 mouse MPTP model and 1 minipig 6-hydroxydopamine model) (Fig. 2B). An approximately equal number of studies used immunosuppressed host animals (81 studies) and immunodeficient host animals (85 studies), while 1 study used immune-tolerised hosts and the remainder (11 studies) did not specify (Fig. 2C).

A)

B)

C)

Fig. 2 Study characteristics. Within the 76 articles included, 178 separate transplant studies were identified. The pie charts above show the proportions of these studies using A) different stem cells, B) host species and Parkinsonian models, and 3) the immunological status of the hosts.

Studies reporting transplanted cell survival and/or differentiation

Of the 178 transplant studies, the number of cells surviving the transplant procedure was directly reported (or calculable from the ranges/figures provided) in 52 studies, while the number of cells differentiating into dopaminergic neurons was directly reported (or calculable) in 129 studies (Fig. 3). Since the number of cells transplanted was reported (or calculable) in all studies, this meant that the cell survival (as a percent of cells transplanted) could be calculated in 52 studies, cell differentiation (as a percent of cells transplanted) could be calculated in 129 studies.

Fig. 3 Studies reporting transplanted cell survival and/or differentiation. Within the 76 articles included, 178 separate transplant studies were identified. The pie charts above show the proportions of these studies reporting A & B) transplanted cell survival and A & C) transplanted cell dopaminergic differentiation.

Overview of cell survival and differentiation

The overall survival of the cells transplanted in shown in Fig. 4A. This ranged from <1% to 500% of cells transplanted, with a median of 51% of transplanted cells surviving in the brain. Of note, 16 studies reported more cells surviving than were transplanted (i.e. >100%) indicating some level of cell proliferation after transplantation. The overall dopaminergic differentiation of the cells ranged from 0% to 46% of cells transplanted with a median of 3% (Fig. 4B). Interestingly, the dopaminergic differentiation was higher in proliferative grafts (with survival >100%) rather than non-proliferative grafts (with survival <100%) (Fig. 4C: Survival unknown: 4.7±0.8%; Survival <100%: 4.8±0.9%; Survival >100%: 9.8±1.9%; Cohort, $F_{(2,126)} = 3.76$, $P<0.05$).

Fig. 4 Overview of cell survival and differentiation. Of the 178 transplant studies extracted, 52 studies reported the number of cells surviving and 129 studies reported the number of dopaminergic neurons in the grafts. Thus, the scatter plots represent A) cell survival (as a percent of cells transplanted), B) dopaminergic differentiation (as a percent of cells transplanted) and C) dopaminergic differentiation (by survival cohort). Each point represents a separate transplant study, and the line indicates the median. Data were analysed by one-way ANOVA (C) with *post-hoc* Tukey's and *P<0.05.

Impact of cell type, host species and host immunological status on graft differentiation

Since *in situ* dopaminergic differentiation of transplanted stem cell-derived progenitors is a critical factor for effective brain repair, we next investigated whether some of the main experimental parameters affected this outcome. With regard to stem cell types used, there was a small but significantly higher dopaminergic differentiation in grafts derived from iPSCs compared with those derived from ESCs (Fig. 5A: ESCs: 4.1±0.6%; iPSCs: 6.9±1.1%; Cell type, $t_{(127)} = 2.33$, $P<0.05$). With regard to the host species, grafts in mice tended to have greater dopaminergic differentiation than those in rats or non-human primates (Fig. 5B: Rat: 5.0±0.8%; Mouse: 8.2±1.1%; NHP: 1.4±0.4%; Species, $F_{(2,126)} = 3.33$, $P<0.05$). However, this was only statistically significant on *post-hoc* analysis between mice and primates ($P<0.05$). In terms of the immunological status of the host animals, grafts in immunodeficient animals were significantly larger than those in immunosuppressed animals (Fig. 5C: Supressed: 2.8±0.6%; Deficient: 7.8±1.0%; Immunological status, $t_{(121)} = 4.12$, $P<0.0001$).

Fig. 5 Impact of cell type, host species and host immunological status on graft differentiation. Of the 178 transplant studies extracted, 129 studies reported the number of dopaminergic neurons in the grafts. Thus, the scatter plots represent dopaminergic differentiation) by A) cell type, B) host species and C) host immunological status. Each point represents a separate transplant study, and the line indicates the median. Data were analysed by *t*-test (A and C) or one-way ANOVA with *post-hoc* Tukey's (B) and *$P<0.05$, ****$P<0.0001$.

Further stratification of differentiation by cell type, host species and host immunological status

Since the data above suggested that stem cell type, host species and immunological status could affect dopaminergic differentiation, we next sought to further stratify the data to investigate this more. Since dopaminergic differentiation in non-human primates was very consistent and all animals were immunosuppressed, these were not considered in this section of the analysis.

We first compared ESC and iPSC-derived dopaminergic differentiation in immunosuppressed and immunodeficient rats (Fig. 6A). We found that there was no significant effect of cell type on differentiation (Cell type, $F_{(1,88)} = 1.23$, $P>0.05$), but there was a significant effect of immunological status with greater differentiation in immunodeficient animals (Immunological status, $F_{(1,88)} = 9.80$, $P<0.01$) and a significant interaction between the two (Cell type x Immunological status, $F_{(1,88)} = 4.04$, $P<0.05$). Subsequent *post-hoc* analysis confirmed that dopaminergic differentiation was greatest in iPSC-derived transplants in immunodeficient rats. We were unable to do a similar comparison for mice as all studies where differentiation was reported used immunodeficient mice. Rather, we next compared ESC and iPSC-derived dopaminergic differentiation between immunodeficient rats and mice (Fig. 6B). Interestingly, when all rodents were immunodeficient, we found that there was no significant effect of cell type or species on differentiation, and no interaction between the two.

Fig. 6 Further stratification of differentiation data in rodents. Of the 178 transplant studies extracted, 121 studies reported the number of dopaminergic neurons in the grafts in rats or mice. Thus, the scatter plots represent dopaminergic differentiation by A) cell type and immunological status in rats, and B) cell type in immunodeficient rats and mice. Each point represents a separate transplant study, and the line indicates the median. Data were analysed two-way ANOVA with *post-hoc* Newman Keuls and *P<0.05, **P<0.01.

Assessment of graft functionality through behavioural testing

The most clinically relevant outcome of effective engraftment in Parkinsonian animals is amelioration of motor dysfunction. Therefore, we evaluated whether or not behavioural testing was included in the selected articles and studies therein (Supplementary Table 1). Of the 178 different transplant studies, 114 included at least one motor function test, while 64 did not do any behavioural testing (Fig. 7A). The most commonly-used behavioural test was amphetamine-induced rotation (used in 95 studies) followed by the cylinder test (used in 22 studies). In the studies using amphetamine-induced rotation, recovery was reported in 81% (Fig. 7B) and, not surprisingly, these had a significantly greater number of cells differentiating compared with the 19% of studies where amelioration of rotational asymmetry did not occur (Fig. 7C; Recovery, $F_{(1,79)} = 2.35$, P<0.05).

Fig. 7 Assessment of graft functionality through behavioural testing. Of the 178 transplant studies extracted, 114 studies included behavioural testing (A). The most widely-used test was amphetamine-induced rotation with recovery in 81% of studies (B). The scatter plot represents dopaminergic differentiation in studies where amelioration of rotational asymmetry did, or did not, occur. Note that the lowest number of cells in the Yes column was 160. Each point represents a separate transplant study, and the line indicates the median. Data were analysed by *t*-test (C) and *$P<0.05$.

Impact of other factors on graft survival and/or differentiation.

To visualise some other trends in the data, we plotted graft survival and differentiation as a function of day of dopaminergic differentiation at the time of transplantation (for protocols using dual SMAD inhibition only; Fig. S1), week of sacrifice after transplantation (Fig. S2) and year of publication (Fig. S3). As expected, these show that most studies transplanted cells between day 16-19 of dopaminergic differentiation and sacrificed the animals around 6 months after transplantation. Interestingly, there was a significant positive correlation

between year of publication and dopaminergic differentiation *in situ* in the brain (Fig. 3B; Regression, $F_{(1,127)}$ = 14.50, R=0.32, $P<0.001$), suggesting that differentiation has improved overall in the two decades since the first publication. However, scrutiny of Table 1 indicates that this is driven by a high differentiation rate in just two articles each with several individual studies [75, 76]. When these were omitted from the analysis there was no correlation between year of publication and differentiation. We also plotted graft survival and differentiation as a function of the number of cells transplanted (for rodent studies only, Fig. S4). Interestingly, this showed a significant negative correlation between the number of cells transplanted and the final percentage of cells surviving (Fig. S4Aii; Regression, $F_{(1,49)}$ = 8.37, R=0.38, $P<0.01$) and differentiating (Fig. S4Bii; Regression, $F_{(1,117)}$ = 15.39, R=0.34, $P<0.001$), suggesting that larger transplants have an overall poorer engraftment outcome. Finally, we plotted graft differentiation as a function of graft survival and found a significant positive correlation (Fig. S5; Regression, $F_{(1,49)}$ = 7.65, R=0.37, $P<0.01$), suggesting that if grafts survived well, they also tended to differentiate well.

Discussion

Stem cell-derived brain repair for Parkinson's offers the possibility of life-long reconstruction of the degenerated dopaminergic input to the striatum after a single surgical transplant [6]. Relative to fetal tissue transplants, stem cells can provide unlimited numbers of the required phenotype of dopaminergic neurons for allogenic transplantation, with the iPSCs offering the additional potential of autologous transplantation. The clinical success of ESC and/or iPSC-derived brain repair will ultimately depend on both the safety and efficacy of the approach, and the field is awaiting the outcome of the iPSC trial (Kyoto Trial: UMIN000033564) and the three ESC trials (BlueRock Trial: NCT04802733; STEM-PD: NCT05635409; S.Biomedics Trial: NCT05887466) in this regard. However, even though stem cell-derived brain repair has already entered clinical trial, we observed that the *in situ* survival and

dopaminergic differentiation reported in preclinical studies was highly variable. Given the importance of these parameters in determining the efficacy of brain repair, we sought to complete this systematic review to shed light on the current status of these in human ESC or iPSC studies in Parkinsonian models.

With regard to survival of the stem cell-derived progenitors in the brain after transplantation, we were able to determine this in about a third of studies. In these studies, it was highly variable (ranging from <1% to 500% of cells transplanted) but relatively high overall (median of 51%). Several studies reported a greater number of cells surviving than were transplanted (reflective of *in situ* proliferation), and this was not confined to historical studies or to transplants of early-stage progenitors. Given the tumorigenicity risks posed by proliferative cells in grafts, it will be important to continue refining pre-transplant screening to reduce the number of such cells transplanted into the brain, but also to consider post-transplant failsafes such as the use of stem cell lines expressing suicide genes [69, 92].

The ability of transplanted dopaminergic progenitors to differentiate and mature into dopaminergic neurons after transplantation into the Parkinsonian brain underpins any functional recovery that will be afforded by brain repair. This was reported by a large proportion of the studies identified and was also found to be highly variable (ranging from 0% to 46% of cells transplanted), but unlike the high survival rate, differentiation was low overall (median of 3%). Interestingly, dopaminergic differentiation was higher in studies with proliferative grafts (survival >100%) which contrasts with suggestions that the dopaminergic yield may be lower in proliferative grafts due to an overabundance of unspecified cells [69]. Numerous factors can affect the post-transplant differentiation of dopaminergic progenitors (as outlined in the Methods section) and it was not possible to consider all of these

parameters here. Nevertheless, we did stratify the differentiation data on some of these factors and found some interesting trends.

The initial stratification suggested that cell line, host species and host immunological status were all factors that affected dopaminergic transplantation, and subsequent analysis (in rodents only) suggested that transplanting progenitors into immunosuppressed animals negatively impaired their ability to differentiate into mature neurons compared with transplanting them into immunodeficient animals. If this were to translate to human studies, it might suggest that differentiation, and therefore clinical efficacy, may be impaired in donor-to-patient allogenic transplantation which requires the use of long-term immunosupression. Conversely, the higher differentiation rate in immunodeficient animals – the vast majority of which were athymic T-cell deficient nude rats or mice – might suggest that autologous transplantation, which will not initiate any T-cell response, may have a better clinical outcome.

Another interesting, and somewhat surprising, aspect of the data was the trend for both survival and dopaminergic differentiation to be negatively correlated with the number of cells transplanted. Overall, transplanting high numbers of cells led to a lesser proportion surviving and differentiating in the brain. This could reflect cellular competition for trophic support, micronutrients, oxygen and synaptic connections with consequent cell death leading to activation of the innate immune system which is known to impair graft differentiation [93]. In addition to all transplants being into immunosuppressed animals, this could also partly explain the relatively low differentiation in non-human primates all of whom had the largest numbers of cells transplanted (~3.5M of which only 1.4% differentiated). Again, if one were to speculate on the translational relevance of this, it might suggest that micro-transplantation

– with multiple deposits of smaller numbers of cells – might have a better outcome than large deposits.

Although four clinical trials are ongoing to test the safety, tolerability and efficacy of both ESC and iPSC-derived transplants, this systematic review highlights the variability in survival and differentiation in the published pre-clinical literature. It also suggests that inclusion of proliferative cells within the transplant is still an ongoing issue that could undermine graft safety, while the relatively low differentiation (and negative correlation between differentiation and cells transplanted) could undermine the efficacy of this approach. Overall, this systematic review indicates that there remains scope for improvement in the differentiation of stem cell-derived dopaminergic progenitors in order to maximize the therapeutic potential of this approach for patients.

Acknowledgements

Our research in this field is supported by research grants from the Michael J Fox Foundation for Parkinson's Research (Grant Numbers: 17244 and 023410) and Science Foundation Ireland (Grant Numbers: 19/FFP/6554).

References

1. Bjorklund, A. and M. Parmar, *Neuronal Replacement as a Tool for Basal Ganglia Circuitry Repair: 40 Years in Perspective.* Front Cell Neurosci, 2020. **14**: p. 146.

2. Perlow, M.J., et al., *Brain grafts reduce motor abnormalities produced by destruction of nigrostriatal dopamine system.* Science, 1979. **204**(4393): p. 643-7.

3. Bjorklund, A., R.H. Schmidt, and U. Stenevi, *Functional reinnervation of the neostriatum in the adult rat by use of intraparenchymal grafting of dissociated cell suspensions from the substantia nigra.* Cell Tissue Res, 1980. **212**(1): p. 39-45.

4. Madrazo, I., et al., *Transplantation of fetal substantia nigra and adrenal medulla to the caudate nucleus in two patients with Parkinson's disease.* N Engl J Med, 1988. **318**(1): p. 51.

5. Lindvall, O., et al., *Human fetal dopamine neurons grafted into the striatum in two patients with severe Parkinson's disease. A detailed account of methodology and a 6-month follow-up.* Arch Neurol, 1989. **46**(6): p. 615-31.

6. Barker, R.A. and A. Bjorklund, *Restorative cell and gene therapies for Parkinson's disease.* Handb Clin Neurol, 2023. **193**: p. 211-226.

7. Martin, G.R., *Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells.* Proc Natl Acad Sci U S A, 1981. **78**(12): p. 7634-8.

8. Evans, M.J. and M.H. Kaufman, *Establishment in culture of pluripotential cells from mouse embryos.* Nature, 1981. **292**(5819): p. 154-6.

9. Thomson, J.A., et al., *Embryonic stem cell lines derived from human blastocysts.* Science, 1998. **282**(5391): p. 1145-7.

10. Studer, L., *Derivation of dopaminergic neurons from pluripotent stem cells.* Prog Brain Res, 2012. **200**: p. 243-63.

11. Kim, J.H., et al., *Dopamine neurons derived from embryonic stem cells function in an animal model of Parkinson's disease.* Nature, 2002. **418**(6893): p. 50-6.

12. Ben-Hur, T., et al., *Transplantation of human embryonic stem cell-derived neural progenitors improves behavioral deficit in Parkinsonian rats.* Stem Cells, 2004. **22**(7): p. 1246-55.

13. Yu, J., et al., *Induced pluripotent stem cell lines derived from human somatic cells.* Science, 2007. **318**(5858): p. 1917-20.

14. Takahashi, K. and S. Yamanaka, *Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors.* Cell, 2006. **126**(4): p. 663-76.

15. Takahashi, K., et al., *Induction of pluripotent stem cells from adult human fibroblasts by defined factors.* Cell, 2007. **131**(5): p. 861-72.

16. Hargus, G., et al., *Differentiated Parkinson patient-derived induced pluripotent stem cells grow in the adult rodent brain and reduce motor asymmetry in Parkinsonian rats.* Proc Natl Acad Sci U S A, 2010. **107**(36): p. 15921-6.

17.   Wernig, M., et al., *Neurons derived from reprogrammed fibroblasts functionally integrate into the fetal brain and improve symptoms of rats with Parkinson's disease.* Proc Natl Acad Sci U S A, 2008. **105**(15): p. 5856-61.

18.   Page, M.J., et al., *The PRISMA 2020 statement: an updated guideline for reporting systematic reviews.* BMJ, 2021. **372**: p. n71.

19.   Kirkeby, A., et al., *Predictive Markers Guide Differentiation to Improve Graft Outcome in Clinical Translation of hESC-Based Therapy for Parkinson's Disease.* Cell Stem Cell, 2017. **20**(1): p. 135-148.

20.   Park, C.H., et al., *In vitro and in vivo analyses of human embryonic stem cell-derived dopamine neurons.* J Neurochem, 2005. **92**(5): p. 1265-76.

21.   Brederlau, A., et al., *Transplantation of human embryonic stem cell-derived cells to a rat model of Parkinson's disease: effect of in vitro differentiation on graft survival and teratoma formation.* Stem Cells, 2006. **24**(6): p. 1433-40.

22.   Roy, N.S., et al., *Functional engraftment of human ES cell-derived dopaminergic neurons enriched by coculture with telomerase-immortalized midbrain astrocytes.* Nat Med, 2006. **12**(11): p. 1259-68.

23.   Martinat, C., et al., *Cooperative transcription activation by Nurr1 and Pitx3 induces embryonic stem cell maturation to the midbrain dopamine neuron phenotype.* Proc Natl Acad Sci U S A, 2006. **103**(8): p. 2874-9.

24.   Sonntag, K.C., et al., *Enhanced yield of neuroepithelial precursors and midbrain-like dopaminergic neurons from human embryonic stem cells using the bone morphogenic protein antagonist noggin.* Stem Cells, 2007. **25**(2): p. 411-8.

25.   Iacovitti, L., et al., *A protocol for the differentiation of human embryonic stem cells into dopaminergic neurons using only chemically defined human additives: Studies in vitro and in vivo.* Brain Res, 2007. **1127**(1): p. 19-25.

26.   Ko, J.Y., et al., *Human embryonic stem cell-derived neural precursors as a continuous, stable, and on-demand source for human dopamine neurons.* J Neurochem, 2007. **103**(4): p. 1417-29.

27.   Chiba, S., et al., *Noggin enhances dopamine neuron production from human embryonic stem cells and improves behavioral outcome after transplantation into Parkinsonian rats.* Stem Cells, 2008. **26**(11): p. 2810-20.

28.   Yang, D., et al., *Human embryonic stem cell-derived dopaminergic neurons reverse functional deficit in parkinsonian rats.* Stem Cells, 2008. **26**(1): p. 55-63.

29.   Geeta, R., et al., *One year survival and significant reversal of motor deficits in parkinsonian rats transplanted with hESC derived dopaminergic neurons.* Biochem Biophys Res Commun, 2008. **373**(2): p. 258-64.

30.   Ko, J.Y., et al., *Conditions for tumor-free and dopamine neuron-enriched grafts after transplanting human ES cell-derived neural precursor cells.* Mol Ther, 2009. **17**(10): p. 1761-70.

31.   Cai, J., et al., *Dopaminergic neurons derived from human induced pluripotent stem cells survive and integrate into 6-OHDA-lesioned rats.* Stem Cells Dev, 2010. **19**(7): p. 1017-23.

32. Kriks, S., et al., *Dopamine neurons derived from human ES cells efficiently engraft in animal models of Parkinson's disease.* Nature, 2011. **480**(7378): p. 547-51.

33. Cho, E.G., et al., *MEF2C enhances dopaminergic neuron differentiation of human embryonic stem cells in a parkinsonian rat model.* PLoS One, 2011. **6**(8): p. e24027.

34. Kikuchi, T., et al., *Survival of human induced pluripotent stem cell-derived midbrain dopaminergic neurons in the brain of a primate model of Parkinson's disease.* J Parkinsons Dis, 2011. **1**(4): p. 395-412.

35. Rhee, Y.H., et al., *Protein-based human iPS cells efficiently generate functional dopamine neurons and can treat a rat model of Parkinson disease.* J Clin Invest, 2011. **121**(6): p. 2326-35.

36. Kirkeby, A., et al., *Generation of regionally specified neural progenitors and functional neurons from human embryonic stem cells under defined conditions.* Cell Rep, 2012. **1**(6): p. 703-14.

37. Doi, D., et al., *Prolonged maturation culture favors a reduction in the tumorigenicity and the dopaminergic function of human ESC-derived neural cells in a primate model of Parkinson's disease.* Stem Cells, 2012. **30**(5): p. 935-45.

38. Daadi, M.M., et al., *Dopaminergic neurons from midbrain-specified human embryonic stem cell-derived neural stem cells engrafted in a monkey model of Parkinson's disease.* PLoS One, 2012. **7**(7): p. e41120.

39. Emborg, M.E., et al., *Intracerebral transplantation of differentiated human embryonic stem cells to hemiparkinsonian monkeys.* Cell Transplant, 2013. **22**(5): p. 831-8.

40. Kang, X., et al., *Dopamine release from transplanted neural stem cells in Parkinsonian rat striatum in vivo.* Proc Natl Acad Sci U S A, 2014. **111**(44): p. 15804-9.

41. Wakeman, D.R., et al., *Survival and integration of neurons derived from human embryonic stem cells in MPTP-lesioned primates.* Cell Transplant, 2014. **23**(8): p. 981-94.

42. Grealish, S., et al., *Human ESC-derived dopamine neurons show similar preclinical efficacy and potency to fetal neurons when grafted in a rat model of Parkinson's disease.* Cell Stem Cell, 2014. **15**(5): p. 653-65.

43. Yang, F., et al., *Activated astrocytes enhance the dopaminergic differentiation of stem cells and promote brain repair through bFGF.* Nat Commun, 2014. **5**: p. 5627.

44. Doi, D., et al., *Isolation of human induced pluripotent stem cell-derived dopaminergic progenitors by cell sorting for successful transplantation.* Stem Cell Reports, 2014. **2**(3): p. 337-50.

45. Steinbeck, J.A., et al., *Optogenetics enables functional analysis of human embryonic stem cell-derived grafts in a Parkinson's disease model.* Nat Biotechnol, 2015. **33**(2): p. 204-9.

46. Grealish, S., et al., *Monosynaptic Tracing using Modified Rabies Virus Reveals Early and Extensive Circuit Integration of Human Embryonic Stem Cell-Derived Neurons.* Stem Cell Reports, 2015. **4**(6): p. 975-83.

47.   Han, F., et al., *Human induced pluripotent stem cell-derived neurons improve motor asymmetry in a 6-hydroxydopamine-induced rat model of Parkinson's disease.* Cytotherapy, 2015. **17**(5): p. 665-79.

48.   Chen, Y., et al., *Chemical Control of Grafted Human PSC-Derived Neurons in a Mouse Model of Parkinson's Disease.* Cell Stem Cell, 2016. **18**(6): p. 817-826.

49.   Nishimura, K., et al., *Estradiol Facilitates Functional Integration of iPSC-Derived Dopaminergic Neurons into Striatal Neuronal Circuits via Activation of Integrin alpha5beta1.* Stem Cell Reports, 2016. **6**(4): p. 511-524.

50.   Niclis, J.C., et al., *Efficiently Specified Ventral Midbrain Dopamine Neurons from Human Pluripotent Stem Cells Under Xeno-Free Conditions Restore Motor Deficits in Parkinsonian Rodents.* Stem Cells Transl Med, 2017. **6**(3): p. 937-948.

51.   Niclis, J.C., et al., *A PITX3-EGFP Reporter Line Reveals Connectivity of Dopamine and Non-dopamine Neuronal Subtypes in Grafts Generated from Human Embryonic Stem Cells.* Stem Cell Reports, 2017. **9**(3): p. 868-882.

52.   Kikuchi, T., et al., *Idiopathic Parkinson's disease patient-derived induced pluripotent stem cells function as midbrain dopaminergic neurons in rodent brains.* J Neurosci Res, 2017. **95**(9): p. 1829-1837.

53.   Kikuchi, T., et al., *Human iPS cell-derived dopaminergic neurons function in a primate Parkinson's disease model.* Nature, 2017. **548**(7669): p. 592-596.

54.   Wakeman, D.R., et al., *Cryopreservation Maintains Functionality of Human iPSC Dopamine Neurons and Rescues Parkinsonian Phenotypes In Vivo.* Stem Cell Reports, 2017. **9**(1): p. 149-161.

55.   Cardoso, T., et al., *Target-specific forebrain projections and appropriate synaptic inputs of hESC-derived dopamine neurons grafted to the midbrain of parkinsonian rats.* J Comp Neurol, 2018. **526**(13): p. 2133-2146.

56.   Wang, Y.K., et al., *Human Clinical-Grade Parthenogenetic ESC-Derived Dopaminergic Neurons Recover Locomotive Defects of Nonhuman Primate Models of Parkinson's Disease.* Stem Cell Reports, 2018. **11**(1): p. 171-182.

57.   Adler, A.F., et al., *hESC-Derived Dopaminergic Transplants Integrate into Basal Ganglia Circuitry in a Preclinical Model of Parkinson's Disease.* Cell Rep, 2019. **28**(13): p. 3462-3473 e5.

58.   Yu, S.J., et al., *NanoCsA improves the survival of human iPSC transplant in hemiparkinsonian rats.* Brain Res, 2019. **1719**: p. 124-132.

59.   Zygogianni, O., et al., *In Vivo Phenotyping of Familial Parkinson's Disease with Human Induced Pluripotent Stem Cells: A Proof-of-Concept Study.* Neurochem Res, 2019. **44**(6): p. 1475-1493.

60.   Gantner, C.W., et al., *Viral Delivery of GDNF Promotes Functional Integration of Human Stem Cell Grafts in Parkinson's Disease.* Cell Stem Cell, 2020. **26**(4): p. 511-526 e5.

61.   Goggi, J.L., et al., *Dopamine transporter neuroimaging accurately assesses the maturation of dopamine neurons in a preclinical model of Parkinson's disease.* Stem Cell Res Ther, 2020. **11**(1): p. 347.

62.   Hoban, D.B., et al., *Impact of alpha-synuclein pathology on transplanted hESC-derived dopaminergic neurons in a humanized alpha-synuclein rat model of PD*. Proc Natl Acad Sci U S A, 2020. **117**(26): p. 15209-15220.

63.   Tiklova, K., et al., *Single cell transcriptomics identifies stem cell-derived graft composition in a model of Parkinson's disease*. Nat Commun, 2020. **11**(1): p. 2434.

64.   Doi, D., et al., *Pre-clinical study of induced pluripotent stem cell-derived dopaminergic progenitor cells for Parkinson's disease*. Nat Commun, 2020. **11**(1): p. 3369.

65.   Song, B., et al., *Human autologous iPSC-derived dopaminergic progenitors restore motor function in Parkinson's disease models*. J Clin Invest, 2020. **130**(2): p. 904-920.

66.   Hiller, B.M., et al., *Mitomycin-C treatment during differentiation of induced pluripotent stem cell-derived dopamine neurons reduces proliferation without compromising survival or function in vivo*. Stem Cells Transl Med, 2021. **10**(2): p. 278-290.

67.   Yoo, J.E., et al., *Trophoblast glycoprotein is a marker for efficient sorting of ventral mesencephalic dopaminergic precursors derived from human pluripotent stem cells*. NPJ Parkinsons Dis, 2021. **7**(1): p. 61.

68.   Piao, J., et al., *Preclinical Efficacy and Safety of a Human Embryonic Stem Cell-Derived Midbrain Dopamine Progenitor Product, MSK-DA01*. Cell Stem Cell, 2021. **28**(2): p. 217-229 e7.

69.   de Luzy, I.R., et al., *Human stem cells harboring a suicide gene improve the safety and standardisation of neural transplants in Parkinsonian rats*. Nat Commun, 2021. **12**(1): p. 3275.

70.   Xiong, M., et al., *Human Stem Cell-Derived Neurons Repair Circuits and Restore Neural Function*. Cell Stem Cell, 2021. **28**(1): p. 112-126 e6.

71.   Shrigley, S., et al., *Grafts Derived from an alpha-Synuclein Triplication Patient Mediate Functional Recovery but Develop Disease-Associated Pathology in the 6-OHDA Model of Parkinson's Disease*. J Parkinsons Dis, 2021. **11**(2): p. 515-528.

72.   Elabi, O.F., et al., *Human Embryonic Stem Cell-Derived Dopaminergic Grafts Alleviate L-DOPA Induced Dyskinesia*. J Parkinsons Dis, 2022. **12**(6): p. 1881-1896.

73.   Lane, E.L., et al., *Spontaneous Graft-Induced Dyskinesias Are Independent of 5-HT Neurons and Levodopa Priming in a Model of Parkinson's Disease*. Mov Disord, 2022. **37**(3): p. 613-619.

74.   Aldrin-Kirk, P., et al., *A novel two-factor monosynaptic TRIO tracing method for assessment of circuit integration of hESC-derived dopamine transplants*. Stem Cell Reports, 2022. **17**(1): p. 159-172.

75.   Hiller, B.M., et al., *Optimizing maturity and dose of iPSC-derived dopamine progenitor cell therapy for Parkinson's disease*. NPJ Regen Med, 2022. **7**(1): p. 24.

76.   de Luzy, I.R., et al., *Identifying the optimal developmental age of human pluripotent stem cell-derived midbrain dopaminergic progenitors for transplantation in a rodent model of Parkinson's disease*. Exp Neurol, 2022. **358**: p. 114219.

77.   Moriarty, N., et al., *Understanding the Influence of Target Acquisition on Survival, Integration, and Phenotypic Maturation of Dopamine Neurons within Stem Cell-Derived Neural Grafts in a Parkinson's Disease Model.* J Neurosci, 2022. **42**(25): p. 4995-5006.

78.   Brot, S., et al., *Long-Term Evaluation of Intranigral Transplantation of Human iPSC-Derived Dopamine Neurons in a Parkinson's Disease Mouse Model.* Cells, 2022. **11**(10).

79.   Kirkeby, A., et al., *Preclinical quality, safety, and efficacy of a human embryonic stem cell-derived product for the treatment of Parkinson's disease, STEM-PD.* Cell Stem Cell, 2023. **30**(10): p. 1299-1314 e9.

80.   Maimaitili, M., et al., *Enhanced production of mesencephalic dopaminergic neurons from lineage-restricted human undifferentiated stem cells.* Nat Commun, 2023. **14**(1): p. 7871.

81.   Naderi, S., et al., *Cryopreserved clinical-grade human embryonic stem cell-derived dopaminergic progenitors function in Parkinson's disease models.* Life Sci, 2023. **329**: p. 121990.

82.   Lopez-Ornelas, A., et al., *Human Embryonic Stem Cell-Derived Immature Midbrain Dopaminergic Neurons Transplanted in Parkinsonian Monkeys.* Cells, 2023. **12**(23).

83.   Rajova, J., et al., *Deconvolution of spatial sequencing provides accurate characterization of hESC-derived DA transplants in vivo.* Mol Ther Methods Clin Dev, 2023. **29**: p. 381-394.

84.   Nakamura, R., et al., *A defined method for differentiating human iPSCs into midbrain dopaminergic progenitors that safely restore motor deficits in Parkinson's disease.* Front Neurosci, 2023. **17**: p. 1202027.

85.   Ramos-Acevedo, R., et al., *Embryoid Body Cells from Human Embryonic Stem Cells Overexpressing Dopaminergic Transcription Factors Survive and Initiate Neurogenesis via Neural Rosettes in the Substantia Nigra.* Brain Sci, 2023. **13**(2).

86.   Hills, R., et al., *Neurite Outgrowth and Gene Expression Profile Correlate with Efficacy of Human Induced Pluripotent Stem Cell-Derived Dopamine Neuron Grafts.* Stem Cells Dev, 2023. **32**(13-14): p. 387-397.

87.   Lyu, Y., et al., *MiR-210-5p promotes the differentiation of human induced pluripotent stem cells into dopaminergic neural precursors by targeting SMAD4 and SUFU and treats parkinsonian rats.* Exp Gerontol, 2023. **179**: p. 112243.

88.   Daadi, E.W., et al., *Combining physical & cognitive training with iPSC-derived dopaminergic neuron transplantation promotes graft integration & better functional outcome in parkinsonian marmosets.* Exp Neurol, 2024. **374**: p. 114694.

89.   Fu, C.L., et al., *A cell therapy approach based on iPSC-derived midbrain organoids for the restoration of motor function in a Parkinson's disease mouse model.* Heliyon, 2024. **10**(2): p. e24234.

90.   Park, S., et al., *Preclinical and dose-ranging assessment of hESC-derived dopaminergic progenitors for a clinical trial on Parkinson's disease.* Cell Stem Cell, 2024. **31**(1): p. 25-38 e8.

91.	Comini, G., et al., *Survival and maturation of human induced pluripotent stem cell-derived dopaminergic progenitors in the Parkinsonian rat brain is enhanced by transplantation in a neurotrophin-enriched hydrogel.* J Neural Eng, 2024.

92.	Liang, Q., et al., *Linking a cell-division gene and a suicide gene to define and improve cell therapy safety.* Nature, 2018. **563**(7733): p. 701-704.

93.	Wenker, S.D., et al., *Cell therapy for Parkinson's disease: Functional role of the host immune response on survival and differentiation of dopaminergic neuroblasts.* Brain Res, 2016. **1638**(Pt A): p. 15-29.