Diel Partitioning in Microbial Phosphorus Acquisition in the Sargasso Sea

Muratore, Daniel[1,†], Gilbert, Naomi E.[2,3,†], LeCleir, Gary R.[2], Wilhelm, Steven W.[2*], Weitz, Joshua S.[4,5,6*],

† Authors contributed equally to this manuscript
1 Santa Fe Institute, Santa Fe, NM, USA
2 Department of Microbiology, The University of Tennessee, Knoxville, TN, USA
3 Lawrence Livermore National Laboratory, Livermore, CA, USA
4 Department of Biology, University of Maryland, College Park, MD, USA
5 Department of Physics, University of Maryland, College Park, MD, USA
6 Institute de Biologie, École Normale Supérieure, Paris, France

E-mail: jsweitz@umd.edu (J.S.W), wilhelm@utk.edu (S.W.W)

Abstract

The daily cycle of photosynthetic primary production at the base of complex marine food webs is often limited by the availability and uptake of scarce nutrients. According to temporal niche partitioning theory, competition for scarce resources can be alleviated insofar as the intensity of nutrient uptake and assimilation activities are distributed heterogeneously across organisms over periodic input cycles. Recent analysis of community transcriptional dynamics in the nitrogen-limited suptropical North Pacific gyre revealed evidence of temporal partitioning of nitrogen uptake and assimilation between eukaryotic phytoplankton, cyanobacteria, and heterotrophic bacteria over day-night cycles. Here, we present results from a Lagrangian metatranscriptomic time series survey in the Sargasso Sea and demonstrate temporally partitioned phosphorus uptake in this phosphorus-limited environment. In the Sargasso, heterotrophic bacteria, eukaryotic phytoplankton, and cyanobacteria express genes for phosphorus assimilation during the morning, day, and dusk, respectively. These results support the generality of temporal niche partitioning as an emergent mechanism structuring uptake of limiting nutrients in open ocean microbial ecosystems.

Introduction

In the ocean, communities of microorganisms set ecological rhythms of primary production and respiration in response to the daily light cycle, fueling the base of the marine food web and modulating biogeochemical cycles [7, 13, 16, 20, 34]. Diel oscillations in carbon fixation among photosynthetic plankton cascade through community metabolism and gene expression, resulting in emergent ecosystem-level cycles in biomass and biomass molecular composition [2–4, 12, 27, 28]. Diel transcriptional and metabolic organization also expands beyond carbon metabolism. Field studies in nitrogen-starved ocean regions [36] have identified diel transcriptional cycles in nitrogen uptake [38] and nitrogen fixation [39]. Heterotrophic taxa from the same region have also been reported to express organic nutrient transporters on a diel

cycle [27–29]. Organisms taking turns in fulfilling specific ecological niches or taking up 12
resources has been termed temporal niche partitioning in terrestrial ecology [18,30]. 13
Niche partitioning over diel cycles has been ascribed to predatory behavior in marine 14
animals [19] and invoked in microbial community successional dynamics in aquatic 15
environments [8,27]. Diel cycles identified across diverse mechanisms for satisfying 16
nitrogen requirements among multiple taxa in a nitrogen-limited system suggest that 17
ecosystems facing limitation for other nutrients may have similar strategies. 18

The Sargasso Sea is an open ocean environment where plankton are considered to be 19
persistently phosphorus-stressed at the surface [9,34,36]. In the subtropical 20
northwestern Atlantic, concentrations of dissolved phosphate can be sub-nanomolar [40], 21
sufficient to induce phosphate stress in primary producers [9,36]. Diverse strategies 22
have evolved to mitigate phosphorus limitation among marine microbes. For example, 23
sulfono- and betaine lipids have been shown to replace phospholipids as a method of 24
reducing cell phosphorus quotas [6,26,37]. Utilizing phosphonates, organic 25
phosphorus-containing molecules, as an alternative phosphorus source has also been 26
described in heterotrophic [33] and photosynthetic bacteria [32]. In the Sargasso Sea, 27
dissolved organic phosphorus has been measured to average almost 95% of total 28
dissolved phosphorus [23]. High copy numbers of the C-P lyase *phnJ* gene and strong 29
responses in a proxy for C-P lyase activity have also been observed *in situ* [31]. This 30
evidence reinforces the ecological relevance of phosphorus limitation in the Sargasso Sea, 31
which we hypothesized subjects phosphorus uptake to similar pressures with respect to 32
competition as in the case of nitrogen uptake, demonstrated previously within 33
nitrogen-limited systems [27]. Indeed, a field study of *Trichodesmium* gene expression 34
found evidence of diel phosphate and phosphonate utilization consistent with this 35
hypothesis [10]. 36

Here, we tested the hypothesis that a phosphorus-limited ocean region will partition 37
phosphorus uptake over the diel cycle within a Lagrangian metatranscriptomic time 38
series survey, sampling every 4 hours over the course of 5 days. Our community 39
metatranscriptomic analyses demonstrate widespread temporal partitioning of diverse 40
phosphorus uptake and incorporation genes, and suggests a diel cascade of phosphorus 41
uptake from heterotrophic bacteria at sunrise, to eukaryotic phytoplankton during the 42
day, to cyanobacteria at dusk. The results reinforce the generality of temporal niche 43
partitioning as a possible mechanism of competition alleviation in open ocean microbial 44
ecosystems. 45

Results 46

We analyzed 58 metatranscriptomes collected during the Lagrangian sampling 47
expedition on the R/V *Atlantic Explorer* (AE1926) from October 12-17, 2019 48
(Figure 1a) in the vicinity of the Bermuda Atlantic Time Series (BATS) study location 49
in the Sargasso Sea to examine diel changes in microbial community-level gene 50
expression (SI Data S1). Using a nonparametric method with multiple testing 51
correction set at a 10% false discovery rate [35], we evaluated whether gene transcripts 52
had 24-hour periodicity in their abundance patterns (see Methods). In all, we evaluated 53
97,829 different genes from eukaryotic phyla, cyanobacterial genera *Prochlorococcus* and 54
Synechococcus, and heterotrophic bacterial orders for diel periodicity. Our analysis 55
found 11,194 significantly diel genes (SI Data S2). Diel genes spanned 21 phyla of 56
eukaryotic phytoplankton (9.33% of total genes), the cyanobacterial groups 57
Prochlorococcus and *Synechococcus* (37.4% of total genes), and heterotrophic bacteria 58
representing 6 major classes comprised of 44 orders (13.2% of total genes). A similar 59
analysis conducted on data from a cruise in the North Pacific Subtropical Gyre 60
examined 64,011 genes [27], and found a similar proportion (8.94% vs 11.4% reported 61

Figure 1. Analysis of community-level diel gene expression. a) Lagrangian cruise track with BATS station annotated in red and sampling locations annotated in black. Initial and final samples are labelled with local date and time. b) Example gene transcript abundance time series for diel genes belonging to the transporter KEGG pathway. Individual black lines represent the transcript abundance of one gene, the blue line represents a generalized additive model smoothing among all genes. Panels indicate the classification of genes by peak time as determined by the peak time analysis described in Methods. Shaded vertical bars indicate hours after sunset and before sunrise (18:00-06:00 local time). c) Mean pairwise differences in peak timing for genes with diel expression across at least 4 distinct taxa. Key genes in nitrogen assimilation and phosphorus assimilation are indicated. Orange point color indicates a gene with significantly (FDR=10%) synchronized transcription. Green points indicate a gene without synchronized transcription, labeled 'asynchronous'.

here) of genes with diel transcript abundances. 62

We estimated the peak time of day for each gene determined to have diel periodicity. 63
Eukaryotic phytoplankton, cyanobacteria, and heterotrophic bacteria had diel 64
expression patterns of different genes occurr throughout all times of day, with 65
eukaryotic phytoplankton expressing the greatest number of diel genes at dusk, 66
heterotrophs during the day, and cyanobacteria divided between dawn and dusk (SI 67
Data S3). Example gene expression time series for diel genes with annotations as 68
ABC-type transporters are shown in Figure 1b. 69

Next we assessed the extent to which the diel timing was synchronized among genes 70
with similar putative function. Genes were functionally annotated using the Kyoto 71

Encyclopedia of Genes and Genomes (KEGG) orthology assignments [1]. The diel genes spanned 3,647 ortholgues. We further analyzed the 849 KEGG orthologues with diel expression in at least four different taxa (eukaryotic phyla/cyanobacterial genera/heterotrophic bacterial orders). For each of these orthologues, we calculated the mean difference in peak time between all pairs of taxa with diel expression of that orthologue. We used an empirical hypothesis testing framework to assess whether that difference was significantly smaller than would be expected by chance given the data set (see Methods). At a false discovery rate of 10%, we identified 324 out of 849 orthologues with significantly synchronized timing across taxa. Significantly synchronized genes included cell division and replication related functions ($ftsH$,$dnaK$), photosynthesis-related genes ($psbD$,$psbA$), and carbon metabolism genes related to the tricarboxylic acid (TCA) cycle and glycine metabolism ($sdhA$,$coxA$) (Figure 1c, SI Data S4).

A pattern also emerged from genes with high degrees of asynchrony across taxa. In particular, genes encoding the uptake and assimilation of nutrients had high asynchrony. We found asynchronous expression of some nitrogen assimilation-related genes, such as components of the GS-GOGAT system and its regulatory proteins as well as the ammonium amt uptake receptor (Figure 1c). Asynchronized expression of these genes is possibly related to temporal niche partitioning, which may help to reduce the severity of competition for ammonium [27]. These results may suggest that this form of nitrogen partitioning is common to the oligotrophic ocean. However, we find an even higher degree of asynchrony in genes involved in the uptake and incorporation of phosphorus. The phosphonate transporter components $phnC$ and $phnD$, inorganic phosphorus transporter $pstS$, phosphate stress regulatory genes $phoH$ and $phoD$, inorganic pyrophosphatase ppa, and polyphosphate kinases $ppk1$ and $ppk2$ all had among the highest degrees of asynchrony in the data set (Figure 1c). A higher degree of asynchrony for phosphorus uptake functions may be further indication of the relevance of diel temporal partitioning as an emergent ecological strategy to overcome limitation of shared scarce resources.
We then examined the taxon-specific transcriptional dynamics of genes involved in the uptake of inorganic and organic phosphorus species to look for specific evidence of uptake partitioning among different functional groups of organisms. We extracted KEGG orthologues specifically annotated as either inorganic phosphate transporters ($pstABC$,$pstS$ - KEGG orthologues K02036,K02037,K02038,K02040) or organic phosphonate transporters ($phnCDE$,$phnK$,$phnS$ - KEGG orthologues K02041, K02042, K02044, K05781, K11081) and analyzed the relative peak timings of different taxonomic groups. Among these transporters, $pstS$, $phnC$, and $phnK$ had significantly diel transcript expression in at least 4 different taxa, and none of the transporters had synchronized expression in our synchronicity analysis. Assembling the gene expression time series of all phosphorus-uptake related transporters among all taxa show different timings in uptake function among cyanobacteria, eukaryotic phytoplankton, and heterotrophic bacteria (see Figure 2). The inorganic phosphorus transporter gene $pstS$ had diel transcription in Rhodospiralles and another alphaproteobacterial order peaking at 08:00 (roughly corresponding to sunrise) along with transcripts assigned to the phylum Dinophyceae, while transcripts assigned to diatoms peaked at 12:00 (noon) and both *Prochlorococcus* and *Synechococcus* had peak expression at 20:00 (roughly corresponding to dusk) (Figure 2). Meanwhile, organic phosphonate transporters are widely transcribed by heterotrophic bacteria, primarily alphaproteobacteria, throughout the day. No eukaryotic organisms had significantly diel transcription of phn phosphonate transporters. *Prochlorococcus* and *Synechococcus* had diel expression of the phosphonate transporter subunits $phnC$ and $phnD$ (*Prochlorococcus* only), but

Prochlorococcus had peak transcript abundance of these genes during the day and at dusk, concomitant with inorganic phosphorus transporters, while *Synechococcus* expressed *phnC* at 00:00 (midnight), after transcript abundance of *pstS* began to decline. Rhodospiralles had peak transcript abundance of both *pstS* and *phnD* at the same time (08:00), while the alphaproteobacterial order Rhodobacterales did not have diel expression of any inorganic phosphate transporters, but did have diel expression of *phnC* and *phnK* during at 16:00 (Figure 2). Notably, the taxa that did have diel expression of phosphorus transporter systems, regardless of inorganic or organic species, tended to have peak transcript abundance of all phosphorus transport genes at around the same time of day, with the exception of *Synechococcus*, which appeared to exhibit a temporal cascade from primarily inorganic phosphorus transporter transcripts at dusk to phosphonate transporter transcripts at midnight. Therefore, apart from *Synechococcus*, timing of diel expression for phosphorus uptake transporters is not influenced by phosphorus being in organic versus inorganic forms. These results suggest that diel partitioning of phosphorus uptake is not entirely due to extrinsic diel cycles in phosphorus speciation that may be induced by e.g., photolysis of phosphorus-containing dissolved organic matter.

Discussion

Here, we tested the hypothesis that complex microbes would temporally partition phosphorous uptake and utilization gene expression in a phosphorus-stressed open ocean oligotrophic ecosystem in the North Atlantic, paralleling previous findings on nitrogen uptake from a primarily nitrogen-limited ecosystem in the North Pacific [27]. Using community metatranscriptome data from a Lagrangian field campaign in the Sargasso Sea, we identified diel gene transcription of phosphorus uptake and assimilation genes across diverse microorganisms. We found diel gene expression was not synchronized across the community, and was in fact even less synchronized than genes with diel expression across diverse taxa related to nitrogen uptake and assimilation. Studying the dynamics of phosphate and phosphonate transporters in more detail, we identified a diel cascade in phosphate transporter expression starting with heterotrophic bacteria in the morning, eukaryotic phytoplankton during the day, and cyanobacteria at dusk. Likewise, we found diel gene transcript abundances for phosphonate transporters with the heterotrophic bacterial order Rhodospiralles expressing phosphonate transporters in the morning (coinciding with phosphate transporters), Rhodobacter during the day, *Prochlorococcus* at dusk (coinciding with its phosphate transporters), and *Synechococcus* overnight (following its phosphate transporters). Our results suggest temporal partitioning of uptake for both phosphorus species, and coinciding timing of phosphate and phosphonate uptake on a taxon-by-taxon basis further suggests that substrate-level specialization is not necessarily sufficient to avoid competition for this scarce resource [25], and further suggests diel cycles in transporter gene transcription are not directly tied to changes in the relative pools of different phosphorus species. Complementary evidence for diel temporal niche partitioning for distinct nutrients (N in the North Subtropical Pacific [27] and P in the Sargasso as shown here) may be indicative of complex, community-level metabolic plasticity across ocean basins. Diel temporal niche partitioning to mitigate competition for limiting resources in oligotrophic environments represents an additional mechanism to resolve the paradox of the plankton and the maintenance of complex, marine microbial communities [15].

Figure 2. Temporal dynamics of phosphorus acquisition gene transcription. Each line plot shows the gene transcript abundance time series of a specific phosphorus-uptake related gene for a specific eukaryotic phylum or bacterial order (or cyanobacterial genus). Line color indicates the taxa expressing the transcript. Panels are divided horizontally by time of peak expression as calculated in methods, with 08:00 representing 8am local time, 12:00 noon local time, and so on. Shaded bars indicate hours between sunset and sunrise (18:00-06:00 local). Dashed lines are added to correspond to the peak time of genes in each panel. Panels are divided vertically by the species the gene is involved in transporting. The phosphate uptake category includes phosphate transporter subunits *pstABC,pstS* (KEGG K02036,K02037,K02038,K02040). The phosphonate uptake category includes the phosphonate transporter subunits *phnCDE* and putative phosphatase uptake subunits *phnK,phnS* (KEGG K02041, K02042, K02044, K05781, K11081).

Materials and Methods 169

Sample Collection and Processing 170

Fieldwork was conducted from 12 October to 19 October, 2019, in the Sargasso Sea 171
near the location of the BATS station. A drogue was deployed before the first cast to 172
facilitate a Langrangian sampling stratey. Water column sampling took place every 4 173
hours for a period of approximately 5 days sampling 5 m depth. See Supporting 174
Information for details on RNA collection and extraction, library preparation and 175
metatranscriptome processing. 176

Gene Expression Diel Periodicity and Synchronization Analysis

Please see Supporting Information for more details on timeseries analysis. Briefly, variance-stabilized RNA read counts were subjected to RAIN nonparametric diel analysis as described in [27]. Peak rank time calculations were performed and synchronicity analysis was conducted as described in [27].

Metatranscriptome Preparation, Sequencing, and Processing

Seawater was filtered through a 0.2 micron pore size Sterivex filtration unit and immediately transferred to a -80 C freezer. RNA was extracted using a publicly available phenol-chloroform based protocol [11] and DNA was reduced using the Turbo DNA-free kit (Ambion). Metatranscriptome libraries were prepared by reducing ribosomal RNA using the QIAGEN's FastSelect kit (Bacteria) and sequenced (2 x 151 nt) using the low-input protocol for total RNA on the Illumina Novaseq S4 platform under the DOE Joint Genome Institute (JGI) Community Sequence Proposal ID 505733. For samples with <10 ng of RNA total, the ultra-low input protocol was used. Raw read filtering and trimming was done using BBDuk v38.67 and BBMap v38.84 from the BBtools packages [5]. Trimmed filtered reads were concatenated across samples and assembled using MEGAHIT v1.2.9 [21]. MetaGeneMark v3.38 was used to call open reading frames (ORFs). For the cellular community, trimmed filtered reads were mapped to the combined assembly using BBMap v38.84 [5] using default parameters and tabulated using featureCounts [22]. The ORF protein sequences were annotated using eggNOG-mapper v2.1.4 (with DIAMOND blastp alignment [14]) for functional annotation, and aligned to the PhyloDB database (https://github.com/allenlab/PhyloDB) using the software package EUKulele [17] for taxonomic annotation.

Diel Transcript Abundance Analysis

Prior to timeseries analysis, all metatranscriptome read mappings were transformed using the DESeq2 variance stabilizing transformation [24]. Methods for periodicity determination and peak time estimation are described in detail in [27]. Briefly, for each time series a linear trend was removed to account for moving averages. Then, the rain [35] method was applied and an FDR control at 10% were set to assess significance. For significantly diel time series, peak time was estimated by calculating the mean rank of each time of day across all samples and selecting the largest.

Transcript Synchrony Analysis

Methods are described in detail in [27]. Briefly, only KEGG orthologues with diel transcription in at least 4 different taxa were analyzed. For each orthologue, the pairwise difference in estimated peak time was calculated for each pair of taxa with diel expression of that orthologue. The mean of this value across all pairs was used as a test statistic and significance was assessed using a monte carlo simulated null ensemble of all peak times from the dataset. Significance was set at an FDR=10% threshold.

Data and code availability

Sequence data are publicly available under the DOE JGI Community Sequencing Proposal ID 505733 (`https://genome.jgi.doe.gov/portal/Infvirtimeseries/Infvirtimeseries.info.html`)

Code and processed sequence data used for analyses are available at ²²⁰
`https://github.com/d-muratore/phosphorus_partitioning`. ²²¹

Supplemental files available for this manuscript are described below: ²²²

Supplemental Data 1: Ship navigation data with sampling locations and sample ²²³
environmental data for all metatranscriptome samples. ²²⁴

Supplemental Data 2: Results of RAIN diel periodicity analysis. Columns include ²²⁵
KEGG orthologue tested, taxonomic assignment of KEGG orthologue, RAIN test ²²⁶
p-value, and a reject/fail to reject decision on the null hypothesis that the transcript ²²⁷
does not have a 24-hour diel oscillation. ²²⁸

Supplemental Data 3: Peak time and functional and taxnomic annotation for genes ²²⁹
with diel periodicity. ²³⁰

Supplemental Data 4: Results of KEGG orthologue synchronicity analysis. Columns ²³¹
show the KEGG Orthologue number of the gene tested, the number of taxa in the ²³²
dataset with diel expression of this KO, the pathway assignment of the KO, the average ²³³
difference in peak rank time between all organisms with diel expression of the KO, the ²³⁴
associated empirical p-value (10,000 Monte Carlo simulations) and the reject/fail to ²³⁵
reject decision of the null hypothesis that the transcript is not synchronized. ²³⁶

Acknowledgments ²³⁷

The authors would like to thank the captain and crew of the RV Atlantic Explorer for ²³⁸
conducting cruise AE1926. This work was funded by NSF grant OCE-1829641 to ²³⁹
S.W.W., Simons Foundation grant 735077 to S.W.W, NSF grant OCE-1829636 to ²⁴⁰
J.S.W., Simons Foundation grant 721231 to J.S.W., and the Blaise Pascal Institute ²⁴¹
Chair of Excellence award at the Institut de Biologie of the École Normale Supérieure to ²⁴²
J.S.W. The work (proposal: 10.46936/10.25585/60001286) conducted by the U.S. ²⁴³
Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a DOE ²⁴⁴
Office of Science User Facility, is supported by the Office of Science of the U.S. ²⁴⁵
Department of Energy under contract no. DE-AC02-05CH11231. We thank Christine ²⁴⁶
Sun and Matthew Sullivan for their assistance. Work at LLNL was performed under the ²⁴⁷
auspices of US Department of Energy Office of Science contract DE-AC52-07NA27344. ²⁴⁸
Institution Paper Number LLNL-JRNL-860896-DRAFT. ²⁴⁹

References

1. K. F. Aoki-Kinoshita and M. Kanehisa. *Gene Annotation and Pathway Mapping in KEGG*, pages 71–91. Humana Press, Totowa, NJ, 2007.

2. K. W. Becker, J. R. Collins, B. P. Durham, R. D. Groussman, A. E. White, H. F. Fredricks, J. E. Ossolinski, D. J. Repeta, P. Carini, E. Armbrust, et al. Daily changes in phytoplankton lipidomes reveal mechanisms of energy storage in the open ocean. *Nature communications*, 9(1):1–9, 2018.

3. K. W. Becker, M. J. Harke, D. R. Mende, D. Muratore, J. S. Weitz, E. F. DeLong, S. T. Dyhrman, and B. A. Van Mooy. Combined pigment and metatranscriptomic analysis reveals highly synchronized diel patterns of phenotypic light response across domains in the open oligotrophic ocean. *The ISME Journal*, 15(2):520–533, 2021.

4. A. K. Boysen, L. T. Carlson, B. P. Durham, R. D. Groussman, F. O. Aylward, F. Ribalet, K. R. Heal, A. E. White, E. F. DeLong, E. V. Armbrust, A. E. Ingalls, and J. Bowman. Particulate metabolites and transcripts reflect diel oscillations of microbial activity in the surface ocean. *mSystems*, 6(3):e00896–20, 2021.

5. B. Bushnell. Bbmap: a fast, accurate, splice-aware aligner. Technical report, Lawrence Berkeley National Lab.(LBNL), Berkeley, CA (United States), 2014.

6. P. Carini, B. A. S. V. Mooy, J. C. Thrash, A. White, Y. Zhao, E. O. Campbell, H. F. Fredricks, and S. J. Giovannoni. Sar11 lipid renovation in response to phosphate starvation. *Proceedings of the National Academy of Sciences*, 112(25):7767–7772, 2015.

7. R. Cavicchioli, W. J. Ripple, K. N. Timmis, F. Azam, L. R. Bakken, M. Baylis, M. J. Behrenfeld, A. Boetius, P. W. Boyd, A. T. Classen, et al. Scientists' warning to humanity: microorganisms and climate change. *Nature Reviews Microbiology*, 17(9):569–586, 2019.

8. F. Dini-Andreote, M. de Cássia Pereira e Silva, X. Triado-Margarit, E. O. Casamayor, J. D. Van Elsas, and J. F. Salles. Dynamics of bacterial community succession in a salt marsh chronosequence: evidences for temporal niche partitioning. *The ISME Journal*, 8(10):1989–2001, 2014.

9. S. Duhamel, J. M. Diaz, J. C. Adams, K. Djaoudi, V. Steck, and E. M. Waggoner. Phosphorus as an integral component of global marine biogeochemistry. *Nature Geoscience*, 14(6):359–368, 2021.

10. K. R. Frischkorn, S. T. Haley, and S. T. Dyhrman. Coordinated gene expression between trichodesmium and its microbiome over day–night cycles in the north pacific subtropical gyre. *The ISME journal*, 12(4):997–1007, 2018.

11. N. Gilbert, H. Pound, and S. Wilhelm. Rna extraction from sterivex using phenol:chloroform - cutting off the filter from the sterivex unit for low biomass samples. *protocols.io*, 07 2020. https://dx.doi.org/10.17504/protocols.io.bh75j9q6.

12. M. J. Harke, K. R. Frischkorn, S. T. Haley, F. O. Aylward, J. P. Zehr, and S. T. Dyhrman. Periodic and coordinated gene expression between a diazotroph and its diatom host. *The ISME journal*, 13(1):118–131, 2019.

13. S. Honjo, S. J. Manganini, R. A. Krishfield, and R. Francois. Particulate organic carbon fluxes to the ocean interior and factors controlling the biological pump: A synthesis of global sediment trap programs since 1983. *Progress in Oceanography*, 76(3):217–285, 2008.

14. J. Huerta-Cepas, K. Forslund, L. P. Coelho, D. Szklarczyk, L. J. Jensen, C. Von Mering, and P. Bork. Fast genome-wide functional annotation through orthology assignment by eggnog-mapper. *Molecular biology and evolution*, 34(8):2115–2122, 2017.

15. G. E. Hutchinson. The paradox of the plankton. *The American Naturalist*, 95(882):137–145, 1961.

16. D. M. Karl. A sea of change: biogeochemical variability in the north pacific subtropical gyre. *Ecosystems*, 2(3):181–214, 1999.

17. A. I. Krinos, S. K. Hu, N. R. Cohen, and H. Alexander. Eukulele: Taxonomic annotation of the unsung eukaryotic microbes. *arXiv preprint arXiv:2011.00089*, 2020.

18. N. Kronfeld-Schor and T. Dayan. Partitioning of time as an ecological resource. *Annual Review of Ecology, Evolution, and Systematics*, 34(1):153–181, 2003.

19. K. O. Lear, N. M. Whitney, J. J. Morris, and A. C. Gleiss. Temporal niche partitioning as a novel mechanism promoting co-existence of sympatric predators in marine systems. *Proceedings of the Royal Society B*, 288(1954):20210816, 2021.

20. L. Legendre and J. Le Fèvre. Microbial food webs and the export of biogenic carbon in oceans. *Aquatic Microbial Ecology*, 9(1):69–77, 1995.

21. D. Li, C.-M. Liu, R. Luo, K. Sadakane, and T.-W. Lam. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*, 31(10):1674–1676, 01 2015.

22. Y. Liao, G. K. Smyth, and W. Shi. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, 30(7):923–930, 11 2013.

23. M. W. Lomas, A. L. Burke, D. A. Lomas, D. W. Bell, C. Shen, S. T. Dyhrman, and J. W. Ammerman. Sargasso sea phosphorus biogeochemistry: an important role for dissolved organic phosphorus (dop). *Biogeosciences*, 7(2):695–710, 2010.

24. M. I. Love, W. Huber, and S. Anders. Moderated estimation of fold change and dispersion for rna-seq data with deseq2. *Genome Biol*, 15(12):550, 2014.

25. T. Masuda, K. Inomura, J. Mareš, T. Kodama, T. Shiozaki, T. Matsui, K. Suzuki, S. Takeda, C. Deutsch, O. Prášil, et al. Coexistence of dominant marine phytoplankton sustained by nutrient specialization. *Microbiology spectrum*, 11(4):e04000–22, 2023.

26. B. A. S. V. Mooy, G. Rocap, H. F. Fredricks, C. T. Evans, and A. H. Devol. Sulfolipids dramatically decrease phosphorus demand by picocyanobacteria in oligotrophic marine environments. *Proceedings of the National Academy of Sciences*, 103(23):8607–8612, 2006.

27. D. Muratore, A. K. Boysen, M. J. Harke, K. W. Becker, J. R. Casey, S. N. Coesel, D. R. Mende, S. T. Wilson, F. O. Aylward, J. M. Eppley, et al. Complex marine microbial communities partition metabolism of scarce resources over the diel cycle. *Nature Ecology & Evolution*, pages 1–12, 2022.

28. E. A. Ottesen, C. R. Young, J. M. Eppley, J. P. Ryan, F. P. Chavez, C. A. Scholin, and E. F. DeLong. Pattern and synchrony of gene expression among sympatric marine microbial populations. *Proceedings of the National Academy of Sciences*, 110(6):E488–E497, 2013.

29. E. A. Ottesen, C. R. Young, S. M. Gifford, J. M. Eppley, R. Marin, S. C. Schuster, C. A. Scholin, and E. F. DeLong. Multispecies diel transcriptional oscillations in open ocean heterotrophic bacterial assemblages. *Science*, 345(6193):207–212, 2014.

30. A. Shmida and S. Ellner. Coexistence of plant species with similar niches. *Vegetatio*, 58(1):29–55, 1984.

31. O. A. Sosa, T. J. Burrell, S. T. Wilson, R. K. Foreman, D. M. Karl, and D. J. Repeta. Phosphonate cycling supports methane and ethylene supersaturation in the phosphate-depleted western north atlantic ocean. *Limnology and Oceanography*, 65(10):2443–2459, 2020.

32. O. A. Sosa, J. R. Casey, D. M. Karl, and S.-J. Liu. Methylphosphonate oxidation in *Prochlorococcus* strain mit9301 supports phosphate acquisition, formate excretion, and carbon assimilation into purines. *Applied and Environmental Microbiology*, 85(13):e00289–19, 2019.

33. O. A. Sosa, D. J. Repeta, S. Ferrón, J. A. Bryant, D. R. Mende, D. Karl, E. F. DeLong, et al. Isolation and characterization of bacteria that degrade phosphonates in marine dissolved organic matter. *Frontiers in microbiology*, 8:1786, 2017.

34. C. A. Stock, J. P. Dunne, S. Fan, P. Ginoux, J. John, J. P. Krasting, C. Laufkötter, F. Paulot, and N. Zadeh. Ocean biogeochemistry in gfdl's earth system model 4.1 and its response to increasing atmospheric co2. *Journal of Advances in Modeling Earth Systems*, 12(10):e2019MS002043, 2020. e2019MS002043 2019MS002043.

35. P. F. Thaben and P. O. Westermark. Detecting rhythms in time series with RAIN. *J. Biol. Rhythm*, 29(6):391–400, 2014.

36. L. J. Ustick, A. A. Larkin, C. A. Garcia, N. S. Garcia, M. L. Brock, J. A. Lee, N. A. Wiseman, J. K. Moore, and A. C. Martiny. Metagenomic analysis reveals global-scale patterns of ocean nutrient limitation. *Science*, 372(6539):287–291, 2021.

37. B. A. Van Mooy, H. F. Fredricks, B. E. Pedler, S. T. Dyhrman, D. M. Karl, M. Koblížek, M. W. Lomas, T. J. Mincer, L. R. Moore, T. Moutin, et al. Phytoplankton in the ocean use non-phosphorus lipids in response to phosphorus scarcity. *Nature*, 458(7234):69–72, 2009.

38. A. Vislova, O. A. Sosa, J. M. Eppley, A. E. Romano, and E. F. DeLong. Diel oscillation of microbial gene transcripts declines with depth in oligotrophic ocean waters. *Frontiers in Microbiology*, 10, 2019.

39. S. T. Wilson, F. O. Aylward, F. Ribalet, B. Barone, J. R. Casey, P. E. Connell, J. M. Eppley, S. Ferron, J. N. Fitzsimmons, C. T. Hayes, et al. Coordinated regulation of growth, activity and transcription in natural populations of the unicellular nitrogen-fixing cyanobacterium crocosphaera. *Nature microbiology*, 2(9):1–9, 2017.

40. J. Wu, W. Sunda, E. A. Boyle, and D. M. Karl. Phosphate depletion in the western north atlantic ocean. *Science*, 289(5480):759–762, 2000.