1   **An ALS Assembly Modulator Signature in Peripheral Blood Mononuclear Cells: Implications**

2   **for ALS Pathophysiology, Therapeutics, and Diagnostics**

4   Shao Feng Yu[1], Maya Michon[1], Anuradha F. Lingappa[1], Kumar Paulvannan[1], Dennis Solas[1], Kim

5   Staats[2], Justin Ichida[2], Debendranath Dey[1], Jeffrey Rosenfeld[3] and Vishwanath R. Lingappa[1,4,5]

[1] Prosetta Biosciences, Inc. 670 5th St San Francisco, CA 94107

[2] University of Southern California, Los Angeles, CA 90033

[3] Loma Linda Medical Center, Loma Linda, CA 92354

[4] Department of Physiology, University of California, San Francisco, CA 94143

[5] To whom correspondence should be directed (vlingappa@prosetta.com)

Abstract

Assembly modulators are a new class of allosteric site-targeted therapeutic small molecules,

some of which are effective at restoring nuclear localization of TDP-43 in ALS cellular models,

and which display efficacy in a variety of ALS animal models. These compounds have been

shown to bind selectively to a small subset of protein disulfide isomerase (PDI), a protein

implicated in ALS pathophysiology. The targeted subset of PDI is found within a novel, transient

and energy-dependent multi-protein complex that includes other important members of the

ALS interactome, such as TDP-43, RanGTPase, and selective autophagy receptor p62/SQSTM1.

We demonstrate here that a similar multi-protein complex drug target is present in PBMCs as

isolated by energy-dependent drug resin affinity chromatography (eDRAC) and characterized by

mass spectrometry and by Western blot (WB). Signature alterations in the composition of the

multi-protein complex in PBMCs from ALS patients compared to PBMCs from healthy individuals

were identified by WB of eDRAC bound proteins, thereby extending earlier literature suggesting

PBMC dysfunction in ALS. Changes in the PBMC drug target in ALS patients compared to healthy

individuals include diminished p62/SQSTM1 and appearance of a 17kDa post-translationally

modified form of RanGTPase. These changes are not readily apparent from analysis of whole

cell extracts, as the individual protein components within the drug target multi-protein complex

comprise only small percentages of the total of those component proteins in the extract.

Furthermore, whole blood from ALS patients shows a distinctive degradation of total

RanGTPase not observed in blood from healthy individuals. This degradation appears to be

rescued by treatment of whole blood from ALS patients for 72 hrs with ALS-active assembly

modulator small molecules. Our findings are consistent with the hypothesis that ALS is

fundamentally a disorder of homeostasis that can be detected early, prior to disability, in blood

by the methods described, and restored to the healthy state by assembly modulator drug

treatment.

Introduction

Amyotrophic Lateral Sclerosis (ALS) is a devastating neurodegenerative disease that

strikes approximately 6,000 American and 150,000 people worldwide annually[1,2]. Approximately

10% of ALS is familial, with a large number of specific gene mutations implicated. The vast

majority of ALS cases however, are sporadic, meaning that that there is no family history of the

disease and therefore seems likely due to an environmental cause, perhaps with epigenetic

influence[3]. Specific environmental factors have been implicated in causing ALS, but remain

controversial and inconclusive[4]. While the pathophysiology of ALS is not well understood,

aggregates of the protein TDP-43, independently implicated in the HIV life cycle[5], are almost

invariably found in the brains of patients with ALS and fronto-temporal dementia (FTD)[6].

Despite this commonality, the clinical presentation as well as progression of ALS, is highly

variable[7].

With regards to TDP-43 pathophysiology, controversy exists as to the triggering event.

Some studies have implicated mislocalization of TDP-43 from the nucleus in healthy cells to the

cytoplasm, which is observed to variable degrees in fibroblasts from many, but not all, cases of

ALS[8]. An alternative view is that, in response to stress, TDP-43 forms aggregates in the

cytoplasm where it is synthesized, prior to transport to the nucleus[9]. A further controversy

exists as to whether the relevant TDP-43 aggregates in the cytoplasm are within, or outside of,

stress granules[10]. While the TDP-43 aggregates have generally been viewed as a trigger of motor

neuron death, it is possible that they are an epiphenomenon whose elimination *per se* may not

necessarily be therapeutic, by analogy to the failed attempts to remedy Alzheimer's Disease by

elimination of $A\beta$ aggregates[11]. Regardless of their natural history and role in the

pathophysiology of ALS, the presence of TDP-43 aggregates are pathognomonic for the

disease[12,13].

Recently we presented our studies on the role of a newly appreciated dimension of gene

expression, termed protein assembly, in the pathophysiology of diverse diseases including

ALS[14,15,16]. These compounds were found by a novel method in which viruses were used as

"trufflehounds" to reveal drug targets inaccessible by genomic or conventional proteomics[15,16].

We have identified several structurally unrelated small molecule chemotypes by virtue of their

ability to modulate host machinery catalytic for viral capsid formation, that we term protein

assembly modulators. These compounds are strikingly efficacious both for the specific viral

disease by which they were identified, and also for a subset of non-viral diseases. Among the

protein assembly modulators with activity against HIV, one prevents ALS-associated

neurodegeneration and its correlates in various cellular and animal models of ALS[14]. A subseries

of analogs shows diminished activity against HIV and improved activity against ALS. These

compounds are broadly effective at reversing both TDP-43 mislocalization and stress-induced

TDP-43 aggregation, in both familial and sporadic ALS and fronto-temporal dementia (FTD)

cellular models. Efficacy has also been demonstrated in various transgenic animals expressing

human ALS-causing mutations including in TDP-43, FUS, c9orf72, and SOD1, by a variety of

assays[14]. In view of this breadth of efficacy, we hypothesized the drug target to be a step

common to most, if not all, ALS and FTD. Advanced analogs of the ALS-selective active subseries

of one of these chemotypes are close to completion of a target product profile (TPP), prior to

initiation of investigational new drug (IND)-enabling studies in anticipation of human clinical

trials in the near future.

The studies to be described here had several motivations. First, to explore the

hypothesis that the biochemical pathways involved in ALS pathophysiology may involve layers of

regulation yet to be discovered[17,18], in which the novel multi-protein complex targets of these

compounds might be involved. How those levels of control are integrated with one another,

and are communicated from one organ system to others, remain largely unknown. It may be

that the novel assembly modulator targets we have discovered are potential candidates for

involvement in such regulation. By corollary, what happens to other feedback controls when

one level of regulation fails, remains to be determined. The argument has been made based on

gene knockout experiments that more ancient back up pathways may be activated when

primary regulatory controls fail[19,20]. Having a drug that impinges on a biochemical pathway in a

novel way is a powerful means of probing biological regulation. Much of this regulation depends

on information that does not reside simply in the sequence of the genome[18,20,21]. Thus the

relevance of the compounds we describe, with their unconventional multi-protein complex

target (energy-dependent for formation, transient in existence, composed of miniscule subsets

of the individual component proteins in the cell), and novel mechanism of action at allosteric

sites[22,23 24,25]. These tools may provide a new understanding of the molecular basis for

homeostasis, and its dysregulation, as exemplified in ALS.

A second motivation for the present studies was to understand the meaning of the

observed dysfunction of PBMCs in ALS[26,27]. These findings can be interpreted in very different

ways. On the one hand, PBMC dysfunction might be due to ALS being a systemic disease whose

most severe manifestation is motor neuron death, but which also manifests in less dramatic

fashion in other organ systems, including in blood. Alternatively, the signature observed in

PBMCs might reflect a role of blood, which sees essentially every cell in the body on a regular

basis, spreading the message that there is a disorder affecting a particular cell type, in this case,

motor neurons. Yet another possibility is that the disorder affecting one organ system (motor

neurons) could have a "domino effect" manifesting in other organ systems (PBMCs), e.g. as a

secondary consequence of feedback regulation intended to monitor and maintain

homeostasis[17,28]. Prion-type spread observed for various diseases may be an example of such

disordered communication[28–30]. To better understand these alternatives, we asked whether

assembly modulator drugs might detect a version of the disordered drug target in PBMCs from

ALS patients.

Finally, apart from intellectually interesting questions of feedback and mechanism of

action[17,28,30], there is a practical desire to advance these compounds to the clinic, given their

remarkable therapeutic properties in cellular and animal models of ALS[31]. Having a readily

accessible signature of ALS in PBMCs would facilitate ALS drug advancement in innumerable

ways, most importantly by making possible early detection of ALS (prior to severe disability).

Compounds designed to restore homeostasis would be expected to be most effective when

administered early, before motor neuron death. Or put another way, treatments that restore

homeostasis would be expected to be very different from those needed to regrow and replace

cells after they have died. A signature in PBMCs would also provide a biomarker for easy

assessment of therapeutic efficacy, and to determine when therapy could be stopped or might

need to be restarted.

Results

Blinded blood samples were obtained from ALS patients and healthy individuals, under

an IRB-approved protocol. The PBMCs were isolated, and extracts prepared as described in

methods, and analyzed for key proteins relevant to ALS. Differences observed in the protein

pattern between PBMCs of healthy and ALS patients, by total protein silver stain were subtle

(see for example, **Figure 1A**).

However when PBMC extracts were applied to the ALS-active drug resin described

previously[31] versus control resin (lacking the drug ligand), washed and bound material analyzed,

a distinctive subset of proteins was observed, also largely shared between healthy individuals

and ALS patients, but with a few possible differences (black arrows in **Figure 1B**).

Previous studies of assembly modulator small molecules[15,16] have shown that the

proteins comprising the multi-protein complex drug target represent single digit percent of the

total of those proteins in the starting extract. As expected therefore, the depleted extracts differ

very subtly from the starting material, as demonstrated for PBMC DRAC flow through here (see

also **Supplemental Figure 1D**).

Figure 1

Legend to Figure 1. A. Silver stain of starting PBMC extract 10kxg/10min supernatant from a

representative healthy individual (#52) and an ALS patient (#78-2). B. Silver stain of free drug eluate from

ALS-active assembly modulator drug resin (column 455 THIQ) or control resin (column 134 lacking the

drug ligand). Black arrows indicate positions at which healthy vs ALS patient samples show a suggestive

difference in the free drug eluate protein pattern.

Mass spectrometry analysis of free drug eluates from PBMCs applied to the ALS-active drug

resin (**Figure 2**) was carried out and compared to the pattern previous revealed from mouse

brain (MoBr). A substantial proportion of tissue-specific proteins (~1/2 to 2/3 depending on the

specific sample) were observed (see **Supplemental Figure 2**). Despite the high tissue specificity,

a core set of 33 proteins shared between both PBMCs and brain were of interest. Of these, 26

were identified as members of the ALS protein interactome. Notably, this set included P4HB, a

member of the PDI family, two known ALS-causing proteins (TUBA4A and VCP)[32], along with

proteins implicated in the KEGG ALS interactome[33] (ATP5A1, HSPA5, TUBA1A, TUBA4A, and

VCP). Furthermore, 24 of these 33 proteins exhibited high-confidence interactions with known

ALS-causing or disease-modifying proteins[32]. The 26 shared ALS interactome proteins also

included multiple members of several protein families and complexes: two ATP synthase

subunits (ATP5A1 and ATP6V1A), six chaperonins containing T-complex components (CCT2,

CCT3, CCT4, CCT5, CCT7, and CCT8), three thioredoxin domain proteins (P4HB, PRDX1, and

PRDX2), and four members of the 14-3-3 family (YWHAB, YWHAE, YWHAH, and YWHAQ) (**Figure**

2A).

MCL clustering analysis was performed on the 33 shared proteins using the STRING

database, which integrates information from various databases cataloging known and predicted

protein-protein interactions. The analysis generated groups called "clusters," characterized by a

higher density of predicted associations within the clusters compared to those between

clusters. Notably, 31 out of the 33 proteins formed a single, highly interconnected cluster,

suggesting potential functional relationships between them. Two proteins, VAT1 and ALDH1A1,

were excluded from this cluster, as there are no predicted interactions between these proteins

and the other 31 proteins (**Figure 2B**). These findings are summarized in **Figure 2C.**

Legend to Figure 2. A. Spectral counts for proteins identified in THIQ drug resin eluate from MoBr and

human PBMC extracts. Colors represent the number of spectral counts for a given protein in each

sample, ranging from light green (low counts) to dark green (high counts). The right panel indicates "hits"

for proteins present in the ALS interactome, where yellow is a hit. B. STRING database protein-protein

interaction network with MCL clustering. Each circle ("node") represents a protein, and each line

("edge") represents a predicted interaction between two nodes. Different colors represent distinct

clusters identified by the MCL algorithm, with red indicating the largest cluster and yellow and green

representing clusters containing only a single protein. C. ALS interactome proteins identified by MS-MS

are shown in venn diagram. Causative and disease modifier proteins are in **bold**.

From the MS-MS analysis it was clear that the multi-protein complex isolated from

PBMCs on the ALS-active drug resin appears related to that observed in MoBr, based on the 27

shared proteins identified that are in the ALS protein interactome observed from both sources.

However no clear differences between healthy and ALS patients were observed by MS-MS.

Recognizing that MS-MS analysis as carried out here can fail to identify specific proteins

for a number of reasons, including covalent post-translational modifications (PTMs) that alter

peptide mass, or result in failure of peptide cleavage with trypsin and/or ionization, we analyzed

the DRAC bound material by WB to identify particular proteins of interest. WB has its own

limitations, partially but not completely overlapping with those of MS-MS. A shared limitation

with MS-MS is that a covalent post-translational can mask an epitope making it undetectable by

an epitope-specific antibody. Nevertheless, if a difference between healthy and ALS patients in a

protein within the drug resin-bound target complex were identified either by WB or MS-MS, it

could serve as a signature of the disease. From WB of the DRAC bound material two proteins

were strikingly different between most healthy individuals studied and most ALS patients. First,

p62/SQSTM1, a known regulator of autophagy, an important host defensive mechanism, was

greatly diminished in ALS patient PBMC DRAC free drug eluates compared to that from most

healthy individuals. Conversely, a 17kDa post-translationally modified fragment of RanGTPase

was detected in ALS patients, but not in most healthy individuals, in DRAC free drug eluates. A

representative example is shown in **Figure 3**.

Figure 3.

Legend to Figure 3. PBMC extracts were prepared from ALS patients (ALS) or healthy individuals (H) as

described in methods and applied to drug resin versus control resin, washed and bound proteins

analyzed by SDS-PAGE transferred to 0.2u PVDF membrane and WB for the indicated proteins (below)

with quantitation of bands using Image J software (above). Shown is DRAC bound material from a

healthy control (#52) and a sporadic ALS patient (#78) analyzed by WB for p62/SQSTM1, RanGTPase, and

PDI. The highest value in each healthy/ALS graph pair is normalized to 100% for proportional

comparison to the lower value.

Out of 13 ALS patients and 6 healthy controls assessed in a blinded fashion, all 13 ALS

patients had a strikingly low values for p62 in the drug resin bound material. Of the 6 healthy

controls, 2 were low. For Ran 17 kDa fragment, all 13 ALS patients had a higher value than the

parallel reference healthy control. Among the 6 healthy controls, the two with low p62 also had

high Ran 17 kDa fragment in the eDRAC bound material. In contrast to striking loss of p62 and

appearance of RanGTPase 17kDa fragment, only a modest difference in the direct drug-binding

protein, PDI, was observed between ALS patients and healthy controls.

Thus, taken together, the eDRAC signatures observed appear to have a sensitivity

approaching 100% and a specificity of approximately 66% for ALS, albeit from a modest sample

size. Further studies are needed to understand how these differences come about and what

their significance is to the underlying pathophysiology of ALS. Nevertheless, together with the

clinical picture, they suggest a basis for early detection of ALS using as an affinity ligand, a small

molecule that is therapeutic for the disease in a diversity of animal models studied[14]. **Table 1**

summarizes these findings together with key demographic features of the individuals studied.

The eDRAC signature in PBMCs, like that observed previously in other cells and tissues,

represent very small subsets of the total of the proteins in the target multi-protein complex

(**Supplemental Figure 3**).

	% p62	% 17kDa Ran Fragment	sex	age at onset	age dx	age at presentation	date of death	disease course	sample date (mo/day/yr)
#52 (healthy)	100 (650)	1 (2)	F			41			7/31/18
#40 (ALS)	71 (462)	5 (12)	M	68	71	72	N/A	LMN predominant - early bulbar/trach slow, progressive course	1/24/19
#50 (ALS)	30 (193)	12 (32)	M	46	47	59	10/21/22	Generalized slowly progressive disease	7/31/18 1/24/19
#51 (ALS)	64 (419)	8 (22)	M	38	39	40	3/1/23	UMN predominant generalized disease with bulbar compromise	7/31/18
#68 (ALS)	3 (16)	100 (261)	F	78	79	79	N/A	primary bulbar presentation with mostly UMN features	2/19/19
#54 (ALS)	6 (41)	76 (197)	M	51	52	52	2/5/19	rapidly progressive, bulbar involving disease	8/7/18
#55 (healthy)	10 (63)	86 (224)	F			55			
#52 (healthy)	80 (645)	0 (0)	F			41			7/31/18
#45 (ALS)	13 (64)	61 (63)	M	42	43	43	N/A		5/22/18 2/28/19
#78 (ALS)	6 (35)	100 (103)	F	45	46	46	10/7/20	LMN predominant, generalized disease	5/28/19 7/18/19
#52 (healthy)	99 (700)	8 (8)	F			41			7/31/18
#87 (healthy)	100 (706)	5 (5)	M			36			12/19/19
#52 (healthy)	100 (710)	0 (0)	F			41			7/31/18
#43 (ALS)	12 (85)	100 (44)	F	68	71	72	12/19/18	UMN predominant; generalized disesease with significant bulbar compromise	12/19/18
#44 (healthy)	15 (105)	27 (12)	F			74			
#46 (ALS)	54 (385)	11 (5)	M	53	54	54	N/A	bulbar onset then generalized disease	5/23/18
#47 (healthy)	56 (400)	7 (3)	M			58			
#53 (ALS)	0 (0)	68 (30)	F	67	68	71	2/12/19	limb onset generalized disease, possible concurrent Parkinson's disease	7/31/18
#26 (healthy)	100 (558)	0 (0)	M			64			1/22/18
#34 (ALS)	1 (3)	7 (2)	M	40	42	42	N/A	monomelic presentation VERY slow LMN generalization, bulbar sparing	2/15/18
#69 (ALS)	2 (13)	100 (24)	M	80	82	82	5/8/21	LMN predominant, generalized disease	2/19/19
#70 (ALS)	13 (71)	7 (2)	M	42	42	43	N/A	generalized progressive disease	2/28/19

Legend to Table 1. 13 ALS patients and 6 healthy controls were analyzed over 4 separate experiments

and the data tabulated, together with key demographic and clinical history information. For each

experiment the highest value p62 was set at 100% and the others scaled proportionally. Likewise, in each

experiment the highest value 17kDa Ran fragment was set at 100% and the others scaled proportionally.

The quantified WB band value is also shown in parentheses. Healthy individuals are noted in green, and

ALS patients in red, except for the two healthy individuals displaying the ALS phenotype, whose values

are in black.

Next, photocrosslinking was carried out with PBMC extracts to which was added a drug

analog in which the attachment to resin was replaced by attachment to a biotin and diazirine

moiety such that exposure to UV light results in a covalent linkage to the drug binding

protein(s). Under native conditions associated proteins of the entire multi-protein complex can

be precipitated with streptavidin beads. If the sample is first denatured, then only the direct

drug-binding protein will be precipitated with streptavidin beads under the conditions

described. Figure 4 demonstrates that PDI is the direct drug-binding protein. It should be noted

that P4HB found by MS-MS is the gene name for the same PDI family member detected by the

PDI antibody by WB.

Legend to Figure 4. Quantitation of PDI in streptavidin precipitates from PBMC extracts prepared from

ALS patients (ALS) or healthy individuals (H), under native vs denatured conditions after

photocrosslinking with PAV-073 analog (THIQ) or control crosslinker lacking the drug ligand. Both control

and THIQ crosslinkers contain biotin and diazirine as described in methods and previously[31].

277   We then assessed whether the ALS patient vs healthy control correlations involving p62

278 and Ran 17kDa fragment, as shown in **Figure 3** and **Table 1**, were accentuated over time with

279 progression of ALS. As can be seen in **Figure 5**, with progression of disease, the amount of

280 PBMC p62 bound to the resin, which is much higher in healthy individuals (panel A), diminishes

281 with advancement of ALS. In parallel, the amount of the RanGTPase 17kDa fragment, which is

282 missing in PBMCs from healthy individuals, increases with ALS disease progression.

284   Notably, the diagnosis of ALS was made in pt 78 very early in her disease progression.

285 She was able to drive herself to the clinic and walk in with a cane, at the time that the initial

286 blood sample was taken, which nevertheless revealed the dual signatures of falling p62 and

287 appearance of Ran 17kDa fragment, in contrast to parallel analysis of a healthy control. As will

288 be seen these features appear to be exacerbated with disease progression.

289   **Figure 5**.

Legend to Figure 5. PBMC extracts were prepared as previously, applied to drug resins and bound proteins analyzed by WB for specific proteins of interest. Bands corresponding to the correct molecular weights were quantified. Blood samples from three ALS patients were assessed over time for both p62 and RanGTPase 17kDa fragment bound to DRAC. Timing of blood draws were 6 weeks apart for patient 78, 6 months apart for patient 50, and 9 months apart for patient 45.

If, as we suggest, ALS represents a breakdown in homeostasis where protective

mechanisms to reboot a cellular operating system are ineffective, certain predictions follow. For

one, there may be a backup mechanism activated in an attempt to correct the failure of the first

line of defense, although by definition, in patients who progress to ALS, that backup has failed

to correct the problem. By analogy, when your computer is frozen (with appearance of the

"colored wheel of doom"), simply attempting to undo the key stroke that caused it to freeze will

not be effective. Instead, it is necessary to reboot the operating system. Perhaps assembly

modulators provide precisely such a biochemical reboot of the cellular operating system. In

view of the distinctive ALS-associated appearance of a 17kDa fragment of RanGTPase, as noted

in **Figure 5** above, degradation of RanGTPase, seemed like a plausible backup feedback

response. This backup feedback response would be engaged if less draconian feedback methods

failed.

A second prediction is that if, as we hypothesize, assembly modulation is a fundamental

mechanism of regulation of gene expression, and its activation by these compounds is

therapeutic for ALS, treatment of whole blood from ALS patients with the drugs should result in

cessation of the backup RanGTPase degradation feedback loop, upon "reboot" of homeostatic

control.

To test these hypotheses, we took blinded blood samples from ALS patients and healthy

controls and treated the whole blood with two different, structurally unrelated ALS therapeutic

assembly modulator compounds versus vehicle control (DMSO). Since RanGTPase is implicated

in ALS pathophysiology and in the PBMC signature presented in Figures 3 and 5 we chose to

assess a readily measurable potential feedback response: degradation of total RanGTPase, even

though only a small subfraction of RanGTPase is hypothesized to be involved in ALS

pathophysiology. We assessed the stability of RanGTPase with and without 72 hrs of assembly

modulator treatment. After the 72 hr incubation of whole blood, RanGTPase stability in extracts

was assessed in a 1hr incubation (**Figure 6**). In some samples, RanGTPase was observed to be

stable over the course of a 1 hour incubation. These proved to be the healthy controls, when

unblinded. However in blood from all of the ALS patients, the levels of RanGTPase after the 1hr

incubation of whole blood extracts were greatly diminished, consistent with what would be

expected for degradation of all RanGTPase as an extreme feedback response to the progression

of ALS. Upon treatment with either of advanced analogs of two ALS assembly modulator

chemotypes (called 073 and 667) for 72hrs, a striking diminution of RanGTPase degradation on

subsequent 1hr extract incubation was observed, with no change in the level of RanGTPase

observed in healthy subjects.

Figure 6

Legend to Figure 6. 10mls of whole blood from ALS patients or healthy controls was collected under an

IRB-approved protocol and shipped overnight at room temperature in EDTA-anticoagulated (purple top)

tubes without any patient-specific confidential information. An aliquot of whole blood was taken,

incubated in 1% SDS for one hour at 37°C, prepared for SDS-PAGE, and set aside. Another aliquot of

whole blood was mixed 1:1 with glucose-supplemented MEM and incubated for 72hrs at 37°C in the

presence of vehicle or protein assembly modulator drugs 073 and 667, at the end of which samples were

processed as previously described, and analyzed by SDS-PAGE. Black bars refer to the sample prepared

for SDS-PAGE immediately, while gray bars refer to the samples incubated prior to preparation for SDS-

PAGE.

Table 2

Analog representative of chemotype	Molecular weight (g/mol)	log P	H-bond donors	H-bond acceptors	Topological polar surface area (Å²)	Tanimoto similarity coefficient	
PAV-073	443.543	4.8	1	5	49	<20%	Legend to Table 2. Two structurally unrelated (Tanimoto similarity coefficient <20 %) assembly modulator chemotypes used in the experiments shown in Figure 6 are compared for drug-like properties.
PAV-667	441.915	6.4	0	7	73.7	<20%	

Discussion

The studies presented here demonstrate the presence, by energy-dependent DRAC and

photocrosslinking, of a multi-protein complex in PBMCs that is related to that observed

previously by the same methods in Tg ALS SOD G93A model MoBr and ALS patient-derived

fibroblasts (PDFs) that display both TDP-43 mislocalization and stress-induced agregation[14]. As

observed previously for the drug target in MoBr and PDFs, the multi-protein complex drug

target in PBMCs has a small subset of PDI as the direct drug-binding protein. The complex

present in ALS patient PBMCs also shares with the target in MoBr the presence of 26 proteins

implicated in the ALS interactome (Figure 2C). The precise role of this multi-protein complex

drug target in ALS pathophysiology remains to be determined. However, based on the efficacy

of structurally unrelated ALS assembly modulators in various cellular and animal models, we

hypothesize that the drug target serves a role in maintenance of homeostasis, whose loss

results in ALS. The presence of p62 in the complex in PBMCs from healthy individuals, and its

striking diminution in ALS patient PBMCs, exacerbated with disease progression (Figure 5), are

consistent with this hypothesis.

Likewise, the appearance of a 17kDa post-translationally modified form of RanGTPase in

the drug target isolated from ALS patient PBMCs (Figure 3 and Table 1) suggests a complex

biology connected to TDP-43 mislocalization, in view of the crucial role of RanGTPase in nucleo-

cytoplasmic transport. These changes appear very early in ALS, as suggested by their detection

in patient #78 prior to onset of significant disability (Figure 4). In the three cases where a

second sample was assessed 6 weeks to 9 months after the first blood draw, both signatures

were seen to be exacerbated with disease progression. These signatures had a sensitivity of

100%, detected in PBMCs of all 13 sporadic ALS patients studied, and a specificity of 66% for

distinguishing ALS patients from healthy individuals. While a larger number of cases will need to

be assessed, this early degree of discrimination, using as an affinity ligand a drug shown

therapeutically active in a diversity of animal models, is promising as a novel path to an ALS

biomarker and highly relevant to ALS therapeutics.

This target multi-protein complex is implicated in ALS for four reasons. First, the drug

that served as the ligand to bind and isolate the target is therapeutic in a wide range of ALS

cellular and animal models[31]. Second, the composition of the target multi-protein complex

includes a number of proteins of the ALS interactome, 26 of which are shared with the

corresponding complex in MoBr. Third, the isolated target from ALS patient PBMCs is altered in

protein composition compared to that from most healthy controls by two metrics: diminution of

p62/SQSTM1 and appearance of 17kDa fragment of Ran GTPase. Finally, separate from the

specific target in PBMCs, a profound degradation of total RanGTPase is observed in whole blood

from ALS patients but not in healthy controls, and is substantially rescued after 72hrs of

treatment with ALS-active assembly modulators.

We interpret these findings to suggest that ALS is, fundamentally, a disease of disrupted

homeostasis, for which assembly modulators may be therapeutic. By analogy to another

therapeutic area where assembly modulation treatment may be effective[16], it is notable that

the vast majority of individuals don't have cancer for most of their lives. However, a small

number will develop cancer, at least in part due to failure of homeostatic host defenses. So also

host defenses protect most of us most of the time from the consequences of various central

nervous system (CNS) insults, including those causing ALS. Host defenses, including restoration

of homeostasis, usually result in resolution of a CNS insult. Sometimes however, those host

defenses fail. As a result, homeostasis is not restored, and progression to motor neuron death is

accelerated. It seems reasonable to hypothesize that when initial feedback loops fail to restore

homeostasis, more and more severe feedback responses are triggered. Thus, mass degradation

of RanGTPase, as observed here in whole blood from ALS patients (Figure 6), may represent a

final desperate attempt to restore defective nucleocytoplasmic transport that results in TDP-43

mislocalization. Autophagy, in part mediated by p62/SQSTM1, is an example of a biochemical

reboot of the cellular operating system. When the reboot fails, e.g. in part due to loss of p62

from the dysfunctional target, the consequence in a subset of cases may be ALS. This persistent

deviation from homeostasis is a stress to which motor neurons appear highly sensitive, and

eventually, succumb. A drug capable of restoring homeostasis is likely most effective before

motor neurons have died, although perhaps synaptic plasticity could synergize allowing some

degree of recovery, once homeostasis has been restored. Regardless, it is surely a truism of

medicine that the earlier a disease is detected, the more effective most therapeutic

interventions are likely to be. Hence the importance of early detection to enable early

treatment, which the PBMC signatures noted here may provide.

The results of the whole blood experiments (Figure 6) are consistent with the hypothesis

that the degradation of total RanGTPase observed, is an extreme feedback response to the

inability more proximal and focused feedback loops to restore the target complex, comprised of

a small subset of RanGTPase, back to the healthy state. Treatment with either of two

structurally unrelated assembly modulators results in correction of the RanGTPase degradation

observed in whole blood, to a significant extent. We hypothesize that treatment for a longer

period would have completely restored RanGTPase levels to normal. Whether assembly

modulator drug treatment serves to fully reboot the system, e.g. restore p62 to the target

complex as has been observed for assembly modulation using a structurally unrelated molecule

in a different therapeutic area[15], remains to be determined and may vary on a case-by-case

basis. Likewise, further studies are needed to determine whether periodic or ongoing drug

treatment are required. Regardless, drug-induced changes in target composition, as implied by

stabilization of RanGTPase in ALS patient blood treated with assembly modulators, is consistent

with restoration of homeostasis. Whether this is reflective of concomitant changes in motor

neurons that prevent further motor neuron loss, remains to be investigated.

Our findings are significant for a number of reasons. First, these studies provide a

window into novel regulatory feedback loops involved in cellular homeostasis. The precise

relationship of the target multi-protein complex in healthy individuals vs ALS patients remains

to be more fully understood. One hypothesis is that the target multi-protein complex in healthy

individuals is the "normal" assembly machine involved in integration of diverse events of

homeostasis. These may include TDP-43 localization, in which RanGTPase likely plays a critical

role, perhaps along with C9orf72 and other ALS-implicated gene products. Other relevant

pathways, including autophagic removal of dysfunctional multi-protein complexes, are naturally

integrated into the physiological feedback that collectively bring about homeostasis. ALS may be

an example where a "stuck" pathway triggers more and more desperate feedback attempts to

reboot the system without success, culminating in the mass degradation of RanGTPase

observed in the whole blood experiment shown in Figure 6. In this model, the assembly

modulator compound action can be thought of as serving as an external rather than internal

reboot of the cellular operating system. A future prediction to be tested, is that longer-term

treatment of blood results in complete normalization of RanGTPase – and restoration of

p62/SQSTM1 -- in the target complex present in PBMCs. By virtue of having therapeutic small

molecules as affinity ligands for target identification, the pathways discovered can be

productively illuminated by future studies, including by fractionation and reconstitution of

functional activities in extracts.

Second, these data, together with the studies shown previously, are consistent with the

hypothesis that, despite the heterogeneity of ALS clinical presentation and time course of

progression, there is a common underlying pathway that is the basis of the disease – and that

assembly modulators are therapeutic at that underlying shared level. While we hypothesize that

restoration of autophagy is a crucial part of this pathway, re-establishment of homeostasis likely

involves far more than just restoration of autophagy. In this regards the involvement of

miniscule subsets of the specific gene products found together in the target multi-protein

complex and the role of protein degradation, is notable. These targets could not have been

detected by powerful molecular biological tools such as CRISPR or siRNA knockdown studies,

because of the heterogeneous roles of small subsets of many, if not most, of the proteins

comprising the target multi-protein complex. Thus the concept of protein "moonlighting"[34,35]

may be more pervasive than is currently generally recognized. Conversely, the level of

organization of gene expression accessed by protein assembly modulation may be therapeutic

for diverse diseases in ways not generally appreciated today.

Finally, it should be noted that many if not most patients come to medical attention long

after the most effective period of treatment is lost. Drugs to stop the disease process may still

be effective, but rely on maintenance of the underlying framework of feedback loops to restore

homeostasis, once disease progression is arrested. Sometimes however, those feedback loops

fail and homeostasis is not restored, whether or not the primary disease process has been

blocked. We hypothesize ALS is one such example, making it extremely difficult to treat. The

methods described here provide a means of detecting ALS early, prior to onset of severe

disability, when it is most treatable – and provide a potentially effective treatment in the form

of the small molecule ligands themselves, that were used for disease detection. It is notable

that pt 78 came to medical attention early in her clinical course, prior to serious disability, and

yet, the signature changes (lowered p62 and elevated RanGTPase 17kDa fragment) were

nevertheless observed, and sadly, progressed with worsening of ALS, culminating in her death,

approximately 2 years after initial detection of the Ran 17kDa fragment as shown in Figure 5,

together with the reciprocal change in p62/SQSTM1, seen exacerbated over time. To date, these

findings have been observed in 13 of 13 ALS patients and are lacking in 4 of 6 healthy controls,

by the methods described. While the specificity of these signatures are relatively low, seen in 2

of 6 healthy controls, the high sensitivity implied by detection in 13 of 13 ALS patients, together

with the clinical history, make this diagnostic approach promising. PBMCs are a readily

accessible tissue. Used as affinity ligands, these small molecules could enable early diagnosis of

ALS, prior to extensive motor neuron death, when treatment would be more effective.

Remarkably, the same compounds, or their close analogs, may prove to be effective ALS

therapeutics.

Methods

Preparation of PBMC extracts[36]

Whole blood collected in purple-top EDTA tubes from healthy individuals or ALS patients under

Loma Linda Medical Center IRB# 5160136 was shipped at room temperature by overnight

courier to the San Francisco lab where it was diluted 1:1 with phosphate-buffered saline and

layered on ficoll and centrifuged for 15 minutes at 1,000 xg at room temperature with brake

off. PBMCs were collected at the interface, diluted with PBS 1:1 and centrifuged at 600 xg /15

minutes. The supernatant was aspirated and the pellet dissolved in p-body buffer (PBB)

consisting of 10mM Tris pH 7.5 ,10mM NaCl, 6mM MgAc, 1mM EDTA, and 0.35% Triton-X-100

and centrifuged at 4oC 10,000xg/10min. The 10kxg supernate was collected, aliquoted and

flash frozen in liquid nitrogen and stored at -80^{o}C until use.

Assembly modulator compounds

Each assembly modulator chemotype is structurally quite different from one another,

depending on the target to which it is directed, the allosteric site involved, etc. The two ALS-

active assembly modulator chemotypes used here, typified by compounds PAV-073 and PAV-

667 are compared for drug-like properties in Table 2. The structure of PAV-073, a member of

the THIQ chemotype, and its modification to allow its use as an affinity ligand for drug resin

affinity chromatography or photocrosslinking, is described in reference 14.

Drug Resin affinity chromatography

A variation on the theme of methods described in previous publications was used here[14–16]. For

brain from wildtype or SOD1 mutant animals, tissue was homogenized in cold low salt buffer

(10mM HEPES pH 7.6, 10mM NaCl, 1mM MgAc with 0.35% Tritonx100) then centrifuged at

10,000 xg for 10 minutes at 4°C. The post-mitochondrial supernatant was adjusted to a

concentration of approximately 10 mg/ml and equilibrated in a physiologic column buffer (50

mM HEPES ph 7.6, 100 mM KAc, 6 mM MgAc, 1 mM EDTA, 4mM TGA). For DRAC of both MoBr

and human PBMC extracts, the 10k sup samples being compared were first normalized either by

absorbance at 280nm or by Coomassie brilliant blue or silver staining pattern. Samples were

then supplemented with an energy cocktail (to a final concentration of 1mM rATP, 1mM rGTP,

1mM rCTP, 1mM rUTP, and 5 μg/mL creatine kinase) in the presence of physiological salts

(50mM HEPES pH 7.6, 100mM Kac, MgAc 5mM, 1mM DTT) and incubated for one hour at either

4° C or 22° C degrees on 30 μl of affigel resin coupled to THIQ compound or a 4% agarose matrix

(control). The input material was collected after flow through the column and the resin was

then washed with 3 ml column buffer. The resins were eluted for 2 hours then overnight at 22°C

then 4°C in 100μl column buffer containing 100μM of the cognate compound. Eluates were run

on WB or sent for MS-MS for analysis.

Chemical photocrosslinking

A variation of protocols as published previously[14] was used. Extract from MoBr were prepared

as described above then adjusted to a protein concentration of approximately 3 mg/ml in

column buffer (Hepes pH 7.5 50mM, Kac 100mM, MgAc 5mM, DTT or TGA 1mM) containing

0.1% triton. 1% DMSO or 100µM PAV-073 was added to 6µl of extract, then 3µM of PAV-073

photocrosslinker or a negative control crosslinker (comprising of the biotin and diazirine

moieties without compound) were added. The extract was incubated for 20 minutes then

exposed to UV at 365nM wavelength for 10 minutes then left on ice for one hour. After

crosslinking, samples were divided in two 20 µl aliquots and one set was denatured by adding

20 µL of column buffer 4µl of 10% SDS, 0.5 µl 1M dithiothreitol (DTT), and heating to 100^oC for 5

minutes. Both native and denatured aliquots were then diluted in 800 µl column buffer

containing 0.1% triton. 5 µl of magnetic streptavidin beads (Pierce) were added to all samples

and mixed for one hour at room temperature to capture all biotinylated proteins and co-

associated proteins. Samples were placed on a magnetic rack to hold the beads in placed and

washed three times with 800 µl of column buffer containing 0.1% triton. After washing, beads

were resuspended in 80 µl of gel loading buffer containing SDS and analyzed by SDS-PAGE

followed by transfer to PVDF membrane for WB.

WB Procedure

SDS-PAGE gels were transferred in Towbin buffer (25mM Tris, 192mM glycine, 20% w/v

methanol) to polyvinylidene fluoride membrane, blocked in 1% bovine serum albumin (BSA) in

PBS, incubated overnight at 4^oC in a 1:1 dilution of 100µg/mL affinity-purified primary rabbit

IGG to PDI (Abcam 96744) in 1% BSA in PBS containing 0.1% Tween-20 (PBST). Membranes were

then washed twice in PBST and incubated for two hours at room temperature in a 1:5,000

dilution of secondary anti-rabbit or anti-mouse antibody coupled to alkaline phosphatase in

PBST. Membranes were washed two more times in PBST then incubated in a developer solution

prepared from 100 µL of 7.5 mg/mL 5-bromo-4-chloro-3-indolyl phosphate dissolved in 60%

dimethyl formamide (DMF) in water and 100µl of 15 mg/ml nitro blue tetrazolium dissolved in

70% DMF in water, adjusted to 50mL with 0.1 Tris (pH 9.5) and 0.1 mM magnesium chloride.

Membranes were scanned and the integrated density of protein band was measured on ImageJ.

Averages and the standard deviation between repeated experiments were calculated and

plotted on Microsoft Excel. Primary antibodies for RanGTPase (Abcam ab31118), p62/SQSTM1

(Abcam, ab56416), were used at 1:1,000 dilution by the same procedure. Secondary goat

antibodies to mouse (Abcam ab97020) or rabbit (Abcam ab6722) were used at 1:5,000 dilution.

Tandem mass spectrometry

Samples were processed by SDS PAGE using a 10% Bis-tris NuPAGE gel with the 2-(N-

morpholino)ethanesulfonic acid buffer system. The mobility region, including all proteins, was

excised and washed with 25 mM ammonium bicarbonate followed by 15mM acetonitrile.

Samples were reduced with 10 mM dithoithreitol and 60^o C followed by alkylation with 50 mM

iodoacetamide at room temperature. Samples were then digested with trypsin (Promega)

overnight (18 hours) at 37^o C then quenched with formic acid and desalted using an Empore SD

plate. Half of each digested sample was analyzed by LC-MS/MS with a Waters NanoAcquity

HPLC system interfaced to a ThermoFisher Q Exactive. Peptides were loaded on a trapping

column and eluted over a 75 µM analytical column at 350 nL/min packed with Luna C18 resin

(Phenomenex). The mass spectrometer was operated in a data dependent mode, with the

Oribtrap operating at 60,000 FWHM and 15,000 FWHM for MS and MS/MS respectively. The

fifteen most abundant ions were selected for MS/MS.

Data was searched using a local copy of Mascot (Matrix Science) with the following parameters:

Enzyme: Trypsin/P; Database: SwissProt Human (conducted forward and reverse plus common

contaminants); Fixed modification: Carbamidomethyl (C) Variable modifications: Oxidation (M),

Acetyl (N-term), Pyro-Glu (N-term Q), Deamidation (N/Q) Mass values: Monoisotopic; Peptide

Mass Tolerance: 10 ppm; Fragment Mass Tolerance: 0.02 Da; Max Missed Cleavages: 2. The data

was analyzed by spectral count methods.

Diagnosis of ALS. A diagnosis of ALS was made in all patients by a combination of their

neurological exam and their electrodiagnostic testing. The examination was required to reveal

signs of upper and lower motor neuron pathology. Although the phenotypic spectrum of ALS is

quite variable, upper and lower motor neuron signs were represented in all patients and a brief

phenotypic description is included in Table 1.   Electrodiagnostic data included multifocal

evidence of denervation and reduced motor neuron recruitment.

Whole blood incubation and analysis. 100μl aliquots of EDTA-anticoagulated whole blood is

dispensed into 24 well plates containing 100μl of room temperature 80% PBS with 10mM

glucose and then treated with DMSO to 0.01% or drug in DMSO to the same dilution with final

concentration of 5μM drug. After 72hrs samples are collected and adjusted to 1% SDS. One

aliquot was prepared in gel loading buffer as starting material. Another aliquot added to energy

cocktail (1mM rNTPs with 0.2µg/ml CK, 50mM Hepes pH 7.6 100mK KAc, 6mM MgAc and 4mM

TGA) at total volume 100µl and incubated at 37°C for 1hr, and then prepared in gel loading

buffer. Samples were analyzed by SDS-PAGE and WB for full length RanGTPase.

Statistics

Graphs were generated and statistical analysis were preformed using GraphPad prism version

10.4.1. For Figure 3, unpaired t-tests were conducted to assess statistical significance in the

amount of protein bound to the THIQ resin for each starting material. In figure 6, a 2way ANOVA

was used, followed by Tukey's multiple comparisons tests. P-values of selected comparisons

were shown on the graph, including changes in RanGTPase levels before and after treatment

within each condition and differences in RanGTPase levels between DMSO and drug treatments

post-treatment. Statistical significance is indicated as follows: nsa ($P > 0.05$), * ($P \leq 0.05$), ** ($P \leq$

0.01), *** ($P \leq 0.001$), **** ($P \leq 0.0001$).

Ethical Approval

Whole blood collected in purple-top EDTA tubes from healthy individuals or ALS patients under

Loma Linda Medical Center IRB# 5160136

References

1. Al-Chalabi, A. & Hardiman, O. The epidemiology of ALS: a conspiracy of genes, environment

and time. *Nat. Rev. Neurol.* **9**, 617–628 (2013).

2. Chiò, A. *et al.* Prognostic factors in ALS: A critical review. *Amyotroph. Lateral Scler. Off. Publ.*

World Fed. Neurol. Res. Group Mot. Neuron Dis. **10**, 310–323 (2009).

3. Rizea, R. E., Corlatescu, A.-D., Costin, H. P., Dumitru, A. & Ciurea, A. V. Understanding

Amyotrophic Lateral Sclerosis: Pathophysiology, Diagnosis, and Therapeutic Advances. *Int. J.*

Mol. Sci. **25**, 9966 (2024).

4. Hernan-Godoy, M. & Rouaux, C. From Environment to Gene Expression: Epigenetic

Methylations and One-Carbon Metabolism in Amyotrophic Lateral Sclerosis. *Cells* **13**, 967

(2024).

5. Cabrera-Rodríguez, R. *et al.* TDP-43 Controls HIV-1 Viral Production and Virus Infectiveness.

Int. J. Mol. Sci. **24**, 7658 (2023).

6. Gandhi, J. *et al.* Protein misfolding and aggregation in neurodegenerative diseases: a review

of pathogeneses, novel detection strategies, and potential therapeutics. *Rev. Neurosci.* **30**,

339–358 (2019).

7. Grad, L. I., Rouleau, G. A., Ravits, J. & Cashman, N. R. Clinical Spectrum of Amyotrophic

Lateral Sclerosis (ALS). *Cold Spring Harb. Perspect. Med.* **7**, (2017).

8. Chou, C.-C. *et al.* TDP-43 pathology disrupts nuclear pore complexes and nucleocytoplasmic

transport in ALS/FTD. *Nat. Neurosci.* **21**, 228–239 (2018).

9. Baradaran-Heravi, Y., Van Broeckhoven, C. & van der Zee, J. Stress granule mediated protein

aggregation and underlying gene defects in the FTD-ALS spectrum. *Neurobiol. Dis.* **134**,

104639 (2020).

10. Gasset-Rosa, F. *et al.* Cytoplasmic TDP-43 De-mixing Independent of Stress Granules Drives

Inhibition of Nuclear Import, Loss of Nuclear TDP-43, and Cell Death. *Neuron* **102**, 339-

357.e7 (2019).

11. Kim, C. K. *et al.* Alzheimer's Disease: Key Insights from Two Decades of Clinical Trial Failures.

J. Alzheimers Dis. JAD **87**, 83–100 (2022).

12. Aggad, D., Vérièpe, J., Tauffenberger, A. & Parker, J. A. TDP-43 toxicity proceeds via calcium

dysregulation and necrosis in aging Caenorhabditis elegans motor neurons. *J. Neurosci. Off.*

J. Soc. Neurosci. **34**, 12093–12103 (2014).

13. Buratti, E. & Baralle, F. E. Multiple roles of TDP-43 in gene expression, splicing regulation,

and human disease. *Front. Biosci. J. Virtual Libr.* **13**, 867–878 (2008).

14. Yu, S. F. *et al.* Protein Assembly Modulation: A New Approach to Amyotrophic Lateral

Sclerosis (ALS) Therapeutics. *J. Exp. Neurol.* **5**, 210–230 (2024).

15. Michon, M. *et al.* A pan-respiratory antiviral chemotype targeting a transient host multi-

protein complex. *Open Biol.* **14**, 230363 (2024).

16. Lingappa, A. F. *et al.* Small molecule protein assembly modulators with pan-cancer

therapeutic efficacy. *Open Biol.* **14**, 240210 (2024).

17. Bechtel, W. & Bich, L. Situating homeostasis in organisms: maintaining organization through

time. *J. Physiol.* **602**, 6003–6020 (2024).

18. Tomkins, G. M. The metabolic code. *Science* **189**, 760–763 (1975).

19. Marty, L. *et al.* The NADPH-dependent thioredoxin system constitutes a functional backup

for cytosolic glutathione reductase in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 9109–

9114 (2009).

20. Noble, D. From Genes to Whole Organs: Connecting Biochemistry to Physiology. in *Novartis*

Foundation Symposia (eds. Bock, G. R. & Goode, J. A.) vol. 239 111–128 (Wiley, 2001).

21. Pardee, A. B. Regulatory Molecular Biology. *Cell Cycle* **5**, 846–852 (2006).

22. Pardee, A. B. & Reddy, G. P.-V. Beginnings of feedback inhibition, allostery, and multi-protein

complexes. *Gene* **321**, 17–23 (2003).

23. Fenton, A. W. Allostery: an illustrated definition for the 'second secret of life'. *Trends*

Biochem. Sci. **33**, 420–425 (2008).

24. Müller-Schiffmann, A. *et al. A Pan-Respiratory Antiviral Chemotype Targeting a Transient*

Host Multiprotein Complex. http://biorxiv.org/lookup/doi/10.1101/2021.01.17.426875

(2021) doi:10.1101/2021.01.17.426875.

25. Lingappa, A. F. *et al.* Small Molecule Assembly Modulators with Pan-Cancer Therapeutic

Efficacy. *bioRxiv* 2022.09.28.509937 (2022) doi:10.1101/2022.09.28.509937.

26. Nardo, G. *et al.* Amyotrophic lateral sclerosis multiprotein biomarkers in peripheral blood

mononuclear cells. *PloS One* **6**, e25545 (2011).

27. Luotti, S. *et al.* Diagnostic and prognostic values of PBMC proteins in amyotrophic lateral

sclerosis. *Neurobiol. Dis.* **139**, 104815 (2020).

28. Pongrácová, E., Buratti, E. & Romano, M. Prion-like Spreading of Disease in TDP-43

Proteinopathies. *Brain Sci.* **14**, 1132 (2024).

29. Duyckaerts, C., Clavaguera, F. & Potier, M.-C. The prion-like propagation hypothesis in

Alzheimer's and Parkinson's disease. *Curr. Opin. Neurol.* **32**, 266–271 (2019).

30. Wenzhi, Y., Xiangyi, L. & Dongsheng, F. The prion-like effect and prion-like protein targeting

strategy in amyotrophic lateral sclerosis. *Heliyon* **10**, e34963 (2024).

31. Feng Yu, S. *et al. Protein Assembly Modulation: A New Approach to ALS Therapeutics.*

http://biorxiv.org/lookup/doi/10.1101/2023.07.24.550252 (2023)

doi:10.1101/2023.07.24.550252.

32. Dervishi, I. *et al.* Protein-protein interactions reveal key canonical pathways, upstream

regulators, interactome domains, and novel targets in ALS. *Sci. Rep.* **8**, 14732 (2018).

33. Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M. & Ishiguro-Watanabe, M. KEGG for

taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res.* **51**, D587–D592

(2023).

34. Copley, S. D. Moonlighting is mainstream: paradigm adjustment required. *BioEssays News*

Rev. Mol. Cell. Dev. Biol. **34**, 578–588 (2012).

35. Jeffery, C. J. Multitalented actors inside and outside the cell: recent discoveries add to the

number of moonlighting proteins. *Biochem. Soc. Trans.* **47**, 1941–1948 (2019).

36. Jia, Y. *et al.* A Modified Ficoll-Paque Gradient Method for Isolating Mononuclear Cells from

the Peripheral and Umbilical Cord Blood of Humans for Biobanks and Clinical Laboratories.

Biopreservation Biobanking **16**, 82–91 (2018).

Supplemental Figure 1.

Legend to Supplemental Figure 1. Input extracts for Figures 3-6. Silver stain (left and middle panels) and CBB stain (right panel) of input extracts used for DRAC (Figure 3 and 5), photocrosslinking (Figure 4), and RanGTPase degradation (Figure 6) assays.

Supplemental Figure 2.

Legend to Supplemental Figure 2. Spectral counts for entire dataset of MoBr and PBMC THIQ resin eluates. Colors represent the number of spectral counts for a given protein in each sample, ranging from light green (low counts) to dark green (high counts). The right panel indicates whether a given protein was in one of both of the datasets, according to threshold values of >4 spectral counts in the drug eluate and <4 spectral counts in the control column eluate.

 Supplemental Figure 3.

Legend to Supplemental Figure 3. Confirmation that only a small subset of PDI is present in the DRAC-

identified target and that very little additional pre-existing PDI is competent for formation of new target

without *de novo* synthesis. PBMC extracts were prepared as previously, applied to drug resins and bound

proteins analyzed by WB for PDI. The band corresponding to the correct molecular weight was

quantified as shown. Samples from three ALS patients and one healthy control are shown. Column 459

represents the THIQ drug resin and column 134 the control drug resin. As can be seen, approximately

10% of PDI is in the complex bound to the drug resin in all cases, with no significant binding to the

control resin in any case. The flowthrough from the first column (drug or control) was applied to a

second copy of the drug resin, demonstrating substantial depletion from the first column and that,

despite losses, a substantial fraction of the flow through from the control column remains competent to

bind to the drug resin column.

Abbreviations

ALS – Amyotrophic Lateral Sclerosis

BSA – bovine serum albumin

CFPSA – Cell-free protein synthesis and assembly

Col – column

Ctrl - control

CNS – Central Nervous System

DTT - dithiothreitol

DRAC – drug resin affinity chromatography

Dx - diagnosis

eDRAC – energy-dependent drug resin affinity chromatography

FTD – Fronto-Temporal Dementia

H – healthy individual

HEPES – 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

IND – investigational new drug

LMN – lower moter neuron

MoBr – mouse brain

MS-MS – tandem mass spectrometry

ns – non-significant

PBMC – Peripheral blood mononuclear cell

PBS – phosphate-buffered saline

PDF – patient-derived fibroblast

PDI – protein disulfide isomerase

PTM – post-translationally modified

PSMW – Prestained molecular weight markers

PVDF - polyvinylidene difluoride

SAP – streptavidin precipitation

SDS-PAGE – polyacrylamide gel electrophoresis in sodium dodecyl sulfate

TDP-43 – Transactive Response DNA-binding Protein 43

TGA – thioglycolic acid

TPP – target product profile

UMN – upper motor neuron

WB – western blot

*   - p value < 0.05

**   - p value < 0.01

***   - p value < 0.001

****   - p value < 0.0001

Dataset

The relevant dataset has been submitted to the DRYAD database and can be accessed upon

publication at the following: DOI: 10.5061/dryad.0rxwdbsbm

Competing Interests

765   VRL is CEO and AFL COO of Prosetta Biosciences

Funding Statement

768   This work was supported by Target ALS (grant# IL-2023-C4-L1), DOD CDMRP (grant #.

769   W81XWH2210721), and Prosetta Biosciences.

Acknowledgements

772   We thank David Hanzel and the anonymous reviewers for careful reading of the manuscript and

773   numerous helpful suggestions.