GENOME TARGETED ENRICHMENT AND SEQUENCING OF HUMAN-

INFECTING *CRYPTOSPORIDIUM* spp.

Bayona-Vásquez NJ[1*], Sullivan AH[2,3], Beaudry MS[2,4], Khan A[5], Baptista RP[6,7], Petersen

KN[8], Bhuiyan MIU[2], Brunelle B[4], Robinson G[9,10], Chalmers RM[9,10], Alves-Ferreira

EVC[11], Grigg ME[11], Kissinger JC[3,12,13], Glenn TC[2,3,12].

[1]Division of Natural Science and Mathematics, Oxford College of Emory University,

Oxford, GA, 30054, USA

[2]Department of Environmental Health Science, College of Public Health, University of

Georgia, Athens, GA, 30602, USA

[3]Institute of Bioinformatics, University of Georgia, Athens, GA, 30602, USA

[4]Daicel Arbor Biosciences, Ann Arbor, MI, 48103, USA

[5]Animal Parasitic Disease Laboratory, Agricultural Research Service, US Department of

Agriculture, Beltsville, MD, 20705, USA

[6]Center for Infectious Diseases, Houston Methodist Research Institute, Houston, TX,

77030, USA

[7]Department of Medicine, Weill Cornell Medical College, New York, USA

[8]Odum School of Ecology, University of Georgia, University of Georgia, Athens, GA,

30602, USA

[9]*Cryptosporidium* Reference Unit, Public Health Wales, Microbiology and Health

Protection, Singleton Hospital, Swansea SA2 8QA, UK

[10]Swansea University Medical School, Singleton Park, Swansea, SA2 8PP, UK

[11]Laboratory of Parasitic Diseases, National Institutes of Allergy and Infectious Diseases,

National Institutes of Health, Bethesda, MD, 20892, USA

[12]Department of Genetics, University of Georgia, Athens, GA, 30602, USA

[13]Center for Tropical and Emerging Global Diseases, University of Georgia, Athens, GA,

30602, USA

*Corresponding author

Bayona-Vásquez and Sullivan contributed equally.

Abstract

Cryptosporidium spp. are parasites that cause severe illness in vulnerable human

populations. Obtaining pure and sufficient *Cryptosporidium* DNA from clinical and

environmental samples is a challenging task. Oocysts shed in available fecal samples can

be limited in quantity, require purification (biased towards dominant strains), and yield

limited DNA (<40 fg/oocyst). Here, we use updated genomic sequences from a broad

diversity of human-infecting *Cryptosporidium* species (*C. cuniculus*, *C. hominis*, *C.*

meleagridis, *C. parvum*, *C. tyzzeri*, and *C. viatorum*) to develop and validate a set of

100,000 RNA baits (CryptoCap_100k) with the aim of enriching *Cryptosporidium* spp.

DNA from varied samples. Compared to unenriched libraries, CryptoCap_100k increases

the percentage of reads mapping to target genome sequences, increases the depth and

breadth of genome coverage and the reliability of detecting species and mixed infections

within a sample, and allows assessment of genetic variation via SNP calling, while

decreasing costs.

Introduction

Cryptosporidiosis, a disease characterized by mild to severe gastrointestinal symptoms, is caused by protist parasites in the genus *Cryptosporidium*. Infection can lead to adverse health outcomes in vulnerable human populations (infants, the elderly, immunocompromised, and pregnant)[1–6]. An estimated 7.5M global cases of cryptosporidiosis occur in children aged 0–24 months, and ~133,422 deaths occur annually[7–9]. In children under five years, cryptosporidiosis is the fourth leading cause of diarrheal-related deaths globally[6,10,11]. In the United States, *Cryptosporidium* is the third leading cause of zoonotic gastroenteritis and the leading cause of waterborne disease outbreaks[12]. In low- and middle-income countries, *Cryptosporidium* spp. is the second most common pathogen in infants from 0–11 months[10].

There are at least 39 recognized species within the genus *Cryptosporidium*[13,14], with a wide range of host specificity and likelihood of infecting humans. Thus, *Cryptosporidium* spp. vary in public health importance[13]. At the genome sequence level, a cluster of species shares greater than 80% identity across much of their genomes[15–18], including *C. cuniculus*, *C. hominis*, *C. meleagridis*, *C. parvum*, *C. tyzzeri,* and *C. viatorum*.

Several molecular methods have been developed to gain insight into the genetic diversity of *Cryptosporidium*, often relying on single-marker typing of conserved (e.g., rRNA-SSU) or polymorphic genes (e.g., *gp60* or *hsp70*)[2,19]. Other approaches include multi-locus sequence typing (MLST)[20–23] and, more recently, whole-genome sequencing (WGS). The disadvantages of MLST include unreliability of markers in different regions due to limited information about the genetic variability of *Cryptosporidium*[23], and thus, assays are not universal. Challenges with methods such as WGS are constrained by the

need to purify oocysts[24], given the limited amount of target DNA from fecal clinical

samples (which also have an abundance of other organisms' DNA) [25,26]. These challenges

have led to the use of purification methods combined with costly deep sequencing[27,28].

For example, to obtain sufficient DNA, oocysts may be propagated through animals,

which has been shown to change *Cryptosporidium* populations through selection of

individual subtypes or recombination[14,29]. Some WGS analyses have relied on whole-

genome amplification protocols to overcome low concentrations of target DNA[14,25,26,30].

A helpful alternative that still requires semi-purified oocyst suspensions[4,14,31], and that

may be prone to uneven amplification and the production of chimeric DNA fragments.

Other promising protocols for WGS directly from human stool samples do not require

whole-genome amplification but rely on a sufficient sample volume and an adequate

concentration of oocysts, achieving WGS data for ~53% of samples[32], showing medium

success rates. Thus, due to the limitations of existing approaches, additional methods that

can characterize *Cryptosporidium* genomes from limited amounts of DNA, if possible,

directly from fecal samples, which can be implemented easily and reliably, would be

beneficial.

Host-derived and environmental samples can also contain more than one species or

genotype of *Cryptosporidium* spp.[33–38]. Mixed infections may be limited to specific

geographic areas, particularly in endemic regions[39]. Most reported human mixed-species

infections contain *C. parvum* and *C. hominis*[34,36,40]; still, mixed infections are more

commonly represented by multiple strains of a single species[16]. Their prevalence has

been reported to range from 5% in the US[41] to 12% in Scotland[42], and up to 33% in

Peru[34]. However, their frequency and prevalence by different species or strains are

challenging to estimate, given the reliance on limited reference databases and PCR-

generated data that may fail to detect low-abundance types/species within mixed

infection samples[19]. Thus, methods that enable the detection and characterization of

mixed infections are necessary to gain a better understanding of cryptosporidiosis.

Hybridization sequence capture has well-known advantages[43–45] relevant to studying

Cryptosporidium spp. and other parasites: it is feasible to design thousands of baits to

capture genomes, such as the 9 Mb genome of *Cryptosporidium* spp.; baits can tolerate

mismatches (thus regions with < 20% sequence divergence may be captured); and

successful enrichment reduces the amount of sequencing needed (thereby reducing

sequencing costs in comparison with non-enriched samples). The first *Cryptosporidium*

whole-genome bait set (CryptoCap_75k[46]) was designed using the *C. parvum* IOWAII

incomplete genome assembly[47], and is reported to perform well when each sample is

enriched individually. However, library pools that contain diverse species showed bias

during enrichment for *C. parvum* over other distantly related species[46]. Because fecal

samples can contain unknown or mixed *Cryptosporidium* species, a second-generation

bait set is needed to enrich genomes for a broader spectrum of *Cryptosporidium* species

and consider highly polymorphic genomic regions. Enriching pools of sample libraries,

rather than single libraries, is also desirable for time- and cost-efficiency.

To ameliorate the current challenges in the genomic study of human-infecting

Cryptosporidium spp., we developed, validated, and critically assessed CryptoCap_100k.

Motivated by CryptoCap_75k[46], we advance this approach by creating a bait set that also

captures highly diverse genomic regions (e.g., including all subtelomeric regions) and a

wide diversity of human-infecting species. CryptoCap_100k was designed (without

consideration of CryptoCap_75k) from the most complete publicly available genome

sequences from six *Cryptosporidium* species (Table 1). We validated the sensitivity and

specificity of CryptoCap_100k *in silico* and *in vitro* with simulated and mock samples,

multiple bait dilutions, and library preparation methods. We tested the reliability of these

methods in pure oocysts and in clinical samples and demonstrated that CryptoCap_100k

is an efficient and economical method to obtain genomic information for

Cryptosporidium. This genomic data facilitates inter-sample population genomics

analyses and intra-sample genetic variation via deconvolution analyses, opening new

opportunities to understand the complex biology of these important pathogens.

<u>Results</u>

CryptoCap_100k Bait Design & Characteristics

We designed a set of 100,000 120-mer RNA baits targeting six *Cryptosporidium*

species, including 73,672 baits for full coverage of *C. parvum* and 4,447 baits to cover

divergent regions among *C. hominis, C. cuniculus, and C. tyzzeri*. Additional baits were

designed to cover *C. meleagridis* and *C. viatorum*. Finally, we added 130 baits for genes

of particular interest (Table 1).

Mapping the baits to twelve *Cryptosporidium* genome sequences (Table 2,

Supplementary Notes #1) revealed ~90.46% breadth of genome coverage at 1X for target

species, but much lower values for distant species not considered in bait design (7.33% at

0.08X; Table 2; Fig. S1).

When comparing the multi-species design (CryptoCap_100k) to the single-species

design (CryptoCap_75k) (Supplementary Notes #2), despite slightly higher percentage of

target reads for CryptoCap_75k than CryptoCap_100k (64.73% vs. 55.6% baits,

respectively), both bait sets showed similar depth and breadth of coverage (Table 2; p-

values > 0.05; Figs. S2–S3).

In Silico Simulations of Enrichment and Sequencing

In silico simulations exhibited breadth of genome coverage of up to 99.9% and a

depth of coverage of 7.6X (for ~200,000 paired reads), minimal off-target hybridization

to more than a thousand non-*Cryptosporidium* genome sequences (including the human

genome), and reliable mapping fidelity to the database containing ten genome sequences

from ten *Cryptosporidium* species (10-CrypGS), which translated to high accuracy in

species assignment for each simulated *Cryptosporidium* species (Supplementary Notes

#3–7, Supplementary Tables S1–S6, Supplementary Figures S4–S13, and Supplementary

Results).

In Vitro Simulation of Single Infections and Mixed Infections

DNA from oocysts of *C. parvum* and *C. meleagridis* (Supplementary Notes #8;

Table S7) showed improved read mapping (95.8% vs. 68.1%, p-value = 0.094; Fig. S14),

similar breadth (85.2% vs. 81.6%, p-value = 0.75), and greater depth of coverage in

enriched vs. unenriched libraries (17.3X vs. 5X, respectively; p-value = 0.01; Fig. S15–

S16). Species assignment was largely accurate for both library types, with a low

proportion of reads from *C. parvum* oocysts assigned to *C. cuniculus* and *C. hominis* (Fig.

S17; Table S8).

The proportion of reads obtained for each species (*C. parvum* and *C. meleagridis*)

from the unenriched mock mixed infection sample (Supplementary Notes #9; Table S9)

showed that initially the mixture was unequal (60.6% *C. parvum* and 39.4% *C.*

meleagridis; Fig. 1A) and enrichment slightly increased the bias (64.3% *C. parvum* and

35.7% *C. meleagridis*; Fig. 1A; Table S9). In total, 95.7% and 81.4% reads mapped to

10-CrypGS, for enriched and unenriched libraries, respectively (p-value = 0.50), with

very accurate species assignment for both library types (> 97%; Fig. 1B). Species

assignment with *gp60* performed well for *C. parvum*, but less accurately for *C.*

meleagridis (Fig. 1C; Supplementary Results). Species assignments with *18S rRNA* were

~70% correct (Fig. 1D; Supplementary Results).

LOD: In Vitro Sensitivity Test with Different Inputs of C. parvum DNA

Dilutions of target DNA allowed the determination of the limit of detection (LOD) in

a background community. For NEB-iTru libraries (Supplementary Notes #10.1), the

percentage of reads mapped to *C. parvum* correlated with the initial concentration of

target DNA, with inputs below 0.0001 ng having very low percentages, with differences

between 1 ng and ≤ 0.01 ng, and 0.1 ng and ≤ 0.01 ng being statistically significant (p-

values ≤ 0.040; Fig. S18). The average depth of genome coverage was 4X for ≤ 0.01 ng,

29X for 0.1 ng, and 126X for 1 ng, with significant differences between 1 ng and all other

dilutions (p-values < 0.0001; Fig. 2A; normalized values Fig. S19). The breadth of

genome coverage was significantly different between comparisons below and above 0.01

ng (p-values ≤ 0.02; Fig. 2A; Supplementary Results; normalized values Fig. S19). For

unenriched libraries, accurate species assignment is observed only for 1 ng and 0.1 ng

inputs (Fig. 2B). For enrichments, *C. parvum* is accurately detected in inputs ≥ 0.001 ng

(Fig. 2B).

For iNextEra libraries (Supplementary Notes #10.2; Table S11, Fig. S19), the average

percentage of reads mapped increased with target input in enriched libraries (Fig. S20).

The percentage of mapped reads was lower for inputs of ≤ 0.01 ng vs. 1 ng (p-values ≤

0.015) and for inputs ≤ 0.001 ng vs. 0.1 ng (p-values ≤ 0.05). Depth and breadth of

genome coverage tended to increase as target DNA increased (Figs. 2C and S21).

Although depth did not show statistically significant differences among dilutions

(normalized values per 100,000 reads did: Supplementary Results, Fig. S21) for libraries

with ≤ 0.1 ng we recovered depths ≤ 4X whereas for those with 1 ng average depth was

34.8X (Fig. 2C). Breadth of genome coverage between the two highest input samples

(i.e., 1 ng and 0.1 ng, mean 57%–58%) differed from all other input samples (p values ≤

0.017, Fig. 2C).

The average percentage of reads mapped to the *Bos taurus* genome sequence (DNA

used as background diluent for *C. parvum*) decreased as *C. parvum* input increased in

enriched libraries. With mapped reads only being significantly different for inputs of 1 ng

vs. all the other input values (0.004 ≤ p-values ≤ 0.021). In unenriched libraries, mapping

percentages to *Bos taurus* were high across all dilutions, averaging 90.5% (Table S11,

Fig. S22; Supplementary Results).

Accurate species assignment was obtained in unenriched samples with inputs ≥ 0.1

ng, in single-enriched samples with ≥ 0.01 ng, and in double-enriched samples

consistently with ≥ 0.001 ng, and less consistently in samples of 0.0001 ng (Fig. 2D).

LOD: In Vitro Sensitivity Test with Different Dilution Levels of Baits

Bait dilutions enabled the determination of the minimum quantity that could be used

in a capture reaction. For both library types, no difference was observed among bait

dilutions (NEB-iTru: full down to one-sixteenth; iNextEra: one-quarter vs. one-eighth) in

the number of reads obtained, the percentage of reads mapped to *C. parvum,* and the

depth and breadth of genome coverage (Supplementary Notes #11.1–11.2; p-values ≥

0.05; Fig. S23–S29; Table S10; Supplementary Results).

Enriching Seven Human Fecal DNA Samples with the CryptoCap_100k Bait Set

From seven clinical patients' samples (Supplementary Notes #12.1, Table S12, Fig.

S30), the percentage of reads mapped increased from 0.02–0.15% (unenriched) to 0.37–

18.17% (enriched), with a fold change of 9.25–479X (Table S12, Fig. S31, p-value =

0.02). Species assignment revealed that in unenriched libraries, the few mapped reads

were assigned to non-human infecting *C. muris* (Fig. S32 right), similar to the

background effect observed in the LOD experiment. In contrast, in enriched libraries,

DOCK and UKH101 are predominantly assigned to *C. hominis* and *C. cuniculus*, while

EC4 and UKP196 to *C. parvum* (Fig. S32 left). No reads mapped to the *gp60* database,

and very few mapped to the *18S rRNA* database with assignment to species not common

in human infections (Fig. S33).

Enriching 100 Human Fecal DNA Samples with the CryptoCap_100k Bait Set

NEB-iTru libraries

For fecal DNA obtained from 100 patients when using the NEB-iTru protocol

(Supplementary Notes #12.2.1, Table S13; Fig. S34), on average, 35.6% of enriched vs.

0.08% of unenriched retained reads mapped to the *C. parvum* genome sequence (p-value

< 0.0001; Fig. 3A; Table S13; Fig. S35). Libraries were enriched with *C. parvum* by an

average factor of 2,095X (7.20–5,097X; Table S13). The depth and breadth of genome

coverage were significantly higher for enriched samples, with average depth of 18X and

breadth of 31% vs. 0.04X and 0.02% in unenriched samples (Fig. 3B; normalized

parameters per 100,000 reads can be found in Supplementary Results and Fig. S36).

For unenriched libraries, qPCR C_T values were not a good predictor of the percentage

of reads mapping to 10-CrypGS (R^2 = 0.012, p-value = 0.36; Fig. 3C). In enriched

libraries the qPCR C_T values were good predictor of the 15.5% of the variation in the

percentage of mapping reads with a statistically significant effect showing that for every

increase of one qPCR C_T unit, the percentage decreased by 2.81% (R^2 = 0.155, p-value =

5.11e-5; ±18%; Fig. 3D; Table S13).

Species assignments for unenriched libraries resemble the complex background from

LOD experiments (cf. Figs. 2B and 3C). Following enrichment, most hits matched *C.*

parvum (63 samples have >50% of hits to *C. parvum*; Fig. 3E), but many samples have

moderate proportions of reads assigned to other species, including those with limited

evidence to infect humans (Fig. 3F; see Supplementary Results).

iNextEra Libraries

For the 91 clinical iNextEra libraries (Supplementary Notes #12.2.2, Tables 3 and

S14; Fig. S37), on average, 0.03% of unenriched reads, 15.85% of single-enriched reads,

and 64.98% of double-enriched reads mapped to the *C. parvum* genome sequence

(Supplementary Notes #12.2.2, p-values < 0.0001, Fig. 4A, Table S14, Fig. S37;

Supplementary Results), yielding an average enrichment factor of 1,112X from

unenriched to single-enriched, and an extra 9.24X from single- to double-enriched (Table

S14). The mean depth of genome coverage was significantly higher for double-enriched

samples (95.5X) than for single-enriched libraries (32.1X) and unenriched libraries

(0.05X) (p-values < 0.005; Fig. 4B; normalized values per 100,000 reads in

Supplementary Results and Fig. S39, Table S14). The mean breadth of genome coverage

was significantly higher for double- and single-enriched samples (48% and 54%,

respectively) than for unenriched samples (2.6%, p-values < 0.0001).

The percentage of reads mapped was negatively correlated with qPCR C_T values for

all library types (Figs. 4C–4E). In unenriched and double-enriched libraries, the linear

model fails to detect C_T values as predictors of percentage of reads mapped (p-value >

0.05; Figs. 4C and 4E), but in single-enriched libraries qPCR C_T has a significant but

small effect on percentage of reads mapping ($R^2 = 0.006$, p-value = 0.016), that is for

every increase of one qPCR C_T unit, the percentage decreased by 1.29% (Fig 4D).

Double-enrichments allowed for the retrieval of a higher percentage of target reads, even

at higher C_T values, than the other two library types (Figs. 4C–4E).

Species assignments were similar to those from NEB-iTru libraries in unenriched and

single-enriched libraries (cf. Figs. 3E vs. 4F, and Figs. 3F vs. Fig. 4G). Double-enriched

libraries compared to single- and unenriched libraries exhibit an increased proportion of

hits to the most frequently observed human-infecting species (*C. parvum* and *C. hominis*;

Fig. 4H). Enriched libraries mapped most frequently along either *C. parvum*

chromosomes for UKP samples or *C. hominis-C. cuniculus* chromosomes for UKH

samples (Fig. S40), aligning with the species identified in the multi-locus typing study in

the same samples[48].

<u>iNextEra vs. NEB-iTru Libraries</u>

iNextEra and NEB-iTru libraries performed similarly (Table S15) but with

noteworthy differences. In single-enriched libraries, NEB/iTru samples obtained and

retained more reads following quality filtering, and a higher percentage of reads mapped

to the *C. parvum* genome sequence than iNextEra samples. The average depth of

coverage is similar for both protocols (p-value = 0.7); however, the breadth of coverage

was significantly higher for iNextEra libraries (p-value < 0.0001; Supplementary Notes

#12.2.3, Table S15; Fig. S41–S45; Supplementary Results).

Population Genomics Analyses of Clinical Samples

Enrichment drastically improved SNP recovery: obtaining 94,783–127,102 SNPs for

double- and single-enriched vs. 15,430 for unenriched in raw VCF files (Figs. S46–S51),

with large variations in read depth (DP; Fig. S52; Supplementary Results). Filtered VCF

files contained: 76 and 77 samples, and 12,498 and 13,578 SNPs for single- and double-

enrichments, respectively (Figs. S53–S56). For unenriched libraries, filtering was based

on read depth and biallelic variants only, yielding 91 samples and 11,226 SNPs (Figs.

S57–S58; Supplementary Notes #13.4; Supplementary Results).

Population genomics analyses (Supplementary Notes #13.5), such as principal

component analysis (PCA), phylogenetic networks, genetic distance trees, and Structure,

are highly concordant in enriched datasets. Both datasets suggest hierarchical structure (K

= 2; Figs. 5, S59–S68), with distinct clustering of UKP (*C. parvum*) and UKH (*C.*

hominis) samples, aligning with the species assignment analyses from above and previous

multi-locus typing results[48] (cf. Figs. 4H and 5A–5B). Discrimination not found in

unenriched libraries (Fig. 5C). Minor distinctions between single- and double-enriched

libraries are evident in sample UKH149, which is positioned in the trees as more closely

related to *C. parvum* than to *C. hominis*. However, in double-enriched libraries, UKH149

clusters much closer to all other *C. parvum* samples than in single enrichments (Figs. 5A

and 5B).

High pairwise F_{ST} values between UKP and UKH samples were estimated in enriched

(0.94 in double vs. 0.98 in single enrichments), but not in unenriched libraries (0.084).

These values are balanced across all chromosomes, showing no locus bias (Figs. S69–

S71). Pairwise genetic distances within samples of *C. parvum* reveal two to three clusters

in enriched libraries (Fig. S72–S73; Supplementary Results). For *C. hominis*, enrichments

reveal that the three samples are highly distinct from each other, even more than the

differences shown between clusters of *C. parvum* (Figs. S72–S73). Genetic distances

using unenriched libraries fail to reflect any biologically meaningful pattern (Fig. S74).

Deconvolution analyses (Supplementary Notes #13.6.1) revealed different numbers of

inferred strains (inferred K) among library types (Figs. 5A–5C). Sample UKH149 was

detected as having two strains by both enrichment methods (Table S16; Fig. 5B), and two

more samples were detected as having two strains for double-enriched libraries (UKP347

and UKP391) with effective Ks above 1.2 and a proportion of the least dominant

haplotype larger than 10% (Table S16; Fig. 5C, Table S16; Supplementary Results).

Three samples (UKH149, UKP347, and UKP391) exhibit the most differences between

capture methods. When assessing allelic read depth (AD) and genotype quality (GQ),

UKH149 and UKP347 had statistically larger AD and GQ values in double-enrichments

compared to single-enrichments (p-values < 0.0001; Figs. S75–S78, Table S17–S18).

Sample UKP391 had similar AD (p-value = 0.64; Fig. S79), but GQ was statistically

larger in double than in single enrichments (p-values < 0.0001; Fig. S79, Table S19).

Genetic diversity, measured when ploidy was set to 2 in GATK (Supplementary

Notes #13.6.5) revealed that the nucleotide diversity (π) of polymorphic sites within

UKH samples (considering UKH149: single-enriched mean $\pi = 0.066$; double-enriched

mean $\pi = 0.335$; not considering UKH149: single-enriched mean $\pi = 0.005$; double-

enriched mean $\pi = 0.0204$) and within UKP samples (single-enriched mean $\pi = 0.0122$;

double-enriched mean $\pi = 0.0187$) was statistically different between double- and single-

enrichments (p-values < 0.0001); and also between UKH and UKP samples (p-values <

0.0002; Fig. S80). High levels of heterozygosity were observed in samples with high

values of effective K in DEploid. These values were consistent among chromosomes,

showing no locus bias (Figs. S81–S86). Double enrichments showed a higher mean

heterozygosity compared to single enrichments (0.009 vs. 0.002), while unenriched

libraries showed a mean heterozygosity 1 for all samples (Fig. S86).

Discussion

Obtaining sufficient *Cryptosporidium* DNA for genomic studies has been a challenge

in the quest to understand *Cryptosporidium* global diversity and its links to transmission,

infection, pathogenesis, and host response. We developed and evaluated a hybridization

capture system to obtain *Cryptosporidium* genomic data from limited amounts of input

DNA and evaluated its performance in a variety of settings.

Experiments based on simulations and on DNA from purified oocysts demonstrate

that CryptoCap_100k retains nearly the whole genome of target species with minimal

bias (Figs. 1, S15–S17). However, obtaining pure oocysts requires increased effort and

time spent per sample, and via propagation, strains may be altered [14,29,31]. Thus, we aimed

to eliminate the need for purifying oocysts to obtain usable genomic data.

When the starting materials is fecal DNA, CryptoCap_100k increases on-target reads,

even at high C_T values, enables the pooling of ~10 libraries per capture reaction, and bait

dilution increases affordability. This translates into inexpensive genomic data that can be

applied to an extensive library of samples, including archived or fresh samples, from

diverse origins such as stool, environmental sources, oocysts, or existing extracted DNA.

Key findings reveal that CryptoCap_100k: i) is highly specific and achieves a higher

percentage of on-target reads than unenriched libraries in clinical fecal samples with both

NEB-iTru (~437X increase) and iNextera (~500X in single-enriched and ~2,167X in

double-enriched) libraries, ii), recovers ~85% of the genome without deep sequencing, in

DNA samples from pure oocysts, iii) recovers ~50% of the genome in fecal DNA

samples, iv) facilitates robust species assignment and SNP calling for population

genomics and deconvolution of mixed-infections within samples, and v) double-

enrichments should be prioritized in samples with very low target DNA.

Development and Assessment of CryptoCap_100k

With a foundation similar to the previous whole-genome sequencing bait capture for

C. parvum (CryptoCap_75k)[46], CryptoCap_100k used more complete genome sequences

that included the 16 subtelomeric regions (i.e., *C. parvum* IOWA-ATCC[49]) and included

additional baits to account for variation among *Cryptosporidium* species that commonly

infect humans. Performance between CryptoCap_75k[46] and CryptoCap_100k is similar *in*

silico (Table 2, Figs. S1–S3); however, CryptoCap_100k allows for the pooling of ~10

libraries before enrichment, decreasing processing costs and time.

When assessing target *Cryptosporidium* species, CryptoCap_100k recovered large

breadths of the genome (~95% *in silico*, ~85% from pure oocysts, and ~50% from fecal

samples; Tables S1, S7, and S15) with low sequencing efforts. A higher breadth of

coverage was not achieved by using more baits or performing double enrichments. These

results suggest that breadth is a property dependent on target input, as none of the

samples with $C_T > 28$ reached a breadth $\geq 50\%$ (Table S14). Still, CryptoCap_100k

enables obtaining genomic data for downstream analyses for 76%–85% of clinical

samples.

To make recommendations about how to use CryptoCap_100k best, we defined two

distinct limits of detection (LOD) for: i) minimum target genomic equivalents and ii)

minimum bait concentration. Given that a single haploid sporozoite contains ~9 fg of

DNA (with four sporozoites per oocyst) and is typically found in a complex community,

we diluted *C. parvum* DNA in a consistent background of calf DNA and determined the

data quality threshold. Enriched libraries exhibit a high percentage of target reads,

substantial breadth and depth of genome coverage, and accurate species assignment with

inputs as low as 0.01 ng of target DNA (~1,000 sporozoites or ~250 oocysts equivalents;

Figs. 2A–2B), but higher input of target DNA significantly reduces non-target reads (Fig.

S22). Non-target reads are likely caused by non-specific interactions of beads and/or

biotin with background DNA when target DNA is low[50]. These parameters improved

further in double-enriched iNextEra libraries with 0.001 ng of target DNA (~100

sporozoites or ~25 oocyst equivalents; Fig. 2D). These findings validate that

CryptoCap_100k can obtain genomic information at low DNA levels without purifying

oocysts. The LOD assay with bait dilutions suggests that there are no differences in

estimated parameters between bait dilutions (full amount down to a sixteenth, Figs. S23–

S27), offering a cost-effective strategy for large-scale and effective use.

Validating CryptoCap_100k in Clinical Samples

We successfully tested CryptoCap_100k on a small initial set of seven clinical

samples. Three of these had also been analyzed with CryptoCap_75k[46], which identified

UK101 as *C. hominis* and UK196 as *C. parvum*[46], both of which agreed with our analyses

(Fig. S32). CryptoCap_75k also identified sample EC4 as *C. canis* based on the *18S*

rRNA marker, but did not obtain sufficient genome-wide read coverage to assess SNP

diversity [46]. In our analyses, which lack a reference genome for *C. canis*, we identified

this sample as largely *C. parvum*, with a profile of likely mixed ancestry (Fig. S32), but

this was not validated with SNP data.

Several studies recommend sequencing only samples with C_T values < 16[30,51], and

although our set of 100 clinical samples had an average C_T value of 25.4, we were able to

recover significantly more target reads, higher averages of breadth and depth of genome

coverage, and better accuracy of species assignment in enriched vs. unenriched libraries

with both the NEB-iTru and iNextEra protocols (Fig. 3; Fig. 4; Table S13–S14). In the

linear regression models for C_T values and target reads, enrichments show less loss of

target reads at higher C_T values compared to unenriched libraries (Figs. 3C–3D, 4C–4E).

Double enrichments consistently resulted in significant increases of target reads and

depth of genome coverage (Figs. 4A–4B), as well as more accurate species assignment

than single-enrichments (Figs. 4F–4H).

CryptoCap_100k Enables the Study of Population Genomics

CryptoCap_100k is a reliable tool for high-confidence SNP calling of thousands of

variants. Enrichment methods resolved species-level (*C. parvum* and *C. hominis*)

clustering, which matched the multi-locus typing results[48] (Figs. 4H, 5A–5B). Beyond

detecting different species, we were able to assess the diversity within each species.

Enrichments show that *C. parvum* haplotypes are highly similar and have low

nucleotide diversity. Still, two clusters within the species are detected (Figs. S68 and

S73), which may correspond to two circulating but highly similar variants, or samples

with some levels of mixed infections; however, the latter was not confirmed by our

deconvolution analyses. Genomic studies have revealed two distinct clusters for *C.*

parvum in the UK[25]. Our samples may represent two less divergent lineages within just

one of these clusters, highlighting the prevalence of clonality and the dominance of some

lineages in the region (Figs. 5A and 5B).

We also identified two distinct clusters of *C. hominis* separated by the first PC axis,

which explains 56-57% of the variance (UKH148 + [UKH149 and UKH150]; Figs. 5A–

5B). Two subspecies of *C. hominis* have been detected in the UK, *C. h. hominis*, and *C.*

h. aquapotentis, with some evidence of gene flow and recombination between them in the

region[26]. The highly divergent *C. hominis* samples in our study may reflect the two

subspecies, given their comparable levels of variation.

Given that more than 90% of cryptosporidiosis cases in humans are caused by *C.*

parvum and *C. hominis*[52], our clinical results were representative. Because ~74% of the

baits in our set match to *C. parvum*, it is relevant to assess CryptoCap_100k performance

for other species. In our study, we demonstrate that CryptoCap_100k enables the

collection of genomic data for *C. hominis* in clinical samples, from which only ~3% of

the baits had been designed specifically, but we expected a larger set to work thanks to

the 96.8% percent similarity with *C. parvum* genome sequences [53] and the mismatch

tolerance of the baits. Our evidence suggests that CryptoCap_100k works appropriately

in other species: i) *in silico* analyses reveal proper hybridization of our baits to other

target species (Fig. S7); ii) *in vitro* tests using pure oocyst of *C. meleagridis* show proper

species assignment and a high percentage of target reads (Fig. 1B); and iii) the number of

baits for some species, such as *C. meleagridis* and *C. viatorum* included in

CryptoCap_100k is even higher than the number for *C. hominis*, which worked

successfully (Table 1).

CryptoCap_100k Enables the Detection of Mixed Infections

Cryptosporidium infections can contain multiple strains, subspecies, or species.

Current methods for detecting mixed infections include amplicon, PCR-RFLP, and

MLST approaches[34,54,55]. However, these methods have inherent biases due to

preferential amplification and can miss mixed infections that are not comprised of *C.*

parvum and *C. hominis*[34,35,54]. Regarding mixed-species infections, CryptoCap_100k was

able to detect these in *in silico* simulations (Fig. S13); in *in vitro* pools of *C. parvum* and

C. meleagridis, with a slight bias towards *C. parvum*; and in clinical samples (i.e., EC4,

UKH149, UKP381, and UKH151; Figs. 4H and S32). These results were confirmed for

sample UKH149 through deconvolution analyses; however, double-enrichments exhibit

higher proportions of *C. parvum* compared to single-enrichments. These higher

proportions of *C. parvum* can be caused by minor enrichment biases due to the high

number of baits designed from *C. parvum*[46].

Mixed infections with strains of the same species were detected in UKP347 and

UKP391 (Fig. 5A) in double-enriched but not in single-enriched libraries. These

discordances were associated with higher allelic depth (AD) and/or improved genotype

quality (GQ) in double-enriched libraries (Figs. S75–S79), which likely allowed for

greater resolution and ability to detect imbalances in within-host allele frequency by

DEploid.

In summary, CryptoCap_100k results in a cost reduction of approximately 98%

relative to deep sequencing of fecal samples. However, cost savings will vary depending

on the abundance of *Cryptosporidium* DNA in the sample (Supplementary Results and

Table S20). CryptoCap_100k can recover >80% of the genome at high depth from pure

oocyst samples (Figs. S13–S14) and achieve as high as 3,690X fold-change enrichment

478 in fecal samples, without the need for oocyst purification, allowing high-accuracy species

479 assignment (Table S14). The quality of the retrieved genomic data allows for thousands

480 of SNPs to be called, useful for population genomic and mixed-infection studies. Double

481 enrichment improves coverage for low-input DNA samples, but this outcome can be

482 matched with single enrichments and higher sequencing effort. Studies using the

483 CryptoCap_100k will enhance our understanding of the population genetics of human-

484 infecting *Cryptosporidium* species and their relationship to epidemiology. Generating

485 economical and efficient genome-level data enables the growth of reference resources

486 and the refinement of analyses and methods for subtyping.

488 **Methods**

489 *CryptoCap_100k Bait Design & Characterization*

490  The genome sequences of six *Cryptosporidium* species were chosen for bait

491 design (Table 1), along with 55 *18S rRNA* sequences and eight *gp60* sequences (Data

492 Availability). All sequences were soft-masked for simple and low-complexity repeats

493 using RepeatMasker 4.1.1 (http://www.repeatmasker.org/). We aimed to design 100,000

494 120-nucleotide baits. Baits that were ≤ 50% soft-masked for repeats were retained. These

495 parameters resulted in 435,323 raw baits. Then, VSEARCH 2.15.0[56] was used to cluster

496 baits at various levels of sequence overlap and identity (Table 1), allowing for

497 mismatches to reduce the number of baits while maintaining diversity. The genome order

498 was designed to prioritize genome sequences based on their completeness and likelihood

499 of human infection, with *C. parvum* IOWA-ATCC listed first for all clustering

500 procedures. Then, *C. hominis*, *C. cuniculus*, and *C. tyzzeri* were collapsed at 50%

overlap, and 90% identity relative to *C. parvum*; and *C. meleagridis* and *C. viatorum*

were collapsed at 45% overlap and 85% identity relative to *C. parvum*. From *C.*

viatorum, a subset of baits was randomly selected. Raw baits were obtained for the *18S*

rRNA targets and collapsed at 50% overlap and 95% identity (Table 1).

We characterized CryptoCap_100k (Supplementary Notes #1) by using twelve

genome sequences available for ten species of *Cryptosporidium* (Data Availability).

BWA 0.7.17[57] was used to map the CryptoCap_100k bait sequences to each genome

sequence. Alignment files were generated and converted using Samtools 1.10[57,58]. The

bamCoverage function from DeepTools 3.3.1[59] was used to produce bigwig files to

visualize coverage tracks calculated as the number of reads per bin in IGV-Web app

version 1.10.8[60]. To obtain alignment statistics, the depth option from Samtools v1.1.0

and the -*a* flag were used to get all sites and create a depth file[58]. The mean depth, the

proportion of baits mapped to each reference, and the breadth of genome coverage were

estimated. The breadth of coverage, as reported throughout this manuscript, was

determined by estimating the number of bases covered in the alignment over the

reference length, with a minimum coverage depth of 1X.

We also compared the CryptoCap_100k bait set to CryptoCap_75k[46]

(Supplementary Notes #2) by estimating the same parameters as above, and using

Welch's two-sample t-test for comparisons.

In Silico Simulations of Enrichment with CryptoCap_100k & Sequencing

To validate data quality from CryptoCap_100k before bait synthesis, a simple

infection scenario, error-free and bias-free, was simulated. We used twelve available

genome sequences from ten *Cryptosporidium* species to simulate the enrichment and

sequencing process with CryptoCap_100k, assuming two different ranges of insert size

fragments (Supplementary Notes #3). To simulate hybridization, we used the coordinates

of the mapped baits in relation to each reference genome, and we extended the position

by several bases to "capture" larger fragments than the bait length using Samtools[58]. To

simulate sequencing, ART 2016.06.05[61] was used to reproduce ~200,000 paired-end 150

bp fastq reads from the extended coordinates in each reference genome. Then, reads were

mapped to their respective genome sequences using BWA 0.7.17[57], and we estimated

depth and breadth of coverage. To validate the accuracy of species assignment, we used a

reference database containing ten *Cryptosporidium* species genome sequences (10-

CrypGS), a *gp60* database[2], and an *18S rRNA* database[52,62,63], and used BBMap 38.90[64] to

map simulated fastq reads to each and explore the proportion of hits to each

Cryptosporidium species.

To assess the sensitivity of the baits to off-target DNA, we assessed the affinity of

our baits to 1,008 genome sequences[65], including the human genome. Because *in vitro*

libraries were diluted in calf thymus DNA (see *In Vitro Test of Limits of Detection with*

Different Levels of C. parvum Dilution), we also assessed affinity to the *Bos taurus*

genome sequence (GCA_002263795.3). For both analyses, we used BWA 0.7.17[57] to

map the bait sequences, and we estimated the number of hits as well as the quality of the

mapping.

To explore the ability to discriminate different species within mixed-species

infection samples, we created four simulated samples by concatenating the fastq files

already simulated for each species above. The four *Cryptosporidium* mixed infection

samples contained at equal proportions: i) reads from twelve genome sequences from ten

species; ii) reads from ten genome sequences from ten species; iii) reads from five

genome sequences and species that share high percent identity; and iv) reads from three

genome sequences and species commonly found in human infections. Then, we mapped

these mixed reads to 10-CrypGS, the *gp60*, and the *18S rRNA* databases. See

Supplementary Notes #3–7 for details.

In Vitro Enrichment & Validation with CryptoCap_100k Baits with Pure Oocsts' DNA

Pure oocyst DNA was obtained from three *C. parvum* samples from Dr. Michael

Grigg (National Institutes of Health, US) and *C. meleagridis* DNA isolate TU1867 from

BEI Resources (BEI Resources NR2521 Manassa, VA). Metagenomic libraries were

prepared with the New England BioLabs Ultra FS II library kit (see *Metagenomic*

Libraries), and aliquots of these were left unenriched or enriched with one-tenth of the

recommended volume of CryptoCap_100k (see *Hybridization Capture Enrichments*).

Libraries were sequenced using an Illumina MiSeq with a PE250 kit.

In Vitro Simulation of Mixed Infections

To test the ability to detect mixed infection samples, after NEB/iTru library

preparation (see *In Vitro Enrichment & Validation with CryptoCap_100k Baits with Pure*

Oocysts' DNA) but before bait capture, two indexed libraries were pooled at equimolar

ratios to simulate a mixed infection. After pooling, an aliquot was enriched with one-

tenth of the recommended volume of CryptoCap_100k, and another remained

unenriched. These were sequenced in an Illumina MiSeq using a PE250 kit. We

estimated the number of reads per sample after demultiplexing, expecting roughly 50%

for each species.

In Vitro Test of Limits of Detection with Different Levels of C. parvum Dilution

574   To determine the CryptoCap_100k limits of detection (LOD) of *Cryptosporidium*,

we performed two types of library preparations, NEB-iTru and iNextEra[66], on samples

with known and varied input DNA. For NEB-iTru libraries, we diluted *C. parvum*

genomic DNA in a background of calf thymus DNA (Sigma-Aldrich, St. Louis, MO). A

45-ng aliquot of calf thymus DNA was pipetted into five tubes. In the first tube, 5 ng of

C. parvum genomic DNA was added to a final volume of 50 µL. Then, three 1:10 serial

dilutions were performed. The fifth tube corresponded to a negative control with just the

calf DNA. Metagenomic libraries using the NEB-iTru protocol and 10 µL of each

dilution were generated (see *Metagenomic Libraries*). In this experiment, up to 7 libraries

were pooled, an aliquot was left unenriched, and independent aliquots were enriched with

full, one-half, a quarter, an eighth, or a sixteenth of the recommended volume of

CryptoCap_100k (see *Hybridization Capture Enrichments*). Pools were sequenced at

Novogene using a HiSeq X with a PE150 kit. The number of reads, percentage of reads

mapped to the *C. parvum* genome sequence, and depth and breadth of genome coverage

were estimated.

589   For iNextEra libraries, 90 ng aliquots of calf DNA were pipetted into six tubes. In

the first tube, 10 ng of *C. parvum* genomic DNA was added in a final volume of 100 µL.

From this, four 1:10 serial dilutions were performed. The sixth tube corresponded to the

negative control (calf DNA only). iNextEra library preparation was performed with 10

μL inputs from these dilutions (see *Metagenomic Libraries*). Up to 6 libraries were

pooled, and aliquots were left unenriched or enriched with one-quarter or one-eighth of

the recommended baits. Aliquots of single-enriched libraries were double-enriched with

one-quarter of the recommended baits (see *Hybridization Capture Enrichments*). Pools of

unenriched, single-enriched, and double-enriched libraries were sequenced at Novogene

using a NovaSeq with a PE150 kit. The number of reads, the percentage of reads mapped

to the *C. parvum* genome sequence, the depth and breadth of genome coverage, and

species assignment based on the mapping of reads against 10-CrypGS were analyzed

among dilutions.

In Vitro Test of Limits of Detection with Different Levels of Bait Dilution

To determine the sensitivity or limits of detection (LOD) when using a lower

concentration of baits than recommended, we modulated the quantity of baits added to

the capture reaction in one experiment with NEB-iTru libraries and one experiment with

iNextEra libraries.

For both library types, the bait dilution sensitivity assay considered pools of the

metagenomic libraries from the *In Vitro Test of Limits of Detection with Different Levels*

of C. parvum Dilution experiment. For NEB-iTru libraries, we tested five different bait

quantities: full volume (5.5 μL, per the manufacturer's specifications), and dilutions in

water at one-half, a quarter, an eighth, and a sixteenth. For iNextEra libraries, we tested a

quarter and an eighth bait dilutions. Sequencing was performed as described in the

section above. The number of reads, the percentage of reads mapped to the *C. parvum*

genome sequence, and the depth and breadth of genome coverage were compared among

bait dilutions.

In Vitro Enrichment & Validation with CryptoCap_100k Baits with Clinical Samples

Two clinical datasets were used for real-world validation. First, Dr. Michael

Grigg (National Institutes of Health, US) and Prof. Rachel Chalmers's laboratory (Public

Health Wales—PHW— Microbiology and Health Protection, UK) provided DNA

extracted (Qiagen PowerSoil Kit) from seven de-identified patient fecal samples. These

samples were obtained under an NIAID IRB-approved protocol NCT00006150. The

NEB-iTru library protocol was used, pools made, aliquots were either left unenriched or

enriched with one-tenth of the recommended volume for baits, and libraries were

sequenced in an Illumina HiSeq using a PE150 kit.

Second, 100 DNA extractions (Qiagen QIAamp Fast DNA Stool kit) from clinical

samples from Prof. Rachel Chalmers's laboratory (PHW) were used. This work was

carried out in accordance with a material transfer agreement between UGA and PHW,

which did not require ethical approval. These samples had previously been screened

using a multi-locus typing system [48] to detect the species present, and these previous

results are reflected in the sample names used here (*C. parvum* = UKP and *C. hominis* =

UKH). The laboratory workflow used for these clinical samples in this study is depicted

in Fig. 6. A real-time qPCR assay was used to quantify *Cryptosporidium*, using primers

for the *18S rRNA* gene and slightly modified probes [67] as specified in Supplementary

Notes #12.2. Two types of genomic libraries were prepared for each sample (NEB-iTru

and iNextEra; see *Metagenomic Libraries*). Nine DNA samples were depleted when used

for iTru libraries, thus not used for iNextEra libraries. Aliquots of pools of ~12 libraries

were left unenriched or enriched. iTru-NEB libraries were enriched with one-tenth and

iNextEra libraries were enriched with one-eighth of the recommended volume of

CryptoCap_100k. For iNextEra libraries, we performed additional double enrichments on

single-enriched aliquots using one-quarter of the recommended volume of baits (see

Hybridization Capture Enrichments with CryptoCap_100k). All libraries were sequenced

using an Illumina NovaSeq X with a PE150 kit (Fig. 6).

Metagenomic Libraries

For iTru-NEB libraries, the New England BioLabs (NEB) Ultra FS II library kit

(New England BioLabs, Ipswich, MA) protocol was used according to the manufacturer,

with enzymatic fragmentation lasting 5–11 minutes, based on the integrity of DNA in gel

electrophoresis. A 12- or 14-cycle PCR with iTru primers[66] was used to add unique dual-

indexes to each ligation product. Products were cleaned with 1.25X SpeedBeads[68] and

quantified with a Qubit 4.0 Fluorometer (Thermofisher, Waltham, MA).

iNextEra library preparation was used to decrease the time, cost, and amount of

input DNA needed. In brief, 2.5 µL of 10 ng/µL (25 ng) of sample DNA was used for a

tagmentation reaction with 3 µL of 2X TMP buffer and 0.5 µL of Transposome (BLT;

CAT # 20015880, Illumina Inc.), incubated at 53 ℃ for 30 minutes. Then, PCR reactions

contained 9.5 µL of Q5 Buffer, 1.25 µL of 10 mM dNTPs, 0.5 µL of Q5 High Fidelity

Polymerase (New England Biolabs M0491L), 40.25 µL of nuclease-free water, 2.5 µL of

iNextEra5 indexed primer at 5 µM, 2.5 µL of iNextEra7 indexed primer at 5 µM[66], and 6

µL of tagmented product. Cycling conditions were: 3 minutes at 72 ℃, 3 minutes at 98

°C, followed by ten cycles of 98 °C for 45 seconds, 62 °C for 30 seconds, and 72 °C for 2

minutes, with a final extension at 72 °C for 1 minute.

Hybridization Capture Enrichments with CryptoCap_100k

Metagenomic libraries were pooled (e.g., Fig. 6) in preparation for enrichment.

For different experiments, the input concentration per pool of samples varied (i.e., for

clinical samples using NEB/ iTru, the input was between 25–80 ng/μL; for clinical

samples using iNextEra, it was 103 ng/μL; and for the LOD experiments, it was 26–59

ng/μL). The enrichment process per pool followed manufacturers' standard protocols

(mybaits v5, Daicel Arbor Biosciences, Ann Arbor, MI), with wash and hybridization

temperatures at 63 °C and modifications of bait quantities as described above. After

enrichments, P5/P7 PCRs with 16–22 cycles were performed, followed by 1–1.25X

SpeedBeads clean-ups (Sigma Aldrich, St. Louis, MO).

For the 100-clinical sample set and the LOD experiments, double enrichments

were performed on the iNextEra single-captured pools (e.g., Fig. 6). For these, aliquots of

the cleaned single-captured pools were pooled and used independently as input for a

second capture using one-quarter of the recommended baits. For post-double-enrichments

P5/P7 PCRs with 12–16 cycles were performed, followed by 1X SpeedBeads clean-up.

All PCRs were performed using the KAPA HiFi Kit (Roche, Basel, Switzerland). All

P5/P7 PCRs consisted of 98°C for 2 minutes.; then, cycles of: 98°C for 20 sec., 60°C for

30 sec., 72°C for 1 min.; followed by 72°C for 5 min.

In Vitro Bioinformatic Processing and Analyses

Bioinformatic and data analysis code are available in the Supplementary Notes.

Sequencing data were processed using Trimmomatic 0.39[69] , and retained reads were

mapped to specific genome sequences or databases using BBMap 38.9[64]. Samtools[58] was

used to sort bam files. We determined mapping statistics, including genome depth and

breadth of coverage. The results of the mapping to 10-CrypGS were used to assess the

proportion of hits to the different *Cryptosporidium* species and produce species

assignment plots. For iNextEra libraries from clinical samples, we also generated bigwig

files, which were visualized using the IGV-Web app version 2.2.7.

Cleaned reads from pure oocysts libraries and *in vitro* mixed-infection libraries

were mapped to the corresponding target references (i.e., *C. parvum* samples to *C.*

parvum IOWA-ATCC, *C. meleagridis* samples to chromosome-level assembly[70], also to

10-CrypGS, to the *gp60* database, and the *18S rRNA* database—used in simulations). The

sequencing reads from the LOD dilution experiments (i.e., input DNA and bait dilutions)

were mapped to *C. parvum* IOWA-ATCC and 10-CrypGS. iNextEra reads from LOD

were also mapped to *Bos taurus* (GCA_002263795.3). The cleaned sequence reads from

clinical samples were mapped to *C. parvum* IOWA-ATCC and 10-CrypGS. For the set of

100 clinical samples, qPCR C_T values were used in the *parsnip 1.1.1*[71] to define linear

regression models predicting the percentage of reads mapped against 10-CrypGS

(Supplementary Notes #12).

Population Genomics and Mixed Infection Analyses of Clinical Samples

To examine the genetic variation in natural populations of *Cryptosporidium* from the

set of 91 clinical samples prepared using the iNextEra protocol, we performed genotype

calling (Fig. S87). In short, reads from each treatment (unenriched, single-enriched, and

double-enriched) were mapped to the *C. parvum* IOWA-ATCC genome sequence using

BWA 0.7.17[57] , and the results were used in GATK 4.6.0.0[72]. Duplicate reads and

haplotypes were determined using Haplotypecaller with GVCF mode and merged into

databases for each library type to perform joint genotyping using GenotypeGVCFs.

Importantly, HaplotypeCaller was run with --sample-ploidy set to 1. A first filtering

approach was performed to obtain SNPs with QD < 2, FS > 60, MQ <40, MQRankSum <

-12.5, and ReadPosRankSum < -8.0.

The resulting VCFs were explored for missing data and coverage depth per sample

and chromosome using vcfR 1.15.5[73]. Then, filtering using dartR 2.9.7[74] considered only

SNPs with coverage (DP) larger than 5 and lower than 200, that were biallelic in the

dataset, and with less than 40% of missing data. Individuals with > 80% missing data

were removed.

To explore the diversity and divergence among samples, filtered VCF files for each

treatment were used for population genomic analyses. These included Principal

Components Analyses performed with SNPRelate 1.40.0[75] and ade4 1.7-23[76]. We used

phangorn 2.12.1[77]and SplitsTree 6.4.11[78] to compute pairwise genetic distances and

visualize phylogenetic networks. We estimated Nei's genetic distances between samples

and visualized them in dendrograms with bootstrap support using poppr 2.9.6[79] and

plotted these with ape 5.8-1[80]. We tested K = 1–5 clusters, each with three replicates, in

Structure 2.3.4[81], run through structure_threader 1.3.10[82], which utilizes Structure

Harvester 0.6.94[83]. We used 10,000 chains for burn-in, 100,000 chains for MCMC, and

set ploidy to 1, with other options left as default. We used readr 2.1.5[84] and starmie

0.1.2[85] to graph Evanno's results and to plot Structure assignment plots. We estimated

pairwise F_{ST} statistics between UKP and UKH samples using hierfstat 0.5-11[86] and

mapped these values along *C. parvum* chromosomes using qqman 0.1.9[87]. We compared

Allelic depth (AD) and genotype quality (GQ) values between VCF files of the different

treatments and used ggridges 0.5.6[88] to plot the frequency of values across samples.[90]

To explore the diversity within each clinical sample, we deconvoluted multiple

infecting strains using DEploid 0.7.1[89]with no reference panel. The effective and inferred

number of strains (K) was plotted. Additionally, using filtered VCFs from the GATK

4.6.0.0[72] run with the setting --sample-ploidy set to 2, we estimated nucleotide diversity

(π) of SNPs in VCFTools 0.1.16[90] for double and single enrichments that were subset to

contain only UKH (*C. hominis*) or UKP (*C. parvum*) samples; and, we assessed and

plotted heterozygosity levels per sample. Data analyses were performed in R Studio

2023.06.0. Least-square comparisons of the means between experimental groups (e.g.,

unenriched and enriched, single-capture vs. double-capture, etc.) were performed to

calculate p-values using a confidence interval of 0.95 with the package: *lsmeans*[91], which

provides a t-test output when comparing two variables, or performs a Tukey method

adjustment when comparing a family of three or more estimates. Other packages used

were *tidyverse 2.0.0*[92], *ggplot2 3.5.2*[93], and *plotly 4.10.4*[94].

<u>Acknowledgments</u>

T.G. and J.C.K. disclose support from the National Institutes of Health under

award number NIH R01AI148667. M.G. discloses support from the Intramural Research

Program of the National Institute of Allergy and Infectious Diseases (NIAID) at the

National Institutes of Health. We thank the authors of CryptoCap_75k[46] for sharing

unpublished results that facilitated the work and comparisons presented here. We also

thank Prisha Sharma and Fiifi Agyabeng-Dadzie for sample processing, Christopher

Fitzgerald for sample tracking and processing, and Mustafa Nural for sample database

help. This study was supported in part by resources and technical expertise from the

Georgia Advanced Computing Resource Center, a partnership between the University of

Georgia's Office of the Vice President for Research and Office of the Vice President for

Information Technology.

<u>Author Contributions</u>

N.J.B-V., A.H.S., M.S.B., A.K., R.P.B., M.E.G., J.C.K., and T.C.G. contributed

to conceptualization, methodology, investigation, and writing—review and editing.

M.S.B., R.P.B., and B.B. designed and oversaw the production and QC of the baits.

M.S.B., K.N.P., M.I.U.B., contributed to investigation and writing—review and editing.

N.J.B-V. and A.H.S. performed validation, data curation, formal analyses, and with

T.C.G. writing—original draft. G.R., R.M.C., E. A-F, M.E.G., and J.C.K. contributed

samples and analyses to the manuscript. N.J.B-V, M.S.B, A.K., R.P.B., M.E.G., J.C.K.,

and T.C.G contributed to funding acquisition. All authors read and approved the final

version of the manuscript.

<u>Competing Interests</u>

B.B. and M.B. are employed by Diacel Arbor Biosciences, which manufactures and

sells the baits described herein. The CryptoCap_100k bait set is available for commercial

purchase (Design ID: D10006Crypto, Cat# #308508.V5, #308548.V5, and #308596.V5,

Arbor Biosciences, MI, USA).

Data Availability

The six genome sequences that supported the CryptoCap_100k bait design were

from GenBank: GCA_015245375.1, GCA_004337795.1, GCA_001483515.1,

GCA_007210665.1, GCA_004337835.1, and GCA_001593445.1. The extra sub-

telomeric sequences from *C. parvum* were: MZ892386.1, MZ892388.1, and

MZ892387.1.

The *18S rRNA* sequence data that supported the CryptoCap_100k bait design

were from GenBank with the EU250845, LC089976, DQ288166, EF641022, GU951714,

AF108864, LC012016, GQ121020, HQ397716, KR296812, EF641014, FJ435960,

JX416368, KP099082, AY120903, AY120902, JX644908, AY324641, AY324638,

AF159113.1, JQ413438, AF108861, GQ121021, KP730304, AY737602, AY731235,

JX294358, AY268584, KR819168, AY120904.1, KU608308, AF112576, EU553551,

JF710247, KP004205, JQ002555, KT235702, AB909498.1, AF316630, AY504515,

AY504516, AF262325, GQ227479, EU827297, AF093500, AF093497, AY954886,

HM116385, DQ650344, KY490554, KC305650, EF547155, FJ769050, HM243547, and

AY524773 accession codes.

The *gp60* sequence data that supported the CryptoCap_100k bait design were

from GenBank with the FJ490060, FJ490087, FJ490058, JX412915, JX412926,

JX412925, KC204983, and KC204984 accession codes.

The twelve *Cryptosporidium* genome sequences used for CryptoCap_100k bait

set characterization and *in silico* simulations were from GenBank with the

GCF_00000165345.1, GCA_004337835.1, GCA_004337795.1, GCA_007210665.1,

GCF_001865345.1, GCA_001865355.1, GCA_001593455.1, GCF_000006515.1,

GCA_000165345.1, GCA_001483515.1, and GCA_002223825.1 accession codes *C.*

hominis UdeA01 and *C. parvum* IOWA-ATCC were removed for those analyses with

only ten genome sequences reference database (10-CrypGS).

Generated fastq files from all *in silico* and *in vitro* experiments for mock and

clinical samples are available in GenBank BioProject number PRJNA1061798. Clinical

samples were scrubbed of human DNA upon submission by NCBI using their Human

Read Removal Tool.

<u>Computer Code</u>

The code and data used for all sections of this paper are described in the

Supplementary Notes, also available in Dryad, doi: 10.5061/dryad.gtht76j0h.

References

1. Hunter, P. R. & Nichols, G. Epidemiology and clinical features of *Cryptosporidium* infection in immunocompromised patients. *Clin Microbiol Rev* **15**, 145–154 (2002).

2. Khan, A., Shaik, J. S. & Grigg, M. E. Genomics and molecular epidemiology of *Cryptosporidium* species. *Acta Trop* **184**, 1–14 (2018).

3. Wang, J. *et al.* Long-term treatment with green tea polyphenols modifies the gut microbiome of female sprague-dawley rats. *J Nutr Biochem* **56**, 55–64 (2018).

4. Bouzid, M., Hunter, P. R., Chalmers, R. M. & Tyler, K. M. *Cryptosporidium* Pathogenicity and Virulence. *Clin Microbiol Rev* **26**, 115–134 (2013).

5. Chalmers, R. M. & Katzer, F. Looking for *Cryptosporidium*: the application of advances in detection and diagnosis. *Trends Parasitol* **29**, 237–251 (2013).

6. Khalil, I. A. *et al.* Morbidity, mortality, and long-term consequences associated with diarrhoea from *Cryptosporidium* infection in children younger than 5 years: a meta-analyses study. *Lancet Glob Health* **6**, e758–e768 (2018).

7. Gilbert, I. H. *et al.* Safe and effective treatments are needed for cryptosporidiosis, a truly neglected tropical disease. *BMJ Global Health* **8**, e012540 (2023).

8. Sow, S. O. *et al.* The Burden of *Cryptosporidium* Diarrheal Disease among Children < 24 Months of Age in Moderate/High Mortality Regions of Sub-Saharan Africa and South Asia, Utilizing Data from the Global Enteric Multicenter Study (GEMS). *PLOS Neglected Tropical Diseases* **10**, e0004729 (2016).

9. GBD Results. *Institute for Health Metrics and Evaluation* https://vizhub.healthdata.org/gbd-results.

10. Kotloff, K. L. *et al.* Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *The Lancet* **382**, 209–222 (2013).

11. Bhutta, Z. A. & Black, R. E. Global maternal, newborn, and child health--so near and yet so far. *N Engl J Med* **369**, 2226–2235 (2013).

12. Cryptosporidiosis NNDSS Summary Report for 2019 | Water-related Topics | Healthy Water | CDC. https://www.cdc.gov/healthywater/surveillance/cryptosporidium/cryptosporidium-2019.html (2021).

13. Feng, Y., Ryan, U. M. & Xiao, L. Genetic Diversity and Population Structure of *Cryptosporidium*. *Trends Parasitol* **34**, 997–1011 (2018).

14. Morris, A., Robinson, G., Swain, M. T. & Chalmers, R. M. Direct Sequencing of *Cryptosporidium* in Stool Samples for Public Health. *Front Public Health* **7**, 360 (2019).

15. Nader, J. L. *et al.* Evolutionary genomics of anthroponosis in *Cryptosporidium*. *Nat Microbiol* **4**, 826–836 (2019).

16. Baptista, R. P., Cooper, G. W. & Kissinger, J. C. Challenges for *Cryptosporidium* Population Studies. *Genes (Basel)* **12**, 894 (2021).

17. Baptista, R. P. *et al.* Long-read assembly and comparative evidence-based reanalysis of *Cryptosporidium* genome sequences reveal expanded transporter repertoire and duplication of entire chromosome ends including subtelomeric regions. *Genome Res* **32**, 203–213 (2022).

18. Šlapeta, J. Cryptosporidiosis and *Cryptosporidium* species in animals and humans: a thirty colour rainbow? *Int J Parasitol* **43**, 957–970 (2013).

19.   Ruecker, N. J., Hoffman, R. M., Chalmers, R. M. & Neumann, N. F. Detection and
resolution of *Cryptosporidium* species and species mixtures by genus-specific nested PCR-
restriction fragment length polymorphism analysis, direct sequencing, and cloning. *Appl
Environ Microbiol* **77**, 3998–4007 (2011).

20.   Wang, Y. *et al.* Population genetics of *Cryptosporidium meleagridis* in humans and birds:
evidence for cross-species transmission. *International Journal for Parasitology* **44**, 515–521
(2014).

21.   Yang, Z. *et al.* Multilocus sequence typing and population genetic structure of
Cryptosporidium cuniculus in rabbits in Heilongjiang Province, China. *Infection, Genetics
and Evolution* **64**, 249–253 (2018).

22.   Tang, Y. *et al.* Development of a multilocus sequence typing tool for high-resolution
subtyping and genetic structure characterization of *Cryptosporidium ubiquitum*. *Infection,
Genetics and Evolution* **45**, 256–261 (2016).

23.   Uran-Velasquez, J. *et al.* Multilocus Sequence Typing helps understand the genetic
diversity of *Cryptosporidium hominis* and *Cryptosporidium parvum* isolated from Colombian
patients. *PLOS ONE* **17**, e0270995 (2022).

24.   Hawash, Y. DNA extraction from protozoan oocysts/cysts in feces for diagnostic PCR.
Korean J Parasitol **52**, 263–271 (2014).

25.   Corsi, G. I. *et al.* Recent genetic exchanges and admixture shape the genome and
population structure of the zoonotic pathogen *Cryptosporidium parvum*. *Molecular Ecology*
32, 2633–2645 (2023).

26.   Tichkule, S. *et al.* Global Population Genomics of Two Subspecies of *Cryptosporidium hominis* during 500 Years of Evolution. *Molecular Biology and Evolution* **39**, msac056 (2022).

27.   Gan, M., Liu, Q., Yang, C., Gao, Q. & Luo, T. Deep Whole-Genome Sequencing to Detect Mixed Infection of *Mycobacterium tuberculosis*. *PLOS ONE* **11**, e0159029 (2016).

28.   Kaupke, A., Gawor, J., Rzeżutka, A. & Gromadka, R. Identification of pig-specific *Cryptosporidium* species in mixed infections using Illumina sequencing technology. *Exp Parasitol* **182**, 22–25 (2017).

29.   Widmer, G. *et al.* Population structure of natural and propagated isolates of *Cryptosporidium parvum*, *C. hominis* and *C. meleagridis*. *Environmental Microbiology* **17**, 984–993 (2015).

30.   Guo, Y. *et al.* Isolation and Enrichment of *Cryptosporidium* DNA and Verification of DNA Purity for Whole-Genome Sequencing. *Journal of Clinical Microbiology* **53**, 641–647 (2015).

31.   Guo, Y. *et al.* Comparative genomic analysis reveals occurrence of genetic recombination in virulent *Cryptosporidium hominis* subtypes and telomeric gene duplications in *Cryptosporidium parvum*. *BMC Genomics* **16**, 320 (2015).

32.   Hadfield, S. J. *et al.* Generation of whole genome sequences of new *Cryptosporidium hominis* and *Cryptosporidium parvum* isolates directly from stool samples. *BMC Genomics* **16**, 650 (2015).

33.   Ruecker, N. J. *et al.* Tracking Host Sources of *Cryptosporidium* spp. in Raw Water for Improved Health Risk Assessment. *Appl Environ Microbiol* **73**, 3945–3957 (2007).

34.   Cama, V. *et al.* Mixed *Cryptosporidium* Infections and HIV. *Emerg Infect Dis* **12**, 1025–1028 (2006).

35.   Jiang, J., Alderisio, K. A. & Xiao, L. Distribution of *Cryptosporidium* Genotypes in Storm Event Water Samples from Three Watersheds in New York. *Appl Environ Microbiol* **71**, 4446–4454 (2005).

36.   McLauchlin, J., Amar, C., Pedraza-Díaz, S. & Nichols, G. L. Molecular Epidemiological Analysis of *Cryptosporidium* spp. in the United Kingdom: Results of Genotyping *Cryptosporidium* spp. in 1,705 Fecal Samples from Humans and 105 Fecal Samples from Livestock Animals. *J Clin Microbiol* **38**, 3984–3990 (2000).

37.   Ng-Hublin, J. S. Y., Combs, B., Mackenzie, B. & Ryan, U. Human cryptosporidiosis diagnosed in Western Australia: a mixed infection with *Cryptosporidium meleagridis*, the *Cryptosporidium* mink genotype, and an unknown *Cryptosporidium* species. *J Clin Microbiol* **51**, 2463–2465 (2013).

38.   Sharma, P., Sharma, A., Sehgal, R., Malla, N. & Khurana, S. Genetic diversity of *Cryptosporidium* isolates from patients in North India. *Int J Infect Dis* **17**, e601-605 (2013).

39.   Reed, C., Sturbaum, G. D., Hoover, P. J. & Sterling, C. R. *Cryptosporidium parvum* Mixed Genotypes Detected by PCR-Restriction Fragment Length Polymorphism Analysis. *Appl Environ Microbiol* **68**, 427–429 (2002).

40.   Korpe, P. S. *et al.* Case-Control Study of *Cryptosporidium* Transmission in Bangladeshi Households. *Clin Infect Dis* **68**, 1073–1079 (2019).

41.   Tanriverdi, S., Arslan, M. Ö., Akiyoshi, D. E., Tzipori, S. & Widmer, G. Identification of genotypically mixed *Cryptosporidium parvum* populations in humans and calves. *Molecular and Biochemical Parasitology* **130**, 13–22 (2003).

42.   Mallon, M. *et al.* Population Structures and the Role of Genetic Exchange in the Zoonotic Pathogen *Cryptosporidium parvum*. *J Mol Evol* **56**, 407–417 (2003).

43.   Gaudin, M. & Desnues, C. Hybrid Capture-Based Next Generation Sequencing and Its Application to Human Infectious Diseases. *Frontiers in Microbiology* **9**, (2018).

44.   Glenn, T. C. & Faircloth, B. C. Capturing Darwin's Dream. *Mol Ecol Resour* **16**, 1051–1058 (2016).

45.   Quek, Z. B. R. & Ng, S. H. Hybrid-Capture Target Enrichment in Human Pathogens: Identification, Evolution, Biosurveillance, and Genomic Epidemiology. *Pathogens* **13**, 275 (2024).

46.   Khan, A. *et al.* Phylogenomic reconstruction of *Cryptosporidium* spp. captured directly from clinical samples reveals extensive genetic diversity. 2024.04.17.589752 Preprint at https://doi.org/10.1101/2024.04.17.589752 (2024).

47.   Abrahamsen, M. S. *et al.* Complete genome sequence of the apicomplexan, *Cryptosporidium parvum*. *Science* **304**, 441–445 (2004).

48.   Robinson, G., Elwin, K. & Chalmers, R. M. *Cryptosporidium* Diagnostic Assays: Molecular Detection. in *Cryptosporidium: Methods and Protocols* (eds Mead, J. R. & Arrowood, M. J.) 11–22 (Springer, New York, NY, 2020). doi:10.1007/978-1-4939-9748-0_2.

49.   Baptista, R. de P., Xiao, R., Li, Y., Glenn, T. C. & Kissinger, J. C. New T2T assembly of *Cryptosporidium parvum* IOWA II annotated with Legacy-Compatible Gene identifiers. *Sci Data* **12**, 1039 (2025).

50.   Simon, M. D. & Machyna, M. Principles and Practices of Hybridization Capture Experiments to Study Long Noncoding RNAs That Act on Chromatin. *Cold Spring Harb Perspect Biol* **11**, a032276 (2019).

51.   Dąbrowska, J., Sroka, J. & Cencek, T. Investigating *Cryptosporidium* spp. Using Genomic, Proteomic and Transcriptomic Techniques: Current Progress and Future Directions. *International Journal of Molecular Sciences* **24**, 12867 (2023).

52.   Xiao, L. Molecular epidemiology of cryptosporidiosis: an update. *Exp Parasitol* **124**, 80–89 (2010).

53.   Agyabeng-Dadzie, F., Xiao, R. & Kissinger, J. C. *Cryptosporidium* Genomics — Current Understanding, Advances, and Applications. *Curr Trop Med Rep* **11**, 92–103 (2024).

54.   Gatei, W. *et al.* Multilocus sequence typing and genetic structure of *Cryptosporidium hominis* from children in Kolkata, India. *Infect Genet Evol* **7**, 197–205 (2007).

55.   Hadfield, S. J., Robinson, G., Elwin, K. & Chalmers, R. M. Detection and Differentiation of *Cryptosporidium* spp. in Human Clinical Samples by Use of Real-Time PCR ▽ . *J Clin Microbiol* **49**, 918–924 (2011).

56.   Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. VSEARCH: a versatile open source tool for metagenomics. *PeerJ* **4**, e2584 (2016).

57.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

58.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

59.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**, W160-165 (2016).

60.   Robinson, J. T. *et al.* Integrative genomics viewer. *Nat Biotechnol* **29**, 24–26 (2011).

61.   Huang, W., Li, L., Myers, J. R. & Marth, G. T. ART: a next-generation sequencing read simulator. *Bioinformatics* **28**, 593–594 (2012).

62.   Paparini, A. *et al.* Comparison of Sanger and next generation sequencing performance for genotyping *Cryptosporidium* isolates at the 18S rRNA and actin loci. *Exp Parasitol* **151–152**, 21–27 (2015).

63.   Puiu, D., Enomoto, S., Buck, G. A., Abrahamsen, M. S. & Kissinger, J. C. CryptoDB: the *Cryptosporidium* genome resource. *Nucleic Acids Res* **32**, D329-331 (2004).

64.   Bushnell, B. BBMap: A Fast, Accurate, Splice-Aware Aligner. (2014).

65.   Lindgreen, S., Adair, K. L. & Gardner, P. P. An evaluation of the accuracy and speed of metagenome analysis tools. *Sci Rep* **6**, 19233 (2016).

66.   Glenn, T. C. *et al.* Adapterama I: universal stubs and primers for 384 unique dual-indexed or 147,456 combinatorially-indexed Illumina libraries (iTru & iNext). *PeerJ* **7**, e7755 (2019).

67.   Stroup, S. E. *et al.* Real-time PCR detection and speciation of *Cryptosporidium* infection using Scorpion probes. *J Med Microbiol* **55**, 1217–1222 (2006).

68.   Rohland, N. & Reich, D. Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture. *Genome Res* **22**, 939–946 (2012).

69.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

70.   Penumarthi, L. R., Baptista, R. P., Beaudry, M. S., Glenn, T. C. & Kissinger, J. C. A new chromosome-level genome assembly and annotation of *Cryptosporidium meleagridis*. *bioRxiv* 2024.02.16.580748 (2024) doi:10.1101/2024.02.16.580748.

71.   Kuhn, M. & Davis, V. parsnip: A Common API to Modeling and Analysis Functions. (2025).

72.   McKenna, A. *et al.* The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

73.   Knaus, B. J. & Grünwald, N. J. vcfr: a package to manipulate and visualize variant call format data in R. *Molecular Ecology Resources* **17**, 44–53 (2017).

74.   Mijangos, J. L., Gruber, B., Berry, O., Pacioni, C. & Georges, A. dartR v2: An accessible genetic analysis platform for conservation, ecology and agriculture. *Methods in Ecology and Evolution* **13**, 2150–2158 (2022).

75.   Zheng, X. *et al.* A high-performance computing toolset for relatedness and principal component analysis of SNP data. *Bioinformatics* **28**, 3326–3328 (2012).

76.   Dray, S. & Dufour, A.-B. The ade4 Package: Implementing the Duality Diagram for Ecologists. *Journal of Statistical Software* **22**, 1–20 (2007).

77.   Schliep, K. P. phangorn: phylogenetic analysis in R. *Bioinformatics* **27**, 592–593 (2011).

78.   Huson, D. H. SplitsTree: analyzing and visualizing evolutionary data. *Bioinformatics* **14**, 68–73 (1998).

79.   Kamvar, Z. N., Tabima, J. F. & Grünwald, N. J. Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**, e281 (2014).

80.   Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**, 526–528 (2019).

81.   Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using multilocus genotype data. *Genetics* **155**, 945–959 (2000).

82.   Pina-Martins, F., Silva, D. N., Fino, J. & Paulo, O. S. Structure_threader: An improved method for automation and parallelization of programs structure, fastStructure and MavericK on multicore CPU systems. *Molecular Ecology Resources* **17**, e268–e274 (2017).

83.   Earl, D. A. & VonHoldt, B. M. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genet Resour* **4**, 359–361 (2012).

84.   Wickham, H., Hester, J. & Bryan, J. readr: Read Rectangular Text Data. R package version 2.1.5. *https://github.com/tidyverse/readr* https://github.com/tidyverse/readr (2024).

85.   Tonkin-Hill, G. & Lee, S. Starmie: Population structure model inference and visualisation. (2016).

86.   Goudet, J. hierfstat, a package for r to compute and test hierarchical F-statistics. *Molecular Ecology Notes* **5**, 184–186 (2005).

87.   Turner, S. D. qqman: an R package for visualizing GWAS results using Q-Q and manhattan plots. *Journal of Open Source Software* **3**, 731 (2018).

88.   Wilke, C. ggridges: Ridgeline Plots in 'ggplot2'. (2024).

89.   Zhu, S. J., Almagro-Garcia, J. & McVean, G. Deconvolution of multiple infections in Plasmodium falciparum from high throughput sequencing data. *Bioinformatics* **34**, 9–15 (2018).

90.   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).

91.   Lenth, R. V. Least-Squares Means: The R Package lsmeans. *Journal of Statistical Software* **69**, 1–33 (2016).

92.   Wickham, H. *et al.* Welcome to the Tidyverse. *Journal of Open Source Software* **4**, 1686 (2019).

93.   Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2016).

94. Plotly Technologies Inc. Collaborative data science. https://plot.ly (2015).

Tables

Table 1. Summary of *Cryptosporidium* species considered, genome version used, and criteria for designing CryptoCap_100k bait set.

Identifier	Reference Genome	Criteria (Identity: Overlap)	Number of Baits
Whole Genome			
C. parvum	GCA_015245375.1	Reference	73,672
C. hominis	GCA_001483515.1	90%:50%	2,851
C. cuniculus	GCA_004337835.1	90%:50%	738
C. tyzzeri	GCA_007210665.1	90%:50%	858
C. meleagridis	GCA_001593445.1	85%:45%	9,034
C. viatorum	GCA_004337795.1	85%:45%	11,774
***C. viatorum* (select genes)**		50%:95%	943
18S rRNA		50%:95%	63
gp60		NA	67
		Total Number of Baits	100,000

GC Content	30.1%
Soft Masked	2.8% (baits ranged from 0-50% masked)

Table 2. Bait characterization for CryptoCap_100k and CryptoCap_75k by mapping all baits to all available reference genome sequences. Mean depth, percent breadth of genome coverage, and percentage of baits mapped to each species' genome sequence. We break down by species included and excluded from bait design.

Species Information		CryptoCap_100k 100,000 baits (This study)			CryptoCap_75k 68,966 baits (Khan and Alves-Ferreira et al., 2024)		
		Mean depth	% Breadth of coverage	% Baits mapped	Mean depth	% Breadth of coverage	% Baits mapped
Species included in CryptoCap_100k bait design	*C. parvum* IOWA II	1.086	97.55	84.98	0.909	90.89	100
	C. parvum IOWA ATCC	1.089	97.83	85.41	0.907	90.63	99.96
	C. hominis UdeA01	1.095	97.75	85.36	0.904	90.33	99.31
	C. hominis 30976	1.097	97.88	85.64	0.903	90.25	99.39
	C. cuniculus UKCU2	0.853	77.86	70.09	0.751	74.93	86.50
	C. tyzzeri UGA55	1.074	96.75	83.66	0.900	89.95	98.75
	C. meleagridis UKMEL1	0.997	91.42	80.45	0.793	79.23	91.27
	C. viatorum (partial inclusion) ABER_CVIA_1.0	0.714	66.68	60.99	0.529	51.40	64.36
Species not included in CryptoCap_100k bait design	*C. andersoni* 30847	0.008	0.52	1.08	0.005	0.45	0.87
	C. baileyi TAMU-09Q1	0.030	2.84	3.45	0.025	2.43	3.70
	C. muris RN66	0.008	0.60	1.12	0.005	0.47	0.91
	C. ubiquitum 39726	0.262	25.39	24.81	0.239	23.86	31.69

Table 3. Summary statistics from enriching 100 human fecal DNA samples with two library preparation methods using CryptoCap_100k baits and obtaining Illumina PE150 reads. Source tables are Tables S13 and S14.

Average values	iTru Libraries			iNextEra Libraries		
	Not enriched	Single enriched		Not enriched	Single enriched	Double enriched
% retained after quality filtering	61.53	83.87		89.70	90.79	94.48
% reads on target *C. parvum*	0.04	35.62		0.033	15.85	64.98
% reads on target 10-CrypGS	0.11	41.23		0.100	17.90	65.78
Enrichment factor *	N/A	2,095		N/A	1,112	10,700**
Depth per 100k *	0.00	1.12		0.001	0.563	2.702
Breadth per 100k *	0.00	2.25		0.065	1.79	5.99
Number of samples	73	100		91	91	91

*(matching *C. parvum*)

** This value corresponds to the enrichment factor of double-enriched samples compared to non-enriched counterparts, which is not displayed in Tables S13 and S14.

Figures

Figure 1. Species assignment plots based on mapping reads from unenriched and enriched libraries for one mock mixed-infection sample. (A) The number of raw paired reads obtained for each species in the mix, *C. parvum* in green and *C. meleagridis* in blue, for both library types. The pink lines indicate the expected number of reads for each sample for a 50-50 proportion. Relative abundance of species assigned from mapping the sequence reads to (B) the ten-genome sequence reference database (10-CrypGS), (C) the *gp60* database, and (D) the *18S rRNA* database.

Figure 2. Summary statistics for sequence data from serially diluted *C. parvum* samples to assess detection levels given different input amounts. (A) *C. parvum* genome depth and breadth of coverage for diluted DNA samples prepared with NEB/ iTru libraries (B) species assignment of diluted *C. parvum* samples prepared with NEB-iTru protocol and unenriched (*U*) or single-enriched (*S*) mapped to the ten *Cryptosporidium* genome sequences reference (10-CrypGS); (C) *C. parvum* genome depth and breadth coverage for diluted DNA samples prepared with iNext/NextEra libraries, (D) species assignment of diluted *C. parvum* samples prepared with iNextEra protocol and unenriched (*U*), single-enriched (*S*), or double-enriched (*D*) mapped to the ten *Cryptosporidium* genome sequences reference (10-CrypGS). In the species assignment plots, the first letter in the sample name corresponds to the library type, and the second letter corresponds to the level of input of DNA (A–G). The direction of the arrow indicates high to low input (A → G).

Figure 3. Summary statistics for sequence data from the 100 clinical samples prepared with the NEB protocol and iTru primers, comparing libraries enriched with CryptoCap_100k bait set versus unenriched libraries for validation. (A) Scatter plot of percentage of reads mapped from one round of enrichment compared to unenriched reads for each sample (x-axis labels not shown but in order from left to right sample 1 to sample 100). (B) Genome depth and breadth of coverage for each sample in enriched libraries compared to unenriched. Linear regression model for predicting the percentage of mapped reads to a reference database containing ten *Cryptosporidium* genome sequences (10-CrypGS) given a C_T value from qPCR for (C) unenriched libraries and (D) enriched libraries with CryptoCap_100k. Species assignment when mapped to 10-CrypGS for (E) libraries left unenriched, and (F) enriched libraries with CryptoCap_100k. Sample names for plot 3E are not shown.

Figure 4. Summary statistics for sequence data from 91 clinical samples prepared with the iNextEra protocol and iNext primers, comparing libraries enriched with CryptoCap_100k bait set versus unenriched libraries for validation. (A) Boxplot of percentage of reads mapped to *C. parvum* from double-enriched [D], single-enriched [S], and unenriched libraries[U]. (B) Genome depth and breadth of coverage for each sample in double-enriched, single-enriched, and unenriched libraries. Linear regression model for predicting the percentage of mapped reads to a reference database containing ten *Cryptosporidium* genome sequences (10-CrypGS) given a C_T value from qPCR for (C) unenriched libraries and (D) single-enriched libraries with CryptoCap_100k, and (E) double-enriched libraries with CryptoCap_100k. Species assignment of each sample prepared with iNextEra protocol with (F) no further enrichment, (G) single-enriched with CryptoCap_100k, and (H) double-enriched with CryptoCap_100k; when mapped to a reference database containing ten

Cryptosporidium genome sequences (10-CrypGS). Sample names for plot 4F are not shown.

Figure 5. Population genomic analyses from called SNP variants of clinical samples prepared with the iNextEra protocol and iNext primers. Libraries were double-, single-enriched, and unenriched, respectively (A, B, and C). Each panel includes: Principal Components Analyses, phylogenetic distance networks, UPGMA Phylogenetic trees of Nei's genetic distances, assignment plots from Structure analyses for K=2 (selected based on Evanno's method—unenriched samples could not be analyzed due to high levels of missing data), and effective (gray bars) and inferred (blue circles) number of strains (K) detected in each sample by the DEploid software.

Figure 6. Laboratory workflow used for the analysis of a set of clinical samples from the UK. The genomic DNA of each patient's sample was extracted, and *Cryptosporidium* was validated via qPCR. Each DNA extract was used for two types of library prep, NEB Ultra FSII and iNextEra library prep. Unenriched library preparation aliquots were used for sequencing. Also, aliquots from unenriched libraries were pooled according to library type, and enrichments were performed with CryptoCap_100k on these pools. Aliquots of these products, referred to as single-enriched products, were used for sequencing. Aliquots of single-enriched libraries, which used the iNextEra protocol, were pooled, and enrichments were performed again with CryptoCap_100k. Aliquots of these products, named double-enriched products, were used for sequencing. Colors represent those used throughout the manuscript to differentiate unenriched,

single-enriched, and double-enriched products. Created in BioRender. Bayona N (2025)

https://BioRender.com/yysbjqh.