1 **Investigating the 'Atypical Rhythm Risk' hypothesis in children with**

2 **developmental language disorder using an EEG rhythmic speech paradigm**

4 Mahmoud Keshavarzi[1], Susan Richards, Georgia Feltham, Lyla Parvez and Usha Goswami

5
6
7
8 Centre for Neuroscience in Education, Department of Psychology, University of Cambridge,
9 Cambridge, CB2 3EB, United Kingdom
21 [1]Corresponding Author: mk919@cam.ac.uk, Mahmoud.keshavarzi.ir@ieee.org

Abstract

Sensitivity to rhythmic and prosodic cues in speech has been described as a precursor of language acquisition. Consequently, atypical rhythmic processing during infancy and early childhood has been considered a risk factor for developmental language disorders. This is the 'atypical rhythm risk' (ARR) hypothesis. The neural processing of rhythmic speech has not yet been explored in children with developmental language disorder (DLD). Here we utilise EEG and a rhythmic speech paradigm previously utilised for dyslexia to investigate the ARR hypothesis in 9-year-old children with and without DLD. We investigate angular velocity, power, ERPs, phase-amplitude coupling (PAC) and phase-phase coupling (PPC), at three frequency bands, delta, theta and low gamma. Both groups demonstrated significant and equivalent phase entrainment in the delta and theta bands, but only the control children showed significant phase entrainment in the low gamma band. Further, while angular velocity in the delta and theta bands was equivalent by group, there was a significant gamma-band difference. The children with DLD also exhibited significantly more theta and gamma power compared to the control children. Regarding PAC and PPC, both groups showed significant and equivalent coupling strength. However, group resultant phase differences showed that low-frequency phase (delta and theta) affected gamma oscillations differently by group. The EEG data show important differences between children with and without DLD in the neural mechanisms underpinning the processing of rhythmic speech. The findings are discussed in terms of auditory theories of DLD.

Keywords

Developmental language disorder, rhythmic audio-visual speech, EEG, neural phase entrainment, atypical rhythm risk

1. Introduction

A growing number of cognitive behavioural studies have investigated rhythmic and/or musical processing in children with developmental language disorder (DLD), and have shown that atypical rhythm processing is associated with speech and language processing problems (e.g., Corriveau et al., 2007; Corriveau & Goswami, 2009; Przybylski et al., 2013; Cumming et al., 2015a,b; Richards & Goswami, 2015, 2019; Sallat & Jentschke, 2015; Bedoin et al., 2016). This apparent link between rhythm processing and language processing has been described by the Atypical Rhythm Risk (ARR) hypothesis (Ladányi et al., 2020). Following a review of the literature, Ladányi et al. (2020) proposed that individuals with atypical rhythm processing are at higher risk for developmental speech/language disorders, including developmental dyslexia, DLD and stuttering. Disordered development of oral language (DLD) affects between 3 – 7% of children in all cultures (Bishop et al., 2017). The clinical diagnosis for such children was previously Specific Language Impairment (SLI), however the multinational and multidisciplinary CATALISE study recommended a change in terminology to DLD (Bishop et al., 2017). Children given a diagnosis of DLD have a range of language difficulties that cause functional impairments in their daily lives, yet have no known biomedical aetiology. The core dimensions of difficulty are problems with syntax and grammar, problems with word semantics, impairments in verbal memory and problems with phonology (see Leonard, 2014; Bishop, 2013). The possibility that atypical *neural* processing of rhythm is associated with DLD has not yet been investigated, and is the focus of the current study.

At the sensory/neural level, the ARR hypothesis is captured by Temporal Sampling (TS) theory (Goswami, 2011; 2015, 2022). TS theory was originally proposed to explain the rhythmic impairments found in children with developmental dyslexia, a disorder of written language processing that presents with core phonological impairments (Stanovich, 1988; Swan & Goswami, 1997; Snowling, 2000). Prior TS-driven behavioural studies of rhythmic

processing by children with dyslexia and children with DLD have shown many similarities,

such as impaired sensory discrimination of amplitude rise times (important acoustic cues to

rhythm, Corriveau et al., 2007; Corriveau & Goswami, 2009; Fraser et al., 2010; Beattie &

Manis, 2012; Cumming et al., 2015a; Richards & Goswami, 2015, 2019) and relative

insensitivity to syllable stress patterns in words and phrases (Cumming et al., 2015a,b; Richards

& Goswami, 2015, 2019). TS theory suggests that the sensory/neural processing of rhythm is

impaired because acoustic processing of the amplitude envelope of both speech and musical

signals is atypical in children with developmental disorders of language, particularly at

amplitude modulation rates <10 Hz that are known to carry information about rhythm and (for

speech) syllable stress patterns (see Greenberg, 2006). The amplitude envelope of speech is

known to be tracked cortically, possibly via entrainment of endogenous low-frequency

oscillations <10 Hz (corresponding to the electrophysiological delta and theta bands, 0.5-4 Hz

and 4-8 Hz, respectively) to the speech envelope (Giraud & Poeppel, 2012). Regarding the

neural side of TS theory, prior TS studies with dyslexic children have documented neural

differences compared to typically-developing (TD) children during the processing of both

speech and non-speech rhythmic stimuli (Power et al., 2013; Keshavarzi et al., 2022; Peter et

al., 2022). For speech stimuli, children's neural processing of rhythm has been investigated

using the syllable repetition task (Power et al., 2012). In this task, EEG is recorded while

children with dyslexia listen to the speech syllable "ba" presented audio-visually at a repetition

rate of 2 Hz (Power et al., 2013; Keshavarzi et al., 2022). Both 9-year-old (Keshavarzi et al.,

2022) and 13-year-old (Power et al., 2013) children with dyslexia show atypical delta-band

phase synchronisation in the syllable repetition task, exhibiting a different preferred phase to

children without dyslexia for the delta band only. The theta and alpha bands are not affected.

Both Power et al. (2013) and Keshavarzi et al. (2022) reported that neural responding in the

delta band was aligned to a different temporal point for each group, despite the highly

predictable rhythmic input. These studies suggest that there is an optimal (or preferred) phase

of entrainment regarding the accurate and efficient processing of rhythmic inputs, and that this

preferred phase differs in the dyslexic brain. To date, the rhythmic syllable repetition task has

not been administered to children with DLD. We provide such a study here.

Prior neural studies of children with DLD have not to date employed rhythmic

processing tasks nor investigated the TS framework. The absence of neural rhythm studies is

surprising, as adult neural studies show that temporal prediction of the location of stressed

syllables facilitates syntactic processing (Kotz & Schmidt-Kassow, 2015). Linguistic analyses

show that adults produce a stressed syllable on average every 500ms, a "beat rate" of 2Hz

(Dauer, 1983), a finding that underpins the choice of repetition rate in the current task. Prior

DLD studies have instead primarily utilised fMRI and have sought structural and functional

differences that may throw light on the aetiology of this developmental disorder. For example,

there has been interest in whether hemispheric asymmetries during linguistic processing are

different when children with DLD are compared to TD children (Bishop, 2013). While

laterality per se has not proved conclusive (Parker et al., 2022), one of the most consistent

findings in the functional fMRI literature with DLD children is that left frontal and temporal

cortical areas show reduced activation during language processing (Asaridou & Watkins, 2022,

for review). Children with DLD also show microstructural differences in dorsal anatomical

pathways that connect frontal and temporal areas, as well as structurally-atypical basal ganglia

(Badcock et al., 2012; Lee et al., 2013; Watkins et al., 2002). Although both the temporal areas

and the basal ganglia are important for speech processing, speech motor learning and temporal

prediction (see Kotz & Schmidt-Kassow, 2015), fMRI studies have not revealed which neural

mechanisms associated with these areas could be affected in DLD. There are also prior EEG

and MEG studies with DLD children, but again these studies have not adopted a rhythmic nor

a speech tracking focus, and have not thrown light on underlying mechanisms (e.g. Shafer et

al., 2001). Many of the prior child EEG studies concern the overlap between DLD diagnoses

and epileptic seizures (e.g. Mehta et al., 2015; Systad et al., 2019). However, a series of EEG

studies with German-learning infants has shown that those at family risk for DLD show

impaired neural ERP markers of language learning, such as a delayed N400 (thought to index

semantic integration, Friedrich & Friederici, 2005, 2006). ERP studies with English-learning

infants at family risk for DLD using a non-speech tone pair task have shown that at-risk infants

have ERPs with larger amplitudes than TD infants when the tones follow each other rapidly in

time (gap of 70 ms, Benasich et al., 2006), and also show a longer latency (in the N250) which

is right-lateralized. Meanwhile, prior MEG studies of DLD using speech processing tasks are

consistent with the functional fMRI literature in showing atypical left hemisphere temporal

function (Helenius et al., 2009, 2014).

The TS framework (TSF) differs from these prior fMRI and MEG literatures in

predicting impairments in right-lateralised neural mechanisms during language processing for

children with DLD (Goswami, 2022, for review). However, the TSF is not based on laterality

differences, rather the right hemisphere predictions arise from the nature of the neural

mechanisms of speech encoding that are proposed to be impaired. These mechanisms involve

the oscillatory encoding of slower amplitude modulation (AM) information in the speech

amplitude envelope, also called cortical speech tracking. Slower AM information in speech is

thought to be preferentially processed in the right hemisphere, while faster AM information is

thought to be processed in the left hemisphere (Boemio & Poeppel, 2005; see also Rios-Lopez

et al., 2020, 2022, for child data). For linguistic inputs, these hemispheric preferences appear

to change during the acquisition of literacy when orthographic representations are acquired

(Cutini et al., 2016; Lallier et al., 2017). TS theory proposes atypical sensory/neural processing

of the slower rates of AM nested in the speech amplitude envelope (AE) for children with DLD

(Goswami, 2011, 2015, 2022). The AE is the slow-varying energy contour of speech that aids

the perception of rhythm (Greenberg, 2006). The AE contains a range of AM patterns, created

by hierarchical nesting of AMs at different temporal rates that are functionally connected by

phase-phase and phase-power relations (Ghitza & Greenberg, 2009; Giraud & Poeppel, 2012;

Goswami & Leong, 2013; Gross et al., 2013). For example, computational speech modelling

studies show that the phase relations between these different AM rates (delta-theta, theta-

beta/low gamma) provide systematic statistical cues to both phonological units such as stressed

vs unstressed syllables, syllables, and onset-rimes (acoustic-emergent phonology, Leong &

Goswami, 2015) and also to aspects of grammar such as plural marking (Flanagan & Goswami,

2018). By TS theory, therefore, impaired cortical tracking of one or more of the AM rates nested

in the AE, or of the phase-amplitude or phase-phase relations between different AM bands,

could cause atypical development of one or more aspects of language.

Recent infant data support the linguistic predictions made by TS theory. In the

Cambridge UK BabyRhythm project, 120 infants were followed longitudinally from the age of

2 months, and EEG was recorded while the infants listened to either sung speech (nursery

rhymes), to the rhythmic syllable repetition task, or to a matched drumbeat (nonspeech)

rhythmic repetition task (Attaheri et al., 2022a,b; Ni Choisdealbha et al., 2022, 2023;

Keshavarzi et al., 2023). In the syllable repetition task, infants showed evidence for a group

preferred phase as early as 2 months of age, and by 6 months individual phase consistency was

present over both left- and right-temporal regions as well as over parietal-occipital regions (Ni

Choisdealbha et al., 2023). Individual phase angles predicted language acquisition at 18 months

as measured by an infant-led vocabulary task (in this task, the infant selects one of two pictures

that matches a spoken word, Ni Choisdealbha et al., 2023). In the rhythmic drumbeat

(nonspeech) task, neural phase angle at 9 months predicted nonword repetition at 24 months, a

phonological measure (Ni Choisdealbha et al., 2023). Accordingly, delta-band phase angles to

both speech and nonspeech rhythmic stimuli predict later language outcomes for a TD sample.

For the nursery rhyme analyses, significant cortical tracking of the low-frequency envelope

information was present in both delta and theta bands by 4 months of age, with no significant

tracking present for faster bands (alpha band, see Attaheri et al., 2022a). Phase-amplitude

coupling (PAC) between both delta and theta phase and beta and gamma amplitudes was

present from 4 months, and showed no change over development for this sample. When

language outcomes were assessed, only delta band cortical tracking and theta-gamma PAC

predicted later vocabulary and phonology measured at 24 months (Attaheri et al., preprint).

Grammar measures were included in the UK BabyRhythm study, but did not show sufficient

individual differences for neural prediction analyses (Rocha et al., 2024). Nevertheless, these

prior TSF-driven infant data suggest that both delta-band phase angle and theta-gamma PAC

in the current study may be associated with DLD status.

There is also one prior study of rhythmic speech processing using a modulation band

manipulation with children with DLD, which would predict atypical gamma band speech

tracking. Goswami et al. (2016) used a nursery rhyme recognition task based on speech that

was either low-pass filtered (< 4 Hz, providing delta band information only) or band-pass

filtered (~33Hz, providing faster low gamma band modulations only) so that the speech input

was degraded and difficult to recognise. The children's task was to recognise the matching

nursery rhyme by making a selection in a picture-based multiple choice format. Goswami et

al. reported that 9-year-old children with DLD were worse than TD age-matched children when

selecting bandpass-filtered targets, but not when selecting low-pass filtered targets. Children

with both DLD and dyslexia (DLD children with phonological impairments) were worse than

TD children at recognising the nursery rhymes in both filtered speech conditions. All groups

of children found it easier to recognise the low-pass filtered rhymes (e.g. children with DLD

recognised 89% of the low-pass filtered rhymes compared to 42% of the bandpass filtered

rhymes, while the TD children recognised 94% and 59% respectively). When relations with

language development were investigated in the full sample of 9-year-olds (N = 95), accuracy

in the bandpass-filtered speech condition was the stronger predictor of language outcomes,

accounting for 12% of unique variance in the Clinical Evaluation of Language Fundamentals

(CELF-III, Wiig, Semel & Secord, 2017) receptive language measures and 10% of unique

variance in CELF expressive language scores. By contrast, accuracy in the low-pass filtered

speech condition accounted for 9% of unique variance in the phonological measure (rhyme

awareness), but only 3% of unique variance in the CELF ($p < 0.05$). To interpret their data,

Goswami et al. (2016) suggested that although prior TSF studies of children with dyslexia and

children with DLD indicate a shared sensory difficulty in processing temporal modulation

patterns in speech, the neural temporal integration windows that are most impaired may differ

for each disorder. Children with DLD may have more difficulties with faster temporal

integration windows in speech-based tasks (gamma band difficulties).

Accordinly, we explored a number of *a priori* predictions in the current study. First, on

the basis of the shared behavioural rhythmic features of dyslexia and DLD and our prior TSF

dyslexia data, we expected atypical delta-band entrainment (reduced phase consistency) and a

different group preferred phase for the participants with DLD compared to the age-matched

TD control children. However, given the filtered speech findings discussed above (Goswami

et al., 2016), and in contrast to the earlier rhythmic syllable repetition studies of children with

dyslexia (Power et al., 2013; Keshavarzi et al., 2022), gamma band oscillations were measured

in the current study as well. Given Goswami et al.'s (2016) finding of rhythmic speech

processing difficulties at faster modulation rates, a second prediction was possible gamma band

oscillatory group differences. Third, on the basis of our TSF infant data, we tentatively

predicted atypical theta-gamma PAC for the children with DLD. The prior study of children

with dyslexia that explored angular velocity (Keshavarzi et al., 2022) had also reported that the

group angular velocity in the delta band for the children with dyslexia was almost twice the

rate of the TD control group, suggestive of different changes in phase across time for the

dyslexic group. Accordingly, a final tentative prediction was that angular velocity in the delta

band may differ between children with and without DLD. We note that the prior literature on

sensory impairments in rapid auditory non-speech (tone) processing in infants at family risk

for DLD would also predict gamma band impairments for the DLD children (Ortiz-Mantilla &

Benasich, 2013).

2. Methods and materials

2.1. Participants

Sixteen typically developing children (mean age of 109 ± 13 months) and sixteen children with

DLD (mean age of 108 ± 11 months) took part in the study. All participants were taking part in

an ongoing study of auditory processing in DLD and had English as the main language spoken

at home (see Parvez et al., 2024). All participants exhibited normal hearing when tested with

an audiometer. In a short hearing test across the frequency range $0.25 - 8$ kHz (0.25, 0.5, 1, 2,

4, 8 kHz), all children were sensitive to sounds within the 20 dB HL range. DLD status was

assessed by administering two expressive and two receptive subtests of the CELF (Wiig, Semel

& Secord, 2017). The subtests were recalling sentences, formulating sentences, and (depending

on age) two of the word structure, sentence comprehension, word classes and semantic

relationships subtests. Children who scored below the mean (7 or less when the mean score is

10) on at least 2 of these subtests were included as DLD. The control children also received

either one or two of the CELF tasks, recalling sentences (N = 16) and formulating sentences

(N = 11) and scored in the normal range. The Picture Completion subtest of the Wechsler

Intelligence Scale for Children (WISC IV, Wechsler, 2016) was used to assess non-verbal

intelligence (NVIQ) and did not differ between groups. Group performance for the tasks

administered to both groups is shown in Table 1.

Table 1. Details of the participating children, showing Scaled Scores (ScS, population mean
= 10) and Standard Scores (SS, population mean = 100).

	DLD	Age-Matched Control (TD)
N	16	16
Age (months) [A]	108.3 (11.3)	108.9 (13.3)
Recalling Sentences ScS [B]	5.3 (1.7)	10.9 (2.3)
Formulating Sentences ScS [B]	4.4 (1.9)	10.8 (1.7)
WISC Picture Completion ScS [A]	7.9 (3.0)	9.9 (2.8)

248 Note. **A**: TD and DLD groups are not significantly different; **B**: DLD group significantly worse than TD group, *p*
249 < 0.01. WISC: Wechsler Intelligence Scale for Children.
250

2.2. Experimental set up and stimuli

The experimental set up and stimuli used in the EEG study were identical to those described

by Keshavarzi et al. (2022, 2024). After putting the EEG cap, the children were seated in an

electrically shielded soundproof room. They watched (on the screen) and listened (through

earphones) to audio-visual stimuli and the EEG data were collected at a sampling rate of 1 kHz

using a 128-channel EEG system (HydroCel Geodesic Sensor Net). The stimuli were a talking

head presenting rhythmic sequences of the syllable "ba" at a rate of 2 Hz with both visual and

additory information present. Visual cues began 68 ms before the onset of the stimulus "ba" as

in natural speech. Each trial (sequence) consisted of sequence of 14 syllable "ba" with both

visual and auditory information present (See Figure 1). One of the 14 syllables (either 9th, 10th

or 11th syllable) in each trial could be out of time (rhythmic violation), in a random order. The

temporal delay was calibrated individually and adaptively for each child, so that the oddball

was detected around 79.4% of the time by a button press. Each child was presented with 90

trials, 15 of which did not have a violation and were presented randomly as catch trials. Each

trial contained three periods (see Figure 1): entrainment period, violation period, and return-

to-isochrony period. Children were instructed to concentrate on the lips of the female face on

the screen and to listen to the auditory stimuli. They were also asked to press a target key on

the keyboard when one of the syllables was out of time. The total time for the EEG recording

(excluding the set-up) was about 15 min. Via the adaptive procedure, the success rate in the

behavioural task was equated for all participants.

Figure 1. Experimental stimuli. Panel A shows the waveform of a single syllable "ba", panel B
indicates an example trial (sequence of "ba" stimuli) with the violation at position of 10th
stimulus, and panel C shows the spectrogram corresponding to sequence in panel B. This figure
reproduced with permission from Keshavarzi et al., 2022.

2.3. EEG data pre-processing and analyses

The EEG data were first referenced to Cz and then band-passed filtered into 0.2 – 48 Hz using

a zero phase FIR filter with low cutoff (−6 dB) of 0.1 Hz and high cutoff (−6 dB) of 48.1 Hz

(EEGLab Toolbox; Delorme and Makeig, 2004). The bad channels were identified and

interpolated using spherical interpolation (EEGLab Toolbox). The Independent Component

Analysis (provided by EEGLab Toolbox) was run for each participant and the calculated

independent components were then evaluated very carefully to remove artefactual components

such as eye blinks and eye movements. The pre-proceed EEG data were downsampled to

100 Hz and band-pass filtered into delta (0.5 – 4 Hz), theta (4 – 8 Hz) and low gamma band

(25 – 40 Hz) frequency bands (MNE Library-Python, Gramfort et al., 2013). The data were

then epoched into individual trials, in the time interval of 0.5 s before the onset of the first "ba"

stimulus and 4 s after that. In this study, the analyses only were focused on the EEG data

corresponding to stimuli 4th-8th in the entrainment period of each trial to ensure that

rhythmicity has occurred.

2.4. Computation of angular velocity

The angular velocity, $w(t)$, for signal $s(t)$ over time interval $\Delta t = t_1 - t_2$ is calculated as:

$$w(t) = \frac{\Delta\varphi_t}{\Delta t} \qquad\qquad (1)$$

where $\Delta\varphi_t$ refers to the changes in the instantaneous phase ($\varphi(t)$) of the signal over the time

interval Δt. The instantaneous phase is calculated as:

$$\varphi(t) = \arctan\left(\frac{s_H(t)}{s(t)}\right) \qquad\qquad (2)$$

where $s_H(t)$ is the Hilbert transform of signal $s(t)$, and is calculated as:

$$s_H(t) = \frac{1}{\pi}\int_{-\infty}^{\infty} \frac{s(\tau)}{t-\tau}\, d\tau \qquad\qquad (3)$$

2.5. Computation of phase entrainment

To check the phase entrainment for each group in each frequency band, we performed the

following steps (Keshavarzi et al., 2022):

(1) Calculating instantaneous phases of all 128 EEG channels at the times corresponding to

onsets of the 5 "ba" stimuli (4th-8th) in each trial.

(2) Computing mean phase for each trial by averaging across the phase values obtained for all

EEG channels and for the 5 "ba" stimuli in step 1. This resulted in a single-phase value for

each trial.

(3) Considering a single unit vector whose angle is determined by the phase value obtained in

step 2 for each trial.

(4) Calculating the mean vector for each child by averaging across the unit vectors in step 3.

This resulted in a single vector for each child, called *child resultant vector*.

(5) Considering a single unit vector whose angle is determined by the phase of *child preferred*

vector in the vector space for each child.

(6) Calculating the mean vector for each group by averaging across the unit vectors obtained

in step 5. This results in a single vector, called *group resultant vector*, for each group.

2.6. Computation of event related potential

To obtain the time-domain ERP for each group and each frequency band, the following steps

were conducted: (1) Calculating the mean voltage amplitude for each trial by averaging across

the voltage amplitudes of all 128 pre-processed for that each trial; (2) Calculating the mean

voltage amplitude for each participant by averaging across the mean voltage obtained (in step

1) for all trials of that participant; (3) Calculating the ERP for each group by averaging across

the mean voltage obtained (in step 2) for all participants in that group.

2.7. Computation of the band-power

To investigate the power of neural response over the entrainment period used for analysis, we

calculated the power of EEG responses for each child and each group using in four steps: (1)

Calculating the power of each EEG channel separately for each trial; (2) Calculating the power

for each trial by averaging across power obtained (in step 1) for channels; (3) Calculating the

power for each participant by averaging across power obtained (in step 2) for all trials of that

participant; (4) Calculating the power for each group by averaging across power values

obtained (in step 2) for all children in that group.

2.8. Computation of cross frequency phase-amplitude coupling

Cross frequency phase-amplitude coupling refers to the modulation of amplitude of signal at a

high frequency band by the phase of low frequency oscillation. Here we quantified the strength

of this type of modulation using modulation index (*MI*; Tort et al. 2008; Hülsemann, 2019):

$MI = \frac{KL\,(U,X)}{logB}$ (4)

where B (=18) is the number of bins, U refers to the uniform distribution, X is the distribution

of the data, and $KL\,(U,X)$ is Kullback–Leibler distance which is calculated as:

$KL\,(U,X) = logB - H(P)$ (5)

where $H(.)$ is the Shannon entropy and P is the vector of normalized averaged amplitudes per

phase bin which is calculated as:

$P(j) = \frac{\bar{a}}{\sum_{k=1}^{N}\bar{a}_k}$, $j = 1, ..., N$ (6)

where $\bar{a}$ is the average amplitude of each bin, k refers to running index for the bins. Note that

P is a vector with N elements.

2.9. Computation of cross frequency phase-phase coupling

Cross-frequency phase-phase coupling (PPC) refers to the phase synchrony between

oscillations in two frequency bands. PPC between two oscillations is a value between 0 and 1,

calculated as:

$PPC = \frac{1}{T}|\sum_{t=1}^{T} e^{-j\Delta\varphi(t)}|$ (7)

where $|.|$ refers to the absolute value operator, T is the number of time samples, and $\Delta\varphi(t)$ is

the phase difference, and is calculated as:

$\Delta\varphi(t) = \varphi_1(t) - \varphi_2(t)$ (8)

where $\varphi_1(t)$ is the instantaneous phase of first signal and $\varphi_2(t)$ is the instantaneous phase of

the second signal.

3. Results

3.1. Phase entrainment consistency in each group

Given our prior TSF dyslexia data, we expected consistent phase entrainment in each group in

the delta and theta bands. To assess the phase entrainment consistency in each group and in

each frequency band, we applied the Rayleigh test of uniformity to the child preferred phases

separately for each group and for each frequency band (see Figure 2). For the control children,

we found consistency in all bands: delta band ($z = 5.14$, $p = 0.004$, see Figure 2A), theta band

($z = 3.70$, $p = 0.022$, see Figure 2B), and low gamma band ($z = 3.33$, $p = 0.033$, see Figure 2C).

For the children with DLD, we also found significantly consistent phase entrainment in the

delta band ($z = 8.31$, $p = 8.35 \times 10^{-5}$, see Figure 2D) and in the theta band ($z = 6.00$, $p = 0.0016$,

see Figure 2E). However, phase consistency in the low gamma band was absent ($z = 0.01$, $p =$

0.99, see Figure 2F). This result indicates that the DLD brain does not show significant

entrainment to rhythmic input in the gamma band.

Figure 2. Entrainment phase distribution across children in each group (rose plots) and group resultant vectors (red lines). Note the absence of phase consistency in the low gamma band for the children with DLD.

3.2. Comparing the strength of phase entrainment between groups

A priori, we had expected the strength of the consistency of phase entrainment to differ by

group in the delta band. The length of a *child resultant vector* can be considered as a criterion

of the strength of phase entrainment across different trials for that child. Indeed, longer *child*

resultant vectors indicate higher consistency of phase entrainment. To compare the strength of

phase consistency in control children with children with DLD, we applied Wilcoxon rank sum

tests based on the length of the *child resultant vectors* for the two frequency bands showing

significant entrainment, delta and theta. Contrary to prediction, there was no significant

difference between the two groups for either band (delta: $z = -0.70$, $p = 0.49$, Figure 3A; theta: z

$= -0.55$, $p = 0.58$, Figure 3B). We also checked these comparisons after removing outliers. A

data point was considered as an outlier if the corresponding power value was greater than a 1.5

interquartile range above the upper quartile or below the lower quartile of the population data

in that group. There were also no significant group differences after outlier removal.

386
387   **Figure 3. The length of child resultant vectors in the delta (A) and theta (B) bands in the**
388   **two groups.** The grey circles denote the length of the child resultant vectors for individual
389   participants, the black disks denote the mean values for each group, and the horizontal line on
390   each box shows the median.
391

3.3. Comparing preferred phase and angular velocity between groups

A priori we expected group differences in both preferred phase in the delta band and in angular velocity in the delta band. To compare preferred phase between groups, the Watson-Williams test was applied to the delta and theta band data. The low gamma band was excluded, as it did not show consistent phase entrainment for the children with DLD. The Watson-Williams test assumes von Mises distributions, although this test is fairly robust against deviations from this assumption (Zar, 1999; Berens, 2009). In our data, the phase distribution for both delta and theta bands, as well as for both groups (TD and DLD), followed von Mises distributions (Watson's U2 test, $p > 0.05$). Contrary to prediction, there was no significant difference in terms of preferred phase between the groups, neither in the delta band (Watson-Williams test, $p = 0.84$) nor in the theta band (Watson-Williams test, $p = 0.46$).

We also explored the behaviour of preferred phase across time separately for each group and each frequency band by computing angular velocities. Here the gamma band data could be included, as angular velocity (see Section 2.4) analyses do not require phase consistency. To this end, we calculated the group preferred phase for each frequency band prior to and after the occurrence of a stimulus (stimuli 4th–8th) in the entrainment period over the time interval of –500 ms to 500 ms. This time interval corresponds to two repetitions of the syllable "ba". Figure 4 shows the group preferred phase for the two groups and for delta, theta, and low gamma bands. For both groups, delta-band and theta-band group preferred phases appear to follow a quasiperiodic pattern, with a frequency of around 2 Hz for the delta band (angular velocity of 4 rad/s, see Figure 4A) and a frequency of around 4 Hz for the theta band (angular velocity of 8 rad/s, see Figure 4B). For the low gamma band, however, the pre-stimulus group preferred phase for children with DLD appears to be different from that of the control children. We therefore employed a two-sample t-test to find the time bins with a statistical difference in terms of angular velocity. The result suggested a significant difference between the two groups

over the time interval of –440 ms to –240 ms (two-sample t-test, p = 0.017, see Figure 4C).

This suggests that phase changes differently across time for the children with DLD, in the

gamma band only.

Figure 4. The group preferred phase (radians) as a function of the delay (ms) relative to
the onset of "ba" stimuli. The blue and red curves are related to control (N = 16) and DLD
group (N = 16), respectively. Both curves appear to follow a quasiperiodic pattern with a
frequency of around 2 Hz in the delta band (A) and a quasiperiodic pattern with a frequency
rate of around 4 Hz in the theta band (B). The curves related to the low gamma band (C) reveal
a pre-stimulus difference between the two groups. In particular, the pre-stimulus angular
velocity in this band over the time interval of –440 ms to –240 ms (marked with green colour)
differs significantly by group.

3.4. Comparing ERPs between groups

Figure 5 depicts the ERP curves for each group. We calculated the ERP in the time interval of

–500 ms to 500 ms separately for each group and each frequency band, as described in Section

2.6. Although differences between group could be expected in terms of pre- and post-stimulus

peaks in ERPs from visual inspection of Figure 5, no significant group difference was found in

any frequency band (delta, theta, low gamma).

Figure 5. Event related potentials for delta (A), theta (B) and low gamma (C) bands. The blue and red curves denote the ERPs for control and DLD groups, respectively. The shaded areas denote the standard error of mean for each group.

3.5. Comparison of band-power between groups

To compare the band-power between groups, Wilcoxon rank sum tests were applied for each frequency band after removing outliers. We calculated the average power over the entrainment period in each frequency band separately for each group, using steps described in Section 2.7. Figure 6 shows the average power versus frequency separately for each group and for delta (see Figure 6A), theta (see Figure 6B), and low gamma (see Figure 6C) bands. A significant difference between control children and children with DLD was found in theta ($z = -2.01$, $p = 0.04$; Figure 6B; outliers [control] N = 3, outliers [DLD] N = 2;) and in low gamma ($z = -2.64$, $p = 0.008$; Figure 6C; outliers [control] N = 2, outliers [DLD] N = 3) bands. However, there

451 was no difference in band-power for the delta band ($z = -1.45$, $p = 0.15$; Figure 6A; outliers

452 [control] N = 2, outliers [DLD] N = 2).

454
455 **Figure 6. Delta-band (A), theta-band (B), and low gamma-band (C) power in the**
456 **entrainment period.** The shaded areas denote the standard error of the mean for each group.
457 Please note the numerical power differences in the y axis by band.
458

3.6. Comparing cross-frequency PAC between groups

A priori, we had tentatively predicted a group difference in theta-gamma PAC between groups.

Delta-theta PAC, delta-low gamma PAC and theta-low gamma PAC were computed over the

entrainment period separately for each group by computing the respective modulation indices.

We then applied Wilcoxon rank sum tests to compare these modulation indices between the

two groups. No significant differences were found for delta-theta PAC ($z = -1.34$, $p = 0.18$,

Figure 7A) or delta-low gamma PAC ($z = -1.07$; $p = 0.28$, Figure 7B), nor for theta-low gamma

PAC (Wilcoxon rank sum test, $z = -0.85$; $p = 0.40$, Figure 7C). Similar non-significant results

were found following outlier removal. Contrary to prediction, therefore, theta-gamma PAC was

not different in the children with DLD for the rhythmic syllable repetition task.

Figure 7. Modulation index was used as a measure for delta-theta (A), delta-low gamma (B), and theta-low gamma (C) phase-amplitude coupling. The error bars denote the standard error of mean.

3.7. Comparing cross-frequency PPC between groups

Phase-phase coupling was also compared between groups. Delta-theta PPC, delta-low gamma PPC and theta-low gamma PPC were computed over the entrainment period separately for each group, as described in Section 2.9. We then applied two-sample t-tests to compare the two groups. As for PAC, no significant group differences were found, delta-theta PPC ($p = 0.89$, Figure 8A), delta-low gamma PPC ($p = 0.26$, Figure 8B), theta-low gamma PPC ($p = 0.11$, Figure 8C). Group PPC remained statistically equivalent after outlier removal.

Figure 8. Delta-band (A), theta-band (B), and low gamma-band (C) phase-phase coupling. The blue and red curves denote the PPCs for control and DLD groups, respectively.

Finally, we calculated the phase difference (see Equation 8) for delta-theta, delta-low gamma, and theta-low gamma oscillations for each child at the onsets of stimuli (4th–8th) during the entrainment period. The resultant phase difference for each group was then computed by averaging across the phase differences obtained for children in that group, and separately for delta-theta, delta-low gamma, and theta-low gamma oscillations (see Figure 9). We applied Watson-Williams tests to compare the group resultant phases between the groups. All measures for both groups (TD and DLD) followed von Mises distributions (Watson's U2 test, $p > 0.05$). In this case, low-frequency resultant phase (delta and theta) did show significant differences between the two groups when coupling with the higher-frequency gamma band, delta-low gamma ($p = 0.04$), theta-low gamma ($p = 3.28 \times 10^{-4}$). Accordingly, the synchronisation of low-frequency phases (delta, theta) with high-frequency phase (low gamma) appears to operate differently for children with DLD. These differences are clearly visible in Figure 9. However, regarding delta-theta resultant phase differences, the two groups were not significantly different ($p = 0.11$).

Figure 9. Delta-theta, delta-low gamma, and theta-low gamma resultant phase differences at the onsets of stimuli (4th–8th) during the entrainment period for control (A, B, C) and DLD (D, E, F) children. The blue and red vectors denote the resultant phase difference for

505 individual children in control and DLD groups, respectively. The black vectors show the group
506 resultant phase difference.
507

4. Discussion

The current study investigated the ARR hypothesis of developmental disorders of language at

the neural level, by examining different mechanistic aspects of the oscillatory encoding of a

highly constrained rhythmic speech input, the repetition of the syllable "ba" at a 2 Hz rate.

Despite the similarities in behavioural and sensory rhythm processing tasks that have been

documented for children with DLD and children with dyslexia (Richardson et al., 2004;

Corriveau et al., 2007; Thomson & Goswami, 2008; Corriveau & Goswami, 2009; Fraser et

al., 2010; Goswami et al., 2011; Beattie & Manis, 2012; Goswami et al., 2013; Cumming et

al., 2015a,b; Richards & Goswami, 2015, 2019), the neural mechanisms that were atypical for

the children with DLD in this paradigm were quite different from those previously documented

for children with dyslexia (Power et al., 2013; Keshavarzi et al., 2022). In particular, the

expectation of impaired delta band responses for children with DLD was not supported. The

children with DLD showed equivalent phase entrainment strength to TD children in the delta

band, they showed equivalent preferred delta phase, and they showed equivalent delta band

angular velocity. ERPs in the delta band did not differ by group, and nor did delta band power

in the entrainment period. Similar results were found for the theta band regarding these

parameters, with the exception that theta band power during the entrainment period was

significantly elevated in DLD children (Figure 6). These data suggest that the delta and theta

band mechanisms that are thought to underpin rhythmic speech encoding are operating in an

essentially similar way in the DLD brain when compared to the brains of age-matched TD

children, at least in the rhythmic syllable repetition task paradigm.

By contrast, low gamma band oscillatory responses in the rhythmic syllable repetition

task showed some differences in the DLD brain. Low gamma band phase entrainment in the

children with DLD was absent in the current paradigm, suggesting that phase consistency in

this band was not attained despite the highly predictable input. There was also a faster angular

velocity in the gamma band for the DLD children, suggesting that gamma phase changed

differently over time when compared to TD children. Furthermore, when low-frequency phases

had to synchronise with high-frequency gamma responses in our task, there were group

differences in resultant phase compared to the TD controls, for both delta-low gamma and

theta-low gamma coupling (Figure 9). Although the strength of phase-amplitude and phase-

phase coupling in the two groups of children was not significantly different (Figure 8), the

resultant phase differences suggest that this similar synchronisation strength was generated by

a different underlying relationship between low-frequency phase and high-frequency phase in

the DLD brain. One way of interpreting the consequences for speech processing is that the

interactions or connectivity between neural networks at these different rates is impaired,

namely the theoretical notion of *communication through coherence* (Fries, 2015). Fries (2009,

2015) proposed that neuronal inputs that arrive at random phases of the excitability cycle of

related rhythmic neuronal responses will have lower effective connectivity, with consequently

negative effects for cognition. Fries used examples from gamma band activity during visual

cognition (primarily in animals) to illustrate communication through coherence, but the

fundamental principle that he was proposing is relevant here, which is that the synchronization

of brain oscillations is a key mechanism for information encoding and perceptual binding. In

our data, the gamma rhythmic response to the highly predictable speech input does not attain

phase consistency, making it more difficult for low-frequency responses (delta, theta) to

synchronise their phases to gamma phase. This could suggest reduced coherence during speech

processing, negatively affecting the precision of linguistic encoding. In communication

through coherence theory, gamma-band coherence enables communication to be not only

effective but also precise. Although some aspects of our data suggest that the low-frequency

responses (delta and theta) and their phase relations are operating in the same way as for TD

children during the rhythmic speech task (with equivalent PAC and PPC strength), an optimal

phase relation with gamma band responses in the temporal domain cannot be established,

thereby impairing speech processing. This is mechanistically quite different from our previous

findings with children with developmental dyslexia in this paradigm, where delta phase was

atypical (Power et al., 2013; Keshavarzi et al., 2022). Nevertheless, for dyslexic children also,

PAC mechanisms were explored and appear to be intact (delta-beta PAC, see Power et al., 2016;

Keshavarzi et al., 2024).

A related view of the importance of gamma band processing for speech encoding

concerns the parsing of phonemic information in the speech signal (Giraud & Poeppel, 2012;

Poeppel, 2014). For example, it is known that speech contains crucial information required for

successful linguistic decoding of the signal that is carried at multiple temporal scales, such as

intonation-level or prosodic information at the scale of 500–2000 ms (argued to be encoded by

delta band responses), syllabic information that is correlated to the acoustic envelope of speech

at the scale of 150–300 ms (encoded by theta band responses), and rapidly-changing featural

information at the scale of 20–80 ms (encoded by gamma band responses). One mechanism for

encoding is suggested to be the sliding and resetting of neural temporal windows that match

information in the signal via oscillatory responses, implemented as phase locking to key

features of the input via phase-resetting of the intrinsic oscillations. While more recent work

has shown that acoustic feature information is also encoded by the slower delta and theta band

responses (for adults, children and infants, see Di Liberto et al., 2015, 2018, 2023), the idea

that gamma band responses encode phoneme-relevant information has been supported by

transcranial alternating current stimulation (tACS) studies using gamma band stimulation. For

example, Marchesotti et al. (2020) reported that 20 minutes of tACS stimulation at ~30Hz

significantly improved phonemic awareness in a behavioural task for French adults with

dyslexia. The data presented here may suggest that the ~30 Hz response to acoustic feature

information in speech is atypical in the DLD brain, which could impair some aspects of

phoneme awareness. However, given that most phonetic features are present in natural speech

over much longer time windows (see Menn et al., 2023), and given that the cortex maintains

representations of phonetic information even after it is lost from sensory input (Gwilliams et

al., 2022), the idea that the gamma band differences reported here prevent children with DLD

from representing phonemic information seems unlikely. Accordingly, we prefer to interpret

the data using communication through coherence theory. Our findings appear to indicate

reduced co-ordination of low-frequency oscillatory phase with gamma band responses in the

DLD brain, reflecting altered neuronal network interactions during linguistic processing that

may affect encoding precision and thereby impair speech perception at all phonological levels,

not only the phoneme level. Converging studies with natural speech rather than the rhythmic

speech paradigm utilised here are required to assess these different possibilities.

The gamma band differences reported here are also interesting with respect to an

alternative auditory theory of DLD, the rapid auditory processing (RAP) theory (Tallal, 1980,

2004). RAP theory proposed that basic auditory processing of brief, rapidly successive acoustic

changes is impaired in children with DLD, drawing on data from a nonspeech task using brief

(75 ms) pairs of tones that differed in pitch and succeeded each other rapidly in time (e.g., gaps

of 8, 15, 30, 60 ms, see Tallal & Piercy, 1973). The children's task was to decide whether a

high tone preceded a low tone in a pair or vice versa. Children with DLD were worse than TD

controls in the tone task when the gap between stimuli was short, but were not different to TD

controls when the gaps were longer (greater than 150 ms). Regarding speech processing, RAP

difficulties were expected to affect phonetic discrimination (thought to require processing at

rapid timescales around 40 ms, for example discriminating /b/ from /d/) via impaired

discrimination of formant transitions (rapid acoustic changes in frequency and intensity).

Further, for children with DLD, intervention studies based on RAP can lead to improved

language skills (e.g., Stevens et al., 2008, although see also Gillam et al., 2008).

608   Regarding neural data with DLD children, a nonspeech MEG study by Heim and

colleagues indicated altered temporal organization of gamma-band oscillatory activity when

children with DLD processed rapid tone sequences (Heim et al., 2011). Children listened to

pairs of tones separated by 70 ms, and judged the pitch of the second tone. Heim et al. reported

significantly reduced amplitude and phase-locking of early (45–75 ms) oscillations in the

gamma-band range (29–52 Hz) for the second stimulus. ERP studies of infants at family risk

for DLD using brief rapidly-presented tones have reported differences in auditory evoked

potentials indexing change detection compared to TD infants, and in resting state gamma

power, which in some studies predicted later language skills (English, language measured at 3,

4 and 5 years, Benasich et al., 2008; Choudhury & Benasich, 2011; Gou et al., 2011; Italian,

language measured at 20 months, Cantiani et al., 2016, 2019). Of particular interest regarding

the current dataset, an investigation of the same English-exposed infants studied by Choudhury

and Benasich (2011) at 6 and 12 months showed that the infant change detection response for

rapidly-presented tone pairs was mainly determined by low frequency power and phase-locking

responses (2-3 Hz, Hamalainen et al., 2019). Regarding the ARR hypothesis, it is logically

possible that while children with both DLD and dyslexia share a difficulty in processing AE

rise times and show difficulties in behavioural speech rhythm tasks, the rates of temporal

integration that are impaired neurally may differ for each developmental disorder. These prior

infant studies and the current gamma band data are consistent with Tallal's (2004) claim that

auditory processing in temporal integration windows of around 40ms (25 Hz) is adversely

affected in children with DLD. However, low frequency phase-phase mechanisms may

contribute to this impaired processing. Further RAP studies based on processing natural speech

rather than pairs of tones are required to investigate this further, and to explore potential links

to phoneme representation.

The current study has a number of limitations. Firstly, the rhythmic speech task is not

a natural language task. Further investigation of the ARR hypothesis using natural speech

listening tasks is required in order to establish whether the current gamma band and resultant

phase differences are present during natural language processing. Second, the sample size is

small, with 16 DLD children. However, it is representative of the existing neural DLD literature

(e.g., Heim et al. 2011 studied 17 children with DLD, Choudhury & Benasich 2011 studied 17

infants at risk for DLD). This reflects the difficulty of engaging families regarding

neuroimaging.

In conclusion, here we find that the 9-year-old DLD brain does not show the delta-band

deficits predicted by TS theory to underpin impaired rhythmic speech processing. Instead, the

children with DLD exhibited an absence of phase consistency in the gamma band when

processing highly predictable rhythmic speech, and showed different angular velocity in the

gamma band compared to TD children. The children with DLD also showed significantly

greater gamma and theta power during the task, and a difference in resultant phase when low-

frequency phases (delta, theta) coupled with low gamma. These latter data are consistent with

some prior neural nonspeech investigations of children with DLD. Overall, the data suggest

reduced coherence between slower and faster networks during speech processing, which would

negatively affect the precision of linguistic encoding. Further studies employing natural

continuous speech are required to investigate this interpretation in more depth.

Data and Code Availability

Data and code will be made available on request.

Author contributions

M.K. conceptualisation, methodology, data collection, data analyses, visualisation, writing – original draft; S.R. data collection, writing – review & editing; G.F. data collection, writing – review & editing; L.P. data collection, writing – review & editing; U.G. funding acquisition, project administration, supervision, conceptualisation, methodology, writing – original draft.

Conflict of interest

The authors declare no conflicts of interest.

Acknowledgements

The authors would like to thank all the children, families and schools involved in the study. This research was funded by a donation to UG from the Yidan Prize Foundation. The sponsor played no role in the study design, data interpretation nor writing of the report.

References

Asaridou, S. S., & Watkins, K. E. (2022). Neural basis of speech and language impairments in development: the case of developmental language disorder. (In K. Cohen Kadosh, Ed.; *The Oxford Handbook of Developmental Cognitive Neuroscience*, pp. C19.S1–C19.S9). Oxford University Press.

Attaheri, A., Panayiotou, D., Phillips, A., Ní Choisdealbha, A., Di Liberto, G., Rocha, S., Brusini, P., Mead, N., Flanagan, S., Olawole-Scott, H., & Goswami, U. (2022). Cortical tracking of sung speech in adults vs infants: a developmental analysis. *Frontiers in Neuroscience*, 16, 842447. https://doi.org/10.3389/fnins.2022.842447

Attaheri, A., Ní Choisdealbha, A., Di Liberto, G., Rocha, S., Brusini, P., Mead, N., Olawole-Scott, H., Boutris, P., Gibbon, S., Williams, I., Grey, C., Flanagan, S., M., & Goswami, U. (2022). Delta- and theta-band cortical tracking and phase-amplitude coupling to sung speech by infants. *Neuroimage*, 247, 118698. https://doi.org/10.1016/j.neuroimage.2021.118698

Attaheri, A., Ní Choisdealbha, A., Rocha, S., Brusini, P., Di Liberto, G., Mead, N., Olawole-Scott, H., Boutris, P., Gibbon, S., Williams, I., Grey, Alfaro e Oliveira, M., Brough, C., Flanagan, S., M., & Goswami, U. (preprint). Infant low-frequency EEG cortical power, cortical tracking and phase-amplitude coupling predicts language a year later. bioRxiv 2022.11.02.514963. https://doi.org/10.1101/2022.11.02.514963

Badcock, N. A., Bishop, D. V. M., Hardiman, M. J., Barry, J. G., & Watkins, K. E. (2012). Co-localisation of abnormal brain structure and function in specific language impairment. *Brain and Language, 120*, 3, 310-320. https://doi.org/10.1016/j.bandl.2011.10.006

Beattie, R., & Manis, F. (2012). Rise time perception in children with reading and combined reading and language difficulties. *Journal of Learning Disabilities*, *46*(3), 200-209. https://doi.org/10.1177/0022219412449421

Bedoin, N., Brisseau, L., Molinier, P., Roch, D., & Tillmann, B. (2016). Temporally Regular Musical Primes Facilitate Subsequent Syntax Processing in Children with Specific Language Impairment. *Frontiers in Neuroscience*, 10, 245. https://doi.org/10.3389/fnins.2016.00245

Benasich, A.A., Gou, Z., Choudhury, N., & Harris, K.D. (2008). Early cognitive and language skills are linked to resting frontal gamma power across the first three years. *Behavioural Brain Research*, 195, 215-222. https://doi.1016/j.bbr.2008.08.049

Bishop, D. V. M. (2013). Uncommon Understanding: Development and Disorders of Language Comprehension in Children. Psychology Press, 288, eBook ISBN9781315804699. https://doi.org/10.4324/9781315804699

Bishop, D., Snowling, M., Thompson, P., & Greenhalgh, T. (2017). Phase 2 of CATALISE: a multinational and multidisciplinary Delphi consensus study of problems with language development: Terminology. *Journal of Child Psychology and Psychiatry*, 58(10), 1068-1080. https://doi.org/10.1111/jcpp.12721

Blomert, L., & Mitterer, H. (2004). The fragile nature of the speech-perception deficit in dyslexia: Natural vs. synthetic speech. *Brain and Language*, 89(1), 21-26. https://doi.org/10.1016/S0093-934X(03)00305-5

Boemio, A., Fromm, S., Braun, A., & Poeppel, D. (2005). Hierarchical and asymmetric temporal sensitivity in human auditory cortices. *Nature Neuroscience*, 8, 389-395. https://doi.org/10.1038/nn1409

Cantiani, C., Riva, V., Piazza, C., Bettoni, R., Molteni, M., Choudhury, N., Marino, C., & Benasich, A. A., (2016). Auditory discrimination predicts linguistic outcome in Italian infants with and without familial risk for language learning impairment. *Developmental Cognitive Neuroscience, 20*, 23-34. https://doi.org/10.1016/j.dcn.2016.03.002

Cantiani, C., Ortiz-Mantilla, S., Riva, V., Piazza, C., Bettoni, R., Musaccia, G., Molteni, M., Marino, C., & Benasich, A. A., (2019). Reduced left-lateralized pattern of event-related EEG oscillations in infants at familial risk for language and learning impairment. *Neuroimage: Clinical, 22*, 101778. https://doi.org/10.1016/j.nicl.2019.101778

Choudhury, N. & Benasich, A. A. (2011). Maturation of auditory evoked potentials from 6 to 48 months: Prediction to 3 – and 4 – year language and cognitive abilities. *Clinical Neurophysiology*, 122, 320-338. https://doi.org/10.1016/j.clinph.2010.05.035

Corriveau, K. H., & Goswami, U. (2009). Rhythmic motor entrainment in children with speech and language impairments: Tapping to the beat. *Cortex*, 45, 119-130. https://doi.org/10.1016/j.cortex.2007.09.008

Corriveau, K., Pasquini, E., & Goswami, U. (2007). Basic auditory processing skills and specific language impairment: A new look at an old hypothesis. *Journal of Speech*, *Language*, *and Hearing Research*, 50, 647-666. https://doi.org/10.1044/1092-4388(2007/046)

Cumming, R., Wilson, A., & Goswami, U. (2015). Basic auditory processing and sensitivity to prosodic structure in children with specific language impairments: A new look at a perceptual hypothesis. *Frontiers in Psychology*, 6, 972. https://doi.org/10.3389/fpsyg.2015.00972

Cumming, R., Wilson, A., Leong, V., Colling, L., & Goswami, U. (2015). Awareness of Rhythm Patterns in Speech and Music in Children with Specific Language Impairments. *Frontiers in Human Neuroscience*, 9. https://doi.org/10.3389/fnhum.2015.00672

Cutini, S., Szűcs, D., Mead, N., Huss, M., & Goswami, U. (2016). Atypical right hemisphere response to slow temporal modulations in children with developmental dyslexia. *NeuroImage*, 143, 40-49. https://doi.org/10.1016/j.neuroimage.2016.08.012

Dauer, R. M. (1983). Stress-timing and syllable-timing reanalyzed. *Journal of Phonetics*, 11(1), 51–62. https://doi.org/10.1016/S0095-4470(19)30776-4

Delorme, A., & Makeig, S. (2004). EEGLAB: an open-source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, 134(1), 9-21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Di Liberto, G. M., Attaheri, A., Cantisani, G., Reilly, R. B., Ní Choisdealbha, A., Rocha, S., Brusini, P., & Goswami, U. (2023). Emergence of the cortical encoding of phonetic features in the first year of life. Nature Communications, 14, 7789. https://doi.org/10.1038/s41467-023-43490-x

Di Liberto, G. M., Peter, V., Kalashnikova, M., Goswami, U., Burnham, D., & Lalor, E. C., (2018). Atypical cortical entrainment to speech in the right hemisphere underpins phonemic deficits in dyslexia. *NeuroImage*, 175, 70-79. https://doi.org/10.1016/j.neuroimage.2018.03.072

Di Liberto, G.M., O'Sullivan, J.A., & Lalor, E.C. (2015). Low-frequency cortical entrainment to speech reflects phoneme-level processing. *Current Biology*, 25, 2457-2465. https://doi.org/10.1016/j.cub.2015.08.030

Flanagan, S., & Goswami, U. (2018). The role of phase synchronisation between low frequency amplitude modulations in child phonology and morphology speech tasks. *Journal of the Acoustical Society of America*, 143(3), 1366. https://doi.org/10.1121/1.5026239

Fraser, J., Goswami, U., & Conti-Ramsden, G. (2010). Dyslexia and specific language impairment: The role of phonology and auditory processing. *Scientific Studies of Reading*, 14, 8-29. https://doi.org/10.1080/10888430903242068

Friedrich, M., & Friederici, D. (2005). Phonotactic Knowledge and Lexical-Semantic Processing in One-year-olds: Brain Responses to Words and Nonsense Words in Picture Contexts. *Journal of Cognitive Neuroscience*, 17(11), 1785-1802. https://doi.org/10.1162/089892905774589172

Friedrich, M., & Friederici, D. (2006). Early N400 development and later language acquisition. *Psychophysiology*, 43, 1, 1-12. https://doi.org/10.1111/j.1469-8986.2006.00381.x

Fries P. (2015). Rhythms for Cognition: Communication through Coherence. *Neuron*, 88(1), 220-35. https://doi.org/10.1016/j.neuron.2015.09.034

Fries, P. (2009). The Model- and the Data-Gamma. *Neuron*, 64(5), 601-612. https://doi.org/10.1016/j.neuron.2009.11.024

Ghitza, O., & Greenberg, S. (2009). On the possible role of brain rhythms in speech perception: Intelligibility of time-compressed speech with periodic and aperiodic insertions of silence. *Phonetica*, 66, 113-126. https://doi.org/10.1159/000208934

Gillam, R.B., Loeb, D.F., Hoffman, L.M., Bohman, T., Champlin, C.A., Thibodeau, L., Widen, J., Brandel, J., & Friel-Patti, S. (2008). The efficacy of Fast ForWord language intervention in school-age children with language impairment: A randomized controlled trial. *Journal of Speech, Language, and Hearing Research*, 51(1), 97-119. https://doi.org/10.1044/1092-4388(2008/007)

Giraud, A-L., & Poeppel, D. (2012). Cortical oscillations and speech processing: Emerging computational principles and operations. *Nature Neuroscience*, 15, 511-517. http://dx.doi.org/10.1038/nn.3063

Goswami, U. (2011). A temporal sampling framework for developmental dyslexia. *Trends in Cognitive Sciences*, 15(1), 3-10. https://doi.org/10.1016/j.tics.2010.10.001

Goswami, U. (2015). Sensory theories of developmental dyslexia: Three challenges for research. *Nature Reviews Neuroscience*, 16, 43-54. http://dx.doi.org/10.1038/nrn3836

Goswami, U. (2022). Language acquisition and speech rhythm patterns: an auditory neuroscience perspective. *Royal Society Open Science*, 9(7), 211855. https://doi.org/10.1098/rsos.211855

Goswami, U., & Leong, V. (2013). Speech rhythm and temporal structure: Converging perspectives?. *Laboratory Phonology*, 4(1), 67-92. https://doi.org/10.1515/lp-2013-0004

Goswami, U., Cumming, R., Chait, M., Huss, M., Mead, N., Wilson, A. M., Barnes, L., & Fosker, T. (2016). Perception of filtered speech by children with developmental dyslexia and children with specific language impairments. *Frontiers in Psychology*, 7, 791. https://doi.org/10.3389/fpsyg.2016.00791

Goswami, U., Mead, N., Fosker, T., Huss, M., Barnes, L., & Leong, V. (2013b). Impaired perception of syllable stress in children with dyslexia: A longitudinal study. *Journal of Memory and Language*, 69 (1), 1-17. https://doi.org/10.1016/j.jml.2013.03.001

Goswami, U., Wang, H-L. S., Cruz, A., Fosker, T., Mead, N., & Huss, M. (2011). Language-universal sensory deficits in developmental dyslexia: English, Spanish, and Chinese. *Journal of Cognitive Neuroscience*, 23, 325-337. https://doi.org/10.1162/jocn.2010.21453

Gou, Z., Choudhury, H., & Benasich, A.A. (2011). Resting frontal gamma power at 16, 24 and 36 months predicts individual differences in language and cognition at 4 and 5 years. *Behavioural Brain Research*, 220, 263-270. https://doi.org/j.bbr.2011.01.048

Gramfort, A., Luessi, M., Larson, E., Engemann, D.A., Strohmeier, D., Brodbeck, C., Goj, R., Jas, M., Brooks, T., Parkkonen, L., & Hämäläinen, M. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in neuroscience*, 267. https://doi.org/10.3389/fnins.2013.00267

Greenberg, S. (2006). A multi-tier framework for understanding spoken language. In S. Greenberg & W. Ainsworth (Eds.), *Listening to speech: An auditory perspective*. Lawrence Erlbaum Associates.

Gross, J., Hoogenboom, N., Thut, G., Schyns, P., Panzeri, S., Belin, P., & Garrod, S. (2013). Speech rhythms and multiplexed oscillatory sensory coding in the human brain. *PLOS Biology*, 11(12), e1001752. https://doi.org/10.1371/journal.pbio.1001752

Gwilliams, L., King, J. R., Marantz, A., & Poeppel, D. (2022). Neural dynamics of phoneme sequences reveal position-invariant code for content and order. *Nature Communications*, 13, 6606. https://doi.org/10.1038/s41467-022-34326-1

Hämäläinen, J. A., Ortiz-Mantilla, S. and Benasich, A., 2019. Change detection to tone pairs during the first year of life–Predictive longitudinal relationships for EEG-based source and time-frequency measures. *NeuroImage*, 198, 83-92. https://doi.org/10.1016/j.neuroimage.2019.05.034

Heim, S., Friedman, J.T., Keil, A., Benasich, A.A. (2011). Reduced Sensory Oscillatory Activity during Rapid Auditory Processing as a Correlate of Language-Learning Impairment. *Journal of Neurolinguistics*, 24(5), 539-555. https://doi.org/10.1016/j.jneuroling.2010.09.006

Helenius, P., Parviainen, T., Paetau, R., & Salmelin, R. (2009). Neural processing of spoken words in specific language impairment and dyslexia. *Brain*, 132, 7, 1918-1927. https://doi.org/10.1093/brain/awp134

Helenius, P., Sivonen, P., Parviainen, T., Isoaho, P., Hannus, S., Kauppila, T., Salmelin, R., & Isotalo, L. (2014). Abnormal functioning of the left temporal lobe in language-impaired children. *Brain and Language*, 130, 11-18. https://doi.org/10.1016/j.bandl.2014.01.005

Hülsemann, M., Naumann, E., & Rasch, B. (2019). Quantification of Phase-Amplitude Coupling in Neuronal Oscillations: Comparison of Phase-Locking Value, Mean Vector Length, Modulation Index, and Generalized-Linear-Modeling-Cross-Frequency-Coupling. *Frontiers in Neuroscience*, 13, 573. https://doi.org/10.3389/fnins.2019.00573

Keshavarzi, M., Mandke, K., Macfarlane, A., Parvez, L., Gabrielczyk F., Wilson, A., & Goswami, U. (2022). Atypical Delta-band Phase Consistency and Atypical Preferred Phase in Children with Dyslexia during Neural Entrainment to Rhythmic Audio-Visual Speech. *NeuroImage: Clinical*, 35, 103054. https://doi.org/10.1016/j.nicl.2022.103054

Keshavarzi, M., Mandke, K., Macfarlane, A., Parvez, L., Gabrielczyk F., Wilson, A., & Goswami, U. (2024). Atypical beta-band effects in children with dyslexia in response to rhythmic audio-visual speech. *Clinical Neurophysiology*, 1388-2457, https://doi.org/10.1016/j.clinph.2024.02.008

Keshavarzi, M., Ní Choisdealbha, A., Attaheri, A., Rocha, S., Brusini, P., Gibbon, S., Boutris, P., Mead, N., Olawole-Scott, H., Ahmed, H., Flanagan, S., Mandke, K., & Goswami. U. (2024). Decoding speech information from EEG data with 4-, 7- and 11-month-old infants: Using convolutional neural network, mutual information-based and backward linear models. *Journal of Neuroscience Methods*, 403, 110036. https://doi.org/10.1016/j.jneumeth.2023.110036

Kotz, S., & Schmidt-Kassow, M. (2015). Basal ganglia contribution to rule expectancy and temporal predictability in speech. *Cortex*, *68*, 48-60. https://doi.org/10.1016/j.cortex.2015.02.021

Ladányi, E., Persici, V., Fiveash, A., Tillmann, B., & Gordon, R. L. (2020). Is atypical rhythm a risk factor for developmental speech and language disorders?. *Wiley Interdisciplinary Reviews: Cognitive Science*, 11(5), e1528. https://doi.org/10.1002/wcs.1528

Lallier, M., Molinaro, N., Lizarazu, M., Bourguignon, M., & Carreiras, M. (2017). Amodal Atypical Neural Oscillatory Activity in Dyslexia: A Cross-Linguistic Perspective. *Clinical Psychological Science*, 5(2), 379-401. https://doi.org/10.1177/2167702616670119

Lee, J. C., Nopoulos, P. C., & Tomblin, J. B. (2013). Abnormal subcortical components of the corticostriatal system in young adults with DLI: a combined structural MRI and DTI study. *Neuropsychologia*, 51(11), 2154-2161. https://doi.org/10.1016/j.neuropsychologia.2013.07.011

Leonard, L.B. (2014). Children with specific language impairment. *Cambridge, MA: MIT Press.* https://doi.org/10.7551/mitpress/9152.001.0001.

Leong, V., & Goswami, U. (2015). Acoustic-emergent phonology in the amplitude envelope of child-directed speech. *PLOS ONE*, 10(12), e0144411. https://doi.org/10.1371/journal.pone.0144411

Manis, F. R., McBride-Chang, C., Seidenberg, M. S., Keating, P., Doi, L. M., Munson, B., & Petersen, A. (1997). Are speech perception deficits associated with developmental dyslexia?. *Journal of experimental child psychology*, 66(2), 211-235. https://doi.org/10.1006/jecp.1997.2383

Marchesotti, S., Nicolle, J., Merlet, I., Arnal, L. H., Donoghue, J. P., & Giraud, A-L. (2020). Selective enhancement of low gamma activity by tACS improves phonemic processing and reading accuracy in dyslexia. *PLoS Biology*, 18(9), e3000833. https://doi.org/10.1371/journal.pbio.3000833

Mehta, B., Chawla, V. K., Parakh, M., Parakh, P., Bhandari, B., & Gurjar, A. S. (2015). EEG Abnormalities in Children with Speech and Language Impairment. *Journal of Clinical and Diagnostic Research*, 9(7), CC04. https://doi.org/10.7860/JCDR/2015/13920.6168

Menn, K. H., Männel, C., & Meyer, L. (2023). Phonological acquisition depends on the timing of speech sounds: Deconvolution EEG modeling across the first five years. *Science Advances*, 9(44), eadh2560. https://doi.org/10.1126/sciadv.adh2560

Ní Choisdealbha, A., Attaheri, A., Rocha, S., Brusini, P., Flanagan, S., M., Mead, N., Gibbon, S., Olawole-Scott, H., Williams, I., Grey, C., Boutris, P., Ahmed, H., & Goswami, U. (2022). Neural detection of changes in amplitude rise time in infancy. *Developmental Cognitive Neuroscience*, 54, 101075. https://doi.org/10.1016/j.dcn.2022.101075

Ní Choisdealbha, A., Attaheri, A., Rocha, S., Mead, N., Olawole-Scott, H., Brusini, P., Gibbon, S., Boutris, P., Grey, C., Hines, D., Williams, I., Flanagan, S., & Goswami, U. (2023). Neural phase angle from two months when tracking speech and non-speech rhythm linked to language performance from 12 to 24 months. *Brain and Language*, 243, 105301. https://doi.org/10.1016/j.bandl.2023.105301

Ortiz-Mantilla, S., & Benasich, A. (2013). Neonatal electrophysiological predictors of cognitive and language development. *Developmental Medicine & Child Neurology*, 55(9), 781-782. https://doi.org/10.1111/dmcn.12207

Parker, A. J., Woodhead, Z. V. J., Carey, D. P., Groen, M. A., Gutierrez-Sigut, E., Hodgson, Hudson, J., Karlsson, E. M., MacSweeney, M., Payne, H., Simpson, N., Thompson, P. A., Watkins, K. E., Egan, C., Grant, J. H., Harte, S., Hudson, B. T., Sablik, M., Badcock, N. A., & Bishop, D. V. M. (2022). Inconsistent language lateralisation Testing the dissociable language laterality hypothesis using behaviour and lateralised cerebral blood flow. *Cortex*, 154, 105-134. https://doi.org/10.1016/j.cortex.2022.05.013

Parvez, L., Keshavarzi, M., Richards, S., Di Liberto, G. M., & Goswami, U. (2024). Atypical speech production of multisyllabic items by children with developmental language disorder (DLD) indicate prosodic difficulties. *PsyArXiv*. https://doi.org/10.31234/osf.io/52r6h

Peter, V., van Ommen, S., Kalashnikova, M., Mazuka, R., Nazzi, T., & Burnham, D. (2022). Language specificity in cortical tracking of speech rhythm at the mora, syllable, and foot levels. *Scientific Reports*, 12, 13477. https://doi.org/10.1038/s41598-022-17401-x

Poeppel, D. (2014). The neuroanatomic and neurophysiological infrastructure for speech and language. *Current Opinion in Neurobiology*, 28, 142-149. https://doi.org/10.1016/j.conb.2014.07.005

Power, A. J., Colling, L. C., Mead, N., Barnes, L., & Goswami, U. (2016). Neural encoding of the speech envelope by children with developmental dyslexia. *Brain & Language*, 160, 1-10. https://doi.org/10. 1016/j.bandl.2016.06.006

Power, A. J., Mead, N., Barnes, L., & Goswami, U. (2012). Neural entrainment to rhythmically presented auditory, visual, and audio-visual speech in children. *Frontiers in Psychology*, 3, 216. https://doi.org/10.3389/fpsyg.2012.00216

Power, A. J., Mead, N., Barnes, L., & Goswami, U. (2013). Neural entrainment to rhythmic speech in children with developmental dyslexia. *Frontiers in Human Neuroscience*, 7(777), 1-19. https://doi.org/10.3389/fnhum.2013.00777

Przybylski, L., Bedoin, N., Krifi-Papoz, S., Herbillon, V., Roch, D., & Léculier, L., Kotz, S. A., & Tillmann, B. (2013). Rhythmic auditory stimulation influences syntactic processing in children with developmental language disorders. *Neuropsychology*, 27(1), 121-131. https://doi.org/10.1037/a0031277

Richards, S., & Goswami, U. (2015). Auditory processing in specific language impairment (SLI): Relations with the perception of lexical and phrasal stress. *Journal of Speech, Language, and Hearing Research*, 58(4), 1292-1305. https://doi.org/10.1044/2015_jslhr-l-13-0306

Richards, S., & Goswami, U. (2019). Impaired recognition of metrical and syntactic boundaries in children with developmental language disorders. *Brain Sciences*, 9(2), 33. https://doi.org/10.3390/brainsci9020033

Richardson, U., Thomson, J. M., Scott, S. K., & Goswami, U. (2004). Auditory processing skills and phonological representation in dyslexic children. *Dyslexia*, 10, 215-233. https://doi.org/10.1002/dys.276

Ríos-López, P., Molinaro, N., Bourguignon, M., & Lallier, M. (2020). Development of neural oscillatory activity in response to speech in children from 4 to 6 years old. *Developmental Science*, 23(6), e12947. https://doi.org/10.1111/desc.12947

Rios-Lopez, P., Molinaro, N., Bourguignon, M., & Lallier, M. (2022). Right-hemisphere coherence to speech at pre-reading stages predicts reading performance one year later. *Journal of Cognitive Psychology*, 34, 2, 179-193. https://doi.org/10.1080/20445911.2021.1986514

Rocha, S., Ní Choisdealbha, A., Attaheri, A., Mead, N., Olawole-Scott, H., Grey, C., Williams, I., Gibbon, S., Boutris, P., Brusini, P., & Goswami, U. (in press). Language Acquisition in the Longitudinal Cambridge UK BabyRhythm Cohort. *Collabra: Psychology*, 10(1). https://doi.org/10.1525/collabra.92998

Sallat, S., & Jentschke, S. (2015). Music perception influences language acquisition: Melodic and rhythmic-melodic perception in children with specific language impairment. *Behavioural Neurology*, 2015, 606470. https://doi.org/10.1155/2015/606470

Serniclaes, W., van Heghe, S., Mousty, P., Carré, R., & Sprenger-Charolles, L. (2004). Allophonic mode of speech perception in dyslexia. *Journal of Experimental Child Psychology*, 87, 336-361. https://doi.org/10.1016/j.jecp.2004.02.001

Shafer, V. L., Schwartz, R. G., Morr, M. L., Kessler, K. L., Kurtzberg, D., & Ruben, R. J. (2001) Neurophysiological Indices of Language Impairment in Children. *Acta Oto-Laryngologica*, 121(2), 297-300. https://doi.org/10.1080/000164801300043929

Snowling, M. J. (2000). *Dyslexia* (2nd ed.). Oxford: Blackwell Publishers.

Stanovich, K. (1988). Explaining the differences between the dyslexic and the garden-variety poor reader: The Phonological-Core Variable-Difference Model. *Journal of Learning Disabilities*, 21, 590-604. https://doi.org/10.1177/002221948802101003

Stevens, C., Fanning, J., Coch, D., Sanders, L., & Neville, H. (2008). Neural mechanisms of selective auditory attention are enhanced by computerized training: electrophysiological evidence from language-impaired and typically developing children. *Brain research*, 1205, 55-69. https://doi.org/10.1016/j.brainres.2007.10.108

Swan, D., & Goswami, U. (1997). Phonological awareness deficits in developmental dyslexia and the phonological representations hypothesis. Journal of Experimental Child Psychology, 66(1), 18-41. https://doi.org/10.1006/jecp.1997.2375

Systad, S., Bjørnvold, M., Sørensen, C., & Halaas Lyster, S. S. (2019). The value of electroencephalogram in assessing children with speech and language impairments. *Journal of Speech, Language, and Hearing Research*, 62(1), 153-168. https://doi.org/10.1044/2018_JSLHR-L-17-0087

Tallal, P. (1980). Auditory temporal perception, phonics, and reading disabilities in children. *Brain and Language*, 9, 182-198. https://doi.org/10.1016/0093-934X(80)90139-X

Tallal, P. (2004). Improving language and literacy is a matter of time. *Nature Reviews Neuroscience*, 5, 721-728. https://doi.org/10.1038/nrn1499

Tallal, P. (2004). Improving language and literacy is a matter of time. Nature Reviews Neuroscience, 5, 721-728. https://doi.org/10.1038/nrn1499

Tallal, P., & Piercy, M. (1973). Defects of non-verbal auditory perception in children with developmental aphasia. *Nature*, 241, 468-469. https://doi.org/10.1038/241468a0

Thomson, J. M., & Goswami, U. (2008). Rhythmic processing in children with developmental dyslexia: Auditory and motor rhythms link to reading and spelling. *Journal of Physiology-Paris*, 102, 120-129. https://doi.org/10.1016/j.jphysparis.2008.03.007

Tort, A. B. L., Kramer, M. A., Thorn, C., & Kopell, N. J. (2008). Dynamic cross-frequency couplings of local field potential oscillations in rat striatum and hippocampus during performance of a T-maze task. *Proceedings of the National Academy of Sciences*, 105(51) 20517-20522. https://doi.org/10.1073/pnas.081052410

Watkins, K. E., Vargha-Khadem, F., Ashburner, J., Passingham, R. E., Connelly, A., Friston, K. J., Frackowiak, R. S. J., Mishkin, M., & Gadian, D. G. (2002). MRI analysis of an inherited speech and language disorder: structural brain abnormalities. *Brain*, 125(3), 465-478. https://doi.org/10.1093/brain/awf057

Wechsler, D. (2016). Wechsler Intelligence Scale for Children, 4th Edn. London, UK: Pearson Assessment.

982   Wiig, E. H., Semel, E., Secord, W. A. (2017). Clinical Evaluation of Language Fundamentals
983   - Fifth Edition. Pearson.