1

Proteome dynamics reveal Leiomodin 1 as a key regulator of myogenic differentiation

Ellen Späth[1*], Svenja C. Schüler[1,2*], Ivonne Heinze[1], Therese Dau[1], Alberto Minetti[1], Maleen

Hofmann[1], Julia von Maltzahn[1,3#] and Alessandro Ori[1#]

Affiliations

1 - Leibniz Institute on Aging - Fritz Lipmann Institute (FLI), Jena, Germany

2 - Département de Pharmacologie - Physiologie, Université de Sherbrooke, QC, Canada

3 - Brandenburg Technische Universität Cottbus-Senftenberg, Universitätsplatz 1, 01968 Senftenberg,

Germany, Faculty of Health Sciences and Faculty for environment and natural sciences

[*] Contributed equally

[#] Correspondence should be addressed to: alessandro.ori@leibniz-fli.de or julia.vonmaltzahn@b-tu.de

Abstract

During myogenic differentiation the cellular architecture and proteome of muscle stem cells

and myoblasts undergo extensive remodeling. These molecular processes are only partially

understood and display alterations in disease conditions as well as during aging resulting in

impaired regeneration. Here, we used mass spectrometry to quantify the temporal dynamics

of more than 6000 proteins during myogenic differentiation. We identified the actin nucleator

leiomodin 1 (LMOD1) among a restricted subset of cytoskeletal proteins increasing in

abundance in early phases of myogenic differentiation. We show that LMOD1 is already

expressed by muscle stem cells *in vivo* and displays increased abundance during skeletal

muscle regeneration, especially during early regeneration suggesting that LMOD1 is important

for induction of myotube formation. Of note, knockdown of LMOD1 in primary myoblasts and

during skeletal muscle regeneration severely affects myogenic differentiation, while

overexpression accelerates and improves the formation of myotubes suggesting that LMOD1

is a critical component regulating myogenic differentiation. Mechanistically, we show that

LMOD1 physically and functionally interacts with the deacetylase sirtuin1 (SIRT1), a regulator

of myogenic differentiation, especially at the onset of myogenic differentiation. We

demonstrate that LMOD1 influences SIRT1 localization and the expression of a subset of its

target genes. Consistently, depletion or pharmacological inhibition of SIRT1 partially rescues

the impairment of myogenic differentiation observed after knockdown of LMOD1. Our work

identifies a new regulator of myogenic differentiation that might be targeted to improve muscle

regeneration in aging and disease.

Keywords: muscle stem cell, satellite cell, cytoskeleton, myogenesis, LMOD1, SIRT1, mass

spectrometry, myogenic differentiation, regeneration

Main Text

Muscle stem cells (MuSCs), also known as satellite cells, possess a remarkable ability to regenerate adult skeletal muscle following injury (Sambasivan et al. 2011; Murphy et al. 2011). These cells are typically quiescent, reside between the basal lamina and the myofiber (Mauro 1961; Sousa-Victor et al. 2015), and are characterized by the expression of the paired box transcription factor 7 (PAX7) (Seale 2003; Lepper, Partridge, and Fan 2011; von Maltzahn et al. 2013; Zammit et al. 2006). In response to physiological and regeneration-inducing stimuli, MuSCs become activated, begin to self-renew, and undergo myogenic differentiation (Fu, Wang, and Hu 2015; Schmidt et al. 2019). During myogenic differentiation after injury - a process recapitulating myogenesis during development - MuSCs proliferate, differentiate, and finally fuse to damaged myofibers or form new myofibers. These processes are tightly regulated by the expression of different myogenic regulatory factors (MRFs) such as MYF5, MYOD, myogenin and MRF4 (Chang and Rudnicki 2014; Schmidt et al. 2019; Tiffin et al. 2003). Additionally, external cues from the surrounding microenvironment play a crucial role in regulating myogenic differentiation (Bentzinger, Wang, Dumont, et al. 2013; Hung et al. 2023). These cues involve different signaling pathways including NOTCH, WNT, BMPs, FGFs, and Hedgehog (Fre et al. 2005; Baghdadi, Castel, et al. 2018; Baghdadi, Firmino, et al. 2018; Bentzinger, Wang, von Maltzahn, et al. 2013; Friedrichs et al. 2011; Larraín et al. 1997; Voronova et al. 2013). Epigenetic mechanisms and chromatin remodeling regulate the accessibility of transcription factors to regulatory regions of the DNA and, thereby, the expression of myogenic genes (Saccone and Puri 2010; Massenet et al. 2021). Histone deacetylases (HDACs) and methyltransferases, e.g., SIRT1 (Ryall et al. 2015), HDAC4 (Moresi et al. 2012; Finke et al. 2022) and SMYD3 (Codato et al. 2019) are implicated in maintaining muscle integrity and differentiation. The cytoskeleton plays crucial roles in regulating cell shape, motility and signal transduction (Pollard and Cooper 2009; Higuchi-Sanabria et al. 2018). This is especially important during fusion of myoblasts to form myotubes (Abmayr and Pavlath 2012). Perturbations in signalling pathways, mechanosensitivity, and interactions with the extracellular matrix (ECM) were shown to significantly affect the cytoskeletal architecture of MuSCs in disease and during aging (Chakkalakal et al. 2012; Hwang and Brack 2018; Lukjanenko et al. 2016). For instance, in Duchenne muscular dystrophy (DMD), mutations in the dystrophin gene can disrupt the connection between the actin cytoskeleton and the ECM, impairing the function of MuSCs (Winder, Gibson, and Kendrick-Jones 1995; Lu-Nguyen et al. 2017; Mournetas et al. 2021). Similarly, mutations in plectin, a protein that connects various cytoskeletal components, result in muscle diseases known as plectinopathies, which are characterized by compromised muscle cell integrity (Winter and Wiche 2013). Of note, also age-related sarcopenia can be caused by changes in cytoskeletal proteins, resulting in a decline in myofiber quantity and functionality (Budai,

Balogh, and Sarang 2018; Larsson et al. 2019). Changes in actin dynamics, e.g., a
dysregulation of ARP2/3 and N-WASP, responsible for forming branched actin networks and
cell motility, can impair the response to muscle injury (Lai and Wong 2020). Additionally, the
distribution of desmin, an essential protein for structural integrity, was shown to affect the
mechanical properties and resilience of myofibers (Meyer, Schenk, and Lieber 2013). Aging
also affects muscle regulatory factors like MyoD and myogenin, disrupting the timely
expression of cytoskeletal and contractile proteins required for proper differentiation of MuSCs
(Sousa-Victor et al. 2014).
Here, we set out to identify new regulators of myogenic differentiation by investigating
alterations in protein dynamics during myogenic differentiation using mass spectrometry.
Thereby, we identified LMOD1 among a restricted group of cytoskeletal proteins that increases
in abundance during early myogenic differentiation. We confirmed the presence of LMOD1 in
MuSCs *in vivo* and show that LMOD1 is transiently upregulated during skeletal muscle
regeneration and that reducing LMOD1 expression after injury impairs skeletal muscle
regeneration. LMODs are known as powerful actin filament nucleator but their role in myogenic
differentiation has not been investigated so far (Chereau et al. 2008; Boczkowska et al. 2015;
Fowler and Dominguez 2017; Tolkatchev, Gregorio, and Kostyukova 2022). Through
knockdown and overexpression experiments, we show that LMOD1, but not the closely related
LMOD2, is essential for proper myogenic differentiation. Mechanistically, we show that
LMOD1 interacts with the deacetylase SIRT1, a known regulator of myogenic differentiation
(Fulco et al. 2003; Ryall et al. 2015), and thereby influence its subcellular localization and the
expression of a subset of target genes.

Proteome dynamics during myogenic differentiation

To comprehensively and unbiasedly examine the protein dynamics during myogenic differentiation, we performed an *in vitro* five-day differentiation time course experiment using mouse primary myoblasts from five different mice (Figure 1A) and collected samples at each day of differentiation for label-free quantitative proteomics (Table S1). Principal Component Analysis (PCA) based on protein intensities obtained from mass spectrometry data (6098 quantified proteins in total) revealed a progressive separation of the different cell populations by the day of differentiation (Figure 1B). We confirmed the expected dynamics of key myogenic regulators by qRT-PCR and mass spectrometry (Schmidt et al. 2019), validating our experimental setup and proper myogenic differentiation (Figures 1C and 1D). Next, we compared protein dynamics by clustering the obtained proteomics data (Table S1). The 3839 protein groups significantly affected in at least one measured time point of differentiation relative to undifferentiated myoblasts were assigned to six clusters by k-means clustering (Figure 1E). Each cluster displayed distinct dynamics and was enriched for specific KEGG pathways (Figure 1F). Notably, cluster 2 included proteins that progressively increased in abundance during differentiation and were closely associated with muscle contraction and function as well as cytoskeletal dynamics e.g. Actinin Alpha 2 (ACTN2), Tropomyosin alpha-1 and -3 (TPM1, TPM3) Leiomodin 1 (LMOD1) and Dystrophin (DMD) (Figure 1G). These findings confirm the importance of cytoskeletal dynamics during myogenic differentiation but also raise questions about the molecular mechanism responsible for this shift in the proteomic landscape.

Figure 1: Analysis of the proteome shows dynamic alterations during myogenic differentiation
A. Experimental workflow for analyzing the five-day differentiation time course using mass spectrometry (Table S1) and qRT-PCR. Primary myoblasts isolated from five individual mice were seeded and differentiated for up to five days. **B**. Principal component analysis (PCA) of proteomics data. Ellipses represent a 95 % confidence interval for each day of differentiation. **C**. qRT-PCR analysis showing the relative mRNA expression of *Pax7*, *MyoD*, and *Myogenin*, normalized to *Gapdh* and day 0. **D**. Mass spectrometry-based quantification of PAX7, MYOD and MYOGENIN protein levels normalized to day 0. **E**. Pie chart for proteome dynamics. 6098 protein groups were subjected to k-means clustering analysis, 2259 were unaffected, while 3839 protein groups showed significant changes (Absolute AVG log2 ratio>0.58 and Q value<0.25) in expression levels in at least one-time point compared to day 0. Protein groups were classified into six clusters based on their expression dynamics using k-means clustering. **F.** Clusters showing distinct protein abundance dynamics during myogenic differentiation and KEGG pathways enriched (FDR <0.05) in each cluster (Table S1). **G.** Proteins from cluster 2 were chosen according to KEGG pathway annotation: striated muscle contraction and function, motor protein activity and cytoskeletal dynamics. The protein-protein interaction network was visualized with STRING (Szklarczyk et al. 2011), edge confidence: medium 0.4. For (**C**) and (**D**): In all bar plots, each symbol represents a biological replicate, and the error bars indicate the standard deviation. (SD). One-way, ANOVA. *: q value ≤ 0.05, **: q value ≤ 0.01, ***: q value ≤ 0.001.

LMOD1 levels increase during early myogenic differentiation, in aged MuSCs and regenerating skeletal muscle

Given the role of cytoskeleton remodeling as one of the drivers of myogenic differentiation, we focused on this cluster of proteins and specifically looked for proteins that increased early during the differentiation process. Among these, we found two proteins encoded by paralog genes, namely leiomodin 1 (LMOD1) and leiomodin 2 (LMOD2) (Figure 2A), which share 67.05 % sequence similarity, although varying in their number of amino acids and expression across tissues (Qualmann and Kessels 2009; Fowler and Dominguez 2017). Interestingly, the expression pattern of LMOD1 and LMDO2 during myogenic differentiation displayed differences: while LMOD1 showed a significant increase at the beginning (day 1) of myogenic differentiation, LMOD2 increased later in the process (Figure 2A) suggesting that LMOD1 might be a key protein in driving early myogenic differentiation. To assess the *in vivo* relevance of our finding, we queried two proteomic datasets of freshly isolated MuSCs and four different skeletal muscles (gastrocnemius, G; soleus, S; tibialis anterior, TA; extensor digitorum longus, EDL) (Schüler et al. 2021). We found LMOD2 to be the most abundant leiomodin in whole skeletal muscle, as expected (Yuen et al. 2022; Tsukada et al. 2010), while the overall abundance of LMOD1 was lower since this protein has been mainly associated with smooth muscle cells (Nanda and Miano 2012; Conley et al. 2001; Nanda et al. 2018) (Figure 2B). Surprisingly, we detected LMOD1, but not LMOD2, in freshly isolated MuSCs, and noted that the protein levels of LMOD1 increased in MuSCs isolated from old mice (Figure 2C and Figure S1B). To independently validate this finding, we performed immunofluorescence stainings on cryosections of TA muscles from young and geriatric mice. Consistent with previous publications, we observed a reduction in the number of PAX7+ cells with increasing age (Brack, Bildsoe, and Hughes 2005; Shefer et al. 2006; Yamakawa et al. 2020) (Figure S1C). Furthermore, we confirmed the age-related increase of LMOD1 abundance in MuSCs (Figure 2D and Figure S1B). Of note, this was accompanied by a significant increase in the number of LMOD1+/PAX7+ cells in sections of geriatric compared to young mice (Figure 2E). Interestingly, we found LMOD1 to be mainly located in the cytoplasm of PAX7+ cells in TA muscles from young mice (Figure 2D and S1D). However, in TA muscles from geriatric mice, we found a significantly higher number of PAX7+ cells that displayed nuclear localization of LMOD1 (Figure 2F), suggesting that LMOD1 accumulates and mis-localizes in MuSCs with increasing age and could be one contributing factor to the reduced ability of MuSCs in aged mice to differentiate.

Finally, we investigated the expression of LMOD1 during regeneration of skeletal muscle *in vivo.* Therefore, we performed a regeneration time course experiment following cardiotoxin (CTX) injury and assessed LMOD1 expression by immunoblot and immunofluorescence analyses in young mice (Figure 2G and Figure 2H). We observed low abundance of LMOD1

in uninjured TA muscle but found a progressive increase in abundance up to day 7 after injury
and a decrease thereafter (Figure 2G). Immunofluorescence staining confirmed that LMOD1
was highly abundant in myofibers at day 5 after injury when mostly newly formed regenerating
myofibers are present, hinting at a functional role of LMOD1 in early myogenic differentiation.
Of note, LMOD1 abundance was barely detectable at day 7 and later time points suggesting
that LMOD1 abundance decreases during maturation of myofibers (Figure 2H). In conclusion,
these findings demonstrate that LMOD1 abundance is not only increased at the onset of
myogenic differentiation *in vitro* but is also expressed by MuSCs under homeostatic conditions
in vivo and in MuSC and newly formed myofibers following injury, indicating its importance for
myofiber differentiation and skeletal muscle regeneration.

Figure 2: LMOD1 increases during early myogenic differentiation and is upregulated in aged MuSCs
A. Protein quantification of LMOD1 and LMOD2 during myogenic differentiation, based on mass spectrometry data and normalized to the protein amount of day 0. Error bars indicate SD, and colored dots indicate the mean value of the biological replicates (n = 5). **B.** LMOD1 and LMOD2 protein abundance estimated from mass spectrometry

data in different skeletal muscles (gastrocnemius, G; soleus, S; tibialis anterior, TA; extensor digitorum longus, EDL). Displayed values are averages of n = 5 samples from individual mice. Mass spectrometry data from (Schüler et al. 2021). **C**. LMOD1 and LMOD2 protein quantity in MuSCs obtained from young (n = 10), old (n = 8) and geriatric (n = 5) mice, nd = not detected. Mass spectrometry data from (Schüler et al. 2021). **D**. Representative immunofluorescence images of LMOD1 (purple), PAX7 (orange), and Hoechst (blue) of TA sections from n = 3 individual mice per age group: young (3-month-old) and old (18-months-old), geriatric (33-months-old) mice. Scale bar: 5 µm. **E.** Ratio of PAX7+ and LMOD1+ cells normalized to the cells positive for PAX7+ (Figure S2C) and **F.** Quantification of the percentage of PAX7+ cells showing nuclear LMOD1 localization. Related to Figure S1. For (**E**) and (**F**): One-way, ANOVA. *: q value ≤ 0.05, **: q value ≤ 0.01, ***: q value ≤ 0.001. In all bar plots, each black dot represents a biological replicate, and the error bars indicate the SD. **G**. Muscle regeneration time course: Immunoblot of LMOD1 levels in regenerating TA muscle after injury. Quantification of LMOD1 levels normalized to Ponceau and relative to the LMOD1 signal at day 0. Each symbol represents a biological replicate, and error bars represent SD. One-way ANOVA; **: p-value ≤ 0.01, ***: p-value ≤ 0.001. **H**. Representative immunofluorescence images of LMOD1 (purple), PAX7 (orange), LAMININ (grey), and Hoechst (blue) of TA sections at different days post-injury from young mice (3 months old). The white arrowhead shows PAX7+ cells at day 0, white arrows indicate newly formed LMOD1+ myofibers at day 5. Scale bar: 50 µm.

Figure S1: LMOD1 in MuSCs and skeletal muscle
A. LMOD1 protein coverage in proteomics data from MuSCs. Multiple unique (proteotypic) peptides were identified. Data from (Schüler et al. 2021). **B**. Immunofluorescence images of TA sections from different mice (n = 3) per age group of young (3 months old) and geriatric (33 months old) mice. PAX7 (orange), LMOD1 (purple) and Hoechst (blue). **C**. Number of PAX7+ cells per mm^2 cells in muscle sections from young, old and geriatric mice and **D**. Quantification of PAX7+ cells showing cytoplasmic LMOD1 localization. For (**C**) and (**D**): One-way ANOVA; *: p-value ≤ 0.05, ***: p-value ≤ 0.001, ns: not significant. In all bar plots, each black dot represents a biological replicate, and the error bars indicate the SD. Related to Figure 2.

Knockdown of *Lmod1* impairs myogenic differentiation, while overexpression of *Lmod1* accelerates myotube formation

The early increase of abundance of LMOD1 during myogenic differentiation suggests a potential functional role for this protein in promoting the initiation of myogenic differentiation. To test this hypothesis, we used a small interfering RNA (siRNA) to reduce LMOD1 expression and analyzed the impact on myoblast proliferation (Figure S2A) and differentiation (Figure 3A). We confirmed the successful knockdown of *Lmod1* mRNA by qRT-PCR (Figure S2B and S2F) and reduced levels of LMOD1 protein by immunoblot analysis (Figure S2C and S2G). After 48 hours of proliferation, we observed a slight (26 %) but significant decrease in the number of proliferating Ki67-positive myoblasts upon knockdown of LMOD1 (Figure S2D and S2E). During myogenic differentiation, the cell population becomes heterogeneous with a distinct expression of myogenic markers (such as MyoD, myogenin, and devMHC) determining cellular identity thereby controlling the potential to differentiate and ultimately form new myofibers (Figure 3B). We characterized the cell populations after knockdown of LMOD1 based on the expression of myogenic markers using immunofluorescence staining at day three of differentiation (Figure 3C and S2M-Q). We found that the reduced levels of LMOD1 resulted in an increased frequency of cells with a pronounced spherical phenotype, also identified as non-elongated, MHC-positive cells (Figure 3C and 3E). Moreover, a significant reduction in the number of nuclei per myotube was observed upon knockdown of LMOD1, indicating a potential role in the fusion process (Figure 3D). Furthermore, we observed significantly shorter myotubes upon knockdown of LMOD1, supporting our previous finding that the cells are unable to properly differentiate (Figure 3F).

Given the structural similarities between LMOD1 and LMOD2, we set out to determine if individual knockdown in primary myoblasts induces similar phenotypes during myogenic differentiation (Figure 3G). Knockdown of both *Lmod1* and *Lmod2* mRNA was validated by qRT-PCR (Figure S2I) and confirmed at the protein level by immunoblot (Figure S2J). Interestingly, the knockdown of LMOD2 did not lead to any noticeable phenotypes compared to primary myoblasts transfected with a scramble control siRNA (Figure 3H and S2K). However, we observed a significant reduction in the number of nuclei per myotube (Figure 3I), increased frequency of non-elongated MHC-positive cells (Figure 3J), and increased number of shorter myotubes (Figure 3K) in the *Lmod1/2* double knockdown. These data suggest that LMOD1 is required for myogenic differentiation in primary myoblasts, while LMOD2 appears to be dispensable.

To examine how reduced levels of LMOD1 affect skeletal muscle regeneration *in vivo*, we injured the TA-muscle of 3-months-old mice using CTX, followed by injection of a self-delivering siRNA targeting Lmod1 (Figure 3L) at day 3 after injury, a time point when MuSCs start to differentiate. Subsequently, TA-muscles were isolated at day 5 after injury, the time

when LMOD1 was previously shown to be highly expressed in newly formed myofibers.
Knockdown of *Lmod1* led to a significant reduction in the number of cells expressing the early
differentiation marker myogenin (Figure 3M and 3N), suggesting a critical role for LMOD1 in
the transition of MuSCs from proliferation to differentiation. Together, these findings indicate
that LMOD1 is not only important for myofiber formation but may also play a key regulatory
role in the differentiation of MuSCs into myoblasts and myocytes during the muscle
regeneration process stressing its role in controlling myogenic differentiation.
To test whether LMOD1 alone is sufficient to promote myogenic differentiation, we generated
six independent primary myoblast cultures that were stably transduced with a virus to
overexpress either LMOD1 or GFP (Figure 3O). Overexpression (OE) of Lmod1 mRNA by
qRT-PCR (Figure S2L) and protein by immunoblot (Figure S2M) was confirmed by comparison
to a GFP OE control. Immunofluorescence analysis (Figure 3P and S2O) after one day of
differentiation showed that overexpression of LMOD1 promotes myogenic differentiation, as
shown by a higher number of nuclei per myotube (Figure 3Q). Additionally, we observed an
accelerated formation of fully differentiated myotubes (Figure 3Q) and an increase in myotube
length (Figure S2N) in LMOD1 OE cells compared to the GFP OE control construct. To further
characterize the molecular alterations induced by overexpression of LMOD1 during myogenic
differentiation, we generated proteomics data from LMOD1 overexpressing and GFP control
cells at multiple time points of differentiation (Table S2). Using gene set enrichment analysis,
we found that at day 1 of differentiation, cells overexpressing LMOD1 exhibit higher levels of
proteins that are up-regulated at the beginning of differentiation, such as HSPB3, ALDH3A1,
SMTNL2, ACTG1, when compared to the GFP control group (Figure S2P) (Table S2). This
proteome signature corroborates the observation that LMOD1 overexpression is sufficient to
promote myogenic differentiation.

273
Figure 3: Knockdown of *Lmod1* impairs myogenic differentiation and reduces muscle regeneration, while
overexpression enhances it.
A. Schematic of the *Lmod1* knockdown experiment after 3 days of differentiation. siRNA directed against *Lmod1*
or scramble (siCtrl) control was used to transfect primary myoblasts isolated from individual mice. Differentiation
was induced by a change to differentiation medium at 0h. **B.** Overview of quantified different cell types based on
their expression of myogenic markers; Myogenin (red), Myosin heavy chain (MHC) (green), Hoechst (blue). Scale
bar: 50 µm. (a) reserve cells: Hoechst+/Myogenin-/MHC- and undifferentiated. (b) Myogenin+/MHC-: Myogenin
positive, MHC negative, just started to differentiate. (c) Myogenin+/MHC+: co-express bot in the process of
differentiation. (d) Myogenin-/MHC+: non elongated. (e) Fully differentiated myotubes: quantified by nuclei count
to assess the fusion process. **C.** Representative immunofluorescence images after three days of differentiation and
siCtrl or siLmod1 transfection; MHC (green), nuclei (blue). Scale bar: 100 µm. **D - F.** Quantification of the number
of nuclei per myotube (**D**), Myogenin-/MHC+ cells (**E**) and length of differentiated myotubes (in µm) (**F**). Paired t-
test; *: p-value ≤ 0.05, **: p-value ≤ 0.01. **G.** Schematic of the *Lmod2* and *Lmod1/Lmod2* double knockdown
experiment after 3 days of differentiation. siRNA against *Lmod1*, *Lmod2*, both or scramble (siCtrl) was used to
transfect primary myoblasts. Differentiation was induced by a change to differentiation medium at 0h. **H.**
Representative immunofluorescence images of siLmod2 and siLmod1/Lmod2 double knockdown after three days
of differentiation; devMHC (green), nuclei (blue). Scale bar: 50 µm. **I - K.** Quantification of the number of nuclei per
myotube (**I**), Myogenin-/MHC+ cells (**J**), Length of differentiated myotubes (in µm) (**K**). One-way ANOVA; *: p-value
≤ 0.05, ns: not significant. **L.** Experimental schematic for analysis of *in vivo* CTX-induced injury of TA muscles
combined with injection of self-delivering siRNAs at 3 days post injury. n = 4 mice per group, 3 months old.
M. Representative immunofluorescence images of MYOGENIN (green), LAMININ (grey), and Hoechst (blue) of
TA-sections 5 days post-injury from young mice (3 months old). **N.** Quantification of the number of Myogenin+ cells
normalized per area. Unpaired t-test; *: p-value ≤ 0.05. **O.** Illustration of the experimental setup. Primary myoblasts

isolated from individual mice stably overexpressing (OE) LMOD1 (purple) or GFP (green) were seeded and either collected during proliferation (3d proli), after 1 day (1d diff), 3 days (3d diff) or 5 days (5d diff) of differentiation. Differentiation was induced by a change to differentiation medium at 0h. Cells were harvested for immunofluorescence analysis (IF), qRT-PCR, immunoblot (WB) or mass spectrometry (MS). **P.** Representative immunofluorescence images of primary myoblasts after one day of differentiation, showing either stable expression of LMOD1 or GFP. devMHC (green), MYOGENIN (red) and nuclei (blue). Scale bar is 50 µm. **Q** and **R.** Quantification of nuclei per myotube of GFP OE or LMOD1 OE cells (**Q**) and cells expressing Myogenin+/MHC+ defined as just differentiated cells (**R**) after 1 day of differentiation. Paired t-test *: p-value ≤ 0.05. Related to Figure S2. In all bar plots, each symbol represents a biological replicate, and the error bars indicate the SD.

307
308 **Figure S2: Validation of knockdown efficiency and statistical analysis of siLmod1/Lmod2 knockdown and**
309 **LMOD1 overexpression**

310
A. Schematic of the *Lmod1* knockdown experiment under proliferating conditions. siRNA directed against *Lmod1*
or scramble (siCtrl) control was used to transfect primary myoblasts isolated from individual mice. **B.** qRT-PCR
analysis showing the relative expression of *Lmod1* after siLmod1 knockdown compared to siCtrl transfected cells,
normalized to *Gapdh* expression levels. **C.** Immunoblot and quantification of LMOD1 after transfections with
siLmod1 and siCtrl and two days of proliferation normalized to GAPDH. Paired t-test: *: p-value ≤ 0.05.
D. Immunofluorescence analysis of primary myoblasts stained for the proliferation marker Ki67 (red) and Hoechst
(blue). Scale bar: 100 μm. **E** Quantification of cell populations identified by immunofluorescence staining: reserve
cells being Hoechst+/Ki67- and cells being Hoechst+/Ki67+ in siCtrl and siLmod1 transfected conditions and the
total counted cells per condition. **F.** and **G.** Validation of the siLmod1 knockdown compared to siCtrl transfected
cells under differentiation conditions. qRT-PCR for *Lmod1* normalized to *Gapdh* expression levels (**F**) and
immunoblot analysis with quantification of LMOD1 signal normalized to GAPDH (**G**). Paired t-test *: p-value ≤ 0.05,
***. **H.** Quantification of cell populations identified by immunofluorescence staining: reserve (Hoechst+/Myogenin-
/MHC-) cells, Myogenin+/MHC- cells, Myogenin+/MHC+, fully differentiated myotubes and total counted cells under
differentiating conditions. **I.** and **J.** Validation of the *siLmod1* or *siLmod2* (single) or *siLmod1/Lmod2* (double)
knockdown compared to siCtrl transfected cells under differentiation conditions. qRT-PCR for *Lmod1* and *Lmod2*
normalized to *Gapdh* expression levels (**I**) and representative immunoblot with quantification of LMOD2 normalized
to GAPDH (**J**). One-way ANOVA *: p-value ≤ 0.05 or numbers are indicated. ns: not significant. **K.** Quantification
of cell populations (in %) identified by immunofluorescence staining after siLmod1 (single), siLmod2 (single) and
siLmod1/2 (double) knockdown under differentiating conditions: reserve (Hoechst+/Myogenin-/MHC-) cells,
Myogenin+/MHC- cells, Myogenin+/MHC+, fully differentiated myotubes and total counted cells under
differentiating conditions. Related to Figures 3I and 3J. **L** and **M.** Validation of LMOD1 OE compared to GFP OE
cells after one day of differentiation. Relative *Lmod1* expression assessed by qRT-PCR was compared to GFP OE
cells and normalized to *Gapdh* expression levels (**L**) and immunoblot with quantification of LMOD1 normalized to
GAPDH (**M**). Paired t-test: *: p-value ≤ 0.05. **N** and **O.** Quantification of cell populations identified by
immunofluorescence staining after LMOD1 OE compared to GFP OE: reserve (Hoechst+/Myogenin-/MHC-) cells,
Myogenin+/MHC- cells, Myogenin+/MHC+, fully differentiated myotubes, total counted cells after and myotube
length (in μm) (**N**). Paired t-test: numbers are indicated, ns: not significant. Related to Figures 3N and 3O. **P.** Gene
set enrichment analysis (GSEA) was based on a gene set containing proteins that significantly increased at one
day of differentiation compared to 0h/proliferating primary myoblasts (AVG Log2 Ratio > 0.58 and Q value < 0.05,
189 proteins) from the proteomic data generated in (Figure 1A). The GSEA was performed using this gene set on
the comparison LMOD1 OE vs. GFP OE at 1 day of differentiation (Table S2). In all bar plots, each symbol or black
dot represents a biological replicate, and the error bars indicate the SD.
343
344
LMOD1 interacts with SIRT1 and SIRT2
To elucidate the mechanism by which LMOD1 contributes to myogenic differentiation, we
performed a proximity-dependent biotinylation experiment (BioID) for LMOD1 to identify its
interaction partners in HEK293T cells (Figure 4A). Given their sequence similarity but distinct
functions during myogenic differentiation, we analyzed the interaction partners of LMOD1 and
LMOD2 in parallel and focused on hits retrieved exclusively by LMOD1. Therefore, we fused
the promiscuous biotin ligase (BirA*) C-terminal to Lmod1 or Lmod2. A cell line expressing
only the sequence for BirA* was used as a control (BirA*-Ctrl) to account for non-specific
biotinylation. The tetracycline-dependent expression of the respective fusion proteins was
validated by immunofluorescence (Figure 4B and 4C) and anti-FLAG immunoblot (Figure
S3A). Biotinylation activity, after the addition of exogenous biotin, was assessed using
streptavidin-HRP immunoblot (Figure S3A) and immunofluorescence staining (Figure 4B and
4C). Subsequently, biotinylated proteins were captured using streptavidin enrichment and
analyzed by mass spectrometry (Table S3). PCA analysis demonstrated a distinct separation
between C-terminal fused LMOD1 and LMOD2 expression constructs (Figure 4D). Using both

LMOD1 and LMOD2, we found a significant enrichment compared to the BirA*-Ctrl for known

interaction partners (Fowler and Dominguez 2017; Boczkowska et al. 2015) involved in actin

filament nucleation such as Tropomodulins (TMODs) and Tropomyosins (TPMs) (Figure S3B-

D), validating our constructs. Interestingly, a direct comparison of streptavidin-enriched

proteins from LMOD1-BirA* vs. LMOD2-BirA* revealed a subset of interaction partners that

specifically interact with LMOD1. These include the histone deacetylases sirtuin 1 (SIRT1)

and sirtuin 2 (SIRT2), as well as the RuvB Like AAA ATPase 1/2 (RUVBL1/2) (Figure 4E and

4F). Sirtuins are involved in various biological processes, including DNA regulation,

metabolism, longevity, and myogenesis (Horio et al. 2011; Abdel Khalek et al. 2014;

Vachharajani et al. 2016). In particular, one study revealed that mice with muscle-specific

inactivation of Sirt1 displayed repression of the myogenic program (Ryall et al. 2015). Since

SIRT1 was already identified as a potential mediator of myogenesis, we wanted to validate its

preferential interaction with LMOD1 by co-immunoprecipitation independently. Therefore, we

transfected SIRT1 as a GFP fusion protein in HEK293T expressing LMOD1-BirA* or LMOD2-

BirA* and performed pull-down experiments using a GFP trap. Immunoblot analysis of the

GFP-trap eluates using anti-BirA antibodies confirmed a co-precipitation between LMOD1 and

SIRT1, while LMOD2 co-precipitated with SIRT1 to a lower extent (Figure 4G). To further

validate the interaction of LMOD1 and SIRT1 in primary myoblasts and myogenic

differentiation, we performed *in situ* detection of LMOD1 and SIRT1 using proximity ligation

assays (PLA) (Figure 4H). In proliferating myoblasts, when the abundance of LMOD1 is low,

only a weak PLA signal indicating close proximity of SIRT1 and LMOD1 was detectable both

in the cytoplasm and the nucleus. However, at the beginning of myogenic differentiation, when

the abundance of LMOD1 increases, an increase in PLA signals indicating close proximity of

LMOD1 and SIRT1 was detectable almost exclusively in the cytoplasm. Together these data

show that LMOD1 preferentially interacts with the deacetylase SIRT1, suggesting a potential

functional interaction between these two proteins in modulating myogenic differentiation.

Figure 4: LMOD1 interacts with SIRT1

A. BioID workflow. Lmod1 and Lmod2 were C-terminally fused to a promiscuous biotin ligase (BirA*) and expressed

in HEK293T cells. BirA* alone served as a control (Ctrl) to assess non-specific biotinylation. Overexpression of

fusion proteins was induced by addition of tetracycline. Exogenous biotin was introduced to label interaction

partners in close proximity. Biotinylated proteins were captured using streptavidin enrichment, followed by mass

spectrometry analysis to identify and quantify proximal interactors. **B.** and **C.** Immunofluorescence analyses of

LMOD1-BirA*-FLAG and LMOD2-BirA*-FLAG in HEK293T cells 4 days after seeding without addition of any

substance (-tet/-bio), with addition of only tetracycline for 4 days (+tet/-bio) or with addition of both tetracycline for

4 days and biotin for 1 day (+tet/+bio); FLAG (green), Streptavidin (red), nuclei (blue). Scale bar: 20 µm. **D.** Principal

component analysis (PCA) of the BioID data. Ellipses represent 95 % confidence intervals. **E.** Volcano plot of

proteins enriched by streptavidin pull-down and analyzed by mass spectrometry from LMOD1-BirA* and LMOD2-

BirA*. n = 4 biological replicates (Table S3) **F.** Quantification of selected unique interaction partners of LMOD1-

BirA* and LMOD2-BirA* in comparison to BirA*-Ctrl. Each symbol represents a biological replicate, error bars

indicate the SD. One-way ANOVA, *: p-value ≤ 0.05, **: p-value ≤ 0.01. **G.** Validation of the SIRT1-LMOD1

interaction. Cells expressing either GFP or SIRT1-GFP were used for co-immunoprecipitation using GFP-trap. The

eluates from the GFP-trap were analyzed by immunoblot using antibodies directed to BirA* or GFP. For

quantification, BirA* was normalized to SIRT1-GFP intensity. **H.** Representative images of proximity ligation assay

(PLA) (red) in primary myoblasts during proliferation and after one day of differentiation. Scale bar: 10 µm.

Figure S3: BioID for LMOD1 and LMOD2

A. Immunoblot of BirA* fusion proteins performed on lysates from HEK293T cells stably transfected with LMOD1-
BirA*-FLAG, LMOD2-BirA*-FLAG or Ctrl-BirA*-FLAG following 24 h incubation with (+tet) or without (−tet)
tetracycline. Middle panel, streptavidin-HRP blot following induction of BirA* fusion proteins with tetracycline and
supplementation of biotin for 24 h. Ponceau stainings were used as loading control. HRP: horseradish peroxidase.
B. Volcano plot of proteins enriched by streptavidin pull-down and analyzed by mass spectrometry from LMOD1-
BirA* and **C.** LMOD2-BirA* compared to BirA*-Ctrl. Significantly enriched (AVG Log2 Ratio > 0.58 and Q value <
0.05), known interaction partners for LMOD1 and LMOD2 are highlighted in red (Table S3). **D.** Quantification of
selected known interaction partners of LMOD1-BirA* and LMOD2-BirA* in comparison to BirA*-Ctrl. Each symbol
represents a biological replicate, and error bars indicate the SD. One-way ANOVA, *: p-value ≤ 0.05, **: p-value ≤
0.01.

LMOD1 and SIRT1 show dynamic subcellular localization during myogenic
differentiation

Next, we investigated how LMOD1 might influence SIRT1 activity and thereby myogenic

differentiation. SIRT1 has been shown to localize both in the cytoplasm and nucleus of neural

precursor cells, murine pancreatic beta cells, rat cardiomyocytes, and vascular endothelial

cells, while LMODs have been mainly associated with cytoplasmic localization (Hisahara et

al. 2008; Tanno et al. 2010; Tong et al. 2013; Tanno et al. 2007; Conley 2001; Nauen et al.

2020). To determine the localization of SIRT1 and LMOD1 and a potential change in

localization during myogenesis, we performed co-immunostainings for LMOD1 and SIRT1 in

primary myoblasts to determine their localization at different stages of myogenic differentiation

(Figure 5A). Overall, the total fluorescence intensity of LMOD1 significantly increased during

differentiation (Figure S4A), while the signal intensity of SIRT1 slightly decreased (Figure

S4B). However, under proliferating conditions, SIRT1 and LMOD1 predominantly localized in

the nucleus (Figure 5A). Interestingly, after initiation of differentiation, both SIRT1 and LMOD1

were clearly detectable in the cytoplasm, indicating their translocation into the cytoplasm. At

a later time point (2d of differentiation), the localization of SIRT1 can once again be observed

in the nucleus, while LMOD1 remains in the cytoplasm. Using the fluorescent signal intensity,

we quantified the cytoplasmic/nuclear ratios for both SIRT1 and LMOD1 and found that both

proteins change their intracellular distribution during myogenic differentiation suggesting that

their localization in the cell affects the differentiation process (Figures 5B and 5C). Of note, we

also observed a correlation between their cytoplasmic-nuclear partitioning, particularly during

proliferation and at day one of differentiation (Figure 5D and 5E). The correlation between their

respective localization was reduced at day 2 of differentiation, with SIRT1 displaying mainly a

nuclear localization while LMOD1 remained predominantly cytoplasmic (Figure 5F). We

confirmed these dynamic changes of sub-cellular localization for SIRT1 and LMOD1 by

cellular fractionation experiments followed by immunoblot analysis (Figure S4C).

Since SIRT1 protein abundance decreased with each day of differentiation as shown by our

proteomics data and immunoblot analysis (Figure 5G and S4D), we hypothesized that the

LMOD1-mediated change in SIRT1 location is especially required during early differentiation.

Therefore, we repeated the cellular fractionation experiments focusing on the onset of

myogenic differentiation with additional time points and by using LMOD1 OE cells. Next, we

compared the nuclear fraction of the LMOD1 OE cells to the control cells to determine if

LMOD1 alone can alter SIRT1 subcellular localization (Figure 5H-K). In particular, at 6 and 12

hours of differentiation, SIRT1 is highly abundant in the nucleus in control cells. However,

these dynamics shift and the SIRT1 abundance decreases in the nucleus when LMOD1 is

overexpressed (Figure 5H and 5I). In line with this observation, in LMOD1 overexpressing

cells the cytoplasmic SIRT1 abundance increased at 0h (proliferation) and 6h differentiation,

suggesting a shift of SIRT1 protein to the cytoplasm upon LMOD1 OE compared to the control

already under proliferating conditions (Figure 5J and 5K). Together these data demonstrate

that SIRT1 and LMOD1 dynamically change their subcellular localization and that

overexpression of LMOD1 favors the cytoplasmic localization of SIRT1 at early stages of

myogenic differentiation.

Figure 5: Overexpression of LMOD1 influences the subcellular localization of SIRT1.

A. Representative immunofluorescence staining of LMOD1 and SIRT1 at different timepoints of myogenic differentiation; LMOD1 (purple), SIRT1 (yellow), nuclei (Hoechst, blue). Scale bar: 20 µm. **B - C.** Cytoplasmic to nuclear SIRT1 (**B**) and LMOD1 (**C**) ratio at different days of differentiation, n = 50 cells were analyzed per biological replicate per time point. **D - F.** Correlation of the cytoplasmic to nuclear ratio of LMOD1 and SIRT1 at 2 days of proliferation (R^2 = 0.47) (**D**), 1 day differentiation (R^2 = 0.42) (**E**) and 2 days of differentiation (R^2 = 0.08) (**F**). **G.** SIRT1 protein abundance upon LMOD1 OE at different timepoints, 3 days proliferation (3dp), 1 day differentiation (1dd), 3 days differentiation (3dd), 5d differentiation (5dd). Each symbol represents a biological replicate (n = 4, primary myoblasts); error bars indicate SD. **H - K.** Representative immunoblot of SIRT1 in the nuclear (**H** and **I**) and cytoplasmic (**J** and **K**) fraction, comparing GFP and LMOD1 OE at early time points of differentiation (0h: undifferentiated/proliferating, 6h, 12h, 24h of differentiated primary myoblasts). SIRT1 signal was analyzed in n = 3 immunoblots and was normalized to Ponceau. In bar plots, each symbol represents a biological replicate, and the error bars indicate the SD. Paired t-test *: p-value ≤ 0.05, ns: not significant.

Figure S4: Sub-cellular localization of LMOD1 and SIRT1 during myogenic differentiation

A - B. Total immunofluorescence intensity (nucleus and cytoplasm) of LMOD1 (**A**) and SIRT1 (**B**) during different timepoints of myogenic differentiation. **C.** Representative immunoblot of the nuclear-cytoplasmic fractionation experiment in primary myoblasts at different timepoints: 2d proliferation, 1d differentiation and 2d differentiation. SIRT1 and LMOD1 signal was analyzed in (n = 3) immunoblots and was normalized to Ponceau. ALDOA signal was used as a cytoplasmic marker, and Histone 4 (H4) was used as a nuclear marker. Immunoblot analyses of LMOD1 and SIRT1 intensity for each cellular compartment are depicted in the barplots. One-way ANOVA *: p-value ≤ 0.05, ns: not significant. **D.** Immunoblot showing LMOD1 and SIRT1 protein abundance at different timepoints of myogenic differentiation (0h = proliferating conditions, 6h differentiation, 12h differentiation, 24h differentiation). Quantification of immunoblots for LMOD1 and SIRT1 shown in barplots normalized to GAPDH. For all bar plots, each symbol represents a biological replicate, and the error bars indicate the SD.

Reduction or inhibition of SIRT1 can partially restore the impaired myogenic differentiation caused by knockdown of LMOD1

Given the interaction of LMOD1 and SIRT1, we hypothesized that LMOD1 regulates SIRT1-mediated transcription of genes that are critical for myogenic differentiation. To test this hypothesis, we investigated the impact of reduced SIRT1 levels using both a genetic (siRNA targeting *Sirt1*) and a pharmacological approach (25 µM EX527, small molecule inhibitor specific for SIRT1) during myogenic differentiation (Figure 6A) (Peck et al. 2010; Broussy, Laaroussi, and Vidal 2020). Primary myoblasts were transfected with siLmod1 and siSirt1 or incubated with EX527, at the time of induction of differentiation. After three days of differentiation, immunofluorescence staining was performed and analyzed (Figure 6B). We confirmed the significant decrease in LMOD1 and SIRT1 levels upon siRNA-mediated knockdown through immunoblot analysis (Figure S5A) and qRT-PCR (Figure S5B). Our results show that both genetic and pharmacological inhibition of SIRT1 signaling effectively promoted myogenic differentiation (Figure 6B and 6F) leading to an increased number of

nuclei per myotube in the siLmod1/siSirt1 double knockdown compared to the knockdown of
LMOD1 alone, indicating an improvement in myotube fusion after knockdown of SIRT1
(Figure 6C, 6G and Figure S5C and S5D). Moreover, both interventions were able to
counteract the negative effects observed after knockdown of LMOD1 alone (Figure 6D and
6H). Previous studies have demonstrated that *Sirt1* knockout mice exhibited premature
differentiation of MuSCs (Ryall et al. 2015). As a measure of myogenic differentiation, we
measured the length of myotubes and observed an elongation after knocking down SIRT1
together with LMOD1 (Figure 6E and 6I).
Next, we performed RNA-sequencing to identify downstream effectors that could explain the
improvements in myogenic differentiation observed after double knockdown of Lmod1 and
Sirt1 (Table S4). Therefore, we performed a knockdown of *Lmod1* and *Sirt1* two days prior to
differentiation, followed by a second siRNA transfection at the initiation of differentiation to
ensure a high knockdown efficiency. After one day of differentiation, cells were harvested to
quantify early transcriptional changes in the myogenic program after knockdown of *Lmod1*
and *Sirt1* using RNAseq (Figure 6J). Differential gene expression analysis confirmed a
significant reduction of *Lmod1* and *Sirt1* mRNA in our experimental setup (Figure S5E). Next,
we identified 246 genes being significantly and oppositely regulated after knockdown of either
Lmod1 or *Sirt1* (Table S4). Interestingly, among these genes, we identified a subset of known
SIRT1 target genes that were previously defined by ChIP-Seq analysis of MuSCs from WT
and Sirt1[mKO] mice (Ryall et al. 2015) (Table S4). These include key factors involved in cell
signaling and differentiation, such as *Fgf6*, *Tgfa*, *Mcr*, *Lifr*, and *Camk1d* (Figure 6K). Taken
together, our data show that genetic or pharmacological inhibition of SIRT1 signaling can
partially reverse the impaired myogenic differentiation caused by knockdown of Lmod1 and
that affecting LMOD1 level is sufficient to alter the expression of a subset of SIRT1 target
genes, highlighting the role of the LMOD1-SIRT1 axis in myogenic differentiation.

Figure 6: Reduced SIRT1 signaling can partially reverse siLmod1-induced impaired myogenic differentiation

A. Illustration of the experimental setup. Primary myoblasts were co-transfected with *Lmod1*-specific siRNA (siLmod1) and siRNA against Sirt1 (siSirt1) (depicted in orange) or treated with SIRT1 inhibitor EX527 (depicted in pink), simultaneously to the induction of differentiation. **B./F.** Representative immunofluorescence images after *Lmod1* and *Sirt1* knockdown with siRNA at day 3 of differentiation (**B**) or after *Lmod1* knockdown and addition of 25 μM EX527 at day 3 of differentiation (**F**); MYOGENIN (red), devMHC (green), nuclei (Hoechst, blue). **C - E.** Quantification of the number of nuclei per myotube (**C**), percentage of Myogenin-/MHC+ cells (**D**) and measured length of differentiated myotubes (**E**) after siRNA transfection of SIRT1 and LMOD1. One-way ANOVA *: p-value ≤ 0.05, **: p-value ≤ 0.01. **G - I.** Quantification of the number of nuclei per myotube (**G**), percentage of Myogenin-/MHC+ cells (**H**) and measured length of differentiated myotubes (**I**) after siRNA knockdown of LMOD1 and EX527 treatment to inhibit SIRT1. One-way ANOVA *: p-value ≤ 0.05, **: p-value ≤ 0.01. **J.** Illustration of the experimental setup for the RNA-Seq experiment. Primary myoblasts were first seeded and then transfected with siRNA against LMOD1 or SIRT1 or siCtrl. After two days of incubation, the primary myoblasts were transfected a second time with siRNA and induced to differentiate simultaneously. After one day of differentiation, cells were harvested, and RNA was isolated for library preparation and RNAseq analysis. **K.** Heatmap indicating oppositely differentially expressed SIRT1 target genes from siSirt1 vs siCtrl an siLmod1 vs siCtrl with a log2 FC ratio > 0.58. SIRT1 target genes identified from ChipSeq experiment published in (Table S4) (Ryall et al. 2015). For all bar plots, each symbol represents a biological replicate, and the error bars indicate the SD.

Figure S5: Reduced SIRT1 signaling can partially reverse siLmod1-induced impaired myogenic differentiation

A. Immunoblot validation of the LMOD1 and SIRT1 single knockdown and double knockdown (siLmod1/siSirt1) or a scrambled siRNA. Primary myoblasts were transfected at the initiation of differentiation and harvested after three days of differentiation. One-way ANOVA: *: p-value ≤ 0.05. **: p-value ≤ 0.01. **B.** qRT-PCR analysis showing the relative expression of *Sirt1* and *Lmod1* in siLmod1 or siSirt1 single knockdown or double knockdown compared to siCtrl treated cells, normalized to *Gapdh* expression levels. Paired t-test *: p-value ≤ 0.05 **: p-value ≤ 0.01. **C** and **D.** Percentage of cell populations found in immunofluorescence staining: reserve (Hoechst+/Myogenin-/MHC-) cells, Myogenin+/MHC- cells, Myogenin+/MHC+, fully differentiated myotubes and total counted cells under differentiating conditions after siRNA mediated knockdown of LMOD1 and SIRT1 (**C.** related to Figure 6C - E) or

siRNA mediated knockdown of LMOD1 and EX527 treatment to inhibit SIRT1 (**D.** related to Figure 6G - I.) **E.**
Normalized RNA-seq read counts of *Lmod1* and *Sirt1* after one day of differentiation after siCtrl and siLmod1 and
siCtrl and siSirt1 knockdown (Table S4). Paired t-test *: p-value ≤ 0.05 ***: p-value ≤ 0.001. For all bar plots, each
symbol represents a biological replicate, and the error bars indicate the SD. **F.** Working model: During the
proliferation stage, SIRT1 is found in the nucleus, repressing the expression of myogenic regulating factors.
However, at the onset of myogenic differentiation, LMOD1 interacts with SIRT1 and affects its subcellular
localization, leading to decreased levels of SIRT1 in the nucleus. This reduction in SIRT1 levels results in the de-
repression of MRFs and the initiation of differentiation.
572

Discussion

Myogenic differentiation involves the coordinated alterations of gene expression and reorganization of the cellular architecture. Here, we present a temporally resolved analysis of the proteome of primary mouse myoblasts undergoing differentiation *in vitro.* With a depth of more than 6000 proteins, our analysis covers three times more proteins compared to a previous study performed on an immortalized myoblast cell line (C2C12 cells) (Tannu et al. 2004; Kislinger et al. 2005). By using five independent primary myoblast lines, we were able to take into account the variability that typically affects primary cultures and derive robust and reproducible proteome signatures (Hindi et al. 2017; K. H. Kim, Qiu, and Kuang 2020). Our data highlight hundreds of changes in energy metabolism, RNA and protein synthesis as well as cytoskeletal organization that occur with specific temporal dynamics during myogenic differentiation. To demonstrate the relevance of our data, we decided to focus on LMOD1, a cytoskeletal protein whose role in myogenic differentiation has not yet been investigated. Although it is known that actin cytoskeletal dynamics are involved in myogenesis and that LMOD1 contributes to actin filament nucleation in smooth muscle cells (Guerin and Kramer 2009; Heng and Koh 2010; Xie et al. 2020; Chereau et al. 2008), the biological significance and functional role of LMOD1 during myogenic differentiation remained so far elusive. We found that LMOD1 is expressed by MuSCs, that its expression increases during myogenic differentiation *in vitro* and *in vivo*, and that reduction of LMOD1 levels significantly affects myotube formation and the number of myonuclei per myotube. Furthermore, the timing of induction of LMOD1 expression during skeletal muscle regeneration suggests a role in differentiation and fusion rather than in myofiber maturation. This is further supported by our finding that knockdown of *Lmod*1 after injury of skeletal muscle results in reduced regeneration. Consistently, our knockdown studies revealed that the reduction of LMOD1 severely affects the myogenic differentiation and myonuclear numbers. Cells that expressed Myogenin started differentiating into proper myotubes, indicated by the quantification of Myogenin+/MHC- and Myogenin+/MHC+ cells even after knockdown of LMOD1. However, a subset of cells positive for MHC but negative for Myogenin did not progress in differentiation, possibly due to loss or insufficient activation of myogenin expression. Loss of myogenin has been shown to lead to deregulated mTORC1 signaling and results in faster cell cycle entry in MuSCs (Ganassi et al. 2020). Since the knockdown of LMOD1 leads to an increase in the percentage of Myogenin-/MHC+ cells, the observed differentiation phenotype might be at least partially explained by a direct or indirect effect of LMOD1 on myogenin expression.

We found that LMOD1 selectively interacts with SIRT1. So far, there is no direct evidence suggesting that LMOD1 and SIRT1 have a specific binding domain where they can directly bind to each other. However, the comparison of primary sequences between LMOD1 and LMOD2 (Fowler and Dominguez 2017) reveals a 103 amino acid sequence unique for LMOD1,

which could represent a possible binding interface. It is also conceivable that other proteins or
molecules might be involved in mediating the interaction between LMOD1 and SIRT1, e.g.,
the RuvB-like AAA ATPase 1/2 (RUVBL1/RUVBL2). RUVBL1 participates in various
complexes, including INO80, NuA4, SWR1, TIP60-P400, PAQosome (R2TP), as a
transcriptional and/or chromatin modifier, with or without its homologue RUVBL2 (Gorynia et
al. 2011; P. B. Chen et al. 2015; Lakisic et al. 2016).
SIRT1 is a protein deacetylase that can localizes in the nucleus and cytoplasm (Bai and Zhang
2016; Moynihan et al. 2005; I.-Y. Chen et al. 2006), and was shown to play a critical role in
muscle differentiation and metabolism (Ryall et al. 2015; Diaz-Ruiz et al. 2015; Fulco et al.
2003, 2008). Myogenesis involves the shuttling of various proteins and myogenic regulators
between the nucleus and cytoplasm, regulated by various signals and interactions (Grifone et
al. 2021; Nguyen et al. 2022). We have observed that the localization of SIRT1 changes
dynamically during myogenic differentiation, and that the interaction between SIRT1 and
LMOD1 increases during the initial stages of myogenic differentiation. Additionally, we showed
that overexpression of LMOD1 affects the subcellular localization of SIRT1, leading to reduced
levels of SIRT1 in the nucleus. We hypothesize that relocation of SIRT1 to the cytoplasm,
mediated by LMOD1, is critical for the initiation of myogenic differentiation, resulting in the de-
repression of MRFs including myogenin. Alternatively, LMOD1 might sequester SIRT1 within
the cytoplasm, preventing its re-entry into the nucleus, which in turn results in the expression
of MRF to initiate myogenic differentiation (Figure S5F).
Previous studies have demonstrated that reducing the protein levels of SIRT1 can enhance
the differentiation of MuSCs (Ryall et al. 2015). Consistently, our data suggest that the
negative impact of knockdown of *Lmod1* on myogenic differentiation can be partially restored
by co-depletion of *Sirt1*. Our RNAseq analysis highlighted a subset of genes which were
oppositely up- or downregulated after knockdown of either LMOD1 or SIRT1, including
secreted factors and receptors that could potentially mediate this restoration of myogenic
differentiation when LMOD1 and SIRT1 levels are reduced. The identified factors include *Fgf6*,
Tgfa, *Camk1d*, *Mcr*, *Lifr*, *Isl2* and *Tgm2,* which were previously shown to be involved in
proliferation and differentiation of MuSCs (Armand, Laziz, and Chanoine 2006; Knight and
Kothary 2011; Yamane et al. 1998; Dos Santos et al. 2023; Hunt et al. 2013) by modulating
signaling pathways such as mTOR, Wnt or IGF1-PI3K/Akt-Foxo signaling (Wang et al. 2022;
Cui et al. 2020; Zhang et al. 2018). Further investigation of possible downstream effectors or
modulators identified by RNAseq could reveal additional mechanisms by which LMOD1
interferes with the myogenic program.
Finally, we have discovered that the abundance of LMOD1 is approximately four times higher
in MuSCs from geriatric mice compared to the abundance in MuSCs from young mice.
Specifically, LMOD1 tends to accumulate in the nucleus of MuSCs as mice age. Disruption of

the actin organization of the cytoskeleton and dynamics in aging and disease such as
Duchenne muscular dystrophy (DMD) or Nemaline myopathy (NM) were shown to impact the
expression of myogenic regulators and different signaling pathways (Lai and Wong 2020;
Eliazer et al. 2019; Y. J. Kim et al. 2022). Studies have shown that SIRT1 maintains H4K16 in
a deacetylated state in quiescent MuSCs, which leads to transcriptional repression (Ryall et
al. 2015). Given the SIRT1-LMOD1 interaction that we have discovered, we speculate that the
nuclear accumulation of LMOD1 might affect gene expression and chromatin remodeling in
aged MuSCs. However, it remains to be investigated whether the accumulation of LMOD1 is
a compensatory mechanism or is a causative factor leading to the dysregulation and
impairment of MuSCs during aging. Identifying the molecular mechanism and potential
downstream targets could be a promising strategy for discovering potential therapeutic targets
for promoting muscle regeneration and improving muscle health in older adults.

Acknowledgments

The authors gratefully acknowledge support from the FLI Core Facilities Proteomics, Sequencing, Imaging, and the Mouse Facility. A.O. is supported by the German Research Council (Deutsche Forschungsgemeinschaft, DFG) via the Research Training Group ProMoAge (GRK 2155), the Else Kröner Fresenius Stiftung (award number: 2019_A79), the Fritz-Thyssen Foundation (award number: 10.20.1.022MN), the Chan Zuckerberg Initiative Neurodegeneration Challenge Network (award numbers: 2020-221617, 2021-230967 and 2022-250618), and the NCL Stiftung. J.vM. received funding from the Deutsche Forschungsgemeinschaft (MA-3975/2-1), the Deutsche Krebshilfe and the Carl Zeiss Foundation (DKH-JvM-861005). S.C.S. was supported by a postdoctoral fellowship by the German Research Foundation (DFG, 505064275). The FLI is a member of the Leibniz Association and is financially supported by the Federal Government of Germany and the State of Thuringia.

Author contributions

Conceptualization: ES, SCS, JvM, AO
Data curation: ES, SCS, AO
Investigation: ES, IH, MH, AM, JvM
Methodology: ES, SCS, TD, JvM
Project administration: JvM, AO
Data analysis: ES, MH, AO
Supervision: JvM, AO
Visualization: ES, AO
Writing – original draft: ES, JvM, AO
Writing – review & editing: SCS

Declaration of interest

Authors declare no competing interests.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mice

All wild-type mice were C57BL/6J obtained from Janvier or internal breeding at the Leibniz Institute on Aging – Fritz Lipmann Institute (FLI) using the Janvier strain. All animals were kept in groups in a specific pathogen-free animal facility with a constant 12 h light/dark cycle and fed *ad libitum*. Young mice were aged 2-6 months, old mice were aged 18-20 months and geriatric animals were aged 24-33 months. All animal experiments were approved by the Thüringer Landesamt für Verbraucherschutz (Germany) under Reg.-Nr. FLI-17-014 and FLI-22-011.

Cardiotoxin muscle injury and siRNA injection of the TA muscle

Young mice (3-months old) were anaesthetized by inhaling isoflurane using the Tec 7 Isoflurane (798932, Covetrus) Anesthesia Vaporizer (Datex Ohmeda). The TA muscle was injured once by injecting 50 µl of 10 µM cardiotoxin (L8102, Latoxan) in sterile 0.9% NaCl (12391112, Medpex) using an insulin syringe with a 29 G needle. To manage pain, the mice were given Metacam (1 mg/kg, 798932, Covetrus) subcutaneously for three days.

Self-delivering Accell siRNA ON-TARGETplus Non-targeting Pool (D-001910-10-50, Dharmacon) or Accell siRNA ON-TARGETplus targeting *Lmod1* (E-051082-00-0050, Dharmacon) were injected into the CTX-damaged TA-muscles on day 3 after injury to induce knockdown *in vivo*. TA muscles were further analyzed by quantifying the number of myogenin+ cells per area.

FACS isolation of MuSCs

MuSCs were isolated from the hindlimb muscles of adult male mice of age 8 - 12 weeks. The muscles were dissected and collected in PBS, minced using scissors and digested in 2.5 g/mL collagenase B (Sigma) and 1 g/mL dispase (Sigma) for 30 min at 37 °C. Together with 12 mL of isolation medium (HAMs F10 + 20 % FBS) the digested muscle pieces were transferred into a 15 mL tube to allow the sedimentation of larger undigested muscle chunks. The sample was then filtered using a 0.74 mm cell strainer into a 50 mL tube, spun down at 450x *g* for 5 min at room temperature and resuspended in 500 mL isolation medium. The sample was then incubated on ice for 15 min with the antibodies used for FACS analysis, as indicated below. Afterwards, 15 mL PBS was added, and the sample was spun down for 5 min at 450x *g* at room temperature followed by resuspension in 1mL PBS. After filtering the cells with a 0.35 mm cell strainer staining with SYTOX Blue Dead Cell Stain (S34857, Thermo Fisher) was performed to distinguish between dead and alive cells. A BD FACSAria III was used for sorting the cells. To FAC sort MuSCs, gating was set to alpha7-integrin+ (anti-Alpha 7 Integrin 647

(clone:R2F2), 67-0010-05, The University Of British Columbia AbLab, 1:500), CD11b-(CD11b-
PE, 553311, BD Bioscience, 1:500), CD31- (CD31-PE, 553373, BD Bioscience,1:500), CD45-
(CD45-PE, 553081, BD Bioscience,1:500) and Sca1- (Ly-6A/E-PE, 553108, BD Bioscience,
1:500).

Cell culture
FACS isolated primary myoblasts were cultured on collagen (#354236, Corning) coated (1h
at RT with 0.167 mg/mL) plates in Ham's F-10 Nutrient Mix (Thermo Fisher #31550031)
supplemented with 10 % FBS (#10270106, Thermo Fisher), 1 % Penicillin-Streptomycin
(#15140-122, Thermo Fisher) and 2.5 ng/mL bFGF (#13256029, Thermo Fisher) in a 37 °C
incubator with 95 % humidity and 5 % CO_2. For the proliferation assay, primary myoblasts
were seeded onto collagen-coated 24-well plates (734-2325, VWR) (25,000 cells per well) and
cultured for 48 h in the myoblast growth medium. For differentiation assays on 24-well plates,
75,000 primary myoblasts were seeded and cultivated for 72 h in the differentiation medium.
Primary myoblasts were always harvested using 0.25 % Trypsin. For location stainings, cells
were grown on collagen-coated 4-well culture slides (80426, Ibidi). For proliferation conditions,
5,000 cells were seeded per well, and 12,000 cells were seeded per well for differentiation
conditions. The cells were then incubated at 37 °C.
HEK293T cells (Flp-In 293 T-Rex cells) were obtained from Thermo Fisher (R78007).
HEK293T cells were grown in Dulbecco's modified Eagle's medium (#D6429, Sigma) with high
glucose (5 g/l) supplemented with 10 % heat-inactivated FBS and supplementation with 100
mg/mL Zeocin (#R250-01, Thermo Fisher) and 15 mg/mL Blasticidin (#R210-01, Thermo
Fisher). After generation of a stable line cells were supplemented with 100 mg/mL Hygromycin
(#10687-010, Thermo Fisher) and 15 mg/mL Blasticidin. Cells were cultured in a 37 °C
incubator with 95 % humidity and 5 % CO_2. For immunofluorescence staining of HEK293T,
cells were grown on 50 µg/mL Poly-D-Lysine solution (Gibco™ A3890401) coated autoclaved
coverslips (Carl Roth, YX03.1). Each coverslip was placed in an individual well of a 12-well
Lab solute plate (with a specified number), and 25,000 cells were seeded per well.
The Platinum-E (Plat-E) cells, a potent retrovirus packaging cell line were cultured in DMEM
high glucose (#D6429, Sigma) supplemented with 10 % FBS, 1 %Penicillin-Streptomycin in a
37 °C incubator with 95 % humidity and 5 % CO_2.

siRNA transfection of cells with Lipofectamine® RNAiMAX Reagent
Knockdown experiments using small interfering RNAs (siRNA) (ON-TARGETplus Non-
targeting Pool (D-001810-10-05, Dharmacon), SMART:pool ONTARGETplus Lmod1 siRNA
(9530015K06Rik | SM-Lmod), SMART:pool ONTARGETplus Lmod2 siRNA (C-Lmod),
SMART:pool ONTARGETplus Sirt1 siRNA (AA673258 | SIR2L1 | Sir2 | Sir2a | Sir2alpha)

against a target mRNA were performed. For siRNA transfection, the Lipofectamine®
RNAiMAX Kit (13778075, ThermoFisher) was used according to the manufacturer's protocol.
777

Generation of overexpression cell lines

The murine stem cell virus (MSCV) retroviral expression system (634401, Takarabio) was
used for overexpression studies. The MSCV vectors are optimized for introducing and
expressing target genes in stem cells such as MuSCs. The vector pENTR233.1-Lmod1
(100016603, Dharmacon, spectinomycin resistant) and GFP were blunt end cloned in the
donor vector cloneJET^{TM} (K1232, Thermo Fisher, ampicillin-resistant) to finally generate
pMSCV.puro plasmids (634401, Takarabio, ampicillin-resistant) encoding Lmod1 and GFP,
respectively (Table 1).

Plat-E cells were transfected with 20 µg of the generated pMSCV.puro plasmids using
polyethylenimine (PEI). 60 uL PEI (1 mg/mL) were added to the plasmid mixture (3:1 ratio of
PEI:DNA), briefly mixed by inverting the tube and incubated for 20 min at RT. Subsequently,
the transfection complex was added dropwise to a 10 cm plate and incubated for 5-6 h in 3.5
% CO2, 37 °C. The medium was changed to standard proliferation medium (DMEM, 10 %
FBS and 1 % Pen-Strep) and incubated until the next day to collect the retrovirus. For further
infection of primary myoblasts, 1 Mio cells were seeded in a p10 plate and infected using a
1:5 ratio of retrovirus to growth medium (F10, 20 % FBS, bFGF, 1 % Pen-Strep) mixed with 5
µl of Protamine Sulfate (8 µg/mL). Infection of primary myoblasts was repeated the next day
as previously described and incubated at 37 °C. After 48 h post-transfection, primary
myoblasts were selected with 1.25 µg/mL puromycin for up to one week. Medium was changed
every 2 - 3 days.

In vitro **proximity labeling for BioID protocol**

HEK293T cells (Flp-In 293 T-Rex cells) expressing BirA*-LMOD1 and BirA*-LMOD2, were
generated as described by (Mackmull et al. 2017). The cells were selected using 15 µg/mL
Blasticidin (GibcoTM, R21001) and 100 µg/mL Zeocin™ (Thermo Fisher Scientific, R25001).
Once stable cell lines were generated, the cells were selected using 100 µg/mL Hygromycin B
(Thermo Fisher Scientific, 10687010) and 15 µg/mL Blasticidin (GibcoTM, R21001).

Stable HEK293T cell lines expressing fusion proteins containing BirA* were seeded in 150
mm dishes at the density of approximately 500K cells. To induce expression of BirA* fusion
proteins, cells were treated with 1 µg/mL tetracycline (Sigma Aldrich, 87128) for four days.
Biotin (50 µM) (Sigma Aldrich, B4501) was added to the cells 24 h prior to harvest to biotinylate
proximal proteins.

Table 1: Primers for cloning

Primer name	Sequence (5'-3')
Lmod1 sequencing primer (forward)	CTT GTG GCC GTT TAC GTC
Lmod1 sequencing primer (reverse)	CCCACTTGCTTGCTTTCATC
GFP sequencing primer (forward)	TAATACGACTCACTATAGGG
GFP sequencing primer (forward)	CGACTCACTATAGGGAGAGCGGC
pMSCV.puro sequencing primer (forward)	CGAGACCTCATCACCCAGG
T7 (forward)	TAATACGACTCACTATAGGG
pJet1.2 sequencing primer (forward)	CGACTCACTATAGGGAGAGCGGC
Lmod1_pcr_(forward)	CACCATGTCCAAAGTAGCTAAGTACCG
Lmod1.wSTOP (reverse)	TTATTGAAGTAGCTTGGGCACC
Lmod1.woSTOP (reverse)	TTGAAGTAGCTTGGGCACC
Lmod2.pcr (forward)	CACCATGTCTACGTTTGGCTACAGAAG
Lmod2.woSTOP (reverse)	TCTCAGAGCTTCGGGAACTTC
Lmod2.wSTOP (reverse)	TTATCTCAGAGCTTCGGGAACTTC

Skeletal muscle cryostat sections and staining

TA muscles were isolated and frozen sections (14 µm) were made using a Leica Cryostat CM

3050 (Meyer Instruments) at -21 °C and placed on glass slides. Sections were stored at -80

°C. For immunostaining, muscle cryosections were fixed with 4 % paraformaldehyde (v/v,

CP10.1, Roth, pH 7.4). for 7 min and then permeabilized with 0.1 M Glycine (1042011000,

VWR) in PBS, (pH 7.4) for 5 min and after one wash with PBS 0.5 %Triton-X was added (v/v,

3051.3, Roth) for 10 min. After another washing step with PBS sections were incubated for 20

min at 65 °C in pre-heated Citrate Antigen Retrieval buffer (10 mM Sodium citrate, 0.05 %

Tween 20, pH 6.0). Slides were re-equilibrated by washing 3x with PBS for 5 min and

incubated for 30 min using MOM blocking (1:40) and subsequently blocked in blocking solution

(5 % horse serum (v/v) in PBS) at RT. The sections were incubated in primary antibody Pax7

(Pax7, DSHB, Hybridoma mouse IgG1, undiluted), myogenin (DSHB, F5D, hybridoma cell

supernatant, undiluted), Lmod1 (Proteintech, 15117-1-AP, 1:500) and Laminin (LSBio, LS-

C96142, 1:500) overnight at 4 °C in a wet chamber. Sections were washed 3x with 0.1 %

Triton in PBS (pH 7.4) for 5 min and secondary antibodies anti-chicken IgG (Alexa Fluor® 488,

A-11039), anti-mouse IgG (Alexa Fluor® 546, A-21123, Thermo Fisher, 1:1000) and anti-

rabbit IgG (Alexa Fluor® 647, A-31573, Thermo Fisher, 1:1000), were applied for 1 h at RT.

Nuclei were stained with Hoechst (bisBenzimideH 33258, B2261 Sigma, 0.02 µg/µl in PBS,

1:5000) for 5 min. After washing sections 3x with 0.1 % Triton in PBS, sections were mounted

in Permafluor mountain medium (TA-006-FM, Thermo Fisher). Immunofluorescence

microscopy was performed with the Zeiss Axio Scan Z.1 using a Plan-Apochromat 20x/0.8

M27 Objective (Zeiss).

Immunofluorescence staining of primary myoblasts

Primary myoblasts from proliferation or differentiation assays were fixed with 2 % formaldehyde (v/v in PBS, CP10.1, Roth) for 5 min at RT, washed twice with PBS for 5 min and were permeabilized with permeabilization solution (0.1 % Triton-X-100 (v/v, 3051.3, Roth), 0.1 M Glycine (1042011000, VWR) in PBS) for 5 min at RT. Cells were incubated for 1 h in blocking solution (5 % horse serum, v/v, 26050-088, Thermo Fisher) in PBS at RT. Incubation with primary antibodies was carried out in 5 % horse serum in PBS overnight at 4 °C. Subsequently, the cells were washed three times with PBS for 5 min and incubated with the secondary antibodies in blocking solution for 1 h at RT in the dark. Next, the cells were washed once with PBS for 5 min, stained with Hoechst (bisBenzimide H 33258, B2261 Sigma, 1:5000 in PBS) at RT for 5 min and washed twice with PBS for 5 min. The cells were stored at 4 °C in PBS in the dark until they were analyzed. For location stainings, the removable chamber was detached, and cells were mounted in permafluor mounting medium (TA-006-FM, Thermo Fisher) and coverslips (631-1574, VWR). The cells were then analyzed using an Axio Observer microscope (Carl Zeiss) or with an Axio Imager (Z2 using a Plan-Apochromat 63 x / 0.8 M27 Objective) and analyzed with Zen 2 Blue Edition software (Carl Zeiss Microscopy GmbH).

Primary antibodies used for immunofluorescence analysis (Table 3): Hybridoma mouse IgG1 myogenin (DSHB, F5D, hybridoma cell supernatant, undiluted), Hybridoma mouse IgG1 devMHC (DSHB, MF20, hybridoma cell supernatant, undiluted), anti-ki67 (ab15580, rabbit IgG 1:500), Lmod1 (Proteintech, 15117-1-A, rabbit, IgG, 1:500) and Sirt1 (Proteintech, 60303-1-Ig, mouse, IgG2b, 1:400). Secondary antibodies: anti-mouse IgG2b (Alexa Fluor® 488, A-21141, Thermo Fisher, 1:1000), anti-rabbit IgG (Alexa Fluor® 546, A10040, Thermo Fisher, 1:1000), anti-mouse IgG (Alexa Fluor® 546, A-21123, Thermo Fisher, 1:1000)

Immunofluorescence staining of HEK293T cells

The cells were washed 3x with 1 x PBS and fixed in 4 % formaldehyde (v/v) in PBS for 10 minutes at room temperature. After that, cells were washed again 3x with 1 x PBS and permeabilized using a permeabilization buffer (0.7 % Triton X-100 in 1 x PBS) for 15 minutes at room temperature, followed by two washing steps with PBS for 5 minutes each. The samples were incubated with a blocking solution (10 % (w/v) BSA, 10 % (v/v) Triton X-100, 5 % (v/v) goat serum) for 1h at room temperature. The coverslips were incubated with primary antibody anti-FLAG (Sigma-Aldrich, mouse, F1804, 1:500) at 4 °C overnight. After washing 3x with PBS, the secondary fluorescence-labeled antibody anti-mouse IgG (Alexa Fluor® 488, A-21121, Thermo Fisher, 1:1000) in blocking solution and anti-biotin (Streptavidin AlexaFuor® 568, S11226, Thermo Fisher, 1:2000) in blocking solution were incubated for 1 h at RT. After washing once with PBS nuclei were stained with Hoechst (bisBenzimideH 33258, B2261

Sigma, 0.02 µg/µl in PBS, 1:5000) at RT for 5 minutes. and were finally washed twice for 5

minutes with PBS. Finally, the samples were washed twice with PBS for 5 minutes, and cells

were mounted in Permafluor mounting medium (TA-006-FM, Thermo Fisher) and coverslips

(631-1574, VWR). Immunofluorescence microscopy was performed with an Axio Imager (Z2

using a Plan-Apochromat 63 x / 0.8 M27 Objective) and analyzed with Zen 2 Blue Edition

software (Carl Zeiss Microscopy GmbH).

Proximity ligation assay

Primary myoblasts were cultured in a collagen-coated removable 12-well chamber (81201,

Ibid) and fixed in 2 % PFA for 5 minutes at RT. Subsequently, the proximity ligation assay

(PLA) (#DUO92008, Sigma-Aldrich) was performed according to the manufacturer's protocol.

Anti-Sirt1 antibodies and anti-Lmod1 antibodies were incubated overnight at 4 °C.

Immunofluorescent pictures were taken with an Axio Imager (Z2 using a Plan-Apochromat 63x

/ 0.8 M27 Objective) and analyzed with Zen 2 Blue Edition software (Carl Zeiss Microscopy

GmbH).

Harvesting and lysis of cells for immunoblotting

To harvest the cultured cells from proliferation or differentiation assay, the medium was

removed and the cells were washed with PBS at RT. Then, 50 µl RIPA buffer (150 mM Sodium

Chloride (P029.2, Roth), 1 % Triton X-100 (v/v, 3051.3, Roth), 0.5 % Sodium Deoxycholate

(89904, Thermo Fisher), 0.1 % SDS (w/v, 75746-250G, Sigma), 50 mM Tris (4855.2, Roth),

pH8) was added to each well of the 24 well plate. The cells were carefully scraped off the plate

by using a cell scraper. Cells were incubated on ice for 20 min and mixed by vortexing every

5 min. Cell lysis was completed by sonication using Diagenode's Bioruptor® Plus 10x for 1

min (1 min 'on', 30 sec 'off').

Primary antibodies used for immunoblotting analysis: anti-BirA (Novus Biologicals, 5B11c3-3,

1:500), anti-GFP (Santa Cruz, sc-9996, 1:1000), anti-mouse (Agilent Dako, P0448, 1:1500)

anti-Lmod1 (Proteintech, 15117-1-AP, 1:500), anti-Lmod2 (Atlas Antibodies, HPA051039,

1:500), anti-Sirt1 (Proteintech, 60303-1-Ig, 1:1000), anti-Aldoa (Proteintech, 67453-1-Ig,

1:20.000), anti-H4 (Cell Signaling, #2935, 1:1000), anti-Gapdh (SantaCruz, sc-365062,

1:200). Secondary antibodies: anti-rabbit (Agilent Dako, P0447, 1:2000), anti-mouse (Agilent

Dako, P0448, 1:1500).

Table 3: Antibody-List

Antibody name	Source	Product number
Monoclonal rat IgG2a Sca1-FITC	eBioscience	11-5981-85RRID:AB_465334
Monoclonal rat IgG2a Sca1-PE (Ly-6A/E-PE)	BDBioscience	553108RRID:AB_394629
Hybridoma mouse IgG1 Pax7	DSHB	Pax7RRID:AB_528428
Polyclonal rabbit IgG Lmod1	Proteintech	15117-1-AP
Polyclonal human IgG Lmod2	Atlas Antibodies	HPA051039
Polyclonal chicken IgG Laminin	LSBio	LS-C96142
Hybridoma mouse IgG1 myogenin	DSHB	F5D
Hybridoma mouse IgG1 devMHC	DSHB	MF20
Polyclonal rabbit IgG Ki-67	Abcam	ab15580
Monoclonal mouse IgG2b Sirt1	Proteintech	60303-1-Ig
bisBenzimideH 33258 (Hoechst)	Sigma-Aldrich	B2261
Monoclonal mouse M2 Flag	Sigma-Aldrich	F1804
Monoclonal mouse IgG GFP	Santa Cruz	Sc-9996
Monoclonal mouse IgG BirA	Novus Biologicals	5B11c3-3
Monoclonal mouse IgG1 Gapdh	SantaCruz	sc-365062
Monoclonal mouse IgG H4 (L64C1)	Cell Signaling	#2935
Monoclonal mouse IgG Aldoa	Proteintech	67453-1-Ig
anti-mouse IgG Alexa Fluor® 488	Thermo Fisher	A-21121
anti-mouse IgG Alexa Fluor® 546	Thermo Fisher	A-21123
anti-mouse IgG2b Alexa Fluor® 488	Thermo Fisher	A-21141
anti-mouse IgG2b Alexa Fluor® 488	Thermo Fisher	A-21242
anti-rabbit IgG Alexa Fluor® 488	Thermo Fisher	A-21206
anti-rabbit IgG Alexa Fluor® 546	Thermo Fisher	A10040
anti-rabbit IgG Alexa Fluor® 647	Thermo Fisher	A-31573
anti-chicken IgG Alexa Fluor® 488	Thermo Fisher	A-11039
Anti-Streptavidin Alexa Fluor ® 568	Thermo Fisher	S11226
Anti-Mouse Immunoglobulins/HRP	Agilent Dako	P0448
Anti-Rabbit Immunoglobulins/HRP	Agilent Dako	P0447

Protein quantification assays

The EZQ™ Protein Quantitation Kit (R33200, Thermo Fisher) and the Invitrogen™ Qubit™ 3

Fluorometer (15387293, Invitrogen), together with Qubit™ assay tubes (Q32856, Invitrogen),

were used to quantify protein samples according to the manufacturer's protocols. The TECAN

microplate reader (1405005206, Tecan INFINITE M1000 PRO, excitation 485 nm, emission

580 nm) was used to analyze the EZQ assay.

GFP Trap

For the generation of cell lines transiently expressing Sirt1 fused to GFP or GFP control, plasmids were generated using Gateway Technology (Invitrogen). Four 10 cm dishes of HEK293T cells (3 million cells/dish) were used for each candidate, and prior to transfection, the medium was replaced with a transfection medium (DMEM with 1 % FBS, without antibiotics). Each plate was transfected with 5 µg plasmid DNA and 15 µg Polyethylenimine (PEI 25K™, Polysciences, 23966-100), both previously prepared in OptiMEM (Gibco, 11520386). The transfection medium was changed to 5 mL DMEM with 10 % FBS and 1 % Pen-Strep 6 h post-transfection. Additionally, 5 µl (1 mg/mL) of Tetracycline was to the media and cells were incubated for 48 h at 3.5 % CO_2, 37 °C. Subsequently, cells were harvested by trypsinization, and 20 million cells per pellet were collected for GFP-Trap pull-down. Each pellet was lysed in 200 µl of lysis buffer (20 mM Tris pH 7.5, 150 mM NaCl, 1 mM EDTA, 0.3 % Triton, 10 % Glycerol) containing protease and phosphatase inhibitors. Tubes were placed on ice for 30 min with vortexing every 10 min, followed by brief sonication in a Bioruptor Plus for 5 cycles (30 sec on/30 sec off) at the highest setting and afterwards centrifuged at 20000x *g*, for 10 min at 4 °C. Lysate-supernatants were transferred to a pre-cooled tube and to each of them 300 µl dilution buffer (10 mM Tris pH 7.5, 150 mM NaCl, 0.5 mM EDTA), was added. 25 µl of GFP Trap beads (ChromoTek GFP-Trap® Agarose, Proteintech, gta) were equilibrated in 0.5 mL of ice-cold dilution buffer and were spun down at 100x *g* for 5-10 sec at 4 °C. Beads were washed two more times with 500 µl dilution buffer. Cell lysates were added to equilibrated GFP Trap beads and incubated for 1 h at 4 °C with constant mixing. Tubes were spun at 100x *g* for 5-10 sec at 4 °C. GFP Trap beads were washed with 500 µl ice-cold dilution buffer, followed by two washes with wash buffer (10 mM Tris pH 7.5, 150 mM NaCl, 0.05 % P40 Substitute, 0.5 mM EDTA). To elute proteins, 2 x 30 µl of 2x SDS-sample buffer (120 mM Tris pH 6.8, 20 % glycerol, 4 % SDS, 0.04 % bromophenol blue) was added to the GFP Trap beads and boiled for 5 min at 95 °C. Subsequently, the beads were collected by centrifugation at 2500x *g* for 2 min and supernatant was collected to perform an SDS-PAGE.

Nuclear cytoplasmic fractionation experiment

Nuclear cytoplasmic fractionation was performed seeding 3 x 10^6 primary myoblasts. Prior cell harvest, cells were washed twice with 10 mL of PBS at RT to remove any residual media. Then, they were trypsinized with 0.25 % Trypsin and resuspended in 8 mL of myoblasts growth medium. After centrifuging at 500x *g* for 5 minutes, the supernatant was discarded, and the cells were resuspended in 5 mL of ice-cold PBS. At this stage, the total number of cells was counted. For hypotonic lysis, the cells were spun at 4 °C and 500x *g*. The resulting pellet was resuspended in 500-700 µl of Buffer A (10 mM Tris pH 7.5) containing protease inhibitor. The cells were incubated on ice for 20-30 minutes, and their swelling was monitored. Then, they were lysed by passing them through a 27G needle 15-30 times. The process was stopped when 70-80 % of the cells were lysed to prevent nuclei breakdown. Nuclei were manually counted to calculate the ratio of nuclei to total cells. The lysed cells were centrifuged at 1000x *g* at 4 °C to separate nuclei, and the supernatant was collected as the cytoplasmic fraction. The pellet of nuclei was resuspended in 60-100 µl of Buffer B (0.25 M sucrose, 50 mM Tris pH 7.5, 25 mM Kcl, 5 mM MgCl2, 2 mM DTT) with protease inhibitor and the nuclei were manually counted using a hemocytometer.

RNA isolation

RNA isolation was performed with TriFast (Peqlab) according to the manufacturer's protocol. Primary myoblasts were either directly harvested into TriFast or cell pellets were resuspended and lysed by pipetting.

cDNA synthesis

cDNA was synthesized by reverse transcription using 600 ng of isolated RNA. The GoScript™ Reverse Transcriptase kit (A5001, Promega) was used according to the manufacturer's protocol. Finally, the cDNA was diluted 1:10 and stored at -20 °C until further analysis.

qRT-PCR

Quantitative real-time PCR was performed on a CFX384 Touch Real-Time PCR Detection System (Biorad). One qRT-PCR reaction mix consists of 3 µl cDNA, 5 µl iQ SYBR™ Green Supermix (1708882, Biorad), 1 µl sterile ddH2O and 0.5 µl of 10 µM forward and reverse primer (Metabion, ordered as 100 µM stock in sterile ddH2O, sequences listed in Table 2). Technical triplicates were performed for each sample. For triplicates having a Cq-SEM (calculated by Bio-Rad CFX Maestro 1.1) higher than 0.4, the replicate with the largest difference was excluded. The obtained Ct-values of target genes and housekeeping genes were used to calculate relative expression according to the $2^{-\Delta\Delta Ct}$ formula (Pfaffl 2001).

975 **Table 2: Primers for qRT-PCR**

Primer name	Sequence (5'-3')
GAPDH (forward)	ATGCCAGTGAGCTTCCCGTC
GAPDH (reverse)	CATCACCATCTTCCAGGAGC
Lmod1 (forward)	AAAAGACAGGAGTGTCAAGAAC
Lmod1 (reverse)	CCCCAGAACCTATGCCCTC
Lmod2 (forward)	ACCTTATCCCGATTTGCTGAAG
Lmod2 (reverse)	ACCTTGAGCATGTCTGCAATG
MyoD (forward)	CTACAGTGGCGACTCAGAT
MyoD (reverse)	CACTGTAGTAGGCGGTGTC
Myogenin (forward)	CAGTACATTGAGCGCCTAC
Myogenin (reverse)	AAGGCAACAGACATATCCTC
Pax7 (forward)	TCTTACTGCCCACCCACCTA
Pax7 (reverse)	CACGTTTTTGGCCAGGTAAT
Sirt1 (forward)	TGATTGGCACCGATCCTCG
Sirt1 (reverse)	CCACAGCGTCATATCATCCAG
18S (forward)	CCATCCAATCGGTAGTAGCG
18S (reverse)	GTAACCCGTTGAACCCCATT

RNA Sequencing

Primary myoblasts were seeded and pre-treated with siRNA against Lmod1 and Sirt1 for two
days under proliferating conditions. A second transfection with siLmod1 and siSirt1 was
performed simultaneously with the induction of differentiation. After one day of differentiation,
the cells were collected in TriFast and RNA was isolated using the procedure described above.
The concentration and integrity of the isolated RNA was determined using Qubit 3.0
Fluorometer (Thermo Fisher), Bioanalyzer 2100, and Tapestation RNA (Agilent). To prepare
the poly-A library, TruSeq RNA Library Prep Kit v2 (Illumina, RS-122-2001) was used
according to the manufacturer's protocol. The process started with the enrichment of poly-A-
containing mRNA using magnetic beads, followed by first-strand synthesis of cDNA using
SuperScript III Reverse Transcriptase (Thermo Fisher). The second strand synthesis was
performed using the second strand master mix (TruSeq RNA Library Prep Kit v2 (Illumina)),
and the DNA was cleaned up using Agencourt AMPure XP Beads (Beckman Coulter,
A63881). After end repair and dA-tailing, adaptors were ligated, and the library was further
enriched using the PCR Master Mix (TruSeq RNA Library Prep Kit v2 (Illumina)). The
concentration and quality of the library were determined using Qubit 3.0 Fluorometer (Thermo
Fisher), Bioanalyzer 2100, and Tapestation DNA (Agilent). Finally, the libraries were pooled
and RNA-Seq and data analysis was performed by Azenta US using Illumina NovaSeq (USA;
www.GENEWIZ.com).

Whole proteome analysis

For proteomics analysis of myoblasts or Lmod1 overexpression cell lines, samples were sonicated (Bioruptor Plus, Diagenode, Belgium) for 10 cycles (60 sec ON/30 sec OFF) at high setting, at 20 °C, followed by boiling at 95 °C for 5 min. Reduction with 10 mM Dithiothreitol (DTT) (6908.3, Roth) for 15 min was followed by alkylation with iodoacetamide (IAA, final concentration 15 mM) for 30 min at room temperature in the dark. Protein amounts were estimated following an SDS-PAGE gel of 10 µL of each sample against an in-house cell lysate of known quantity. 30 µg of each sample was used for digestion. Proteins were precipitated overnight at -20 °C after addition of 4x volume of ice-cold acetone. The following day, the samples were centrifuged at 20800x g for 30 min at 4 °C and the supernatant was carefully removed. Pellets were washed twice with 300 µL ice-cold 80 % (v/v) acetone in water and then centrifuged at 20800x g at 4 °C for 10 min. After removing the acetone, pellets were air-dried before addition of 25 µL of digestion buffer (1M Guanidine, 100 mM HEPES, pH 8). Samples were resuspended with sonication as explained above, then LysC (Wako) was added at 1:100 (w/w) enzyme:protein ratio and digestion proceeded for 4 h at 37 °C under shaking (1000 rpm for 1 h, then 650 rpm). Samples were then diluted 1:1 with MilliQ water and trypsin (Promega) added at at 1:100 (w/w) enzyme:protein ratio. Samples were further digested overnight at 37 °C under shaking (650 rpm). The day after, digests were acidified by the addition of TFA to a final concentration of 10 % (v/v), heated at 37 °C and then desalted with Waters Oasis® HLB µElution Plate 30 µm (Waters Corporation, MA, USA) under a soft vacuum following the manufacturer instruction. Briefly, the columns were conditioned with 3x100 µL solvent B (80 % (v/v) acetonitrile; 0.05 % (v/v) formic acid) and equilibrated with 3x100 µL solvent A (0.05 % (v/v) formic acid in Milli-Q water). The samples were loaded, washed 3 times with 100 µL solvent A, and then eluted into 0.2 mL PCR tubes with 50 µL solvent B. The eluates were dried down using a speed vacuum centrifuge (Eppendorf Concentrator Plus, Eppendorf AG, Germany). Dried samples were stored at -20 °C until analysis.

Enrichment and digestion of biotinylated proteins (BioID)

The protocol was used as described in Bartolome et al (Bartolome et al. 2023). In short, pellets corresponding to 20 Mio cells were resuspended in 4.75 mL lysis buffer (50 mM Tris pH 7.5; 150 mM NaCl; 1 mM EDTA; 1mM EGTA; 1 % (v/v) Triton X-100; 1 mg/ml aprotinin; 0.5 mg/ml leupeptin; 250 U turbonuclease; 0.1 % (w/v) SDS) and incubated for 1 h at 4 °C. Streptavidin Sepharose High Performance (GE Healthcare) was acetylated by the addition of 10 mM sulfo-NHS acetate (Thermo Fisher Scientific) for 30 min twice and then equilibrated in lysis buffer. For each lysate 80 µl of equilibrated beads were used. Upon loading, beads were washed 5 times with 600 µl 50 mM AmBic. On-bead digest was performed with 200 µl LysC (5 ng/µl) at

37 °C overnight. First elution step was achieved using two times 150 μl of 50 mM AmBic. After

pooling both fractions, peptides were further digested off-beads by adding 1 μg trypsin and

incubating at 37 °C for 3 h. Biotinylated peptides were eluted in a second elution step using

two times 150 μl 20 % TFA (Biosolve) in acetonitrile (Biosolve). Both fractions were pooled

and neutralized to pH 8.0 by adding 50 μl 200 mM HEPES and sodium hydroxide as

necessary. Peptides were digested further off-beads by adding 1 μg trypsin at 37 °C for 3 h.

Digested peptides from the two elution steps were desalted using Waters Oasis® HLB

μElution Plate 30 μm (Waters) according to the manufacturer's instructions.

1041

LC-MS/MS analysis

Peptides were reconstituted in 10 μl reconstitution buffer at a concentration of 1 μg/μl, 0.5 μl

of the HRM kit (Ki-3002-1, Biognosys) was added in a dilution recommended by the

manufacturer and 1 μl were injected for measurement. Peptides were separated using the

UltiMate 3000 UPLC system (Thermo Fisher Scientific, for myoblasts) or nanoAcquity UPLC

system (Waters, for BioID and Lmod1 overexpressing cell lines) fitted with trapping

(nanoAcquity Symmetry C18, 5μm, 180 μm x 20 mm) and an analytical column (nanoAcquity

BEH C18, 1.7μm, 75μm x 250mm). The outlet of the analytical column was coupled directly

to Q-exactive HF (Thermo Fisher Scientific, for myoblasts) or Orbitrap Exploris 480 (Thermo

Fisher Scientific, for BioID and Lmod1 overexpressing cell lines) using the Proxeon nanospray

source. For myoblast experiment, solvent A was water, 0.1 % FA and solvent B was 80 %

(v/v) acetonitrile, 0.08 % FA. Peptides were eluted via a non-linear gradient from 1 % to 62.5

% B in 131 min. Total runtime was 150 min, including clean-up and column re-equilibration.

The peptides were introduced into the mass spectrometer via a Pico-Tip Emitter 360 μm OD

x 20 μm ID; 10 μm tip (New Objective) and a spray voltage of 2.2 kV was applied. The RF ion

funnel was set to 60 %.

Data Dependent Acquisition (DDA) settings were as follows: Full scan MS spectra with mass

range 350-1650m/z were acquired in the Orbitrap with a resolution of 60,000 FWHM. The

filling time was set at a maximum of 20ms with an AGC target of $3x10^6$ ions. A Top15 method

was employed to select precursor ions from the full scan MS for fragmentation (minimum AGC

target of $1x 10^3$ ions, normalized collision energy of 27 %), quadrupole isolation (1.6 m/z) and

measurement in the Orbitrap (resolution 15,000 FWHM, fixed first mass 120 m/z). The

fragmentation was performed after the accumulation of $2x10^5$ ions or after filling time of 25ms

for each precursor ion (whichever occurred first). Only multiply charged (2+ -7+) precursor

ions were selected for MS/MS. Dynamic exclusion was employed with a maximum retention

period of 20s. Isotopes were excluded.

For Data Independent Acquisition (DIA), 1 μg of reconstituted peptides were loaded using the

same setup and LC conditions used for DDA. MS conditions were modified as follows: Full

scan MS spectra with mass range 350-1650 m/z were acquired in profile mode in the Orbitrap with resolution of 120,000 FWHM. The filling time was set at a maximum of 60 ms with an AGC target of 3×10^6 ions. DIA scans were acquired with 40 mass window segments of differing widths across the MS1 mass range. The default charge state was set to 3+. HCD fragmentation (stepped normalized collision energy; 25.5, 27, 30 %) was applied and MS/MS spectra were acquired with a resolution of 30,000 FWHM with a fixed first mass of 200 m/z after accumulation of 3×10^6 ions or after a filling time of 35ms (whichever occurred first). Data was acquired in profile mode. For data acquisition and processing Tune version 2.9 Q-Exactive HF

For BioID project experiment and, solvent A was water, 0.1 % FA and solvent B was 80 % (v/v) acetonitrile, 0.08 % FA. Peptides were eluted via a non-linear gradient from 3 % to 40 % B in 90 min. Total runtime was 115 min, including clean-up and column re-equilibration. The peptides were introduced into the mass spectrometer via a Pico-Tip Emitter 360 µm OD x 20 µm ID; 10 µm tip (New Objective) and a spray voltage of 2.2 kV was applied. The RF ion funnel was set to 30 %. For Data Independent Acquisition (DIA), MS conditions were modified as follows: Full scan MS spectra with mass range 350-1650 m/z were acquired in profile mode in the Orbitrap with resolution of 120,000 FWHM. The filling time was set at a maximum of 60 ms with an AGC target of 3×10^6 ions (300 %). DIA scans were acquired with 34 mass window segments of differing widths across the MS1 mass range. The default charge state was set to 2+. HCD fragmentation (stepped normalized collision energy; 25.5, 27, 30 %) was applied and MS/MS spectra were acquired with a resolution of 30,000 FWHM with a fixed first mass of 200 m/z after accumulation of 3×10^6 ions (3000 %) or after a filling time of 40ms (whichever occurred first). Data was acquired in profile mode. For data acquisition and processing Tune 3.1.

For Lmod1 overexpressing cell line experiment, solvent A was water, 0.1 % FA and solvent B was acetonitrile, 0.1 % FA. Peptides were eluted via a non-linear gradient from 3 % to 40 % B in 120 min. Total runtime was 145 min, including clean-up and column re-equilibration. The peptides were introduced into the mass spectrometer via a Pico-Tip Emitter 360 µm OD x 20 µm ID; 10 µm tip (New Objective) and a spray voltage of 2.2 kV was applied. The RF ion funnel was set to 30 %. For Data Independent Acquisition (DIA), 1 µg of reconstituted peptides were loaded using the same setup and LC conditions used for DDA. MS conditions were modified as follows: Full scan MS spectra with mass range 350-1650 m/z were acquired in profile mode in the Orbitrap with resolution of 120,000 FWHM. The filling time was set at a maximum of 60 ms with an AGC target of 3×10^6 ions. DIA scans were acquired with 40 mass window segments of differing widths across the MS1 mass range. The default charge state was set to 3+. HCD fragmentation (stepped normalized collision energy; 25.5, 27, 30 %) was applied and MS/MS spectra were acquired with a resolution of 30,000 FWHM with a fixed first mass of 200 m/z

1107   after accumulation of 3x 10^6 ions or after a filling time of 35 ms (whichever occurred first). Data
1108   was acquired in profile mode. For data acquisition and processing Tune 2.0.
1109

Proteomics data analysis
For myoblasts DIA search against a library was performed. Acquired data were processed
using Spectronaut Professional v13 (Biognosys AG). For library creation, the DDA and DIA
raw files were searched with Pulsar (Biognosys AG) against the mouse UniProt database (*Mus*
musculus, v. 160106, 16,747 entries) with a list of common contaminants (247 entries)
Swissprot database appended, using default settings. For library generation, default BGS
factory settings were used.
DIA data were then uploaded and searched against this spectral library using Spectronaut
Professional (v.10 and v.13 Biognosys) and default settings. Relative quantification was
performed in Spectronaut for each pairwise comparison using a two-sided t-test performed at
the precursor level followed by multiple testing correction and default settings, except: Major
Group Quantity = median peptide quantity; Major Group Top N = OFF; Minor Group Quantity
= median precursor quantity; Minor Group Top N = OFF; Data Filtering = Q value sparse;
Normalization Strategy = Local normalization; Row Selection = Q value complete.
For Lmod1 overexpression cell lines, DIA raw data were analyzed using the directDIA pipeline
in Spectronaut v.14 (Biognosysis, Switzerland) with BGS settings besides the following
parameters: Protein LFQ method= QUANT 2.0, Proteotypicity Filter = Only protein group
specific, Major Group Quantity = Median peptide quantity, Minor Group Quantity = Median
precursor quantity, Data Filtering = Qvalue, Normalizing strategy = Local Normalization. The
data were searched against an in-house database (*Mus Musculus*, 16,747 entries) and a
contaminants (247 entries) Swissprot database. The data were searched with the following
variable modifications: Oxidation (M) and Acetyl (Protein N-term). A maximum of 2 missed
cleavages for trypsin and 5 variable modifications were allowed. The identifications were
filtered to satisfy FDR of 1 % on peptide and protein level. Relative quantification was
performed in Spectronaut for each paired comparison using the replicate samples from each
condition. The data (candidate table) and data reports (protein quantities) were then exported,
and further data analyses and visualization were performed with Rstudio using in-house
pipelines and scripts.
BioID DIA raw data were analyzed using the directDIA pipeline in Spectronaut with v.15. The
data were searched against a specific species (*Homo sapiens,* v. 160112, 20,375 entries) and
a contaminants (247 entries) Swissprot database. The data were searched with the following
modifications: Oxidation (M), Acetyl (Protein N-term) and Biotin_K. A maximum of 2 missed
cleavages for trypsin and 5 variable modifications were allowed. The identifications were
filtered to satisfy FDR of 1 % on peptide and protein level. Relative quantification was

performed in Spectronaut using LFQ QUANT 2.0 method with Global Normalization, precursor

filtering percentile using fraction 0.2 and global imputation. The data (candidate table) and

data reports (protein quantities) were then exported, and further data analyses and

visualization were performed with Rstudio using in-house pipelines and scripts.

1148

1149   Clustering and gene set enrichment analysis

Proteomics data for the myoblast differentiation time course were filtered for protein groups

quantified by at least two unique (proteotypic) peptides. All the differentiation time points

(1d:5d) were compared to undifferentiated myoblast (0d) and protein groups that showed an

absolute AVG Log2 Ratio > 0.58 and Q value < 0.25 at any time point were selected for

clustering analysis. Determination of an optimal number of clusters, clustering, and enrichment

analysis for each cluster was performed with ClueR (Yang et al. 2015) using the AVG Log2

Ratios to 0d as input.

A gene set was created to analyze the LMOD1 overexpression cell line using all the protein

groups differentially abundant in 1d vs. 0d myoblast differentiation time course (AVG Log2

Ratio > 0.58 and Q value < 0.05). This gene set was used as input for a gene set enrichment

analysis based on AVG Log2 Ratios from the comparison of LMOD1 OE vs. GFP OE

performed with WebGestalt (Liao et al. 2019).

1162

1163   Statistical analysis

Mouse and cell line experiments were performed at least in biological triplicates, and numbers

are indicated in the figure legends. The results are shown as the mean ± SEM unless indicated

otherwise in the figure legends. Statistical significance was calculated using the GraphPad

Prism software using the statistical test indicated in the figure legends, with *: p 0.05, **: p 0.01,

***: p 0.001, ****: p 0.0001 and ns (not significant) or nd (not detectable). RNA-seq approaches

were conducted in biological replicates (n = 5) using primary myoblasts isolated from individual

mice per experimental group, and statistical analysis is described in the respective section.

Data availability

1172

1173   The mass spectrometry proteomics data have been deposited to the MassIVE

1174   (https://massive.ucsd.edu) repository with the following dataset identifier:

1175

1176   Proteomic dataset: MSV000093379

1177   Username: MSV000093379_reviewer

1178   Password: ES_Lmod1

1179

1180   RNA-Seq data have been deposited on Gene Expression Omnibus (GEO) international public

1181   repository at: GSE254443

1182   Password: epkbcyyinjmxzmd

REFERENCES

Abdel Khalek, Waed, Fabienne Cortade, Vincent Ollendorff, Laure Lapasset, Lionel Tintignac, Béatrice Chabi, and Chantal Wrutniak-Cabello. 2014. "SIRT3, a Mitochondrial NAD⁺-Dependent Deacetylase, Is Involved in the Regulation of Myoblast Differentiation." *PloS One* 9 (12): e114388.

Abmayr, Susan M., and Grace K. Pavlath. 2012. "Myoblast Fusion: Lessons from Flies and Mice." *Development* 139 (4): 641–56.

Armand, Anne-Sophie, Iman Laziz, and Christophe Chanoine. 2006. "FGF6 in Myogenesis." *Biochimica et Biophysica Acta* 1763 (8): 773–78.

Baghdadi, Meryem B., David Castel, Léo Machado, So-Ichiro Fukada, David E. Birk, Frederic Relaix, Shahragim Tajbakhsh, and Philippos Mourikis. 2018. "Reciprocal Signalling by Notch-Collagen V-CALCR Retains Muscle Stem Cells in Their Niche." *Nature* 557 (7707): 714–18.

Baghdadi, Meryem B., Joao Firmino, Kartik Soni, Brendan Evano, Daniela Di Girolamo, Philippos Mourikis, David Castel, and Shahragim Tajbakhsh. 2018. "Notch-Induced miR-708 Antagonizes Satellite Cell Migration and Maintains Quiescence." *Cell Stem Cell* 23 (6): 859–68.e5.

Bai, Wenlong, and Xiaohong Zhang. 2016. "Nucleus or Cytoplasm? The Mysterious Case of SIRT1's Subcellular Localization." *Cell Cycle* 15 (24): 3337–38.

Bartolome, Aleksandar, Julia C. Heiby, Domenico Di Fraia, Ivonne Heinze, Hannah Knaudt, Ellen Späth, Omid Omrani, et al. 2023. "ProteasomeID: Quantitative Mapping of Proteasome Interactomes and Substrates for in Vitro and in Vivo Studies." *bioRxiv*. https://doi.org/10.1101/2022.08.09.503299.

Bentzinger, C. Florian, Yu Xin Wang, Nicolas A. Dumont, and Michael A. Rudnicki. 2013. "Cellular Dynamics in the Muscle Satellite Cell Niche." *EMBO Reports* 14 (12): 1062–72.

Bentzinger, C. Florian, Yu Xin Wang, Julia von Maltzahn, Vahab D. Soleimani, Hang Yin, and Michael A. Rudnicki. 2013. "Fibronectin Regulates Wnt7a Signaling and Satellite Cell Expansion." *Cell Stem Cell* 12 (1): 75–87.

Boczkowska, Malgorzata, Grzegorz Rebowski, Elena Kremneva, Pekka Lappalainen, and Roberto Dominguez. 2015. "How Leiomodin and Tropomodulin Use a Common Fold for Different Actin Assembly Functions." *Nature Communications* 6 (1): 1–12.

Brack, Andrew S., Heidi Bildsoe, and Simon M. Hughes. 2005. "Evidence That Satellite Cell Decrement Contributes to Preferential Decline in Nuclear Number from Large Fibres during Murine Age-Related Muscle Atrophy." *Journal of Cell Science* 118 (Pt 20): 4813–21.

Broussy, Sylvain, Hanna Laaroussi, and Michel Vidal. 2020. "Biochemical Mechanism and Biological Effects of the Inhibition of Silent Information Regulator 1 (SIRT1) by EX-527 (SEN0014196 or Selisistat)." *Journal of Enzyme Inhibition and Medicinal Chemistry* 35 (1): 1124–36.

Budai, Zsofia, Laszlo Balogh, and Zsolt Sarang. 2018. "Altered Gene Expression of Muscle Satellite Cells Contributes to Agerelated Sarcopenia in Mice." *Current Aging Science* 11 (3): 165–72.

Chakkalakal, Joe V., Kieran M. Jones, M. Albert Basson, and Andrew S. Brack. 2012. "The Aged Niche Disrupts Muscle Stem Cell Quiescence." *Nature* 490 (7420): 355–60.

Chang, Natasha C., and Michael A. Rudnicki. 2014. "Satellite Cells: The Architects of Skeletal Muscle." *Current Topics in Developmental Biology* 107:161–81.

Chen, Ieng-Yi, Jacqueline Lypowy, Jayashree Pain, Danish Sayed, Stan Grinberg, Ralph R.

Alcendor, Junichi Sadoshima, and Maha Abdellatif. 2006. "Histone H2A.z Is Essential
for Cardiac Myocyte Hypertrophy but Opposed by Silent Information Regulator 2α*."
The Journal of Biological Chemistry 281 (28): 19369–77.

Chen, Poshen B., Hsiuyi V. Chen, Diwash Acharya, Oliver J. Rando, and Thomas G. Fazzio.
2015. "R Loops Regulate Promoter-Proximal Chromatin Architecture and Cellular
Differentiation." *Nature Structural & Molecular Biology* 22 (12): 999–1007.

Chereau, David, Malgorzata Boczkowska, Aneta Skwarek-Maruszewska, Ikuko Fujiwara,
David B. Hayes, Grzegorz Rebowski, Pekka Lappalainen, Thomas D. Pollard, and
Roberto Dominguez. 2008. "Leiomodin Is an Actin Filament Nucleator in Muscle Cells."
Science 320 (5873): 239–43.

Codato, Roberta, Martine Perichon, Arnaud Divol, Ella Fung, Athanassia Sotiropoulos, Anne
Bigot, Jonathan B. Weitzman, and Souhila Medjkane. 2019. "The SMYD3
Methyltransferase Promotes Myogenesis by Activating the Myogenin Regulatory
Network." *Scientific Reports* 9 (1): 1–14.

Conley, C. A. 2001. "Leiomodin and Tropomodulin in Smooth Muscle." *American Journal of
Physiology. Cell Physiology* 280 (6): C1645–56.

Conley, C. A., K. L. Fritz-Six, A. Almenar-Queralt, and V. M. Fowler. 2001. "Leiomodins:
Larger Members of the Tropomodulin (Tmod) Gene Family." *Genomics* 73 (2): 127–39.

Cui, Can, Shunshun Han, Xiaoxu Shen, Haorong He, Yuqi Chen, Jing Zhao, Yuanhang Wei,
et al. 2020. "ISLR Regulates Skeletal Muscle Atrophy via IGF1-PI3K/Akt-Foxo Signaling
Pathway." *Cell and Tissue Research* 381 (3): 479–92.

Diaz-Ruiz, Alberto, Marta Gonzalez-Freire, Luigi Ferrucci, Michel Bernier, and Rafael de
Cabo. 2015. "SIRT1 Synchs Satellite Cell Metabolism with Stem Cell Fate." *Cell Stem
Cell* 16 (2): 103–4.

Dos Santos, Matthieu, Akansha M. Shah, Yichi Zhang, Svetlana Bezprozvannaya, Kenian
Chen, Lin Xu, Weichun Lin, et al. 2023. "Opposing Gene Regulatory Programs
Governing Myofiber Development and Maturation Revealed at Single Nucleus
Resolution." *Nature Communications* 14 (1): 1–14.

Eliazer, Susan, Jonathon M. Muncie, Josef Christensen, Xuefeng Sun, Rebecca S. D'Urso,
Valerie M. Weaver, and Andrew S. Brack. 2019. "Wnt4 from the Niche Controls the
Mechano-Properties and Quiescent State of Muscle Stem Cells." *Cell Stem Cell* 25 (5):
654–65.e4.

Finke, Daniel, Leonard M. Schanze, Friederike Schreiter, Michael M. Kreußer, Hugo A.
Katus, Johannes Backs, and Lorenz H. Lehmann. 2022. "Histone Deacetylase 4
Deletion Broadly Affects Cardiac Epigenetic Repression and Regulates Transcriptional
Susceptibility via H3K9 Methylation." *Journal of Molecular and Cellular Cardiology* 162
(January):119–29.

Fowler, Velia M., and Roberto Dominguez. 2017. "Tropomodulins and Leiomodins: Actin
Pointed End Caps and Nucleators in Muscles." *Biophysical Journal* 112 (9): 1742–60.

Fre, Silvia, Mathilde Huyghe, Philippos Mourikis, Sylvie Robine, Daniel Louvard, and Spyros
Artavanis-Tsakonas. 2005. "Notch Signals Control the Fate of Immature Progenitor
Cells in the Intestine." *Nature* 435 (7044): 964–68.

Friedrichs, Melanie, Florian Wirsdöerfer, Stefanie B. Flohé, Sabine Schneider, Manuela
Wuelling, and Andrea Vortkamp. 2011. "BMP Signaling Balances Proliferation and
Differentiation of Muscle Satellite Cell Descendants." *BMC Cell Biology* 12 (June):26.

Fulco, Marcella, Yana Cen, Po Zhao, Eric P. Hoffman, Michael W. McBurney, Anthony A.
Sauve, and Vittorio Sartorelli. 2008. "Glucose Restriction Inhibits Skeletal Myoblast
Differentiation by Activating SIRT1 through AMPK-Mediated Regulation of Nampt."

Developmental Cell 14 (5): 661–73.

Fulco, Marcella, R. Louis Schiltz, Simona Iezzi, M. Todd King, Po Zhao, Yoshihiro Kashiwaya, Eric Hoffman, Richard L. Veech, and Vittorio Sartorelli. 2003. "Sir2 Regulates Skeletal Muscle Differentiation as a Potential Sensor of the Redox State." *Molecular Cell* 12 (1): 51–62.

Fu, Xin, Huating Wang, and Ping Hu. 2015. "Stem Cell Activation in Skeletal Muscle Regeneration." *Cellular and Molecular Life Sciences: CMLS* 72 (9): 1663–77.

Ganassi, Massimo, Sara Badodi, Kees Wanders, Peter S. Zammit, and Simon M. Hughes. 2020. "Myogenin Is an Essential Regulator of Adult Myofibre Growth and Muscle Stem Cell Homeostasis." *eLife* 9 (October). https://doi.org/10.7554/eLife.60445.

Gorynia, Sabine, Tiago M. Bandeiras, Filipa G. Pinho, Colin E. McVey, Clemens Vonrhein, Adam Round, Dmitri I. Svergun, Peter Donner, Pedro M. Matias, and Maria Arménia Carrondo. 2011. "Structural and Functional Insights into a Dodecameric Molecular Machine - the RuvBL1/RuvBL2 Complex." *Journal of Structural Biology* 176 (3): 279–91.

Grifone, Raphaëlle, Audrey Saquet, Manon Desgres, Claudia Sangiorgi, Caterina Gargano, Zhenlin Li, Dario Coletti, and De-Li Shi. 2021. "Rbm24 Displays Dynamic Functions Required for Myogenic Differentiation during Muscle Regeneration." *Scientific Reports* 11 (1): 9423.

Guerin, Colleen M., and Sunita G. Kramer. 2009. "Cytoskeletal Remodeling during Myotube Assembly and Guidance: Coordinating the Actin and Microtubule Networks." *Communicative & Integrative Biology* 2 (5): 452–57.

Heng, Yi-Wen, and Cheng-Gee Koh. 2010. "Actin Cytoskeleton Dynamics and the Cell Division Cycle." *The International Journal of Biochemistry & Cell Biology* 42 (10): 1622–33.

Higuchi-Sanabria, Ryo, Joseph W. Paul, Jenni Durieux, Camila Benitez, Phillip A. Frankino, Sarah U. Tronnes, Gilberto Garcia, Joseph R. Daniele, Samira Monshietehadi, and Andrew Dillin. 2018. "Spatial Regulation of the Actin Cytoskeleton by HSF-1 during Aging." *Molecular Biology of the Cell* 29 (21): 2522–27.

Hindi, Lubna, Joseph D. McMillan, Dil Afroze, Sajedah M. Hindi, and Ashok Kumar. 2017. "Isolation, Culturing, and Differentiation of Primary Myoblasts from Skeletal Muscle of Adult Mice." *Bio-Protocol* 7 (9). https://doi.org/10.21769/BioProtoc.2248.

Hisahara, Shin, Susumu Chiba, Hiroyuki Matsumoto, Masaya Tanno, Hideshi Yagi, Shun Shimohama, Makoto Sato, and Yoshiyuki Horio. 2008. "Histone Deacetylase SIRT1 Modulates Neuronal Differentiation by Its Nuclear Translocation." *Proceedings of the National Academy of Sciences of the United States of America* 105 (40): 15599–604.

Horio, Yoshiyuki, Takashi Hayashi, Atsushi Kuno, and Risa Kunimoto. 2011. "Cellular and Molecular Effects of Sirtuins in Health and Disease." *Clinical Science* 121 (5): 191–203.

Hung, Margaret, Hsiao-Fan Lo, Grace E. L. Jones, and Robert S. Krauss. 2023. "The Muscle Stem Cell Niche at a Glance." *Journal of Cell Science* 136 (24). https://doi.org/10.1242/jcs.261200.

Hunt, Liam C., Aradhana Upadhyay, Jalal A. Jazayeri, Elizabeth M. Tudor, and Jason D. White. 2013. "An Anti-Inflammatory Role for Leukemia Inhibitory Factor Receptor Signaling in Regenerating Skeletal Muscle." *Histochemistry and Cell Biology* 139 (1): 13–34.

Hwang, Ara B., and Andrew S. Brack. 2018. "Muscle Stem Cells and Aging." *Current Topics in Developmental Biology* 126:299–322.

Kim, Kun Ho, Jiamin Qiu, and Shihuan Kuang. 2020. "Isolation, Culture, and Differentiation of Primary Myoblasts Derived from Muscle Satellite Cells." *Bio-Protocol* 10 (14): e3686.

Kim, Yu Jin, Min Jeong Cho, Won Dong Yu, Myung Joo Kim, Sally Yunsun Kim, and Jae Ho Lee. 2022. "Links of Cytoskeletal Integrity with Disease and Aging." *Cells* 11 (18). https://doi.org/10.3390/cells11182896.

Kislinger, Thomas, Anthony O. Gramolini, Yan Pan, Khaled Rahman, David H. MacLennan, and Andrew Emili. 2005. "Proteome Dynamics during C2C12 Myoblast Differentiation." *Molecular & Cellular Proteomics: MCP* 4 (7): 887–901.

Knight, James Dr, and Rashmi Kothary. 2011. "The Myogenic Kinome: Protein Kinases Critical to Mammalian Skeletal Myogenesis." *Skeletal Muscle* 1 (September):29.

Lai, Wing-Fu, and Wing-Tak Wong. 2020. "Roles of the Actin Cytoskeleton in Aging and Age-Associated Diseases." *Ageing Research Reviews* 58 (March):101021.

Lakisic, Goran, Alice Lebreton, Renaud Pourpre, Olivia Wendling, Emanuele Libertini, Elizabeth J. Radford, Morwenna Le Guillou, et al. 2016. "Role of the BAHD1 Chromatin-Repressive Complex in Placental Development and Regulation of Steroid Metabolism." *PLoS Genetics* 12 (3): e1005898.

Larraín, J., J. Alvarez, J. R. Hassell, and E. Brandan. 1997. "Expression of Perlecan, a Proteoglycan That Binds Myogenic Inhibitory Basic Fibroblast Growth Factor, Is down Regulated during Skeletal Muscle Differentiation." *Experimental Cell Research* 234 (2): 405–12.

Larsson, Lars, Hans Degens, Meishan Li, Leonardo Salviati, Young Il Lee, Wesley Thompson, James L. Kirkland, and Marco Sandri. 2019. "Sarcopenia: Aging-Related Loss of Muscle Mass and Function." *Physiological Reviews* 99 (1): 427–511.

Lepper, Christoph, Terence A. Partridge, and Chen-Ming Fan. 2011. "An Absolute Requirement for Pax7-Positive Satellite Cells in Acute Injury-Induced Skeletal Muscle Regeneration." *Development* 138 (17): 3639–46.

Liao, Yuxing, Jing Wang, Eric J. Jaehnig, Zhiao Shi, and Bing Zhang. 2019. "WebGestalt 2019: Gene Set Analysis Toolkit with Revamped UIs and APIs." *Nucleic Acids Research* 47 (W1): W199–205.

Lukjanenko, Laura, M. Juliane Jung, Nagabhooshan Hegde, Claire Perruisseau-Carrier, Eugenia Migliavacca, Michelle Rozo, Sonia Karaz, et al. 2016. "Loss of Fibronectin from the Aged Stem Cell Niche Affects the Regenerative Capacity of Skeletal Muscle in Mice." *Nature Medicine* 22 (8): 897–905.

Lu-Nguyen, Ngoc, Alberto Malerba, Linda Popplewell, Fred Schnell, Gunnar Hanson, and George Dickson. 2017. "Systemic Antisense Therapeutics for Dystrophin and Myostatin Exon Splice Modulation Improve Muscle Pathology of Adult Mdx Mice." *Molecular Therapy. Nucleic Acids* 6 (March):15–28.

Mackmull, Marie-Therese, Bernd Klaus, Ivonne Heinze, Manopriya Chokkalingam, Andreas Beyer, Robert B. Russell, Alessandro Ori, and Martin Beck. 2017. "Landscape of Nuclear Transport Receptor Cargo Specificity." *Molecular Systems Biology* 13 (12): 962.

Maltzahn, Julia von, Andrew E. Jones, Robin J. Parks, and Michael A. Rudnicki. 2013. "Pax7 Is Critical for the Normal Function of Satellite Cells in Adult Skeletal Muscle." *Proceedings of the National Academy of Sciences of the United States of America* 110 (41): 16474–79.

Massenet, Jimmy, Edward Gardner, Bénédicte Chazaud, and F. Jeffrey Dilworth. 2021. "Epigenetic Regulation of Satellite Cell Fate during Skeletal Muscle Regeneration." *Skeletal Muscle* 11 (1): 4.

Mauro, A. 1961. "Satellite Cell of Skeletal Muscle Fibers." *The Journal of Biophysical and Biochemical Cytology* 9 (2): 493–95.

Meyer, Gretchen A., Simon Schenk, and Richard L. Lieber. 2013. "Role of the Cytoskeleton

in Muscle Transcriptional Responses to Altered Use." *Physiological Genomics* 45 (8): 321–31.

Moresi, Viviana, Michele Carrer, Chad E. Grueter, Oktay F. Rifki, John M. Shelton, James A. Richardson, Rhonda Bassel-Duby, and Eric N. Olson. 2012. "Histone Deacetylases 1 and 2 Regulate Autophagy Flux and Skeletal Muscle Homeostasis in Mice." *Proceedings of the National Academy of Sciences of the United States of America* 109 (5): 1649–54.

Mournetas, Virginie, Emmanuelle Massouridès, Jean-Baptiste Dupont, Etienne Kornobis, Hélène Polvèche, Margot Jarrige, Alan R. L. Dorval, et al. 2021. "Myogenesis Modelled by Human Pluripotent Stem Cells: A Multi-Omic Study of Duchenne Myopathy Early Onset." *Journal of Cachexia, Sarcopenia and Muscle* 12 (1): 209–32.

Moynihan, Kathryn A., Andrew A. Grimm, Marie M. Plueger, Ernesto Bernal-Mizrachi, Eric Ford, Corentin Cras-Méneur, M. Alan Permutt, and Shin-Ichiro Imai. 2005. "Increased Dosage of Mammalian Sir2 in Pancreatic Beta Cells Enhances Glucose-Stimulated Insulin Secretion in Mice." *Cell Metabolism* 2 (2): 105–17.

Murphy, Malea M., Jennifer A. Lawson, Sam J. Mathew, David A. Hutcheson, and Gabrielle Kardon. 2011. "Satellite Cells, Connective Tissue Fibroblasts and Their Interactions Are Crucial for Muscle Regeneration." *Development* 138 (17): 3625–37.

Nanda, Vivek, and Joseph M. Miano. 2012. "Leiomodin 1, a New Serum Response Factor-Dependent Target Gene Expressed Preferentially in Differentiated Smooth Muscle Cells." *The Journal of Biological Chemistry* 287 (4): 2459–67.

Nanda, Vivek, Ting Wang, Milos Pjanic, Boxiang Liu, Trieu Nguyen, Ljubica Perisic Matic, Ulf Hedin, et al. 2018. "Functional Regulatory Mechanism of Smooth Muscle Cell-Restricted LMOD1 Coronary Artery Disease Locus." *PLoS Genetics* 14 (11): e1007755.

Nauen, David W., Michael C. Haffner, Juyun Kim, Qizhi Zheng, Hao Yin, Angelo M. DeMarzo, Vasiliki Mahairaki, Carlo Colantuoni, J. Geoffrey Pickering, and Tory P. Johnson. 2020. "Putative Autoantigen Leiomodin-1 Is Expressed in the Human Brain and in the Membrane Fraction of Newly Formed Neurons." *Pathogens* 9 (12). https://doi.org/10.3390/pathogens9121036.

Nguyen, Mai Thi, You Han Won, Tae Won Kwon, and Wan Lee. 2022. "Twinfilin-1 Is an Essential Regulator of Myogenic Differentiation through the Modulation of YAP in C2C12 Myoblasts." *Biochemical and Biophysical Research Communications* 599 (April):17–23.

Peck, Barrie, Chun-Yuan Chen, Ka-Kei Ho, Paolo Di Fruscia, Stephen S. Myatt, R. Charles Coombes, Matthew J. Fuchter, Chwan-Deng Hsiao, and Eric W-F Lam. 2010. "SIRT Inhibitors Induce Cell Death and p53 Acetylation through Targeting Both SIRT1 and SIRT2." *Molecular Cancer Therapeutics* 9 (4): 844–55.

Pfaffl, M. W. 2001. "A New Mathematical Model for Relative Quantification in Real-Time RT-PCR." *Nucleic Acids Research* 29 (9): e45.

Pollard, Thomas D., and John A. Cooper. 2009. "Actin, a Central Player in Cell Shape and Movement." *Science* 326 (5957): 1208–12.

Qualmann, Britta, and Michael M. Kessels. 2009. "New Players in Actin Polymerization--WH2-Domain-Containing Actin Nucleators." *Trends in Cell Biology* 19 (6): 276–85.

Ryall, James G., Stefania Dell'Orso, Assia Derfoul, Aster Juan, Hossein Zare, Xuesong Feng, Daphney Clermont, et al. 2015. "The NAD(+)-Dependent SIRT1 Deacetylase Translates a Metabolic Switch into Regulatory Epigenetics in Skeletal Muscle Stem Cells." *Cell Stem Cell* 16 (2): 171–83.

Saccone, Valentina, and Pier Lorenzo Puri. 2010. "Epigenetic Regulation of Skeletal

Myogenesis." *Organogenesis* 6 (1): 48–53.

Sambasivan, Ramkumar, Roseline Yao, Adrien Kissenpfennig, Laetitia Van Wittenberghe,
Andràs Paldi, Barbara Gayraud-Morel, Hind Guenou, Bernard Malissen, Shahragim
Tajbakhsh, and Anne Galy. 2011. "Pax7-Expressing Satellite Cells Are Indispensable
for Adult Skeletal Muscle Regeneration." *Development* 138 (17): 3647–56.

Schmidt, Manuel, Svenja C. Schüler, Sören S. Hüttner, Björn von Eyss, and Julia von
Maltzahn. 2019. "Adult Stem Cells at Work: Regenerating Skeletal Muscle." *Cellular and*
Molecular Life Sciences: CMLS 76 (13): 2559–70.

Schüler, Svenja C., Joanna M. Kirkpatrick, Manuel Schmidt, Deolinda Santinha, Philipp
Koch, Simone Di Sanzo, Emilio Cirri, Martin Hemberg, Alessandro Ori, and Julia von
Maltzahn. 2021. "Extensive Remodeling of the Extracellular Matrix during Aging
Contributes to Age-Dependent Impairments of Muscle Stem Cell Functionality." *Cell*
Reports 35 (10): 109223.

Seale, Patrick. 2003. *Pax7 Is Required for Muscle Satellite Cell Specification and*
Regenerative Myogenesis. McMaster University.

Shefer, Gabi, Daniel P. Van de Mark, Joshua B. Richardson, and Zipora Yablonka-Reuveni.
2006. "Satellite-Cell Pool Size Does Matter: Defining the Myogenic Potency of Aging
Skeletal Muscle." *Developmental Biology* 294 (1): 50–66.

Sousa-Victor, Pedro, Laura García-Prat, Antonio L. Serrano, Eusebio Perdiguero, and Pura
Muñoz-Cánoves. 2015. "Muscle Stem Cell Aging: Regulation and Rejuvenation." *Trends*
in Endocrinology and Metabolism: TEM 26 (6): 287–96.

Sousa-Victor, Pedro, Susana Gutarra, Laura García-Prat, Javier Rodriguez-Ubreva, Laura
Ortet, Vanessa Ruiz-Bonilla, Mercè Jardí, et al. 2014. "Geriatric Muscle Stem Cells
Switch Reversible Quiescence into Senescence." *Nature* 506 (7488): 316–21.

Szklarczyk, Damian, Andrea Franceschini, Michael Kuhn, Milan Simonovic, Alexander Roth,
Pablo Minguez, Tobias Doerks, et al. 2011. "The STRING Database in 2011: Functional
Interaction Networks of Proteins, Globally Integrated and Scored." *Nucleic Acids*
Research 39 (Database issue): D561–68.

Tanno, Masaya, Atsushi Kuno, Toshiyuki Yano, Tetsuji Miura, Shin Hisahara, Satoko
Ishikawa, Kazuaki Shimamoto, and Yoshiyuki Horio. 2010. "Induction of Manganese
Superoxide Dismutase by Nuclear Translocation and Activation of SIRT1 Promotes Cell
Survival in Chronic Heart Failure." *The Journal of Biological Chemistry* 285 (11): 8375–
82.

Tanno, Masaya, Jun Sakamoto, Tetsuji Miura, Kazuaki Shimamoto, and Yoshiyuki Horio.
2007. "Nucleocytoplasmic Shuttling of the NAD+-Dependent Histone Deacetylase
SIRT1." *The Journal of Biological Chemistry* 282 (9): 6823–32.

Tannu, Nilesh S., Vamshi K. Rao, Ritcha M. Chaudhary, Francesco Giorgianni, Abdelwahab
E. Saeed, Yong Gao, and Rajendra Raghow. 2004. "Comparative Proteomes of the
Proliferating C2C12 Myoblasts and Fully Differentiated Myotubes Reveal the Complexity
of the Skeletal Muscle Differentiation Program." *Molecular & Cellular Proteomics: MCP*
3 (11): 1065–82.

Tiffin, N., R. D. Williams, J. Shipley, and K. Pritchard-Jones. 2003. "PAX7 Expression in
Embryonal Rhabdomyosarcoma Suggests an Origin in Muscle Satellite Cells." *British*
Journal of Cancer 89 (2): 327–32.

Tolkatchev, Dmitri, Carol C. Gregorio, and Alla S. Kostyukova. 2022. "The Role of Leiomodin
in Actin Dynamics: A New Road or a Secret Gate." *The FEBS Journal* 289 (20): 6119–
31.

Tong, Chao, Alex Morrison, Samantha Mattison, Su Qian, Mark Bryniarski, Bethany Rankin,

Jun Wang, D. Paul Thomas, and Ji Li. 2013. "Impaired SIRT1 Nucleocytoplasmic Shuttling in the Senescent Heart during Ischemic Stress." *FASEB Journal: Official Publication of the Federation of American Societies for Experimental Biology* 27 (11): 4332–42.

Tsukada, Takehiro, Christopher T. Pappas, Natalia Moroz, Parker B. Antin, Alla S. Kostyukova, and Carol C. Gregorio. 2010. "Leiomodin-2 Is an Antagonist of Tropomodulin-1 at the Pointed End of the Thin Filaments in Cardiac Muscle." *Journal of Cell Science* 123 (Pt 18): 3136–45.

Vachharajani, Vidula T., Tiefu Liu, Xianfeng Wang, Jason J. Hoth, Barbara K. Yoza, and Charles E. McCall. 2016. "Sirtuins Link Inflammation and Metabolism." *Journal of Immunology Research* 2016 (January):8167273.

Voronova, Anastassia, Erin Coyne, Ashraf Al Madhoun, Joel V. Fair, Neven Bosiljcic, Catherine St-Louis, Grace Li, et al. 2013. "Hedgehog Signaling Regulates MyoD Expression and Activity." *The Journal of Biological Chemistry* 288 (6): 4389–4404.

Wang, Dongdong, Dandan Zhao, Yuan Li, Tingjun Dai, Fuchen Liu, and Chuanzhu Yan. 2022. "TGM2 Positively Regulates Myoblast Differentiation via Enhancing the mTOR Signaling." *Biochimica et Biophysica Acta, Molecular Cell Research* 1869 (3): 119173.

Winder, S. J., T. J. Gibson, and J. Kendrick-Jones. 1995. "Dystrophin and Utrophin: The Missing Links!" *FEBS Letters* 369 (1): 27–33.

Winter, Lilli, and Gerhard Wiche. 2013. "The Many Faces of Plectin and Plectinopathies: Pathology and Mechanisms." *Acta Neuropathologica* 125 (1): 77–93.

Xie, Xin, S. Raza Mahmood, Tamara Gjorgjieva, and Piergiorgio Percipalle. 2020. "Emerging Roles of Cytoskeletal Proteins in Regulating Gene Expression and Genome Organization during Differentiation." *Nucleus* 11 (1): 53–65.

Yamakawa, Hiroyuki, Dai Kusumoto, Hisayuki Hashimoto, and Shinsuke Yuasa. 2020. "Stem Cell Aging in Skeletal Muscle Regeneration and Disease." *International Journal of Molecular Sciences* 21 (5). https://doi.org/10.3390/ijms21051830.

Yamane, A., K. Takahashi, M. Mayo, H. Vo, L. Shum, M. Zeichner-David, and H. C. Slavkin. 1998. "Induced Expression of myoD, Myogenin and Desmin during Myoblast Differentiation in Embryonic Mouse Tongue Development." *Archives of Oral Biology* 43 (5): 407–16.

Yang, Pengyi, Xiaofeng Zheng, Vivek Jayaswal, Guang Hu, Jean Yee Hwa Yang, and Raja Jothi. 2015. "Knowledge-Based Analysis for Detecting Key Signaling Events from Time-Series Phosphoproteomics Data." *PLoS Computational Biology* 11 (8): e1004403.

Yuen, Michaela, Lisa Worgan, Jessika Iwanski, Christopher T. Pappas, Himanshu Joshi, Jared M. Churko, Susan Arbuckle, et al. 2022. "Neonatal-Lethal Dilated Cardiomyopathy due to a Homozygous LMOD2 Donor Splice-Site Variant." *European Journal of Human Genetics: EJHG* 30 (4): 450–57.

Zammit, Peter S., Frederic Relaix, Yosuke Nagata, Ana Pérez Ruiz, Charlotte A. Collins, Terence A. Partridge, and Jonathan R. Beauchamp. 2006. "Pax7 and Myogenic Progression in Skeletal Muscle Satellite Cells." *Journal of Cell Science* 119 (Pt 9): 1824–32.

Zhang, Kuo, Yuying Zhang, Lijie Gu, Miaomiao Lan, Chuncheng Liu, Meng Wang, Yang Su, et al. 2018. "Islr Regulates Canonical Wnt Signaling-Mediated Skeletal Muscle Regeneration by Stabilizing Dishevelled-2 and Preventing Autophagy." *Nature Communications* 9 (1): 1–16.