1   **Exonuclease action of replicative polymerase gamma drives**

2   **damage-induced mitochondrial DNA clearance**

7   Akshaya Seshadri[1,2] and Anjana Badrinarayanan[1*]

12   [1]National Centre for Biological Sciences-TIFR, Bangalore; [2]SASTRA University, Thanjavur

14   Correspondence to: anjana@ncbs.res.in

29   **Keywords:** mtDNA damage, PolG, Mip1, DNA replication, proofreading

Abstract

Mitochondrial DNA (mtDNA) replication is essential for mitochondrial function. This is carried out by a dedicated DNA polymerase gamma, with 5'-3' polymerase and 3'-5' proofreading/ exonuclease activity. Perturbations to either properties can have pathological consequences. Predominant sources for replication stress are DNA lesions, such as those induced by oxidative damage. How mtDNA lesions affect the polymerase activity and mtDNA stability *in vivo* is not fully understood. To address this, we induce mtDNA-specific damage in *S. cerevisiae*. We observe that mtDNA damage results in significant mtDNA loss. This loss occurs independent of cell cycle progression or cell division, suggesting an active mechanism for damaged mtDNA clearance. We implicate the 3'-5' exonuclease activity of the mtDNA polymerase in this clearance, with rates of loss being affected by cellular dNTP levels. Overall, our findings reveal context-dependent, selective regulation of two critical but opposing functions of polymerase gamma to ensure mitochondrial genome integrity.

Introduction

Faithful replication and maintenance of mitochondrial DNA (mtDNA) is essential for mitochondrial function and cell survival under respiratory growth conditions. Most eukaryotic cells contain multiple copies of their mtDNA, packaged into distinct mtDNA nucleoids (Chen & Butow, 2005). The number of nucleoids increases as a function of cell volume (Seel et al., 2023). Replication of mtDNA within these nucleoids occurs independently of nuclear genome replication (Bogenhagen & Clayton, 1977), (Sasaki et al., 2017), and is carried out by a dedicated DNA polymerase, Polymerase gamma (PolG) (Genga et al., 1986), (Lestienne, 1987), (Copeland & Longley, 2014). PolG belongs to the A-family of polymerases, which includes *E. coli* DnaI and the T7 DNA polymerase (Kaguni, 2004). In metazoans, PolG is a trimeric holoenzyme: a PolGα subunit with 5'-3' polymerase and 3'-5' exonuclease domains, and a dimeric PolGβ that increases the processivity of PolGα, via regulation of both polymerase and exonuclease activity (Oliveira et al., 2015). The budding yeast ortholog of PolG (Mip1) is a processive enzyme encoding the 5'-3' polymerase and 3'-5' exonuclease domains in a single polypeptide (Viikov et al., 2011). *In vitro,* PolG (without exonuclease activity) displays an error rate of 15×10^{-6} per nucleotide (Longley et al., 2001). The synthesis-associated error rate is reduced to $<8 \times 10^{-6}$ per nucleotide by the exonuclease domain, that recognizes mis-paired 3' termini during synthesis (Foury & Vanderstraeten, 1992), (Kaguni & Olson, 1989). *In vivo,* mutations in exonuclease function can increase mtDNA mutation rates by several 100-fold (Foury et al., 2004) and mice models of exonuclease-deficient PolG have displayed elevated mtDNA mutations, cardiomyopathy and premature ageing (Copeland & Longley, 2014).

In addition to maintaining replication fidelity at the site of DNA synthesis, the exonuclease domain of the mitochondrial polymerase plays important roles in other contexts as well. Given the polyploid nature of mtDNA, mutant DNA molecules typically constitute only a small fraction of the total mtDNA. It has been suggested that elimination of the damaged mtDNA copies, combined with the replication of the intact ones could be a mechanism by which mtDNA integrity and function is maintained (Shokolenko & Alexeyev, 2015). In several instances, the exonuclease domain has been implicated in such clearance. For example, linear mtDNA fragments (generated by action of restriction endonucleases) are subject to degradation by the exonuclease domain of PolG (N Nissanka et al., 2018),(Peeva et al., 2018). Furthermore, in conditions of extreme starvation in yeast, it has been shown that exonuclease domain of Mip1 results in mtDNA clearance, likely as a salvage pathway to replenish dNTP levels

(Medeiros et al., 2018). Interestingly, in Trypanosoma, one of the six mtDNA polymerases was
recently shown to function as an exonuclease only, as it lacked the catalytic activities associated
with polymerase function (Delzell et al., 2022).

It remains unclear as to how the exonucleolytic function, that is typically restricted to only a
few bases around the terminal nucleotides in a primer strand, is extended to these different
capacities, while still maintaining a balance between the polymerase and exonuclease activity.
Indeed, imbalance in the same can have direct consequences on replication fidelity as well as
total amount of DNA synthesis carried out by the polymerase (Szczepanowska & Foury, 2010).
Even within the context of mtDNA replication, the polymerase is likely to encounter
replication-stalling lesions (such as those generated by oxidative damage) frequently during
DNA synthesis, in part due to the reactive oxygen species rich environment of the mitochondria
(Anderson et al., 2020). This makes mtDNA susceptible to damage significantly more than its
nuclear counterpart (Yakes & Van Houten, 1997), with point mutation rates of mtDNA
reported to be ~10-fold higher than the nuclear genome (Parsons et al., 1997), (Haag-Liautard
et al., 2008). How the polymerase responds to such DNA lesions *in vivo,* and how
polymerase/exonuclease balance is maintained to ensure mtDNA integrity in such a scenario
remains unknown.

We previously developed a tool to induce mtDNA-specific damage using a mitochondrially-
targeted bacterial toxin, DarT (mtDarT) (Dua et al., 2022). This toxin generates adducts on
single-stranded DNA, creating replication-stalling lesions specifically on mtDNA (Dua et al.,
2022; Jankevicius et al., 2016; Lawarée et al., 2020). In fermentative conditions, where yeast
cells can proliferate without mtDNA or its associated functions, such mtDNA damage resulted
in reorganization of the mitochondrial network and asymmetric segregation of mtDNA-less
mitochondria, with an acceleration in this process observed in cells carrying a mutation in the
exonuclease domain of PolG (Mip1 henceforth) (Dua et al., 2022). In this study, we utilized
mtDarT to understand how such lesions affect mtDNA maintenance and Mip1 function *in vivo*
in *S. cerevisiae,* in conditions where oxidative phosphorylation (and hence mitochondrial
function) is essential for cell proliferation. We specifically monitored the impact of mtDNA
lesions on mtDNA copy number. We found that mtDNA damage resulted in mtDNA loss. This
loss was independent of cell cycle progression or cell division, suggesting an active mechanism
of mtDNA clearance. In support, we uncovered a role for the 3'-5' exonuclease of Mip1 in this
mtDNA loss under damage, with cellular dNTP levels regulating the rates of clearance. We

propose a model where Mip1 encounter with a DNA lesion could switch its function from
synthesis to degradation, thus resulting in the clearance of the lesion-containing mtDNA.
Together, our work highlights the selective regulation of polymerase/ exonuclease balance for
the maintenance of mtDNA integrity under persistent DNA damage.

Results

mtDNA damage results in mtDNA loss

To study the effect of mtDNA damage on mtDNA maintenance, we utilized a tool that was
previously characterised by our group to specifically damage mtDNA in *S. cerevisiae* (Dua et
al., 2022). For this, we expressed mitochondrially-targeted DarT (mtDarT) from a β-estradiol-
inducible promoter in yeast cells grown in non-fermentable glycerol-containing medium
(YPG), with cells maintained in mid-exponential growth throughout the course of the
experiment (Dua et al., 2022) (Fig. EV1A). We first confirmed the expression of mtDarT in
YPG medium via Western blotting and found robust induction of mtDarT 4 and 8 hrs after
addition of β-estradiol (Fig. EV1B). Furthermore, cells expressing mtDarT were compromised
in survival, as assessed by their growth on YPG plates carrying β-estradiol. This was in contrast
to empty vector control cells, which did not show a growth defect on β-estradiol containing
YPG plates (Fig. EV1C). We then proceeded to monitor the fate of mtDNA in cells subjected
to damage, via SYBR Green I (SGI) staining of mtDNA nucleoids in live cells (Jajoo et al.,
2016). As a control, we compared mtDarT-expressing cells with cells carrying an empty vector
and subjected to the same experimental regime (Fig. EV1A).

In the absence of damage, pulse-staining with SGI resulted in the labelling of mtDNA nucleoids
(Jajoo et al., 2016), that were seen as distinct cytoplasmic puncta in the cell (Fig. 1A). These
puncta were localized within the mitochondria (Fig. EV1D), and no nuclear DNA staining was
observed (Fig. EV1D). As a control, SGI staining of ρ^0 cells (which lack mtDNA) resulted in
nuclear DNA staining and no cytoplasmic puncta were observed (Fig. EV1E, and as reported
previously (Baruffini et al., 2010)), suggesting that the cytoplasmic puncta we were visualizing
were indeed mitochondrial nucleoids. In the absence of damage, multiple mtDNA foci were
observed in all cells (empty vector, Fig. 1A). We found that the number of nucleoids per cell
increased as a function of cell area, in line with previous observations showing mtDNA
nucleoid scaling with cell size (Seel et al., 2023) (empty vector (0 hr), Spearman's correlation
coefficient (ρ) 0.76, CI: 12.43503, 13.18894) (Fig. 1B). The percentage of cells with mtDNA

foci as well as the relationship of the number of nucleoids with cell area was maintained in empty vector cells, even after 8 hrs of β-estradiol treatment (Fig. 1B-C).

In cells carrying the mtDarT expression vector, we observed a modest increase in the percentage of cells without mtDNA foci even without damage induction, likely due to leaky expression of mtDarT (mtDarT (0hr) (Fig.1A-C). This increase was significantly pronounced upon induction of mtDarT, with fewer nucleoids per cell observed when compared to the empty vector control (Fig 1A-B, mtDarT). At 8 hrs post-induction, the relationship between number of nucleoids and cell area was significantly perturbed (Spearman's correlation coefficient (ρ) 0.53, CI: 8.851388, 9.779604) (Fig. 1B), and the percentage of cells without mtDNA foci increased gradually, with $32\pm4\%$ cells without mtDNA foci observed after 8 hrs of mtDarT induction (Fig. 1C). Indeed, majority of cells had fewer than 10 nucleoids per cell 8 hrs after damage induction (Fig. 1D). Consistent with the depletion of mtDNA in these conditions, we found that mtDarT-expressing cells frequently displayed nuclear DNA staining as well (Fig. 1A). Together, these observations suggest that mtDNA damage results in mtDNA loss under non-fermentative growth conditions.

mtDNA loss occurs independent of cell cycle progression

What cellular mechanisms might result in the observed loss of mtDNA? We first considered the possibility that asymmetric partitioning of damaged mtDNA during division could result in the generation of ρ^0 cells as we had previously seen in fermentative growth conditions (Dua & Badrinarayanan, 2023). In this scenario, cells at each division would receive fewer numbers of nucleoids (or no nucleoids at all). To test this possibility, we arrested cells in either G1/S or G2/M stage of the cell cycle and then induced mtDarT to follow the fate of mtDNA under damage in non-dividing cells. For this, we first pre-treated $\Delta bar1$ cells (*MAT*a) with α-factor which resulted in a G1/S phase arrest (Fig. 2A), following which we induced mtDNA damage in this arrested population. As a control, damage was also induced in an asynchronously growing cell culture for comparison. We observed that the number of nucleoids was reduced in arrested cells after 8 hrs of damage, with few to no nucleoids in these highly elongated cells (Fig. 2B). Additionally, the percentage of cells with mtDNA foci was comparable between asynchronous (72.4%) and arrested (70.5%) populations after 8 hrs of damage (Fig. 2C). Similarly, the relationship between cell size and number of nucleoids was also perturbed in the arrested cells expressing mtDarT ($\rho = 0.44$ (CI: 12.65272, 15.20398) (Fig. 2D). In contrast, empty vector cells had densely packed nucleoid foci dispersed throughout the cell (Fig. 2B)

and the relationship between cell size and number of nucleoids per cell was also maintained (ρ
= 0.87 (CI: 33.81381, 40.35107) (Fig. 2D). Thus, G1/S arrested cells showed mtDNA loss
under damage in respiratory growth conditions (Fig. 2C-E). To eliminate any potential
influence from the cell cycle stage under arrest, we separately investigated mtDNA loss in
nocodazole-induced G2/M-arrested cells (Fig. EV2A). Upon mtDarT expression, here too we
observed a decrease in the percentage of cells with mtDNA foci (Fig. EV2B), similar to that
observed in G1/S arrested cells. Taken together, these observations indicate that cell cycle
progression (and cell division) does not contribute to mtDNA loss under mtDNA damage.

mtDNA loss is dependent on Mip1 exonuclease

An important source for mtDNA loss could be mitophagy, which can drive the clearance of
dysfunctional mitochondria (with mutant mtDNA). Such dysfunction-induced mitochondrial
clearance has been reported in case of *Drosophila* cells carrying a mutation on mtDNA (Kandul
et al., 2016). To assess if this occurs in case of mtDNA damage in budding yeast, we deleted
the receptor essential for mitophagy in budding yeast (Atg32) (Kanki et al., 2009). We observed
that mitophagy was not required for mtDNA loss under damage, with a non-significant
difference in the percentage of cells with mtDNA nucleoids between wild type and Atg32-
deleted cells even after 8 hrs of damage induction (Fig. EV3A). Furthermore, we did not
observe a significant change in Cox4 protein levels under damage conditions, suggesting that
mitochondrial content did not significantly change upon damage induction (Fig. EV3B). These
observations are also consistent with our previous findings that mitophagy (which is typically
observed in much longer timescales) may not play a role under mtDNA damage in yeast (Dua
et al., 2022).

Since mtDNA loss in response to mtDNA damage occurred even in non-dividing cells, we
considered the possibility of an active mtDNA clearance mechanism, such as a nuclease-
mediated DNA degradation under damage. In *S. cerevisiae,* Nuc1, homolog to the mammalian
EndoG, is the major mitochondrial endonuclease (Dzierzbicki et al., 2012). In order to check
if Nuc1 was responsible for clearing out mtDNA under damage, we deleted Nuc1 and tested
the effect of the same on mtDNA nucleoid numbers. We observed that Nuc1 was not essential
for mtDNA loss under damage (Fig. EV3C).

Another nuclease that has been connected with mtDNA clearance is the exonuclease domain
of the mitochondrial replicative polymerase (PolG/ Mip1). In mammalian cells, PolG

exonuclease domain contributes to degradation of linear mtDNA fragments (Nissanka et al.,
2018), (Peeva et al., 2018). Similarly, in yeast cells facing extreme starvation, the exonuclease
domain of Mip1 has been implicated in mtDNA clearance (Medeiros et al., 2018). Whether
such activity is also relevant in the context of mtDNA damage is unknown. We thus assessed
whether the exonuclease activity of Mip1 could also result in mtDNA degradation under
conditions of mtDNA damage.

Based on previously characterized mutations, we generated an exonuclease-deficient Mip1 via
mutating 2 catalytic residues in the ExoI motif of the enzyme, Aspartate 171 (Asp171) to
Alanine 171 (Ala171) and Glutamate 173 (Glu173) to Alanine 173 (Ala173) (*mip1*$^{exo-}$
henceforth) (Foury & Vanderstraeten, 1992b), (Medeiros et al., 2018), (Longley et al., 1998)
(Fig. 3A). Both Mip1 and *mip1*$^{exo-}$ were expressed from their endogenous locus on the
chromosome. Western blot experiments confirmed that their expression in mtDarT-induced
samples were comparable (Fig. EV3D). We did note that *mip1*$^{exo-}$ cells expressing mtDarT
showed an elevated percentage of cells without mtDNA foci prior to damage induction (Fig.
3C). However, upon mtDarT induction, *mip1*$^{exo-}$ cells maintained nucleoids even after 8 hrs of
damage, and damage-induced mtDNA depletion was not observed as was seen in case of wild
type cells (Fig. 3B-E). Although there was no loss of nucleoids, the relationship between
number of nucleoids and cell area was still perturbed in this background (Spearman's
correlation coefficient (ρ) 0.44, CI: 13.54795, 14.56954) (Fig. 3D, mtDarT). These
observations suggest that the exonuclease activity of the mitochondrial replicative polymerase
drives mtDNA loss under DNA damage. The compromised scaling of nucleoid numbers with
increasing cell size even in the absence of exonuclease activity of Mip1 suggests that
replication progression could be additionally compromised under mtDNA damage.

Exonuclease activity of Mip1 drives mtDNA loss under damage

To test whether exonuclease activity was directly driving mtDNA clearance under damage, we
complemented the *mip1*$^{exo-}$ strain with ectopic expression of the exonuclease domain in this
background. For this, we generated a variant of *mip1* (Mip1-CΔ263) expressed from the *MIP1*
promoter on a low copy replicating plasmid where the last 263 amino acids in the C-terminus
of Mip1 are deleted. This mutant has been extensively characterized previously, and *in vitro*
studies have shown that it is deficient in polymerization activity, but still retains the ability of
polymerase to bind DNA, as well as an intact exonuclease domain and function (Trasviña-

Arenas et al., 2019; Young et al., 2021). We confirmed the expression of the truncated protein

via western blot and that the protein colocalized with mitochondrial nucleoids (Fig. EV4A -B).

Similar to wild type cells (Fig. 4A), we observed that expression of Mip1-CΔ263 in *mip1^{exo-}*

background resulted in a significant decrease in the percentage of cells containing mtDNA foci

upon mtDNA damage induction (Fig. 4B). So as to confirm that exonuclease activity from the

Mip1-CΔ263 construct was indeed contributing to the observed mtDNA loss, we introduced

the exonuclease mutation in the truncation construct. Western blot of both Mip1-CΔ263 and

the exonuclease deficient Mip1-CΔ263 (*mip1^{exo-}*-CΔ263) showed similar expression levels,

and this protein also colocalized with mtDNA (Fig. EV4A-B). Furthermore, under damage, we

no longer observed mtDNA loss occurring in the absence of exonuclease activity from this

exonuclease-deficient construct when expressed in *mip1^{exo-}* strain (Fig. 4C). Based on these

results, we conclude that Mip1-CΔ263 does indeed harbour exonuclease activity that drives the

observed mtDNA loss under damage.

To further assess the role of the exonuclease domain of Mip1 in mtDNA clearance, we

generated two additional mutants in highly conserved regions of the polymerase domain that

should be compromised in polymerase activity, while still retaining exonuclease function.

These variants (Mip1^{G651S}) and (Mip1^{H734Y}) are located in the thumb and finger sub-domains

and are reported to compromise catalysis and selectivity in nucleotide binding respectively

(Kasiviswanathan et al., 2009). Both mutants are orthologous to PolG mutations, G848S and

H932Y, falling in regions of high amino-acid homology among several organisms including

budding yeast (Fig.S4D) (Stumpf et al., 2010). We found that both mutants were expressed in

cells and colocalized with mtDNA (Fig. EV4C, E). These mutants were able to complement

mip1^{exo-}, with mtDNA loss being observed upon damage induction (Fig. 4D-E). Together, these

results support a direct role for the exonuclease domain of Mip1 in mtDNA clearance under

conditions of mtDNA damage.

Rates of mtDNA loss are influenced by dNTP levels

How does Mip1 exonuclease activity result in mtDNA clearance under damage? Given that the

polymerase and exonuclease domains are both present on the same protein in case of Mip1, we

wondered whether exonuclease-mediated clearance occurred on mtDNA that is being actively

replicated. Studies in *S. cerevisiae* have shown that elevation in dNTP levels can increase Mip1

polymerase function (Lecrenier & Foury, 1995; Medeiros et al., 2018; Vanderstraeten et al.,

1998; Viikov et al., 2012). We thus hypothesized that increasing polymerase activity (and
hence access) on mtDNA should impact the rates of mtDNA loss under damage. To assess this,
we deleted Sml1, a negative regulator of ribonucleotide reductase (RNR) activity in *S.*
cerevisiae, that results in an increase in cellular dNTP levels (Zhao et al., 1998), and assessed
the rates of mtDNA clearance under damage in this background.

We confirmed that the expression of Mip1 in Δ*sml1* background was comparable to that in wild
type cells under damage (Fig. EV5A). Furthermore, Δ*sml1* cells grew similarly to wild type
cells in non-fermentable media (Fig. EV5B), and had comparable number of mtDNA nucleoids
in the absence of damage (Fig. 5A). Upon mtDarT induction, Δ*sml1* cells showed a significant
increase in mtDNA loss (Fig. 5B-C). In comparison to wild type, the rates of mtDNA loss were
higher in Δ*sml1* cells. After 8 hrs of damage induction, the percentage of cells without mtDNA
foci were found to be 55±5% and 32±4% in Δ*sml1* and wild type cells respectively (Fig. 5B).
Importantly, this elevated mtDNA loss in Δ*sml1* cells was dependent on the exonuclease
activity of Mip1, with no loss being observed in Δ*sml1 mip1^{exo-}* cells (Fig. 5D).

In sum, our observations implicate the exonuclease domain of Mip1 in mtDNA clearance under
conditions of mtDNA damage, with dNTP levels affecting the rates of this mtDNA loss.

Discussion
Taken together, our study places the exonuclease activity of PolG as a central mediator of
mtDNA loss under mtDNA damage conditions. Thus, the clearance of damaged mtDNA copies
employs the very apparatus that replicates it. Such a mechanism of damaged mtDNA clearance
would restrict the degradative function of the polymerase to the damaged mtDNA alone, and
circumvent the requirement to have an additional pathway to sense and respond to the damage
more globally. In support, we did not observe a significant contribution from mitophagy in this
process. While we cannot rule out the possibility that this damage clearance mechanism may
contribute to mtDNA loss at much longer timescales, our observations are consistent with other
reports suggesting that mitophagy may not play a significant role in mtDNA damage clearance
in case of yeast cells (as opposed to metazoan systems) (Dua et al., 2022) (Abeliovich, 2023).
Indeed, such drastic DNA loss might only be tolerated in a multicopy genome, allowing cells
to lose most copies yet retain the capacity to restore homeostasis post-damage removal, in line
with the 'disposable genome' hypothesis for mitochondria (Shokolenko & Alexeyev, 2015).

The modest loss of mtDNA nucleoids from leaky expression of mtDarT observed in the absence of Mip1 exonuclease function suggests that mtDNA replication may be additionally compromised. However, our data strongly support a major role for exonuclease action in the mtDNA loss observed upon damage induction. For example, in the *mip1^exo-* mutant where polymerase is still active and present, cells do not lose mtDNA upon mtDarT expression (Fig. 3C). This highlights that a possible perturbation of replication alone under damage does not immediately reflect as significant mtDNA loss, at least in the timescales in which we track mtDNA.

How are contrasting functions of Mip1 polymerase and exonuclease selectively regulated in this context-dependent manner (replication vs degradation)? We do not yet fully understand the mechanism by which substrates for Mip1 exonuclease activity (such as nicks or breaks) would be generated at sites of mtDarT damage. Sources for this can include events such as collisions at stalled replisomes, breaks due to the instability of exposed ssDNA, or additional damage triggered by oxidative stress. Based on our observations of accelerated mtDNA clearance under damage in Δ*sml1* cells, it is tempting to hypothesize that the hand-off from synthesis to degradation might occur at the site of replication stalling. Such an arrangement would ensure that the exonucleolytic action of Mip1 is regulated to prevent promiscuous mtDNA clearance in the absence of DNA damage. In the context of heteroplasmy (Nissanka & Moraes, 2020), where only a fraction of mtDNA would typically carry a mutation, selective clearance would allow the retention of intact genomes over the mutant copies. In support, studies in *Drosophila* cells have shown preferential replication of non-mutated mtDNA copies (Hill et al., 2014). More recently, reports in budding yeast have also shown that mutant mtDNA are compartmentalized via modulation of the cristae morphology (Jakubke et al., 2021). It is possible that exonuclease activity of Mip1 on these sequestered mtDNA would additionally contribute to the selective clearance of damaged genomes. How Mip1 accesses damaged mtDNA, how much DNA is degraded by a single exonuclease, and the dynamics of mtDNA resynthesis upon removal of the stress are exciting and outstanding questions.

Exonuclease activity of the replicative polymerase is canonically considered in the context of proofreading, where only few mismatched nucleotides from newly synthesized DNA are removed (Kunkel & Mosbaugh, 1989). However, more recent evidences suggest that the exonuclease domain of the mitochondrial replicative polymerase may play more versatile roles. Exonuclease action has previously been observed on linear mtDNA fragments (Nissanka et al., 2018),(Peeva et al., 2018) and under conditions of extreme starvation (where dNTP levels are

depleted) (Medeiros et al., 2018). Exonuclease activity is also required for mtDNA ligation
during replication (Macao et al., 2015), absence of which results in accumulation of mtDNA
fragments. A previous study from our group (Dua et al., 2022) and our current study both
highlight the key role of Mip1 exonuclease in mtDNA loss under damage. In growth conditions
with a fermentable carbon source (where yeast cells can continue to grow and divide even in
the absence of functional mitochondria/ mtDNA), we had previously found that Mip1
exonuclease activity played a role in mtDNA clearance; in its absence, clearance rates were
accelerated via asymmetric mtDNA segregation and cell division, resulting in rapid generation
of ρ^0 cells (Dua et al., 2022). When the possibility of viable cell divisions is compromised (such
as in the present study using a non-fermentable carbon source, where mitochondrial function
is essential for cell proliferation), Mip1 exonuclease appears to be the sole driver of this
clearance during mtDNA stress.

The implication of exonuclease function in mtDNA clearance under conditions of mtDNA
damage suggests that it is a central player in the maintenance of mtDNA integrity as well,
beyond the regulation of replication-associated errors. Indeed, the choice to repair or degrade
damaged mtDNA could be driven by the type and quantity of damage. For example, given the
absence of active nucleotide excision repair in mitochondria (Alexeyev et al., 2013; Clayton et
al., 1974), it is possible that the exonuclease-mediated pathway for mtDNA degradation may
be the first response to persistent, replication-stalling lesions. Other contexts can include
mtDNA deletions or mutations, where selective degradation of the damaged copies would be
a reliable mechanism for ensuring mitochondrial genome integrity maintenance.

Methods

Yeast strains, media and growth conditions

Strains, plasmids and oligos used in this study are listed in the Reagents and tools table. Cells were grown at 30ºC with 180 rpm shaking, in YP (yeast extract 1%, peptone 2%) medium. All experiments were performed in non-fermentable conditions using glycerol (2%) as the carbon source. Dextrose (2%, fermentable medium) was used for preparatory experiments including transformations etc. All experiments were performed in log-phase of growth (OD600, between 0.2-0.6). Appropriate antibiotics and inducers were added to the medium as required. Strains generated were haploid prototrophs belonging to CEN.PK background (Van Dijken et al., 2000). For generating strains with C-terminal tagging or deletion of a genomic locus, PCR-based targeted homologous recombination strategy was used as described in (Longtine et al., 1998). Strains were verified by PCR, followed by fluorescence microscopy and/ or western blotting for validating the insertion of epitope tags.

Plasmid construction

Plasmids expressing empty vector and mtDarT: Details of plasmids expressing the empty vector (pNABC557) and mtDarT (pNABC561) can be found in Dua et al., 2022.

Mip1-CΔ263 (pNABC775): The plasmid containing the Mip1-CΔ263 sequence (pNABC775), expressed under the pMip1 promoter, was generated by PCR amplification of the pMip1-Mip1-CΔ263 fragment (ending at Ser991 in Mip1) from wild-type yeast genomic DNA. The primers used were AS188 and AS202. A stop codon was introduced after residue 991. This fragment was then Gibson cloned into pNABC555 between SacI and BamHI sites, removing the cyc1 promoter.

Mip1-CΔ263-mNeongreen (pNABC777): A mNeongreen-tagged version of Mip1-CΔ263 (pNABC777) was created by Gibson cloning the Mip1-CΔ263 fragment (amplified using AS182 and AS163 from wild-type yeast genomic DNA) with a fragment containing a 10 amino acid linker followed by mNeongreen (amplified using AS184 and AS185). The combined fragment was then inserted into pNABC555 at the XbaI site.

Mip1-CΔ263-mCherry (pNABC943): To construct the mCherry-tagged Mip1-CΔ263 (pNABC943), the pMip1-Mip1-CΔ263 fragment (ending at Ser991 in Mip1) was amplified from wild-type yeast genomic DNA using primers AS188 and AS202. This fragment was then

Gibson cloned with a fragment containing a 10 amino acid linker followed by mCherry

(amplified using AS200 and AS201) into pNABC555 between SacI and BamHI sites, removing

the cyc1 promoter.

Plasmids expressing $mip1^{exo-}$-CΔ263 (pNABC944) and $mip1^{exo-}$-CΔ263-mCherry

(pNABC945) were constructed similarly to pNABC777 and pNABC943. The yeast genomic

DNA used for amplifying the Mip1 sequence was from NABC573 ($mip1^{exo-}$).

Mip1-G651S (pNABC946): The plasmid containing mip1-G651S under pMip1 was generated

by PCR amplification of the pMip1-mip1-G651S fragment (ending at Ser991 in Mip1) from

wild-type yeast genomic DNA using primers AS271 and AS272. A stop codon was introduced

at residue 991. The fragment was then Gibson cloned into pNABC555 between SacI and

BamHI sites, removing the cyc1 promoter.

Mip1-G651S-mNeongreen (pNABC947): The plasmid containing mip1-G651S tagged with

mNeongreen (pNABC947) was generated similarly. The pMip1-mip1-G651S fragment

(ending at Ser991) was amplified using primers AS271 and AS272 from wild-type yeast

genomic DNA. This fragment was Gibson cloned with a 17 amino acid linker followed by

mNeongreen (amplified using AS225 and AS192) and inserted into pNABC555 between SacI

and BamHI sites.

pNABC948 (Mip1-H734Y): This plasmid, expressing mip1-H734Y under pMip1, was

generated by PCR amplification of the pMip1-mip1-H734Y fragment (ending at Ser991 in

Mip1) from wild type yeast genomic DNA using primers AS273 and AS274. A stop codon was

introduced at residue 991. The fragment was then Gibson cloned into pNABC555 between

SacI and BamHI sites, removing the cyc1 promoter.

pNABC949 (Mip1-H734Y-mNeongreen): To generate the mNeongreen-tagged version of

Mip1-H734Y (pNABC949), the pMip1-mip1-H734Y fragment (ending at Ser991 in Mip1)

was amplified using primers AS273 and AS274 from wild type yeast genomic DNA. This

fragment was then Gibson cloned with a 10 amino acid linker followed by mNeongreen into

pNABC555 between SacI and BamHI sites. All plasmids described were sequenced to confirm

the presence of relevant sequence and/or mutations.

Plasmids for expression of the empty vector (pNABC557), mtDarT (pNABC561), Mip1-

CΔ263 (pNABC775), $mip1^{exo-}$-CΔ263 (pNABC944), $mip1^{G651S}$ (pNABC946) and $mip1^{H734Y}$

(pNABC948) were freshly transformed into yeast cells before each experiment. This was to

prevent mtDNA instability that might arise from the continuous maintenance and expression

of both the toxin and Mip1 truncations/ mutants.

Yeast strain construction

Protrophic Cen.PK strain was used for all constructions.

Mip1-mCherry (NABC772): pNABC549 was used to amplify the mCherry-HygR cassette

together with overhangs for Mip1, AS79 and AS80. Transformants were screened and

confirmed for Mip1 C-ter tagging using AS78 and AS81.

Mip1-mNeongreen (NABC773): pNABC553 was used to amplify the mNeonGreen-KanR

cassette together with overhangs for Mip1, AS79 and AS80. Transformants were screened and

confirmed for Mip1 C-ter tagging using AS78 and AS81.

mip1^{exo-}-**mNeongreen (NABC680)**: Same as NABC773, but in NABC573 cells.

Δsml1 **Mip1-mNeongreen (NABC766)**: Same as NABC773, but in NABC639 cells.

Cox4-mCherry (NABC569): pNABC549 was used to amplify the mCherry-HygR cassette

together with overhangs for Cox4, ND328 and ND329. Transformants were screened and

confirmed for Cox4 C-ter tagging using ND327 and AS81.

Δsml1 **(NABC639)**: pNABC552 was used to amplify the HygR cassette together with

overhangs for Sml1, ND489 and ND490. Transformants were screened and confirmed for

Δsml1 using ND488 and AS81.

Δnuc1 **(NABC641)**: pNABC551 was used to amplify the KanR cassette together with

overhangs for Nuc1, AS92 and AS93. Transformants were screened and confirmed for *Δnuc1*

using AS91 and AS81

Δatg32 **(NABC676)**: pNABC552 was used to amplify the HygR cassette together with

overhangs for Atg32, ND162 and ND163. Transformants were screened and confirmed for

Δatg32 using ND161 and AS81.

Mip1 exonuclease-deficient mutant (NABC573): Exonuclease-deficient Mip1 mutant

(*mip1^{exo-}*) was constructed as previously described (Stuart et al., 2006), (Medeiros et al., 2018).

The mutant harbors two tandem mutations, D171A and E173A, in the ExoI motif of the

exonuclease domain. The mutations were introduced by site-directed mutagenesis where the

endogenous copy of Mip1 was deleted and subsequently transformed with a gene fragment

containing Mip1 D171A, E173A double-mutation. Insertion of the mutant fragment was

confirmed by sequencing. Deletion of Mip1 renders a cell ρ0. Therefore, mtDNA was

reintroduced into *mip1^exo-* mutants by crossing with wild type cells of opposite mating-type.

Haploid spores were dissected to obtain *mip1^exo-* mutants with mtDNA.

mtDNA damage induction

The protocol for inducing mtDNA damage was adopted from (Dua et al., 2022). Briefly, cells

carrying empty vector control plasmid (pNABC557) or mtDarT plasmid (pNABC561), with

NAT resistance, were transformed into yeast cell cells using the standard Lithium acetate

protocol. Transformants with the plasmid were selected by plating on media containing

Nourseothricin (NTC). Colonies were inoculated in YPG cultures containing NTC for plasmid

maintenance. Upon reaching early log-phase (OD~0.2-0.4), samples were collected (0 hr).

Subsequently, the cultures were treated with 100nM β-estradiol to induce expression. Inducer

was maintained at all times during the experiment. Samples were collected at 4, 8 hrs post-

induction as indicated for specific experiments. Cells were maintained in mid-exponential

phase growth throughout the experiment via back-dilution at appropriate time intervals.

Survival assay

460 Survival assay to compare growth between different strains was performed by serially diluting

461 log phase cultures (starting from OD600 of 0.06, diluted 10x, until 6×10^{-9}). 6μl of the diluted

462 culture was spotted on agar plates with respective selection markers. The plates were incubated

463 at 30°C, and growth was monitored.

Live cell staining of mtDNA

mtDNA were stained in live cells using SYBR Green I. The protocol was adopted from [28].

In brief, 1 ml aliquots of cultures were taken at specified time points, pelleted and washed in

1X PBS. This was followed by resuspension of cells in 1X PBS followed by addition of 0.5μl

of 10,000X SYBR Green I. After gently mixing the contents, cells were immediately spun

down and washed with 1X PBS. The pellet was resuspended in appropriated volume of 1X

PBS. 1μl of diluted culture was spotted on 1% water agarose pads for microscopy.

For DAPI staining, 1ml cultures were spun down and the pellets were resuspended in 1X PBS.

0.5ul of 10mg/ml DAPI was added to the cell suspension above. The dye was resuspended

gently into the cell pellet and the mixture was immediately spun down. The pellet was washed

with 1X PBS, twice, before resuspending it in appropriate volume of 1X PBS. 1μl of diluted

culture was spotted on 1% water agarose pads for microscopy.

Western blot

Trichloroacetic acid (TCA) extraction method, as previously described, was used (Dua et al., 2022). Dilution and details of the antibodies can be found in Table EV4. The protein bands were detected using chemiluminescence substrate based assay. Separately, in each case a second protein gel was probed as loading control, and visualized with Coomassie brilliant blue staining solution. For quantification of relative proteins levels, band intensities of protein of interest were obtained from ImageJ. Values were normalized to the intensity of the loading control.

G1/S arrest using α-factor

To arrest yeast cells in G1/S phase using α-factor (mating pheromone), Δ*bar1* (*MAT*a) strain was used. Overnight cultures of cells expressing empty vector or mtDarT were grown to appropriate OD (early log phase, 0.2-0.4), after which they were pelleted. The cell pellet was carefully resuspended in a pre-warmed YPG medium containing α-factor (final concentration of 25ng/ml, from 10μg/ml stock solution in Methanol) and NAT. The culture was placed in an incubator for 3 hrs for arrest to occur completely. After 3 hrs, majority of cells were arrested in G1/S phase as revealed by the formation of schmoo cells with oblong morphology (Rosebrock, 2017). Upon reaching this stage, inducer (β-estradiol) was added and samples were collected to monitor the mtDNA status. To maintain continuous arrest, cells were pelleted and resuspended in media containing fresh α-factor every 3 hrs.

G2/M arrest using Nocodazole

Overnight cultures of cells expressing empty vector or mtDarT were grown to appropriate OD (early log phase, 0.2-0.4), after which they were pelleted. The cell pellet was carefully resuspended in pre-warmed YPG medium containing Nocodazole (final concentration of 9μl/ml, from 1.5mg/ml stock solution) and NAT. The culture was placed in an incubator for 3 hrs for arrest to occur completely. After 3 hrs, majority of cells were arrested in G2/M phase as revealed by the dumbbell shape of these cells (Rosebrock, 2017). Upon reaching this stage, inducer (β-estradiol) was added and samples were collected to monitor the mtDNA status. To maintain continuous arrest, cells were pelleted and resuspended in media containing fresh Nocodazole every 3 hrs.

Flow cytometry

Cell cycle analysis using flow cytometry was performed following the protocol from (Haase & Reed, 2002). Cultures were collected and centrifuged at 3000 rpm, for 3 mins at room temperature. After discarding the media, the cells were fixed in 70% ethanol overnight at 40ºC. Following fixation, cells were pelleted and then washed in 1ml of 50mM Sodium citrate (pH 7.5). Cells were then re-suspended in 500μl RNAse A buffer (500μl sodium citrate buffer + 5μl RNase A) and incubated with shaking (350 rpm) at 50ºC for 2 hrs. Cells were then pelleted and re-suspended in Proteinase K solution (20μl enzyme (2mg/ml) + 180μl sodium citrate buffer) and incubated at 37ºC for 1 hr, with shaking (350 rpm). Following Proteinase K treatment, cells were pelleted, washed and re-suspended in 1ml of sodium citrate buffer. 100μl of this sample was made upto 1ml with sodium citrate buffer containing 1μM SYTOX Green. The sample was sonicated on ice, at 20% amplitude for 10 secs with two 5 secs pulses with a gap of 5 secs. Flow cytometry analysis was performed on Accuri C6+ flow cytometer.

Multiple sequence alignment (MSA)

MSA of PolG and related orthologs was performed using COBALT: multiple alignment tool (Papadopoulos & Agarwala, 2007). The protein sequences were obtained from Uniprot (Bateman et al., 2023). Sequences are colour-coded based on the ClustalX criteria (conservation and position) (Thompson et al., 1997). The displayed consensus sequence was obtained by setting the threshold at 50%.

Fluorescence microscopy and image analysis

Fluorescence imaging of budding yeast cells was carried out on a wide-field, epifluorescence microscope (Nikon Eclipse Ti2-E), with motorized XY-stage, 60X plan apochromat objective (NA 1.41) and a pE4000 light source (CoolLED). Focus was maintained using an infrared-based Perfect focus system. Z scanning was done using motorized Z-control. Z-stacks were acquired across 7μm of the cell in steps of 0.3μm. Images were acquired with Hamamatsu Orca Flash 4.0 camera using NIS-elements software (version 5.01).

Image analysis

Following acquisition, image stacks were deconvolved in NIS-elements software using 3D-Lucy-Richardson method. A maximum projection of the deconvolved stacks was generated in ImageJ for fluorescence image analysis. Total number of nucleoids was obtained using the 3D-maxima finder plugin in ImageJ. Cell segmentation was performed using Cell-ACDC program (Padovani et al., 2022) generating cell ROI's for single cell based analyses. ROI's with

incorrect segmentation or with abnormal SGI staining were manually excluded from the
analysis. Number of nucleoids per cell was obtained by custom macros written using the
Mitochondria analyser plugin (Chaudhry et al., 2020) in ImageJ where the number of foci was
overlayed on cell ROI's to obtain number of nucleoid foci per cell. Cell area was calculated
using Analyse Particles function in ImageJ. Correlation density plots were plotted using
RStudio (Version 1.3.1093).

For colocalization analysis, line profiles were created on composite images using ImageJ to
generate intensity values. The values obtained for each channel was overlayed and plotted
using GraphPad Prism (8.4.2) software. Quantification of percentage of cells with nucleoids
was done using mother cells alone to avoid inconsistencies that might arise from incomplete
mtDNA segregation in bud cells. Correlation plots were plotted ignoring the zero values for ρ^0
cells and density plots for mt-nucleoid number scaling per cell were plotted for all the cells
(inc. ρ^0) in a field. For these plots, all the cells (mother, bud) in the image field were used.
All microscopy data are deposited in the Biostudies Portal (accession number S-BIAD1418).
557

Statistical analysis
Statistical analyses were performed using GraphPad Prism (8.4.2) after assessing normality
using Shapiro-Wilk test. For comparisons between two groups, parametric two-tailed unpaired
t-tests were applied for normally-distributed data, while non-parametric Wilcoxon's tests were
applied for non-normal data. To compare differences in percentages of cells with nucleoid foci
across multiple time points, a repeated measures one-way ANOVA was employed. This was
followed by use of Tukey's multiple comparisons test to derive exact P values for differences
between 2 groups of data. Since the percentages for empty vector control were nearly 100%,
non-parametric tests were applied to account for lack of variance. For assessing correlations,
Spearman's correlation coefficient was obtained through custom scripts on R. P values ≤ 0.05
were considered to be significant, P values ≤ 0.05 were indicated by (*), P values ≤ 0.01 by
(**), P values ≤ 0.001 by (***) and P values ≤ 0.0001 by "****". P values > 0.05 were
considered non-significant. Details of the test for each dataset including results obtained can
be accessed in Dataset EV1.

Data availability

All the source data from this publication have deposited in the Biostudies Portal and can be accessed via https://www.ebi.ac.uk/biostudies/bioimages/studies/S-BIAD1418.

Acknowledgements

The authors are grateful to Dr. Nitish Dua and Prof. Sunil Laxman for sharing of reagents. The authors thank Dr. Nitish Dua, Dr. Asha Mary Joseph and Mr. Aditya Kamat for helpful discussions and feedback on the manuscript. The authors would like to thank Dr. Francesco Padovani for assistance with CellACDC customizations. The authors would like to thank Mr. Anton Iyer for help with coding. This work was supported by fellowships from CSIR (AS) as well as grants from HFSP RPG (AB) (RGP0038/2021-102) and intramural funding from NCBS-TIFR (AB) (Department of Atomic Energy, Government of India, Project Identification No. RTI 4006).

Disclosure Statement and Competing Interests

The authors declare no competing interests.

References

References

Abeliovich, H. (2023). Mitophagy in yeast: known unknowns and unknown unknowns.
Biochemical Journal, *480*(20), 1639. https://doi.org/10.1042/BCJ20230279

Alexeyev, M., Shokolenko, I., Wilson, G., & LeDoux, S. (2013). The maintenance of
mitochondrial DNA integrity--critical analysis and update. *Cold Spring Harbor*
Perspectives in Biology, *5*(5). https://doi.org/10.1101/CSHPERSPECT.A012641

Anderson, A. P., Luo, X., Russell, W., & Yin, Y. W. (2020). Oxidative damage diminishes
mitochondrial DNA polymerase replication fidelity. *Nucleic Acids Research*, *48*(2),
817–829. https://doi.org/10.1093/NAR/GKZ1018

Baruffini, E., Ferrero, I., & Foury, F. (2010). In vivo analysis of mtDNA replication defects
in yeast. *Methods (San Diego, Calif.)*, *51*(4), 426–436.
https://doi.org/10.1016/J.YMETH.2010.02.023

Bateman, A., Martin, M. J., Orchard, S., Magrane, M., Ahmad, S., Alpi, E., Bowler-Barnett,
E. H., Britto, R., Bye-A-Jee, H., Zhang, J. et al. (2023). UniProt: the Universal Protein
Knowledgebase in 2023. *Nucleic Acids Research*, *51*(D1), D523–D531.
https://doi.org/10.1093/NAR/GKAC1052

Bogenhagen, D., & Clayton, D. A. (1977). Mouse L cell mitochondrial DNA molecules are
selected randomly for replication throughout the cell cycle. *Cell*, *11*(4), 719–727.
https://doi.org/10.1016/0092-8674(77)90286-0

Chaudhry, A., Shi, R., & Luciani, D. S. (2020). A pipeline for multidimensional confocal
analysis of mitochondrial morphology, function, and dynamics in pancreatic β-cells.
American Journal of Physiology - Endocrinology and Metabolism, *318*(2), E87–E101.
https://doi.org/10.1152/AJPENDO.00457.2019/ASSET/IMAGES/LARGE/ZH10022082
780008.JPEG

Chen, X. J., & Butow, R. A. (2005). The organization and inheritance of the mitochondrial
genome. *Nature Reviews Genetics*, *6*(11), 815–825. https://doi.org/10.1038/nrg1708

Clayton, D. A., Doda, J. N., & Friedberg, E. C. (1974). The absence of a pyrimidine dimer
repair mechanism in mammalian mitochondria. *Proceedings of the National Academy of*
Sciences of the United States of America, *71*(7), 2777–2781.
https://doi.org/10.1073/PNAS.71.7.2777

Copeland, W. C., & Longley, M. J. (2014). Mitochondrial genome maintenance in health and
disease. *DNA Repair*, *19*, 190–198. https://doi.org/10.1016/J.DNAREP.2014.03.010

Delzell, S. B., Nelson, S. W., Frost, M. P., & Klingbeil, M. M. (2022). Trypanosoma brucei
Mitochondrial DNA Polymerase POLIB Contains a Novel Polymerase Domain Insertion
That Confers Dominant Exonuclease Activity. *Biochemistry*, *61*(23), 2751–2765.
https://doi.org/10.1021/ACS.BIOCHEM.2C00392

Dua, N., & Badrinarayanan, A. (2023). Mechanisms regulating mitochondrial DNA quality
control. *Journal of Translational Genetics and Genomics*, *7*(2), 110–125.
https://doi.org/10.20517/JTGG.2023.08

Dua, N., Seshadri, A., & Badrinarayanan, A. (2022). DarT-mediated mtDNA damage induces
dynamic reorganization and selective segregation of mitochondria. *Journal of Cell
Biology*, *221*(10). https://doi.org/10.1083/JCB.202205104/213451

Dzierzbicki, P., Kaniak-Golik, A., Malc, E., Mieczkowski, P., & Ciesla, Z. (2012). The
generation of oxidative stress-induced rearrangements in Saccharomyces cerevisiae
mtDNA is dependent on the Nuc1 (EndoG/ExoG) nuclease and is enhanced by
inactivation of the MRX complex. *Mutation Research*, *740*(1–2), 21–33.
https://doi.org/10.1016/J.MRFMMM.2012.12.004

Foury, F., Hu, J., & Vanderstraeten, S. (2004). Mitochondrial DNA mutators. *Cellular and
Molecular Life Sciences*, *61*(22), 2799–2811. https://doi.org/10.1007/S00018-004-4220-
Y/METRICS

Foury, F., & Vanderstraeten, S. (1992a). Yeast mitochondrial DNA mutators with deficient
proofreading exonucleolytic activity. *The EMBO Journal*, *11*(7), 2717–2726.
https://doi.org/10.1002/J.1460-2075.1992.TB05337.X

Foury, F., & Vanderstraeten, S. (1992b). Yeast mitochondrial DNA mutators with deficient
proofreading exonucleolytic activity. *The EMBO Journal*, *11*(7), 2717–2726.
https://doi.org/10.1002/J.1460-2075.1992.TB05337.X

Genga, A., Bianchi, L., & Foury, F. (1986). A nuclear mutant of Saccharomyces cerevisiae
deficient in mitochondrial DNA replication and polymerase activity. *Journal of
Biological Chemistry*, *261*(20), 9328–9332. https://doi.org/10.1016/S0021-
9258(18)67658-6

Haag-Liautard, C., Coffey, N., Houle, D., Lynch, M., Charlesworth, B., & Keightley, P. D.
(2008). Direct Estimation of the Mitochondrial DNA Mutation Rate in Drosophila
melanogaster. *PLOS Biology*, *6*(8), e204.
https://doi.org/10.1371/JOURNAL.PBIO.0060204

Haase, S. B., & Reed, S. I. (2002). Improved Flow Cytometric Analysis of the Budding Yeast
Cell Cycle. *Cell Cycle*, *1*(2), 117–121. https://doi.org/10.4161/cc.1.2.114

Hill, J. H., Chen, Z., & Xu, H. (2014). Selective propagation of functional mitochondrial
DNA during oogenesis restricts the transmission of a deleterious mitochondrial variant.
Nature Genetics, *46*(4), 389–392. https://doi.org/10.1038/NG.2920

Jajoo, R., Jung, Y., Huh, D., Viana, M. P., Rafelski, S. M., Springer, M., & Paulsson, J.
(2016). Accurate concentration control of mitochondria and nucleoids. *Science (New
York, N.Y.)*, *351*(6269), 169–172. https://doi.org/10.1126/SCIENCE.AAA8714

Jakubke, C., Roussou, R., Maiser, A., Schug, C., Thoma, F., Bunk, D., Hörl, D., Leonhardt,

H., Walter, P., Osman, C. et al. (2021). Cristae-dependent quality control of the
mitochondrial genome. *Science Advances*, *7*(36).
https://doi.org/10.1126/SCIADV.ABI8886

Jankevicius, G., Ariza, A., Ahel, M., & Ahel, I. (2016). The Toxin-Antitoxin System DarTG
Catalyzes Reversible ADP-Ribosylation of DNA. *Molecular Cell*, *64*(6), 1109–1116.
https://doi.org/10.1016/j.molcel.2016.11.014

Kaguni, L. S., & Olson, M. W. (1989). Mismatch-specific 3'----5' exonuclease associated
with the mitochondrial DNA polymerase from Drosophila embryos. *Proceedings of the*
National Academy of Sciences of the United States of America, *86*(17), 6469.
https://doi.org/10.1073/PNAS.86.17.6469

Kaguni, Laurie S. (2004). DNA polymerase γ, the mitochondrial replicase. *Annual Review of*
Biochemistry, *73*, 293–320.
https://doi.org/10.1146/ANNUREV.BIOCHEM.72.121801.161455

Kandul, N. P., Zhang, T., Hay, B. A., & Guo, M. (2016). Selective removal of deletion-
bearing mitochondrial DNA in heteroplasmic Drosophila. *Nature Communications 2016*
7:1, *7*(1), 1–11. https://doi.org/10.1038/ncomms13100

Kanki, T., Wang, K., Cao, Y., Baba, M., & Klionsky, D. J. (2009). Atg32 is a mitochondrial
protein that confers selectivity during mitophagy. *Developmental Cell*, *17*(1), 98.
https://doi.org/10.1016/J.DEVCEL.2009.06.014

Kasiviswanathan, R., Longley, M. J., Chan, S. S. L., & Copeland, W. C. (2009). Disease
Mutations in the Human Mitochondrial DNA Polymerase Thumb Subdomain Impart
Severe Defects in Mitochondrial DNA Replication. *The Journal of Biological*
Chemistry, *284*(29), 19501. https://doi.org/10.1074/JBC.M109.011940

Kunkel, T. A., & Mosbaugh, D. W. (1989). Exonucleolytic Proofreading by a Mammalian
DNA Polymerase γ. *Biochemistry*, *28*(3), 988–995. https://doi.org/10.1021/bi00429a011

Lawarée, E., Jankevicius, G., Cooper, C., Ahel, I., Uphoff, S., & Tang, C. M. (2020). DNA
ADP-Ribosylation Stalls Replication and Is Reversed by RecF-Mediated Homologous
Recombination and Nucleotide Excision Repair. *Cell Reports*, *30*(5), 1373-1384.e4.
https://doi.org/10.1016/j.celrep.2020.01.014

Lecrenier, N., & Foury, F. (1995). Overexpression of the RNR1 gene rescues Saccharomyces
cerevisiae mutants in the mitochondrial DNA polymerase-encoding MIP1 gene. *MGG*
Molecular & General Genetics, *249*(1), 1–7. https://doi.org/10.1007/BF00290229

Lestienne, P. (1987). Evidence for a direct role of the DNA polymerase gamma in the
replication of the human mitochondrial DNA in vitro. *Biochemical and Biophysical*
Research Communications, *146*(3), 1146–1153. https://doi.org/10.1016/0006-
291X(87)90767-4

Longley, M. J., Nguyen, D., Kunkel, T. A., & Copeland, W. C. (2001). The Fidelity of

Human DNA Polymerase γ with and without Exonucleolytic Proofreading and the p55
Accessory Subunit. *Journal of Biological Chemistry*, *276*(42), 38555–38562.
https://doi.org/10.1074/JBC.M105230200

Longtine, M. S., Iii, K., Demarini, D. J., Shah, N. G., Wach, A., Brachat, A., Philippsen, P.,
& Pringle, J. R. (1998). Additional Modules for Versatile and Economical PCR-based
Gene Deletion and Modification in Saccharomyces cerevisiae. □□□□□ *Yeast*, *14*, 953–
961. https://doi.org/10.1002/(SICI)1097-0061(199807)14:10

Macao, B., Uhler, J. P., Siibak, T., Zhu, X., Shi, Y., Sheng, W., Olsson, M., Stewart, J. B.,
Gustafsson, C. M., & Falkenberg, M. (2015). The exonuclease activity of DNA
polymerase γ is required for ligation during mitochondrial DNA replication. *Nature*
Communications, *6*. https://doi.org/10.1038/NCOMMS8303

Matthew J. Longley, Philip A. Ropp, Susan E. Lim, and, & Copeland*, W. C. (1998).
Characterization of the Native and Recombinant Catalytic Subunit of Human DNA
Polymerase γ: Identification of Residues Critical for Exonuclease Activity and
Dideoxynucleotide Sensitivity. *Biochemistry*, *37*(29), 10529–10539.
https://doi.org/10.1021/BI980772W

Medeiros, T. C., Thomas, R. L., Ghillebert, R., & Graef, M. (2018). Autophagy balances
mtDNA synthesis and degradation by DNA polymerase POLG during starvation.
Journal of Cell Biology, *217*(5), 1601–1611. https://doi.org/10.1083/JCB.201801168

Nissanka, N, Bacman, S., Plastini, M., & Moraes, C. T. (2018). The mitochondrial DNA
polymerase gamma degrades linear DNA fragments precluding the formation of
deletions. *Nature Communications*, *9*(1). https://doi.org/10.1038/S41467-018-04895-1

Nissanka, Nadee, & Moraes, C. T. (2020). Mitochondrial DNA heteroplasmy in disease and
targeted nuclease-based therapeutic approaches. *EMBO Reports*, *21*(3).
https://doi.org/10.15252/EMBR.201949612

Oliveira, M. T., Haukka, J., & Kaguni, L. S. (2015). Evolution of the Metazoan
Mitochondrial Replicase. *Genome Biology and Evolution*, *7*(4), 943.
https://doi.org/10.1093/GBE/EVV042

Padovani, F., Mairhörmann, B., Falter-Braun, P., Lengefeld, J., & Schmoller, K. M. (2022).
Segmentation, tracking and cell cycle analysis of live-cell imaging data with Cell-
ACDC. *BMC Biology*, *20*(1). https://doi.org/10.1186/S12915-022-01372-6

Papadopoulos, J. S., & Agarwala, R. (2007). COBALT: constraint-based alignment tool for
multiple protein sequences. *Bioinformatics (Oxford, England)*, *23*(9), 1073–1079.
https://doi.org/10.1093/BIOINFORMATICS/BTM076

Parsons, T. J., Muniec, D. S., Sullivan, K., Woodyatt, N., Alliston-Greiner, R., Wilson, M.
R., Berry, D. L., Holland, K. A., Weedn, V. W., Holland, M. M. et al. (1997). A high
observed substitution rate in the human mitochondrial DNA control region. *Nature*
Genetics, *15*(4), 363–368. https://doi.org/10.1038/NG0497-363

Peeva, V., Blei, D., Trombly, G., Corsi, S., Szukszto, M. J., Rebelo-Guiomar, P., Gammage,
P. A., Kudin, A. P., Becker, C., Kunz, W. S. et al. (2018). Linear mitochondrial DNA is
rapidly degraded by components of the replication machinery. *Nature Communications*,
9(1). https://doi.org/10.1038/s41467-018-04131-w

Rosebrock, A. P. (2017). Synchronization and Arrest of the Budding Yeast Cell Cycle Using
Chemical and Genetic Methods. *Cold Spring Harbor Protocols*, *2017*(1),
pdb.prot088724. https://doi.org/10.1101/PDB.PROT088724

Sasaki, T., Sato, Y., Higashiyama, T., & Sasaki, N. (2017). Live imaging reveals the
dynamics and regulation of mitochondrial nucleoids during the cell cycle in Fucci2-
HeLa cells. *Scientific Reports 2017 7:1*, *7*(1), 1–12. https://doi.org/10.1038/s41598-017-
10843-8

Seel, A., Padovani, F., Mayer, M., Finster, A., Bureik, D., Thoma, F., Osman, C., Klecker, T.,
& Schmoller, K. M. (2023). Regulation with cell size ensures mitochondrial DNA
homeostasis during cell growth. *Nature Structural & Molecular Biology*, *30*(10), 1549–
1560. https://doi.org/10.1038/S41594-023-01091-8

Shokolenko, I. N., & Alexeyev, M. F. (2015). Mitochondrial DNA: A disposable genome? In
Biochimica et Biophysica Acta - Molecular Basis of Disease (Vol. 1852, Issue 9, pp.
1805–1809). Elsevier. https://doi.org/10.1016/j.bbadis.2015.05.016

Stuart, G. R., Santos, J. H., Strand, M. K., Van Houten, B., & Copeland, W. C. (2006).
Mitochondrial and nuclear DNA defects in Saccharomyces cerevisiae with mutations in
DNA polymerase γ associated with progressive external ophthalmoplegia. *Human
Molecular Genetics*, *15*(2), 363–374. https://doi.org/10.1093/HMG/DDI454

Stumpf, J. D., Bailey, C. M., Spell, D., Stillwagon, M., Anderson, K. S., & Copeland, W. C.
(2010). mip1 containing mutations associated with mitochondrial disease causes
mutagenesis and depletion of mtDNA in Saccharomyces cerevisiae. *Human Molecular
Genetics*, *19*(11), 2123. https://doi.org/10.1093/HMG/DDQ089

Szczepanowska, K., & Foury, F. (2010). A cluster of pathogenic mutations in the 3′–5′
exonuclease domain of DNA polymerase gamma defines a novel module coupling DNA
synthesis and degradation. *Human Molecular Genetics*, *19*(18), 3516–3529.
https://doi.org/10.1093/HMG/DDQ267

Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F., & Higgins, D. G. (1997). The
CLUSTAL_X Windows Interface: Flexible Strategies for Multiple Sequence Alignment
Aided by Quality Analysis Tools. *Nucleic Acids Research*, *25*(24), 4876–4882.
https://doi.org/10.1093/NAR/25.24.4876

Trasviña-Arenas, C. H., Hoyos-Gonzalez, N., Castro-Lara, A. Y., Rodriguez-Hernandez, A.,
Sanchez-Sandoval, M. E., Jimenez-Sandoval, P., Ayala-García, V. M., Díaz-Quezada,
C., Lodi, T., Brieba, L. G. et al. (2019). Amino and carboxy-terminal extensions of yeast
mitochondrial DNA polymerase assemble both the polymerization and exonuclease
active sites. *Mitochondrion*, *49*, 166–177. https://doi.org/10.1016/J.MITO.2019.08.005

Van Dijken, J. P., Bauer, J., Brambilla, L., Duboc, P., Francois, J. M., Gancedo, C.,
Giuseppin, M. L. F., Heijnen, J. J., Hoare, M., Pronk, J. T. et al. (2000). An
interlaboratory comparison of physiological and genetic properties of four
Saccharomyces cerevisiae strains. *Enzyme and Microbial Technology*, *26*(9–10), 706–
714. https://doi.org/10.1016/S0141-0229(00)00162-9

Vanderstraeten, S., Brûle, S. Van den, Hu, J., & Foury, F. (1998). The Role of 3′-5′
Exonucleolytic Proofreading and Mismatch Repair in Yeast Mitochondrial DNA Error
Avoidance *. *Journal of Biological Chemistry*, *273*(37), 23690–23697.
https://doi.org/10.1074/JBC.273.37.23690

Viikov, K., Jasnovidova, O., Tamm, T., & Sedman, J. (2012). C-Terminal Extension of the
Yeast Mitochondrial DNA Polymerase Determines the Balance between Synthesis and
Degradation. *PLOS ONE*, *7*(3), e33482.
https://doi.org/10.1371/JOURNAL.PONE.0033482

Viikov, K., Väljamäe, P., & Sedman, J. (2011). Yeast mitochondrial DNA polymerase is a
highly processive single-subunit enzyme. *Mitochondrion*, *11*(1), 119–126.
https://doi.org/10.1016/J.MITO.2010.08.007

Yakes, F. M., & Van Houten, B. (1997). Mitochondrial DNA damage is more extensive and
persists longer than nuclear DNA damage in human cells following oxidative stress.
Proceedings of the National Academy of Sciences of the United States of America, *94*(2),
514–519. https://doi.org/10.1073/PNAS.94.2.514

Young, M. J., Imperial, R. J., Lakhi, S., & Court, D. A. (2021). A non-radioactive DNA
synthesis assay demonstrates that elements of the Sigma 1278b Mip1 mitochondrial
DNA polymerase domain and C-terminal extension facilitate robust enzyme activity.
Yeast (Chichester, England), *38*(4), 262. https://doi.org/10.1002/YEA.3541

Zhao, X., Muller, E. G. D., & Rothstein, R. (1998). A suppressor of two essential checkpoint
genes identifies a novel protein that negatively affects dNTP pools. *Molecular Cell*,
2(3), 329–340. https://doi.org/10.1016/S1097-2765(00)80277-4

Main figure legends

Figure 1: mtDNA damage results in mtDNA clearance

(A) Representative images of mitochondrial nucleoids stained with Sybr Green I (SGI). All experiments were performed in YPG medium. [Top] nucleoids in empty vector and mtDarT cells before addition of β-estradiol (0 hr). [Bottom] nucleoids in empty vector and mtDarT cells after addition of β-estradiol (8 hrs). Dashed lines represent cell boundaries. * indicates mt-nucleoids and # indicates cells devoid of mtDNA staining. **(B)** 2D density correlation plots of the number of nucleoids per cell against the respective cell area for empty vector or mtDarT before [top] and 8 hrs post-induction [bottom]. Spearman's correlation coefficient (ρ) is indicated in the top-left corner of each plots. Data are pooled from 3 independent repeats (n > 287 cells per group, per repeat). **(C)** Percentage of cells containing mtDNA foci in empty vector [left] or mtDarT [right] expressing cells at 0, 4 and 8 hrs after induction. Data are shown from 3 independent repeats (n > 207 cells per group, per repeat). Mean and S.D. are shown. Significance was calculated using Repeated Measures one-way ANOVA and post-hoc tests. *P $\leq$ 0.05. **(D)** Density distributions for the number of nucleoids per cell in empty vector [left] and mtDarT [right] expressing cells before (0 hr) and after (4, 8 hrs) induction. Data are pooled from 3 independent repeats (n > 273 cells per group, per repeat). Scale bar, 8μm here, and in all other images. Statistical tests and exact P values for all graphs are provided in Dataset EV1.

Figure 2: mtDNA loss occurs independent of cell cycle progression

(A) Experimental setup followed for inducing mtDNA damage under cell cycle arrest using α-factor in Δ*bar1* cells. Representative images of nucleoids in asynchronous and arrested cells are shown. Flow cytometry profiles of cell cycle stages in asynchronous and arrested cells are also provided. Dashed lines represent cell boundaries. **(B)** Representative images of nucleoid foci in G1/S arrested cells. [Top] empty vector and mtDarT expressing cells at 0 hr. [Bottom] empty vector and mtDarT expressing cells at 8 hrs post-induction. Dashed lines represent cell boundaries. **(C)** Percentage of cells with mtDNA foci in empty vector (blue) or mtDarT (black) expressing cells at 0, 4 and 8 hrs after induction in [left] asynchronous population and [right] in G1/S arrested cells. Data shown from 3 independent repeats (n > 61 cells per group, per repeat). Mean and S.D. are shown. Significance was calculated using Repeated Measures one-way ANOVA and post-hoc tests. *P $\leq$ 0.05 and **P $\leq$ 0.01. **(D)** 2D density correlation plots of the number of nucleoids per cell against the respective cell area in G1/S arrested cells for empty vector or mtDarT before [top] and 8 hrs post-induction [bottom]. Spearman's correlation

coefficient (ρ) is indicated in the top-left corner of each plots. Data are pooled from 4

independent repeats (n > 50 cells per group, per repeat). **(E)** Density distributions for the

number of nucleoids per cell in empty vector [left] or mtDarT [right] expressing cells before

(0 hr) and after (4, 8 hrs) induction under G1/S arrest. Data are pooled from 4 independent

repeats (n > 50 cells per group, per repeat). Inset in mtDarT shows distribution up to 40

nucleoids/cell. Scale bar, 8μm here, and in all other images.

Figure 3: mtDNA loss is dependent on Mip1 exonuclease

(A) Schematic representing the domain organization of Mip1. Residues mutated to generate

mip1^exo- mutant are indicated. Schematic created with Biorendor.com. **(B)** Representative

images of mt-nucleoid foci in *mip1^exo-* cells. [Top] empty vector and mtDarT expressing cells

at 0 hr. [Bottom] empty vector and mtDarT expressing cells at 8 hrs. Dashed lines represent

cell boundaries. **(C)** Percentage of cells with mtDNA foci in empty vector (blue) or mtDarT

(black) at 0, 4 and 8 hrs after induction for wild type [left] and *mip1^exo-* cells [right]. Data shown

from 4 independent repeats (n > 139 cells per group, per repeat). Mean and S.D. are shown.

Significance was calculated using Repeated Measures one-way ANOVA and post-hoc tests.

*P $\leq$ 0.05. **(D)** 2D density correlation plots of the number of mt-nucleoids per cell against the

respective cell area for empty vector or mtDarT before [top] and 8 hrs post-induction [bottom]

in *mip1^exo-* cells. Spearman's correlation coefficient (ρ) is indicated in the top-left corner of

each plots. Data are pooled from 3 independent repeats (n > 149 cells per group, per repeat).

(E) Density distributions for the number of nucleoids per cell before (0 hr) and after (4, 8 hrs)

induction in empty vector [left] and mtDarT [right] expressing *mip1^exo-* cells. Data are pooled

from 3 independent repeats (n > 88 cells per group, per repeat). Scale bar, 8μm here, and in all

other images.

Figure 4: Exonuclease activity of Mip1 drives mtDNA loss under damage

(A-C) Percentage of cells with mtDNA foci in empty vector (blue) or mtDarT (black) at 0, 4

and 8 hrs after induction for wild type (A), *mip1^exo-* cells expressing Mip1-CΔ263 (B) and

mip1^exo- cells expressing *mip1^exo-* -CΔ263 (C). Data are from 3 independent repeats (n > 127

cells per group, per repeat). Mean and S.D. are shown. Significance was calculated using

Repeated Measures one-way ANOVA and post-hoc tests. *P $\leq$ 0.05 and **P $\leq$ 0.01 **(D-E)**

Percentage of cells with mtDNA foci in empty vector (blue) or mtDarT (black) at 0, 4 and 8

hrs after induction for *mip1^exo-* cells expressing Mip1^G651S (D) and *mip1^exo-* cells expressing

Mip1^H734Y (E). Data shown from 3 independent repeats (n > 119 cells for Mip1^G651S, and n >

73 cells for Mip1^H734Y per group, per repeat). Mean and S.D. are shown. Significance was

calculated using Repeated Measures one-way ANOVA and post-hoc tests. *P $\leq$ 0.05 and **P

$\leq$ 0.01.

Figure 5: Rates of mtDNA loss are influenced by dNTP levels

(A) Density distributions of the number of nucleoids per cell [left] and cell area [right] for wild type (blue) and *Δsml1* (grey) cells in the absence of damage. Data shown from 3 independent repeats (n > 287 cells per group, per repeat). **(B)** Percentage of cells containing mtDNA foci in empty vector (blue) or mtDarT (black) at 0, 4 and 8 hrs after induction in wild type [left] and *Δsml1* [right] cells. Mean and S.D. are shown. Significance was calculated using Repeated Measures one-way ANOVA and post-hoc tests. *P $\leq$ 0.05 and **P $\leq$ 0.01. Data shown from 3 independent repeats (n > 203 cells per group, per repeat). **(C)** Density distributions for the number of nucleoids per cell in empty vector [left] or mtDarT [right] at 0, 4 and 8 hrs after induction in *Δsml1* cells. (n > 303 cells per group, per repeat). **(D)** Percentage of cells containing mtDNA foci in empty vector [left] or mtDarT [right] at 0, 4 and 8 hrs after induction in *Δsml1 mip1^{exo-}* background. Data shown from 3 independent repeats (n > 139 cells per group, per repeat). Mean and S.D. are shown. Significance was calculated using Repeated Measures one-way ANOVA and post-hoc tests.

Expanded view figures

Figure EV1: mtDNA damage results in mtDNA clearance

(A) Experimental setup followed for inducing mtDNA damage using the mtDarT system is shown (Dua et al., 2022). Cells were grown in a non-fermentable carbon source (YPG) and cultures were maintained in mid-exponential growth during the course of the entire experiment. OD600 at the time of imaging was always ensured to be comparable between control (empty vector) and mtDarT-treated cells. Schematic created with Biorendor.com. **(B)** Representative Western blot of mtDarT-3x-FLAG cells before (0 hr) and after (4, 8 hrs) 100nM β-estradiol addition. Whole cell lysate is probed with Coomassie as loading control (n=3 independent repeats). **(C)** Survival of yeast cells with empty vector or mtDarT plasmid, grown on YPG media with and without 100nM β-estradiol. Representative image from 3 independent repeats is shown. Scale bar refers to the increasing dilution of cells from the left spot to the right-most spot. **(D)** Representative images of nucleoids stained with SGI and mitochondria marked with Cox4-mCherry. Dashed lines represent cell boundaries. **(E)** SGI staining in Δ*fzo1* cells (ρ^0, mtDNA absent). Dashed lines represent cell boundaries. Scale bar, 8μm here, and in all other images.

Figure EV2: mtDNA loss occurs independent of cell cycle progression

(A) Representative images of nucleoid foci in G2/M arrested, empty vector and mtDarT expressing cells at 0 hr [top] and 8 hrs after induction [bottom]. Dashed lines represent cell boundaries. **(B)** Percentage of cells containing mtDNA foci in asynchronous [left] or G2/M arrested [right] empty vector and mtDarT cells at 0 and 8 hrs after induction. Data shown from 3 independent repeats (n > 138 cells per group, per repeat). Mean and S.D. are shown. Significance was calculated using Unpaired t test (Two-tailed). $^{**}P \leq 0.01$, $^{***}P \leq 0.001$.). Scale bar, 8μm here, and in all other images.

Figure EV3: mtDNA loss is dependent on Mip1 exonuclease

(A) Percentage of cells with mtDNA foci in wild type [left] or Δ*atg32* cells [right] at 0, 4 and 8 hrs after induction with empty vector (blue) and mtDarT (black). Data shown from 3 independent repeats (n > 144 cells per group, per repeat). Mean and S.D. are shown. Significance was calculated using Repeated Measures one-way ANOVA and post-hoc tests. $^{**}P \leq 0.01$, $^{****}P \leq 0.0001$. **(B)** Western blot of Cox4-mCherry in empty vector and mtDarT cells, before (0 hr) and after (4, 8 hrs) damage induction. Representative western blot image is shown at the [top] and levels are quantified at the [bottom]. n=3 independent repeats. Mean and S.D. are shown. Significance was calculated using Repeated Measures one-way ANOVA

and post-hoc tests. **(C)** Percentage of cells with mtDNA foci in wild type [left] or Δ*nuc1* cells [right] at 0, 4 and 8 hrs after induction with empty vector (blue) and mtDarT (black). Data shown from 3 independent repeats (n > 115 cells per group, per repeat). Mean and S.D. are shown. Significance was calculated using Repeated Measures one-way ANOVA and post-hoc tests. *P ≤ 0.05. **(D)** Western blot of Mip1-mNeonGreen and Mip1^{exo-}-mNeonGreen before (0 hr) and after (4, 8 hrs) damage induction. Representative western blot image is shown at the [top] and levels are quantified at the [bottom]. n=3 independent repeats. Mean and S.D. are shown. Significance was calculated using Repeated Measures one-way ANOVA and post-hoc tests.

Figure EV4: Exonuclease activity of Mip1 drives mtDNA loss under damage

(A) [Left] Localization of Mip1-CΔ263-mNeonGreen (i) or *mip1^{exo-}*-CΔ263-mCherry (ii) and mt-nucleoids stained with DAPI in *mip1^{exo-}* cells. [Right] Line profiles of Mip1-CΔ263-mNeonGreen (i) or Mip1^{exo-}-CΔ263-mCherry (ii) and mt-nucleoids stained with DAPI in *mip1^{exo-}* cells. Line profile for a representative across an ROI is shown (white line in merged panel). Dashed lines represent cell boundaries. **(B)** Western blot of Mip1-mCherry, Mip1-CΔ263-mCherry and Mip1^{exo-}-CΔ263-mCherry. Representative image with loading control (Coomassie) is shown from one of the 3 independent repeats. **(C)** Western blot of Mip1-mNeonGreen, Mip1^{G651S}-mNeonGreen and Mip1^{H734Y}-mNeonGreen. Representative image with loading control (Coomassie) is shown from one of the 3 independent repeats. **(D)** Multiple sequence alignment of Human PolG and related orthologues, including Mip1, performed using COBALT to show the conservation and position of G651 and H734 across organisms. **(E)** [Left] Localization of Mip1^{G651S}-mNeonGreen (i) or Mip1^{H734Y}-mNeonGreen (ii) and mt-nucleoids stained with DAPI in *mip1^{exo-}* cells. [Right] Line profiles of Mip1^{G651S}-mNeonGreen (i) or Mip1^{H734Y}-mNeonGreen (ii) and mt-nucleoids stained with DAPI in *mip1^{exo-}* cells. Line profile for a representative across an ROI is shown (white line in merged panel). Dashed lines represent cell boundaries. Scale bar, 8μm here, and in all other images.

Figure EV5: Rates of mtDNA loss are influenced by dNTP levels

(A) Western blot of Mip1-mNeonGreen in wild type and Δ*sml1* cells before (0 hr) and after (4, 8 hrs) damage induction. Representative western blot image is shown on the [left] and levels are quantified on the [right]. n=3 independent repeats. Mean and S.D. are shown. Significance was calculated using Repeated Measures one-way ANOVA and post-hoc tests. **(B)** Serial dilution growth assay to measure growth of Δ*sml1* cells in comparison to a wild type control.

952   Representative image from 3 independent repeats is shown. Scale bar refers to the increasing

953   dilution of cells from the left spot to the right-most spot.

A. Replicative polymerase Mip1

B. *mip1^exo-*

C.

D. 0 hr

E. *mip1^exo-*

8 hrs

A. Mip1 — wild type — P = 0.0361

B. *mip1^exo-* / Mip1-CΔ263 — *mip1^exo-* + Mip1-CΔ263 — P = 0.0083

C. *mip1^exo-* / *mip1^exo-*-CΔ263 — *mip1^exo-* + *mip1^exo-*-CΔ263

D. *thumb domain (catalysis)* — *mip1^exo-* / Mip1 G651S — *mip1^exo-* + Mip1^G651S — P = 0.0077

E. *finger domain (nucleotide binding, processivity)* — *mip1^exo-* / Mip1 H734Y — *mip1^exo-* + Mip1^H734Y — P = 0.0464

empty vector
mtDarT

A.

Budding yeast cells

mtDarT

empty vector

i) Transform with empty vector or mtDarT

ii) Select transformants

β-estradiol

0hr   4hrs   8hrs

iii) Grow cultures to mid-exponential in YPG

■ Selectable marker
■ mtDarT

iv) Collect 0hr sample and induce with β-estradiol

v) Collect samples 4hrs and 8hrs after induction

B.

β-estradiol
↓

0hr   4hrs   8hrs

mtDarT-3x-FLAG

coomassie

C.

0nM β-estradiol   100nM β-estradiol

empty vector

mtDarT

YPG+NAT

dilution

D.

mitochondria (Cox4-mCherry)

nucleoids (SGI)

red - Cox4-mCherry
green - SGI

E.

Δ*fzo1* (ρ⁰) (SGI)

A.

i) Mip1-CΔ263-mNg | DAPI | green - Mip1-CΔ263-mNg / blue - DAPI

ii) Mip1^exo⁻-CΔ263-mCh | DAPI | green - Mip1^exo⁻-CΔ263-mCh / blue - DAPI

B.

C.

D. G651 / H734

Source of PolG-a sequence

CONSENSUS▶	A I L P Q V V X A G T I T R R A V E P T	M T L Q G X K S X G T D L H S K T A X X
Homo sapiens	A I L P Q V V T A G T I T R R A V E P T 858	M T L Q G R K S R G T D L H S K T A T I 938
Saccharomyces cerevisiae	I I I P K I V P M G T I T R R A V E N A 661	M C L E G T K N E G T D L H T K T A Q I 740
Drosophila melanogaster	A I C P Q V V A C G T L T R R A M E P T 776	M T L S G S K S N G S D M H S I T A K A 856
Mus musculus	A I L P Q V V T A G T I T R R A V E P T 837	M T L Q G R K S R G T D L H S K T A A T 917
Danio rerio	A I L P Q V I P A G T V T R R A V E P T 819	M T L Q G K K S Q G T D L H S R T A D A 899

% conservation

E.

i) Mip1^G651S-mNg | DAPI | green - Mip1^G651S-mNg / blue - DAPI

ii) Mip1^H734Y-mNg | DAPI | green - Mip1^H734Y-mNg / blue - DAPI

A.

Mip1 mNg (mtDarT)

Mip1 mNg in Δ*sml1* (mtDarT)

0hr 4hrs 8hrs   0hr 4hrs 8hrs

coomassie

● Mip1 mNg (mtDarT)
● Mip1 mNg in Δ*sml1* (mtDarT)

B.

wild type

Δ*sml1*

YPG

dilution

Reagents and Tools table

Reagent/Resource	Reference or Source	Identifier or Catalog Number
Experimental Models/Yeast strains		
CEN.PK *(S. cerevisiae)* Mat**a**	(Van Dijken et al., 2000)	NABC567
CEN.PK *(S. cerevisiae)* Matα	(Van Dijken et al., 2000)	NABC568
Δ*bar1 (S. cerevisiae)* Mat**a**	Dr. Sunil Laxman Lab stock	NABC595
Δ*sml1 (S. cerevisiae)* Mat**a**	This study	NABC639
Δ*nuc1 (S. cerevisiae)* Mat**a**	This study	NABC641
Mip1-mCherry *(S. cerevisiae)* Mat**a**	This study	NABC772
Mip1 mNeongreen *(S. cerevisiae)* Mat**a**	This study	NABC773
Mip1^{exo-} *(S. cerevisiae)* Mat**a**	This study	NABC573
Mip1^{exo-}-mNeongreen *(S. cerevisiae)* Mat**a**	This study	NABC680
Δ*sml1* Mip1-mNeongreen *(S. cerevisiae)* Mat**a**	This study	NABC766
Cox4-mCherry *(S. cerevisiae)* Mat**a**	This study	NABC569
Δ*atg32 (S. cerevisiae)* Mat**a**	This study	NABC676
Recombinant DNA		
pFA6a-KO-Hyg	Dr. Sunil Laxman Lab stock	pNABC552
pFA6a-KO-KanMX6	Dr. Sunil Laxman Lab stock	pNABC551
pFA6a-mCherry-Hyg	Dr. Sunil Laxman Lab stock	pNABC549
pFA6a-mNeongreen-Kan	This study	pNABC553
pGEV-3XFLAG-NAT	AB Lab stock, Dua et al, 2022	NABC557
pGEV-Su9-DarT-3XFlag-NAT	AB Lab stock, Dua et al, 2022	pNABC561
p417-cyc1-Kan	Dr. Sunil Laxman Lab stock	pNABC555
pMip1-Mip1-CΔ263 in NABC555 background	This study	pNABC775
Hamamatsu Photonics	This study	pNABC777
pMip1-Mip1-CΔ263-mCherry in NABC555 background	This study	pNABC943
pMip1-mip1^{exo-}- CΔ263 in NABC555 background	This study	pNABC944

Reagents and Tools table (continued)

Reagent/Resource	Reference or Source	Identifier or Catalog Number
pMip1-mip1^{exo-}- CΔ263-mCherry in NABC555 background	This study	pNABC945
pMip1-mip1-G651S in NABC555 background	This study	pNABC946
pMip1-mip1-G651S-mNeonGreen in NABC555 background	This study	pNABC947
pMip1-mip1-H734Y in NABC555 background	This study	pNABC948
pMip1-mip1-H734Y-mNeonGreen in NABC555 background	This study	pNABC949
Antibodies		
Rabbit Anti-Flag	Cell signalling	14793S
Mouse Anti-mCherry	Sigma	SAB2702291
Rabbit Anti-mCherry	Agrisera	AS18 4179
Rabbit Anti-mNeongreen	Agrisera	AS21 4525
Anti-rabbit IgG, HRP-linked Antibody	Cell Signaling Technology	7074S
Anti-mouse IgG, HRP-linked Antibody	Cell Signaling Technology	7076S
Oligonucleotides and other sequence-based reagents		
GGAGAACAGGAGAAAGAAGG	This study	AS78
GGTTGAGCTGGAAAGGGACATTACTATTTCTAGAGAGTACcggatccccgggttaattaa	This study	AS79
ATATAAATACAAATGCGAAAGCTAATGCAGATTTTGCCTAgaattcgagctcgtttaaac	This study	AS80
TTAATTAACCCGGGGATCCG	This study	AS81
CAAACAAGATCACCAAAAGG	This study	AS91
AGTTCCCCACTTATTAGATCCCAAAGTTAAAGTTATCATGcggatccccgggttaattaa	This study	AS92
TTATGAATTAATTTATATTTACAGTTTTTCAGTACATTCAgaattcgagctcgtttaaac	This study	AS93
ATGACGAAATTGATGGTTAGATCTGAATGCATG	This study	AS112
CTAGTACTCTCTAGAAATAGTAATGTCCCTTTC	This study	AS113

Reagent/Resource	Reference or Source	Identifier or Catalog Number
AATCGTCTTTCACAGTTTGC	This study	AS161
ACAAAAGCAAAAAAAATCTGCCAGGAACAGTAAACATATGcggatccccgggttaattaa	This study	AS162
AGTAGGAACGTGTATGTTTGTGTATATTGGAAAAAGGTTAgaattcgagctcgtttaaac	This study	AS163
gctgccgccgctcgcgctgctgccgctgccGCTGTCGATATCGAGATTTGGTTTACC	This study	AS183
ggcagcggcagcagcgcgagcggcggcagcATGGCTTCTTTGCCAGCTACTC	This study	AS184
CCCTCACTAAAGGGAACAAAAGCTGGAGCTCgcagaaagaaccttcaacctcattttgtt	This study	AS188
GATATCGAATTCCTGCAGCCCGGGGGATCCtcaagaattagaagaagcagaacctgaacc	This study	AS192
ggcagcggcagcagcgcgagcggcggcagcATGGTGAGCAAGGGCGA	This study	AS200
gatatcgaattcctgcagcccggggggatccCTACTTGTACAGCTCGTCCATG	This study	AS201
GATATCGAATTCCTGCAGCCCGGGGGATCCCTAgctgtcgatatcgagatttggtt	This study	AS202
ggcagcggcagcagcgcgagcggcggcagcATGGCTTCTTTGCCAGCTACTCATGAATTG	This study	AS225
GAAAATCGTCCCCATGtcCACAATCACTAGAAGA	This study	AS271
TCTTCTAGTGATTGTGgaCATGGGGACGATTTTC	This study	AS272
GAAGGTACAGATTTGtACACGAAGACTGCTCAA	This study	AS273
TTGAGCAGTCTTCGTGTaCAAATCTGTACCTTC	This study	AS274
ACTAAACCCTGTTGGTGTTCCAAATGATGACCACCATCACcggatccccgggttaattaa	This study	ND328
GTAAAAGAGAAACAGAAGGGCAACTTGAATGATAAGATTAgaattcgagctcgtttaaac	This study	ND329
GGCATTAAATTATCGTCGTG	This study	ND307

Reagents and Tools table (continued)

Reagent/Resource	Reference or Source	Identifier or Catalog Number
TCAAAAATTGATGGTGGAAGAAATAAATTTAGACATCGATcggatccccgggttaattaa	This study	ND308
ATTGATTTGAAAAGACCTCATATATTTACAAGAATATCTAgaattcgagctcgtttaaac	This study	ND309
GAAAACTGCCCAAAACTTAG	This study	ND327
GGGATTATATGTAGTTGTTGAGCAACGAGGGACAAGTATGcggatccccgggttaattaa	This study	ND334
ATATAAATACAAATGCGAAAGCTAATGCAGATTTTGCCTAgaattcgagctcgtttaaac	This study	ND335
TATGTAGTTGTTGAGCAACGAGGGACAAGTATGACGAAATTGATGGTTAGATCTGAATGCATGCTG	This study	ND339
CTGGTGGTGTTTGCTGTAGCAACACTCTATAAC	This study	ND340
CGTTATAGAGTGTTGCTACAGCAAACACCACCAG	This study	ND341
CTAGTACTCTCTAGAAATAGTAATGTCCCTTTCCAG	This study	ND342
AGGGACATTACTATTTCTAGAGAGTACTAGGACATGGAGGCCCAGAATACCCTCCTT	This study	ND343
TACAAATGCGAAAGCTAATGCAGATTTTGCCTTCGAGCGTCCCAAAACCTTCTCAAGC	This study	ND344
GGCCAATGATAGGAAAGAAC	This study	ND488
CACTAACCTCTCTTCAACTGCTCAATAATTTCCCGCTATGcggatccccgggttaattaa	This study	ND489
AAATGGAAAGAGAAAAGAAAAGAGTATGAAAGGAACTTTAgaattcgagctcgtttaaac	This study	ND490
Chemicals, Enzymes and other reagents		
SYBR™ Green I Nucleic Acid Gel Stain	Invitrogen	S7563
Molecular Probes SYTOX Green Nucleic Acid Stain	Thermo Fisher Scientific	S7020
AgriseraECL SuperBright	Agrisera	AS16 ECL-S
β-estradiol	Sigma-Aldrich	E2758
Ultrapure agarose	Invitrogen	16500500

Reagents and Tools table (continued)

Reagent/Resource	Reference or Source	Identifier or Catalog Number
α1-Mating Factor acetate salt	Sigma-Aldrich	T6901
Nocodazole	Sigma-Aldrich	487928
Hygromycin B	Thermo Fisher Scientific	10687010
Nourseothricin	Jena Bioscience	AB-102XL
DAPI (4',6-Diamidino-2-Phenylindole, Dihydrochloride)	Invitrogen	D1306
Geneticin™ Selective Antibiotic (G418 Sulfate), Powder	Thermo Fisher Scientific	11811031
RNAse A	Genetix	EN0531
Proteinase K	Invitrogen	AM2548
Softwares		
ImageJ FIJI 2.16.0		https://imagej.net/software/fiji/
Graphpad Prism 8.4.2		https://www.graphpad.com
Cell ACDC	(Padovani et al., 2022)	https://github.com/SchmollerLab/Cell_ACDC
RStudio Version 1.3.1093		https://posit.co/products/open-source/rstudio/
NIS-elements software (version 5.01)		Nikon Instruments Inc.
COBALT: multiple alignment tool	(Papadopoulos & Agarwala, 2007))	https://www.ncbi.nlm.nih.gov/tools/cobalt/re_cobalt.cgi
Others		
Accuri C6+ flow cytometer		BD Biosciences
Nikon Eclipse Ti2-E		Nikon Instruments Inc.
pE-4000		CoolLED
ORCA-Flash4.0 V3 Digital CMOS camera		Hamamatsu Photonics